Behavior-specific occupancy patterns of Pinyon Jays (*Gymnorhinus cyanocephalus*) in three Great Basin study areas and significance for pinyon-juniper woodland management

(Short Title): Behavior-specific Pinyon Jay occupancy patterns

John D. Boone^{1*}, Chris Witt², and Elisabeth M. Ammon¹

¹Great Basin Bird Observatory, Reno, Nevada, United States of America

²U.S. Forest Service Rocky Mountain Research Station, Boise, Idaho, United States of America

*Corresponding author. E-mail boone@gbbo.org

Abstract (Level 1 heading)

The Pinyon Jay is a highly-social, year-round inhabitant of pinyon-juniper woodlands in the western United States. Range-wide, Pinyon Jays have declined ~ 3 – 4% per year for at least the last half-century. At the same time, large acreages of pinyon-juniper woodland have been removed or thinned to improve habitat for Greater Sage-Grouse or other game species across much of the Great Basin, which is home to nearly half of the global population of Pinyon Jays. Occupancy patterns and habitat use of Pinyon Jays have not been well characterized across much of the species' range, and obtaining this information is necessary for better understanding the causes of ongoing declines and determining useful conservation strategies. Our goal of this study was to identify the characteristics of areas used by Pinyon Jays for several critical life history components and to thereby facilitate the inclusion of Pinyon Jay conservation measures in the design of vegetation management projects. To accomplish this, we studied Pinyon Jays in three widely separated study areas using radio telemetry and direct observation, and measured key attributes of their locations and a separate set of randomly-selected control sites using the U. S. Forest Service's Forest Inventory Analysis protocol. Data visualizations, non-metric dimension scaling ordinations, and logistic regressions of the resulting data indicated that Pinyon Jay occupancy was concentrated in a distinct subset of available pinyon-juniper woodland habitat, and further that Pinyon Jays used different habitats, arrayed along elevational and tree-cover gradients, for seed caching, foraging, and nesting. Caching was concentrated in low-elevation, relatively flat areas with low tree cover; foraging occurred at slightly higher elevations with moderate tree cover, and nesting was concentrated in somewhat higher areas with greater tree cover and higher stand density. All three of these Pinyon Jay behavior types were highly concentrated within the lower-elevation band of pinyon-juniper woodland close to the woodland-

shrubland ecotone. Because woodland removal projects in the Great Basin are often concentrated in these same areas, it is critical to incorporate conservation measures informed by Pinyon Jay occupancy patterns into existing woodland management paradigms, protocols, and practices.

Introduction (level 1 heading)

The Pinyon Jay (*Gymnorhinus cyanocephalus*) is a highly-social corvid that inhabits pinyon-juniper and other coniferous woodlands in the interior western U. S. [1-3] (Fig 1). Pinyon Jays form year-round flocks that can range from a few dozens to several hundred members [3-6]. They are perhaps best known for harvesting and caching the seeds, or “pine nuts”, of the pinyon pine (primarily *Pinus monophylla* and *P. edulis*) as their primary food source, though they also consume other conifer seeds and insects [3, 7, 8]. Pinyon Jays are present year-round in parts of at least ten states, but most of their range lies within Bird Conservation Regions (BCRs) 9 (“Great Basin”) and 16 (“Southern Rockies and California Plateau”) [9], with highest densities in central-eastern Nevada and western New Mexico (Fig 2). Since North American Breeding Bird Survey data collection began in 1967, Pinyon Jays have experienced steep and sustained declines averaging 3-4% per year both range-wide and within most of the states and regions they occupy [4, 10]. This equates to a loss of about 85% of the population over 50 years, one of the largest recorded declines among all widely-distributed passerine birds in the interior western United States [10, 11].

Fig 1. Distribution of the Pinyon Jay [4], pinyon pine (*Pinus monophylla* and *P. edulis* combined), and juniper (*Juniperus osteosperma*, *J. occidentalis*, and *J. spoculorum* combined) [5]. Some of the juniper distribution is invisible under the pinyon pine distribution on the map.

Fig 2. Relative density of the Pinyon Jay (purple colors, with darker colors representing higher relative densities [4]) and Bird Conservation Regions (BCRs) 9 (Great Basin) and 16 (Southern Rockies / Colorado Plateau).

Despite this decline, no systematic conservation efforts have been undertaken for the Pinyon Jay, although it is included on the ‘sensitive species’ lists of many federal and state management agencies and avian conservation organizations [12-15], and is the subject of a recent interagency working group conservation strategy [15]. The lack of conservation action for Pinyon Jays may be attributable to several factors. First, despite a rich knowledge of the species’ social behavior, breeding behavior, and spatial memory [6, 16, 17], its occupancy and habitat use patterns are poorly characterized and understudied in most parts of its range. Second, there are no widely-accepted or strongly-supported hypotheses about the causes of Pinyon Jay population declines. Finally, the landscapes inhabited by Pinyon Jays are primarily managed for other priorities. These include improving habitat for game species such as Greater Sage-Grouse (*Centrocercus urophasianus*) and mule deer (*Odocoileus heminus*), creating wildlife corridors, and mitigating fire hazards [3, 18-22].

Pinyon Jay declines could be related, at least in part, to changes in the pinyon-juniper woodlands that comprise most of their habitat [3, 22-24] and much of the forested landscape of the Great Basin (i.e., BCR 9) and Colorado Plateau (i.e., BCR 16) [25-28]. The spatial extent of pinyon-juniper woodlands (most commonly a *Pinus monophylla* - *Juniperus osteosperma* association) in this region has undergone climate-induced fluctuations since the end of the Pleistocene epoch 11,500 years ago [29, 30], but it has been suggested by some authors that over the last 150 years, extension of local woodland range (i.e. “expansion”) and increased tree densities within extant stands (i.e. “infill”) have occurred at atypical and perhaps unprecedented rates, at least in the Great Basin [22, 31-34]. Other authors have questioned this conclusion and suggest that expansion and infill are either localized, part of a historically normal pattern of spatio-temporal woodland dynamics, or represent recoveries from earlier widespread clearing during the western settlement period [22, 35, 36]. Regardless of which of these paradigms is correct, over recent decades extensive acreages of pinyon-juniper woodlands have been clear cut or thinned by resource management agencies and private owners to accomplish various management objectives [3, 37, 38].

In the Great Basin, the primary pinyon-juniper woodland management objective over the last 20 years has been creation or restoration of shrubland habitat by clear cutting stands that are regarded as encroaching into shrublands, often with a goal of benefitting Greater Sage-Grouse [39, 40]. If and how this type of vegetation management affects Pinyon Jays remains undetermined. Answering this question requires a better understanding of Pinyon Jay occupancy patterns and habitat use in the Great Basin, determining the extent of overlap between their preferred habitats and ongoing vegetation management activities, and monitoring or modeling

the effects of those activities. In this study, we compared the habitats used by Pinyon Jays for caching or retrieving seeds, for foraging, and for nesting, to the full range of habitat available within pinyon-juniper woodlands. Our goals were to determine whether Pinyon Jays in the Great Basin used predictable subsets of available habitat for these distinct behavior types. If so, this information could assist managers seeking to incorporate Pinyon Jay conservation measures into existing vegetation management programs for pinyon-juniper woodlands, and help to guide future research [15].

Methods (level 1 heading)

Overview of study design and study region (level 2 heading)

We used a case-control study design [41, 42], with observed locations of Pinyon Jay flocks as case records and a set of pre-existing Forest Inventory Analysis (FIA) plots established by the U.S. Forest Service (USFS) as control sites (see below for details). Pinyon Jay data were collected from 2008 – 2014 in three Great Basin study areas. The control sites were distributed over a broader study region located entirely within the Central Basin and Range Ecoregion [43] and BCR 9 (Fig 2) that encompassed all three Pinyon Jay study areas (Fig 3). Control sites were not assumed to be unoccupied by Pinyon Jays, but were intended to provide a representative characterization of the diversity of pinyon-juniper woodland habitat available within the general study region. Habitat at Pinyon Jay locations and control sites was quantified using the standardized FIA protocol (see below) within circular 0.405-ha (1-acre) plots, which defined the

spatial scale of this analysis. Analyses consisted of ordinations and logistic regressions, supplemented by data visualizations. We recognize the potential bias inherent in case-control sampling designs and followed Manly et al.'s [44] and Keating and Cherry's [42] guidance for analysis.

Fig 3. Locations of three Pinyon Jay study areas and FIA plots that served as control sites. FIA plot locations are approximate (i.e. "fuzzed") to comply with USFS data protection policies. FIA plots are color coded to indicate the Pinyon Jay study area with which they were analytically paired.

Pinyon Jay study areas and timeline (level 2 heading)

Pinyon Jay locations came from multiple Pinyon Jay flocks in these three study areas (Fig 3):

- 1) Eastern Nevada (~ 63,130-ha study area), specifically the foothills between Baker, Nevada and Great Basin National Park, and Steptoe Valley south of the town of Ely, Nevada. This area is mostly comprised of public lands managed by the U.S. Bureau of Land Management, but includes some private property near Baker. Pinyon Jay data were collected from 2/28/2008 – 6/26/2008 and from 5/10/2009 – 8/27/2009.
- 2) Southern Idaho (~ 23,470-ha study area), specifically the area in and around City of Rocks National Reserve and Castle Rocks State Park in Cassia County. This study area contains the northernmost occurrence of pinyon pine in North America [45]. Jurisdictions within the study area are the U.S. National Park Service, Idaho State Parks, U.S. Bureau

of Land Management, and private lands. Pinyon Jay data were collected from 7/20/2012 – 10/5/2012.

3) Central Nevada (~ 89,030-ha study area), specifically the Desatoya Range which lies along the border between Churchill County on the west and Lander County on east. This study area is comprised almost exclusively of U.S. Bureau of Land Management lands, with limited private inholdings. Pinyon Jay data were collected from 3/29/2013 – 6/5/2013.

Study area locations and periods of data collection were governed by three different funding agreements, and therefore seasonality was not standardized across all three areas. All study areas are characterized by mountainous “basin and range” topography, with plant communities dominated by big sagebrush (*Artemisia tridentata*) at lower elevations, pinyon-juniper woodlands at mid-elevations, and various mountain shrub and forest types at high elevations. Pinyon-juniper woodlands range in elevation from ~ 1,500 m – 2,600 m across all study areas, and are usually comprised of varying proportions of *P. monophylla* and *J. osteosperma*. Public lands in all study areas are managed for multiple uses and experience varying levels of livestock grazing and off-road vehicle travel.

Pinyon Jay data collection and processing (level 2 heading)

Pinyon Jay data collection had several distinct components; initial searches, observational surveys, capture and radio-tagging, and radio-telemetry surveys. The observational surveys and radio-telemetry surveys generated the Pinyon Jay locations that are the basis of our analyses.

171

172 Initial searches of potential Pinyon Jay habitat (defined as pinyon-juniper woodlands and
173 visually adjacent shrublands) were conducted during the first 1-2 weeks of field work at a given
174 study area on foot and by vehicle to identify all or most of the Pinyon Jay flocks present.
175 Because Pinyon Jay flocks are visually apparent from long distances, “noisy” (except when at
176 the nest), and spatially segregated from one another, we regarded this a feasible goal. Search
177 patterns used during initial searches were not systematized but were instead tailored to take
178 advantage of local topography and access points. Areas searched were delineated on imagery
179 maps to facilitate thorough coverage of pinyon-juniper woodland across a given study area.
180 Upon detecting a flock, the observer usually maintained contact over a period of 1-3 h on each of
181 several visits to obtain a preliminary and approximate delineation of the flock’s movement
182 patterns and primary activity areas.

183

184 Flocks that were consistently detected during initial searches were then subjected to more
185 intensive study by observational surveys and/or radio-telemetry surveys to obtain a sample of
186 occupied locations. Observational surveys involved establishing visual contact with a flock;
187 observing the flock with binoculars from a distance sufficient to prevent alteration of flock
188 behavior (typically > 75 m); and recording approximately once per hour the point coordinates of
189 the estimated centroid of the flock’s location along with estimated flock size and predominate
190 behavior type (Table 1) whenever possible. The goal of observational surveys was to obtain
191 locations across an entire daylight cycle at least once per week (usually assembled from several
192 observation sessions conducted during different time periods on different days) for each flock
193 over the duration of the data collection period. Coordinates of flocks were usually obtained by

recording observer position with a GPS unit; recording a bearing to the visually-estimated flock centroid with a compass; measuring distance to the flock centroid by rangefinder or estimating distance by eye (for shorter distances < 25 m); and then plotting the flock's estimated point location in GIS based on these parameters. In some cases, coordinates could be obtained directly by GPS after a flock vacated a previously-occupied location. As observers became increasingly familiar with the daily movement patterns of a flock, which tended to be consistent, their efforts were increasingly focused on the portions of the daily activity cycle that were more difficult to characterize.

Table 1. Description of Pinyon Jay behavior types that were recorded during observational and radio-telemetry surveys whenever possible.

Behavior Type	Description
Caching	Birds observed either caching or retrieving previously-cached pine nuts or other similar food items.
Foraging	Birds observed collecting any type of new food item in trees, shrubs, or on the ground, including pinyon nuts, other plant material, or insects.
Nesting	Nest site or sites confirmed by direct observation or by confirmatory behaviors, such as carrying nest materials.
Roosting ¹	Birds present at a location during period(s) of darkness.
Loafing ²	Birds observed resting, usually in trees, while not engaged in any other listed behaviors.
Flyover ²	Entire flock flying in a directional manner, without landing. Short flight movements by one or a few birds while the flock was engaged in one of the other listed behaviors were not considered flyovers.

¹Due to small sample size, this behavior type was used only for ordination analyses.

²Observations for these behavior types were excluded from analysis.

Radio-telemetry surveys were also used to obtain Pinyon Jay locations for some of the studied flocks. Two methods were used to capture Pinyon Jays for radio-tagging; baited walk-in traps and mist nets. Traps were used where Pinyon Jays could be consistently drawn (determined by game cameras) to a supplemental food station baited with shelled peanuts, sunflower seeds, and dried corn. In these areas, a home-made wood and mesh walk-in trap (0.6 x 1.2x 0.9 m) with an

open door was placed at the site, and bait spread periodically inside the trap to habituate birds to entering the trap. When habituation was sufficient (determined by camera traps), capture attempts were made by rigging the door for remote manual release, observing activity at the trap from a blind ~ 25 m from the trap, and releasing the door with a pull cord when Pinyon Jays were inside the trap. Mist-nets were used in areas where walk-in traps were not viable, specially in locations routinely visited by Pinyon Jays where nets could be deployed without being easily detected by birds. Mist nets (60 mm mesh) were arrayed either singly, as doubles, or stacked, depending on the geometry of the woodland opening where they were erected. In some cases, call playback was also used to try to draw birds into the nets. Any Pinyon Jays captured by walk-in trap or mist net were weighed and aged; standard aluminum leg bands (U.S. Geological Survey) were attached; and radio transmitters (Advanced Telemetry Systems model A2450) were glued onto feather stubble clipped to ~ 0.3 cm above skin level in the interscapular area. No more than six individuals from any single flock were radio-tagged, since data were collected to characterize flocks rather than individual birds. Captured birds were handled and processed only by experienced individuals holding a U.S. Fish and Wildlife Service master banding permit. Radio-tagged birds were manually tracked using a handheld three-element Yagi antenna with an Advanced Telemetry Systems R410 or R100 receiver. The goal of radio-telemetry surveys was to collect hourly locations over one entire daylight cycle per week for each flock, either in a single long session or over several sessions of 3 – 5 hours on different days and at different times. Most often, telemetry fixes from one or more radio-tagged flock members were used to approach the flock and establish visual contact, at which point locational and behavior type attributes were collected exactly as described above for observational surveys. Occasionally, flock locations were estimated by biangulation or triangulation of telemetry bearings that were

post-processed using LOAS software (Ecological Software Solutions, LLC). In cases where the flock was not directly observed, behavior type was only recorded when it could be reasonably inferred based on previous observations (i.e. return to a nest colony location) or context (roosting locations at after sunset).

For a given flock, locations could be obtained by either observational surveys, radio-telemetry surveys, or both, but because both approaches ultimately relied on the same observational process, we regarded the resulting data as comparable. Initially, radio-telemetry was the preferred survey approach, but by later data collection periods we determined that visual contact with most flocks could routinely be established without reliance on radio-tagged flock members, and our efforts to capture birds for radio-telemetry were discontinued.

The full set of Pinyon Jay locations recorded during field work were filtered and processed prior to analysis. First, any locations where no behavior type was recorded were deleted. Next, all “flyover” locations and loafing locations (Table 1) were deleted, based on the premise that these could occur throughout and sometimes beyond the flock’s core home range. Finally, locations that represented the same flock engaged in the same behavior at the same location were combined as follows:

- 1) All nesting locations for a single flock were spatially averaged using the Mean Center tool in ArcMap 10.5 (ESRI, Redlands CA), based on the premise that at a given time, a flock has only one nesting area.
- 2) If caching, foraging, or roosting locations were closer together than 71.8 m, they were spatially averaged to generate a single location, as above. This was the smallest threshold

distance sufficient to prevent any overlap between the 0.405-ha habitat plots used to assess habitat (see below), so this process smoothed Pinyon Jay locations to a spatial scale that matched the habitat assessment scale.

After these steps, 152 Pinyon Jay locations with recorded behaviors were retained for analysis (see Table S1).

Data collection procedures described in this section were primarily observational and were not submitted to or approved by an Institutional Animal Care and Use Committee. Field work that involved capture, radio-tagging, and release of Pinyon Jays was conducted under U.S. Department of the Interior Federal Bird Banding Permit # 22912, Idaho Fish and Game Department Collection Permit # 120724, and Nevada Department of Wildlife Collecting Permit # 29948.

Control site selection (level 2 heading)

Control sites were selected from the pre-existing FIA data set of plots visited and measured between 2005-2013. FIA data provide a probabilistic and geographically unbiased assessment of forest / woodland attributes over time and space [46, 47] and a robust dataset to describe habitat for multiple species [48-52]. One FIA plot is randomly positioned within every cell of a sampling grid that covers all public and private forest land in the United States, with an overall density of one plot per 2,428 ha (6,000 acres) within the extent of that sampling frame [53]. Each year, 10% of these plots are surveyed or re-surveyed in the western U.S. using a spatially-

interpenetrating sampling design that avoids conflation of spatial with temporal trends [46]. The criteria used to select the subset of FIA plots that were appropriate control sites for this study were as follows:

- 1) The FIA plot had to be < 200 km from the nearest Pinyon Jay location retained for analysis.
- 2) Presence of pinyon-juniper woodland within the FIA plot had to be confirmed by direct observation of the field assessment crew during the most recent assessment visit. A woodland type classification based solely on remote-sensing data was not sufficient to meet this criterion.
- 3) The site had to be located in both the Central Great Basin Ecoregion [43] and in BCR 9.
- 4) The most recent assessment visit had to have occurred in 2005 or later.

In total, 346 FIA plots from the larger data set met these criteria (Fig 3) and were included in our analysis (see Table S1). Each control site was assigned a regional attribute that corresponded to the closest Pinyon Jay study area (n = 212 for the Eastern Nevada region, n = 81 for the Southern Idaho region, and n = 53 for the Central Nevada region). Control sites were not distributed symmetrically around the Pinyon Jay study areas (especially for Southern Idaho) because their extent was constrained by the distribution of pinyon-juniper woodland and by the USFS site selection process.

Habitat assessment (level 2 heading)

Forested habitat at Pinyon Jay locations was characterized using current FIA field protocols [54]. Site attribute measurements at Pinyon Jay locations on non-forested land followed procedures outlined for a FIA All-Conditions Inventory [55]. For all Pinyon Jay activity sites, FIA plot centers were placed at the Pinyon Jay location point coordinates and assessments were performed by USFS crews fully trained in FIA protocols and procedures. Control sites were assessed as part of routine USFS operations in various years between 2005-2013 [56-58].

FIA plot layout is based on a 0.405 ha circle that defines four subplots of 7.3 m diameter within which actual measurements are made. One subplot is located at the center of the larger circle, and the other three subplots have their centers equally spaced along its circumference (Fig 4). On a standard forested FIA plot, over 120 attributes are measured within each subplot to characterize location, condition, and vegetation [59]. Subplot data are then averaged or summed over subplots as appropriate and extrapolated to generate data at the whole-plot scale. The subset of FIA attributes that were considered for use in this analysis are presented and briefly described in Table 2.

Fig 4. Schematic diagram of a USFS FIA plot layout. Data used in this study consisted of full-plot values produced by either summing or averaging sub-plot data.

Table 2. Attributes describing Pinyon Jay habitat that were considered in this study. Brief descriptions of attributes are provided parenthetically, with complete descriptions of associated methodologies available in [59]. The 2nd, 3rd, and 4th columns indicate whether or not an attribute was used (Y = yes, N = No, NA = categorical attribute, not applicable) for the ordinations, logistic regressions, and data visualizations presented below. Footnotes provide additional explanations about attribute use considerations.

Attribute	Ordination	Logistic Regression	Data Visualization
FIA Attributes			
Elevation (measured in feet at plot center)	Y	Y	Y
Slope (expressed as slope percentage, or {{rise/run} x 100} at plot center)	Y	Y	Y
Habitat Type [60]	NA	N ¹	N
Stand Age (age in years of the oldest pinyon pine or juniper tree on plot, as determined by coring)	Y	Y	Y
Stand Density Index (index of three-dimensional tree density within stand [60])	N ²	N ²	Y ²
Canopy Cover (% by ocular estimation)	Y	N ³	N
Tree Cover (% by ocular estimation)	Y	Y	Y
Shrub Cover (% by ocular estimation)	Y	Y	Y
Forb Cover (% by ocular estimation)	Y	Y	Y
Grass Cover (% by ocular estimation)	Y	Y	Y
Woody Debris (count of pieces of woody debris material along transects for seven different size diameter classes, defined by twig / branch diameter ranges)	Y	Y ⁴	Y ⁴
Distance to Road (km from plot center, assigned using topographic maps; original ordinal ranges converted to range midpoints)	Y	Y	Y
Disturbance Presence (“yes-no”, for any disturbances within the last five years)	NA	Y	N
Disturbance Type (categories of silvicultural treatment or other disturbance occurring within the previous five years)	NA	N ⁵	N
Non-FIA Attribute			
Distance to Edge (Distance to the lower-elevation woodland-shrubland ecotone, measured directly in on imagery in ArcMap 10.5)	N ⁶	N ⁶	Y

¹ All sites used in analysis were characterized by presence of at least some pinon pine and/or juniper trees, but the FIA habitat type distinctions were too fine-grained (33 different types, median number of sites / type = 5) for inclusion in analysis.

²Omitted from analyses due to high correlations with other attributes, but included in data visualizations because of possible synthetic interpretational value.

³Omitted because of high correlation to Tree Cover attribute ($r = 0.68$).

⁴Reduced to a single attribute by Principle Component Analysis that described 86.1% of variation across all original classes.

⁵The level of articulation was too fine for the limited number ($n=68$) of disturbed sites and too unevenly distributed among types, so only the “yes-no” disturbance attribute was retained.

⁶Omitted from ordinations and logistic regression because attribute estimation process was not part of the standardized FIA protocol. Retained in data visualizations for possible interpretive value

320

321 Based on our field observations of Pinyon Jay landscape use over the course of the study, we
322 decide to create one additional non-FIA attribute, “Distance to Edge” (Table 2) which is the
323 shortest linear distance between a Pinyon Jay location or control site and the lower-elevation
324 woodland-shrubland ecotone. First, polylines were digitized in ArcMap 10.5 to delineate the
325 approximate ecotonal boundary based on visual examination of imagery. This process did not
326 delineate woodland – shrubland boundaries inside well-defined woodland patches, but instead
327 defined distinct woodland patches based on their lowest-elevation ecotonal boundary. Then, the
328 shortest distance from each Pinyon Jay location or control site to the polyline was computed
329 using the Near tool in ArcMap 10.5. This value was usually positive, but could be negative if a
330 Pinyon Jay location was in shrubland at a lower elevation than the polyline. Because the
331 Distance to Edge metric was not a FIA attribute, it was used only for data visualizations, not for
332 statistical analysis.

333

334 **Analysis** (level 2 heading)

Data (level 3 heading)

The complete data set used for all analyses and summarizations is provided in Table S1.

Ordination (level 3 heading)

To visualize how the habitat characteristics of Pinyon Jay locations for each behavior type overlapped with control sites, we performed region-specific ordinations on continuous FIA habitat attributes (Table 2) using nonmetric multidimensional scaling (NMDS). NMDS optimizes the non-parametric monotonic relationship among the data and the visualization space, providing the best low-dimensional representation [61-63]. All ordinated attributes were standardized with a z-transformation across all of the regions prior to the NMDS to promote optimization, which was defined as minimizing the stress value [63]. The NMDS was performed with the metaMDS function in the vegan package (v 2.5-5; [64]) in R (v3.5.1; [65]), using the Euclidian distance with two dimensions to force a visualizable representation. The maximum number of iterations was increased from the default of 200 to 10^{10} and the maximum number of runs was increased from 20 to 1,000 to achieve convergent solutions across repeated runs. Defaults for auto-transformation, extended dissimilarities, and weighted average scoring were turned off because they were not applicable to our transformed input data set.

Logistic Regression (level 3 heading)

To evaluate the effects of measured habitat attributes on Pinyon Jay occupancy by behavior type [42, 44, 66] we used logistic regressions for each behavior type that had sufficient data (i.e. caching, foraging, nesting). Roosting location were not analyzed because of low sample size (n=3). Because the ratio of Pinyon Jay locations to control sites was low, the odds ratio output from logistic regression can be treated as an approximation to the resource selection function [42, 67, 68]. However, it is important to recognize that the intercept value of the logistic regression does not estimate the overall use probability, as the Pinyon Jay locations were not randomly selected [42].

Initial evaluation of available predictor attributes (Table 2) consisted of plotting data pairs, evaluating correlations, and preliminary overall model fitting to ensure base-level convergence. Several attributes were eliminated from the analysis due to high correlations with other attributes or highly uneven distribution of values. Others were combined into a single attribute with principle components analysis (Table 2). All continuous attributes used for analysis (Table 2) were converted to metric units and z-transformed to facilitate model fitting and term comparison.

For the logistic regression models, we split the control sites assigned to each region (see above) according to behavior types in proportion to behavior frequency, rather than “reusing” control sites across multiple behavior types. Splitting was randomized and permuted 10,000 times to account for variation in the dataset splitting process. For each permutation, the control data were first split among the behaviors types within each region, then the data for a given behavior type were combined across the three regions and analyzed using a generalized (binomial) mixed model fit with the glmer function in the lme4 package (v1.1-19; [69]) in R (v3.5.1; [65]). The

permutations were then combined to integrate over the sample allocation variation and estimate overall terms. This was done by first discarding any permutations that did not converge due to unreasonable splits of control sites that could occur within the randomization. Then, for each permutation, a single set of fixed effects parameters were drawn from the multivariate normal distribution described by the glmer model fit using the rmvnorm function in the mvtnorm package (v1.0-8; [70]) in R (v3.5.1; [65]). By drawing from the distribution within each permutation, the full set of parameter values across permutations includes both parameter estimation uncertainty and data allocation uncertainty. To avoid distributional assumptions in evaluating the significance of the parameter estimates, we tested whether the estimate significantly differed from 0 by using a permutation-style two-tailed approach [71]. This involved using the empirical cumulative distribution function (calculated using the ecdf function in the stats package in base R (v3.5.1; [65])) to estimate the two tail probabilities with respect to 0, taking the smaller value as the focal tail, and doubling that tail's probability. We combined the estimates of the random effects (one for each successful permutation) across all of the permutations to generate the distribution of values for each random effects term across the uncertainty in control data allocation.

Within-sample classification accuracy of logistic regression models for each Pinyon Jay behavior type was averaged over all 10,000 permutations. Because the number of Pinyon Jay locations for each behavior type was limited, we did not withhold a subset of locations for external validation.

Data visualization (level 3 heading)

To aid in the interpreting statistical results and to highlight univariate patterns of interest, box plots were created for each continuous habitat attribute shown in Table 2, contrasting the distribution of attribute values for behavior-specific Pinyon Jay locations and control points.

Results (level 1 heading)

Data summary (level 2 heading)

Behavior-specific Pinyon Jay locations (n = 152) were obtained from 15 different flocks. Details about distribution of Pinyon Jay behavior type locations among study regions, apportionment of control sites, number of flocks per study area, and methods of data collection used within study areas are summarized in Table 3. Pinyon Jay data were not fully symmetrical among the study areas. Nesting was not recorded in Southern Idaho because the data collection period excluded the breeding season, and our data recording protocol in 2008-2009 for Eastern Nevada did not specify the foraging behavior type. Only caching locations were recorded in all three study areas, and only in the Central Nevada study area were all three major behavior types (caching, foraging, and nesting) recorded.

Table 3. Summary of Pinyon Jay locations after data processing, by behavior type and by region / study area. Shown parenthetically in 2nd - 5th columns are the number of control sites assigned to each unique combination of study region x behavior type within logistic regression models as described in above. The final column summarizes the number of different Pinyon Jay flocks from which data were collected in each study area, with the data collection methods used shown in brackets (T = telemetry surveys, with number of deployed radio tags in parentheses; O = observational surveys).

Study Region / Area	# Caching Sites	# Foraging Sites	# Nesting Sites	# Roosting Sites	# Flocks [Method of Study]
---------------------	-----------------	------------------	-----------------	------------------	----------------------------

Eastern Nevada	12 (106)	0 (0)	12 (106)	3 (N/A)	2 flocks [Flock #1 = T(6); Flock #2 = T(6)]
Southern Idaho	32 (51)	19 (30)	0 (0)	0 (N/A)	2 flocks [Flock #1 = O & T(6); Flock #2 = O & T(2)]
Central Nevada	26 (18)	22 (15)	28 (20)	0 (N/A)	11 flocks [All = O]

Ordination (level 2 heading)

The NMDS ordination found convergent solutions within 28, 388, and 20 runs for the Eastern Nevada, Southern Idaho, and Central Nevada regions, respectively (Fig 5). Each of the ordinations was well-correlated with true distances with stress values of 0.183, 0.127, and 0.161, respectively. Notably, for all three regions, Pinyon Jay locations formed distinct clusters in the NMDS plots that were easily distinguishable from the broader distribution of control sites (Fig 5). Degree of overlap among Pinyon Jay behavior type clusters varied somewhat across study regions, though nesting locations and caching locations were distinct in the two regions where both types of data were available. NMDS results suggested that the locations where Pinyon Jays occurred in the three study areas tended to be different than typical control sites within the corresponding study regions, and that Pinyon Jays tended to use different habitats for different behaviors.

Fig 5. NMDS ordination results for each region, by location type. Note that each region's ordination is independent and the axes cannot be compared among them.

Logistic regression (level 2 heading)

442
 443 Logistic regression models offer potential insights into which habitat covariates underpin the
 444 differentiation seen in the ordinations. We note, however, that within our logistic regressions,
 445 many predictors that had strong and statistically significant effects for a majority of iterations in
 446 the control site allocation process became non-significant when all iterations were combined.
 447 Because of this sensitivity to control site allocation, examination of data visualizations (next
 448 section) may assist in the interpretation of model results. Internal classification accuracy of
 449 behavior-specific Pinyon Jay locations versus control sites (averaged overall all iterations) was
 450 high (0.890 for the caching model, 0.863 for the foraging model, and 0.920 for the nesting
 451 model).

452
 453 Models indicated that Pinyon Jay caching locations were more likely to occur with lower slope,
 454 lower tree cover, increased woody debris, shorter distance to roads, and more disturbance than
 455 control sites (Table 4). There was also a strong effect of decreased elevation, though it was not
 456 significant when averaged across all control allocation iterations. None of the predictors for
 457 Pinyon Jay foraging locations were statistically significant when averaged across all iterations
 458 due to high standard errors, but the largest effect sizes were noted for lower slope, less grass and
 459 forb cover, increased woody debris, and shorter distance to roads compared with control sites
 460 (Table 4). Pinyon Jay nesting locations were more likely to occur at lower elevations with
 461 decreased forb cover and increased woody debris compared to control sites (Table 4). Across all
 462 three Pinyon Jay behavior types, occupancy probability generally increased with lower elevation,
 463 lower slope, lower forb cover, shorter distance to roads, and increased woody debris compared to

control sites (Table 4). Lower tree cover was a marginally significant predictor for caching locations but was not a significant predictor for foraging and nesting locations.

Table 4. Logistic regression results over all control site allocation iterations for the caching, foraging, and nesting location analyses. Fixed effects are reported as mean and standard error of the estimate and the two-tailed p value; random effects are reported as mean and standard error of the estimate and the probability (fraction of permutations) that the term was equal to 0. Bolded terms are significant at $\alpha = 0.05$.

CACHING LOCATIONS			
	Mean	Standard Error	p
Fixed Effect			
Intercept	-3.687	1.162	0.001
Elevation	-0.405	0.585	0.481
Slope	-2.066	0.557	< 0.0001
Stand Age	-0.177	0.432	0.654
Tree Cover	-0.913	0.553	0.075
Shrub Cover	0.251	0.349	0.445
Forb Cover	-0.021	0.3	0.904
Grass Cover	0.377	0.439	0.385
Woody Debris	1.235	0.818	0.010
Distance to Road	-2.712	0.789	< 0.0001
Disturbance	1.971	0.951	0.022
Random Effect			
Eastern Nevada: Intercept	1.838	0.413	0.0001
Southern Idaho: Intercept	-1.320	0.301	0.0001
Central Nevada: Intercept	-0.522	0.288	0.0001
FORAGING LOCATIONS			
Fixed Effect			
Intercept	-5.486	7.418	0.044
Elevation	-1.566	2.790	0.317
Slope	-4.184	6.299	0.058
Stand Age	-2.013	3.641	0.236
Tree Cover	1.931	3.079	0.200
Shrub Cover	1.380	2.479	0.311
Forb Cover	-9.236	14.819	0.080
Grass Cover	-4.970	8.384	0.113
Wood Debris	4.419	9.650	0.053
Distance to Road	-3.326	4.848	0.063
Disturbance	-1.700	75.272	0.555
Random Effect			

Eastern Nevada: Intercept	0.686	1.076	0.290
Southern Idaho: Intercept	N/A	N/A	N/A
Central Nevada: Intercept	-0.707	1.327	0.290
NESTING LOCATIONS			
Fixed Effect			
Intercept	-4.723	2.623	0.021
Elevation	-2.445	1.600	0.004
Slope	-1.249	1.090	0.154
Stand Age	0.528	1.016	0.564
Tree Cover	0.500	1.030	0.570
Shrub Cover	0.261	0.848	0.743
Forb Cover	-7.110	4.247	0.022
Grass Cover	0.123	1.256	0.983
Woody Debris	2.156	1.266	0.003
Distance to Road	-1.118	1.291	0.289
Disturbance	2.606	2.406	0.115
Random Effect			
Eastern Nevada: Intercept	2.116	0.903	0
Southern Idaho: Intercept	-2.088	0.904	0
Central Nevada: Intercept	NA	NA	NA

The random effects components of the three models were all estimated to be non-zero, although the foraging model estimated a zero-value in 29% of allocation iterations, whereas the caching model estimated a zero-value in < 1% of iterations. The additional complexity introduced by a singularity/boundary condition (a zero-value random effect) within many foraging model iterations may have contributed to the variability (and thus reduced significance) seen in the parameter estimates.

Data visualizations (level 2 heading)

Fig 6 shows box plots for all FIA attributes included in logistic regression models combined across all study regions, and Fig 7 shows box plots for two additional habitat attributes (the FIA Stand Density Index attribute and the non-FIA Distance to Edge attribute) that were not included in logistic regressions, but which showed patterns of interest with regard to Pinyon Jay occupancy. Notable patterns in these box plots are as follows:

- 1) Across all study regions, locations used by Pinyon Jays appeared to be a distinct subset available woodland habitat with regard to many habitat attributes. This is consistent with the patterns observed in the NMDS ordination (Fig 5). Additionally, locations used by Pinyon Jays for different behaviors appeared to vary with regard to multiple habitat attributes.
- 2) Compared to available habitat, caching locations were concentrated in lower elevation, lower-slope areas with young woodland stands, low tree cover, and high shrub, forb, and grass cover. The Stand Density Index was typically very low for caching locations, which were also highly concentrated near the woodland-shrubland ecotone (i.e. low Distance to Edge values). In fact, many caching locations occurred in the pure shrubland habitat located down-slope from the woodland-shrubland ecotone (note that these pure shrublands were not represented in the sample of FIA control sites).
- 3) Foraging locations were also concentrated in lower elevation and lower slope areas, but were somewhat higher and steeper than caching locations, with somewhat higher stand densities. Foraging locations had more woody debris than typical control sites, but stand age, tree cover, forb cover, and grass cover were comparable to control sites. Shrub cover in foraging locations was highly variable, but tended to be higher than in control sites.

Foraging locations were close to the woodland-shrubland ecotone, with lower stand density indices, though to a less extent than caching locations

- 4) Nesting locations also tended to occur at lower elevations and lower slopes, but to a lesser degree than foraging and caching locations. Stand age and shrub cover were comparable between nesting locations and control sites, but nesting locations had higher tree cover, lower forb cover, and more woody debris. Stand density was somewhat lower for nesting locations than for control sites, but with considerable overlap. Nesting locations were also concentrated close to the woodland-shrubland ecotone, but to a lesser degree than foraging locations.
- 5) There was a distinct pattern of increasing elevation and slope, increasing distance from edge, and increasing stand density moving from caching locations to foraging locations to nesting locations, but all three behavior types had lower values for these attributes than typical control sites.

Fig 6: Box plots for all continuous attributes used in logistic regression models, as described in Table 2. Y-axis codes are Co = control sites, N = Pinyon Jay nesting locations, F = Pinyon Jay foraging locations, and Ca = Pinyon Jay caching locations. For better visual clarity, the extreme high range of observed Forb Cover values (15 – 25%) is truncated, omitting a small number of Co sites and Ca locations.

Fig 7: Box plots for two habitat attributes not used in logistic regression models, as described in Table 2. Y-axis codes are Co = control sites, N = Pinyon Jay nesting locations, F = Pinyon Jay foraging locations, and Ca = Pinyon Jay caching locations.

For better visual clarity, the extreme high range of observed Distance to Edge values (6,000 – 12,000 m) values are truncated, omitting a small number of Co sites.

Discussion (level 1 heading)

Main findings and significance (level 2 heading)

This study offers the first systematic description of Pinyon Jay occupancy patterns and behavior-specific habitat characteristics in the Great Basin. The combination of ordinations, logistic regressions, and data visualizations presented in this study suggest that Pinyon Jays use pinyon-juniper woodlands selectively. All Pinyon Jay locations in our study areas were concentrated in or near lower-elevation, flatter woodlands and were less common in higher-elevation, steeper woodlands. Additionally, the areas used for different Pinyon Jay behaviors appear to have distinctive (but overlapping) habitat profiles, with caching, foraging, and nesting arrayed sequentially along gradients of increasing slope, elevation, and stand density. Similar patterns of Pinyon Jay caching and nesting activities partitioned along elevation and stocking gradients were observed by Johnson et al. [23] within a pinyon-juniper (*P. edulis* / *Juniperus spp.* association) woodland system in New Mexico, suggesting that habitat partitioning by behavior along an elevational gradient could be present across a broader physiographic range than our study regions.

Pinyon Jay caching locations were concentrated in open woodland stands with high shrub and grass cover, which are similar to the Phase I (early successional) pinyon-juniper woodlands defined in the classification scheme by Miller et al. [33], and sometimes occurred in pure shrublands. This occurrence pattern could have a mutualistic explanation [3]. From the pinyon pine perspective, seedlings likely experience less competition from established trees in more open areas, and seeds placed next to shrubs, rocks, or woody debris in otherwise open areas may benefit from favorable microsite conditions created and maintained by those features [72, 73]. From the Pinyon Jay perspective, pinyon pine seeds cached away from their source of origin may be less likely to be discovered and eaten by small mammals that specialize on pinyon seeds [73, 74].

Pinyon Jay foraging locations generally occurred in older (though still relatively young) stands than caching locations, across a wide range of tree cover values. Given that foraging behavior as defined in our study encompassed the gathering of diverse food items from trees, shrubs, and ground, areas with these characteristics may offer a beneficial combination of pinyon pines in their most productive seed-bearing years [75, 76] interspersed with areas where abundant insect prey is available due to higher shrub or ground cover [77]. The habitat characteristics of Pinyon Jay foraging locations correspond to a mosaic of Phase I and Phase II pinyon-juniper successional stages [33].

Pinyon Jay nesting locations tended to be concentrated in areas with higher tree cover and more woody debris, presumably because of the concealment they offer [23, 78]. However, like caching and foraging locations, nesting locations were concentrated in lower elevation, lower slope areas,

and steeper, higher sites that otherwise offered good concealment for nesting locations appeared to be avoided. Pinyon Jay nesting locations correspond best with the denser portion of the Phase II class of pinyon-juniper woodlands, but may also include some Phase III areas [33].

All Pinyon Jay locations, regardless of behavior type, were concentrated in lower-elevation woodlands (most likely a mix of Phase I and Phase II classes) near the woodland-shrubland ecotone. This could occur because of the longer snow-free season of lower elevations, or the presence of a mosaic of desirable habitat characteristics needed to support different behavior types. Phase I and Phase II woodlands are relatively common at lower elevations where Pinyon Jay locations are concentrated, whereas the proportion of Phase III woodlands tends to increase with increasing elevations based on our field observations.

In addition to providing important information about Pinyon Jay occupancy patterns and habitat use, these findings are potentially significant for vegetation management planning and implementation because Pinyon Jays in our three study areas appear to prefer the same lower-elevation, relatively-open woodlands where most woodland removal management is performed [79, 80]. These vegetation management projects, which are most often conducted to create or improve habitat for Greater Sage-Grouse [81] (but see Miller et al. [22] and Somershoe et al. [15] for others reasons for woodland treatments), have resulted in the removal of an estimated 45,000 ha of pinyon-juniper woodland in the Great Basin portions of Nevada, Utah, and Idaho over the last eight years alone (Witt, unpublished USFS data). The pace of this activity appears to be steady or accelerating, and may therefore be or become a significant factor affecting Pinyon Jay populations, either negatively or positively. Information from other regions suggests that

woodland treatments can have unintended effects on Pinyon Jays. For example, Johnson et al. [82] found that a fuels treatment within pinyon-juniper woodlands in northern New Mexico that reduced tree density by almost 90 percent prompted the local Pinyon Jay flock to avoid the treated area altogether. To date, however, almost no direct monitoring has been conducted in the Great Basin to determine if and how Pinyon Jay flocks respond to vegetation management projects that occur within or close to their home ranges.

Interpretational considerations (level 2 heading)

The findings presented in this study should be interpreted with the following considerations in mind:

- 1) Pinyon Jay data were collected at the three distinct study areas that collectively encompassed 175,630 ha. Although this was a large area, it represents only a small portion of the Great Basin region that we would like to characterize.
- 2) This study combined data from three distinct projects that covered different years and seasons. This is most immediately relevant to interpreting the foraging model, given that seasonal variation in foraging behaviors and locations are plausible.
- 3) We equated “potential habitat” for Pinyon Jays to all pinyon-juniper woodlands lying within 200 km of any Pinyon Jay study area, as sampled by FIA plots. Although inferences from this study can cautiously be extended to the pinyon-juniper woodlands beyond our immediate Pinyon Jay study areas (subject to confirmation in future studies; see Johnson and Sadoti [78] for caveats about model transferability to similar systems) it does not extend to other forest types or regions where Pinyon Jays occur.

- 4) Our analysis does not distinguish between flocks within a study area, and does not allow us to draw inferences about inter-flock variability with regard to behavior-specific occupancy patterns.
- 5) The iterative allocation of control sites among Pinyon Jay behavior types in logistic regression modeling was necessary, but it could have diluted the statistical significance of some important predictors of occupancy. Future modeling efforts where the spatial extents and sample sizes of Pinyon Jay data and control data are better matched should reduce this issue. Models created from data sets with a large sample of Pinyon Jay locations will also allow data to be withheld from the model building process and used for external validation.
- 6) Interpreting the relationship between Pinyon Jay occurrence and distance to road is complex and potentially non-causal. Roads density is typically higher in lower-elevation, flatter areas, and Pinyon Jays prefer these areas for reasons completely unrelated to the proximity of roads. However, one author has suggested the vegetation typically present alongside graded, unpaved roads may provide valuable foraging opportunities for Pinyon Jays [83].
- 7) The Distance to Edge attribute used for data visualizations shows clear contrasts across behavior-specific Pinyon Jay locations and control sites, but it needs to be further investigated using more formal methods for delineating ecotones. Additionally, the patterns seen in our data (Fig 7) were enhanced because control sites by definition excluded the pure or near-pure shrublands that comprised a significant proportion of Pinyon Jay caching locations, and a smaller proportion of foraging locations.

Currently, we are analyzing a separate Pinyon Jay data set derived from a long-term statewide bird monitoring program in Nevada. Because these data were obtained from a broader-scale fully-randomized sampling design and used a fully standardized survey protocol, they should provide a useful independent characterization of Pinyon Jay occupancy patterns in the Great Basin.

Conclusions and recommendations (level 2 heading)

Pinyon Jay populations have been declining precipitously for at least the last half-century, while the pinyon-juniper woodlands that they inhabit in the Great Basin are thought by many to have been expanding at unprecedented rates [84-87]. Given this apparent paradox, identifying the reasons for Pinyon Jay declines is critical for defining constructive conservation actions that ensure the species' long-term viability [3, 15]. This urgency is amplified by the widespread and potentially accelerating woodland management activities in the Great Basin that prioritize creation or preservation of shrublands without a clear understanding of their impacts on Pinyon Jays and other woodland-associated birds. Progress towards a more inclusive management paradigm can be achieved through: a) better knowledge of Pinyon Jay ecology and habitat requirements, b) monitoring of management impacts, c) better understanding of the ecology and dynamics of the woodlands that comprise Pinyon Jay habitat, and d) integration of knowledge obtained from these three areas into existing vegetation management protocols and guidance.

A fundamental need is for more robust, spatially-extensive data characterizing Pinyon Jay occupancy patterns and habitat use as a function of region, season, and behavior type, both

661 across and within the individual flock level. To maximize their value, these data would ideally be
662 gathered using a standardized survey protocol. Pinyon Jays, however, present multiple
663 challenges to the field biologist, study designer, and data analyst [15], and approaches suitable
664 for typical passerine birds may be suboptimal for Pinyon Jays for several reasons. Unlike species
665 where a single breeding pair occupies a clearly-delineated territory at all times during the
666 breeding season, Pinyon Jays occupy (and presumably select) habitat at the flock and subflock
667 level. Pinyon Jays are also year-round residents, and protecting breeding habitat alone may be
668 insufficient for effective conservation. The Pinyon Jay's pattern of habitat use, which involves
669 temporal flock movements across a relatively large home range to accommodate different
670 behaviors and take advantage of seasonally-varying food resources, has potentially profound
671 effects on both the detection properties of a given survey protocol and the ecologically-legitimate
672 interpretation of those detection patterns. As a simple example, a Pinyon Jay flock may be
673 frequently absent from a critically-important subset of its home range, either during portions of
674 the day, or during entire seasons. Similarly, roaming flocks may frequently fly over or loaf in
675 areas of the home range that are not critical to home range quality or viability. To provide
676 accurate and actionable information about the habitat requirements of Pinyon Jays, survey
677 protocols and research study designs need to account for these realities appropriately, operate at
678 scales that reflect actual Pinyon Jay habitat selection patterns, and be guided by a sampling
679 framework that produces well-balanced data suitable for presence / absence modeling. The
680 multi-agency Pinyon Jay Working Group [15] is currently actively investigating options for
681 standardized Pinyon Jay survey protocols. The same group's Pinyon Jay Conservation Strategy
682 [15] also noted that some of the information needed to better characterize Pinyon Jay habitat
683 requirements could be obtained by systematically monitoring Pinyon Jay responses to vegetation

management activities, especially in situations where their pre-treatment presence has been confirmed by baseline or clearance surveys.

With regard to pinyon-juniper woodlands, their structural attributes and other characteristics that might be limiting to Pinyon Jay populations need to be further studied. We suggest that it may be especially important to identify the correlates or profiles of tree stands and landscapes that exhibit a predictable and/or abundant pinyon pine mast [22]. Our preliminary review of FIA data collected in Nevada between 2006-2015 suggests that woodlands matching the structural characteristics Pinyon Jays used for foraging in this study are 5-7 times less extensive than nesting or caching habitat in Nevada (unpublished USFS data). Presence of reliably productive stands within the home range could be especially important to Pinyon Jays during years of more generally depressed pine mast production. Given evidence of reduced mast production in pinyon pine [88], and associated changes in habitat use by Pinyon Jays [24] in some areas affected by climate change, it might be critical to long-term Pinyon Jay conservation to systematically investigate the quality and quantity of good foraging areas.

In addition, more research is needed to better clarify the degree to which woodland expansion and infill is part of a historically “normal” dynamic, versus a problematic departure condition. In the absence of this more holistic understanding, colonization of shrublands by trees tends to be widely regarded as “invasive”, even though at least some of these recently colonized areas appear to be an important component of Pinyon Jay habitat in the Great Basin. Achieving this broader perspective may be a necessary prerequisite to successfully accommodating the needs of

Greater Sage-Grouse, Pinyon Jays, and other sensitive shrubland and woodland bird species within the overall framework of pinyon-juniper woodland management.

Ultimately, accruing information about Pinyon Jays must be incorporated into woodland management paradigms and protocols in the Great Basin (see Ricca et al. [89] for an example) in ways that accommodate both previously-identified and newly-emerging goals within the context of healthy ecosystem function [90]. For the present, Somershoe et al. [15] provide guidance for managers seeking to incorporate Pinyon Jay conservation measures into their vegetation management projects based the extent of current knowledge.

Acknowledgments (level 1 heading)

Field work was conducted by Gustavo Gonzalez, Michael Maples, and John B. Free (in Eastern Nevada), by Murrelet Halterman, Natasha Peters, Larry Teske, and Mercer Owen (in Southern Idaho, and by Mercer Owen and Sue Brunner (in Central Nevada). Their efforts, which were supplemented by the authors, are much appreciated. Statistical analysis was performed by Juniper Simonas of Dapper Stats (www.dapperstats.com).

Wallace Keck, Superintendent of City of Rocks National Reserve and Park Manager for Castle Rock State Park, provided us with extensive information about local Pinyon Jay populations, generously allowed us use of park facilities, with further assistance from Trenton Durfee. Many local landowners permitted us to access to their property, notable among them LeAnn and Kim

Draper, whose property functioned as a capture location and staging area for telemetry efforts. At the Bureau of Land Management, Sandra Brewer and John Wilson provided invaluable assistance and support in Central Nevada. We thank all of these individuals.

References (level 1 heading)

1. Burleigh TD. Birds of Idaho. Caldwell: Caxton Printers Ltd.; 1972.
2. Grenfell Jr. WE, Laudenslayer Jr. WF. The distribution of California birds. San Francisco, CA: Calif. Wildlife/Habitat Relationships Program, Publ. No. 4; California Department of Fish and Game Bulletin, Sacramento, and U.S. Forest Service. 1983.
3. Johnson K, Balda RP. Pinyon Jay (*Gymnorhinus cyanocephalus*). 2020. In: Rodewald PG, Keeny BK, editors. Birds of the World. Version 2.0. NY: Cornell Lab of Ornithology. Available from <https://birdsoftheworld.org> Subscription required.
4. Sauer JR, Niven DK, Hines JE, Ziolkowski, Jr. DJ, Pardieck KL, Fallon JE. The North American Breeding Bird Survey [Internet]. Laurel, MD: USGS Patuxent Wildlife Research Center; 2017. Results and Analysis 1966-2015. Version 2.07. Available from: <https://www.mbr-pwrc.usgs.gov/bbs/>

5. Little Jr. EL. Atlas of United States trees. Volume 1. Conifers and important hardwoods. Washington, DC: U.S. Department of Agriculture, Forest Service; 1971. Misc. Publ. 1146. p. 320
6. Marzluff, JM, Balda RP. The Pinyon Jay: Behavioral Ecology of a Colonial and Cooperative Corvid. London: Poyser; 1992.
7. Balda RP, Bateman GC. Flocking and annual cycle of the Pinyon Jay, *Gymnorhinus cyanocephalus*. *The Condor*. 1971;73: 278-302.
8. Bateman GC, Balda RP. Growth, development, and food habits of young Pinon Jays. *Auk*. 1973;90(1): 39-61.
9. NABCI Bird Conservation Regions [Internet]. Canada: Bird Studies Canada on behalf of the North American Bird Conservation Initiative. [cited 2017 Oct 3]. Available from: <http://www.birdscanada.org/research/gislab/index.jsp?targetpg=bcr>
10. Boone JD, Ammon E, Johnson K. Long-term declines in the Pinyon Jay and management implications for Pinyon-Juniper woodlands. In: Shuford WD, Gill Jr. RE, Handel CM, editors. *Trends and Traditions: Avifaunal Change in Western North America*. Studies of Western Birds 3. Camarillo, CA: Western Field Ornithologists; 2019. p 190-197. Available from: DOI: 10.21199/SWB3.10

- 773 11. Rosenberg KV, Kennedy JA, Dettmers R, Ford RP, Reynolds D, Alexander JD, et al.
774 Partners in Flight Landbird Conservation Plan: 2016 revision of Canada and continental United
775 States. Partners in Flight Science Committee; 2016.
776
- 777 12. U.S. Bureau of Land Management [Internet]. Updated Bureau of Land Management (BLM)
778 Sensitive Species List for Nevada; 2017. Available from: <https://blm.gov/policy/nv-im-2018-003>
779
- 780 13. BirdLife International [Internet]. The IUCN Red List of Threatened Species, *Gymnorhinus*
781 *cyanocephalus* (amended version of 2016 assessment); 2017. Available from:
782 <https://www.iucnredlist.org/species/22705608/110431877>
783
- 784 14. U.S. Fish and Wildlife Service [Internet]. Birds of Conservation Concern 2019. Falls Church,
785 VA: US Dept of the Interior, Fish and Wildlife Service; 2019. Available from:
786 <http://www.fws.gov/birds/management/managed-species/birds-of-conservation-concern.php>
787
- 788 15. Somershoe, SG, Ammon E, Boone JD, Johnson K, Darr M, Witt C, et al. Conservation
789 Strategy for the Pinyon Jay (*Gymnorhinus cyanocephalus*). Denver, CO: Partners in Flight
790 Western Working Group and U. S. Fish and Wildlife Service; 2020. Available from:
791 [https://partnersinflight.org/wp-content/uploads/2019/10/Conservation-Strategy-for-Pinyon-](https://partnersinflight.org/wp-content/uploads/2019/10/Conservation-Strategy-for-Pinyon-Jay_Version-1_February-2020_LowRes.pdf)
792 [Jay_Version-1_February-2020_LowRes.pdf](https://partnersinflight.org/wp-content/uploads/2019/10/Conservation-Strategy-for-Pinyon-Jay_Version-1_February-2020_LowRes.pdf).
793
- 794 16. Marzluff, JM, Balda RP. The advantages of, and constraint forcing, mate fidelity in pinyon
795 jays. Auk.1988;105(2): 286-295.

796

797 17. Marzluff, JM, Balda RP. Resource and climatic variability: influences on sociality of two
798 southwestern corvids. In: Slobodchikoff CN, editor. The ecology of social behavior. [Publication
799 location unknown]: Academic Press; 1988. p 255-283.

800

801 18. Gottfried GJ, Severson KE. Managing pinyon-juniper woodlands. *Rangelands*. 1994;16:234-
802 236.

803

804 19. Miller RF, Tausch RJ, McArthur ED, Johnson DD, Sanderson SC. Age structure and
805 expansion of piñon-juniper woodlands: a regional perspective in the Intermountain West. Ft.
806 Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station;
807 2008. RMRS-RP-69.

808

809 20. Morris LR, Rowe RJ. Historical land use and altered habitats in the Great Basin. *Journal of*
810 *Mammalogy*. 2014;95:1144-1156.

811

812 21. Kramer DW, Sorensen GE, Taylor CA, Cox RD, Gipson PS, Cain III JW. Ungulate
813 exclusion, conifer thinning, and mule deer forage in northeastern New Mexico. *Journal of Arid*
814 *Environments*. 2015;113:29–34.

815

816 22. Miller RF, Chambers JC, Evers L, Williams CJ, Snyder KA, Roundy BA, et. al. The ecology,
817 history, ecohydrology, and management of pinyon and juniper woodlands in the Great Basin and
818 Northern Colorado Plateau of the Western United States. Fort Collins, CO: U. S. Department of

Agriculture, Forest Service, Rocky Mountain Research Station; 2019. Gen tech report. RMRS-GTR-403.

23. Johnson K, Neville TB, Smith JW, Horner MW. Home range- and colony-scale habitat models for Pinyon Jays in piñon-juniper woodlands of New Mexico, USA. *Avian Conservation and Ecology*. 2016;11:6.

24. Johnson K, Sadoti G, Smith J. Weather-induced declines in Piñon tree condition and response of a declining bird species. *Journal of Arid Environments*. 2017;146: 1-9.

25. Goeking SA, Shaw JD, Witt C, Thompson MT, Werstak Jr. CE, Amacher, MC, et. al. New Mexico's forest resources, 2008-2012. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station; 2014. Resource bull. RMRS-RB-18. 144 p.

26. Menlove J, Shaw JD, Witt C, Werstak Jr CE, DeRose RJ, Goeking, SA, et. al. Nevada's forest resources, 2004-2013. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station; 2016. Resource bull. RMRS-RB-22. 167 p.

27. Werstak Jr. CE, Shaw JD, Goeking, SA, Witt C, Menlove J, Thompson MT, et. al. Utah's forest resources, 2003-2012. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station; 2016. Resource bull. RMRS-RB-20. 159 p.

28. Thompson MT, Shaw JD, Witt C, Werstak Jr. CE, Amacher MC, Goeking SA, et. al.
Colorado's forest resources, 2004-2013. Fort Collins, CO: U.S. Department of Agriculture,
Forest Service, Rocky Mountain Research Station; 2017. Resource bull. RMRS-RB-23. 136 p.
29. Betancourt JL. Paleoecology of pinyon-juniper woodlands: summary. In: Everett RL, editor.
Proceedings- Pinyon-juniper conference. USDA Forest Service; 1987. Gen tech report. INT-215.
p 129-139.
30. Thompson RS. Late Quaternary vegetation and climate in the Great Basin. In: Betancourt JL,
et al., editors. Packrat middens: The last 40,000 years of biotic change. Tucson, AZ: University
of Arizona Press; 1990. p. 200-239
31. Cottom WP, Stewart G. Plant succession as a result of grazing and of meadow
desiccation by erosion since settlement in 1862. Journal of Forestry. 1940;38: 613-626.
32. Tausch RJ, West NE, Nabi AA. Tree age and dominance patterns in Great Basin pinyon-
juniper woodlands. Journal of Range Management. 1981;34:259-264.
33. Miller RF, Tausch RJ, MacArthur ED, Johnson DJ, Sanderson SC. Age structure and
expansion of piñon-juniper woodlands: a regional perspective in the Intermountain West. Fort
Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station;
2008. RMRS-RP-69.

34. Morris LR, Rowe RJ. Historical land use and altered habitats in the Great Basin. *Journal of Mammalogy*. 2014;95: 1144-1156.
35. Young JA, Budy, JD. Historical use of Nevada's pinyon-juniper woodlands. *Forest and Conservation History*. 1979;23(3):112-121.
36. Ko DW, Sparrow AD, Weisberg PJ. Land-use legacy of historical tree harvesting for charcoal production in a semi-arid woodland. *Forest Ecology and Management* [Internet]. 2011;261(7):1238-1292. Available from: doi:10.1016/j.foreco.2011.01.007
37. Natural Resources Conservation Service. Sage Grouse Initiative; Success on the range. Washington D.C.: U.S. Department of Agriculture, Natural Resources Conservation Service; 2015.
38. Chambers JC, Beck JL, Bradford JB, Bybee J, Campbell S, Carlson J, et al. Science framework for conservation and restoration of the sagebrush biome: Linking the Department of the Interior's Integrated Rangeland Fire Management Strategy to long-term strategic conservation actions. Part 1. Science basis and applications. Fort Collins, CO: U.S Department of Agriculture, Forest Service, Rocky Mountain Research Station; 2017. Gen tech rep. RMRS-GTR-360. 213 pp.

39. Baruch-Mordo S, Evans JS, Severson JP, Naugle DE, Maestas JD, Kiesecker JM, et al. Saving sage-grouse from the trees: A proactive solution to reducing a key threat to a candidate species. *Biological Conservation*. 2013;167:233–241.
40. Donnelly JP, Tack JD, Doherty KE, Naugle DE, Allred BW, Dreitz VJ. Extending conifer removal and landscape protection strategies from sage-grouse to songbirds, a range-wide assessment. *Rangeland Ecology and Management*. 2017;70:95-105.
41. Lewallen S, Courtright P. Epidemiology in practice: case-control studies. *Community Eye Health*. 1998;11:57-58.
42. Keating KA, Cherry S. Use and interpretation of logistic regression in habitat-selection studies. *The Journal of Wildlife Management*. 2004;68:774-789.
43. Soulard CE. Central Basin and Range Ecoregion. In: Sleeter BM, Wilson TS, Acevedo W, editors. *Status and Trends of Land Change in the Western United States – 1973 – 2000*. U.S. Geological Survey; 2012. Professional Paper 1794-A. Available from: <https://pubs.usgs.gov/pp.1794/a/>
44. Manly BF, McDonald L, Thomas D, McDonald TL, Erickson WP. *Resource Selection by Animals*. 2nd edition. New York: Springer; 2002.

45. Cole KL, Arundel ST. Modeling the climatic requirements for southwestern plant species. In: Starratt S, Bloomquist N, editors. Proceedings of the Twenty-First Annual Pacific Climate Workshop. State of California Department of Water Resources [Internet]. 2007 [Cited 2020 Mar 25]. Available from: <http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.606.6097>.
46. Gillespie AJR. Rationale for a National Annual Forest Inventory Program. *Journal of Forestry*. 1999;97(12): 16-20.
47. Smith WB. Forest Inventory and Analysis: A National Inventory and Monitoring Program. *Environmental Pollution*. 2002;116(1): S233-S242.
48. Zielinski WJ, Truex RL, Dunk JR, Gaman T. Using Forest Inventory data to assess fisher resting habitat suitability in California. *Ecological Applications*. 2006;16 (3):1010-1025.
49. Zarnetske PL, Edwards TC, Moisen GG. Modeling forest bird species' likelihood of occurrence in Utah with Forest Inventory and Analysis and Landfire map products and ecologically based pseudo-absence points. *Proceedings of the Seventh Annual Forest Inventory and Analysis Symposium*; 2005 Oct 3-4; Portland, Maine. Department of Agriculture, Forest Service; 2007. WO-GTR-77. p. 291-306
50. Witt C. Quantification of Lewis's Woodpecker habitat using Forest Inventory and Analysis data. *Proceedings of 2008 Forest Inventory and Analysis Symposium*; 2008 Oct 21-23; Park

- City, Utah. United States Forest Service Rocky Mountain Research Station; 2009. RMRS-P-56CD.
51. Witt C. Characteristics of aspen infected with heartrot: Implications for cavity-nesting birds. Forest Ecology and Management. 2009;260 (6): 101-1016.
52. Witt C, Patterson PL. Big Sagebrush in Pinyon-juniper Woodlands: Using Forest Inventory and Analysis Data as a Management Tool for Quantifying and Monitoring Mule Deer Habitat. Natural Resources and Environmental Issues. 2011;16(27). Available from: <http://digitalcommons.usu.edu/nrei/vol16/iss1/27>
53. Bechtold WA, Patterson PL. The enhanced Forest Inventory and Analysis program-national sampling design and estimation procedures. Department of Agriculture Forest Service, Southern Research Station; 2005. SRS-GTR-80.
54. United States Department of Agriculture. Interior West Forest Inventory and Analysis field procedures version 6.1 [Internet]. Ogden, UT: United States Forest Service, Rocky Mountain Research Station; 2012. Available from: https://www.fia.fs.fed.us/library/field-guides-methods-proc/docs/2014/Core%20FIA%20field%20guide_6-1.pdf
55. United States Department of Agriculture, Forest Service. Interior West Forest inventory and Analysis Region 4 All Condition Inventory Supplemental Field Guide [Internet]. Odgen, UT: United States Forest Service, Rocky Mountain Research Station; 2012. Available from:

https://www.fs.fed.us/rm/ogden/datacollection/pdf/Region_4_All_Condition_Inventory_Supplemental_Field_Guide.pdf

56. United States Department of Agriculture, Forest Service. Interior West Forest Inventory and Analysis field procedures version 4.0 [Internet]. Ogden, UT: United States Forest Service, Rocky Mountain Research Station; 2007. Available from: https://www.fia.fs.fed.us/library/field-guides-methods-proc/docs/core_ver_4-0_10_2007_p2.pdf

57. United States Department of Agriculture, Forest Service. Interior West Forest Inventory and Analysis field procedures version 5.0 [Internet]. Ogden, UT: United States Forest Service, Rocky Mountain Research Station; 2010. Available from: <https://www.fia.fs.fed.us/library/field-guides-methods-proc/docs/Complete%20FG%20Document/NRS%20FG%205.0-Oct%202010-Complete%20Document.pdf>

58. United States Department of Agriculture, Forest Service. Interior West Forest Inventory and Analysis field procedures version 5.1 [Internet]. Ogden, UT: United States Forest Service, Rocky Mountain Research Station; 2011. Available from: https://www.fia.fs.fed.us/library/field-guides-methods-proc/docs/Complete%20FG%20Document/core_ver_5-1_10_2011.pdf

59. Burrill EA, Wilson AM, Turner JA, Pugh SA, Menlove J, Christiansen G, et al. The Forest Inventory and Analysis Database: database description and user guide version 8.0 for Phase 2 [Internet]. U.S. Department of Agriculture, Forest Service; 2018. 946 p. Available from:

<https://www.fia.fs.fed.us/library/database->

documentation/current/ver80/FIADB%20User%20Guide%20P2_8-0.pdf

60. Alexander RR. Forest Vegetation in the Rocky Mountain and Intermountain Regions: Habitat Types and Community Types. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station; 1988. Gen tech report. RM-162.

61. Kruskal JB. 1964. Multidimensional scaling by optimizing goodness-of-fit to a nonmetric hypothesis. *Psychometrika*. 1964;29: 1-28.

62. Kruskal JB. Nonmetric multidimensional scaling: a numerical method. *Psychometrika*. 1964;29:115-129.

63. Kruskal JB, Wish M. Multidimensional Scaling. Newbury Park, CA: Sage Publications; 1978.

64. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. Vegan: community ecology package. R package version 2.5-5 [Internet]. 2019. Available from: <https://CRAN.R-project.org/package=vegan>

65. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing [Internet]. 2018. Available from: <https://www.R-project.org/>

997

998 66. Newlon KR, Saab VA. Nest-site selection and nest survival of Lewis's Woodpecker in aspen
 999 riparian woodlands. *The Condor*. 2011;113:183-193.

1000

1001 67. Hosmer DW, Lemeshow S. Applied logistic regression. New York: Wiley; 1989.

1002

1003 68. Compton BW, Rhymer JM, McCollough M. Habitat selection by wood turtles (*Clemmys*
 1004 *insculpta*): an application of paired logistic regression. *Ecology*. 2002;83: 833-843.

1005

1006 69. Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4.
 1007 *Journal of Statistical Software*. 2004;67:1-48.

1008

1009 70. Genz A, Bretz F, Miwa T, Mi X, Leisch F, Scheipl F, et al. mvtnorm: multivariate normal
 1010 and t distributions. R package version 1.0-8 [Internet]. 2018. Available from: [http://CRAN.R-](http://CRAN.R-project.org/package=mvtnorm)
 1011 [project.org/package=mvtnorm](http://CRAN.R-project.org/package=mvtnorm)

1012

1013 71. Gotelli NJ, Ellison AM. A Primer of Ecological Statistics. Sunderland, MA: Sinauer
 1014 Associates; 2004.

1015

1016 72. Drivas EP, Everett RL. Water relations characteristics of competing singleleaf pinyon
 1017 seedlings and sagebrush nurse plants. *Forest Ecology and Management*. 1988;23: 27-37.

1018

1019 73. Hollander JL, Vander Wall SB. Effectiveness of six species of rodents as

1020 dispersers of singleleaf piñon pine (*Pinus monophylla*). *Oecologia*. 2004;138: 57-65.

1021

1022 74. Vander Wall SB. Food Hoarding in Animals. Chicago, IL: University of Chicago Press;

1023 1990.

1024

1025 75. Evans RA. Management of pinyon-juniper woodlands. Ogden, UT: U. S. Department of

1026 Agriculture, Forest Service, Intermountain Research Station; 1988. Gen tech report. INT-249.

1027

1028 76. Meeuwig RO, Budy JD, Everett RL. *Pinus monophylla* Torr. & Frem., singleleaf pinyon. In:

1029 Burns RM, Honkala, technical coordinators. Silvicultures of Northern America, Volume 1:

1030 Conifers, Agricultural Handbook 654. Washington, D.C.: U. S. Department of Agriculture,

1031 Forest Service; 1990. p. 380-384

1032

1033 77. Wenninger EJ, Inouye RS. Insect community response to plant diversity and productivity in a

1034 sagebrush-steppe ecosystem. *Journal of Arid Environments*. 2008;72:24-33.

1035

1036 78. Johnson K, Sadoti G. Model transferability and implications for woodland management: a

1037 case study of Pinyon Jay nesting habitat. *Avian Conservation and Ecology* [Internet]. 2019;14(2)

1038 17. Available from: <http://www.ace-eco.org/vol14/iss2/art17/>

1039

1040 79. Tausch RJ, Tueller PT. Plant succession following chaining of pinyon-juniper woodlands in

1041 Eastern Nevada. *Journal of Range management*. 1977;30: 44-49.

1042

1043

1044 80. O'Meara TE, Haufler JB, Stelter LH, Nagy JG. Nongame wildlife response to chaining of

1045 pinyon-juniper woodlands. *Journal of Wildlife Management*. 1981;45: 381-389.

1046

1047 81. Commons ML, Baydack RK, Braun CE. Sage grouse response to pinyon-juniper

1048 management. *Proceedings of the conference on ecology and management of piñon–juniper*

1049 *communities in the Interior West*; 1999; Ogden, UT. In: Monsen S, Stevens R comps. USDA

1050 Forest Service. RMRS-P-9. p. 238-239

1051

1052 82. Johnson K, Petersen N, Smith J, Sadoti G. Piñon-juniper fuels reduction treatment impacts

1053 Pinyon Jay nesting habitat. *Global Ecology and Conservation*. 2018;16:1-7.

1054

1055 83. Gabaldon DJ. Factors involved in nest site selection by Pinon Jays [Dissertation]. Flagstaff,

1056 AZ: Northern Arizona University; 1979.

1057

1058 84. Mehringer PJ, Wigand PE. Comparison of late Holocene environments from woodrat

1059 middens and pollen: Diamond Craters, Oregon. In: Betancourt JL, Van Devender TR, Martin PS,

1060 editors. *Packrat Middens: The Last 40,000 Years of Biotic Change*. Tucson, AZ: University of

1061 Arizona Press; 1990. p. 294-325

1062

- 1063 85. Miller RF, Wigand PE. Holocene changes in semiarid pinyon-juniper woodlands.
1064 BioScience. 1994;44(7):465-474.
1065
- 1066 86. Cole KL, Henderson N, Shafer DS. Holocene vegetation and historic grazing impacts at
1067 Capitol Reef National Park reconstructed using packrat middens. Great Basin Naturalist.
1068 1997;57(4): 315-326.
1069
- 1070 87. Biondi F, Bradley M. Long-term survivorship of single-needle pinyon (*Pinus monophylla*) in
1071 mixed-conifer ecosystems of the Great Basin, USA. Ecosphere. 2013;4(10): 1-19.
1072
- 1073 88. Redmond MD, Forcella F, Barger NN. Declines in pinyon pine cone production associated
1074 with regional warming. Ecosphere. 2012;3(12):120. Available from: DOI: 10.1890/ES12-
1075 00306.1
1076
- 1077 89. Ricca MA, Coates PS, Gustafson KB, Brussee BE, Chambers JC, Espinosa SP, Gardner SC,
1078 Lisius SZ, Ziegler P, Delehanty DJ, Casazza ML. A conservation planning tool for Greater Sage-
1079 Grouse using indices of species distribution, resilience, and resistance. Ecological Applications.
1080 2018;28(4):878-896. Available from: DOI: 10.1002/eap.1690
1081
- 1082 90. Chambers JC, Allen CR, Cushman SA. Operationalizing ecological resilience concepts for
1083 managing species and ecosystems at risk. Frontiers in Ecology and Evolution. 2019;7:241.
1084 Available from: DOI: 10.3389/fevo.2019.00241
1085

Supporting Information Captions

S1 Table. All data used for data visualizations, ordinations, and logistic regression analysis, as summarized in Table 2. Each record refers to either a behavior-specific Pinyon Jay location or a control site. Each record includes various locational and data category attributes, including decimal latitude and longitude, along with all habitat attributes considered for inclusion in analysis, as described in Table 2. All attribute headers are sufficiently explicit to be self-explanatory when viewed in conjunction with Table 2. Definitions of codes for habitat types (which were not used in any analysis or data visualization) are available in Alexander 1988 [60]. Missing data are intrinsic to the FIA data set.

bioRxiv preprint doi: <https://doi.org/10.1101/2020.07.31.239367>; this version posted July 31, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. This article is a US Government work. It is not subject to copyright under 17 USC 105 and is also made available for use under a CC0 license.

Esri, HERE, DeLorme, MapmyIndia, © OpenStreetMap contributors, and the GIS user community

Fig 1

bioRxiv preprint doi: <https://doi.org/10.1101/2020.07.31.230367>; this version posted July 31, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. This article is a US Government work. It is not subject to copyright under 17 USC 105 and is also made available for use under a CC0 license.

BCR 9

BCR 16

Fig 2

Fig 4

Fig 5

Fig 6

Fig 7