1 **Effects of Starter Feeds of Different Physical Forms on Rumen Fermentation and** 2 **Microbial Composition for Pre-weaning and Post-weaning Lambs**

3

4 Yong Li,^{*} Yanli Guo,[#] Chengxin Zhang, Xiaofang Cai, Peng Liu, Cailian Li

5

6 College of Animal Science and Technology, Gansu Agricultural University, Lanzhou

7 730070, P.R. China

8

9

10

11

12

13

14

15

16

17

18

19 [#]Address correspondence to Yanli Guo, guoyl@gsau.edu.cn.

20 ^{*}Present address: Yong Li, Zhoukou Vocational and Technical College, Zhoukou, P. R.

21 China.

ABSTRACT

This study aimed to evaluate the effects of starter feeds of different physical forms on rumen fermentation and microbial composition for lambs. Twenty-four eight-day-old male Hu lambs (5.04 ± 0.75 kg body weight) were fed either milk replacer (MR) and pelleted starter feed (PS), or MR and textured starter feed (TS) in pre-weaning (day 8 to 35) and post-weaning (day 36 to 42) lambs. And the MR was fed by bottles to lambs at 2% of body weight at day 8 divided as three equal amounts at 08:00, 14:00 and 20:00 in pre-weaning. And the lambs were readily availed starter feeds and clean fresh water in the whole experiment. Six lambs for each treatment were euthanized at day 21 or 42 for sampling. The results showed the total volatile fatty acids, propionate and butyrate of rumen liquid in TS groups were all higher than them in PS groups respectively for pre-weaning and post-weaning lambs ($P < 0.05$), and the pH of rumen liquid in TS group was lower than it in PS group for post-weaning lambs ($P < 0.05$). Moreover, the pH of rumen and OTUs in TS group had trends to lower than them in PS group for pre-weaned lambs ($P = 0.061$, $P = 0.066$). TS established the predominant Phylum, *Bacteroidetes*, earlier than PS, and increased significantly the relative abundances of *Sharpea* compared to PS at level of genus ($P < 0.05$) for pre-weaning and post-weaning lambs. TS were more benefits to trigger rumen development for lambs.

IMPORTANCE Early use of starter feed could trigger rumen fermentation and establishment of dominant flora, which were in favour of growth and development of rumen for ruminants. The physical form of starter feed is one of the important factors to

promote rumen fermentation and establishment of dominant flora for ruminants of transition. However, limited study on effects of physical forms of starter feeds, especially the texturized starters containing steam-flaked grains, to rumen fermentative pattern and microbial composition for pre-weaning and post-weaning lambs to date. It was necessary to investigate the effects of physical form of starter feed on rumen fermentation and microbial composition for lambs. The significance of our research showed TS were better benefits to promote the rumen fermentation and establishment of dominant flora for lambs, which will greatly enhance our understanding of physical forms of starter feeds, leading to broader studies on rumen development for lambs.

KEYWORDS: starter feeds; physical forms; rumen fermentation; rumen microbial composition; lambs

INTRODUCTION

It is commonly known that a lot of complicated and diverse microbe, such as protozoan, bacteria, archeobacteria and fungi, existed in rumen of ruminant, whose interaction played an important role to maintain stable environments of rumen and health of animals (1). However, at birth, young ruminants did not possess anaerobic microbial population in their rumens. And establishment of rumen microbiota was very necessary to the physiological development of the rumens and the animal's abilities to convert plant feed into products that can be utilized by the animal for maintenance and production (2). Previous studies showed the introduction of solid diet around weaning could promote the establishment of anaerobic microbial ecosystem and formation of fermentation processes,

which were benefits to trigger the growth and development of the rumen (3).

Furthermore, processing of grain and ingredients of starter feeds and chemical and structural composition in dietary could affect fermentative pattern of rumen (4, 5). Feeding starters containing fine particles in form of mash or processed in form of pelleted could trigger rapid ruminal acid production from fermentation of carbohydrates (6), decreased ruminal pH (7). Consumption of starter feed containing highly fermentable carbohydrate by lambs or calves could increase the ruminal concentrations of volatile fatty acids (VFAs), particularly propionate and butyrate (8, 9), which was essential chemical stimulations for the rapid development of rumen epithelium (10). Furthermore, consumption of solid feed could promote the acquisition of anaerobic microbes and establishment of rumen fermentation (11).

Additionally, microbial colonization can also affect rumen development and function during early life, and the diversities and function of rumen microbe during early life was given concerned widely in recent years and next-generation sequencing was widely used to study rumen microbial ecology. Jami et al. and Li et al. found that the rumen microbe of prior to weaning calves has a similar functional capacity as that of a mature ruminant using next generation DNA (2, 12). Chen et al. studied the changes in bacterial diversity associated with epithelial tissue in the beef cow rumen during the transition to a high-grain diet (13). Jiao et al. taxonomic identified the ruminal epithelial bacterial diversity during rumen development in Goats (14). However, limited information is available on how changes in the physical forms of starter feeds, especially the texturized

starters containing steam-flaked grains, influence the rumen fermentative pattern and microbial composition for pre-weaning and post-weaning lambs to date. And there was not sufficient research on diversities and function of rumen microbe of starter feeds of different physical forms used next generation DNA for pre-weaning and post-weaning lambs.

Hence, the objectives of experiment were to elaborate the relation of effect of starter feeds of different physical forms on rumen fermentation and microbial composition for pre-weaning and post-weaning lambs by diversities and functions of rumen microbe (next-generation sequencing, 16S rDNA gene sequencing) so as to explain the reasons that effects of starter feeds of different physical forms on growth and development of rumen for pre-weaning and post-weaning lambs to certain extent.

RESULTS AND ANALYSIS

Rumen fermentative parameters. The total VFAs, propionate and butyrate of rumen in TS groups were all higher than them in PS groups respectively for pre-weaned and post-weaned lambs ($P < 0.05$), and the pH of rumen in TS group was lower than it in PS group for post-weaned lambs ($P < 0.05$, Table 1). Moreover, the pH of rumen in TS group had trends to lower than it in PS group in pre-weaned lambs ($P = 0.061$).

Sequence qualities of 16S rRNA genes and alpha diversities of rumen microbe. Raw sequences were joined together, optimized, quality controlled, then 1,648,195 high quality sequences were obtained. Every sample obtained 82,410 V3-V4 16S r RNA effective tags averagely. The length of effective tags was between 416 bp-426 bp. The

2,248 OTUs were obtained in all, and every sample had 562 OTUs averagely (Fig 1). Rarefaction curve of 16S rRNA gene showed Good' coverage was higher than 0.99. Based on similarity principles of 97% sequences between reads, OTUs coverage of sequences was adequate. As showed Fig 2, the rarefaction curve of 16S rRNA gene had trends to smooth, which showed sequences were reasonable. The span of rank abundance of 16S rRNA gene in the horizontal direction had trends to increase, which showed the abundance of species had trend to increase, and the span of rank abundance of 16S rRNA gene in the vertical direction had trends to smooth, which showed the distribution of species had trend to even. In a word, the sequences could reflect accurately the rumen microbial composition for lambs.

Except for OTUs of PS had trend to higher than them of TS for pre-weaning lambs ($P = 0.066$), the physical form of starter feed did not affect the alpha diversities of rumen bacterial communities for pre-weaning and post-weaning lambs ($P > 0.05$, Table 2).

Beta diversities of rumen microbe. Principal co-ordinates analysis (PCoA) of rumen bacterial OTUs showed contribution rates of PC1 and PC2 to differences among samples were 21.80% and 19.69% respectively, which could reflect adequately the differences among samples. PCoA results showed differences of samples were minute in same groups (Fig 3). Non-metric multi-dimensional scaling analysis (NMDS) of rumen bacterial OTUs (Fig 4) showed stress value lower 0.2 (0.133), which could indicate accurately the data and reflect the significant differences of rumen microbial structure and diversity ($P < 0.05$).

Effects of starter feeds of different physical forms on rumen microbe. Among the Phylum of top 10, the predominant Phylum was all *Bacteroidetes* and *Firmicutes* in every group, and their relative abundances were all higher than 24% (Fig 5). Only the relative abundances of *Bacteroidetes* of TS had trends to higher than them of PS for pre-weaning lambs ($P=0.084$). However, with intake of starter feed, the relative abundances of *Bacteroidetes* increased, and the relative abundances of *Firmicutes* decreased. The relative abundances of *Bacteroidetes* (61.96%) of TS had exceed them of *Firmicutes* (32.08%) and *Proteobacteria* (3.99%) for pre-weaning lambs, and become the first predominant Phylum, which were similar to them *Bacteroidetes* (65.36%) of TS for post-weaning lambs. But the relative abundances of *Bacteroidetes* (57.28%) of PS exceeded them of *Firmicutes* (31.21%) and *Proteobacteria* (1.15%) for post-weaning lambs, which were still lower than them of *Bacteroidetes* of TS for pre-weaning (61.96%) and post-weaning (65.36%) lambs.

It showed 153 rumen microflora were detected at the genus level between TS and PS for pre-weaning lambs, and the relative abundances of 13 rumen microflora were higher than 0.5%. The predominant microflora of PS and TS commonly were *unidentified_Prevotellaceae* (6.49% and 21.85%). The predominant microflora of PS peculiarly was *Lactobacillus* (14.53%), *Succinivibrio* (10.82%) and *unidentified_Cyanobacteria* (6.96%), and the predominant microflora of TS peculiarly was *Sharpea* (4.41%), *Dialister* (3.85%) and *Succinivibrio* (3.18%). The results of rumen microbe showed the PS increased significantly *unidentified_Clostridiales*, *Lactpcoccus*,

Sarcina, *unidentified_Cyanobacteria*, and TS increased significantly *sharpea* and *Oribacterium* compared between PS and TS for pre-weaning lambs at the genus level (Fig 6, LDA>4).

However, the effects of starter feed of two physical forms on rumen microflora were different for post-weaning lambs. And 139 rumen microflora were detected at the genus level between TS and PS for post-weaning lambs, and the relative abundances of 13 rumen microflora were higher than 0.5%. The predominant microflora of PS and TS commonly were *unidentified_Prevotellaceae* (29.57% and 38.49%). The predominant microflora of PS peculiarly was *Dialister* (7.23%), *unidentified_Lachnospiraceae* (6.72%), and the predominant microflora of TS peculiarly were *Sharpea* (7.43%) and *Succinivibrio* (6.87%). The results of rumen microbe showed only the TS increased significantly *sharpea* compared between PS and TS for post-weaned lambs at the genus level (Fig 7, LDA>4).

Functional prediction of rumen microbe. Functional profiles (KEGG level 2 pathways) of the two groups for pre-weaning and post-weaning lambs were all found to be similar in comparison (Fig 8). And the main functions of rumen microbe (the top seven) for lambs were replication and repair (pre-weaning: PS 10.56%, TS 11.03%; post-weaning: PS 11.44%, TS 11.39%), carbohydrate metabolism (pre-weaning: PS 11.04%, TS 10.95%; post-weaning: PS 10.73%, TS 10.72%), translation (pre-weaning: PS 9.86%, TS 10.33%; post-weaning: PS 10.65%, TS 10.67%), membrane transport (pre-weaning: PS 10.15%, TS 8.57%; post-weaning: PS 8.36%, TS 8.29%), amino acid metabolism (pre-weaning:

PS 8.35%, TS 8.31%; post-weaning: PS 8.16%, TS 8.22%), nucleotide metabolism (pre-weaning: PS 4.66%, TS 4.84%; post-weaning: PS 5.00%, TS 4.99%), energy metabolism (pre-weaning: PS 4.26%, TS 4.55%; post-weaning: PS 4.60%, TS 4.65%). Only the “Transporters” and “Fatty acid degradation” predicted function of KEGG level 3 pathways of PS were increased significantly and “Amino acid related enzymes” was decreased significantly for pre-weaning lambs compared to those of TS ($P < 0.05$, Fig 9). And the predicted function of KEGG level 3 pathways between PS and TS for post-weaning lambs still had no significant differences ($P > 0.05$).

DISCUSSION

Rumen pH was the most intuitive indicators reflected the fermentative condition of rumen, which could integrated reflect the rumen microbe, production, absorption and neutralization of organic acid. Furthermore, the acidity of rumen played a dominant role to maintain ruminal environment. Many factors affected the rumen pH, such as diet structure, secretion volume of saliva, speed of intake, VFA in rumen and the rates of production, absorption and excretion of organic acids. But the fundamental reasons of fluctuation of rumen pH were the diet structures and nutrition levels. Murphy and Kennelly indicated that rumen pH changed regularly from 5.0 to 7.5, which resulted from diet nature and time after intake (15). When intake fermentable carbohydrate, a lot of VFA could be produced that led to decrease of pH in rumen. In experiment, rumen pH changed from 5.14 to 5.96, which were in normal scope. However, rumen pH of TS was lower than them of PS respectively for pre-weaned and post-weaned lambs. The possible

reasons were TS contained steam-flaked corn, which included a lot of fermentable carbohydrate and reduced rumen pH. Nejad et al. showed steam-flaked corn increased the gelatinization of corn starch so as to change the degradation form of corn in rumen microbe, increase the surface area of corn and enhance hydrolytic ability of rumen microbe and enzyme to starch granule (16). Hence, TS increased fermentable carbohydrate utilized by rumen microbe, which produced a lot of VFA and reduced the rumen pH.

VFA were main products fermented by carbohydrates in rumen, included the acetate, propionate and butyrate, which were important energy sources of ruminant. Research found VFA were important promoting factors to growth and development of rumen epithelium. Furthermore, among VFA, stimulant action of butyrate was the most effective, following as propionate and acetate (17). In experiment, concentration of total VFA, propionate and butyrate of TS were all higher significantly than them of PS for pre-weaned and post-weaned lambs, which showed TS were better benefits to fermentation and development of rumen. These were consistent with the results of research about calves. Lesmeister et al. reported concentrations of VFA and propionate of TS contained steam flaking corn were higher than them of starter feed contained the whole corn, dry-rolled corn and roasted-rolled corn in serum of calve (18). Pavlata et al. also found concentrations of VFA, acetate and propionate of TS with chopped straw were higher significantly than them of PS (19). These results indicated that texture starter feed contained steam flaking corn could provide more chemical stimulations to development

of rumen, which were better benefits to fermentation and development of rumen for lambs. These were consistent with results of rumen development in experiment.

$\text{NH}_3\text{-N}$ was degradation products of protein nitrogen, non-protein nitrogen of diet and endogenous nitrogen in rumen, which were essential precursors to synthesize bacterial protein of rumen microbe. Maintaining reasonable $\text{NH}_3\text{-N}$ concentration was an important condition to growth and reproduction of rumen microbe. $\text{NH}_3\text{-N}$ concentration not only could reflect the speed of production and utilization, which degraded from nitrogenous substances by rumen microbe, but also could reflect the balances between degradation and synthesis of protein under specific diets to the extent. Many factors could affect the $\text{NH}_3\text{-N}$ concentration in rumen, such as protein quality of diets, emptying rate of chyme and absorption of rumen wall (20). It was reported that 5 mg/100ml $\text{NH}_3\text{-N}$ concentration was the lowest concentration which maintained growth and protein synthesis of rumen microbe. When $\text{NH}_3\text{-N}$ concentration was lower than 5 mg/100ml, the growth of rumen microbe would be suppressed. Moreover, the optimum $\text{NH}_3\text{-N}$ concentration which maintained growth of rumen microbe was 6.3-27.5mg/100ml (21). In current experiment, the starter feed of two different physical forms did not affect the $\text{NH}_3\text{-N}$ concentration of rumen for pre-weaning and post-weaning lambs, but the $\text{NH}_3\text{-N}$ concentration of rumen were under the optimum scopes maintained growth of rumen microbe. Beharka et al. and Pazoki et al. found that the $\text{NH}_3\text{-N}$ concentration in rumen liquid of calves had no significant differences between pelleted and textured starter feed (22, 23). Additionally, Qi and Ga et al. also found that the corn processed by pelletizing,

baking and steaming did not affect significantly the $\text{NH}_3\text{-N}$ concentration in rumen liquid of sheep (24, 25).

OTU could reflect the clustering quantities, and alpha diversities were used to evaluate the abundances and diversities of rumen microbe. Among alpha diversities, chao1 and Ace reflected the abundances of rumen microbe, whose higher chao1 and Ace indicated abundances of flora were greater; shannon and simpson reflected the diversities of rumen microbe, whose greater shannon and lower simpson showed diversities of flora were more (26). In experiment, OTU, chao1, ACE, shannon and simpson of rumen microbe all had no significant differences between two starter feed of different physical forms for pre-weaned and post-weaned lambs, only OTU of rumen microbe of TS had trends to lower than that of PS for pre-weaned lambs. These showed physical forms of starter did not affect the species, abundances and diversities of rumen microbe for pre-weaning and post-weaning lambs, only the species of rumen microbe of TS had trends to lower than that of PS for pre-weaning lambs. The results showed the lambs were easier to adopt the TS during courses of transition from liquid to solid starter feed, and could urge rumen of lambs to establish dominant flora and disappear instantaneous flora.

Intake of starter feed early could change rumen flora for pre-weaning lambs (27). With increasing of ages and intake of starter feed for lambs, *Bacteroidetes*, *Firmicutes* and *Proteobacteria* could become the main dominant flora of higher relative abundances, and the relative abundances of *Bacteroidetes* were increased, and the relative abundances of *Firmicutes* and *Proteobacteria* were reduced (2). When calves intake MR and starter

feed, the *Proteobacteria* of rumen could be replaced by *Bacteroidetes* at ages of 42d, and the relative abundances of *Bacteroidetes* become the highest, which were possible to be related to chemical composition of diets (12). In experiment, the *Firmicutes* and *Proteobacteria* of TS had been replaced by *Bacteroidetes* for pre-weaning lambs, and *Bacteroidetes* had become the first predominant Phylum, and achieve similar relative abundances to them of TS for post-weaning lambs (pre-weaning, 61.96%; post-weaning, 65.36%). However, the *Firmicutes* and *Proteobacteria* of PS were replaced by *Bacteroidetes* for post-weaning lambs, and *Bacteroidetes* become the first predominant Phylum. Furthermore, relative abundances of *Bacteroidetes* of PS (57.28%) for post-weaning lambs were still lower than them of TS for pre-weaning (61.96%) and post-weaning (65.36%) lambs. The results showed the lambs were easier to adopt the TS during courses of transition from liquid to solid starter feed, and could urge rumen of lambs to establish dominant flora.

Among the rumen microflora of ruminants, *Bacteroidetes* and *Firmicutes* were two main dominant floras (28). It was well known that rumen microbe, especially *Bacteroidetes*, played an important role in degradation of starch and protein of diets, synthesizes of protein of rumen microbe, absorption of peptides and amino acids (29). And *Firmicutes* contained many bacterial degraded fibres, such as *Ruminococciis*, *Eubacterium*, *Pseudobutyrvibrio*, *Butyrovibro* and *Oscillibacter*, whose main function were to degrad the cellulose (30). These showed *Bacteroidetes* was better benefits to degrade the concentrated diets, and *Firmicutes* was better benefits to degrade the

roughage diets. Jiang et al. found the relative abundances of *Bacteroidetes* of fermented corn gluten meal were significant higher than them of corn gluten meal, and relative abundances of *Firmicutes* of fermented corn gluten meal were significant lower than them of corn gluten meal (31). In experiment, compared to PS, TS increased the relative abundances of *Bacteroidetes* for pre-weaning (PS, 35.74%; TS, 61.96%) and post-weaning (PS, 57.28%; TS, 65.36%) lambs, decreased the relative abundances of *Firmicutes* for pre-weaning (PS, 37.88%; TS, 32.08%) and post-weaning (PS, 31.21%; TS, 24.03%) lambs. These results were consistent with previous research. These results indicated physical forms of starter feeds affected the structures of rumen microbe at levels of phylum for pre-weaning and post-weaning lamb, and TS was better benefits to promote the fermentation of rumen and absorption of nutrients for lambs. The possible reasons were corns of TS processed by steam flaking, and improved the gelatinization of starch, contained more fermentable carbohydrates, which were better benefits to fermentation of *Bacteroidetes*. These were also consistent with results of fermentative parameters of rumen.

However, Kim et al. found the dominant flora in rumen was *Firmicutes* and *Bacteroidetes* in turn (32, 33). The reasons of differences might be related to composition of diets. Their diets were type of roughage, and the diets were type of concentration in current experiment. Additionally, the research also found when fed diets of higher proportional concentration, the dominant flora in rumen were *Bacteroidetes*; and when fed diets of higher proportional roughage, the dominant flora in rumen were *Firmicutes*

(34). And *Bacteroidetes* were correlated negatively to *Firmicutes* (35). These proved further TS were better effective to fermentation of rumen for pre-weaning and post-weaning lambs.

The main fermented products of *sharpea* in rumen of sheep were lactates, and formation of lactates promoted further fermentation of *sharpea*, urged lactates to change into butyrate, which led to produce lower H₂ compared to traditional fermentation directly from carbohydrate to butyrate and reduced production of CH₄ in rumen (36, 37). In current experiment, compared to PS, TS increased significantly the relative abundances of *sharpea* of rumen for pre-weaning and post-weaning lambs, which were consistent with significant higher concentration of butyrate of TS in rumen liquid. These results showed TS contained steam flaking corn were benefits to improve rumen microflora, urge rumen fermentation for lambs. At same time, Xue et al. also found concentrations of butyrate in rumen liquid were correlated positively to the relative abundances of *sharpea* when they compared effects of higher yield, higher concentrations of milk protein and lower yield, lower concentrations of milk protein on rumen microflora of calves (38). Lin et al. also proved when fed starter feed, *Sharpea* produced lactates was main enriched in rumen of lambs (39).

Additionally, we found PS also increased significantly the relative abundances of *unidentified_Clostridiales*, *Lactpcoccus*, *Lactobacillus*, *Sarcina* and *unidentified_Cyanobacteria* for pre-weaning lambs compared TS in the experiment. *Clostridiales* and *Lactpcoccus* were all the main representative genus of *Firmicutes*,

which could degrade cellulose in rumen. Hence, the higher relative abundances of *Clostridiales* and *Lactpcoccus* of PS were consistent with higher relative abundances of *Firmicutes* of PS (PS, 37.88%; TS, 32.08%) at levels of phylum for pre-weaning lambs. *Lactobacillus* could ferment carbohydrates to produce lactates, which were benefits to health of animals. Hence, *Clostridiales*, *Lactpcoccus* and *Lactobacillus* played decisive roles in digestion and absorption of nutrients in gastrointestinal tract and immunities of animals. *Cyanobacteria* were one of microalgae, which could improve the performances of animals and qualities of meat as feed additives (40). *Sarcina* was related to rumen bloating of lambs (41) and calves (42), which should not exist in digestive tracts of animals.

However, predicted functions of rumen were found to be similar in two groups for pre-weaning and post-weaning lambs in current experiment. Only the “Transporters” and “Fatty acid degradation” of PS were increased significantly and “Amino acid related enzymes” was decreased significantly for pre-weaning lambs compared to those of TS. It was confirmed previously that significant changes of microbial composition might not lead to a shift of function because many microbes shared the same metabolic pathways. Li et al. found that all of the functional classes between two age groups (d14 and d42 of calves) were similar, suggesting that although their phylogenetic composition greatly fluctuated, the rumen microbial communities of pre-ruminant calves maintained a stable function and metabolic potentials (12). These might be the reasons that two group lambs had similar functions in this experiment. Maybe it was necessary to analyse the rumen

microbiome functions using metagenomic and/or metabolomics technologies for completed and integrated understand the impact of rumen function further.

In a word, physical forms of starter feeds affected the fermentation and microbial composition of rumen for pre-weaning and post-weaning lambs. TS were better benefits to improve fermentation environment and establish dominant flora of rumen early, which were in favour of growth and development of rumen for pre-weaned and post-weaning lambs.

MATERIALS AND METHODS

Animals, feeds and experimental design. This experiment was carried out on a local sheep farm (Baiyin Kangrui breeding sheep co., Baiyin, Gansu province). All the experimental protocols performed in this study were approved by the Animal Care Committee of Gansu Agricultural University and the experimental procedures used in this study were in accordance with the recommendations of the University's guidelines for animal research. In this study, twenty-four healthy male Hu lambs, whose average body weight were 5.04 ± 0.75 kg, were separated from their dams at day 8 and moved into a naturally ventilated barn with individual cages (0.8×1.3 m). And the trial lambs were fed either milk replacer (MR) and pelleted starter feed (PS, with a mean particle size of six mm diameter), or MR and textured starter feed (TS, which included coarse mashed steam-flaked corn, also with a mean particle size of six mm diameter) in pre-weaning (day 8 to 35) and post-weaning (day 36 to 42) lambs. And the MR (23% CP and 12% fat, DM basis) was fed by single bottles to lambs at 2% of body weight at day 8 divided as

three equal amounts at 08:00, 14:00 and 20:00 in pre-weaning. After weaning, all lambs continued to be fed starter according to their trial group. And all lambs had free access to readily avail clean fresh water and their respective ad lib starter feed in the whole experiment. The diets (Table 3) were prepared by Gansu Aonong Feed Co., Ltd according to National Research Council recommendations (NRC, 2007). The MR was bought from Beijing Accurate Animal Nutrition Research Center.

Sample collection. Lambs for each treatment were euthanized by captive bolt stunning and exsanguinated in a specialized room of the experimental farm without any transportation at the age of 21 or 42 days. After slaughter, a part of rumen content was used to collect rumen liquid by immediately filtering through four layers of swab and transferring into 10 cm centrifuge tubes, and then stored at -20°C to analyse the total volatile fatty acids (VFAs) and ammonia nitrogen ($\text{NH}_3\text{-N}$); the other part of rumen content was collected for storage at -80°C for rumen bacteria analysis.

Determination of rumen fermentative parameters. After slaughter, the rumen content was mixed and determined pH immediately by PB-10 acidity meter (Zedorius, Kogentin, Germany). Rumen liquid were thawed and analyzed for individual and total VFA concentrations by gas chromatography (AI 3000, Thermo, Germany) (43) and $\text{NH}_3\text{-N}$ by colorimetric method (44) using visible spectrophotometer (Shanghai Jinghua Technology Co. Ltd). Details of VFA determination were as follows:

0.6 μL rumen liquid samples were injected by an auto sampler into an AE-FFAP (30m $\times$ 0.25mm $\times$ 0.33 μm ; Zhongke Antai, Lanzhou, China) . Chromatographic

conditions: temperature of injection entrance 200°C; N₂ flow 2.0 mL•min⁻¹; split ratio 40:1; procedure heating mode (120°C 3 min, 10°C•min⁻¹ to 180°C, kept 1 min); detector temperature 250°C; FID air, H₂ and N₂ flow were 450 mL•min⁻¹, 40 mL•min⁻¹ and 45 mL•min⁻¹ respectively; cylinder heating procedure: from 45°C to 150°C as speed 20°C•min⁻¹, and kept 5min. Finally, the peak integration was performed using Chromeleon® Software.

Total DNA extraction of rumen microbe, illumina sequencing. Rumen content was sent to Novogene Bioinformatics Technology Co., Ltd. (Beijing, China) to extract DNA and sequence of rumen microbe. Details were following:

Total DNA of rumen microbe in rumen content was extracted by thecetyltrimethylammonium bromide method (45) with a bead-beater (Biospec Products; Bartlesville, OK, United States) as described by Gagen et al. (46). The amplification of V₃-V₄ hypervariable region of the 16S rRNA gene was carried out with formwork of each of the DNA samples using the primer set 515F/806R and Phusion® High-Fidelity PCR Master Mix (New England Biolabs, Ipswich, MA, United States) as described by Caporaso et al. (47). When each forward and reverse primer had a 6-bp error-correcting barcode at the 5' terminus, it was seen as unique to each DNA sample. The sequencing for all samples was on an Illumina HiSeq platform by Novogene Bioinformatics Technology Co., Ltd. (Beijing, China) to generate 2 × 250 bp paired end reads.

After the paired-end reads were cut off barcode and primer, they were joined together and formed single sequences using FLASH based on overlapping regions (48).

Sequences with a quality score of <20 and a length of >300 bp or <200 bp were filtered and discarded using Quantitative Insight into Microbial Ecology (QIIME, 49). At same time, the possible chimeric sequences were also identified and removed from the sequences using the usearch61 algorithm in USEARCH 6.1 (50). Operational taxonomic units (OTUs) were clustered as 97% similarity, and chosen the representative sequence according to the algorithmic principle and annotation analysed in SILVA SSU rRNA database (51) used Uparse (Upase v7.0.1001, 52) by Mothur method (53).

Microbial functional prediction. Microbial function predicted byTax4Fun based on 16S Silva database (54). Details were as following:

All the genes 16S rRNA sequences of prokaryote in Kyoto Encyclopedia Genes and Genomes (KEGG) database were extracted, then compared in SILVA SSU Ref NR database(BLAST bitscore >1500)by BLASTN method and established correlation matrix. Finally, functional information of all the genes 16S rRNA sequences of prokaryote annotated in KEGG database were compared with functional information in SILVA database by UProC and PAUDA so as to achieve the goals of microbial functional prediction.

Statistical analyses. Data for rumen fermentative and metabolic parameters, alpha diversity indices (number of OTU, ACE, Chao1, Shannon and Simpson index) of rumen microbe were pooled at each time point for the six slaughtered lambs in each group. Data were analysed as independent sample t-tests (SPSS 20.0, Inc., Chicago, IL, USA). Significance was designated as $P<0.05$ with a trend being between $P\geq 0.05$ and $P<0.10$.

Beta diversity of rumen microbe was analysed by T-test and wilcox-test. Analysis of similarity between groups was calculated by Bray-Curtis. Principal co-ordinates analysis (PCoA) of rumen bacterial OTUs calculated the distance by Bray-Curtis firstly, then drawn the fig by R software (v3.3.0). Difference between groups was analysed by LEfSe (LDA Effect Size), and $LDA > 4$ was different marking of statistics and biology (55).

ACKNOWLEDGMENTS

This work was supported by National Natural Science Foundation of China (31860654). The authors express their kind appreciation to Dr. Xiaojuan Wang and Ligang Yuan from the Gansu Agricultural University for their assistance throughout the experiments.

REFERENCES

1. Bi YL, Zeng SQ, Zhang R, Diao QY, Tu Y. 2018. Effects of dietary energy levels on rumen bacterial community composition in Holstein heifers under the same forage to concentrate ratio condition. BMC Microbiol 18: 69.
2. Jami E, Israel A, Kotser A, Mizrahi I. 2013. Exploring the bovine rumen bacterial community from birth to adulthood. ISME J 7: 1069-1079.
3. Khan MA, Weary DM, von Keyserlingk MAG. 2011. Invited review: Effects of milk ration on solid feed intake, weaning, and performance in dairy heifers. J Dairy Sci 94: 1071-1081.
4. Huntington GB. 1997. Starch utilization by ruminants: From basics to the bunk. J Anim Sci 75: 852-867.

- 442 5. Owens WE, Ray CH, Watts JL, Yancey RJ. 1997. Comparison of success of antibiotic
443 therapy during lactation and results of antimicrobial susceptibility tests for bovine
444 mastitis. J Dairy Sci 80:313-317.
- 445 6. Laarman AH, Ruiz-Sanchez AL, Sugino T, Guan LL, Oba M. 2012. Effects of feeding
446 a calf starter on molecular adaptations in the ruminal epithelium and liver of Holstein
447 dairy calves. J Dairy Sci 95: 2585-2594.
- 448 7. Laarman AH, Oba M. 2011. Short communication: Effect of calf starter on rumen pH
449 of Holstein dairy calves at weaning. J Dairy Sci 94: 5661-5664.
- 450 8. Suarez-Mena FX, Heinrichs AJ, Jones CM, Hill TM, Quigley JD. 2015. Digestive
451 development in neonatal dairy calves with either whole or ground oats in the calf
452 starter 1. J Dairy Sci 98: 3417-3431.
- 453 9. Baldwin RL, McLeod KR, Klotz JL, Heitmann RN. 2004. Rumen development,
454 intestinal growth and hepatic metabolism in the pre and postweaning ruminant. J
455 Dairy Sci 87: E55-E65.
- 456 10. Mirzaei M, Khorvash M, Ghorbani GR, Kazemi-Bonchenari M, Riasi A, Nabipour A,
457 Borne JJG Cvd. 2015. Effects of supplementation level and particle size of alfalfa hay
458 on growth characteristics and rumen development in dairy calves. J Anim Physiol
459 Anim Nutr 99: 553-564.
- 460 11. Dockalova H, Stastnik O, Krivova S, Sedlakova L, Pavlata L. 2016. Effect of
461 different physical forms of starter on feed intake and performance of calves. Mendel
462 Net 23: 201-204.

- 463 12. Li RW, Connor EE, Li C, Baldwin VI, Ransom L, Sparks ME. 2012. Characterization
464 of the rumen microbiota of pre-ruminant calves using metagenomic tools. *Environ*
465 *Microbiol* 14: 129-139.
- 466 13. Chen Y, Penner GB, Li M, Oba M, Guan LL. 2011. Changes in bacterial diversity
467 associated with epithelial tissue in the beef cow rumen during the transition to a
468 high-grain diet. *Appl Environ Microbiol* 77: 5770-5781.
- 469 14. Jiao J, Huang J, Zhou C, Tan Z. 2015. Taxonomic identification of ruminal epithelial
470 bacterial diversity during rumen development in goats. *Appl Environ Microbiol* 81:
471 3502-3509.
- 472 15. Murphy JJ, Kennelly JJ. 1987. Effect of protein concentrate and protein source on the
473 degradability of dry matter and protein in situ. *J Dairy Sci* 70: 1841-1849.
- 474 16. Nejad Ghassemi J, Torbatinejad N, Naserian AA. 2012. Effects of processing of
475 starter diets on performance, nutrient digestibility, rumen biochemical parameters and
476 body measurements of Brown Swiss dairy calves. *Asian-Australas. J Anim Sci* 25:
477 980-987.
- 478 17. Tamate H, McGilliard AD, Jacobson NL, Getty R. 1962. Effect of various dietaries on
479 the anatomical development of the stomach in the calf. *J Dairy Sci* 45: 408-420.
- 480 18. Lesmeister KE, Heinrichs AJ. 2004. Effects of corn processing on growth
481 characteristics, rumen development, and rumen parameters in neonatal dairy calves. *J*
482 *Dairy Sci* 87: 3439-3450.
- 483 19. Pavlata L, Sťastnik O, Křivova S, Dockalova H, Sedlakova L, Mrkvicova E, Doležal

- 484 P. 2017. The effect of different physical forms of starter feed on rumen fermentation
485 indicators and weight gain in calves after weaning. *Acta Vet Brno* 86: 285-291.
- 486 20. Hristov AN, Ropp JK, Hunt CW. 2002. Effect of barley and its amylopectin content
487 on ruminal fermentation and bacterial utilization of ammonia-N in vitro. *Anim Feed*
488 *Sci Tech* 99: 25-36.
- 489 21. Ortega ME, Stem MD, Satter LD. 1979. The effect of rumen ammonia concentrate
490 on dry matter disappearance in situ. *J Dairy Sci* 62: 76 (Abstr).
- 491 22. Beharka AA, Nagaraja TG, Morrill JL, Kennedy GA, Klemm RD. 1998. Effects of
492 form of the diet on anatomical, microbial, and fermentative development of the rumen
493 of neonatal calves. *J Dairy Sci* 81: 1946-1955.
- 494 23. Pazoki A, Ghorbani GR, Kargar S, Sadeghi-Sefidmazgi A, Ghaffari MH, Drackley JK.
495 2017. Growth performance, nutrient digestibility, ruminal fermentation, and rumen
496 development of calves during transition from liquid to solid feed: Effects of physical
497 form of starter feed and forage provision. *Anim Feed Sci Tech* 234: 173-185.
- 498 24. Qi ZL. 2001. Effects of processed corns on rumen fermentation, nutrient digestion
499 and metabolism in sheep. Academic dissertation of master degrees in Inner Mongolia
500 Agricultural University, Hohhot, China.
- 501 25. Ga ED, ZL, Zhang RH, Chen HJ, Zhao FR. 2002. Effects of different processed corns
502 on pH, NH₃-N and VFA in rumen liquid of sheep. *Heilongjiang journal of animal*
503 *science and veterinary medicine* 9: 18-20.
- 504 26. Zeng Y, Gao YH, Peng ZL, Chen SY, Xie XT, Miao JJ, Bai X, Guo CH. 2020. Effects

of yeast culture supplementation in diets on rumen fermentation parameters and
microflora of housed-feeding yak. Chinese journal of animal nutrition 32: 1721-1733.

27. Wang W, Li C, Li F, Wang X, Zhang X, Liu T, Nian F, Yue X, Li F, Pan X, La Y, Mo F,
Wang F, Li B. 2016. Effects of early feeding on the host rumen transcriptome and
bacterial diversity in lambs. Sci Rep 6: 3247-3249.

28. Singh KM, Jakhesara SJ, Koringa PG, Rank DN, Joshi CG. 2012. Metagenomic
analysis of virulence-associated and antibiotic resistance genes of microbes in rumen
of Indian buffalo (*Bubalus bubalis*). Gene 507: 146-151.

29. Castillo-Lopez E, Moats J, Aluthge ND, Ramirez Ramirez HA, Christensen DA,
Mutsvangwa T, Fernando SC. 2018. Effect of partially replacing a barley-based
concentrate with flaxseed-based products on the rumen bacterial population of
lactating Holstein dairy cows. J Appl Microbiol 124: 42-57.

30. Evans NJ, Brown JM, Murray RD, Getty B, Birtles RJ, Hart CA, Carter SD. 2011.
Characterization of novel bovine gastrointestinal tract *Treponema* isolates and
comparison with bovine digital dermatitis treponemes. Appl Environ Microbiol 77:
138-147.

31. Jiang X, Ma GM, Cui ZQ, Li Y, Zhang YG. 2020. Effects of fermented corn gluten
meal on growth performance, plasma metabolites, rumen fermentation and bacterial
community of Holstein calves during the pre-weaning period. Livest Sci 231: 103866.

32. Kim M, Morrison M, Yu ZT. 2011. Phylogenetic diversity of bacterial communities in
bovine rumen as affected by diets and microenvironments. Folia Microbiol 56:

526 453-458.

527 33. Kim M, Felix TL, Loerch SC, Yu ZT. 2014. Effect of Haylage and Monensin
528 Supplementation on Ruminal Bacterial Communities of Feedlot Cattle. *Curr.*
529 *Microbiol* 69: 169-175.

530 34. Fernando SC, Purvis HT, Najjar FZ, Sukharnikov LO, Krehbiel CR, Nagaraja TG,
531 Roe BA, Desilva U. 2010. Rumen microbial population dynamics during adaptation
532 to a high-grain diet. *Appl Environ Microbiol* 76: 7482-7490.

533 35. Rice WC, Galyean ML, Cox SB, Dowd SE, Cole NA. 2012. Influence of wet
534 distillers grains diets on beef cattle fecal bacterial community structure. *BMC*
535 *Microbiol* 12: 25.

536 36. Kamke J, Kittelmann S, Soni P, Li Y, Tavendale MH, Ganesh S, Janssen PH, Shi WB,
537 Froula J, Rubin EM, Attwood GT. 2016. Rumen metagenome and metatranscriptome
538 analyses of low methane yield sheep reveals a *Sharpea*-enriched microbiome
539 characterised by lactic acid formation and utilisation. *Microbiome* 4: 56.

540 37. Kumar S, Treloar BP, The KH, Mckenzie CM, Henderson G, Attwood GT, Waters SM,
541 Patchett ML, Janssen PH. 2018. *Sharpea* and *Kandleria* are lactic acid producing
542 rumen bacteria that do not change their fermentation products when co-cultured with
543 a methanogen. *Anaerobe* 54: 31-38.

544 38. Xue MY, Sun HZ, Wu XH, Guan LL, Liu JX. 2019. Assessment of rumen bacteria in
545 dairy cows with varied milk protein yield. *J Dairy Sci* 102: 5031-5041.

546 39. Lin LM, Xie F, Sun DM, Liu JH, Zhu WY, Mao SY. 2019. Ruminal microbiome-host

547 crosstalk stimulates the development of the ruminal epithelium in a lamb model.
548 Microbiome 7:1-16.

549 40. Madeira MS, Cardoso C, Lopes P., Coelho D, Afonso C, Bandarra NM, Prates JAM.
550 2017. Microalgae as feed ingredients for livestock production and meat quality: A
551 review. *Livest Sci* 205: 111-121.

552 41. Vatn S, Gunnes G, Nybo, K, Juul HM. 2000. Possible involvement of *Sarcina*
553 *ventriculi* in canine and equine acute gastric dilation. *Acta Vet Scand* 41: 333-337.

554 42. Edwards GT, Woodger NG, Barlow AM, Bell SJ, Harwood DG, Otter A, Wight AR.
555 2008. *Sarcina*-like bacteria associated with bloat in young lambs and calves. *Vet. Rec*
556 163: 391-393.

557 43. Isac M, García M, Aguilera J, Alcaide EM. 1994. A comparative study of nutrient
558 digestibility, kinetics of digestion and passage and rumen fermentation pattern in
559 goats and sheep offered medium quality forages at the maintenance level of feeding.
560 *Arch Anim Nutr* 46: 37-50.

561 44. Feng ZC, Gao M. 2010. An improved method for determination of ammonia nitrogen
562 in rumen liquid by colorimetry. *Animal Husbandry and Feed Science* 31: 37.

563 45. Brookman JL, Nicholson MJ. 2005. “Anaerobic fungal populations,” in *Methods in*
564 *Gut Microbial Ecology for Ruminants*, eds Makkar HPS, McSweeney CS. 139-150.
565 Springer, Berlin.

566 46. Gagen EJ, Denman SE, Padmanabha J, Zadbuke S, Al Jassim R, Morrison M,
567 McSweeney CS. 2010. Functional gene analysis suggests different acetogen

568 populations in the bovine rumen and tammar wallaby forestomach. *Appl Environ*
569 *Microbiol* 76: 7785-7795.

570 47. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ,
571 Knight R. 2011. Global patterns of 16S rRNA diversity at a depth of millions of
572 sequences per sample. *P Natl Acad Sci USA* 108: 4516-4522.

573 48. Mago^ˆc T, Salzberg SL. 2011. FLASH: fast length adjustment of short reads to
574 improve genome assemblies. *Bioinformatics* 27: 2957-2963.

575 49. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK,
576 Knight R. 2010. QIIME allows analysis of high-throughput community sequencing
577 data. *Nat Methods* 7: 335-336.

578 50. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: a versatile open
579 source tool for metagenomics. *Peer J* 4: 2584.

580 51. Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naive Bayesian classifier for rapid
581 assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ*
582 *Microbiol* 73: 5261-5267.

583 52. Haas BJ, Gevers D, Earl AM, Feldgarden M, Ward DV, Giannoukos G, Ciulla D,
584 Tabbaa D, Highlander SK, Sodergren E, Methé B, DeSantis TZ. 2011. Human
585 Microbiome Consortium, Petrosino JF, Knight R, Birren BW,. Chimeric 16S
586 rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR
587 amplicons. *Genome Res* 21: 494-504.

588 53. Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon

reads. Nat Methods 10: 996-998.

54. Aßhauer KP, Wemheuer B, Daniel R, Meinicke P. 2015. Tax4Fun: predicting functional profiles from metagenomic 16S rRNA data. Bioinformatics 31: 2882-2884.
55. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C. 2011. Metagenomic biomarker discovery and explanation. Genome Biol 12: 1-18.

610 **Table 1 Rumen fermentative parameters of lambs**

Items	Pelleted	Textured	SEM	<i>P-value</i>
Pre-weaning				
pH	5.96	5.60	0.12	0.061
Total volatile fatty acids (mmol/L)	24.41	30.04	1.72	0.046
)				
Acetate (mmol/L)	13.94	15.56	1.30	0.414
Propionate (mmol/L)	6.69	9.07	0.72	0.046
Butyrate (mmol/L)	2.02	3.16	0.19	0.006
Isobutyrate (mmol/L)	0.34	0.48	0.05	0.110
Valerate (mmol/L)	0.85	1.08	0.21	0.465
Isovalerate (mmol/L)	0.56	0.70	0.09	0.320
Acetate to propionate	2.16	1.76	0.26	0.308
Ammonia nitrogen (mg/100ml)	12.51	14.87	0.93	0.107
Post-weaning				
pH	5.48	5.14	0.07	0.011
Total volatile fatty acids (mmol/L)	65.59	70.72	1.51	0.046
)				
Acetate (mmol/L)	38.89	38.02	1.65	0.739
Propionate (mmol/L)	16.73	21.56	1.49	0.044
Butyrate (mmol/L)	5.79	7.67	0.50	0.023
Isobutyrate (mmol/L)	0.45	0.46	0.04	0.989
Valerate (mmol/L)	3.03	2.20	0.58	0.332
Isovalerate (mmol/L)	0.70	0.81	0.08	0.317
Acetate to propionate	2.52	1.81	0.29	0.153
Ammonia nitrogen (mg/100ml)	12.25	12.75	0.82	0.677

611

612

613

614

615

616

617

618

619 ***Table 2 Alpha diversities of rumen bacterial communities***

Items	Pelleted	Textured	SEM	<i>P-value</i>
Pre-weaning				
OTU	415.80	352.20	40.79	0.066
ACE	421.40	374.85	44.74	0.146
Chao1	415.79	406.93	94.83	0.893
Shannon	3.74	4.09	0.27	0.407
Simpson	0.78	0.84	0.09	0.323
Post-weaning				
OTU	279.00	258.40	48.90	0.557
ACE	282.18	263.49	43.66	0.556
Chao1	272.76	258.52	40.10	0.645
Shannon	3.61	3.44	0.57	0.662
Simpson	0.81	0.79	0.09	0.766

620

621

622

623

624

625

626

627

628

629

630

631

Table 3

Ingredients and chemical composition of the experimental diets

Item	Pelleted	Textured
Ingredients (g/kg)		
Alfalfa hay	50.00	50.00
Corn grain, ground	650.00	-
Corn grain, steam flaked	-	650.00
Wheat bran	50.00	50.00
Expanded soybean	60.00	60.00
Soybean meal	165.00	165.00
Salt	3.00	3.00
Calcium carbonate	11.80	11.80
Vitamin and mineral mix ^a	10.00	10.00
Sweetening agent	0.20	0.20
Total	1000.00	1000.00
Chemical composition (g/kg, DM)		
Apparent digestible energy b(MJ/kg)	13.75	13.82
Dry Matter ^b	892.40	885.00
Crude Protein ^b	214.60	216.40
Ether extract ^b	27.20	27.90
Neutral detergent fiber ^b	128.00	128.80
Non-fiber carbohydrates ^c	625.10	622.20
Starch ^b	115.10	117.10
Calcium ^b	7.50	7.10
Total phosphorus ^b	4.48	4.52

^a Provided per kilogram of premix: Fe 75000 mg, Zn 15000 mg, Cu 3500 mg, Mn 15000 mg, I

500000 mg, Se 50 mg, Co 200 mg, VA 2500000 IU, VD 1000000 IU, VE 1900 IU.

^b The actual values for measurement.

^c Non-fiber carbohydrates=1000 – (crude protein + ether extract + aNDF + ash) (NRC, 2001).

Fig 1 Venn graph

Fig 2 Rarefaction curve and Rank abundance of 16S rRNA gene

Note: A, rarefaction curve; B, rank abundance.

654

655

656 Fig 3 Principal co-ordinates analysis (PCoA) of rumen bacterial OTUs

657

658

659

660 Fig 4 Non-metric multi-dimensional scaling analysis (NMDS) of rumen bacterial OTUs

661

662

Fig 5 Phylum of top 10 of rumen microbe for lambs

Fig 6 Value of linear discriminant analysis (LDA) effect size (LEf Se) on rumen microflora between two starter feeds for pre-weaned lambs

Fig 7 Value of linear discriminant analysis (LDA) effect size (LEf Se) on rumen microflora between two starter feeds for post-weaned lambs

Fig 8 Relative abundances of top 10 function of rumen microflora between two starter feeds for pre-weaning and post-weaned lambs (KEGG level 2 pathways)

Fig 9 Analysis of different metabolism pathways between two starter feeds for pre-weaning lambs (KEGG level 3 pathways)

Stress: 0.133

PS21 TS21

Cladogram

PS21
TS21

a: f_Rikenellaceae
b: f_unidentified_Cyanobacter
c: o_unidentified_Cyanobacter
d: c_unidentified_Cyanobacter
e: f_Lactobacillaceae
f: o_Lactobacillales
g: c_Bacilli
h: f_unidentified_Clostridiales
i: f_Erysipelotrichaceae
j: o_Erysipelotrichales
k: c_Erysipelotrichia

Cladogram

PS42
TS42

a: f__Lachnospiraceae
b: f__Erysipelotrichaceae
c: o__Erysipelotrichales
d: c__Erysipelotrichia
e: f__Acidaminococcaceae