Date: 8/6/2020

Expression and distribution of synaptotagmin isoforms in the zebrafish retina

Diane Henry^{1,3*}, Christina Joselevitch^{1,3*}, Gary G. Matthews^{1,3}, and Lonnie P. Wollmuth^{1,2,3}

¹Department of Neurobiology & Behavior, ²Department of Biochemistry & Cell Biology,

³Center for Nervous System Disorders, Stony Brook University, Stony Brook, NY 11794-5230

*These authors contributed equally to this work.

Abbreviated title: Synaptotagmins in zebrafish retina

Contents: Pages: 49; Figures: 7; Tables: 7.

Word Counts: 10,761 words (Abstract 249; Introduction 732; Discussion 1304).

Correspondence:

Dr. Lonnie P. Wollmuth: lonnie.wollmuth@stonybrook.edu

Tel: (631) 632-4186, Fax: (631) 632-6661

Acknowledgments: This work was supported by a NIH RO1 grant from NEI (EY003821 to

GGM/LPW). We thank Miaomiao He for assistance with protein structure modeling and David

Zenisek and Howard Sirotkin for helpful discussions and/or comments on the manuscript.

Conflict of interest: None.

ABSTRACT

Synaptotagmins belong to a large family of proteins. While various synaptotagmins have been implicated as Ca^{2+} sensors for vesicle replenishment and release at conventional synapses, their roles at retinal ribbon synapses remain incompletely understood. Zebrafish is a widely used experimental model for retinal research. We therefore investigated the homology between human, rat, mouse, and zebrafish synaptotagmins 1 to 10 using a bioinformatics approach. We also characterized the expression and distribution of various synaptotagmin (*syt*) genes in the zebrafish retina using RT-PCR and *in situ* hybridization, focusing on the family members whose products likely underlie Ca^{2+} -dependent exocytosis in the central nervous system (synaptotagmins 1, 2, 5 and 7). We find that most zebrafish synaptotagmins are well conserved and can be grouped in the same classes as mammalian synaptotagmins, based on crucial amino acid residues needed for coordinating Ca^{2+} binding and determining phospholipid binding affinity. The only exception is synaptotagmin 1b, which lacks 34 amino acid residues in the C2B domain and is therefore unlikely to bind Ca^{2+} there. Additionally, the products of zebrafish *syt5a* and *syt5b* genes share identity with mammalian class 1 and 5 synaptotagmins. Zebrafish *syt1*, *syt2*, *syt5* and *syt7* paralogues are found in the zebrafish brain, eye, and retina, excepting *syt1b*, which is only present in the brain. The complementary expression pattern of the remaining paralogues in the retina suggests that *syt1a* and *syt5a* may underlie synchronous release and *syt7a* and *syt7b* may mediate asynchronous release or other Ca^{2+} dependent processes in different types of retinal neurons.

Key Words: retina, ribbon synapses, synaptotagmins, synaptic transmission, Ca^{2+} sensor, exocytosis

INTRODUCTION

Neuronal communication depends critically on Ca^{2+} (Katz & Miledi, 1965; Miledi, 1973) and Ca^{2+} sensors to regulate vesicle release and replenishment (Neher & Sakaba, 2008). The main Ca^{2+} sensors for exocytosis in the central nervous system of vertebrates are synaptotagmins, a large protein family that comprises about 15-17 members in humans (Sudhof, 2002; Craxton, 2004; Gustavsson & Han, 2009). Eight of these isoforms do not bind Ca^{2+} at all (Dai et al., 2004; Hui et al., 2005). The remainder show distinct Ca^{2+} sensitivities and phospholipid binding properties (Li et al., 1995; Sugita et al., 2002; Bhalla et al., 2005; Hui et al., 2005), as well as varying subcellular distributions (Sugita et al., 2001; Takamori et al., 2006; Dean et al., 2012) and cellular expression (Ullrich et al., 1994), potentially conferring neurons with a highly versatile repertoire in terms of synaptic gain, kinetics and transmission bandwidths (Hui et al., 2005; Xu et al., 2007; Chen & Jonas, 2017). However, only synaptotagmins 1, 2, 5 and 7 have been directly implicated in synaptic transmission (Xu et al., 2007; Bacaj et al., 2013).

In the retina, synapses often rely on continuous transmission. To cope with the demands of uninterrupted sensory inputs, the first retinal neurons - photoreceptors and bipolar cells - use specialized presynaptic contacts called ribbon synapses (Matthews & Fuchs, 2010). Ca^{2+} influx is continuous at such synapses and modulates neurotransmitter release around a mean tonic level. Retinal ribbon synapses presumably depend on multiple modes of release to relay information (Jackman et al., 2009; Oesch & Diamond, 2011), and because photoreceptor and bipolar cell ribbon synapses are distinct morphologically and physiologically (Matthews & Fuchs, 2010), it is plausible that they use different sets of synaptotagmins to control neurotransmitter release.

Although synaptotagmin 1 (Ullrich & Sudhof, 1994; Fox & Sanes, 2007; Grassmeyer et

al., 2019), synaptotagmin 2 (Ullrich et al., 1994; Fox & Sanes, 2007; Neumann & Haverkamp, 2013), synaptotagmin 3 (Butz et al., 1999; Berntson & Morgans, 2003) and synaptotagmin 7 (Luo et al., 2015) have been found in the retina, only synaptotagmin 1 (Grassmeyer et al., 2019) and synaptotagmin 7 (Luo et al., 2015) were reported to modulate different aspects of release in retinal ribbon-containing neurons. Furthermore, synaptotagmin 1 is thought to underlie alone both modes of synaptic transmission in mammalian cone but not rod photoreceptors (Grassmeyer et al., 2019). To date, the synaptotagmins controlling transient and sustained release in bipolar cells are unknown.

Zebrafish are a powerful animal model to study the nervous system in general and the retina specifically. First, there are multiple available toolboxes for the manipulation of genes of interest (Kawakami, 2007; Ablain et al., 2015; Kawakami et al., 2016; Niklaus & Neuhauss, 2017; Wierson et al., 2020). Second, zebrafish husbandry is relatively easy, mating has high yields and development is fast when compared to mammalian models. Lastly, zebrafish are experimentally accessible during development and display conserved retinal structures and main cell types (Gestri et al., 2012; Angueyra & Kindt, 2018).

The validity of this animal model for the study of retinal structure and function depends critically on the ability to draw analogies between the retinal anatomy, physiology and molecular biology of zebrafish and mammals. We therefore investigated the homology, expression, and distribution of synaptotagmins in the zebrafish retina. To do so, we took advantage of bioinformatic approaches to identify the genetic, sequence, and potential structural homology between zebrafish, mouse, and human synaptotagmins. We also assayed for the presence of synaptotagmins in the retina using RT-PCR and *in situ* hybridization.

We find that zebrafish synaptotagmins are well conserved and can be grouped in the

93 same classes as mammalian synaptotagmins. The only exception are synaptotagmins 5a and 5b,
 94 which are at an intermediate position between mammalian class 1 and 5 synaptotagmins.
 95 Zebrafish proteins retain key amino acid residues needed for coordinating Ca^{2+} binding and
 96 determining phospholipid binding affinity. Notably, while zebrafish synaptotagmin 1a is
 97 homologous to mammalian synaptotagmin 1 and is expressed in the retina, synaptotagmin 1b is
 98 not, and lacks the residues necessary to function as a Ca^{2+} sensor in its C2B domain. Zebrafish
 99 *syt1*, *syt2*, *syt5* and *syt7* paralogues are found in the zebrafish brain, eye and retina, except *syt1b*,
 100 which is only present in the brain. The remaining paralogues had complementary retinal
 101 expression patterns, suggesting that *syt1a* and *syt5a* may underlie synchronous release and *syt7a*
 102 and *syt7b* may mediate asynchronous release in different types of retinal neurons.

MATERIALS AND METHODS

Animals

All animal procedures were approved by the institutional animal care and usage committee (IACUC) at Stony Brook University and were in concordance with the guidelines established by the National Institutes of Health and by the Statement for the Use of Animals in Ophthalmic and Vision Research from The Association for Research in Vision and Ophthalmology.

Wild-type (WT) zebrafish (*Danio rerio*) were kept at 28.5 °C in aquaria under a 13:11 h light to dark cycle and fed artemia and GEMMA micropellets twice a day. The WT strain used for all experiments was a hybrid WT background consisting of Tubingen long-fin crossed to Brian's WT. Adult animals (age 8-12 months) of either sex were dark-adapted for 2 h and euthanized by immersion in 1 mM tricaine methanesulfonate (SIGMA, cat. no. A5040) in pH 7.0 buffered system water prior to experiments.

RT-PCR analysis

Zebrafish eyes and brains were dissected from adult animals and rapidly frozen by immersion in a mixture of ethanol and dry ice. Total RNA was extracted by the method of Cathala *et al.* (Cathala et al., 1983) and stored in ethanol at -80 °C until use.

Reverse transcription using Superscript IV reverse transcriptase (Invitrogen) was performed according to the manufacturer's protocol. In brief, 250-500 µg of total DNase I-treated total RNA was added to nuclease-free water to a final volume of 11 µL. Subsequently, 1 µL of random hexamers and 1 µL of 10 mM dNTP mix were added to the RNA solution, heated to 65 °C for 5 minutes and then incubated on ice for one minute. After addition of 4 µL of

5x buffer, 1 μ L 10 mM DTT and 1 μ L RNase inhibitor, the sample was incubated at 23°C and 1 μ L of Superscript IV reverse transcriptase was added; synthesis was performed for 1 hour at 55 °C. The enzyme was inactivated at 80 °C for 10 minutes. All reagents were obtained from Invitrogen/Thermo Fisher.

For conventional PCR, 1 μ L reverse-transcribed cDNA was added to 25 μ L of PCR buffer and reactants, using EconoTaq DNA polymerase (Lucigen). The standard amplification protocol consisted of 95°C for 5 minutes, followed by 35 cycles of 95, 55 and 72°C for 1 minute each, and ending with a 72°C-extension for 5 minutes. PCR primers were designed based on GenBank sequences for zebrafish synaptotagmin isoforms. Type-specific primers were designed such that each one amplified a unique fragment either near the ATG start of the respective coding sequence (CDS) or a larger unique fragment in the untranslated (UTR) region. The design of isoform-specific primers for synaptotagmin genes is complicated by the fact that nearly 40% of the gene is comprised of homologous C2A and C2B domains. The primers shown in **Table 1** were designed to amplify the largest unique fragment for each gene. DNA fragments were confirmed by sequencing and alignment. cDNA products of correct sizes were gel-purified and subcloned into the pGEM-T Easy cloning vector (Promega). Selected clones were sequenced on an automatic DNA sequencer and aligned to GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>, RRID:SCR_002760) data to verify the integrity of DNA fragments.

***In situ* hybridization**

Zebrafish eyes were dissected and fixed in 4% paraformaldehyde in phosphate-buffered saline (PBS) overnight at 4 °C. The fixed tissue was washed in PBS and cryoprotected in 30% sucrose before embedding in Shandon M1 media (Thermo Fisher Scientific) and stored at -80 °C until

use. On day one of *in situ* experiment, 25- μ m sections were placed on Superfrost Plus microscope slides (Thermo Fisher Scientific) and dried at 55°C for a minimum of 3 hours. This tissue was then immediately used for *in situ* hybridization.

Table 1. PCR primers.

Zebrafish gene	Direction	Sequence	Location (bp)*	Amplicon Size (bp)	Accession #
<i>syt1a</i>	forward	cattgcattcgttctgtgttac	186-208	420	NM_001327829.1
	reverse	acaggattcaacgtcttcggtggacc	605-579		
<i>syt1b</i>	forward	cgtctcggttcacacagtcttg	209-230	412	NM_001089461.1
	reverse	gtcggctcagtggtttgcgatgaact	620-594		
<i>syt2a</i>	forward	atgaagtggatgtgtgaagaagaagc	1-28	196	XM_691185.7
	reverse	gtggaagctgtccatctcattgagg	196-171		
<i>syt2b</i>	forward	ccagtggcaccctctccagcagg	1242-1264	219	XM_005166154.4
	reverse	caatctatgtggcgaacgtgtg	3'-UTR +150-129		
<i>syt5a</i>	forward	cctaaccacaagatcagaatgcccatg	139-165	294	NM_001103137.1
	reverse	gccaacaatgagctgatttcagt	432-408		
<i>syt5b</i>	forward	cttcattgccgatgtgggcttagg	259-284	309	NM_001020546.2
	reverse	ttgaagaatccaactatcagctggg	567-542		
<i>syt7a</i>	forward	gactacgtcccgtcagcagg	558-579	502	XM_021470598.1
	reverse	gaggaaatgctcgtctgcgtct	1061-1041		
<i>syt7a_splice**</i>	forward	atgtatctcaacaggaggaggagt	1-26	264	See Appendix
	reverse	cggctgacccttaggactgttgg	264-242		
<i>syt7b</i>	forward	ccagacgagagccaccggcgg	529-549	407	NM_001245957.1
	reverse	tcataatccagcacttgcagg	935-915		

* For primers located within the CDS, 1 = ATG. For 3'-UTR primers, 1 = first nucleotide after stop.

** Splice variant missing nucleotides 219-993 identified in zebrafish eye (see Results for more information and Appendix for sequenced fragment); numbering for the primers is therefore unique to this sequence. *Syt7a_splice* nucleotide 264 corresponds to 1038 in XM_021470598.

Isotype-specific probes for zebrafish synaptotagmins present in the eye were synthesized in sense and antisense directions from cDNA obtained by RT-PCR. Riboprobe lengths and locations are listed in **Table 2**. All probes were localized in the 5' end of the gene or 3'-UTR, where sequences are divergent. Synthesis, hybridization, and detection of digoxigenin-labeled probes were carried out according to manufacturer's protocol (Roche), using anti-digoxigenin

antibody conjugated to alkaline phosphatase for detection. A SNAP25b riboprobe (a gift from H. Sirotkin, Moravec et al., 2016) was used as a positive control for tissue and reagents. Slides were viewed and photographed under 20x or 40 x magnification (Zeiss PlanNeofluor objectives, 0.5 and 0.75 N.A., respectively) in a microscope equipped for brightfield (Zeiss Axioskop) attached to a color camera (Infinity 3, Lumenera Corporation, Ottawa, Canada) controlled by Infinity Capture software (Lumenera Corporation, Ottawa, Canada). Photomicrographs of sense and anti-sense pairs generated from the same experiment were acquired as TIFF files at 1936x1456 pixels and post-processed jointly with Adobe Photoshop (<https://www.adobe.com/products/photoshop.html>, RRID:SCR_014199) to correct contrast and brightness.

Table 2. Riboprobes for *in situ* hybridization.

Zebrafish Gene	Direction	Sequence	Location (bp)*	Probe Length (bp)	Accession #
<i>syt1a</i>	forward	ctgcgccaccgctgcgcctgagg	32-321	289	NM_001327829.1
	reverse	gtccttcacgtcctcatgttgatgg			
<i>syt1a</i>	forward	gtgtaataacctcagtgatataagg	3'-UTR +71-1124	1053	NM_001327829.1
	reverse	atcagtcacacgtaacgagaagtaagcc			
<i>syt2a</i>	forward	atgaagtgggaatgtgtgaagaagaagc	1-196	196	XM_691185.7
	reverse	gtggaagcttgctcatctcattgagg			
<i>syt5a</i>	forward	tcaatcttattcatcctgtcagtgaggag	3'-UTR +63-309	247	NM_001103137.1
	reverse	catcttgagtcacagaacaagaggcc			
<i>syt7a</i>	forward	gcacaagactaactgtgctccaggcgg	3'-UTR +64-756	693	XM_021470598.1
	reverse	gactcatgtttacctcttgagaagagg			
<i>syt7b</i>	forward	atgcacctaatacgggaggagcagg	1-202	202	NM_001245957.1
	reverse	gtgacatgttgaggaggaaactctttgg			

* For probes located within the CDS, 1 = ATG. For 3'-UTR probes, 1 = first nucleotide after stop.

Bioinformatics

The orthology between the zebrafish *syt* genes and their human and mouse counterparts was determined by searching the Zebrafish Information Network (Zfin, <http://zfin.org>,

RRID:SCR_002560), NCBI Entrez Gene (<http://www.ncbi.nlm.nih.gov/gene>,
RRID:SCR_002473), NCBI Homologene (<http://www.ncbi.nlm.nih.gov/homologene>,
RRID:SCR_002924) and Ensembl (<http://www.ensembl.org/>, RRID:SCR_002344) databases.
Additionally, the NCBI BLAST Suite (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>,
RRID:SCR_004870) was used to align and compare gene, transcript or protein sequences of
interest and to investigate possible interactors with *in situ* probes or help determine homology.

In addition to percent identity, we also used two scores in Ensembl to evaluate homology:
Gene Order Conservation (GOC) and the Whole Genome Alignment (WGA) scores. The GOC
score, which ranges from 0 to 100, indicates how many of the four closest neighbors of a gene
match between orthologous pairs, and assumes that genes that are descended from the same gene
are likely to be part of a block of genes, all in the same order, in both species. The WGA score
also varies from 0 to 100 and indicates how well query and target genomic regions align to each
other. It assumes that genes which are orthologous to each other will fall within genomic regions
that can be aligned to one another (i.e. percent synteny).

For phylogenetic analysis, protein sequences from NCBI Protein
(<http://www.ncbi.nlm.nih.gov/protein>, RRID:SCR_014312) databases were aligned with Clustal
Omega (<https://www.ebi.ac.uk/Tools/msa/clustalo/>, RRID:SCR_001591, Sievers et al., 2011) or
MUSCLE (<https://www.ebi.ac.uk/Tools/msa/muscle/>, RRID:SCR_011812, Edgar, 2004), both at
default settings for highest accuracy. After alignment, regions with gaps or misaligned were
removed with GBlocks (http://molevol.cmima.csic.es/castresana/Gblocks_server.html,
RRID:SCR_015945, Castresana, 2000) to eliminate ambiguity. A stringent set of parameters was
chosen to eliminate divergent regions and improve reliability: (i) no gaps allowed; (ii) blocks
after gap cleaning ≥ 10 residues; (iii) contiguous nonconserved residues ≤ 4 ; (iv) sequences for

conserved position ≥ 28 ; and (v) sequences for a flanking position ≥ 45 . After curation, 117 amino acids were selected for subsequent evaluation.

Results were imported into JalView (<https://www.jalview.org/>, RRID:SCR_006459, Waterhouse et al., 2009) for visualization and calculation of neighbor joining phylogenetic trees using the BLOSUM62 matrix. Phylograms were imported as Newick files into iTOL (Interactive Tree of Life, <https://itol.embl.de>, RRID:SCR_004473, Ciccarelli et al., 2006; Letunic & Bork, 2019) for manipulation and annotation, and exported as vector-based graphics (i.e. svg or eps files) for final editing in Canvas X (<http://www.canvasgfx.com/en/products/canvas-15>, RRID:SCR_014312).

Prediction of secondary structure and 3-dimensional modeling of zebrafish synaptotagmin sequences was performed with RaptorX (<http://raptorx.uchicago.edu/>, RRID:SCR_018118). Models were visualized and aligned to known crystal structures of mammalian synaptotagmins obtained from the Research Collaboratory for Structural Bioinformatics Protein Data Bank (RCSB PDB, <http://www.rcsb.org/pdb/>, RRID:SCR_012820) using PyMOL (<http://www.pymol.org/>, RRID:SCR_000305) and exported as PNG files at 5000 x 5000 pixel resolution for subsequent editing in Canvas X.

RESULTS

Homology of zebrafish synaptotagmins to their mammalian counterparts

Due to a whole genome duplication that happened at the beginning of the teleost lineage some 320-350 million years ago (Amores et al., 1998; Glasauer & Neuhauss, 2014; Pasquier et al., 2016), zebrafish in general have two synaptotagmin (*syt*) paralogue genes, named *a* and *b*, for each mammalian counterpart. The exceptions are *syt3*, *syt4*, *syt8* and *syt10* (Craxton, 2004, 2010), which have only one gene, probably as a result of deleterious mutations of one of the paralogues that led to loss of function (Postlethwait et al., 2000; Glasauer & Neuhauss, 2014; Pasquier et al., 2016). At present it is unknown whether the remaining zebrafish *syt* genes still act as Ca^{2+} sensors, or whether they have acquired new functions during evolution.

To begin to address this issue, we compiled the genomic identities between zebrafish and human orthologues in the Ensembl database (**Figure 1A**) and compared them to those between mouse and human orthologue pairs (**Figure 1B**). For this and subsequent analyses, we concentrated on synaptotagmins 1 to 10, since synaptotagmins 11-15 do not bind Ca^{2+} (von Poser et al., 1997; Fukuda, 2003a; Bhalla et al., 2008; Craxton, 2010), and there is uncertainty as to whether synaptotagmins 16-17 should be considered members of the synaptotagmin family because they lack a transmembrane domain (Gustavsson & Han, 2009; Wolfes & Dean, 2020). The nomenclature between zebrafish and mammalian orthologue pairs is consistent (i.e., the human orthologue for zebrafish *syt1a* is human SYT1; Craxton, 2004), because zebrafish *syt* genes were named after extensive examination of their syntenic relations and the amino acid sequences of their products (Craxton, 2004, 2010).

Figure 1. Genomic homology between zebrafish, mouse and human synaptotagmin 1 to 10.

(A, B) The percent identity listed in the Ensembl database between zebrafish and human (A) and between mouse and human (B) orthologues (human *SYT* gene numbers are indicated above the bars). *Target*: % of the human sequence matching the zebrafish or mouse sequence; *Query*: % of the zebrafish or mouse sequence matching the human sequence.

(A) The percent identity between zebrafish and human is above 50% (dashed red line), except for *syt7a* (*query*: 62%; *target*: 49%). The only zebrafish gene with >80% identity to its human counterpart (dashed grey line) is *syt1a* (*query*: 82.5%; *target*: 83%). Empty spaces in the graph reflect that zebrafish *syt2b* and *syt8* are not mapped in Ensembl, and that the database does not list any human orthologue gene matches for zebrafish *syt5a* and *syt5b*. Zebrafish has duplicate genes for all synaptotagmins, except for *syt3*, *syt4*, *syt8* and *syt10*.

(B) The percent identity between mouse and human is in most cases above 80% (dashed grey line), except for *Syt7* (*query*: 70%; *target*: 83%) and *Syt8* (*query*: 74%; *target*: 73%).

(C, D) Gene Order Conservation (*GOC*) and Whole Genome Alignment (*WGA*) scores, which vary from 0 to 100, are an additional means to evaluate homology (see *Materials & Methods*). Both scores are much lower for the zebrafish-human (C) than for mouse-human (D) orthologues.

Although the percent identity between zebrafish and human orthologues (Figure 1A) is

lower than that between mouse and human counterparts (**Figure 1B**), in almost all cases identity is >50%, the only exception being *syt7a*. GOC and WGA scores were much lower for zebrafish-human pairs (**Figure 1C**) than those between mouse and human orthologue genes (**Figure 1D**).

Some gene pairs are missing in **Figure 1A** and **Figure 1C**: zebrafish *syt2b* and *syt8* genes have not been mapped in Ensembl, and although *syt5a* and *syt5b* are mapped, human or murine orthologue searches for these paralogues did not yield any hits in either Ensembl or NCBI. Indeed, if these genes are true orthologues as suggested (Craxton, 2004, 2010), they must have diverged considerably, since the amino acid similarity between human synaptotagmin 5 and zebrafish synaptotagmins 5a and 5b is low (between 49 and 51% identity for synaptotagmin 5a, depending on the isoforms compared, and 53% for synaptotagmin 5b). Aligning the translation products of the *syt5a* and *syt5b* paralogues with those of human SYT genes with MUSCLE yielded human synaptotagmin 2 as the closest match for zebrafish synaptotagmin 5a (85% and 61% identity, respectively), and human synaptotagmin 1 as the closest match for zebrafish synaptotagmin 5b (61% identity) and synaptotagmin 8 (between 53 and 59% identity, depending on the isoforms compared). That said, the manually curated orthology from the Zfin database indicates human *SYT5* and murine *Syt5* as the orthologues for *syt5a* and *syt5b* based on both synteny and amino acid sequence, albeit with no quantitative measure of homology.

Presumably, such sequence divergence arises because of the accumulation of mutations over many generations, which may or may not impair protein function (Glasauer & Neuhaus, 2014). On the other hand, one expects functional portions of the translation products to be more conserved amongst different species than those that are not crucial for protein function. Synaptotagmins are a fairly conserved family of proteins that share a common structure (**Figure 2A**), consisting of a short extracellular/intravesicular N-terminus, a transmembrane domain, and

an intracellular portion formed by an intracellular linker, two tandemly arranged C2 domains interspersed by a shorter linker sequence, and a C-terminus (Sudhof, 2002; Wolfes & Dean, 2020). The C2 domains, named C2A and C2B, are responsible for Ca^{2+} , phospholipid and SNARE binding, and are together the main functional components of these proteins.

Figure 2. General synaptotagmin structure.

- (A) Schematic drawing of human synaptotagmin 1. All synaptotagmins share the same basic structure: an intravesicular or extracellular N-terminus (N) of variable length; a transmembrane domain (TM) that anchors the protein to the vesicular or plasma membrane; a variable linker region (L1); two almost identical Ca^{2+} -binding domains (C2A and C2B) arranged in tandem and coupled by a second shorter linker sequence (L2); and a C-terminus (C). The number of amino acid residues of the non-variable regions for human synaptotagmin 1 are shown. The loops responsible for Ca^{2+} binding in the C2 domains (loops 1 and 3) are shown in pink. Drawing not to scale.
- (B) Ribbon model of the 3-dimensional structure of zebrafish synaptotagmin 1a (rainbow colors) generated with RaptorX from the amino acid sequence (isoform 1, accession number in **Table 3**) and superposed on the crystal structure of the C2 domains of human synaptotagmin 1 (shown in grey, PDB: 2R83; Fuson et al., 2007). Loops responsible for Ca^{2+} binding in the C2 domains (loops 1 and 3) of the human synaptotagmin are shown in pink.

The modeled 3-D structure of the C2 domains of zebrafish synaptotagmin 1a (depicted in

rainbow colors in **Figure 2B**) align to the crystal structure of human synaptotagmin 1 (in grey, **Figure 2B**) (PDB: 2R83, Fuson et al., 2007). This is highly suggestive of functional conservation during evolution. To look more closely at the adaptive radiation of the functional domains of zebrafish synaptotagmins, we examined the phylogenetic relations of conserved amino acid sequences in the C2 domains of zebrafish and human synaptotagmins 1 to 10 (**Figure 3**). The NCBI accession #s used for this analysis are compiled in **Table 3** and **Table 4**; sequences from NCBI were chosen because Ensembl is incomplete (i.e. no zebrafish *syt2b* or *syt8* genes and products).

Zebrafish and human orthologue proteins cluster in the same general classes (**Figure 3**) (Sugita et al., 2002). Synaptotagmins 5a and 5b constitute the only exceptions because they fall between class 1 and 5 and could therefore belong to either (overlapping brackets in **Figure 3**). The properties of these classes in mammals and relevant literature are summarized in **Table 5**. Class 1 proteins (synaptotagmins 1 and 2) are fast, low sensitivity and low affinity Ca^{2+} sensors located in vesicles and involved in synchronous synaptic transmission. Class 2 proteins (synaptotagmin 7) are also involved in synaptic transmission but anchor to the plasma membrane and have higher affinity and sensitivity to Ca^{2+} than class 1 synaptotagmins. Class 3 proteins (synaptotagmins 3, 6, 9 and 10) are mostly anchored to the plasma membrane, and regulate endocytosis, postsynaptic receptor trafficking and non-synaptic secretion. Class 4 proteins (synaptotagmins 4 and 11) are insensitive to Ca^{2+} and modulate the activity of other synaptotagmins. Class 5 proteins (synaptotagmin 5) are also fast, low sensitivity/low affinity Ca^{2+} sensors in mammals, but with slower kinetics than class 1 proteins. Finally, class 6 proteins (synaptotagmin 8) are also unable to bind Ca^{2+} and act as inhibitory proteins, as class 4 synaptotagmins.

Figure 3. Phylogenetic comparison of zebrafish and human synaptotagmins. Unrooted phylogram of zebrafish and human synaptotagmins 1 to 10 based on 117s conserved amino acid residues from the C2 domains. Proteins are either shown in color (zebrafish) or in grey (human). Class division indicated by brackets, after (Sudhof, 2002). NCBI accession numbers are shown at the end of the branches and represent different isoforms of each protein. Numbers starting with “NP” are validated sequences (shown in bold for zebrafish proteins, see **Table 3** and **Table 4** for NCBI status of each sequence), while numbers starting with “XP” are sequences predicted by computer algorithms from mRNA or genomic sequences, which still await validation. Only validated or reviewed human sequences were used, while for zebrafish, all isoforms listed at the NCBI database were used due to the limited number of validated sequences available. Most zebrafish synaptotagmins fall within the corresponding mammalian class, except synaptotagmins 5a and 5b. These occupy an intermediate location between class 5 and class 1 synaptotagmins (overlapping brackets). Scale bar corresponds to amino acid similarity.

Our analysis (**Figures 1-3**) and published data on genomic homology (Craxton, 2004, 2010) point to functional conservation of most synaptotagmin proteins in zebrafish. To address this question more rigorously, we looked at the Ca^{2+} binding pockets of the C2A and C2B

domains. We focused specifically on whether zebrafish synaptotagmins could act as Ca²⁺ sensors by examining the amino acid residues responsible for Ca²⁺ coordination in the C2 domains.

Table 3. Accession data for the zebrafish synaptotagmins used for phylogenetic analysis.

Species	Gene	Isoform	aa*	NCBI Accession #**	Status	UniProt #	Ensembl #
<i>Danio rerio</i>	1a	1	419	NP_001314758.1	validated	Q5TZ27	ENSDARP00000141860.1
<i>Danio rerio</i>	1a	X1	419	XP_017210484.1	model	Q5TZ27	-
<i>Danio rerio</i>	1a	X2	416	XP_009298648.1	model	-	-
<i>Danio rerio</i>	1b	1	388	NP_001082930.1	provisional	A3KQ92	ENSDARP00000092016.3
<i>Danio rerio</i>	1b	X1	380	XP_021323234.1	model	-	-
<i>Danio rerio</i>	2a	X1	430	XP_696277.2	model	F1R5C0	ENSDARP00000121161.1
<i>Danio rerio</i>	2a	X2	428	XP_005174169.1	model	-	-
<i>Danio rerio</i>	2a	X3	364	XP_021325567.1	model	-	-
<i>Danio rerio</i>	2b***	X1	436	XP_005166211.1	model	-	-
<i>Danio rerio</i>	2b***	X2	433	XP_688092.3	model	-	-
<i>Danio rerio</i>	3	X1	566	XP_005164031.1	model	I3IS33	ENSDARP00000125697.1
<i>Danio rerio</i>	3	X2	559	XP_021327105.1	model	-	-
<i>Danio rerio</i>	4	1	439	NP_956242.1	provisional	Q6PBU6	ENSDARP00000045478.6
<i>Danio rerio</i>	5a	1	405	NP_001096607.1	provisional	A7MCG9, F1QX49	ENSDARP00000122114.1
<i>Danio rerio</i>	5a	X1	405	XP_009297805.1	model	-	-
<i>Danio rerio</i>	5b	1	446	NP_001018382.2	validated	F1QCC9, Q504A0	ENSDARP00000012595.6
<i>Danio rerio</i>	6a	X1	537	XP_017209198.1	model	F1QA93	ENSDARP00000100090.4
<i>Danio rerio</i>	6a	X2	417	XP_009295199.1	model	E9QFE9	ENSDARP00000120056.1
<i>Danio rerio</i>	6b	X1	504	XP_021335668.1	model	F1QD38	ENSDARP00000048364.5
<i>Danio rerio</i>	7a****	X1	653	XP_021326273.1	model	X1WDZ6	ENSDARP00000128447.1
<i>Danio rerio</i>	7b	1	488	NP_001232886.1	provisional	D4P8S0	ENSDARP00000136207.1
<i>Danio rerio</i>	7b	X1	380	XP_005166426.1	model	-	-
<i>Danio rerio</i>	8*****	1	397	NP_001352218.1	validated	FQ976913	-
<i>Danio rerio</i>	8*****	2	365	NP_001352219.1	validated	FQ976913	-
<i>Danio rerio</i>	8*****	X3	364	XP_021326392.1	model	-	-
<i>Danio rerio</i>	8*****	X4	272	XP_021326393.1	model	-	-
<i>Danio rerio</i>	9a	X1	547	XP_001336058.4	model	E7F1D9	ENSDARP00000003622.7
<i>Danio rerio</i>	9b	1	517	NP_001003985.1	provisional	Q68EH6	ENSDARP00000049991.4
<i>Danio rerio</i>	10	1	553	NP_001076311.1	provisional	Q5RI28, Q5RG04	ENSDARP00000123473.2
<i>Danio rerio</i>	10	X1	606	XP_017210538.2	model	-	-

* Isoform length varies among the different databases. We list here the number of residues for the NCBI accession #s used for the phylogenetic tree.

** Some isoforms are duplicated in the NCBI database and have therefore two or more accession #s. For this analysis, we used one accession number per isoform. Only data from NCBI was used; matching accession # for UniProt and Ensembl were obtained from the NCBI site when available.

*** Called synaptotagmin-2-like (LOC559637) in the NCBI database, but synaptotagmin 2b at Zfin, after manual curation (Craxton, 2004).
 **** Synaptotagmin 7a has splice variants not listed in NCBI database. This is the longest version (syt7a_202 in Ensembl).
 ***** Synaptotagmin 8 was formerly called synaptotagmin-1-like.

Table 4. Accession data for the human synaptotagmins used for phylogenetic analysis.

Species	Gene	Isoform	aa*	NCBI Accession #**	Status	UniProt #	Ensembl #
<i>Homo sapiens</i>	1	1	422	NP_001129277.1	validated	P21579	ENSP00000376932.3
<i>Homo sapiens</i>	1	2	419	NP_001278830.1	validated	J3KQA0	ENSP00000391056.2
<i>Homo sapiens</i>	2	1	419	NP_001129976.1	reviewed	Q8N9I0	ENSP00000356236.1
<i>Homo sapiens</i>	3	1	590	NP_001153800.1	validated	Q9BQG1	ENSP00000468982.1
<i>Homo sapiens</i>	4	1	425	NP_065834.1	validated	Q9H2B2	ENSP00000255224.2
<i>Homo sapiens</i>	5***	1	386	NP_003171.2	validated	O00445, A0A024R4N8	ENSP00000346265.2
<i>Homo sapiens</i>	5***	2	382	NP_001284703.1	validated	O00445	ENSP00000465576.1
<i>Homo sapiens</i>	6	a	510	NP_001240701.1	reviewed	-	ENSP00000476396.1
<i>Homo sapiens</i>	6	b	425	NP_001257734.1	reviewed	Q5T7P8, I6L9C3	ENSP00000358560.1
<i>Homo sapiens</i>	6	c	528	NP_001353153.1	reviewed	-	-
<i>Homo sapiens</i>	6	d	503	NP_001353154.1	reviewed	-	-
<i>Homo sapiens</i>	6	e	443	NP_001353155.1	reviewed	-	-
<i>Homo sapiens</i>	7	1	478	NP_001238994.1	reviewed	O43581	ENSP00000444201.1
<i>Homo sapiens</i>	7	2	403	NP_004191.2	reviewed	O43581	ENSP00000263846.4
<i>Homo sapiens</i>	7	3	447	NP_001287702.1	reviewed	O43581	ENSP00000444568.1
<i>Homo sapiens</i>	7	4	686	NP_001352738.1	reviewed	-	ENSP00000439694.1
<i>Homo sapiens</i>	7	5	479	NP_001357139.1	reviewed	-	-
<i>Homo sapiens</i>	7	6	383	NP_001357140.1	reviewed	-	-
<i>Homo sapiens</i>	8	1	402	NP_001277261.2	reviewed	-	-
<i>Homo sapiens</i>	8	2	401	NP_612634.4	reviewed	Q8NBV8	ENSP00000371394.3
<i>Homo sapiens</i>	8	3	403	NP_001277262.2	reviewed	-	-
<i>Homo sapiens</i>	8	4	400	NP_001277263.2	reviewed	-	-
<i>Homo sapiens</i>	9	1	491	NP_783860.1	validated	Q86SS6	ENSP00000324419.6
<i>Homo sapiens</i>	10	1	523	NP_945343.1	validated	Q6XYQ8	ENSP00000228567.3

* Isoform length varies among the different databases. We list here the number of residues for the NCBI accession #s used for the phylogenetic tree.

** Some isoforms are duplicated in the NCBI database and have therefore two or more accession #s. For this analysis, we used one accession # per isoform. Only data from NCBI was used; matching accession # for UniProt and Ensembl were obtained from the NCBI site when available.

*** Mammalian SYT5 genes and respective products are sometimes called SYT9 and vice-versa (Li et al., 1995; Sugita et al., 2002; Chen & Jonas, 2017). We here adopt the nomenclature convention of the NCBI, Ensembl and Zfin databases, after (Craxton & Goedert, 1995; Hudson & Birnbaum, 1995).

375 **Table 5. Characteristics of neuronal synaptotagmins 1 to 10 in mammals.**

Class*	Syt	S _(Ca) **	PL_b(Ca)***	Brain Elsewhere****	Location	Target	Process	Description
1	1	low ^{1,2}	yes ^{3,4}	14.8	synapses ⁵	synaptic vesicles ^{5,6}	exocytosis ^{7,8}	synchronous release ^{9,10}
1	2	low ²	yes ^{3,4}	1.1	synapses ⁵	synaptic vesicles ⁵	exocytosis ¹⁰	synchronous release ¹⁰
2	7	high ^{1,2}	yes ^{3,4}	1.1	synapses ¹¹ , axons ⁵	plasma membrane ¹¹	exocytosis ^{9,12}	asynchronous release ^{9,12} synaptic facilitation ¹³ vesicle replenishment ¹⁴ suppression of spontaneous release ¹⁵
3	3	high ²	yes ^{3,4}	6.2	synapses ¹⁶ , dendrites ^{5,17}	plasma membrane ¹⁶	endocytosis ¹⁷	postsynaptic GluR trafficking ¹⁷
3	6	high	no ³ , yes ⁴	3.5	synapses ¹⁶ , axons/dendrites ⁵	plasma membrane ¹⁶		
3	9****	high ²	yes ^{3,4}	2.1	axons ⁵	plasma membrane		
3	10	high ²	no ⁵ , yes ⁴	0.9	soma, axons ⁵	plasma membrane ⁵	exocytosis ⁸	IGF-1 secretion ⁸
4	4	none ^{4,18-20}	no ^{3,4,20}	3.4	axons/dendrites ⁵		modulation ⁴	inhibition of synaptotagmin-1-mediated fusion ⁴
5	5*****	intermediate ¹ , low ¹⁰	yes ⁴	9.5	synapses, axons/dendrites ⁵	synaptic vesicles	exocytosis ¹⁰	synchronous release ¹⁰
6	8	none ^{4,5,21}	no ^{3,4}	0.0001	not detected in brain ^{5,22-24} soma ²¹	cytosol ²¹	modulation ⁴	inhibition of synaptotagmin-1-mediated fusion ⁴

* *There are several divisions for the synaptotagmin family in literature. The one we adopt here is from Sudhof (2002).*

** *Ca²⁺ sensitivity.*

378 *** *Ca²⁺-dependent phospholipid binding.*

379 **** *The ratio of RNA reads per kilobase per million reads placed (RPKM) from brain vs. the maximal RPKM elsewhere in the body, calculated from quantitative transcriptome analysis (RNA-Seq) of 27 human tissue samples from 95 individuals (Fagerberg et al., 2014). Most synaptotagmins are more expressed in the brain (i.e. RPKM ratio >1) than elsewhere in the body, except for synaptotagmins 8 and 10, which have very low expression in the brain (RPKM <1).*

382 ***** *Called synaptotagmin 5 in some studies (Li et al., 1995; Dean et al., 2012). Because of this confusion in the literature (Haberman et al., 2003), data from articles that did not make clear which convention was used (NCBI/Ensembl or naming based on papers from Thomas Sudhof's group) were not included in this table.*

385 ***** *Called synaptotagmin 9 in some studies (Bhalla et al., 2005; Xu et al., 2007; Dean et al., 2012). Data from articles that did not make clear which convention was used were not included in this table.*

387 Refs.: ¹(Bhalla et al., 2005); ²(Sugita et al., 2002); ³(Li et al., 1995); ⁴(Bhalla et al., 2008); ⁵(Dean et al., 2012); ⁶(Takamori et al., 2006); ⁷(Geppert et al., 1994);

388 ⁸(Cao et al., 2013); ⁹(Bacaj et al., 2013); ¹⁰(Xu et al., 2007); ¹¹(Sugita et al., 2001); ¹²(Luo et al., 2015); ¹³(Jackman & Regehr, 2017); ¹⁴(Liu et al., 2014);

389 ¹⁵(Luo & Sudhof, 2017); ¹⁶(Butz et al., 1999).

Conservation of crucial residues for Ca²⁺ binding

The interaction between synaptotagmins and Ca²⁺ is electrostatic in nature and creates an electrical switch for phospholipid binding (Ubach et al., 1998). In mammalian synaptotagmin 1, six negatively charged amino acid residues located in loops 1 and 3 of the C2A domain coordinate together the binding of three Ca²⁺ ions (**Figure 4A**), and five negative residues in loops 1 and 3 of the C2B domain coordinate the binding of two Ca²⁺ ions (**Figure 4B**). When not bound to Ca²⁺, the negative surface charge of these loops likely prevents close contact between the C2 domains and negatively charged phospholipid membranes. Binding of Ca²⁺ neutralizes these charges and promotes direct interaction between synaptotagmins and the cell membrane (or vesicular membrane, in the case of membrane-bound synaptotagmins, **Table 5**). Since the amino acid composition of these loops controls both the charge (Ubach et al., 1998) and exact shape (Dai et al., 2004; Qiu et al., 2017) of the Ca²⁺ binding pocket, it is therefore critical for the role of synaptotagmins as Ca²⁺ sensors for exocytosis.

C2 domains in Class 1 synaptotagmins

Ca²⁺ binding to the C2B domain of synaptotagmin 1 is crucial for synchronous release (Mackler & Reist, 2001; Mackler et al., 2002; Nishiki & Augustine, 2004; Shin et al., 2009; Bacaj et al., 2013; Lee et al., 2013), probably because the C2B domain has significantly higher sensitivity to Ca²⁺ (Bradberry et al., 2020) and phospholipid-binding activity (Bai et al., 2004; Li et al., 2006; van den Bogaart et al., 2012; Bradberry et al., 2020) than the C2A domain in this protein. Indeed, all mutations in synaptotagmin 1 that cause diseases in humans target the C2B domain (Baker et al., 2018; Bradberry et al., 2020), while removal of the residues critical for Ca²⁺ binding in the C2A domain is relatively innocuous (Stevens & Sullivan, 2003).

synaptotagmin 1a and 1b (*dre_1a_1* and *dre_1b_1*, accession numbers in **Table 3**) with those of isoforms 1 and 2 of human (*hsa_1_1* and *hsa_1_2*, accession numbers in **Table 4**), rat (*rno_1_1*: NCBI# NP_001028852.2) and mouse (*mmu_1_1*: NCBI# NP_001239270.1 and *mmu_1_2*: NCBI# NP_001239271.1) synaptotagmin 1. Locations of the β -strands in the human sequence are highlighted in grey, after (Fuson et al., 2007); loops 1 and 3 are highlighted in pink, and negative residues responsible for Ca^{2+} binding are in blue (aspartates) or yellow (serine). Consensus between sequences is indicated below the alignments: “*”, identical residue; “:”, conservative replacement; “.”, semi-conservative substitution. All charged residues that coordinate Ca^{2+} in the C2A domain of mammalian synaptotagmins are conserved in both zebrafish paralogues, but 34 residues are missing in a region that includes two β sheets ($\beta 6$ and $\beta 7$) and loop 3 from the C2B domain of zebrafish synaptotagmin 1b (highlighted in red). This paralogue, therefore, is likely unable to bind Ca^{2+} .

Zebrafish synaptotagmin 1a has all 11 residues needed to coordinate Ca^{2+} binding at the same locations in both C2 domains as mammalian synaptotagmin 1, which render the 3-dimensional arrangement of the Ca^{2+} binding pockets very similar to that of its mammalian orthologues (**Figure 4C-D**). Alignment of the amino acid sequences of both C2 domains of zebrafish (*dre*), human (*hsa*), rat (*rno*) and mouse (*mmu*) orthologues confirms that these regions are highly conserved in synaptotagmin 1a (**Figure 4E-F**). This paralogue is therefore likely to function as a Ca^{2+} sensor. On the other hand, synaptotagmin 1b lacks a stretch of 34 amino acids around loop 3 of the C2B domain (area highlighted in red in the alignment in **Figure 4F**). This truncation probably renders this paralogue unable to coordinate Ca^{2+} binding and to function as a sensor for exocytosis.

Table 6 summarizes the homology of the Ca^{2+} binding motifs in both C2 domains between zebrafish, human, rat and mouse orthologues. Interestingly, the Ca^{2+} binding motifs of synaptotagmins 2a and 2b are more homologous to mammalian synaptotagmin 1 than to synaptotagmin 2. The C2B domains of the zebrafish proteins are identical to those of mammalian synaptotagmin 1, while mammalian synaptotagmin 2 has a conservative aspartate (D)-to-glutamate (E) substitution in the C2B domain. Such a substitution is also present in mammalian synaptotagmin 3 (and in all class 3 orthologues, **Table 6**) and may lead to collapse of the last

Ca²⁺ binding site of this domain (Sutton et al., 1999). Therefore, zebrafish synaptotagmins 2a and 2b could probably bind the same number of Ca²⁺ ions as synaptotagmins 1 and 1a, which is one Ca²⁺ ion more than mammalian synaptotagmin 2. Also, since the C2B domain is more important for synaptotagmin 1 function in synchronous release (Mackler et al., 2002; Nishiki & Augustine, 2004), it is likely that synaptotagmin 2a and 2b have similar properties as synaptotagmin 1. Notably, synaptotagmin 2b underlies synchronous release in the zebrafish neuromuscular junction (Wen et al., 2010).

Table 6. Homology of the Ca²⁺ binding motifs.

Protein	Accession #	C2A domain Loop 1	Loop 3	C2B domain Loop 1	Loop 3
Class 1					
dre_1b_1	NP_001082930.1	MDMSGTSD	DFDRESKHD	MDVGGLSD	-----
dre_1b_X1	XP_021323234.1	MDMSGTSD	DFDRESKHD	MDVGGLSD	-----
dre_1a_1	NP_001314758.1	MDMGGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
dre_1a_X1	XP_017210484.1	MDMSGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
dre_1a_X2	XP_009298648.1	MDMSGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
hsa_1_1	NP_001129277.1	LDMGGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
hsa_1_2	NP_001278830.1	LDMGGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
rno_1_1	NP_001028852.2	LDMGGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
mmu_1_1	NP_001239270.1	LDMGGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
mmu_1_2	NP_001239271.1	LDMGGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
		: ** *****	*****	*****	
dre_2b_X1	XP_005166211.1	MDSGGTSD	DYDRESKHD	MDVGGLSD	DYDKIGKND
dre_2b_X2	XP_688092.3	MDSGGTSD	DYDRESKHD	MDVGGLSD	DYDKIGKND
dre_2a_X3	XP_021325567.1	MDSGGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
dre_2a_X1	XP_696277.2	MDSGGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
dre_2a_X2	XP_005174169.1	MDSGGTSD	DFDRESKHD	MDVGGLSD	DYDKIGKND
hsa_2_1	NP_001129976.1	LDMGGTSD	DFDRESKHD	MDVGGLSD	DYDKLGKNE
rno_2_1	NP_036797.2	LDMGGTSD	DFDRESKHD	MDVGGLSD	DYDKLGKNE
mmu_2_1	NP_001342655.1	LDMGGTSD	DFDRESKHD	MDVGGLSD	DYDKLGKNE
		: * *****	* : *****	*****	**** : *** :
Class 2					
dre_7b_1	NP_001232886.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
dre_7b_X1	XP_005166426.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
dre_7a_X1	XP_021326273.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
hsa_7_1	NP_001238994.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
hsa_7_2	NP_004191.2	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
hsa_7_3	NP_001287702.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
hsa_7_4	NP_001352738.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
hsa_7_5	NP_001357139.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
hsa_7_6	NP_001357140.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
rno_7_1	NP_067691.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
mmu_7_alpha	NP_061271.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
mmu_7_beta	NP_775090.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
mmu_7_gamma	NP_775091.2	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
mmu_7_4	NP_001360873.1	KDFSGTSD	DYDRESRND	MDIGGTSD	DKDKLSRND
		*****	*****	*****	*** , *****

Protein	Accession #	C2A domain Loop 1	Loop 3	C2B domain Loop 1	Loop 3
Class 3					
dre_3_X1	XP_005164031.1	KDANGFSD	DFDRFSRHD	MDLTGFSD	DYDCIGHNE
dre_3_X2	XP_021327105.1	KDANGFSD	DFDRFSRHD	MDLTGFSD	DYDCIGHNE
hsa_3_1	NP_001153800.1	KDSNGFSD	DFDRFSRHD	MDLTGFSD	DYDCIGHNE
rno_3_1	NP_061995.1	KDSNGFSD	DFDRFSRHD	MDLTGFSD	DYDCIGHNE
mmu_3_1	NP_001107588.1	KDSNGFSD	DFDRFSRHD	MDLTGFSD	DYDCIGHNE
		** : *****	*****	*****	*****
dre_6b_X1	XP_021335668.1	KDLGSSD	DFDRFSRHD	MDITGYSD	DYDLVGHNE
dre_6a_X1	XP_017209198.1	KDLGSSD	DFDRFSRHD	MDITGYSD	DYDLVGHNE
dre_6a_X2	XP_009295199.1	KDLGSSD	DFDRFSRHD	MDITGYSD	DYDLVGHNE
hsa_6_a	NP_001240701.1	KDFCGSSD	DFDRFSRHD	MDITGYSD	DYDRVGHNE
hsa_6_b	NP_001257734.1	KDFCGSSD	DFDRFSRHD	MDITGYSD	DYDRVGHNE
hsa_6_c	NP_001353153.1	KDFCGSSD	DFDRFSRHD	MDITGYSD	DYDRVGHNE
hsa_6_d	NP_001353154.1	KDFCGSSD	DFDRFSRHD	MDITGYSD	DYDRVGHNE
hsa_6_e	NP_001353155.1	KDFCGSSD	DFDRFSRHD	MDITGYSD	DYDRVGHNE
rno_6_1	NP_071527.1	KDFCGSSD	DFDRFSRHD	MDITGYSD	DYDRVGHNE
mmu_6_2	NP_001263605.1	KDFCGSSD	DFDRFSRHD	MDITGYSD	DYDRVGHNE
mmu_6_1	NP_061270.2	KDFCGSSD	DFDRFSRHD	MDITGYSD	DYDRVGHNE
mmu_6_3	NP_001263606.1	KDFCGSSD	DFDRFSRHD	MDITGYSD	DYDRVGHNE
mmu_6_4	NP_001263609.1	KDFCGSSD	DFDRFSRHD	MDITGYSE	VY-----
		** : *****	*****	*****;	*
dre_9b_1	NP_001003985.1	KDFTGTSD	DFDRFSRHD	MDITGASD	DYDRVGHNE
dre_9a_X1	XP_001336058.4	KDFSGTSD	DFDRFSRHD	MDITGASD	DYDRVGHNE
hsa_9_1	NP_783860.1	KDFSGTSD	DFDRFSRHD	MDITGASD	DYDRVGHNE
rno_9_1	NP_445776.1	KDFSGTSD	DFDRFSRHD	MDITGASD	DYDRVGHNE
mmu_9_1	NP_068689.2	KDFSGTSD	DFDRFSRHD	MDITGASD	DYDRVGHNE
		*** : ****	*****	*****	*****
dre_10_1	NP_001076311.1	KDFTGTSD	DFDRFTSHD	MDITGYSD	DYDRVGHNE
dre_10_X1	XP_017210538.2	KDFTGTSD	DFDRFTSHD	MDITGYSD	DYDRVGHNE
hsa_10_1	NP_945343.1	KDFTGTSD	DFDRFSRHD	MDITGSSD	DYDRVGHNE
rno_10_1	NP_113854.1	KDFTGTSD	DFDRFSRHD	MDITGSSD	DYDRVGHNE
mmu_10_1	NP_061273.1	KDFTGTSD	DFDRFSRHD	MDITGSSD	DYDRVGHNE
		*****	*****; **	***** **	*****
Class 4					
dre_4_1	NP_956242.1	TDEQSLTSD	SFDRFSRDE	ADSSGSPSD	DSDRTSRTP
hsa_4_1	NP_065834.1	MDEQSMSTSD	SFDRFSRDD	SDVSGLSLSD	DSEGRSRNE
rno_4_1	NP_113881.1	MDEQSMSTSD	SFDRFSRDD	SDVSGLSLSD	DSEGRSRNE
mmu_4_1	NP_033334.2	MDEQSMSTSD	SFDRFSRDD	SDVSGLSLSD	DSEGRSRNE
		*****; **	*****;	: * ** *	** : * **
Class 5					
dre_5b_1	NP_001018382.2	MDIGGTSD	DFDRFGKHD	MDVGGLSLSD	DYDKLGSND
dre_5a_1	NP_001096607.1	MDIGGTSD	DFDRFGKHD	MDVGGLSLSD	DYDKLGSND
dre_5a_X1	NP_001096607.1	MDIGGTSD	DFDRFGKHD	MDVGGLSLSD	DYDKLGSND
hsa_5_2	NP_001284703.1	LDLGGSSD	DFDRFSRND	MDVGGLSLSD	DYDKLGKNE
rno_5_1	NP_062223.1	LDLGGSSD	DFDRFSRND	MDVGGLSLSD	DYDKLGKNE
mmu_5_1	NP_001347350.1	LDLGGSSD	DFDRFSRND	MDVGGLSLSD	DYDKLGKNE
hsa_5_1	NP_003171.2	LDLGGSSD	DFDRFSRND	MDVGGLSLSD	DYDKLGKNE
		: *: **: **	*****. . : *	*****	*****. * :
Class 6					
dre_8_X4	XP_021326393.1	MDSGGTSD	DFNRFSEKHD	-----	-----
dre_8_X3	XP_021326392.1	MDSGGTSD	DFNRFSEKHD	MDQVGSSD	DHDKMSRND
dre_8_1	NP_001352218.1	MDSGGTSD	DFNRFSEKHD	MDQVGSSD	DHDKMSRND
dre_8_2	NP_001352219.1	MDSGGTSD	DFNRFSEKHD	MDQVGSSD	DHDKMSRND
hsa_8_1	NP_001277261.2	---GGTVD	NFKRFSGHE	---GLAE	DRSLPLRTE
hsa_8_2	NP_612634.4	---GGTVD	NFKRFSGHE	---GLAE	DRSLPLRTE
hsa_8_3	NP_001277262.2	---GGTVD	NFKRFSGHE	---GLAE	DRSLPLRTE
hsa_8_4	NP_001277263.2	---GGTVD	NFKRFSGHE	---GLAE	DRSLPLRTE
rno_8_1	NP_445777.1	---EGTAD	DFKRFSSEHE	---GLAE	ARGLQLLAE
mmu_8_a	NP_061272.2	---EGTAD	DFKRFSSEHE	---GLAE	ARGLQLRTE
mmu_8_b	NP_001272787.1	---EGTAD	DFKRFSSEHE	---GLAE	ARGLQLRTE
mmu_8_c	NP_001272790.1	---EGTAD	DFKRFSSEHE	---GLAE	ARGLQLRTE
		** *	: *: ** *		

Species used for these alignments: hsa = Homo sapiens; rno = Rattus norvegicus; mmu = Mus musculus; dre = Danio rerio. Sequences are named as species_protein_isoform (i.e. dre_1a_1 = Danio rerio synaptotagmin 1a, isoform 1). The Clustal consensus symbols are depicted under each alignment (“*” = no change; “:” = conservative amino acid substitution; “.” = semi-conservative amino acid substitution).

C2 domains in Class 2 and 3 synaptotagmins

Zebrafish class 2 synaptotagmins have only one semi-conservative amino acid substitution at a non-essential location in the C2B domain in relation to their mammalian orthologues. There are no substitutions in the C2A domain, which has been deemed more important than the C2B domain for controlling exocytosis in this class (Xue et al., 2010; Bacaj et al., 2013; Voleti et al., 2017). This extreme degree of conservation is highly suggestive of functional preservation (Table 5). In fact, the 3-dimensional model of the C2 domains of synaptotagmin 7b aligns well with the crystal structure of the C2 domains of rat synaptotagmin 7 (PDB: 6ANK, Voleti et al., 2017)(Figure 5A). It is interesting to note that the C2B domain of mammalian synaptotagmin 7 can bind 3 Ca^{2+} ions instead of 2, due to the presence of a serine residue in loop 3 (Xue et al., 2010). This makes the C2B domain of these proteins highly similar to the C2A domain as far as Ca^{2+} coordination is concerned, although its functional significance is yet unknown (Xue et al., 2010). This negative residue is also present in the C2B domain of zebrafish class 2 synaptotagmins (highlighted in yellow in Table 6).

Since synaptotagmin 7 was implicated in Ca^{2+} -dependent asynchronous release in mammals (Bacaj et al., 2013; Ablain et al., 2015), probably due to the higher affinity of its Ca^{2+} -bound C2A domain for phospholipids (Voleti et al., 2017), it is likely to perform the same role in the zebrafish. Indeed, synaptotagmin 7b was reported to control asynchronous release at the zebrafish neuromuscular junction (Wen et al., 2010). Other roles for mammalian synaptotagmin 7 in synaptic transmission are listed in Table 5.

Figure 5. Homology of the functional domains of zebrafish synaptotagmin 7b.

- (A) Ribbon model of the 3-dimensional structure of zebrafish synaptotagmin 7b, isoform 1 (rainbow colors) generated with RaptorX from the amino acid sequence (isoform 1, accession number in **Table 3**) and superposed on the crystal structure of the C2 domains of rat synaptotagmin 7 (shown in grey, PDB: 6ANK, Voleti et al., 2017). Figure is rotated 90 (center) and 180 degrees (right) to display the loops responsible for Ca^{2+} binding in the C2 domains (loops 1 and 3) of the rat synaptotagmin (pink).
- (B) Diagram summarizing the three Ca^{2+} -binding sites of the C2A domain, coordinated by loops 1 and 3 (pink), of rat synaptotagmin 7. The negative residues responsible for the coordination of Ca^{2+} binding are shown in blue (for aspartate) or yellow (serine). Redrawn from Voleti *et al.* (Fernandez et al., 2001; 2017).
- (C) Ribbon model of the three-dimensional structure of the C2A domain of zebrafish synaptotagmin 7b generated with RaptorX. Loops 1 and 3 are shown in pink. Residues responsible for Ca^{2+} binding are shown in same colors as in (B).

Both class 2 and class 3 synaptotagmins show two crucial amino acid substitutions in their C2A domains in relation to class 1 synaptotagmins that, in mammals, determine their much higher affinity to phospholipid membranes: the methionine (M)-to-phenylalanine (F) substitution

at the third position of loop 1 and the lysine (K)-to-arginine (R) substitution at the seventh position of loop 3 (Voleti et al., 2017). Although these are conservative replacements, they seem to highly impact Ca^{2+} -dependent phospholipid binding and show cooperative effects: the highest affinity is found when both substitutions are present (i.e. synaptotagmins 7, 7a, 7b, 6, 9a, 9b and mammalian 10). Zebrafish synaptotagmin 10 is an exception here: it has as negatively charged residue (serine) at the seventh position of loop 3 and no serine at the sixth position, which may impact its ability to coordinate Ca^{2+} binding.

C2 domains in Class 4 and 6 synaptotagmins

Class 4 and 6 synaptotagmins are the most divergent (**Figure 3, Table 6**). These proteins are unable to bind Ca^{2+} in mammals and function therefore as inhibitory synaptotagmins in these species (von Poser et al., 1997; Dai et al., 2004; Hui et al., 2005). Zebrafish synaptotagmin 4 misses more aspartates than the mammalian orthologues in the C2A domain and may not bind Ca^{2+} there, because the two missing aspartates coordinate together all Ca^{2+} binding sites in this domain. Since there is evidence that all three Ca^{2+} ions are required for binding with syntaxin (Ubach et al., 1998), this could render this synaptotagmin unable to drive exocytosis. Of note, the amino acid substitutions in the C2A domain are semi-conservative (aspartate (D)-to-serine (S)) or conservative (aspartate (D)-to-glutamate (E)), so there is a chance that these negatively charged residues could still create Ca^{2+} binding sites ((Dai et al., 2004), but see (von Poser et al., 1997; Wang & Zhang, 2017)). The C2B domain of zebrafish synaptotagmin 4 has more aspartates than its mammalian orthologues and may still be able to bind at least one Ca^{2+} ion according to the model in **Figure 4A-B** (Fernandez et al., 2001).

Whether a synaptotagmin will bind Ca^{2+} efficiently depends on both the amino acid

sequence and on the 3-dimensional structure of loops 1 and 3 in each C2 domain (i.e. the orientation of the negative residues in space) (Dai et al., 2004). This 3-dimensional structure, in turn, is determined by the amino acid composition of the loops themselves and of the flanking regions that interact electrically with the loops (Qiu et al., 2017). For instance, although rat and *Drosophila* synaptotagmin 4 have exactly the aspartate (D)-to-serine (S) substitution in loop 1 of the C2A domain, only in the latter does it function as a Ca^{2+} sensor, presumably because loop 1 in rat has a different 3-dimensional structure (Dai et al., 2004). This is an example of homology without orthology; clearly, this protein suffered a change in function during evolution. It remains to be determined whether zebrafish synaptotagmin 4 can still function as a Ca^{2+} sensor.

While mammalian synaptotagmin 8 misses significant portions of loop 1 in both C2A and C2B domains, at least three isoforms of zebrafish synaptotagmin are not only intact but have most Ca^{2+} coordinating residues. The only exception is isoform X4, which is truncated. The first aspartate residue missing in loop 3 of the C2A domain in all orthologues coordinates binding of all 3 Ca^{2+} ions in synaptotagmin 1 (**Figure 4**); a study with mammalian synaptotagmins 1 and 2 suggests that its absence may lead to collapse of the whole Ca^{2+} binding pocket in this domain (Stevens & Sullivan, 2003). It is therefore unlikely that the C2A domain of zebrafish synaptotagmin 8 can bind Ca^{2+} , despite having more conserved residues in relation to other synaptotagmins than its mammalian orthologues. The C2B domain, however, has all 5 aspartates in this species, so it may still potentially coordinate Ca^{2+} binding. Further studies will be needed to address this issue.

C2 domains in Class 5 synaptotagmins

As discussed earlier, zebrafish synaptotagmin 5a and 5b fall between mammalian class 1 and 5.

We kept them in class 5 for this analysis to keep the nomenclature consistent. The low overall genomic homology between zebrafish synaptotagmin 5a and 5b is also reflected in the amino acid composition of their Ca^{2+} coordinating loops. Both synaptotagmin 5a and 5b lack the serine necessary to close the C2A binding pocket and coordinate Ca^{2+} binding in the third site of this domain (Qiu et al., 2017). However, the same semi-conservative serine (S)-to-glycine (G) substitution in human synaptotagmin 5 does not compromise its three Ca^{2+} binding sites. Rather, this substitution leads only to a slight conformational change in loop 3 of the C2A domain, which renders the Ca^{2+} binding pocket more open and decreases Ca^{2+} affinity, making synaptotagmin 5 more synaptotagmin 1-like (Qiu et al., 2017). Further, the zebrafish synaptotagmins have all aspartates in the C2B domain, whereas the mammalian orthologues have a conservative aspartate (D)-to-glutamate (E) substitution in relation to other isoforms, similar to the one found in class 3 synaptotagmins, and may actually bind less Ca^{2+} ions there (Sutton et al., 1999). Therefore, the Ca^{2+} binding motifs of zebrafish synaptotagmins 5a and 5b may be more homologous to those of class 1 synaptotagmins than to those of other class 5 synaptotagmins. These results, added to the fact that mammalian synaptotagmins 1, 2 and 5 were shown to be involved in synchronous release in central synapses (Xu et al., 2007), make zebrafish 5a and 5b good candidates for mediating fast exocytosis.

In summary, the results presented thus far point to a generally good functional homology between zebrafish and mammalian synaptotagmins at the protein level, especially for classes 1, 2, 3 and 5. We next concentrated on the zebrafish isoforms that could likely serve as Ca^{2+} sensors for neurotransmitter release in the retina.

Expression and distribution of synaptotagmin candidates for Ca^{2+} -dependent exocytosis in

the zebrafish retina

Since class 3 synaptotagmins in mammals are mostly involved in processes other than vesicle release proper, such as endocytosis and non-synaptic secretion (**Table 5**), we set out to determine the retinal expression of *syt1l*, *syt2*, *syt5* and *syt7* paralogues. RT-PCR cDNA fragments of all these genes were found in the zebrafish brain, eye, and retina, except for *syt1b* (**Figure 6** and **Table 7**). This paralogue was not detected in the eye or retina but was present in brain.

Figure 6. Retinal expression and distribution of zebrafish *syt1* paralogue genes.

Agarose gel electrophoresis of RT-PCR cDNA products amplified from total RNA from adult zebrafish brain or eye using primers for *syt1a* and *syt1b* paralogues (**Table 1**). The sizes of the molecular markers are indicated on the left (*bp*: base pairs). *Syt1a* is present in both brain and eye, whereas *syt1b* is only present in the brain. The different lanes for each tissue represent tissue RNA + reverse transcriptase (+RT), plus two controls: tissue RNA without reverse transcriptase (-RT), and tissue RNA with no DNA (-DNA).

We also identified expression of a new splice variant of *syt7a* in zebrafish eye (named “*syt7a_splice*” in **Table 7**). This variant has close homology with *syt7a*-like sequences in bony fishes (ex: XP_016324438.1) but misses 774 base pairs in relation to variant XM_021470598.1,

which codes for a long version of the protein (653 amino acids, **Table 3**). Synaptotagmin 7a_{splice} should therefore be around 395 amino acids long. In mammals, the L1 region (**Figure 2**) of synaptotagmin 7 undergoes extensive alternative splicing (Sugita et al., 2001). This leads to dramatic changes in protein structure and several protein isoforms of different lengths (**Table 4**), but this truncation at a linker region is of minor consequence for protein function. The alternatively spliced sequence also codes for L1 and should therefore give rise to a fully functional protein. The sequenced fragment for *syt7a_{splice}* is in the *Appendix*.

Table 7. Expression of *syt* mRNAs in different tissues assayed by RT-PCR.

Zebrafish gene	Brain	Eye	Retina
<i>syt1a</i>	+	+	+
<i>syt1b</i>	+	-	-
<i>syt2a</i>	+	+	+
<i>syt2b</i>	+	+	+
<i>syt5a</i>	+	+	+
<i>syt5b</i>	+	+	+
<i>syt7a</i>	+	+	+
<i>syt7a_{splice}</i>	+	+	+
<i>syt7b</i>	+	+	+

To study protein distribution in the adult zebrafish retina in more detail, we assayed mRNA expression for the different isoforms by *in situ* hybridization on retinal slices. Riboprobes were designed using ATG as the 5' start if sequences were divergent in this area. Alternate primer locations were used if isotypes were not divergent. Since we identified a splice variant in *syt7a*, its riboprobe was located in the 3' UTR to ensure that expression of all *syt7a* variants were detected.

We tested two isotype-specific riboprobes for synaptotagmin 1a (**Table 2**): a shorter probe that targeted only the coding sequence (CDS) and a longer riboprobe that included the 3'-

UTR and could presumably enhance signal in case of low expression levels, as well as further validate the expression pattern seen with original probe (**Figure 7A-B**). Both yielded similar expression patterns, albeit the signal-to-noise ratio was better with the shorter CDS probe. *Syt1a* mRNA could be detected in three locations: (i) the outer plexiform layer (*OPL*), where photoreceptors contact bipolar cells and horizontal cells; (ii) the inner border of the inner nuclear layer (*INL*), where the somas of cone-driven bipolar cells and amacrine cells are located; and (iii) around somas in the ganglion cell layer (*GCL*).

Although we sequenced a fragment consistent with *syt2a* in the zebrafish retina by PCR, our short CDS riboprobe did not yield any signal (**Figure 7C**). This result could mean that either the *syt2a* paralogue is not transcribed, the products of this gene could be targeted for nonsense mediated decay, and/or that the probe was too short to hybridize strongly. Further experiments with a longer isotype-specific probe are needed to disambiguate this result. It may be necessary to confirm that the full-length *syt2a* gene is indeed found in zebrafish retinal tissue. If positive, specific probes, located in either the 3' or 5' UTR should be generated.

Weak labeling for *syt5a* mRNA was seen in the inner segments (IS) of short cones and in the outer half of the INL, where the cell bodies of horizontal cells and of mixed-input bipolar cells (i.e. bipolar cells that contact both rods and cones) and cone-driven bipolar cells are located (**Figure 7D**). The expression pattern of *syt7a* was like that of *syt1a*, which could mean that these synaptotagmins 1a and 7a are expressed in the same neurons in the zebrafish (**Figure 7E**).

Finally, *syt7b* was detected at the inner border of the INL and in the GCL (**Figure 7F**).

The differential expression of *syt1a* and *syt5a* suggest that these isoforms may control synchronous release in distinct neurons in the zebrafish retina, while *syt7a* and *syt7b* would mediate asynchronous release, vesicle replenishment or other Ca^{2+} dependent processes,

probably in different neuronal populations as well.

Figure 7. Retinal expression and distribution of zebrafish *syt* mRNAs assayed by *in situ* hybridization.

For all panels, the left panel shows the negative control with a sense probe. *IS*: photoreceptor inner segments; *ONL*: outer nuclear layer; *OPL*: outer plexiform layer; *INL*: inner nuclear layer; *IPL*: inner plexiform layer; *GCL*: ganglion cell layer.

(A, B) Detection of *syt1a* mRNA using a short CDS riboprobe (A) and a longer 3'-UTR riboprobe (B). *Syt1a* is expressed at highest levels by cells in the *OPL*, inner border of the *INL* and in the *GCL*.

(C) *Syt2a* mRNA was undetectable by a short CDS type-specific riboprobe. An additional riboprobe (perhaps larger fragment in the UTR specific to *syt2a*) needs to be tested.

(D) *Syt5a* mRNA is found at highest levels by inner segments of small cones located at the border with the *ONL*, and by cells at the outer half of the *INL*.

(E) *Syt7a* mRNA can be detected at the *OPL*, inner half of the *INL* and *GCL*.

(F) *Syt7b* is expressed at highest levels by cells at the inner border of the *INL* and by cells in the *GCL*.

DISCUSSION

In this study, we took advantage of bioinformatics to address the similarity between zebrafish and human synaptotagmins. We focused on synaptotagmin 1-10 as their roles are better defined and synaptotagmins 11-17 are unlikely candidates to mediate Ca^{2+} exocytosis in the central nervous system (von Poser et al., 1997; Fukuda, 2003a, b; Bhalla et al., 2008; Craxton, 2010). Because of whole genome duplication, zebrafish generally have two paralogues for each human orthologue, except for *syt3*, *syt4*, *syt8* and *syt10* (Craxton, 2004, 2010). Despite duplication and the evolutionary distance, we find that most zebrafish synaptotagmins share common features with mouse, rat and human orthologues, including key motifs involved in Ca^{2+} binding. Accordingly, we find that *syt1a*, *syt2a*, *syt2b*, *syt5a*, *syt5b*, *syt7a* and *syt7b* genes are expressed in the zebrafish brain, eye and retina. From these, we successfully detected mRNA for *syt1a*, *syt5a*, *syt7a* and *syt7b* in retinal tissue with distinct distribution patterns. This differential expression suggests that these synaptotagmins may control various aspects of the vesicle cycle in distinct subsets of neurons. Overall, these results highlight that zebrafish will be an outstanding model to study the role of synaptotagmins in synaptic transmission in the retina.

Homology of zebrafish synaptotagmins

In the present study, we characterized genetic and protein similarities between zebrafish and human synaptotagmins (**Figures 1-3**). Importantly, zebrafish synaptotagmins cluster mostly into the same classes as their mammalian counterparts (**Figure 3**), except synaptotagmins 5a and 5b, which may be more homologous to class 1 synaptotagmins. In addition, zebrafish synaptotagmins show strong similarities to their mammalian counterparts in terms of their Ca^{2+}

binding motifs (**Figures 4-5**, and **Table 6**). Notable exceptions are zebrafish synaptotagmin 1b, which lacks 34 amino acid residues in its C2B domain, and synaptotagmin 8, which has more crucial residues for Ca^{2+} binding in both C2 domains than the mammalian synaptotagmin 8.

Synaptotagmin 1b is unlikely to act as a Ca^{2+} sensor for exocytosis. In mammals (Dai et al., 2004; Bhalla et al., 2008; Xue et al., 2010; Voleti et al., 2017), Ca^{2+} binding to the C2B domain of synaptotagmin 1 is crucial for synchronous release (Mackler & Reist, 2001; Mackler et al., 2002; Nishiki & Augustine, 2004; Shin et al., 2009; Bacaj et al., 2013; Lee et al., 2013), probably because the C2B domain has significantly higher sensitivity to Ca^{2+} (Bradberry et al., 2020) and phospholipid-binding activity (Bai et al., 2004; Li et al., 2006; van den Bogaart et al., 2012; Bradberry et al., 2020) than the C2A domain in this protein. Indeed, all mutations in synaptotagmin 1 that cause diseases in humans target the C2B domain (Baker et al., 2018; Bradberry et al., 2020), while removal of the residues critical for Ca^{2+} binding in the C2A domain is relatively innocuous (Stevens & Sullivan, 2003).

One could wonder why the *sytlb* gene is expressed at all, since its product lacks the ability to effectively bind Ca^{2+} . Synaptotagmins can bind other synaptotagmins and t-SNARES in a Ca^{2+} -independent manner, thereby displacing the Ca^{2+} -sensitive isoforms and impeding direct contact between them and their targets (Bhalla et al., 2008). It is possible that synaptotagmin 1b may be one such inhibitory protein that regulates the activity of other synaptotagmins in the zebrafish brain, like mammalian synaptotagmins 4 and 8 (von Poser et al., 1997; Dai et al., 2004; Hui et al., 2005; Bhalla et al., 2008).

That said, it is important to note that sequence analysis alone predicts Ca^{2+} binding properties rather poorly. For example, despite the high homology of their C2 Ca^{2+} binding pockets, drosophila synaptotagmin 4 is Ca^{2+} sensitive, while its mammalian orthologue is not

(Dai et al., 2004; Bhalla et al., 2008). Given that the differences in amino acid composition of the C2 domains of Ca^{2+} -sensitive synaptotagmins are rather subtle, the basis for their Ca^{2+} sensitivity and functional differences remain poorly understood (Xue et al., 2010; Voleti et al., 2017). For both zebrafish synaptotagmin 1b and 8 further experiments are thus necessary to determine whether they may or may not function as Ca^{2+} sensors.

Synaptotagmins expressed in the zebrafish retina

Although zebrafish is a powerful model organism, the duplication of its genome can often complicate the identification of homologous genes and their isotypes. Frequently, the available sequence data is not mapped, completely validated, or correctly named in the various databases. This bioinformatic study allowed us to examine the current genomic and mRNA sequences and ensure that current nomenclature for these various synaptotagmins are accurately based on their mammalian orthologues. With focus on those synaptotagmins that are likely candidates for mediating Ca^{2+} -dependent exocytosis in the retina, we were thus able to confidently identify appropriate sequence data for *syt1a/b*, *syt2a/b*, *syt5a/b* and *syt7a/b*. These sequences were then used to design RT-PCR primers and ultimately riboprobes for identification of expressed synaptotagmins in the retina.

We find using RT-PCR that 7 of the 8 studied paralogues are present in retina, the exception being *syt1b*. This raises the possibility that the various synaptotagmins might make differential contributions to retinal physiology. Of these, we could only detect mRNA for *syt1a*, *syt5a*, *syt7a* and *syt7b* by *in situ* hybridization. This divergence might reflect that very short CDS riboprobes needed to guarantee isotype-specificity may limit detection. Longer riboprobes containing sequences from UTRs, which will maintain specificity and hopefully improve signal-

to-noise ratio, will be needed to fully address this issue. Alternatively, it could be that some of the fragments amplified by RT-PCR (i.e from *syt2a* and *syt2b*) do not reflect the presence of a full-length gene in the retina. In this case, zebrafish and mammalian retinas could use different sets of synaptotagmins to control vesicle release in ribbon-type synapses of the OPL, since synaptotagmin 2 was reported in the rat and mouse OPL (Ullrich & Sudhof, 1994; Fox & Sanes, 2007), albeit in the mouse it seems to be excluded from photoreceptor terminals (Fox & Sanes, 2007). Future experiments will have to address this.

The expression of *syt1a* in the OPL is consistent with the immunofluorescence data in rat (Ullrich & Sudhof, 1994) and with the finding that synaptic release from photoreceptors in the mouse is largely controlled by synaptotagmin 1 (Grassmeyer et al., 2019). Additionally, the strong *syt1a* signal detected in the inner half of the INL and in the GCL is very similar to the one for *syt1* mRNA in the mouse (Fox & Sanes, 2007). Together, these results suggest that synaptotagmin 1a may have analogous functions in the zebrafish and control synchronous vesicle release both in ribbon synapses of photoreceptors and in conventional synapses of the retina, such as those of amacrine cells and ganglion cells.

While synaptotagmin 7 was shown to be involved in types of asynchronous release both at conventional (Bacaj et al., 2013; Luo & Sudhof, 2017) and ribbon-type synapses of retinal bipolar cells (Luo et al., 2015), the localization of mRNA signal for *syt7a* and *syt7b* in our study is not consistent with this role in zebrafish bipolar cells. Rather, it suggests that these paralogues are expressed mostly in amacrine cells (inner half of the INL) and ganglion cells (GCL). The weak *syt7a* signal we find in the zebrafish OPL suggests that synaptotagmin 7a may be additionally be present in photoreceptors. However, the role(s) of these proteins in the zebrafish retina still need elucidating, since synaptotagmin 7 was associated with other aspects of

neurotransmission, such as vesicle replenishment (Liu et al., 2014) and synaptic facilitation (Jackman et al., 2016; Jackman & Regehr, 2017).

Finally, the expression pattern of *syt5a* is consistent with cone photoreceptors (ONL and inner segments of short cones) and horizontal and bipolar cells (outer half of the INL). To our knowledge, this pattern does not resemble any data from mammals. This, added to the similarities between the products of zebrafish *syt5* paralogues with both class 1 and class 5 synaptotagmins in mammals warrants further investigation regarding whether these synaptotagmins are directly involved in neurotransmission.

APPENDIX

1. cDNA fragment of *syt7a_splice* (264 nt). ISH primer sequences are shown in bold font.

ATGTATCTCAACAGGGAGGAGGAGTACAGCAAAGGCTCCATCTCTTTGAGCGTGC
TGCTGGTGTCTGTTGGCGGTAAGTGTGTGTGGGGTTTGGCTGGTGGCTCTCTGTGGCG
TCTGTGGATGGTGTCAACGCAAGCTGGGGAAGAGGAATAAACCCGGAGTGGAGTCC
GTCGGGTCTCCAGATTCAGGAAGAGGAAGAGGGGAGAAAAAGCCATCAATAGGA
ATATGGGGAATAAGCCAGCCAACAGTCCTAAGGGTCAGCCG

2. BLASTN alignment to *Sinocyclocheilus anshuiensis* synaptotagmin-7-like (LOC107674837), transcript variant X1 (95% identity). Zebrafish *syt7a_splice* sequence is shown as “query”.

Download ▾ GenBank Graphics ▾ Next ▲ Previous

PREDICTED: Sinocyclocheilus anshuiensis synaptotagmin-7-like (LOC107674837), transcript variant X1, mRNA
Sequence ID: [XM_016468952.1](#) Length: 2774 Number of Matches: 1

Range 1: 21 to 284 [GenBank](#) [Graphics](#) ▾ Next Match ▲ Previous Match

Score	Expect	Identities	Gaps	Strand
423 bits(468)	3e-114	252/264(95%)	0/264(0%)	Plus/Plus

Query 1 ATGTATCTCAACAGGGAGGAGGAGTACAGCAAAGGCTCCATCTCTTTGAGCGTCTGCTG 60
Sbjct 21 ATGTATCTCAACAGGGAGGAGGAGTACAGCAAAGGCTCCGTCCTTTGAGCGTCTGCTG 80

Query 61 GTGTCGTTGGCGGTAAGTGTGTGTGGGGTTTGGCTGGTGGCTCTCTGTGGCGTCTGTGA 120
Sbjct 81 GTGTCGTTGGCGGTAAGCGTGTGTGTGGTGGTGGCTGGTGGTGGTGGTGGTGGTGGTGGT 140

Query 121 TGGTGTCAACGCAAGCTGGGGAAGAGGAATAAACCCGGAGTGGAGTCCGTCGGGTCTCCA 180
Sbjct 141 TGGTGTCAACGCAAGCTGGGGAAGAGGAATAAACCCGGAGTGGAGTCCGTCGGGTCTCCT 200

Query 181 GATTTCAGGAAGAGGAAGAGGGGAGAAAAAGCCATCAATAGGAATATGGGGAAATAGCCA 240
Sbjct 201 GATTTCAGGAAGAGGAAGAGGGGAGAAAAAGCCATCAATAGGAATATGGGGAAATAGCCA 260

Query 241 GCCAACAGTCCTAAGGGTCAGCCG 264
Sbjct 261 GCTAACAGTCCTAAGGGTCAGCCG 284

[Questions/comments](#)

Related Info
[Gene - associat](#)
[Genome Data](#)
genomic conte

REFERENCES

- Ablain J, Durand EM, Yang S, Zhou Y, Zon LI (2015) A CRISPR/Cas9 vector system for tissue-specific gene disruption in zebrafish. *Dev Cell* 32:756-764.
- Amores A, Force A, Yan YL, Joly L, Amemiya C, Fritz A, Ho RK, Langeland J, Prince V, Wang YL, Westerfield M, Ekker M, Postlethwait JH (1998) Zebrafish hox clusters and vertebrate genome evolution. *Science* 282:1711-1714.
- Angueyra JM, Kindt KS (2018) Leveraging zebrafish to study retinal degenerations. *Front Cell Dev Biol* 6:110.
- Bacaj T, Wu D, Yang X, Morishita W, Zhou P, Xu W, Malenka RC, Sudhof TC (2013) Synaptotagmin-1 and synaptotagmin-7 trigger synchronous and asynchronous phases of neurotransmitter release. *Neuron* 80:947-959.
- Bai J, Tucker WC, Chapman ER (2004) PIP2 increases the speed of response of synaptotagmin and steers its membrane-penetration activity toward the plasma membrane. *Nat Struct Mol Biol* 11:36-44.
- Baker K et al. (2018) SYT1-associated neurodevelopmental disorder: a case series. *Brain* 141:2576-2591.
- Berntson AK, Morgans CW (2003) Distribution of the presynaptic calcium sensors, synaptotagmin I/II and synaptotagmin III, in the goldfish and rodent retinas. *J Vis* 3:274-280.
- Bhalla A, Tucker WC, Chapman ER (2005) Synaptotagmin isoforms couple distinct ranges of Ca^{2+} , Ba^{2+} , and Sr^{2+} concentration to SNARE-mediated membrane fusion. *Mol Biol Cell* 16:4755-4764.
- Bhalla A, Chicka MC, Chapman ER (2008) Analysis of the synaptotagmin family during

807 reconstituted membrane fusion. Uncovering a class of inhibitory isoforms. *J Biol Chem*
808 283:21799-21807.

809 Bradberry MM, Courtney NA, Dominguez MJ, Lofquist SM, Knox AT, Sutton RB, Chapman
810 ER (2020) Molecular basis for synaptotagmin-1-associated neurodevelopmental disorder.
811 *Neuron* 107:52-64 e57.

812 Butz S, Fernandez-Chacon R, Schmitz F, Jahn R, Sudhof TC (1999) The subcellular
813 localizations of atypical synaptotagmins III and VI. Synaptotagmin III is enriched in synapses
814 and synaptic plasma membranes but not in synaptic vesicles. *J Biol Chem* 274:18290-18296.

815 Cao P, Yang X, Sudhof TC (2013) Complexin activates exocytosis of distinct secretory vesicles
816 controlled by different synaptotagmins. *J Neurosci* 33:1714-1727.

817 Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in
818 phylogenetic analysis. *Mol Biol Evol* 17:540-552.

819 Cathala G, Savouret JF, Mendez B, West BL, Karin M, Martial JA, Baxter JD (1983) A method
820 for isolation of intact, translationally active ribonucleic acid. *DNA* 2:329-335.

821 Chen C, Jonas P (2017) Synaptotagmins: that's why so many. *Neuron* 94:694-696.

822 Ciccarelli FD, Doerks T, von Mering C, Creevey CJ, Snel B, Bork P (2006) Toward automatic
823 reconstruction of a highly resolved tree of life. *Science* 311:1283-1287.

824 Craxton M (2004) Synaptotagmin gene content of the sequenced genomes. *BMC Genomics* 5:43.

825 Craxton M (2010) A manual collection of Syt, Eyt, Rph3a, Rph3al, Doc2, and Dblc2 genes
826 from 46 metazoan genomes--an open access resource for neuroscience and evolutionary
827 biology. *BMC Genomics* 11:37.

828 Craxton M, Goedert M (1995) Synaptotagmin V: a novel synaptotagmin isoform expressed in rat
829 brain. *FEBS Lett* 361:196-200.

830 Dai H, Shin OH, Machius M, Tomchick DR, Sudhof TC, Rizo J (2004) Structural basis for the
831 evolutionary inactivation of Ca²⁺ binding to synaptotagmin 4. *Nat Struct Mol Biol* 11:844-
832 849.

833 Dean C, Dunning FM, Liu H, Bomba-Warczak E, Martens H, Bharat V, Ahmed S, Chapman ER
834 (2012) Axonal and dendritic synaptotagmin isoforms revealed by a pHluorin-syt functional
835 screen. *Mol Biol Cell* 23:1715-1727.

836 Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space
837 complexity. *BMC Bioinformatics* 5:113.

838 Fagerberg L et al. (2014) Analysis of the human tissue-specific expression by genome-wide
839 integration of transcriptomics and antibody-based proteomics. *Mol Cell Proteomics* 13:397-
840 406.

841 Fernandez I, Arac D, Ubach J, Gerber SH, Shin O, Gao Y, Anderson RG, Sudhof TC, Rizo J
842 (2001) Three-dimensional structure of the synaptotagmin 1 C2B-domain: synaptotagmin 1 as
843 a phospholipid binding machine. *Neuron* 32:1057-1069.

844 Fox MA, Sanes JR (2007) Synaptotagmin I and II are present in distinct subsets of central
845 synapses. *J Comp Neurol* 503:280-296.

846 Fukuda M (2003a) Molecular cloning, expression, and characterization of a novel class of
847 synaptotagmin (Syt XIV) conserved from *Drosophila* to humans. *J Biochem* 133:641-649.

848 Fukuda M (2003b) Molecular cloning and characterization of human, rat, and mouse
849 synaptotagmin XV. *Biochem Biophys Res Commun* 306:64-71.

850 Fuson KL, Montes M, Robert JJ, Sutton RB (2007) Structure of human synaptotagmin 1 C2AB
851 in the absence of Ca²⁺ reveals a novel domain association. *Biochemistry* 46:13041-13048.

852 Geppert M, Goda Y, Hammer RE, Li C, Rosahl TW, Stevens CF, Sudhof TC (1994)

853 Synaptotagmin I: a major Ca^{2+} sensor for transmitter release at a central synapse. *Cell*
854 79:717-727.

855 Gestri G, Link BA, Neuhauss SC (2012) The visual system of zebrafish and its use to model
856 human ocular diseases. *Dev Neurobiol* 72:302-327.

857 Glasauer SM, Neuhauss SC (2014) Whole-genome duplication in teleost fishes and its
858 evolutionary consequences. *Mol Genet Genomics* 289:1045-1060.

859 Grassmeyer JJ, Cahill AL, Hays CL, Barta C, Quadros RM, Gurumurthy CB, Thoreson WB
860 (2019) Ca^{2+} sensor synaptotagmin-1 mediates exocytosis in mammalian photoreceptors.
861 *Elife* 8.

862 Gustavsson N, Han W (2009) Calcium-sensing beyond neurotransmitters: functions of
863 synaptotagmins in neuroendocrine and endocrine secretion. *Biosci Rep* 29:245-259.

864 Haberman Y, Grimberg E, Fukuda M, Sagi-Eisenberg R (2003) Synaptotagmin IX, a possible
865 linker between the perinuclear endocytic recycling compartment and the microtubules. *J Cell*
866 *Sci* 116:4307-4318.

867 Hudson AW, Birnbaum MJ (1995) Identification of a nonneuronal isoform of synaptotagmin.
868 *Proc Natl Acad Sci U S A* 92:5895-5899.

869 Hui E, Bai J, Wang P, Sugimori M, Llinas RR, Chapman ER (2005) Three distinct kinetic
870 groupings of the synaptotagmin family: candidate sensors for rapid and delayed exocytosis.
871 *Proc Natl Acad Sci U S A* 102:5210-5214.

872 Jackman SL, Regehr WG (2017) The mechanisms and functions of synaptic facilitation. *Neuron*
873 94:447-464.

874 Jackman SL, Turecek J, Belinsky JE, Regehr WG (2016) The calcium sensor synaptotagmin 7 is
875 required for synaptic facilitation. *Nature* 529:88-91.

876 Jackman SL, Choi SY, Thoreson WB, Rabl K, Bartoletti TM, Kramer RH (2009) Role of the
877 synaptic ribbon in transmitting the cone light response. *Nat Neurosci* 12:303-310.

878 Katz B, Miledi R (1965) The effect of calcium on acetylcholine release from motor nerve
879 terminals. *Proc R Soc Lond B Biol Sci* 161:496-503.

880 Kawakami K (2007) Tol2: a versatile gene transfer vector in vertebrates. *Genome Biol* 8 Suppl
881 1:S7.

882 Kawakami K, Asakawa K, Hibi M, Itoh M, Muto A, Wada H (2016) Gal4 driver transgenic
883 zebrafish: powerful tools to study developmental biology, organogenesis, and neuroscience.
884 *Adv Genet* 95:65-87.

885 Lee J, Guan Z, Akbergenova Y, Littleton JT (2013) Genetic analysis of synaptotagmin C2
886 domain specificity in regulating spontaneous and evoked neurotransmitter release. *J Neurosci*
887 33:187-200.

888 Letunic I, Bork P (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new
889 developments. *Nucleic Acids Res* 47:W256-W259.

890 Li C, Ullrich B, Zhang JZ, Anderson RG, Brose N, Sudhof TC (1995) Ca²⁺-dependent and -
891 independent activities of neural and non-neural synaptotagmins. *Nature* 375:594-599.

892 Li L, Shin OH, Rhee JS, Arac D, Rah JC, Rizo J, Sudhof T, Rosenmund C (2006)
893 Phosphatidylinositol phosphates as co-activators of Ca²⁺ binding to C2 domains of
894 synaptotagmin 1. *J Biol Chem* 281:15845-15852.

895 Liu H, Bai H, Hui E, Yang L, Evans CS, Wang Z, Kwon SE, Chapman ER (2014)
896 Synaptotagmin 7 functions as a Ca²⁺-sensor for synaptic vesicle replenishment. *Elife*
897 3:e01524.

898 Luo F, Sudhof TC (2017) Synaptotagmin-7-mediated asynchronous release boosts high-fidelity

899 synchronous transmission at a central synapse. *Neuron* 94:826-839 e823.

900 Luo F, Bacaj T, Sudhof TC (2015) Synaptotagmin-7 is essential for Ca²⁺-triggered delayed
901 asynchronous release but not for Ca²⁺-dependent vesicle priming in retinal ribbon synapses. *J*
902 *Neurosci* 35:11024-11033.

903 Mackler JM, Reist NE (2001) Mutations in the second C2 domain of synaptotagmin disrupt
904 synaptic transmission at *Drosophila* neuromuscular junctions. *J Comp Neurol* 436:4-16.

905 Mackler JM, Drummond JA, Loewen CA, Robinson IM, Reist NE (2002) The C(2)B Ca(2+)-
906 binding motif of synaptotagmin is required for synaptic transmission in vivo. *Nature* 418:340-
907 344.

908 Matthews G, Fuchs P (2010) The diverse roles of ribbon synapses in sensory neurotransmission.
909 *Nat Rev Neurosci* 11:812-822.

910 Miledi R (1973) Transmitter release induced by injection of calcium ions into nerve terminals.
911 *Proc R Soc Lond B Biol Sci* 183:421-425.

912 Moravec CE, Samuel J, Weng W, Wood IC, Sirotkin HI (2016) Maternal REST/NRSF regulates
913 zebrafish behavior through SNAP25a/b. *J Neurosci* 36:9407-9419.

914 Neher E, Sakaba T (2008) Multiple roles of calcium ions in the regulation of neurotransmitter
915 release. *Neuron* 59:861-872.

916 Neumann S, Haverkamp S (2013) Characterization of small-field bistratified amacrine cells in
917 macaque retina labeled by antibodies against synaptotagmin-2. *J Comp Neurol* 521:709-724.

918 Niklaus S, Neuhauss SCF (2017) Genetic approaches to retinal research in zebrafish. *J*
919 *Neurogenet* 31:70-87.

920 Nishiki T, Augustine GJ (2004) Dual roles of the C2B domain of synaptotagmin I in
921 synchronizing Ca²⁺-dependent neurotransmitter release. *J Neurosci* 24:8542-8550.

922 Oesch NW, Diamond JS (2011) Ribbon synapses compute temporal contrast and encode
923 luminance in retinal rod bipolar cells. *Nat Neurosci* 14:1555-1561.

924 Pasquier J, Cabau C, Nguyen T, Jouanno E, Severac D, Braasch I, Journot L, Pontarotti P, Klopp
925 C, Postlethwait JH, Guiguen Y, Bobe J (2016) Gene evolution and gene expression after
926 whole genome duplication in fish: the PhyloFish database. *BMC Genomics* 17:368.

927 Postlethwait JH, Woods IG, Ngo-Hazelett P, Yan YL, Kelly PD, Chu F, Huang H, Hill-Force A,
928 Talbot WS (2000) Zebrafish comparative genomics and the origins of vertebrate
929 chromosomes. *Genome Res* 10:1890-1902.

930 Qiu X, Ge J, Gao Y, Teng M, Niu L (2017) Structural analysis of Ca(2+)-binding pocket of
931 synaptotagmin 5 C2A domain. *Int J Biol Macromol* 95:946-953.

932 Shin OH, Xu J, Rizo J, Sudhof TC (2009) Differential but convergent functions of Ca²⁺ binding
933 to synaptotagmin-1 C2 domains mediate neurotransmitter release. *Proc Natl Acad Sci U S A*
934 106:16469-16474.

935 Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert
936 M, Soding J, Thompson JD, Higgins DG (2011) Fast, scalable generation of high-quality
937 protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* 7:539.

938 Stevens CF, Sullivan JM (2003) The synaptotagmin C2A domain is part of the calcium sensor
939 controlling fast synaptic transmission. *Neuron* 39:299-308.

940 Sudhof TC (2002) Synaptotagmins: why so many? *J Biol Chem* 277:7629-7632.

941 Sugita S, Shin OH, Han W, Lao Y, Sudhof TC (2002) Synaptotagmins form a hierarchy of
942 exocytotic Ca(2+) sensors with distinct Ca(2+) affinities. *EMBO J* 21:270-280.

943 Sugita S, Han W, Butz S, Liu X, Fernandez-Chacon R, Lao Y, Sudhof TC (2001) Synaptotagmin
944 VII as a plasma membrane Ca(2+) sensor in exocytosis. *Neuron* 30:459-473.

945 Sutton RB, Ernst JA, Brunger AT (1999) Crystal structure of the cytosolic C2A-C2B domains of
 946 synaptotagmin III. Implications for Ca²⁺-independent snare complex interaction. *J Cell Biol*
 947 147:589-598.

948 Takamori S et al. (2006) Molecular anatomy of a trafficking organelle. *Cell* 127:831-846.

949 Ubach J, Zhang X, Shao X, Sudhof TC, Rizo J (1998) Ca²⁺ binding to synaptotagmin: how
 950 many Ca²⁺ ions bind to the tip of a C2-domain? *EMBO J* 17:3921-3930.

951 Ullrich B, Sudhof TC (1994) Distribution of synaptic markers in the retina: implications for
 952 synaptic vesicle traffic in ribbon synapses. *J Physiol Paris* 88:249-257.

953 Ullrich B, Li C, Zhang JZ, McMahon H, Anderson RG, Geppert M, Sudhof TC (1994)
 954 Functional properties of multiple synaptotagmins in brain. *Neuron* 13:1281-1291.

955 van den Bogaart G, Meyenberg K, Diederichsen U, Jahn R (2012) Phosphatidylinositol 4,5-
 956 bisphosphate increases Ca²⁺ affinity of synaptotagmin-1 by 40-fold. *J Biol Chem* 287:16447-
 957 16453.

958 Voleti R, Tomchick DR, Sudhof TC, Rizo J (2017) Exceptionally tight membrane-binding may
 959 explain the key role of the synaptotagmin-7 C2A domain in asynchronous neurotransmitter
 960 release. *Proc Natl Acad Sci U S A* 114:E8518-E8527.

961 von Poser C, Ichtchenko K, Shao X, Rizo J, Sudhof TC (1997) The evolutionary pressure to
 962 inactivate. A subclass of synaptotagmins with an amino acid substitution that abolishes Ca²⁺
 963 binding. *J Biol Chem* 272:14314-14319.

964 Wang YL, Zhang CX (2017) Putting a brake on synaptic vesicle endocytosis. *Cell Mol Life Sci*
 965 74:2917-2927.

966 Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ (2009) Jalview Version 2--a
 967 multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25:1189-1191.

968 Wen H, Linhoff MW, McGinley MJ, Li GL, Corson GM, Mandel G, Brehm P (2010) Distinct
969 roles for two synaptotagmin isoforms in synchronous and asynchronous transmitter release at
970 zebrafish neuromuscular junction. *Proc Natl Acad Sci U S A* 107:13906-13911.

971 Wierson WA et al. (2020) Efficient targeted integration directed by short homology in zebrafish
972 and mammalian cells. *Elife* 9.

973 Wolfes AC, Dean C (2020) The diversity of synaptotagmin isoforms. *Curr Opin Neurobiol*
974 63:198-209.

975 Xu J, Mashimo T, Sudhof TC (2007) Synaptotagmin-1, -2, and -9: Ca²⁺ sensors for fast
976 release that specify distinct presynaptic properties in subsets of neurons. *Neuron* 54:567-581.

977 Xue M, Craig TK, Shin OH, Li L, Brautigam CA, Tomchick DR, Sudhof TC, Rosenmund C,
978 Rizo J (2010) Structural and mutational analysis of functional differentiation between
979 synaptotagmins-1 and -7. *PLoS One* 5.

980