Up-Regulation of α CaMKII Impairs Cued Fear Extinction and NMDAR-Dependent LTD in the Lateral Amygdala

Shuming An^{a#}, Jiayue Wang^{a#}, Xuliang Zhang^{a#}, Yanhong Duan^a, Junyan Lv^a, Dasheng Wang^a, Huan Zhang^a, Gal Richter-Levin^b, Xiaohua Cao^{a*}

a Key Laboratory of Brain Functional Genomics, MOE & STCSM, East China Normal University, Shanghai 200062, China

b “Sagol” Department of Neurobiology, University of Haifa, Haifa 31905, Israel

These authors contributed equally to this work.

*Correspondence: xhcao@brain.ecnu.edu.cn

Running title: α CaMKII up-regulation impairs fear extinction and LTD

ABSTRACT

Impaired fear extinction is one of the hallmark symptoms of post-traumatic stress disorder (PTSD). The roles of α CaMKII have been not extensively studied in fear extinction and LTD. Here, we found PTSD susceptible mice exhibited significant up-regulation of α CaMKII in the lateral amygdala (LA). Consistently, increasing α CaMKII in LA profoundly not only caused PTSD-like symptoms such as impaired fear extinction and anxiety-like behaviors, but also attenuated NMDAR-dependent LTD at thalamo-LA synapses, reduced GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization. Suppressing the elevated α CaMKII to normal level could completely reverse both PTSD-like symptoms and the impairments in LTD, GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization. Intriguingly, deficits in AMPAR internalization and GluA1-Ser845/Ser831 dephosphorylation were detected not only after impaired fear extinction, but also after attenuated LTD. Our results demonstrate for the first time GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization are molecular links between LTD and fear extinction, and suggest α CaMKII may be a potential molecular determinant of PTSD.

INTRODUCTION

Although some progresses have been made in understanding the molecular and cellular mechanisms of post-traumatic stress disorder (PTSD) recently, effective treatment for PTSD is still lacking. Since impaired fear extinction is one of the core symptoms of PTSD (Michopoulos et al., 2014; Yehuda et al., 2015), and fear extinction is the basis for psychological exposure therapy (M. R. Milad & Quirk, 2012), a deeper understanding of the molecular and cellular substrates underlying fear extinction would have important implications for developing the more effective treatment for PTSD.

At the synaptic level, long-term depression (LTD) has been implicated in fear extinction (Bennett, Arnold, Hatton, & Lagopoulos, 2017). N-methyl-D-aspartate (NMDA) GluN2B receptor antagonist can abolish both LTD at thalamo-lateral amygdala (T-LA) synapses and fear extinction (Dalton, Wu, Wang, Floresco, & Phillips, 2012). Moreover, deletion of kinesin superfamily proteins (KIFs) 21B impairs both hippocampal LTD and contextual fear extinction (Morikawa, Tanaka, Cho, Yoshihara, & Hirokawa, 2018). Besides, aquaporin-4 deficiency facilitates both NMDAR-dependent hippocampal LTD and fear extinction (Wu et al., 2017). Optogenetic delivery of LTD conditioning to the auditory input to LA facilitates cued fear extinction (Nabavi et al., 2014). Taken together, these findings indicate that there may be a link between LTD and fear extinction. NMDAR-dependent α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPA) internalization is involved in fear extinction (Bai, Zhou, Wu, & Dong, 2014; Dalton, Wang, Floresco, & Phillips, 2008; J. Kim et al., 2007; Lin, Mao, Su, & Gean, 2010). Notably, disruption of AMPAR internalization impairs fear extinction (Dalton et al., 2008; J. Kim et al., 2007). Conversely, the promotion of AMPAR internalization facilitates fear extinction (Bai et al., 2014; Lin et al., 2010). It has been well known that AMPAR internalization also participates in LTD (Brebner et al., 2005; Collingridge, Isaac, & Wang, 2004). Thus, we wonder whether AMPAR internalization is a direct link between fear extinction and LTD.

At the molecular level, CaMKII is the major kinase mediating NMDAR-dependent synaptic plasticity, AMPAR trafficking and memory (Collingridge et al., 2004). In mammals, CaMKII has four isoforms, α , β , γ and δ (Colbran & Soderling, 1990; Hell, 2014), and the α isoform is predominantly expressed in the forebrain (Kennedy, McGuinness, & Greengard, 1983). On the

one hand, α CaMKII plays a crucial role in long-term potentiation (LTP) and memory formation (Kerchner & Nicoll, 2008; J. Lisman, Yasuda, & Raghavachari, 2012). On the other hand, α CaMKII is also required for NMDAR-dependent hippocampal LTD. For example, both CaMKII inhibitor and α CaMKII knock out could block LTD in CA1 (Coultrap et al., 2014). Moreover, α CaMKII is activated during LTD expression (J. Y. Delgado et al., 2007; Lu, Isozaki, Roche, & Nicoll, 2010). In α CaMKII-F89G transgenic (TG) mice, α CaMKII overexpression in the forebrain impairs LTD in anterior cingulate and medial prefrontal cortices, and disrupts behavioral flexibility (J. Ma et al., 2015; Wei et al., 2006). However, whether and how α CaMKII in LA affect LTD at T-LA synapses and cued fear extinction are still unknown.

To better illuminate the mechanism of cued fear extinction, thereby understanding the mechanism of PTSD, using the behavioral profiling approach (Ardi, Albrecht, Richter-Levin, Saha, & Richter-Levin, 2016), we identified PTSD susceptible mice with cued fear extinction deficit and anxiety-like behaviors from the trauma-exposed mice. It is worth noting that increased α CaMKII was detected in LA of PTSD susceptible mice. To determine whether increased α CaMKII can cause PTSD-like symptoms, we employed an inducible and reversible chemical-genetic technique to temporally and spatially manipulate α CaMKII level in the forebrain of α CaMKII-F89G TG mice, as well as using adeno-associated viral (AAV) vectors to elevate α CaMKII specifically in LA of C57BL/6J mice. Consistently, up-regulation of α CaMKII induced PTSD-like symptoms including cued fear extinction deficit and anxiety-like behaviors, which could be reversed by suppressing elevated α CaMKII to normal level. In addition, we prove that GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization are the links between cued fear extinction and NMDAR-dependent LTD at T-LA synapses.

RESULTS

PTSD susceptible mice exhibit increased α CaMKII and reduced AMPAR internalization in LA.

PTSD susceptible individuals were identified in UWT-exposed group (23 male mice) and 4-CS/US-exposed group (23 male mice) by employing the behavioral profiling approach described in MATERIALS AND METHODS section. PTSD susceptible mice had persistently higher level of cued freeze responses through extinction trials (Fig. 1B, PS-UWT vs Control, $F_{(4, 85)} = 6.33$, $P < 0.001$; PS-4CS/US vs Control, $F_{(4, 85)} = 4.70$, $P < 0.01$), spent significantly less time in the center area of open field (OF) chamber (Fig. 1C, PS-UWT vs Control, $P < 0.01$; PS-4CS/US vs Control, $P < 0.001$), in the light zone of light/dark box (LD) test (Fig. 1D, PS-UWT vs Control, $P < 0.01$; PS-4CS/US vs Control, $P < 0.001$), and in the open arms of water zero maze (OM) test (Fig. 1E, PS-UWT vs Control, $P < 0.001$; PS-4CS/US vs Control, $P < 0.001$) compared with control mice. Behavioral profiling revealed that only 7 mice each group showed PTSD-like symptoms in 23 mice exposed to UWT or 23 mice exposed repeatedly to US/CS.

LA is a key brain region for fear extinction and anxiety-like behaviors (Erlich, Bush, & Ledoux, 2012; Forster, Novick, Scholl, & Watt, 2012; Grosso, Santoni, Manassero, Renna, & Sacchetti, 2018; Jacques et al., 2019; Jihye Kim et al., 2015; J. Kim et al., 2007; Krabbe, Gründemann, & Lüthi, 2018; Mahan & Ressler, 2012; Ressler, 2010; Schafe, Doyère, & LeDoux, 2005). CaMKII has been shown to be important for memory extinction (Bevilaqua et al., 2006; Burgdorf et al., 2017; Szapiro, Vianna, McGaugh, Medina, & Izquierdo, 2003). Moreover, GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization contribute to fear extinction (Bai et al., 2014; Dalton et al., 2008; Hollis, Sevelinges, Grosse, Zanoletti, & Sandi, 2016; J. Kim et al., 2007; S. Lee et al., 2013; Lin et al., 2010; Talukdar, Inoue, Yoshida, & Mori, 2018). Thus, we investigated levels of CaMKII, GluA1-Ser845/Ser831 phosphorylation and synaptic GluA1/2 expression in LA of PTSD susceptible mice and found α CaMKII and the phosphorylated (p)- α CaMKII at Thr286 (p- α CaMKII-Thr286) were significantly up-regulated in PTSD susceptible mice experienced either UWT or 4-CS/US exposure (Fig. 1FG, PS-UWT vs Control, α CaMKII, $P < 0.01$, p- α CaMKII-Thr286, $P < 0.05$; PS-4CS/US vs Control, α CaMKII, $P < 0.05$, p- α CaMKII-Thr286, $P < 0.05$). However, no significant difference was observed in β CaMKII among the three groups (Fig. 1F, PS-UWT vs Control, $P > 0.05$; PS-4CS/US vs

Control, $P > 0.05$). In addition, PTSD susceptible mice had a significant higher synaptic expression levels in the synaptic GluA1/2 expression and phosphorylated GluA1-Ser845/Ser831 (Fig. 1F, PS-UWT vs Control, GluA1: $P < 0.01$, GluA2: $P < 0.01$, GluA1-Ser831: $P < 0.01$, GluA1-Ser845: $P < 0.01$; PS-4CS/US vs Control, GluA1: $P < 0.001$, GluA2: $P < 0.05$, GluA1-Ser831: $P < 0.01$, GluA1-Ser845: $P < 0.01$). Taken together, these results suggest that PTSD susceptible mice display the significantly higher level of α CaMKII, the lower level of GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization in LA.

Increasing α CaMKII in LA is sufficient to cause PTSD-like phenotypes in both α CaMKII-F89G TG and AAV- α CaMKII mice

To further investigate whether elevated α CaMKII in LA cause PTSD-like phenotypes such as impaired fear extinction and anxiety-like behaviors, we temporally and spatially manipulated α CaMKII overexpression in α CaMKII-F89G TG mice by employing an inducible and reversible chemical-genetic technique described in MATERIALS AND METHODS section. The higher level of α CaMKII and normal morphology in LA were observed in TG mice (Supplemental information, Fig. S1).

Then, cued fear memory recall and cued fear extinction were measured after only 1-CS / US for cued fear conditioning (Fig. 2A). Given that forebrain α CaMKII overexpression impairs fear memory retrieval in our previous study (Cao et al., 2008), to examine the effect of α CaMKII overexpression on cued fear extinction in TG mice, we designed the “normal α CaMKII level during cued fear memory retrieval but elevated α CaMKII level during cued fear extinction period” paradigm by a single i.p. injection of NM-PP1 into both TG mice and WT littermates 15 mins before the first recall test of cued fear memory (Fig. 2A). Under this paradigm, TG mice exhibited normal retrieval of cued fear memory in comparison to that of wild-type littermate (Fig. 2B, TG + i.p. vs WT + i.p., $P > 0.05$). However, during cued fear extinction trials, as shown in Fig. 2B, a significant declining freezing behavior was observed in WT mice but not in TG mice (Fig. 2B, TG + i.p. vs WT + i.p., $F_{(3, 264)} = 10.73$, $P < 0.001$). A *post hoc* analysis revealed that TG mice exhibited significantly higher level of freezing response to the CS in cued fear extinction trial 2, 3 and 4 (Fig. 2B, TG + i.p. vs WT + i.p., $P < 0.05$), suggesting that elevated α CaMKII may impair cued fear extinction. In addition, TG mice spent significantly less time (Fig. 2C-E, TG + i.p. vs WT + i.p., $P < 0.001$) in the center area of OF chamber (Fig. 2C), in the

light zone of LD test (Fig. 2D), and in the open arms of elevated plus maze (EPM) test (Fig. 2E) compared with WT mice. Together, it indicates that increased α CaMKII in LA may cause PTSD-like phenotypes.

To further confirm whether PTSD-like phenotypes in TG mice are due to the overexpression of α CaMKII-F89G protein, we then designed the “normal α CaMKII level during both fear memory recall and extinction period” paradigm by i.p. injection of NM-PP1 15 min before recall test and oral (p.o.) administration throughout the entire fear extinction period (Fig. 2B). Under this “normal α CaMKII level during both fear memory recall and extinction period” paradigm, TG mice had similar freezing response with that in WT mice during cued extinction trials (Fig. 2B, TG + i.p. + o.p. vs WT + ip, $P > 0.05$), suggesting impaired cued fear extinction was rescued by NM-PP1 treatment in TG mice. Moreover, NM-PP1 had no effect on cued fear extinction in WT mice (Fig. 2B, WT + i.p. + o.p. vs. WT + i.p., $P > 0.05$), excluding the possibility that the rescuing effects by NM-PP1 were due to ‘facilitating extinction’ effects. In addition, TG mice with NM-PP1 treatments spent comparable amounts of time (Fig. 2C-E, TG + i.p. + p.o. vs. WT + i.p., $P > 0.05$) in the center area of OF chamber (Fig. 2C), in the light box of LD test (Fig. 2D) and in EPM test (Fig. 2E) compared with WT mice. Furthermore, TG mice without any treatment exhibited normal locomotor activity, exploratory behavior and pain threshold (Supplemental information, Fig. S2). Taken all together, we conclude that increased α CaMKII indeed is sufficient to produce PTSD-like phenotypes including impaired fear extinction and anxiety-like behaviors.

To further examine whether increasing α CaMKII specifically in LA is also sufficient to cause PTSD-like phenotypes, we bilaterally injected viral vectors AAV- α CaMKII (pAAV-TRE- α CaMKII-P2A-EGFP-CMV-rTA) into LA of C57BL/6J mice to overexpress α CaMKII specifically in LA (Fig. 3A). As expected, both α CaMKII and p- α CaMKII-Thr286 expression levels significantly increased in LA of AAV- α CaMKII mice (Fig. 3DE, α CaMKII, $P < 0.001$; p- α CaMKII-Thr286, $P < 0.001$). 24 h after 1-CS/US pairing, we performed cued fear memory test. AAV- α CaMKII mice exhibited impairment of cued fear memory during recall test (Fig. 3C, $P < 0.001$), which is consistent our previous finding that α CaMKII overexpression impairs the retrieval of fear memory (Cao et al., 2008). In addition, AAV- α CaMKII mice showed significantly impaired fear extinction (Fig. 3C, AAV- α CaMKII vs AAV-control, $F_{(4, 81)} = 2.63$, P

< 0.05). A post hoc analysis revealed that AAV- α CaMKII mice exhibited the significant higher freezing responses than AAV-control mice on the 4th extinction trial ($P < 0.05$). In addition to deficits in cued fear extinction, AAV- α CaMKII mice showed anxiety-like behaviors (data not shown). These data suggest that elevated α CaMKII expression specifically in LA is also sufficient to result in PTSD-like symptoms.

Increasing α CaMKII in LA impairs AMPAR internalization and GluA1-Ser845/Ser831 dephosphorylation after cued fear extinction in both α CaMKII-F89G TG and AAV- α CaMKII mice

We quantified the expression of synaptic AMPAR composition subunits (GluA1/2) and GluA1-Ser845/Ser831 phosphorylation in LA before/after cued fear conditioning and extinction trials. After cued fear extinction trials, compared with cued fear conditioning trial, significant decreases in the GluA1/2 synaptic expression and GluA1-Ser845/Ser831 phosphorylation levels could be found only in WT mice (Fig. 2FG, WT + FC vs WT + Ext, GluA1: $P < 0.001$; GluA2: $P < 0.01$; pGluA1-Ser845: $P < 0.05$; pGluA1-Ser831: $P < 0.01$), but not in TG mice (TG + FC vs TG + Ext, GluA1/A2, pGluA1-Ser845/831: $P > 0.05$). Furthermore, the GluA1/2 synaptic expression and phosphorylated GluA1-Ser845/Ser831 in TG mice were significantly higher than that in WT mice after cued fear extinction trials (Fig. 2FG, TG + Ext vs WT + Ext, GluA1/A2: $P < 0.01$; pGluA1-Ser845/Ser831: $P < 0.05$).

Moreover, consistent with the above western blotting data from α CaMKII-F89G TG mice, synaptic GluA1/2 expression, phosphorylated GluA1-Ser845/Ser831 levels were significantly higher in LA of AAV- α CaMKII mice than that in AAV-control mice after cued fear extinction trials (Fig. 3DE, GluA1/2, pGluA1-Ser845/Ser831, $P < 0.01$). Taken all together, these results indicate that increasing α CaMKII specifically in LA disrupts GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization, consequently may impair cued fear extinction in both α CaMKII-F89G TG and AAV- α CaMKII mice.

Increasing α CaMKII impairs NMDAR-dependent LTD at T-LA synapses and NM-PP1 can recover the impairments

To investigate the cellular mechanism of impaired cued fear extinction, we measured the basal synaptic transmission and synaptic plasticity at T-LA synapses in TG mice. No significant difference was observed in input-output curves, synaptic and total GluA1/2 expression of LA

(Supplemental information, Fig. S3AB, TG vs WT, $P > 0.05$), paired-pulse depression (PPD) and synapsin expression (Fig. S3CD, TG vs WT, $P > 0.05$) in LA between TG and WT mice. Moreover, either tetanic or theta burst stimulations induced similar level of LTP at T-LA synapses (Fig. 4AB, TG vs WT, $P > 0.05$). These results indicate that α CaMKII overexpression does not affect basal synaptic transmission and LTP at T-LA synapses.

We then analyzed the effects of the α CaMKII overexpression on LTD at T-LA synapses. 1Hz-LTD in TG slices was significantly reduced (Fig. 4C, TG vs WT, $P < 0.05$) compared to that of WT slices, which could be recovered by 0.5 μ M NM-PP1 (Fig. 4C, TG + NM-PP1 vs TG, $P < 0.05$), while 1Hz-LTD in WT slices was not affected (Fig. 4C, WT + NM-PP1 vs WT, $P > 0.05$). Notably, LTD at the T-LA synapses could be blocked by application of APV (50 μ M) and NM-PP1 (0.5 μ M) (Fig. 4D, TG vs WT, $P > 0.05$), suggesting the LTD at the T-LA synapses is NMDAR-dependent. Besides, 3Hz-LTD was blocked in TG slices (Fig. 4E; TG vs WT, $P < 0.001$), which could also be recovered by NM-PP1 (Fig. 4E, TG + NM-PP1 vs TG, $P < 0.01$), while 3Hz-LTD in WT slices was not affected (Fig. 4E, WT + NM-PP1 vs WT, $P > 0.05$). Likewise, TG mice exhibited deficits in the depotentiation at T-LA synapses (Fig. 4F, TG vs WT, $P < 0.01$). In summary, our results show that α CaMKII overexpression impairs NMDAR-dependent LTD and depotentiation at T-LA synapses in TG mice.

Increasing α CaMKII impairs AMPAR internalization and GluA1-Ser845/Ser831 dephosphorylation during NMDAR-dependent LTD and NM-PP1 can rescue the impairments

Besides the low-frequency stimulation (LFS), brief NMDA exposure can chemically induce NMDAR-dependent LTD (H. K. Lee, K. Kameyama, R. L. Huganir, & M. F. Bear, 1998). In TG slices, NMDA application (30 μ M, 3 min) could elicit a significantly weaker LTD at T-LA synapses than that in WT slices (Fig. 5A, TG + NMDA vs WT + NMDA, $P < 0.01$). Furthermore, 0.5 μ M NM-PP1 could rescue the reduced NMDA-induced LTD in TG slices to normal level (Fig. 5A, TG + NMDA + NM-PP1 vs WT + NMDA, $P > 0.05$; TG + NMDA + NM-PP1 vs TG + NMDA, $P < 0.01$), but had no detectable effects on NMDA-induced LTD in WT slices (Fig. 5A, WT + NMDA + NM-PP1 vs WT + NMDA, $P > 0.05$). These results suggest that increasing α CaMKII in LA attenuates NMDAR-dependent chem-LTD at T-LA synapses in TG mice.

NMDA-induced LTD could elicit more widespread depression of synapse strength and share the similar molecular mechanisms to LFS-LTD such as AMPAR internalization and GluA1-Ser845/Ser831 phosphorylation (Jary Y. Delgado et al., 2007; He, Lee, Song, Kanold, & Lee, 2011; Kollen, Dutar, & Jouvenceau, 2008; H.-K. Lee, K. Kameyama, R. L. Huganir, & M. F. Bear, 1998). To investigate the molecular mechanisms underlying deficit in NMDAR-dependent LTD in TG mice, we examined the amount of some synaptic proteins after NMDA-induced LTD. NMDA application significantly decreased the GluA1/2 synaptic expression and GluA1-Ser845/Ser831 phosphorylation in WT slices (Fig. 5BC, WT+NMDA vs WT, GluA1, GluA2, pGluA1-Ser845: $P < 0.01$; pGluA1-Ser831: $P < 0.001$), but not in TG slice (Fig. 5BC, TG+NMDA vs TG, GluA1/A2, pGluA1-Ser845/831: $P > 0.05$). Besides, the synaptic expression of GluA1/2 and GluA1-Ser845/Ser831 phosphorylation in LA of TG slices were significantly higher than that in LA of WT slices (Fig. 5BC, TG + NMDA vs WT + NMDA, GluA1: $P < 0.05$; GluA2: $P < 0.01$; pGluA1-Ser845/Ser831: $P < 0.05$). Furthermore, NM-PP1 (0.5 μ M) successfully rescued the impairments of AMPAR internalization and GluA1-Ser845/Ser831 dephosphorylation of LA in TG slices (Fig. 5BC, TG+NMDA+NM-PP1 vs TG, GluA1/A2, pGluA1-Ser845/831: $P < 0.01$), with no effect on that of WT slices (Fig. 5BC, WT+NMDA+NM-PP1 vs WT+NMDA, GluA1/A2, pGluA1-Ser845/831: $P > 0.05$). Collectively, it indicates that α CaMKII overexpression leads to impairment of AMPARs internalization and dephosphorylation in LA, which consequently impairs NMDAR-dependent LTD at T-LA synapses.

Increasing α CaMKII reduces protein phosphatase (PP) activity and enhances stargazin expression during NMDAR-dependent LTD and NM-PP1 can recover the abnormalities

Activation of protein phosphatase 1 (PP1) contributes to LTD formation (Isabelle M. Mansuy & Shirish Shenolikar, 2006; Mauna, Miyamae, Pulli, & Thiels, 2011). Moreover, stargazin can be dephosphorylated by PP1 to induce the clathrin-dependent AMPAR endocytosis during NMDAR-dependent LTD (Bats, Groc, & Choquet, 2007; Matsuda et al., 2013). Dephosphorylation of the Thr320 residue on the C-terminal domain of PP1 can enhance PP1 activity during NMDAR-dependent LTD (Dohadwala et al., 1994; Goldberg et al., 1995). Therefore, we investigated PP1-Thr320 phosphorylation (pPP1-Thr320) and stargazin expression in LA fractions of WT and TG slices with NMDA treatment. With NMDA exposure, significant reductions of pPP1-Thr320 and stargazin expression of LA could be found only in WT (Fig. 5DE,

WT+NMDA vs WT, pPP1-Thr320: $P < 0.01$; stargazin: $P < 0.001$) but not in TG slices (TG+NMDA vs TG, pPP1-Thr320, stargazin: $P > 0.05$). Moreover, the PP1-Thr320 phosphorylation and stargazin expression in LA of TG slices were dramatically higher than that in WT slices (Fig. 5DE, TG + NMDA vs WT + NMDA, pPP1-Thr320, stargazin, $P < 0.05$), suggesting that the PP1 activity and stargazin expression were abnormal in TG mice during NMDA-induced LTD. Furthermore, NM-PP1 could recover the abnormalities in PP1 activity and stargazin expression in LA of TG slices (Fig. 5DE, TG + NMDA + NM-PP1 vs TG, pPP1-Thr320: $P < 0.01$; stargazin: $P < 0.01$; TG + NMDA + NM-PP1 vs TG + NMDA, pPP1-Thr320: $P < 0.05$, stargazin: $P < 0.05$) but not affecting that of WT slices (Fig. 5DE, WT + NMDA + NM-PP1 vs WT + NMDA, pPP1-Thr320: $P > 0.05$; stargazin: $P > 0.05$).

Protein phosphatase 2A (PP2A) and calcineurin (PP2B) play important roles in LTD maintenance and induction (Pi & Lisman, 2008; Winder & Sweatt, 2001). A significant augment of PP2A/2B activity could be found in LA of WT slices (Fig. 5F, WT+NMDA vs WT, PP2A: $P < 0.01$; PP2B: $P < 0.001$), but not in LA of TG slices during LTD formation (Fig. 5F, TG+NMDA vs TG, PP2A, PP2B: $P > 0.05$). Besides, PP2A/2B activity was dramatically lower in LA of TG slices than that of WT slices (Fig. 5F, TG + NMDA vs WT + NMDA, PP2A: $P < 0.01$; PP2B: $P < 0.001$), during NMDA-induced LTD. Furthermore, NM-PP1 (0.5 μ M) could also recover PP2A/2B activity down-regulation in LA of TG slices (Fig. 5F, TG + NMDA + NM-PP1 vs TG, PP2A, PP2B: $P < 0.001$; TG + NMDA + NM-PP1 vs TG + NMDA, PP2A, PP2B: $P < 0.01$) without affecting that of WT slices (Fig. 5F, WT + NMDA + NM-PP1 vs WT + NMDA, PP2A, PP2B: $P > 0.05$). Taken together, all these results suggest that α CaMKII overexpression can weaken PP1, PP2A/2B activity and increase stargazin expression in LA fractions during NMDAR-dependent LTD, which may be potential mechanisms of AMPAR internalization and NMDAR-dependent LTD impairments.

DISCUSSION

In the present study, we reveal that PTSD susceptible mice exhibits significant up-regulation of α CaMKII, down-regulation of GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization in LA. Consistently, increasing α CaMKII specifically in LA can cause PTSD-like phenotypes such as fear extinction deficit and anxiety-like behaviors, and impairs AMPAR internalization and dephosphorylation, NMDAR-dependent LTD and depotentiation at T-LA synapses. Furthermore, deficits in AMPAR internalization and dephosphorylation are observed not only after impaired cued fear extinction *in vivo*, but also after attenuated NMDA-induced LTD in TG slices *in vitro*. Additionally, the deficits in AMPAR internalization and dephosphorylation are due to down-regulation of PP1/2A, PP2B activity and increased stargazin in TG mice. Importantly, NM-PP1, a specific inhibitor of the exogenous α CaMKII-F89G, could rescue the above deficits in α CaMKII-F89G TG mice. These data suggest up-regulation of α CaMKII may weaken activity of PP1/2A and PP2B, increase stargazing, thereby impairing AMPAR internalization and dephosphorylation, which consequently impairs LTD and fear extinction.

α CaMKII and memory extinction

CaMKII has been shown to play an important role in the extinction of different memories. Pharmacological inhibition of CaMKII by KN-62 blocked the extinction of step-down passive avoidance performance (Bevilaqua et al., 2006; Szapiro et al., 2003). Similarly, α/β CaMKII inhibitor KN93 significantly attenuated the extinction of cocaine conditioned place preference (Burgdorf et al., 2017). Furthermore, partial reduction of α CaMKII function due to the T286A^{+/-} mutation impaired the extinction of contextual fear and spatial memories (Kimura, Silva, & Ohno, 2008). On the contrary, reduction of α CaMKII by phosphorylation at serine 331 in LA enhances cocaine memory extinction (Rich et al., 2016). Besides, increased activation of CaMKII α in the CPEB3-knockout hippocampus reduced the extinction of spatial memories (Berger-Sweeney, Zearfoss, & Richter, 2006; Huang, Chao, Tsai, Chung, & Huang, 2014). In our study, we found that mouse models of PTSD with cued fear extinction deficit exhibited significant up-regulation of α CaMKII in LA. Furthermore, increasing α CaMKII in LA can cause PTSD-like phenotypes including impaired cued fear extinction.

The causal relationship between elevated α CaMKII and impaired LTD

CaMKII is a major kinase mediating AMPAR trafficking and NMDAR-dependent synaptic plasticity (Collingridge et al., 2004). Specifically, CaMKII can phosphorylate AMPA receptors GluA1 subunits at Ser845/Ser831, which can promote the integration of new AMPA receptors at the postsynaptic density (Barria, Muller, Derkach, Griffith, & Soderling, 1997), further enhancing synaptic transmission. On the contrary, CaMKII has been found to interact with Arc/Arg3.1 gene product to weaken synapses by promoting AMPA internalization (Okuno et al., 2012). Recently, CaMKII has been also shown to phosphorylate GluA1 subunits at Ser567 site to promote P2X2-mediated AMPAR internalization and drive synaptic depression (Pougnnet et al., 2016). In our study, we found that PTSD susceptible mice with blocked fear extinction exhibited significantly higher α CaMKII, lower GluA1-Ser845/Ser831 dephosphorylation and lower AMPA internalization in LA. To investigate whether elevated α CaMKII led to PTSD-like symptoms including impaired fear extinction, changed NMDAR-dependent LTD, GluA1 dephosphorylation and AMPA internalization in LA, we then up-regulated α CaMKII expression in α CaMKII-F89G TG and AAV- α CaMKII infected mice. We found that α CaMKII overexpression in LA caused impairments in GluA1-Ser845/Ser831 dephosphorylation, AMPA internalization, NMDAR-dependent LTD at T-LA synapses and cued fear extinction in TG mice, which could be completely rescued by a specific inhibitor (NM-PP1) of exogenous α CaMKII-F89G. These results suggest there is causality between up-regulated α CaMKII and impaired GluA1-Ser845/Ser831 dephosphorylation, defective AMPA internalization, NMDAR-dependent LTD and cued fear extinction.

The molecular links between LTD and fear extinction

NMDAR mediates both LTD and fear extinction (Bai et al., 2014; Brebner et al., 2005; Dalton et al., 2008; Fox, Russell, Titterness, Wang, & Christie, 2007; Radulovic, Ren, & Gao, 2019). It has been reported that a GluR2-derived peptide (Tat-GluR23Y) blocked AMPAR internalization and impaired NMDAR-dependent LTD both *in vitro* (Bai et al., 2014; Brebner et al., 2005; Dalton et al., 2008) and *in vivo* (Fox et al., 2007). Moreover, NMDA NR2B receptors antagonist (Ro25-6981) blocked AMPAR internalization and disrupted fear extinction (J. Kim et al., 2007). Conversely, systemic administration of d-serine enhanced both AMPAR internalization and fear extinction (Bai et al., 2014). In addition, GluA1-Ser845/Ser831

dephosphorylation also played important roles in NMDAR-dependent LTD (Diering, Heo, Hussain, Liu, & Huganir, 2016) and fear extinction (Hollis et al., 2016; Talukdar et al., 2018). Although the above findings indicate AMPAR internalization and dephosphorylation may be links between fear extinction and LTD, supporting evidence is still lacking. In our current study, deficits in GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization were observed not only after impaired cued fear extinction *in vivo*, but also after attenuated NMDA-induced LTD in α CaMKII-F89G TG slices *in vitro*. Furthermore, a specific inhibitor of the exogenous α CaMKII-F89G (NM-PP1) could completely rescue the deficits in cued fear extinction, NMDA-induced LTD, GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization. Thus, our data demonstrate that deficits in Ser845/GluA1-Ser831 dephosphorylation and AMPAR internalization by elevated α CaMKII are molecular links between impaired NMDAR dependent-LTD and fear extinction. In other words, we demonstrate for the first time that GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization are molecular links between NMDA dependent-LTD and fear extinction.

How does excessive α CaMKII impair AMPAR internalization and dephosphorylation during NMDAR-dependent LTD?

LTD formation requires PPs (PP1, PP2A and PP2B) activation (Kameyama, Lee, Bear, & Huganir, 1998; H. K. Lee et al., 1998). Activated PPs dephosphorylate GluA1-Ser845/Ser831 (Hu, Huang, Yang, & Xia, 2007; I. M. Mansuy & S. Shenolikar, 2006; Winder & Sweatt, 2001), which cause a reduction of open probability or conductance for AMPAR channels and finally contribute to LTD formation. Specifically, PP1 is activated through a Ca^{2+} -PP2B-I1 pathway and has a more predominant role in depressing potentiated synapses, whereas PP2A is activated through PP2B/PP1 cascade or pathways independent on PP2B and mainly depresses naive synapses (Winder & Sweatt, 2001). However, in α CaMKII-F89G TG mice, α CaMKII overexpression could exhibit higher potency in the competition with PP2B for Ca^{2+} /CaM, which might decrease the accessibility of PP2B to Ca^{2+} /CaM and inhibit the activity of the PP2B-I1-PP1 pathway, thereby inhibiting PP1 activity. In addition, high concentration of phosphorylated α CaMKII could saturate the dephosphorylation ability of PP1, and thereby weaken PP1 dephosphorylating GluA1-Ser845 or GluA1-Ser831 (Hu et al., 2007; H. K. Lee, Barbarosie, Kameyama, Bear, & Huganir, 2000; J. E. Lisman & Zhabotinsky, 2001; I. M. Mansuy & S.

Shenolikar, 2006; Winder & Sweatt, 2001). Unlike PP1, PP2A can be directly inactivated by CaMKII through phosphorylating its B' α subunits (Fukunaga et al., 2000; Pi & Lisman, 2008), so excessive CaMKII can weaken PP2A activity. Collectively, one explanation for impairment of NMDAR-dependent LTD is that the excessive α CaMKII can lower activity of PPs, thereby reduce GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization, and consequently impair LTD.

It has been shown that stargazin can be dephosphorylated by PP1 through Ca^{2+} -PP2B-I1 pathway and form a ternary complex with APs to promote AMPAR internalization during NMDAR-dependent LTD (Matsuda et al., 2013; Tomita et al., 2003). Conversely, stargazin can be directly phosphorylated by activated CaMKII and bind to PSD-95 to immobilize AMPARs at synapses, which contributes to LTP (Bats et al., 2007; Opazo et al., 2010). In α CaMKII-F89G TG mice, more stargazin is expressed at the synaptic sites during NMDAR-dependent LTD. Therefore, another explanation for impairment of NMDAR-dependent LTD is that excessive CaMKII weakens AMPAR internalization through directly increasing stargazin phosphorylation and indirectly reducing stargazin dephosphorylation caused by lower PP1 activity, and finally impairs LTD.

CONCLUSION

We have found that PTSD-susceptible mice exhibit the higher α CaMKII expression, and lower GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization in LA. Increasing α CaMKII leads to PTSD-like phenotypes such as impaired fear extinction and anxiety-like behaviors, and impairs LTD at T-LA synapses. Furthermore, diminished GluA1-Ser845/Ser831 dephosphorylation and AMPAR internalization were observed not only after impaired fear extinction *in vivo*, but also after attenuated NMDA-induced LTD in TG slices *in vitro*. Further data suggest that the impairment of NMDAR-dependent LTD is caused by the defective PPs activity and the excessive synaptic stargazin in α CaMKII-F89G TG mice. In summary, α CaMKII may be identified as a powerful regulator of the core symptoms of PTSD and LTD at T-LA synapses, and may be a key molecular determinant of PTSD.

MATERIALS AND METHODS

Animals

Biochemical Characterizations of α CaMKII-F89G TG mice.

α CaMKII-F89G TG mice were donated by Dr. Tsien's lab (Wang et al., 2003). Mutant α CaMKII-F89G was generated with silent mutation (i.e. replacing the Phe-89 with Gly in α CaMKII), so that the ATP-binding pocket of α CaMKII-F89G kinase was enlarged. To selectively block exogenous α CaMKII-F89G and leave endogenous α CaMKII intact, NM-PP1 was designed to fit only this enlarged pocket but not the unmodified pocket of native α CaMKII. By using α CaMKII promoter-driven construct, we were able to overexpress α CaMKII-F89G in the forebrain neurons. The α CaMKII-F89G could be rapidly and selectively manipulated in the mouse forebrain by intraperitoneal (i.p.) injection or noninvasive oral intake of 1-Naphthylmethyl (NM)-PP1. Specifically, a single i.p. injection of NM-PP1(16.57 ng/g) into freely behaving TG mice could completely suppress α CaMKII-F89G in the forebrain regions of TG mice within 15 minutes and the complete suppression could be maintained for 40 min. The oral intake (5 μ M NM-PP1 in drinking water) could result in partial inhibition of α CaMKII-F89G in the TG mice by 6 h (no inhibition for the initial 3 h) and complete inhibition by 24 h. Bath application of 0.5 μ M NM-PP1 in the slices of TG mice could inhibit α CaMKII-F89G but had no effect on native α CaMKII (Wang et al., 2003).

All experimental procedures were conducted according to Animals Act, 2006 (China) and approved by the Institutional Animal Care and Use Committee (IACUC approval ID #M09018) of the East China Normal University. All mice were male and 3-4 months old. C57BL/6J mice were used for Figure 1 and 3. α CaMKII-F89G transgenic mice and wild-type littermates were used for the rest of Figures. The mice were housed in 12 h light/12 h dark cycle (lights on at 7 a.m.) with free access to food and water.

Behavior experiments

Behavioral profiling for identification of PTSD susceptible mice

We applied a behavioral profiling approach (Ardi et al., 2016) to identify PTSD susceptible mice in either underwater trauma (UWT)-exposed mice (G. Ritov, Boltyansky, & Richter-Levin, 2016) or 4 conditioned stimulus /unconditioned stimulus (4-CS/US)-exposed mice (Borghans & Homberg, 2015; Dbiec, Bush, & LeDoux, 2011; Fenster, Lebois, Ressler, & Suh, 2018; Ji et al.,

2014a, 2014b; Mahan & Ressler, 2012; Mohammed R. Milad & Quirk, 2011; Radulovic et al., 2019).

In detail, the C57BL/6J mice were randomly divided into three groups: control group (n = 12), UWT-exposed group (n = 23) and 4-CS/US-exposed group (n = 23).

The control mice without any treatment were kept in home cages for 4 weeks. The UWT-exposed mice were individually allowed to swim freely for 5 s in a water-filled plastic tank, then submerged under water for 35 s using a metal net, next kept in their home cages for 4 weeks (Ardi et al., 2016; G. Ritov et al., 2016).

The 4-CS/US-exposed mice were individually placed in the chamber and allowed to explore the environment freely for 2 min, and then exposed to the conditioned stimulus (CS: 75 dB sound at 2800 Hz) for 30 s. At the last 2 s of tone stimulus, the unconditioned stimulus (US: 0.50 mA footshock, 2 s) was delivered. After 4-CS/US pairings with 2 min intertrial interval, mice were kept in the chamber for 2 min and then stayed in their home cages for 4 weeks.

Three groups were examined in the open field (OF) test, light/dark (LD) test, water zero maze (OM) test, fear conditioning and extinction experiments. The freezing behavior was monitored by Freeze Frame system (Coulbourn Instruments, USA).

We calculated six parameters: two parameters represent the level of locomotor activity and four parameters represent anxiety-like performances from the four experiments. To create the behavior profiles, firstly we referred to the performances of the control group as the behavior of the normal population and determined the distribution of values in the control group. Standard deviations were used to calculate the upper and lower “cut-off values” for each chosen parameter. Secondly, the performances of each mouse in the UWT-exposed group or 4-CS/US-exposed group were compared to the distribution curve of the control group. Each susceptible mouse must exhibit values that are under or above the lower and upper cut-off values in at least four out of the six parameters. “Cut-off values” of six parameters: the center time in the OF test, 560.32 ± 34.25 s; the time in the light box in the LD test, 788.60 ± 58.92 s; the time in the open arms in the OM test, 111.43 ± 8.88 s; the freezing percentage in the last day of cued fear extinction, $31.98\% \pm 3.91\%$; total distance in the OF test, 7059.99 ± 427.80 cm; total distance in the LD test, 9124.67 ± 220.50 cm.

Cued fear extinction

4 weeks after 4-CS/US pairings or 24 h after the 1-CS/US pairing, each mouse was placed into a novel chamber and monitored for 2 min (in the absence of the tone). For the recall test, the cued freezing responses to a 3 min tone (75 dB sound at 2800 Hz) without footshock were measured. Then, 4 cued fear extinctions trials were conducted like the recall test in the next 4 following days. Data were presented as the mean $\pm$ s.e.m. Two-way ANOVA was used for statistical analysis.

Open field

As described previously (Yan et al., 2015), briefly, each mouse was placed in an acrylic open-field chamber (27 cm long $\times$ 27 cm wide $\times$ 38 cm high) for 30 min. The amount of moving distance, the time in the center area, and the number of rearing were measured using a Tru-scan DigBahv-locomotion Activity Video Analysis System (Coulbourn Instruments, USA). Data were presented as the mean $\pm$ s.e.m. One-way ANOVA was used for statistical analysis in Fig. 1C, 2C and Student's t-test in Fig. S2A.

Light/dark test

The box (27 cm long $\times$ 27 cm wide $\times$ 38 cm high) was divided into two equal zones - light zone and dark zone. The light zone was painted white and illuminated by the white light while the dark zone was painted black and not illuminated. These two zones were connected by a door in the middle divider. Mice could shuttle freely between two boxes. The total distance and the time stayed in light zone were delineated by the Tru-scan DigBahv-locomotion Activity Video Analysis System (Coulbourn Instruments, USA) for 30 min. Data were presented as the mean $\pm$ s.e.m. One-way ANOVA was used for statistical analysis in Fig. 1D and 2D.

Water-associated zero maze task

Experimental protocol and device were similar as described previously (Gilad Ritov & Richter-Levin, 2014). This device was composed of an annular platform and a plastic bucket. The annular platform was divided into four equal quadrants - two open arms and two closed arms. The plastic bucket was full of water for 40 cm deep. After 5 min habituation, mouse was put into one of the open arms facing the closed arm for 5 min. The time spent in the open arms and

closed arms were measured by Any-maze system (USA, Stoelting). Data were presented as the mean $\pm$ s.e.m. One-way ANOVA was used for statistical analysis.

Elevated plus maze test

The apparatus consists of two opposed open arms (30 cm $\times$ 5 cm), two opposed closed arms (30 cm $\times$ 5 cm) and one open square (5 cm $\times$ 5 cm) in the center, which was elevated above the floor (50 cm). Each mouse was placed in the center of the plus maze with its face directing to an open arm and allowed to explore for 5 min. The time spent and moving distances in open and closed arms were automatically recorded by Any-maze system (USA, Stoelting). Data were presented as the mean $\pm$ s.e.m. One-way ANOVA was used for statistical analysis.

Animal surgery

To elevate α CaMKII specifically in LA of C57BL/6J mice, we injected pAAV-TRE- α CaMKII-P2A-EGFP-CMV-rTA (AAV- α CaMKII) or pAAV-TRE-P2A-EGFP-CMV-rTA (AAV-control) virus (2.45×10^{-12} and 2.38×10^{-12} vector genomes/ml, respectively, Obio Technology, China) bilaterally into LA (AP, -1.60 mm; ML, ± 3.35 mm; DV, -4.80 mm) of C57BL/6J mice. After the injection, mice were put back into home cages to recover for one month before experiments. AAV- α CaMKII mice were fed with doxycycline solution (1g/L in drinking water) to induce the virus expression throughout the behavior tests.

Dendritic spine analysis

Dendritic spine analysis were performed as previously described (Ming et al., 2018). Briefly, mice were deeply anaesthetized and transcardially perfused. 200 μ m coronal brain sections were cut and collected in 0.1 M PBS. LA neurons were loaded iontophoretically with a 5% Lucifer Yellow solution. Images of basal and apical dendrites of LA pyramidal neurons were scanned using a Leica SP2 confocal microscope at 63 $\times$ under oil immersion. The number of spines per micrometer along the dendritic longitudinal axis was counted as spine density. Data were presented as the mean $\pm$ s.e.m. Student's t-test was used for statistical analysis.

Sensitivity to foot shock

This test was performed according to the methods as published (Duan, Zhou, Ma, Yin, & Cao, 2015). Mice were individually placed in the conditioning chamber to receive 1 s shocks of gradually increasing current intensity by an increment of 0.01 mA (flinching, 0.05-0.1 mA;

vocalization, 0.1-0.2 mA; jumping, 0.45-0.6 mA) with 20 s intervals. The minimum current required to elicit flinching, vocalization and jumping in mice were measured. Data were presented as the mean $\pm$ s.e.m. Student's t-test was used for statistical analysis.

Amygdala slice electrophysiology

Protocols were similar as described previously (J. Kim et al., 2007; T. F. Ma et al., 2013). Mice (3-4 months old) were anaesthetized with sodium pentobarbital and sacrificed by decapitation. Whole brain coronal slices (370 μ m thick for fEPSPs recording) containing the amygdala were cut using a vibroslicer (vibratome 3000) with the cold (4 $^{\circ}$ C) and oxygenated (95% O₂ /5% CO₂) modified artificial cerebrospinal fluid (ACSF) containing (in mM): Choline chloride, 110; KCl, 2.5; CaCl₂, 0.5; MgSO₄, 7; NaHCO₃, 25; NaH₂PO₄, 1.25; D-glucose, 25; pH 7.4. The slices were recovered in an incubation chamber with normal ACSF containing (in mM): NaCl, 119; CaCl₂, 2.5; KCl, 2.5; MgSO₄, 1.3; NaHCO₃, 26.2; Na₂HPO₄, 1.0; D-glucose, 11, pH 7.4; 95% O₂ and 5% CO₂ for 60 min at 31 $^{\circ}$ C, and then returned to room temperature for at least 1 h before recording.

Field excitatory postsynaptic potential recording

A stimulating electrode was placed in the fibers from the internal capsule to activate the thalamic input to the lateral amygdala (T-LA) synapses. A recording electrode was positioned in LA to record field excitatory postsynaptic potential (fEPSP). Test responses were elicited at 0.033 Hz. After obtaining a stable baseline response for at least 15 min, LTP or LTD was induced. LTP was induced by applying 2 trains high frequency stimulation (100 Hz for 1 s) with 10 s interval or 3 trains theta burst stimulation (10 bursts delivered every 200 ms, each burst consisted of 4 pulses at 100 Hz) with 10 s interval. For LTD induction, the standard 1 Hz protocol (1 Hz for 15 min) and 3 Hz protocol (3 Hz for 5 min) were used. Depotentiation was induced by applying 2 trains of high frequency stimulation (100 Hz for 1s) with 10 s interval followed by the standard 1 Hz protocol (1 Hz for 15 min) after 20 min. For chemical-LTD induction, NMDA (Sigma, 30 μ M in ACSF) was infused into the slice chamber for 3 min. Data were presented as the mean $\pm$ s.e.m. Student's t-test (for comparing two different groups with Gaussian distribution) and one-way ANOVA followed by HSD post-hoc test with Bonferroni's correction (for comparing more than two different groups) were used for statistical analysis.

Proteins sample preparation

Combined with the previous protocol (Cui et al., 2011; Yin et al., 2013), synaptosomes were prepared as follows. LA tissues were homogenized in 1.5 ml homogenate-buffer (320 mM sucrose, 5 mM HEPES, pH 7.4) containing freshly added PMSF, PIC and PIC3. Homogenates were centrifuged at 500 g for 5 min to yield insoluble components. Then the supernatant fraction was collected and centrifuged at 10,000 g for 10 min to yield precipitation. The precipitation pellet was resuspended in 2 ml of 0.32 M sucrose, layered onto 2.25 ml of 0.8 M sucrose, and centrifuged at 98,000 g for 15 min using a swinging bucket rotor. Synaptosomes were collected from the 0.8 M sucrose layer and concentrated by centrifugation at 20,800 g for 45 min. Then the precipitation was resuspended in synaptosome lysis buffer (30 mM Tris (pH 8.5), 5 mM magnesium acetate, 8 M Urea, and 4% W/V CHAPS). For total proteins preparation, the LA areas were homogenized with RIPA buffer containing freshly added PMSF, PIC and PIC3 and lysed on ice for 30 min, centrifuged at 10,000 g at 4 °C for 5 min, and total proteins were taken as supernatant. Then the protein samples were stored in a -80°C freezer until used. Protein samples were quantified us by a Pierce BCA Protein Assay kit (Thermo Scientific) after which protein was stored at -20°C.

Western blot

Each sample of protein (5 µg/lane) was separated by 10% SDS-PAGE (P40650, NCM Biotech) and separated at 120 V for 120 minutes. Then the separated proteins were transferred onto a polyvinylidene fluoride (PVDF) membrane. The PVDF membranes were blocked in blocking solution (5% skim milk and 1% BSA) at room temperature for 1h. A reversible Ponceau S staining of the membranes was done to normalize the relative amount of each protein on the membrane (just for synaptosomes). After washing with TBST buffer, the PVDF membranes were immunoblotted with following antibodies: GluA1 antibody (1:2,000, Santa Cruz), GluA2 antibody (1:2,000, Millipore), pGluA1-Ser845 antibody (1:500, Abcam), pGluA1-Ser831 antibody (1:500, Abcam), αCaMKII antibody (1:3,000, Abcam), p-αCaMKII-Thr286 antibody (1:20,000, Santa Cruz), βCaMKII antibody (1:2,000, Invitrogen), β-actin antibody (1:20,000, Sigma), GAPDH antibody (1:20,000, Proteintech), synapsin (SYP) antibody (1:2,000, Proteintech), TfR antibody (1:2,000, Abcam), Tubulin antibody (1:1,000, Millipore) at 4°C for 12h. After washing with TBST buffer, the blots were reacted with an HRP-conjugated secondary antibody

at room temperature for 1 hour. Band intensity on the blot was quantified by the ECL immunoblotting detection system (Bio-rad). Data were shown as mean $\pm$ s.e.m.. Statistical differences were analyzed using post hoc test with Bonferroni's correction following one-way ANOVA.

PP2A activity measurement

PP2A activity was measured by using immunoprecipitation phosphatase assay kit according to the manufacturer's instructions (Catalog # 17-313, Millipore). Statistical differences were analyzed using post hoc test with Bonferroni's correction following one-way ANOVA. Data were shown as mean $\pm$ s.e.m.

PP2B activity measurement

The activity of calcineurin (PP2B) was assayed by using a calcineurin cellular activity assay kit (207007, Millipore) by following the manufacturer's instructions. Statistical differences were analyzed using post hoc test with Bonferroni's correction following one-way ANOVA. Data were shown as mean $\pm$ s.e.m.

Statistical analysis

Statistical significance was assessed by one-way ANOVA, two-way ANOVA analysis of variance or two-tailed, unpaired and paired t-tests, where appropriate. Significant effects in analysis of variances were followed up with Bonferroni post-hoc tests. Results were considered significantly different when $P < 0.05$. All data were presented as means $\pm$ s.e.m. The detail information about statistical analysis was provided in legends.

ACKNOWLEDGMENTS

We thank Dr. Xuechu Zhen and Dr. Liyong Li for their assistance with immunoblotting study, Dr. Kevan M. Shokat of UCSF for providing NM-PP1 inhibitor, and Dr. Bo Wang, Dr. Yihui Cui for their comments on the manuscript. This research was supported by fund from MOST China-Israel cooperation (No: 2016YFE0130500) and National Natural Science Fund of China (No: 31471077 and No: 31771177).

AUTHOR CONTRIBUTIONS:

S.A., J.W., X.Z., Y.D., J.L., D.W., H.Z., G.R.L., and X.C. designed the work; S.A., J.W., X.Z., J.L. and D.W. performed the acquisition of data for the work. S.A., J.W, X.Z., and X.C. analyzed and interpreted data; S.A., J.W, X.Z., G.R.L., and X.C. wrote the manuscript.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ADDITIONAL INFORMATION

Supplementary information (Fig. S1, S2, S3A) accompanies this paper.

REFERENCES

- Ardi, Z., Albrecht, A., Richter-Levin, A., Saha, R., & Richter-Levin, G. (2016). Behavioral profiling as a translational approach in an animal model of posttraumatic stress disorder. *Neurobiol Dis*, 88, 139-147. doi:10.1016/j.nbd.2016.01.012
- Bai, Y., Zhou, L., Wu, X., & Dong, Z. (2014). d-Serine enhances fear extinction by increasing GluA2-containing AMPA receptor endocytosis. *Behavioural Brain Research*, 270, 223-227. doi:<https://doi.org/10.1016/j.bbr.2014.05.025>
- Barria, A., Muller, D., Derkach, V., Griffith, L. C., & Soderling, T. R. (1997). Regulatory phosphorylation of AMPA-type glutamate receptors by CaM-KII during long-term potentiation. *Science*, 276(5321), 2042-2045. doi:DOI 10.1126/science.276.5321.2042
- Bats, C., Groc, L., & Choquet, D. (2007). The Interaction between Stargazin and PSD-95 Regulates AMPA Receptor Surface Trafficking. *Neuron*, 53(5), 719-734. doi:<https://doi.org/10.1016/j.neuron.2007.01.030>
- Bennett, M. R., Arnold, J., Hatton, S. N., & Lagopoulos, J. (2017). Regulation of fear extinction by long-term depression: The roles of endocannabinoids and brain derived neurotrophic factor. *Behav Brain Res*, 319, 148-164. doi:10.1016/j.bbr.2016.11.029
- Berger-Sweeney, J., Zearfoss, N., & Richter, J. (2006). Reduced extinction of hippocampal-dependent memories in CPEB knockout mice. *Learn Mem*, 13, 4-7. doi:10.1101/lm.73706
- Bevilaqua, L. R., Bonini, J. S., Rossato, J. I., Izquierdo, L. A., Cammarota, M., & Izquierdo, I. (2006). The entorhinal cortex plays a role in extinction. *Neurobiol Learn Mem*, 85(2), 192-197. doi:10.1016/j.nlm.2005.10.001
- Borghans, B., & Homberg, J. R. (2015). Animal models for posttraumatic stress disorder: An overview of what is used in research. *World J Psychiatry*, 5(4), 387-396. doi:10.5498/wjp.v5.i4.387
- Brebner, K., Wong, T. P., Liu, L., Liu, Y., Campsall, P., Gray, S., . . . Wang, Y. T. (2005). Nucleus accumbens long-term depression and the expression of behavioral sensitization. *Science*, 310(5752), 1340-1343. doi:10.1126/science.1116894
- Burgdorf, C. E., Schierberl, K. C., Lee, A. S., Fischer, D. K., Van Kempen, T. A., Mudragel, V., . . . Rajadhyaksha, A. M. (2017). Extinction of Contextual Cocaine Memories Requires Ca(v)1.2 within D1R-Expressing Cells and Recruits Hippocampal Ca(v)1.2-

- 1 Dependent Signaling Mechanisms. *J Neurosci*, 37(49), 11894-11911.
2 doi:10.1523/jneurosci.2397-17.2017
- 3 Cao, X., Wang, H., Mei, B., An, S., Yin, L., Wang, L. P., & Tsien, J. Z. (2008). Inducible and
4 selective erasure of memories in the mouse brain via chemical-genetic manipulation.
5 *Neuron*, 60(2), 353-366.
- 6 Colbran, R. J., & Soderling, T. R. (1990). Calcium/calmodulin-dependent protein kinase II. *Curr*
7 *Top Cell Regul*, 31, 181-221.
- 8 Collingridge, G. L., Isaac, J. T., & Wang, Y. T. (2004). Receptor trafficking and synaptic
9 plasticity. *Nat Rev Neurosci*, 5(12), 952-962. doi:10.1038/nrn1556
- 10 Coultrap, S. J., Freund, R. K., O'Leary, H., Sanderson, J. L., Roche, K. W., Dell'Acqua, M. L., &
11 Bayer, K. U. (2014). Autonomous CaMKII Mediates Both LTP and LTD Using a
12 Mechanism for Differential Substrate Site Selection. *Cell Reports*, 6(3), 431-437.
13 doi:10.1016/j.celrep.2014.01.005
- 14 Cui, Y., Jin, J., Zhang, X., Xu, H., Yang, L., Du, D., . . . Cao, X. (2011). Forebrain NR2B
15 overexpression facilitating the prefrontal cortex long-term potentiation and enhancing
16 working memory function in mice. *PLoS One*, 6(5), e20312.
17 doi:10.1371/journal.pone.0020312
- 18 Dalton, G. L., Wang, Y. T., Floresco, S. B., & Phillips, A. G. (2008). Disruption of AMPA
19 receptor endocytosis impairs the extinction, but not acquisition of learned fear.
20 *Neuropsychopharmacology : official publication of the American College of*
21 *Neuropsychopharmacology*, 33(10), 2416-2426. doi:10.1038/sj.npp.1301642
- 22 Dalton, G. L., Wu, D. C., Wang, Y. T., Floresco, S. B., & Phillips, A. G. (2012). NMDA
23 GluN2A and GluN2B receptors play separate roles in the induction of LTP and LTD in
24 the amygdala and in the acquisition and extinction of conditioned fear.
25 *Neuropharmacology*, 62(2), 797-806. doi:10.1016/j.neuropharm.2011.09.001
- 26 Dębiec, J., Bush, D. E., & LeDoux, J. E. (2011). Noradrenergic enhancement of reconsolidation
27 in the amygdala impairs extinction of conditioned fear in rats--a possible mechanism for
28 the persistence of traumatic memories in PTSD. *Depress Anxiety*, 28(3), 186-193.
29 doi:10.1002/da.20803
- 30 Delgado, J. Y., Coba, M., Anderson, C. N., Thompson, K. R., Gray, E. E., Heusner, C. L., . . .
31 O'Dell, T. J. (2007). NMDA receptor activation dephosphorylates AMPA receptor

- 1 glutamate receptor 1 subunits at threonine 840. *J Neurosci*, 27(48), 13210-13221.
- 2 doi:10.1523/JNEUROSCI.3056-07.2007
- 3 Delgado, J. Y., Coba, M., Anderson, C. N. G., Thompson, K. R., Gray, E. E., Heusner, C. L., . . .
- 4 O'Dell, T. J. (2007). NMDA Receptor Activation Dephosphorylates AMPA Receptor
- 5 Glutamate Receptor 1 Subunits at Threonine 840. *The Journal of Neuroscience*, 27,
- 6 13210-13221.
- 7 Diering, G. H., Heo, S., Hussain, N. K., Liu, B., & Huganir, R. L. (2016). Extensive
- 8 phosphorylation of AMPA receptors in neurons. *Proceedings of the National Academy of*
- 9 *Sciences*, 113(33), E4920. doi:10.1073/pnas.1610631113
- 10 Dohadwala, M., da Cruz e Silva, E. F., Hall, F. L., Williams, R. T., Carbonaro-Hall, D. A., Nairn,
- 11 A. C., . . . Berndt, N. (1994). Phosphorylation and inactivation of protein phosphatase 1
- 12 by cyclin-dependent kinases. *Proc Natl Acad Sci U S A*, 91(14), 6408-6412.
- 13 Duan, Y., Zhou, S., Ma, J., Yin, P., & Cao, X. (2015). Forebrain NR2B overexpression
- 14 enhancing fear acquisition and long-term potentiation in the lateral amygdala. *Eur J*
- 15 *Neurosci*, 42(5), 2214-2223. doi:10.1111/ejn.13008
- 16 Erlich, J. C., Bush, D. E. A., & Ledoux, J. E. (2012). The role of the lateral amygdala in the
- 17 retrieval and maintenance of fear-memories formed by repeated probabilistic
- 18 reinforcement. *Frontiers in behavioral neuroscience*, 6, 16-16.
- 19 doi:10.3389/fnbeh.2012.00016
- 20 Fenster, R. J., Lebois, L. A. M., Ressler, K. J., & Suh, J. (2018). Brain circuit dysfunction in
- 21 post-traumatic stress disorder: from mouse to man. *Nat Rev Neurosci*, 19(9), 535-551.
- 22 doi:10.1038/s41583-018-0039-7
- 23 Forster, G. L., Novick, A. M., Scholl, J. L., & Watt, M. J. (2012). The Role of the Amygdala in
- 24 Anxiety Disorders *The Amygdala - A Discrete Multitasking Manager*.
- 25 Fox, C. J., Russell, K., Titterness, A. K., Wang, Y. T., & Christie, B. R. (2007). Tyrosine
- 26 phosphorylation of the GluR2 subunit is required for long-term depression of synaptic
- 27 efficacy in young animals in vivo. *Hippocampus*, 17(8), 600-605.
- 28 doi:10.1002/hipo.20302
- 29 Fukunaga, K., Muller, D., Ohmitsu, M., Bak ́ E., DePaoli-Roach, A. A., & Miyamoto, E. (2000).
- 30 Decreased Protein Phosphatase 2A Activity in Hippocampal Long-Term Potentiation.
- 31 *Journal of Neurochemistry*, 74(2), 807-817. doi:10.1046/j.1471-4159.2000.740807.x

- 1 Goldberg, J., Huang, H. B., Kwon, Y. G., Greengard, P., Nairn, A. C., & Kuriyan, J. (1995).
2 Three-dimensional structure of the catalytic subunit of protein serine/threonine
3 phosphatase-1. *Nature*, 376(6543), 745-753. doi:10.1038/376745a0
- 4 Grosso, A., Santoni, G., Manassero, E., Renna, A., & Sacchetti, B. (2018). A neuronal basis for
5 fear discrimination in the lateral amygdala. *Nat Commun*, 9(1), 1214.
6 doi:10.1038/s41467-018-03682-2
- 7 He, K., Lee, A., Song, L., Kanold, P. O., & Lee, H. K. (2011). AMPA receptor subunit GluR1
8 (GluA1) serine-845 site is involved in synaptic depression but not in spine shrinkage
9 associated with chemical long-term depression. *J Neurophysiol*, 105(4), 1897-1907.
10 doi:10.1152/jn.00913.2010
- 11 Hell, J. W. (2014). CaMKII: claiming center stage in postsynaptic function and organization.
12 *Neuron*, 81(2), 249-265. doi:10.1016/j.neuron.2013.12.024
- 13 Hollis, F., Sevelinges, Y., Grosse, J., Zanoletti, O., & Sandi, C. (2016). Involvement of CRFR1
14 in the Basolateral Amygdala in the Immediate Fear Extinction Deficit. *Eneuro*, 3(5).
15 doi:10.1523/Eneuro.0084-16.2016
- 16 Hu, X.-d., Huang, Q., Yang, X., & Xia, H. (2007). Differential Regulation of AMPA Receptor
17 Trafficking by Neurabin-Targeted Synaptic Protein Phosphatase-1 in Synaptic
18 Transmission and Long-Term Depression in Hippocampus. *The Journal of Neuroscience*,
19 27(17), 4674. doi:10.1523/JNEUROSCI.5365-06.2007
- 20 Huang, W. H., Chao, H. W., Tsai, L. Y., Chung, M. H., & Huang, Y. S. (2014). Elevated
21 activation of CaMKII α in the CPEB3-knockout hippocampus impairs a specific form of
22 NMDAR-dependent synaptic depotentiation. *Front Cell Neurosci*, 8, 367.
23 doi:10.3389/fncel.2014.00367
- 24 Jacques, A., Chaaya, N., Hettiarachchi, C., Carmody, M.-L., Beecher, K., Belmer, A., . . .
25 Johnson, L. R. (2019). Microtopography of fear memory consolidation and extinction
26 retrieval within prefrontal cortex and amygdala. *Psychopharmacology*, 236(1), 383-397.
27 doi:10.1007/s00213-018-5068-4
- 28 Ji, M. H., Jia, M., Zhang, M. Q., Liu, W. X., Xie, Z. C., Wang, Z. Y., & Yang, J. J. (2014a).
29 Dexmedetomidine alleviates anxiety-like behaviors and cognitive impairments in a rat
30 model of post-traumatic stress disorder. *Progress in Neuro-Psychopharmacology &*
31 *Biological Psychiatry*, 54, 284-288. doi:10.1016/j.pnpbp.2014.06.013

- 1 Ji, M. H., Jia, M., Zhang, M. Q., Liu, W. X., Xie, Z. C., Wang, Z. Y., & Yang, J. J. (2014b).
2 Dexmedetomidine alleviates anxiety-like behaviors and cognitive impairments in a rat
3 model of post-traumatic stress disorder. *Prog Neuropsychopharmacol Biol Psychiatry*, 54,
4 284-288. doi:10.1016/j.pnpbp.2014.06.013
- 5 Kameyama, K., Lee, H. K., Bear, M. F., & Huganir, R. L. (1998). Involvement of a postsynaptic
6 protein kinase A substrate in the expression of homosynaptic long-term depression.
7 *Neuron*, 21(5), 1163-1175.
- 8 Kennedy, M. B., McGuinness, T., & Greengard, P. (1983). A calcium/calmodulin-dependent
9 protein kinase from mammalian brain that phosphorylates Synapsin I: partial purification
10 and characterization. *J Neurosci*, 3(4), 818-831.
- 11 Kerchner, G. A., & Nicoll, R. A. (2008). Silent synapses and the emergence of a postsynaptic
12 mechanism for LTP. *Nat Rev Neurosci*, 9(11), 813-825. doi:10.1038/nrn2501
- 13 Kim, J., An, B., Kim, J., Park, S., Park, S., Hong, I., . . . Choi, S. (2015). mGluR2/3 in the
14 Lateral Amygdala is Required for Fear Extinction: Cortical Input Synapses onto the
15 Lateral Amygdala as a Target Site of the mGluR2/3 Action. *Neuropsychopharmacology :*
16 *official publication of the American College of Neuropsychopharmacology*, 40(13), 2916-
17 2928. doi:10.1038/npp.2015.145
- 18 Kim, J., Lee, S., Park, K., Hong, I., Song, B., Son, G., . . . Choi, S. (2007). Amygdala
19 depotentiation and fear extinction. *Proc Natl Acad Sci U S A*, 104(52), 20955-20960.
20 doi:10.1073/pnas.0710548105
- 21 Kimura, R., Silva, A. J., & Ohno, M. (2008). Autophosphorylation of alphaCaMKII is
22 differentially involved in new learning and unlearning mechanisms of memory extinction.
23 *Learn Mem*, 15(11), 837-843. doi:10.1101/lm.1049608
- 24 Kollen, M., Dutar, P., & Jouvenceau, A. (2008). The magnitude of hippocampal long term
25 depression depends on the synaptic location of activated NR2-containing N-methyl-D-
26 aspartate receptors. *Neuroscience*, 154(4), 1308-1317.
27 doi:10.1016/j.neuroscience.2008.04.045
- 28 Krabbe, S., Gründemann, J., & Lüthi, A. (2018). Amygdala Inhibitory Circuits Regulate
29 Associative Fear Conditioning. *Biological Psychiatry*, 83(10), 800-809.
30 doi:<https://doi.org/10.1016/j.biopsych.2017.10.006>

- 1 Lee, H.-K., Kameyama, K., Huganir, R. L., & Bear, M. F. (1998). NMDA Induces Long-Term
2 Synaptic Depression and Dephosphorylation of the GluR1 Subunit of AMPA Receptors
3 in Hippocampus. *Neuron*, 21(5), 1151-1162. doi:[https://doi.org/10.1016/S0896-](https://doi.org/10.1016/S0896-6273(00)80632-7)
4 [6273\(00\)80632-7](https://doi.org/10.1016/S0896-6273(00)80632-7)
- 5 Lee, H. K., Barbarosie, M., Kameyama, K., Bear, M. F., & Huganir, R. L. (2000). Regulation of
6 distinct AMPA receptor phosphorylation sites during bidirectional synaptic plasticity.
7 *Nature*, 405(6789), 955-959. doi:10.1038/35016089
- 8 Lee, H. K., Kameyama, K., Huganir, R. L., & Bear, M. F. (1998). NMDA induces long-term
9 synaptic depression and dephosphorylation of the GluR1 subunit of AMPA receptors in
10 hippocampus. *Neuron*, 21(5), 1151-1162.
- 11 Lee, S., Song, B., Kim, J., Park, K., Hong, I., An, B., . . . Choi, S. (2013). GluA1
12 phosphorylation at serine 831 in the lateral amygdala is required for fear renewal. *Nature*
13 *Neuroscience*, 16, 1436. doi:10.1038/nn.3491
14 <https://www.nature.com/articles/nn.3491#supplementary-information>
- 15 Lin, H. C., Mao, S. C., Su, C. L., & Gean, P. W. (2010). Alterations of excitatory transmission in
16 the lateral amygdala during expression and extinction of fear memory. *Int J*
17 *Neuropsychopharmacol*, 13(3), 335-345. doi:10.1017/S1461145709990678
- 18 Lisman, J., Yasuda, R., & Raghavachari, S. (2012). Mechanisms of CaMKII action in long-term
19 potentiation. *Nature reviews. Neuroscience*, 13(3), 169-182. doi:10.1038/nrn3192
- 20 Lisman, J. E., & Zhabotinsky, A. M. (2001). A model of synaptic memory: A CaMKII/PP1
21 switch that potentiates transmission by organizing an AMPA receptor anchoring
22 assembly. *Neuron*, 31(2), 191-201. doi:Doi 10.1016/S0896-6273(01)00364-6
- 23 Lu, W., Isozaki, K., Roche, K. W., & Nicoll, R. A. (2010). Synaptic targeting of AMPA
24 receptors is regulated by a CaMKII site in the first intracellular loop of GluA1.
25 *Proceedings of the National Academy of Sciences of the United States of America*,
26 107(51), 22266-22271. doi:10.1073/pnas.1016289107
- 27 Ma, J., Duan, Y., Qin, Z., Wang, J., Liu, W., Xu, M., . . . Cao, X. (2015). Overexpression of
28 α CaMKII impairs behavioral flexibility and NMDAR-dependent long-term depression in
29 the medial prefrontal cortex. *Neuroscience*, 310, 528-540.
30 doi:<https://doi.org/10.1016/j.neuroscience.2015.09.051>

- 1 Ma, T. F., Zhou, L., Wang, Y., Qin, S. J., Zhang, Y., Hu, B., . . . Gu, S. L. (2013). A selective
2 M1 and M3 receptor antagonist, penahyclidine hydrochloride, prevents postischemic LTP:
3 involvement of NMDA receptors. *Synapse*, 67(12), 865-874. doi:10.1002/syn.21693
- 4 Mahan, A. L., & Ressler, K. J. (2012). Fear conditioning, synaptic plasticity and the amygdala:
5 implications for posttraumatic stress disorder. *Trends in Neurosciences*, 35(1), 24-35.
6 doi:10.1016/j.tins.2011.06.007
- 7 Mansuy, I. M., & Shenolikar, S. (2006). Protein serine/threonine phosphatases in neuronal
8 plasticity and disorders of learning and memory. *Trends Neurosci*, 29(12), 679-686.
9 doi:10.1016/j.tins.2006.10.004
- 10 Mansuy, I. M., & Shenolikar, S. (2006). Protein serine/threonine phosphatases in neuronal
11 plasticity and disorders of learning and memory. *Trends in Neurosciences*, 29(12), 679-
12 686. doi:<https://doi.org/10.1016/j.tins.2006.10.004>
- 13 Matsuda, S., Kakegawa, W., Budisantoso, T., Nomura, T., Kohda, K., & Yuzaki, M. (2013).
14 Stargazin regulates AMPA receptor trafficking through adaptor protein complexes during
15 long-term depression. *Nature Communications*, 4, 2759. doi:10.1038/ncomms3759
16 <https://www.nature.com/articles/ncomms3759#supplementary-information>
- 17 Mauna, J. C., Miyamae, T., Pulli, B., & Thiels, E. (2011). Protein phosphatases 1 and 2A are
18 both required for long-term depression and associated dephosphorylation of cAMP
19 response element binding protein in hippocampal area CA1 in vivo. *Hippocampus*,
20 21(10), 1093-1104. doi:10.1002/hipo.20823
- 21 Michopoulos, V., Rothbaum, A. O., Jovanovic, T., Alml, L. M., Bradley, B., Rothbaum, B.
22 O., . . . Ressler, K. J. (2014). Association of CRP Genetic Variation and CRP Level With
23 Elevated PTSD Symptoms and Physiological Responses in a Civilian Population With
24 High Levels of Trauma. *American Journal of Psychiatry*, 172(4), 353-362.
25 doi:10.1176/appi.ajp.2014.14020263
- 26 Milad, M. R., & Quirk, G. J. (2011). Fear Extinction as a Model for Translational Neuroscience:
27 Ten Years of Progress. *Annual Review of Psychology*, 63(1), 129-151.
28 doi:10.1146/annurev.psych.121208.131631
- 29 Milad, M. R., & Quirk, G. J. (2012). Fear Extinction as a Model for Translational Neuroscience:
30 Ten Years of Progress. *Annual Review of Psychology*, Vol 63, 63, 129-151.
31 doi:10.1146/annurev.psych.121208.131631

- 1 Ming, C., Shao, D., Fu, Y., Ma, Q., Cui, D., Song, J., . . . Zheng, P. (2018). Key determinants for
2 morphine withdrawal conditioned context-induced increase in Arc expression in anterior
3 cingulate cortex and withdrawal memory retrieval. *Experimental Neurology*, 311.
4 doi:10.1016/j.expneurol.2018.10.009
- 5 Morikawa, M., Tanaka, Y., Cho, H. S., Yoshihara, M., & Hirokawa, N. (2018). The Molecular
6 Motor KIF21B Mediates Synaptic Plasticity and Fear Extinction by Terminating Rac1
7 Activation. *Cell Rep*, 23(13), 3864-3877. doi:10.1016/j.celrep.2018.05.089
- 8 Nabavi, S., Fox, R., Proulx, C. D., Lin, J. Y., Tsien, R. Y., & Malinow, R. (2014). Engineering a
9 memory with LTD and LTP. *Nature*, 511(7509), 348-352. doi:10.1038/nature13294
- 10 Okuno, H., Akashi, K., Ishii, Y., Yagishita-Kyo, N., Suzuki, K., Nonaka, M., . . . Bito, H. (2012).
11 Inverse Synaptic Tagging of Inactive Synapses via Dynamic Interaction of Arc/Arg3.1
12 with CaMKII β . *Cell*, 149(4), 886-898. doi:10.1016/j.cell.2012.02.062
- 13 Opazo, P., Labrecque, S., Tigaret, C. M., Frouin, A., Wiseman, P. W., De Koninck, P., &
14 Choquet, D. (2010). CaMKII Triggers the Diffusional Trapping of Surface AMPARs
15 through Phosphorylation of Stargazin. *Neuron*, 67(2), 239-252.
16 doi:<https://doi.org/10.1016/j.neuron.2010.06.007>
- 17 Pi, H. J., & Lisman, J. E. (2008). Coupled Phosphatase and Kinase Switches Produce the
18 Tristability Required for Long-Term Potentiation and Long-Term Depression. *The*
19 *Journal of Neuroscience*, 28(49), 13132. doi:10.1523/JNEUROSCI.2348-08.2008
- 20 Pougnet, J.-T., Compans, B., Martinez, A., Choquet, D., Hosy, E., & Boué-Grabot, E. (2016).
21 P2X-mediated AMPA receptor internalization and synaptic depression is controlled by
22 two CaMKII phosphorylation sites on GluA1 in hippocampal neurons. *Scientific reports*,
23 6, 31836-31836. doi:10.1038/srep31836
- 24 Radulovic, J., Ren, L. Y., & Gao, C. (2019). N-Methyl D-aspartate receptor subunit signaling in
25 fear extinction. *Psychopharmacology (Berl)*, 236(1), 239-250. doi:10.1007/s00213-018-
26 5022-5
- 27 Ressler, K. J. (2010). Amygdala activity, fear, and anxiety: modulation by stress. *Biological*
28 *Psychiatry*, 67(12), 1117-1119. doi:10.1016/j.biopsych.2010.04.027
- 29 Rich, M. T., Abbott, T. B., Chung, L., Gulcicek, E. E., Stone, K. L., Colangelo, C. M., . . .
30 Torregrossa, M. M. (2016). Phosphoproteomic Analysis Reveals a Novel Mechanism of

CaMKII α Regulation Inversely Induced by Cocaine Memory Extinction versus Reconsolidation. *J Neurosci*, 36(29), 7613-7627. doi:10.1523/jneurosci.1108-16.2016

Ritov, G., Boltyansky, B., & Richter-Levin, G. (2016). A novel approach to PTSD modeling in rats reveals alternating patterns of limbic activity in different types of stress reaction. *Molecular Psychiatry*, 21(5), 630-641. doi:10.1038/mp.2015.169

Ritov, G., & Richter-Levin, G. (2014). Water associated zero maze: a novel rat test for long term traumatic re-experiencing. *Frontiers in behavioral neuroscience*, 8, 1-1. doi:10.3389/fnbeh.2014.00001

Schafe, G. E., Doyère, V., & LeDoux, J. E. (2005). Tracking the Fear Engram: The Lateral Amygdala Is an Essential Locus of Fear Memory Storage. *The Journal of Neuroscience*, 25(43), 10010. doi:10.1523/JNEUROSCI.3307-05.2005

Szapiro, G., Vianna, M. R., McGaugh, J. L., Medina, J. H., & Izquierdo, I. (2003). The role of NMDA glutamate receptors, PKA, MAPK, and CAMKII in the hippocampus in extinction of conditioned fear. *Hippocampus*, 13(1), 53-58. doi:10.1002/hipo.10043

Talukdar, G., Inoue, R., Yoshida, T., & Mori, H. (2018). Impairment in extinction of cued fear memory in syntenin-1 knockout mice. *Neurobiol Learn Mem*, 149, 58-67. doi:10.1016/j.nlm.2018.01.006

Tomita, S., Chen, L., Kawasaki, Y., Petralia, R. S., Wenthold, R. J., Nicoll, R. A., & Bredt, D. S. (2003). Functional studies and distribution define a family of transmembrane AMPA receptor regulatory proteins. *The Journal of cell biology*, 161(4), 805-816. doi:10.1083/jcb.200212116

Wang, H., Shimizu, E., Tang, Y. P., Cho, M., Kyin, M., Zuo, W., . . . Tsien, J. Z. (2003). Inducible protein knockout reveals temporal requirement of CaMKII reactivation for memory consolidation in the brain. *Proc Natl Acad Sci U S A*, 100(7), 4287-4292. doi:10.1073/pnas.0636870100

Wei, F., Wang, G. D., Zhang, C., Shokat, K. M., Wang, H., Tsien, J. Z., . . . Zhuo, M. (2006). Forebrain overexpression of CaMKII abolishes cingulate long term depression and reduces mechanical allodynia and thermal hyperalgesia. *Mol Pain*, 2, 21. doi:10.1186/1744-8069-2-21

Winder, D. G., & Sweatt, J. D. (2001). Roles of serine/threonine phosphatases in hippocampal synaptic plasticity. *Nature Reviews Neuroscience*, 2, 461. doi:10.1038/35081514

- 1 Wu, X., Zhang, J.-T., Li, D., Zhou, J., Yang, J., Zheng, H.-L., . . . Wang, F. (2017). Aquaporin-4
2 deficiency facilitates fear memory extinction in the hippocampus through excessive
3 activation of extrasynaptic GluN2B-containing NMDA receptors. *Neuropharmacology*,
4 *112*, 124-134. doi:<https://doi.org/10.1016/j.neuropharm.2016.06.031>
- 5 Yan, W. J., Tan, Y. C., Xu, J. C., Tang, X. P., Zhang, C., Zhang, P. B., & Ren, Z. Q. (2015).
6 Protective Effects of Silibinin and Its Possible Mechanism of Action in Mice Exposed to
7 Chronic Unpredictable Mild Stress. *Biomolecules & Therapeutics*, *23*(3), 245-250.
8 doi:10.4062/biomolther.2014.138
- 9 Yehuda, R., Hoge, C. W., McFarlane, A. C., Vermetten, E., Lanius, R. A., Nievergelt, C. M., . . .
10 Hyman, S. E. (2015). Post-traumatic stress disorder. *Nature Reviews Disease Primers*, *1*,
11 15057. doi:10.1038/nrdp.2015.57
- 12 Yin, D. M., Chen, Y. J., Lu, Y. S., Bean, J. C., Sathiyamurthy, A., Shen, C., . . . Mei, L. (2013).
13 Reversal of behavioral deficits and synaptic dysfunction in mice overexpressing
14 neuregulin 1. *Neuron*, *78*(4), 644-657. doi:10.1016/j.neuron.2013.03.028

Figure 1

FIGURE LEGENDS:

Figure 1. PTSD susceptible mice with cued fear extinction deficit and anxiety-like behaviors exhibited significant up-regulation of α CaMKII and down-regulation of AMPAR internalization in LA.

(A₁₋₂) Schematic illustration for identifying PTSD susceptible mice following UWT (A₁, PS-UWT) or 4-CS/US pairings (A₂, PS-4CS/US) exposure. (B-E) PTSD susceptible mice exhibited the higher level of freezing responses in the fear extinction (B), and anxiety-like behaviors in OF(C), DL(D), OM(E) tests. PTSD susceptible mice spent significantly more time freezing during extinction (B, two-way ANOVA followed by multiple comparisons with Bonferroni's correction), less time in center area of OF chamber, in the light box of DL test and in the open arms of OM tests (C-E, one-way ANOVA followed by multiple comparisons with Bonferroni's correction) compared to control mice (control, n = 12; PS-UWT, n = 7; PS-4CS/US, n = 7). (F₁) Representative blottings of LA synaptosomal region illustrating significant higher expression in α CaMKII, p- α CaMKII-Thr286, GluA1/2, GluA1-Ser831 /Ser845 phosphorylation in PTSD susceptible mice following stress exposure, but no significant change in β CaMKII expression. (F₂) Ponceau S staining was used as a loading control. (G) Quantifications were based on the average of independent experiment (n = 5 per group). Western blotting in "Control", "PS-UWT" or "PS-4CS/US" groups was performed after fear extinction and all the anxiety-like behavior tests. One-way ANOVA followed by multiple comparisons with Bonferroni's correction. n.s.: not significant, * P < 0.05, ** P < 0.01, *** P < 0.001. Error bars represent s.e.m.

Figure 2

Figure 2. α CaMKII-F89G TG mice exhibited PTSD-like behaviors and impairments in AMPAR internalization.

(A) The schematic of behavioral procedure for cued fear conditioning and extinction trials.

(B) Impaired cued fear extinction in TG mice (two-way ANOVA followed by multiple comparisons with Bonferroni's correction). Intraperitoneal (i.p.) injection and oral (p.o.) administration with NM-PP1 could rescue the cued extinction deficits of TG mice (two-way ANOVA followed by multiple comparisons with Bonferroni's correction). (C-E) The higher level of anxiety-like behaviors in TG mice in the OF(C), DL(D) and EPM(E) tests after cued fear conditioning and extinction (one-way ANOVA followed by multiple comparisons with Bonferroni's correction). (F) Up: Representative blottings of LA synaptosomal fractions illustrating an increase in GluA1/2, phosphorylation level of GluA1-Ser845/Ser831 in both WT and TG mice after cued fear conditioning. Down: Ponceau S staining was used as a loading control. A decrease in GluA1/2, phosphorylation level of GluA1-Ser845/Ser831 in WT mice, but not in TG mice after cued fear extinction (n = 5 per group). (G) Quantifications were based on the average of independent experiment. Western blotting in "WT/TG + FC" or "WT/TG + Ext" groups was performed after fear conditioning or fear extinction following with anxiety-like behavior tests, respectively (one-way ANOVA followed by Bonferroni's multiple comparisons test). n.s.: not significant, * P < 0.05, ** P < 0.01 and *** P < 0.001 versus WT group; \$ P < 0.05, \$\$ P < 0.01 and \$\$\$ P < 0.001 versus WT + FC group; # P < 0.05, ## P < 0.01 and ### P < 0.001 versus TG group; + P < 0.05 and ++ P < 0.01. Error bars represent s.e.m.

Figure 3

Figure 3. Increasing α CaMKII specifically in LA impaired the cued fear extinction and AMPAR internalization in AAV- α CaMKII mice.

(A) Images of coronal brain slice showing the expression of eGFP (green-colored) 6 weeks after bilateral injections of pAAV-TRE- α CaMKII-P2A-EGFP-CMV-rTA virus into LA. Numbers indicate coordinates relative to bregma. Scale bar, 500 μ m. (B) The schematic of behavioral procedure for cued fear extinction trials. (C) Elevating α CaMKII in LA could impair cued fear memory and fear extinction (two-way ANOVA followed by multiple comparisons with Bonferroni's correction). (D₁) Representative blottings of LA synaptosomal fractions illustrating an increased expression of α CaMKII, p- α CaMKII-Thr286, GluA1/2, phosphorylated GluA1-Ser845/Ser831 in LA of AAV- α CaMKII mice than that in AAV-control mice. (D₂) Ponceau S staining was used as a loading control. (E) Quantifications were based on the average of independent experiment (n = 6 per group). Western blotting was performed after fear extinction and all the anxiety-like behavior tests. Statistical differences were evaluated with Student's t test. * P < 0.05, ** P < 0.01, *** P < 0.001 Error bars represent s.e.m.

Figure 4

Figure 4. Increasing α CaMKII impairs NMDAR-dependent LTD at T-LA synapses of α CaMKII-F89G TG mice and NM-PP1 can rescue the impairments.

(A) Similar LTP induced by high frequency stimulations (2 trains of 100 Hz stimulation for 1 s, 10 s interval) in TG slices and WT slices. In this and the subsequent figures, insets show sample traces taken at baseline (1) and the last 10 min recording (2). (B) Normal LTP induced by three trains of theta burst stimulations (TBS, each train consisted of 10 bursts delivered at 5 Hz, each burst consisted of 4 pulses at 100 Hz) in TG slices. (C) Significantly weaker LTD induced in TG slices than that in WT slice after 1 Hz (15 min) stimulation. NM-PP1 (0.5 μ M) recovered the reduced LTD in TG slice to normal level. (D) LTD was abolished in WT and TG slices exposed to both NM-PP1 (0.5 μ M) and APV (50 μ M). The solid line shows the duration of both NM-PP1 and APV application. (E) Strong LTD could be induced by 3 Hz (5 min) stimulation in WT slices but not in TG slices, NM-PP1 (0.5 μ M) rescued the impaired LTD in TG slice. (F) Impaired depotentiation can be observed in TG slices. All of the bar graph summarizing data obtained during last 10 min recording. Statistical differences were evaluated with Student's t test (A, B, D and F) and one-way ANOVA followed by Bonferroni's multiple comparisons test (C and E). n.s.: not significant, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. All values are mean $\pm$ s.e.m.

Figure 5. Increasing α CaMKII impairs AMPAR internalization / dephosphorylation, reduces protein phosphatase (PP) activity, and increases stargazin expression during NMDAR-dependent LTD and NM-PP1 can rescue all impairments.

(A) Attenuated chem-LTD induced by 30 μ M NMDA for 3 min in TG slices. This deficit could be rescued by 0.5 μ M NM-PP1. Right-up panel: bar graph summarizing data obtained during last 10 min recording in the different groups depicted. The following Western blotting was performed 1 hour later after NMDA application. (B) Representative blottings of LA synaptosomal fractions illustrating a reduction in GluA1/2, phosphorylation level of GluA1-Ser845/831 in WT slices after NMDA treatment but not in TG slices. NM-PP1 could rescue these deficits in TG slices. Down: Ponceau S staining was used as a loading control. (C) Quantifications were based on the average of independent experiments (n = 5 per group). (D) Up: Representative blottings of LA synaptosomal fractions illustrating a reduction in phosphorylation level of pPP1-Thr320, indicating an increase in PP1 activity in WT mice after NMDA treatment but not in TG. NM-PP1 rescued such deficit in TG mice. Down: Quantifications were based on the average of independent experiments (n = 4 per group). (E) A remarkably higher level of stargazin in amygdala synaptosomal fractions in TG slices than that in WT slices after NMDA application, NM-PP1 rescued the deficit in TG mice (n = 5 per group). Down: Ponceau S staining was used as a loading control. (F) An increased activity of PP2A and PP2B in WT slices were exhibited after NMDA application but not in TG slices, and NM-PP1 rescued these deficits in TG mice (n = 4 per group). Statistical differences were evaluated with one-way ANOVA followed by multiple comparisons with Bonferroni's correction. n.s.: not significant, * P < 0.05, ** P < 0.01 and *** P < 0.001 versus WT group; # P < 0.05, ## P < 0.01 and ### P < 0.001 versus TG group; + P < 0.05, ++ P < 0.01 and +++ P < 0.001. Error bars represent s.e.m.

Supplemental information

Higher level of α CaMKII but normal morphology in LA of α CaMKII-F89G TG mice

First, we examined the α CaMKII expression level in LA of both α CaMKII-F89G TG and WT mice. Western blotting quantification revealed the expression of synaptic α CaMKII protein in LA of TG mice was 136% of WT littermates (Supplementary Fig. S1A, 1B, $P < 0.05$). Strikingly, the p- α CaMKII-Thr286 in LA of TG mice was 195% of WT littermates ($P < 0.001$). However, no obvious change in β CaMKII expression was observed in LA of TG mice (Supplementary Fig. S1A, B). Moreover, Nissl staining showed no detectable morphological abnormalities in LA of TG mice (Supplementary Fig. S1C). Normal shapes and architecture of dendritic spines could also be found in LA of TG mice (Supplementary Fig. S1D, S1E). These results suggest that the transgenic expression of α CaMKII-F89G increase α CaMKII expression in LA.

Normal locomotor activity and acute pain threshold to footshock in α CaMKII-F89G TG Mice.

Then, to investigate whether α CaMKII overexpression influences basal motor, exploratory behaviors and the foot shock sensitivity, we performed open field and pain threshold tests. No significant difference was observed between TG and WT mice in both locomotor activity (Fig. S2A, $P > 0.05$; Student's t-test) and rearing behavior (Fig. S2A, $P > 0.05$), showing that TG mice exhibit normal locomotor activity and exploratory behavior. Moreover, we quantified the minimum current intensity of foot shock required to induce flinching, vocalizing and jumping in two groups of mice. There was also no significant difference in the threshold of current intensity to trigger flinching, vocalizing and jumping behaviors in TG mice and WT littermates (Fig. S2 B, $P > 0.05$). Taken all together, we can conclude that α CaMKII overexpression indeed impairs cued fear extinction.

A **B** **C** **D**

Fig S1. Higher level of α CaMKII but normal morphology in LA of α CaMKII-F89G TG mice.

(A) Immunoblottings of α CaMKII protein in LA from WT and TG mice (p- α CaMKII-Thr286: p < 0.001; α CaMKII: p < 0.05; β CaMKII: p > 0.05). (B) Densitometric analysis shows a significantly higher expression of α CaMKII and p- α CaMKII-Thr286 in TG than that in WT mice. (C) Parts of Nissl stained coronal slices showing the amygdala of both WT and TG mice. Note no detectable morphological differences between WT and TG mice in the amygdala. Scale bars, 100 μ m. (D) Dendritic spine of LA pyramidal neurons in WT and TG mice. Scale bar, 5 μ m. The spine density (spines / 20 μ m) was comparable between WT and TG mice (p > 0.05). Statistical differences were evaluated with Student's t test, * P < 0.05, ** P < 0.01, *** P < 0.001. All data are shown as mean $\pm$ s.e.m.

Fig S2

Fig S2. Normal locomotor activity and acute pain threshold to footshock in α CaMKII-F89G TG Mice.

(A) Similar moving distance ($P > 0.05$) and rearing behavior ($P > 0.05$) in TG and WT mice during a 15 min of the open field test. **(B)** Normal pain sensitivity to an increasing electric footshock in TG mice ($P > 0.05$). All values are mean $\pm$ s.e.m. Statistical differences were evaluated with Student's t-test.

Fig S3

Fig S3. Normal basal synaptic transmission at T-LA synapses in α CaMKII-F89G TG mice.

(A) No significant difference in the input/output curve at T-LA synapses between WT and TG slices (two-way ANOVA followed by multiple comparisons with Bonferroni's correction). **(B)** Comparable synaptic or total GluA1/2 expression in LA of WT and TG slices (Statistical differences were evaluated with Student's t-test). **(C)** Similar paired-pulse depression at different interpulse intervals in WT and TG amygdala slices (two-way ANOVA followed by multiple comparisons with Bonferroni's correction). **(D)** Comparable expression levels of synapsin in LA of WT and TG amygdala slices (Statistical differences were evaluated with Student's t-test). All data are shown as mean $\pm$ s.e.m.