1 **Title:**

2 **Metabarcoding, direct stomach observation and stable isotope analysis reveal a highly**
 3 **diverse diet for the invasive green crab in Atlantic Patagonia**

4 **Authors--**

5 Georgina Cordone¹, Mariana Lozada², Elisabet Vilacoba³, Bettina Thalinger⁴, Gregorio Bigatti^{5,6},
 6 Darío A. Lijtmaer³, Dirk Steinke⁴, David E. Galván^{1*}

7 * Corresponding author: galvan@cenpat-conicet.gob.ar ; Tel: +54 (280) 488-3184 Ext. 1277 - Fax:
 8 +54 (280) 488-3543

9 ¹ Centro para el Estudio de Sistemas Marinos (CESIMAR), Consejo Nacional de Investigaciones
 10 Científicas y Técnicas (CONICET), Edificio CCT CONICET-CENPAT, Bvd. Brown 2915, (U9120)
 11 Puerto Madryn, Chubut, Argentina.

12 ² Laboratorio de Microbiología Ambiental (IBIOMAR), Consejo Nacional de Investigaciones
 13 Científicas y Técnicas (CONICET), Edificio CCT CONICET-CENPAT, Bvd. Brown 2915, (U9120)
 14 Puerto Madryn, Chubut, Argentina.

15 ³ División Ornitología, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia” MACN-
 16 CONICET. Av. Ángel Gallardo 470 (C1405) Ciudad Autónoma de Buenos Aires, Buenos Aires,
 17 Argentina.

18 ⁴ Centre for Biodiversity Genomics, University of Guelph, 50 Stone Rd E (N1G2W1), Guelph,
 19 Ontario, Canada.

20 ⁵ Laboratorio de Reproducción y Biología Integrativa de Invertebrados Marinos, (LARBIM,
 21 IBIOMAR). Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Edificio CCT
 22 CONICET-CENPAT, Bvd. Brown 2915, (U9120) Puerto Madryn, Chubut, Argentina.

23 ⁶ Universidad Espíritu Santo, Ecuador

DEG, GB, ML, and GC conceived and designed this study. DEG, GB, and GC took the samples. GC conducted the visual analysis and prepare the samples for stable isotope analysis. GC and DEG analyzed direct stomach observation and stable isotope data. ML contributed to preparing the metabarcoding samples and analyzed metabarcoding data. EV visited as an invited researcher the Centre for Biodiversity Genomics to obtain the metabarcoding data. EV, BT, DAL, and DS contributed to the generation of metabarcoding data, molecular and bioinformatic analyses. All authors contributed to manuscript writing.

24 Abstract

25 The European green crab *Carcinus maenas* and, its sister species, *C. aestuarii* are highly
 26 invasive species causing damages to coastal ecosystems and severe economic losses worldwide. *C.*
 27 *maenas* was detected at the Atlantic Patagonian coast twenty years ago. In this work, we studied the
 28 diet of the green crab in a recently invaded location in Nuevo Gulf, using three complementary
 29 techniques: direct stomach observation, stable isotope analysis, and metabarcoding of the gut
 30 content. Direct stomach observation and metabarcoding showed that green crabs have a broad
 31 omnivorous diet, ingesting most of the phyla present in the study area. Gut content metabarcoding
 32 allowed a detailed description of algae diversity and revealed other taxa that went unnoticed in the
 33 visual stomach analysis. Stable isotope analysis showed that the major contribution to crabs' diet
 34 was from the phytoplankton chain (by bivalve consumption) and not directly from algae. This study
 35 approach combining three complementary techniques allowed us to detect some differences in the
 36 diet between sexes, which suggests that male and female crabs are not as ecologically equivalent as
 37 previously thought. Besides, we detected sequences corresponding to *C. aestuarii* suggesting that
 38 the green crab Patagonian population is a hybrid of both sister species. Finally, we highlight
 39 possible direct and indirect interactions of the green crab with the native species that can trigger
 40 negative effects throughout the entire food web.

41 **Keywords:** *Carcinus maenas*, *Carcinus aestuarii*, e-DNA, Rocky shore ecology, Trophic
 42 interactions

43 1. Introduction

44 Invasive species, particularly invasive predators are one of the main causes of worldwide
 45 species decline and biodiversity loss (Clavero and García-Berthou 2005, Doherty et al. 2016). Once
 46 invasive species arrive at a new location, they establish new ecological interactions with resident
 47 species and these new interactions can have both positive or negative effects on native species or
 48 the entire native food web (David et al. 2017). Food webs represent energy and matter flows among

organisms related to each other by consumer-resource relationships; their network structure influences community dynamics and stability (Bascompte 2009, Thébault and Fontaine 2010). The ways biological invasions impact food webs are complex and context-dependent as there are various direct and indirect mechanisms by which an invasive species can affect other taxa (Thomsen et al. 2011a, Thomsen et al. 2011b). Preying upon a native species is a direct mechanism that usually ends up with population decrease, or even species loss (Bruno et al. 2005, David et al. 2017). Exploitative and apparent competition are indirect mechanisms that can also cause the decline of native populations (White et al. 2006, David et al. 2017). On the other hand invaders can display positive effects for resident species. In fact, for species at higher trophic positions, invaders can increase food availability and new habitats (Rodriguez 2006, Thomsen et al. 2010, Thomsen et al. 2014). The invaders' impact depends on the food web's invasibility which is ultimately related to network structure (Hui et al. 2016). Earlier work has shown that food webs with higher diversity are more resistant to invasions; however, on the flip side highly diverse communities generally support more invasive species (Fridley et al. 2007, Tilman et al. 2014, David et al., 2017). By using simulations, Romanuk et al. (2017) have shown that trophic cascades are more frequent when invaders become secondary consumers with abundant prey and only few other predators. Consequently, if we want to accurately predict possible impacts of a newly arrived invasive species, we need to understand its trophic interactions at the invaded ecosystem.

Determining the diet of a particular species is a key requirement to understand its trophic ecology, but it is far from being a simple achievement. There are a series of conceptual and methodological aspects that must be taken into account (e.g Traugott et al. 2013). It is not the same to document which organisms an animal ingests as to list which organisms are targets of a predator or a scavenger. Similarly, the ingestion of a certain item does not assure its digestion and assimilation. A variety of direct and indirect methods are currently used to analyze diet composition (Majdi et al. 2018). Methods include visual observation of feeding events and regurgitations; the inspection of faeces, pellets and stomach contents; and the molecular profiling of stomach contents

75 or faeces using DNA sequencing or immunoassays. DNA-based techniques allow the
 76 characterization of feeding interactions at high taxonomic resolution and are very useful when
 77 visual recognition is not possible (Clare 2014, Unger et al. 2020). In cases of omnivorous species,
 78 where a mixture of DNA from different species is present in gut content, the most effective
 79 approach is metabarcoding (Taberlet et al. 2012, Pompanon et al. 2012, Nielsen et al. 2018).
 80 Metabarcoding couples DNA barcoding (Hebert et al. 2003) with high-throughput sequencing of
 81 community samples (i.e. gut samples) (Nielsen et al. 2018). It not only provides information on
 82 ingested organisms, it also has the advantage of providing complementary information such as the
 83 gastrointestinal parasites of a predator. Aside from such direct methods, indirect tracers of diet
 84 composition are robust and commonly used tools. They include the characterization of biomarkers
 85 such as fatty acids, sterols or pigments present in consumer tissues and the determination of the
 86 isotopic compositions of C, N and S in specific tissues (e.g. Galloway et al. 2015, Stock et al.,
 87 2018). Stable isotope analysis (SIA) is based on predictable differences between the isotopic
 88 signature of an organism and its food resources. It allows for the quantitative estimation of fluxes in
 89 food webs, the determination of a species' trophic level and insights into the patterns of resource
 90 allocation (Boecklen et al. 2011). All these approaches are largely complementary as direct methods
 91 can identify the prey which is ingested, while indirect methods provide information on what is
 92 assimilated. In addition, indirect methods such as SIA provide information at longer time scales
 93 while DNA or visual analysis register consumption within short time periods (typically 6-48h)
 94 (Albaina et al. 2010, Hayden et al. 2014). Majdi et al. (2018) highlighted the potential of combining
 95 direct and indirect methodologies and noted the lack of this kind of studies in current trophic
 96 ecology research.

97 However, there are method-specific biases. Direct visual stomach content analysis provides
 98 indispensable first-hand information, allows quantifying the ingested amount, and for certain cases
 99 distinguishes between secondary predation and random ingestion; however, digestion of prey
 100 material can limit its detection strength, favoring certain taxonomic groups such as shell-bearing

101 organisms. Moreover, it largely depends on the skills of taxonomists involved (Majdi et al. 2018,
102 Traugott et al. 2020). Molecular results can contain errors associated with contamination, biases
103 associated with primer coverage, sequencing errors, as well as bioinformatics errors (Braukmann et
104 al. 2019). On the other hand, metabarcoding is typically regarded as a powerful, high-resolution
105 method (Unger et al. 2020) although it sometimes can lack taxonomic resolution due to gaps in
106 reference databases, particularly for poorly described environments. As for indirect methods, a
107 potential confounding factor of SIA is the variability of trophic discrimination factors (TDF) and
108 turnover rates that may depend on environments, diet, taxon and tissue (Pinnegar and Polunin 1999,
109 Philippsen and Benedito 2013, Hussey et al. 2014, Lefebvre and Dubois 2016). Isotope
110 compositions are highly variable through space and time (Hyndes et al. 2013, Mackey et al. 2015).
111 Therefore, these two factors combined with TDF should be carefully considered when stable
112 isotopes are used in diet studies.

113 The European green crab *Carcinus maenas* (Linnaeus, 1758) is an invasive species native to
114 the North Atlantic Coast. To this day it has reached a worldwide distribution (Carlton and Cohen
115 2003). Not only *C. maenas* have invaded new locations, but its sister species *C. aestuarii*, or
116 hybrids of both, have also been detected at invaded sites (Darling 2011). Concerning its feeding
117 ecology, green crab is considered an aggressive invader due to its voracity and predation on other
118 invertebrates (McDonald et al. 2001). It has been shown that portunid crabs are scavengers and
119 predators, and that green crab is an active carnivore feeding especially on mollusca and crustacea
120 (Ebling et al. 1964, Elner 1981, Grosholz and Ruiz, 1995, McDonald et al. 2001, Chen et al. 2004).
121 It was first detected at the Patagonian coast in 2001, and twelve years later it was registered in the
122 Nuevo Gulf, 250 km north of the point of the first observation (Torres and González-Pisani 2016).
123 The coast of Argentina currently bears a total of 129 introduced and 72 cryptogenic marine species,
124 and it has been estimated that every 178 days a new invasion is detected (Schwindt et al. 2020). The
125 interaction of the green crab population with native as well as introduced species could have
126 unpredictable consequences for the local trophic structure, as well as for ecosystem functioning.

127 Very little is known about green crab ecology at the Argentine coasts, in particular with regards to
128 its local trophic interactions (Hidalgo et al. 2007, Young and Elliott 2020). Moreover, the
129 observation that feeding behaviour is more aggressive in invasive than in native populations
130 (Howard et al. 2018), leads to further uncertainties on how this predator could affect the invaded
131 ecosystem.

132 In this study we aim to shed a light on the diet of the green crab in Atlantic Patagonian
133 coastal environments by using three complementary techniques that have proven useful for diet
134 tracing: visual stomach content inspection, stable isotope analysis, and amplicon sequencing of a
135 gene fragment from stomach content DNA (metabarcoding). We address the following questions to
136 understand possible impacts on the trophic structure of the affected coastal communities in this
137 coastal environment: What are green crabs eating in the affected coastal environments of Atlantic
138 Patagonia? Are there differences in diet between female and male green crabs? What changes can
139 green crabs trigger in local food webs? Does the use of a multidisciplinary approach contribute to a
140 more complete picture?

141 **2. Materials and Methods**

142 **2.1 Sample collection**

143 Green crabs were collected at Punta Este beach in the Nuevo Gulf, Patagonia, Argentina
144 (42°46 ' S, 64°57 ' W). Sampling was done from November 2018 to February 2019 in shallow
145 waters with depths of about 2-3 m at high tide. Crabs were captured manually by SCUBA diving,
146 resulting in a total of 223 specimens. Individuals were separated into two groups. one for traditional
147 visual analysis (183) and the other for metabarcoding (40). After sampling, all specimens were
148 transported to the laboratory and immediately frozen at -20 °C. Crab sizes were registered by
149 measuring carapace width (CW) in cm. Subsequently, crabs were sexed and dissected. For
150 specimens destined for visual analysis, stomach content was extracted and kept in 70% ethanol. For
151 specimens destined for metabarcoding, stomach content was manually collected from frozen

152 samples by letting samples thaw and carefully extracting contents with a plastic pipette tip through
153 the mouth of the crab. Contents were placed in 100% ethanol (Merck) and stored at -80°C until
154 DNA extraction. Muscle from crab chela was extracted and frozen for stable isotope analysis. In
155 addition, five individuals of the mussel *Mytilus edulis* and five of the herbivorous snail *Tegula*
156 *patagonica* were collected to establish isotopic baselines. This was done under the assumption that
157 both the mussel and the herbivorous snail are primary consumers feeding mostly on micro- and
158 macroalgae, respectively.

159 2.2 Visual analysis of gut contents

160 The visual analysis of gut contents was done using a stereomicroscope (Zeiss Stemi, model
161 2000c). Prey was identified to the lowest possible taxonomic level following catalogues and keys of
162 the local fauna (Fauchald 1977, Gosztonyi and Kuba 1996, Boschi et al. 1992, Spivak et al. 2019),
163 as well as through expert consultation and by comparisons with fresh material. Presence of prey
164 items in each stomach was recorded and the frequency of occurrence ($FO_{(vis)}\%$) calculated per prey
165 item as the percentage of non-empty stomachs in which the item was recorded. For statistical
166 analysis, we grouped prey using major groups (Annelida, Arthropoda, Mollusca, Chordata,
167 Echinodermata and Algae) and estimated the confidence interval of the FO by a bootstrap procedure
168 (Tirasin and Jorgensen 1999); using the functions implemented in the R package ‘boot’ (Ripley
169 2005).

170 2.3 Metabarcoding analysis of gut content

171 *DNA extraction and COI amplification*

172 All molecular analyses were carried out under clean-room conditions at the Centre for
173 Biodiversity Genomics (University of Guelph, Canada). First, the ethanol was evaporated from the
174 gut content samples in a fume hood over night, then the remaining pellets were lysed and the DNA
175 extracted using the DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany), following
176 manufacturers’ instructions except for an additional centrifugation step (14000 rpm) for removing

the buffer AW2 from the silica membrane and for elution in two steps using 100 µl of Buffer AE per each. A fume hood control and extraction control were processed along the stomach content samples to check for cross-contamination. The DNA extracts were quantified using Qubit dsDNA HS Assay Kit and a Qubit 3.0 Fluorometer (Thermo Fisher Scientific, MA, USA). For the amplification of gut content DNA, we used a two-step PCR protocol following Elbrecht et al. (2019). PCRs were carried out using the Multiplex PCR Master Mix Plus (Qiagen, Hilden, Germany), 0.5 mM of each primer (IDT, Skokie, Illinois), and molecular grade water (HyPure, GE, Utha, USA) for a total volume of 25 µl. For the first PCR step, amplification of cytochrome oxidase I gene was performed using primers BF3/BR2 (5' CCHGAYATRGCHTTYCCHCG 3' / 5' TCDGGRTGNCCRAARAAYCA 3') and 4 µl of DNA extract. For the second PCR step, 1 µl of the product obtained from the first PCR was used and amplification was performed using fusion primers with individual tags per sample (see Electronic Supplemental Material, Online Resource 1, section 2.3) (Elbrecht and Leese 2017, Elbrecht et al. 2019, Elbrecht and Steinke 2019). Thermocycling conditions for both PCR steps included an initial denaturation at 95°C for 5 minutes, 24 cycles of denaturation at 95°C for 30 seconds, annealing at 50°C for 30 seconds, and extension at 72°C for 50 seconds; and a final extension at 68°C for 10 minutes. After the second PCR, amplification success and band strength were checked on 1.5% agarose gels. For several samples which showed no or only very weak amplification, the two PCRs were repeated, but with 35 cycles instead of 24 in the first PCR. These replicates received individual tag combinations and were prepared for sequencing on an Illumina MiSeq platform along with the extraction controls and seven PCR negative controls:

First, normalization of the second PCR products was performed using the SequalPrep Normalization Plate Kit (Invitrogen, CA, USA), then samples were pooled in one tube and cleaned using SPRIselect following the left side size protocol with a sample-to-volume ratio of 0.76X (Beckman Coulter, CA, USA). Library quantification was performed using Qubit dsDNA HS Assay Kit and a Qubit 3.0 Fluorometer (Thermo Fisher Scientific, MA, USA).

Sequence processing

204 Sequences were preprocessed using JAMP (<https://github.com/VascoElbrecht/JAMP>).
 205 Briefly, sequences were demultiplexed, and forward and reverse paired-end reads were merged.
 206 Primer sequences were removed, and sequences between 380bp and 440 bp (+/- 30bp of expected
 207 fragment length) were kept in the dataset, in order to preserve the most of the possible diversity
 208 present in the sample.

209 The resulting files were processed in two ways: i) the reads were quality filtered and
 210 taxonomy was assigned using the Multiplex Barcode Research And Visualization Environment of
 211 the Barcode of Life platform (mBRAVE/BOLD; Ratnasingham, and Hebert, 2007), and ii) quality
 212 filtering was carried out in JAMP and the resulting haplotypes were grouped into OTUs and blasted
 213 against the NCBI nucleotide database (for details see Supplementary Methods).

214 *Bioinformatic and statistical analysis of molecular data*

215 Taxonomic assignments obtained by the two previous pipelines (mBRAVE/BOLD-based
 216 dataset and blast/NCBI-based dataset) were manually curated. First, genus level taxonomic tables
 217 were screened for locally occurring species and, in the case of the mBRAVE dataset, the quality of
 218 the BOLD references (BINs) was determined to define the most reliable taxonomic level. For
 219 instance, genera which are not described for Argentina, not known from the marine environment,
 220 and/or whose BIN included multiple genera or could be considered unreliable (e.g. composed by
 221 only one sequence) were considered as “unidentified” members of that family/class. The NCBI
 222 dataset was curated at the OTU level, based on % similarity of the first hit to OTU representative
 223 sequences. Only hits with pairwise identity >85% and alignment coverage >75% were kept for
 224 analysis. Of these, representative sequences were assigned to: species (at least 98% similarity with
 225 the first hit), genus (95%), family (90%), and order (85%) (Elbrecht et al. 2017). For analysis of
 226 taxa occurrence, both datasets were combined into one that contained BOLD assignments plus
 227 NCBI assignments of taxa not detected in BOLD. The results were used to calculate the frequency
 228 of occurrence ($FO_{(met)}$) at different taxonomic levels in the same way as $FO_{(vis)}$ (see section 2.2). We
 229 decided to use occurrence data and not quantitative data such as read count because gut content

material is usually at various states of digestion and the sequence counts are less likely to contain a meaningful quantitative signal compared to e.g. faecal matter (Nakahara et al. 2015, Deagle et al. 2019). Although green crab inhabits the intertidal zone, there had been no previous evidence of terrestrial intakes (Cohen et al. 1995). Therefore, terrestrial taxa such as most insects in the metabarcoding data were not taken into account for further analyses. Complete tables of taxonomic assignments are provided in Electronic Supplemental Material (Online Resource 2: Table A1 and Online Resource 3: Table A2).

2.4 Statistical analysis of diet description

To test if sample size was sufficient to obtain a representative description of the diet, we constructed cumulative trophic diversity curves using the quadrats methods following Koen Alonso et al. (2002) for both datasets (i.e. visual and metabarcoding analyses). We calculated trophic diversity as the Brillouin index (H_z) and considered diversity curves as asymptotic if the two previous values (i.e. $n-1$ and $n-2$) were in the range $H_z \pm 0.05 H_z$ for phylum and genus classification (Pielou 1966, Koen Alonso et al. 2002). Once this condition was met the number of stomachs in our study ($n_{(vis)} = 183$ and $n_{(met)} = 40$) was considered sufficient to describe the green crab diet.

To test differences between male and female green crab diet, we performed a distance-based multivariate Welch t-test (Wd*-test) both for the metabarcoding (met) and the visual (vis) analyses. The Wd*-test is a robust alternative to PERMANOVA when heteroscedasticity or imbalanced data are present (Alekseyenko et al. 2016, Hamidi et al. 2019). The tests were performed using a Bray-Curtis similarity matrix implemented in vegan and the MicEco R package (Oksanen et al. 2019, Russel 2020).

Calculation of alpha and beta diversity metrics, ordinations, and other statistical analyses based on OTUs were performed using only the NCBI dataset, using the R packages vegan and phyloseq (McMurdie and Holmes, 2013, Oksanen et al. 2019). Briefly, OTU richness was calculated with the specnumber function in vegan, and a Kruskal-Wallis non-parametric test was

used to analyze whether there were differences in prey richness between sexes. For beta diversity analyses, a similarity matrix based on OTUs was built using the Unifrac similarity index, a phylogenetic-based method which introduces the genetic distance between OTUs to estimate dissimilarity between any two communities (Lozupone and Knight 2005). A phylogenetic tree based on genetic distance was built in QIIME™ from the OTU representative sequences, using the tool phylogeny align-to-tree-mafft-fasttree. This command uses FastTree version 2, which constructs approximately maximum-likelihood trees from large datasets and calculates local support values using the Shimodaira-Hasegawa test (Price et al. 2010). This tree was then used in phyloseq to calculate the index for each pair of samples, which is defined as the fraction of the branch length of the tree leading to OTUs unique to a particular sample over total branch length (Lozupone and Knight 2005). The index was used in its unweighted version, which does not take into account OTU abundances for index calculation. From the Unifrac pairwise distance matrix, NMDS ordinations were constructed. The whole process is wrapped in phyloseq with the command ordinate (physeq, method="NMDS", distance="unifrac"). In addition, the tree with support values based on 1,000 resamples was used to visualize the shared OTUs between males and females.

All analyses were carried out in R (R Core Team, 2020).

2.5 Stable isotope analysis

Muscle samples of both green crabs and the two baseline species (*M. edulis* and *T. patagonica*) were dried for 48 h at 60 °C, ground into a fine powder and ≈ 1.25 μg per sample was sent to the Stable Isotopes Center at the University of New Mexico for C and N stable isotope determination. Internal standards used in the laboratory were blue grama, buckeye, cocoa powder, casein, tuna muscle, graphite, whey, elemental protein, serine and IAEA 1N. Standard deviations (SDs) of reference materials were 0.05‰ for $\delta^{13}\text{C}$ and 0.07‰ $\delta^{15}\text{N}$. A total of 36 muscle samples were sent, of which 26 corresponded to green crabs (10 males, 11 females and 5 duplicates), five to *M. edulis* and five to *T. patagonica*. The error estimated through duplicate samples for green crabs muscle was 0.14‰ for both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values.

We estimated the trophic position (TP) of the green crabs using the two baselines (i.e. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of mussels and snails) and the Bayesian approach proposed by Quezada-Romegialli et al. (2018) (See equations in ESM: Online Resource 1: eq. A1, eq. A2 and eq. A3). The discrimination factors used for TP estimations were $1.80 \pm 0.40 \text{‰}$ ($\delta^{13}\text{C}$) and $2.35 \pm 1.26\text{‰}$ ($\delta^{15}\text{N}$) following published values for omnivorous crustaceans (Parker et al. 1989, Vanderklist and Ponsard 2003, Rudnick and Resh 2005, Yokoyama et al. 2005). For TP estimations sex was taken into account (i.e. independent estimations were undertaken for female and male crabs). We performed pairwise comparisons of posterior estimated TP as the probability of TP_A to be less than or equal to TP_B ($\text{TP}_A \leq \text{TP}_B$) for comparisons between sexes. We similarly performed pairwise comparisons of the proportional contribution of each baseline (α). The Bayesian probability was considered to be significant when $p > 0.95$ as an analogous of the frequentist p-value. Estimations of TP and α were done using the ‘TwoBaselinesFull’ model in the R package ‘tRophic Position’ (Quezada-Romegialli et. al. 2018).

The trophic niche of green crabs was described in terms of isotopic niche characteristics following the standard ellipse approach (Jackson et al. 2011). We calculated the standard ellipse area for males and females separately considering 40% of the data and corrected by sample size (SEAc) and the Bayesian standard ellipses area (SEAb) using the ‘SIBER’ and “rjags” packages on R (Jackson et al. 2011). Subsequently, we calculated the percentage of overlap between females and males SEA as the area at the interception divided by the SEA of male or female respectively. To test differences between SEAb we calculated 100,000 posterior iterations of SEAb based on our data set and estimated the probability that the bigger SEAb was higher than the smaller SEAb (i.e. we estimated the probability of male SEAb > female SEAb) (Jackson et al. 2011).

3. Results

What are green crabs eating in coastal environments of Patagonia Argentina?

We considered the number of stomachs analyzed ($n_{\text{(vis)}} = 183$ and $n_{\text{(met)}} = 40$) sufficient to provide a good description of the diet, because the trophic diversity curves were asymptotic and all

(n-1) and (n-2) iterations were within (Hz – 0.05) range (Fig. A1). Overall, 18.58% of the visually analyzed stomachs were empty and 28.41% contained material that was impossible to identify due to its high degree of digestion. In the metabarcoding analysis, one sample did not result in high-quality reads and four samples did not have any DNA that could be assigned to prey (i.e. they only resulted in host/green crab DNA). Therefore, the first sample was excluded from the analysis and the other four were considered as empty stomachs resulting in 10.25% of empty guts for metabarcoding.

The visual analysis of the gut content showed that green crabs at Nuevo Gulf waters had a very broad diet, with 25 prey items (Table 1). Identifications were possible due to prey remains (typically hard parts) found in gut contents. Polychaetes were identified by jaws and mollusks by opercle and radula. We also found some well preserved organisms such as tanaidaceans and amphipods, but finding an almost fully intact organism was very unusual (see Fig. A2). We detected the presence of different prey taxa such as mollusks, algae, arthropoda, annelida, fishes and echinodermata. The main prey items were Mytilidae (bivalves) with a FO_(vis) of 35.57% followed by the gastropod *T. patagonica* (12.75%).

Metabarcoding analysis of gut contents showed a similar picture to the visual analysis (i.e. a very wide omnivorous diet) but detected a higher number of phyla and provided a more detailed description at the genus level. In general, FO_(met) were higher than FO_(vis), indicating that the level of detection using metabarcoding was higher than through traditional analysis. Here, Arthropoda was the dominant phylum in the crab's gut, with a FO_(met) of 94.28%, followed by Ochrophyta (i.e. Phaeophyta) and Mollusca. The main prey items were amphipods of the Gammaridae family (82.86%), an unidentified species of Heterodonta (45.71%) and algae (*Dictyota* and an unidentified member of Ralfsiales, both with 40.00%) (Table 1).

Are there differences in diet between females and males in Patagonia Argentina?

We could observe only slight differences in prey items between males and females, evidenced by the overlapping of CIs of the simulated FO_(vis). The main differences were a higher

frequency of arthropods and a lower frequency of molluscs in the female gut contents compared to male contents (Fig. 1A). However, these differences were not significant ($Wd^*_{(vis)} = 3.14$ and $p\text{-value}_{(vis)} = 0.058$). The $FO_{(met)}$ showed less differences between male and female main prey groups (Fig. 1B); and we did not find significant differences in their diets ($Wd^*_{(met)} = 1.84$ and $p\text{-value}_{(met)} = 0.098$). Similarly, ordinations based on OTUs and phylogenetic distances showed a high degree of overlap between males and females, although some grouping can be seen in the plot (Fig. 2A).

The main difference between males and females was in their trophic diversity. Metabarcoding OTU richness was significantly higher in males than in females (7.4 ± 3.4 and 4.7 ± 2.2 for males and females, respectively; Kruskal-Wallis chi-squared = 6.208, $p\text{-value} = 0.013$) (Fig. 2B). From a total of 70 OTUs considered prey in the NCBI dataset, only 19 were shared between males and females (27%) (Fig. 2C). These corresponded roughly to the most frequently occurring OTUs, explaining the lack of clear grouping in ordinations. However, a great proportion (53%, 37 OTUs) were male-only OTUs while only 14 (20%) were female-only (Fig. 2C). These corresponded to a variety of different OTUs with low frequency, but they are responsible for the higher richness in male diet when they are taken together.

Stable isotope analysis (SIA) results were in accordance with metabarcoding analysis. Males showed a broader trophic niche than females, being $SEAc$ bigger for males ($SEAc_{males} = 1.03$ and $SEAc_{females} = 0.51$, Fig. 4.A). The Bayesian reconstruction of the SEA showed that the probability of this difference was $p = 0.05$ (Fig. 4B). However, it should be noted that some overlap at standard ellipses between males and females was present (11.89% of male SEA and 24.59% of female SEA) (Fig 4A and Fig. A3 in ESM: Online Resource 1). With respect to trophic position (TP), female crabs showed a posterior estimated TP that was slightly higher than that for males ($TP_{males} = 3.02$, $TP_{females} = 3.16$ and $TP_{overall} = 3.10$, Fig 3A) but pairwise comparisons did not result in significant differences for TP or for alpha (α) posterior estimates (Table A3). The posterior estimates of the proportional contribution of each baseline showed the phytoplankton pathway of energy to be more important than the macroalgae pathway in the nutrition of green crabs ($\alpha_{males} = 0.74$, $\alpha_{females} = 0.88$

360 and $\alpha_{\text{overall}} = 0.86$, Fig 3B). This result is in accordance with bivalves (filter feeders) being the main
361 prey item.

362 The metabarcoding analysis also showed other complementary results that are not diet-
363 related but also relevant. First, sequences from both *C. maenas* and *C. aestuarii* were detected in
364 most samples, highlighting the possibility of the presence of a hybrid crab population in Patagonian
365 coastal waters. Second, we detected sequences assigned to the genus *Carcinonemertes*, a nemertean
366 parasite known to infect populations of this crab (Torchín et al 1996).

367 4. Discussion

368 What are green crabs eating in the coastal environments of Atlantic Patagonia?

369 Green crabs in the study site at the Patagonian Atlantic coast display a broad omnivorous
370 diet, consuming the majority of phyla present at the study area. The most representative phyla
371 reported for the sampling site (Mollusca, Arthropoda and Phaeophyceae, Rechimont et al. 2013)
372 were also the prey items with the highest occurrences in our dietary samples. Rhodophyta and
373 Annelida were also found frequently but with a lower occurrence rate. These results are consistent
374 with previous studies in other areas, where green crabs have been classified as opportunist
375 omnivorous, feeding on a wide range of prey species (Elner 1981, Singh 1991, Pardal et al. 2006).
376 Other green crab populations, both native and invasive, have shown a strong preference for
377 mollusks (bivalves and gastropods) (Young and Elliot 2020). In this study, mollusks were among
378 the main prey items based on visual examinations, whilst metabarcoding indicated species of
379 amphipods of the Gammaridae family, which were present in more than 80% of the samples, as
380 main prey item. Metabarcoding also revealed the presence of other prey items such as Nemertea,
381 Porifera and Ascidiacea, which were not detected in visual analysis of stomach content.

382 The molecular method also provided a detailed description of algal diversity in the green
383 crab diet. Although algae have been previously detected, the description achieved by visual analysis

was usually very poor (Cohen et al. 1995, Pardal et al. 2006, Wilcox and Rochette 2015). This is due to the little amount of intact algae that is usually found and the difficulties associated with taxonomic recognition. DNA analysis clearly represents an improvement for the identification of taxa that are notoriously difficult to recognize in degraded samples such as gut contents (Smith et al. 2005, García-Robledo et al. 2013, Taberlet et al. 2018). Although algae were frequently detected in our crabs' diet, stable isotope analysis revealed that the major dietary contribution stemmed from phytoplankton chains (i.e. canalized by bivalve consumption). Green crabs at Atlantic Patagonia feed in an opportunist manner, with bivalves and amphipods being their preferred prey; in this manner they also ingest some algal material, but this does not significantly contribute to their nutrition.

Are there differences in diet between female and male green crabs?

Various studies have shown that there are no differences between male and female foraging, as well as no differences in prey handling time and that therefore the two sexes are ecological equivalents (Spooner et al. 2007, Young and Elliot 2020). However, we observed a more complex situation in Patagonian waters. Males and females displayed similar diets without differences in trophic position (TP) but males had a broader diet than females (i.e. higher trophic diversity). This difference was evidenced by both the isotopic niche and the metabarcoding OTU richness. The fact that males have a higher trophic diversity is probably related to differences in behavior between the sexes, e.g. females are usually found in deeper waters (Klassen and Locke 2007). Crab's chela could also be playing a role in this process, as it was previously shown and later confirmed by modeling that master chela size was a good predictor of feeding rate and energy intake (Elnor 1980, Howard et al. 2018). Such higher feeding efficiency could be indirectly influencing prey diversity in males, and suggests that females and males may not be ecological equivalents after all. However, dietary overlap between the sexes remains high and the differences are rather subtle. New dietary studies focusing on seasonality and the reproductive season will be very helpful in exploring these differences in Atlantic Patagonia.

410 **What changes can green crabs trigger in local food webs?**

411 Green crabs from newly established populations have shown to be superior foragers to long-
 412 established invaders (Rossong et al. 2011). Therefore, we expect that the most dramatic changes in
 413 local food webs will occur in the early stages of the invasion until a new equilibrium state with new
 414 relative abundances has been reached. However, such a new food web state can bring serious
 415 consequences for native populations and even lead to extirpation of some species. The potential
 416 impact of green crabs as predators at the study site (rocky shore of Punta Este) can be illustrated by
 417 comparing results of an earlier survey that was part of the Census of Marine Life, NAGISA project.
 418 In the mentioned study, a total of 35 species of benthic invertebrates and 29 macroalgae were
 419 recorded (Rechimont et al. 2013), while in the present study we found 31 benthic invertebrates and
 420 23 macroalgae in the stomachs of green crabs. Hidalgo et al. (2007) have observed that green crabs
 421 are bigger and feed with greater voracity than native predators on Patagonia rocky shores. They
 422 have provoked negative impacts worldwide such as damaging populations of *M. edulis*, American
 423 oysters (*Crassostrea virginica*), or rock crabs (*Cancer irroratus*), among others. They have even
 424 caused major economic losses for different fisheries (DeGraaf and Tyrrel 2004, Miron et al. 2005,
 425 Sigurdsson and Rochette 2013, Leignel et al. 2014, Young and Elliot 2020). In addition, bird mass
 426 mortality has been attributed to the transmission of parasites by green crabs (Camphuysen et al.,
 427 2002). Coastal birds at Patagonia are already incorporating green crabs to their diet, a fact
 428 confirmed by a recent study on the kelp gull (*Larus dominicanus*) that showed green crabs as one of
 429 their prey items (Yorio et al. 2020). Carcinophagus fish (e.g. *Mustelus schmitti*) are likely to
 430 incorporate green crabs into their diet as well, although this has not been confirmed yet (Belleggia
 431 et al. 2012, Alessandra Pasti pers. comm. July 2020). The fact that the nemertean parasite
 432 *Carcinonemertes* was observed in one of our samples highlights the possible risk of dispersion of
 433 these parasites, which had not been reported before for Argentina (Bigatti and Signorelli 2018), in
 434 green crabs and other species of native crabs.

Community structure at intertidal Patagonian rocky shores is driven by harsh physical conditions to which organisms are exposed to, rather than by consumer pressure (Bertness et al. 2006, Hidalgo et al. 2007). Invasive green crabs could have a negative impact on the community structure not only by the increase in predation pressure, but also by interfering with processes that maintain biodiversity such as facilitation (Silliman et al. 2011). That means green crabs could trigger both direct and indirect effects on Patagonian food webs. We predict that changes will begin with their preferred prey: mollusks, especially bivalves, as well as crustaceans, and then scale up to the entire food web through a cascade effect (White et al. 2006, Davies et al. 2017) (Figure 5). We also predict that high predation on foundation species such as mitylids will constitute a loss of refuge for other intertidal species, which could result in a decline in biodiversity (Hidalgo et al. 2007). Changes at Nuevo Gulf coasts have already been observed such as a dramatic decrease of *Perumytilus purpuratus* populations and *Brachidontes rodriguezii* (Mendez et al., 2019), however, the causes for this mass mortality are unknown and if it relates to the green crab invasion or environmental factors is currently unknown. In addition, other species of common native crabs are much more difficult to find at the intertidal since green crabs arrival, suggesting that green crabs could have a negative interaction with other species of local crabs.

Does the use of a multidisciplinary approach contribute to a more complete picture?

The inclusion of new techniques to trophic ecology such as stable isotope analyses and metabarcoding can add new results that would otherwise go unnoticed when using traditional techniques such as visual inspection of gut contents. Both visual and metabarcoding analysis contribute to the knowledge of diet richness and provide quantitative information about it. Stable isotope analysis allows for estimates of species trophic levels and trophic pathways (i.e. which sources contribute the most to species nutrition). The conjunction of techniques used in our work provided information on different aspects of trophic ecology, such as interspecific prey-predator interactions and intrapopulation differences (Figure 5). In particular, we observed slight differences between male and female diets related to niche breadth and we achieved a more detailed taxonomic classification of certain taxa due to the use of various techniques. The methodology used in this

work allowed us to better understand the trophic relations of the green crab in a recently invaded habitat, highlighting the importance of monitoring coastal ecosystems to detect changes that can affect these marine habitats, which provide very important ecosystem services.

Acknowledgements

We would like to thank Florencia Ríos, Manuela Funes, Juan Pablo Livore, and Maria Martha Mendez for their fieldwork support. We would like to thank Agustina Ferrando, Cynthia Ibarra, Pablo Macchi and, German Cheli for expert assessment on taxonomic identifications. We would like to thank Antonela Leiva for facilitating the *Larus dominicanus* photograph.

Funding

This research was supported by Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina) and Sistema Nacional de Datos Genómicos (SNDG, Argentina). Fieldwork campaigns and stable isotope analysis were funded by Fondo para la Investigación Científica y Tecnológica (FonCyT) under the grants PICT 2017-2403, PICT 2018-0903 and PICT 2018-0969. The metabarcoding analysis was funded by the Natural Sciences and Engineering Research Council of Canada and the Canada First Research Excellence Fund.

Ethical approval

All applicable institutional and/or national guidelines for the care and use of animals were followed. Permissions for sample collection were granted from the Secretaría de Pesca and Secretaría de Conservación de Áreas Protegidas from Chubut Province. Exportation permissions were granted in 2019 by the Dirección Nacional de Biodiversidad (Secretaría de Ambiente y Desarrollo Sustentable) under ANEXO II, resolution number: IF-2019-40388897-APN-DNBI#SGP.

Author Contributions

DEG, GB, ML, and GC conceived and designed this study. DEG, GB, and GC took the samples. GC conducted the visual analysis and prepare the samples for stable isotope analysis. GC

486 and DEG analyzed direct stomach observation and stable isotope data. ML contributed to preparing
487 the metabarcoding samples and analyzed metabarcoding data. EV visited as an invited researcher
488 the Centre for Biodiversity Genomics to obtain the metabarcoding data. EV, BT, DAL, and DS
489 contributed to the generation of metabarcoding data, molecular and bioinformatic analyses. All
490 authors contributed to manuscript writing.

491 **Conflict of Interest**

492 The authors declare that they have no conflict of interest.

493 **REFERENCES**

494 Albaina A, Fox C J, Taylor N, Hunter E, Maillard M, and Taylor MI (2010) A TaqMan real-
495 time PCR based assay targeting plaice (*Pleuronectes platessa* L.) DNA to detect predation by the
496 brown shrimp (*Crangon crangon* L.) and the shore crab (*Carcinus maenas* L.)—assay development
497 and validation. Journal of Experimental Marine Biology and Ecology, 391(1-2), 178-189.

498 Alekseyenko AV (2016) Multivariate Welch t-test on distances. Bioinformatics, 32(23),
499 3552-3558.

500 Alonso MK, Crespo EA, García NA, Pedraza SN, Mariotti PA, and Mora NJ (2002) Fishery
501 and ontogenetic driven changes in the diet of the spiny dogfish, *Squalus acanthias*, in Patagonian
502 waters, Argentina. Environmental Biology of Fishes, 63(2), 193-202.

503 Bascompte J (2009) Disentangling the web of life. Science, 325: 416 – 419

504 Belleggia M, Figueroa DE, Sánchez F, and Bremec C (2012) The feeding ecology of
505 *Mustelus schmitti* in the southwestern Atlantic: geographic variations and dietary shifts.
506 Environmental Biology of Fishes, 95(1), 99-114.

507 Bertness MD, Crain CM, Silliman BR, Bazterrica MC, Reyna MV, Hildago F, and Farina JK
508 (2006) The community structure of western Atlantic Patagonian rocky shores. Ecological
509 Monographs, 76(3), 439-460.

510 Bigatti G, and Signorelli J (2018) Marine invertebrate biodiversity from the Argentine Sea,
511 south western Atlantic. ZooKeys, (791), 47.

512 Boecklen WJ, Yarnes CT, Cook BA, and James AC (2011) On the use of stable isotopes in
513 trophic ecology. Annual review of ecology, evolution, and systematics, 42, 411-440.

514 Boschi EE, Fischbach CE, and Iorio MI (1992) Catálogo ilustrado de los crustáceos
515 estomatópodos y decápodos marinos de Argentina.

516 Braukmann TW, Ivanova NV, Prosser SW, Elbrecht V, Steinke D, Ratnasingham S, de
517 Waard JR, Sones JE, Zakharov EV, and Hebert PD (2019) Metabarcoding a diverse arthropod mock
518 community. Molecular ecology resources, 19(3), 711-727.

519 Bruno JF, Fridley JD, Bromberg KD, and Bertness MD (2005) Insights into biotic
520 interactions from studies of species invasions. Species invasions: Insights into ecology, evolution,
521 and biogeography, 13-40.

522 Carlton JT, and Cohen AN (2003) Episodic global dispersal in shallow water marine
523 organisms: the case history of the European shore crabs *Carcinus maenas* and *C. aestuarii*. Journal
524 of Biogeography, 30(12), 1809-1820.

525 Camphuysen CJ, Berrevoets CM, Cremers HJWM, Dekinga A, Dekker R, Ens BJ, van der
526 Have TM, Kats RKH, Kuiken T, Leopold MF, van der Meer J, and Piersma, T (2002) Mass
527 mortality of common eiders (*Somateria mollissima*) in the Dutch Wadden Sea, winter 1999/2000:
528 starvation in a commercially exploited wetland of international importance. Biological
529 conservation, 106(3), 303-317.

530 Chen RB, Watanabe S, and Yokota M (2004) Feeding habits of an exotic species, the
531 Mediterranean green crab *Carcinus aestuarii*, in Tokyo Bay. Fisheries science, 70(3), 430-435.

532 Clare EL (2014) Molecular detection of trophic interactions: emerging trends, distinct
533 advantages, significant considerations and conservation applications. Evolutionary applications,
534 7(9), 1144-1157.

535 Clavero M, and García-Berthou E (2005) Invasive species are a leading cause of animal
536 extinctions. Trends in ecology and evolution, 20(3), 110.

537 Cohen, A. N., Carlton, J. T., and Fountain, M. C. (1995). Introduction, dispersal and potential
538 impacts of the green crab *Carcinus maenas* in San Francisco Bay, California. *Marine Biology*,
539 122(2), 225-237.

540 Darling JA (2011) Interspecific hybridization and mitochondrial introgression in invasive
541 *Carcinus* shore crabs. *Plos One*, 6(3), e17828.

542 David P, Thebault E, Anneville O, Duyck PF, Chapuis E, and Loeuille N (2017) Impacts of
543 invasive species on food webs: a review of empirical data. In *Advances in ecological research* (Vol.
544 56, pp. 1-60). Academic Press.

545 Deagle BE, Thomas AC, McInnes JC, Clarke LJ, Vesterinen EJ, Clare EL, Kartzinel TR, and
546 Eveson JP (2019) Counting with DNA in metabarcoding studies: How should we convert sequence
547 reads to dietary data?. *Molecular ecology*, 28(2), 391-406.

548 DeGraaf JD, and Tyrrell MC (2004) Comparison of the feeding rates of two introduced crab
549 species, *Carcinus maenas* and *Hemigrapsus sanguineus*, on the blue mussel, *Mytilus edulis*.
550 *Northeastern Naturalist*, 11(2), 163-167.

551 Doherty TS, Glen AS, Nimmo DG, Ritchie EG, and Dickman CR (2016) Invasive predators
552 and global biodiversity loss. *Proceedings of the National Academy of Sciences*, 113(40), 11261-
553 11265.

554 Ebling FJ, Kitching JA, Muntz L, and Taylor CM (1964) Experimental observations of the
555 destruction of *Mytilus edulis* and *Nucella lapillus* by crabs. *Journal of Animal Ecology*, 33, 73-82.

556 Elbrecht V, Braukmann TW, Ivanova NV, Prosser SW, Hajibabaei M, Wright M, Zakharov
557 EV, Hebert PDN, and Steinke D (2019) Validation of COI metabarcoding primers for terrestrial
558 arthropods. *PeerJ*, 7, e7745.

559 Elbrecht V, and Steinke D (2019) Scaling up DNA metabarcoding for freshwater
560 macrozoobenthos monitoring. *Freshwater Biology*, 64(2), 380-387.

561 Elbrecht V, Vamos EE, Meissner K, Aroviita J, and Leese F (2017) Assessing strengths and
562 weaknesses of DNA metabarcoding based macroinvertebrate identification for routine stream
563 monitoring. *Methods in Ecology and Evolution*, 8(10), 1265-1275.

564 Elbrecht V, and Leese F (2017) Validation and development of COI metabarcoding primers
565 for freshwater macroinvertebrate bioassessment. *Frontiers in Environmental Science*, 5, 11.

566 Elner R.W (1981) Diet of green crab *Carcinus maenas* (L.) from Port Hebert, southwestern
567 Nova Scotia. *J. Shellfish Res.* 1, 89–94.

568 Elner RW (1980) The influence of temperature, sex and chela size in the foraging strategy of
569 the shore crab, *Carcinus maenas* (L.). *Marine and Freshwater Behaviour and Phy*, 7(1), 15-24.

570 Fauchald K (1977) The polychaete worms. Definitions and keys to the orders, families and
571 genera. Natural History Museum of Los Angeles County, Science Series.

572 Fridley JD, Stachowicz JJ, Naeem S, Sax DF, Seabloom EW, Smith MD, Stohlgren TJ,
573 Tilman D, and Holle BV (2007) The invasion paradox: reconciling pattern and process in species
574 invasions. *Ecology*, 88(1), 3-17.

575 Galloway AW, Brett MT, Holtgrieve GW, Ward EJ, Ballantyne AP, Burns CW, Kainz MJ,
576 Müller-Navarra DC, Persson J, Ravet JL, Strandberg U, Taipale SJ, and Alhgren G (2015) A fatty
577 acid based Bayesian approach for inferring diet in aquatic consumers. *PloS one*, 10(6), e0129723.

578 García-Robledo C, Erickson DL, Staines CL, Erwin TL, and Kress WJ (2013) Tropical
579 plant–herbivore networks: reconstructing species interactions using DNA barcodes. *PLoS One*,
580 8(1), e52967.

581 Gosztonyi AE, and Kuba L (1996) Atlas de huesos craneales y de la cintura escapular de
582 peces costeros patagónicos. Informe Técnico: Plan de Manejo integrado de la zona costera
583 patagónica. Fundación Patagonia Natural.

584 Grosholz ED, and Ruiz GM (1995) Spread and potential impact of the recently introduced
585 European green crab, *Carcinus maenas*, in central California. *Marine Biology*, 122(2), 239-247.

586 Hamidi B, Wallace K, Vasu C, and Alekseyenko AV (2019) W d* \$ W_ {d}^{*} \$-test:
587 robust distance-based multivariate analysis of variance. *Microbiome*, 7(1), 1-9.

588 Hayden B, Harrod C, and Kahilainen KK (2014) Dual fuels: Intra annual variation in the
589 relative importance of benthic and pelagic resources to maintenance, growth and reproduction in a
590 generalist salmonid fish. *Journal of Animal Ecology*, 83(6), 1501-1512.

591 Hebert PD, Cywinska A, Ball SL, and Dewaard JR (2003) Biological identifications through
592 DNA barcodes. Proceedings of the Royal Society of London. Series B: Biological Sciences,
593 270(1512), 313-321.

594 Hidalgo FJ, Baron PJ, and Orensanz JML (2005) A prediction comes true: the green crab
595 invades the Patagonian coast. Biological Invasions, 7(3), 547-552.

596 Hidalgo FJ, Silliman BR, Bazterrica MC, and Bertness MD (2007) Predation on the rocky
597 shores of Patagonia, Argentina. Estuaries and Coasts, 30(5), 886-894.

598 Howard BR, Barrios-O'Neill D, Alexander ME, Dick JT, Therriault TW, Robinson TB, and
599 Côté IM (2018) Functional responses of a cosmopolitan invader demonstrate intraspecific
600 variability in consumer-resource dynamics. PeerJ, 6, e5634.

601 Hui C, Richardson DM, Landi P, Minoarivelo HO, Garnas J, and Roy HE (2016) Defining
602 invasiveness and invasibility in ecological networks. Biological Invasions, 18(4), 971-983.

603 Hussey NE, MacNeil MA, McMeans BC, Olin JA, Dudley SF, Cliff G, Wintner SP,
604 Fennessy ST, and Fisk AT (2014) Rescaling the trophic structure of marine food webs. Ecology
605 letters, 17(2), 239-250.

606 Hyndes G, Hanson C, Vanderklift M (2013) The magnitude of spatial and temporal variation
607 in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ differs between taxonomic groups: implications for food web studies. Estuar.
608 Coast. Shelf Sci. 119, 176–187.

609 Jackson AL, Inger R, Parnell AC, and Bearhop S (2011) Comparing isotopic niche widths
610 among and within communities: SIBER—Stable Isotope Bayesian Ellipses in R. Journal of Animal
611 Ecology, 80(3), 595-602.

612 Jakob Russel (2020) MicEco: Various functions for microbial community data. R package
613 version 0.9.9

614 Klassen GJ, and Locke A (2007) A biological synopsis of the European green crab, *Carcinus*
615 *maenas*. Moncton, New Brunswick, Canada: Fisheries and Oceans Canada.

616 Lefebvre S, and Dubois S (2016) The stony road to understand isotopic enrichment and
617 turnover rates: insight into the metabolic part. *Vie Et Milieu-life And Environment*, 66(3-4), 305-
618 314.

619 Leignel VSJH, Stillman JH, Baringou S, Thabet R, and Metais I (2014) Overview on the
620 European green crab *Carcinus spp.* (Portunidae, Decapoda), one of the most famous marine
621 invaders and ecotoxicological models. *Environmental Science and Pollution Research*, 21(15),
622 9129-9144.

623 Lozupone C, and Knight R (2005) UniFrac: a new phylogenetic method for comparing
624 microbial communities. *Applied and environmental microbiology*, 71(12), 8228-8235.

625 Mackey AP, Hyndes GA, Carvalho MC, and Eyre BD (2015) Physical and biogeochemical
626 correlates of spatio-temporal variation in the $\delta^{13}\text{C}$ of marine macroalgae. *Estuarine, Coastal and*
627 *Shelf Science*, 157, 7-18.

628 McDonald PS, Jensen GC, and Armstrong DA (2001) The competitive and predatory
629 impacts of the nonindigenous crab *Carcinus maenas* (L.) on early benthic phase Dungeness crab
630 *Cancer magister* Dana. *Journal of Experimental Marine Biology and Ecology*, 258(1), 39-54.

631 McMurdie PJ, and Holmes S (2013) Phyloseq: An R package for reproducible interactive
632 analysis and graphics of microbiome census data. *PLoS ONE* 8(4):e61217.

633 Mendez MM, Livore JP, Márquez F, Gotlieb E, and Bigatti G (2019) Drástica pérdida de
634 cobertura en mejillinares del golfo nuevo. 3° Congreso Argentino de Malacología. 4 al 6 de
635 diciembre de 2019. Bahía Blanca, Argentina.

636 Miron G, Audet D, Landry T, and Moriyasu M (2005) Predation potential of the invasive
637 green crab (*Carcinus maenas*) and other common predators on commercial bivalve species found on
638 Prince Edward Island. *Journal of Shellfish Research*, 24(2), 579-586.

639 Nakahara F, Ando H, Ito H, Murakami A, Morimoto N, Yamasaki M, Takayanagi A, and
640 Isagi Y (2015) The applicability of DNA barcoding for dietary analysis of sika deer. *DNA Barcodes*,
641 3(1).

642 Nielsen JM, Clare EL, Hayden B, Brett MT, and Kratina P (2018) Diet tracing in ecology:
643 Method comparison and selection. *Methods in Ecology and Evolution*, 9(2), 278-291.

644 Oksanen J, Blanchet, FG, Kindt R, Legendre P, O'hara RB, Simpson GL, Solymos P, Henry
645 M, Szoecs S, Szoecs E, and Wagner H (2019) Vegan: community ecology package. R package
646 version 2.5-6. URL <http://CRAN.R-project.org/package=vegan>.

647 Pardal M, Baeta A, Marques J, and Cabral H (2006) Feeding ecology of the green crab,
648 *Carcinus maenas* (L., 1758) in a temperate estuary, Portugal. *Crustaceana*, 79(10), 1181-1193.

649 Parker PL, Anderson RK, and Lawrence A (1989) A $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ tracer study of nutrition
650 in aquaculture: *Penaeus vannamei* in a pond growout system. In *Stable isotopes in ecological*
651 *research* (pp. 288-303). Springer, New York, NY.

652 Philippsen JS, and Benedito E (2013) Discrimination factor in the trophic ecology of fishes:
653 a review about sources of variation and methods to obtain it. *Oecologia Australis*, 17(2), 205-2016.

654 Pielou EC (1966) The measurement of diversity in different types of biological collections.
655 *Journal of theoretical biology*, 13, 131-144.

656 Pinnegar JK, and Polunin NVC (1999) Differential fractionation of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ among
657 fish tissues: implications for the study of trophic interactions. *Functional ecology*, 13(2), 225-231.

658 Pompanon F, Deagle BE, Symondson WO, Brown DS, Jarman SN, and Taberlet P (2012)
659 Who is eating what: diet assessment using next generation sequencing. *Molecular ecology*, 21(8),
660 1931-1950.

661 Price MN, Dehal PS, and Arkin AP (2010) FastTree 2—approximately maximum-likelihood
662 trees for large alignments. *PloS one*, 5(3), e9490.

663 Quezada-Romegialli C, Jackson AL, Hayden B, Kahilainen KK, Lopes C, and Harrod C
664 (2018) tRophicPosition, an R package for the Bayesian estimation of trophic position from
665 consumer stable isotope ratios. *Methods in Ecology and Evolution*, 9(6), 1592-1599.

666 R Core Team (2020) R: A language and environment for statistical computing. R Foundation
667 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

668 Ratnasingham S, and Hebert PD (2007) BOLD: The Barcode of Life Data System
669 (<http://www.barcodinglife.org>). Molecular ecology notes, 7(3), 355-364.

670 Rechimont ME, Galvan DE, Sueiro MC, Casas G, Piriz ML, Diez ME, Primost M, Zabala
671 MS, Márquez F, Brogger M, Alfaya JF, and Bigatti G (2013) Benthic diversity and assemblage
672 structure of a north Patagonian rocky shore: a monitoring legacy of the NaGISA project. Marine
673 Biological Association of the United Kingdom. Journal of the Marine Biological Association of the
674 United Kingdom, 93(8), 2049.

675 Ripley B (2005) Boot: Bootstrap R (S-Plus) Functions (Canty). R package Version 1.2-24.

676 Rodriguez LF (2006) Can invasive species facilitate native species? Evidence of how, when,
677 and why these impacts occur. Biol Invasions 8: 927–939

678 Romanuk TN, Zhou Y, Valdovinos FS, and Martinez ND (2017) Robustness trade-offs in
679 model food webs: invasion probability decreases while invasion consequences increase with
680 connectance. In Advances in Ecological Research (Vol. 56, pp. 263-291). Academic Press.

681 Rudnick D, and Resh V (2005) Stable isotopes, mesocosms and gut content analysis
682 demonstrate trophic differences in two invasive decapod crustacea. Freshwater Biology, 50(8),
683 1323-1336

684 Russel J (2020) MicEco: Various functions for microbial community data. R package
685 version 0.9.9.

686 Schwindt E, Carlton JT, Orensanz JM, Scarabino F, and Bortolus A (2020) Past and future of
687 the marine bioinvasions along the Southwestern Atlantic. Aquatic Invasions, 15(1).

688 Sigurdsson GM, and Rochette R (2013) Predation by green crab and sand shrimp on settling
689 and recently settled American lobster postlarvae. J. Crustac. Biol. 33, 10–14

690 Silliman BR, Bertness MD, Altieri AH, Griffin JN, Bazterrica MC, Hidalgo FJ, Crain CM,
691 and Reyna MV (2011) Whole-community facilitation regulates biodiversity on Patagonian rocky
692 shores. PLoS One, 6(10), e24502.

693 Singh R (1991) Natural Diet of, and Shelter-Use by, the Green Crab, *Carcinus maenas* (L.)
694 in Nova Scotia. Master's Thesis, University of New Brunswick, Fredericton, NB, Canada,

695 Smith PJ, McVeagh SM, Allain V, and Sanchez C (2005) DNA identification of gut contents
696 of large pelagic fishes. *Journal of Fish Biology*, 67(4), 1178-1183.

697 Spivak ED, Farías NE, Ocampo EH, Lovrich GA, and Luppi TA (2019) Annotated catalogue
698 and bibliography of marine and estuarine shrimps, lobsters, crabs and their allies (Crustacea:
699 Decapoda) of Argentina and Uruguay (Southwestern Atlantic Ocean). *Frente Marítimo*, 26, 1-164.

700 Spooner EH, Coleman RA, and Attrill MJ (2007) Sex differences in body morphology and
701 multitrophic interactions involving the foraging behaviour of the crab *Carcinus maenas*. *Marine*
702 *Ecology*, 28(3), 394-403.

703 Stock BC, Jackson AL, Ward EJ, Parnell AC, Phillips DL, and Semmens BX (2018)
704 Analyzing mixing systems using a new generation of Bayesian tracer mixing models. *PeerJ*, 6,
705 e5096.

706 Taberlet P, Bonin A, Coissac E, and Zinger L (2018) *Environmental DNA: For biodiversity*
707 *research and monitoring*. Oxford University Press.

708 Taberlet P, Coissac E, Pompanon F, Brochmann C, and Willerslev E (2012) Towards next
709 generation biodiversity assessment using DNA metabarcoding. *Molecular ecology*, 21(8), 2045-
710 2050.

711 Thébault E, and Fontaine C (2010) Stability of ecological communities and the architecture
712 of mutualistic and trophic networks. *Science*, 329(5993), 853-856.

713 Thomsen MS, Olden JD, Wernberg T, Griffin JN, and Silliman BR (2011a) A broad
714 framework to organize and compare ecological invasion impacts. *Environmental research*, 111(7),
715 899-908.

716 Thomsen MS, Wernberg T, Olden JD, Griffin JN, and Silliman BR (2011b) A framework to
717 study the context-dependent impacts of marine invasions. *Journal of Experimental Marine Biology*
718 *and Ecology*, 400(1-2), 322-327.

719 Thomsen MS, Wernberg T, Altieri A, Tuya F, Gulbransen D, McGlathery KJ, Holmer M, and
720 Silliman BR (2010) Habitat cascades: the conceptual context and global relevance of facilitation

721 cascades via habitat formation and modification. Integrative and Comparative Biology, 50(2), 158-
722 175.

723 Tilman D, Isbell F, and Cowles JM (2014) Biodiversity and ecosystem functioning. Annual
724 review of ecology, evolution, and systematics, 45, 471-493.

725 Tirasin EM, and Jørgensen T (1999) An evaluation of the precision of diet description.
726 Marine Ecology Progress Series, 182, 243-252.

727 Torchin M E, Lafferty KD, and Kuris AM (1996) Infestation of an introduced host, the
728 European green crab, *Carcinus maenas*, by a symbiotic nemertean egg predator, *Carcinonemertes*
729 *epialti*. The Journal of parasitology, 449-453.

730 Torres PJ, and González-Pisani X (2016) Primer registro del cangrejo verde, *Carcinus*
731 *maenas* (Linnaeus, 1758), en Golfo Nuevo, Argentina: un nuevo límite norte de distribución en
732 costas patagónicas. Ecología austral, 26(2), 134-137.

733 Traugott M, Thalinger B, Wallinger C, and Sint D (2020) Fish as predators and prey: DNA
734 based assessment of their role in food webs. Journal of Fish Biology.

735 Traugott M, Kamenova S, Ruess L, Seeber J, and Plantegenest M (2013) Empirically
736 characterising trophic networks: what emerging DNA-based methods, stable isotope and fatty acid
737 analyses can offer. In Advances in ecological research (Vol. 49, pp. 177-224). Academic Press.

738 Unger SD, Williams LA, Diaz L, and Jachowski CB (2020) DNA barcoding to assess diet of
739 larval eastern hellbenders in North Carolina. Food Webs, 22, e00134.

740 Vanderklift MA, and Ponsard S (2003) Sources of variation in consumer-diet $\delta^{15}\text{N}$
741 enrichment: a meta-analysis. Oecologia, 136(2), 169-182.

742 White EM, Wilson JC, and Clarke AR (2006) Biotic indirect effects: a neglected concept in
743 invasion biology. Diversity and distributions, 12(4), 443-455.

744 Wilcox M, and Rochette R (2015) Does claw morphology of the green crab *Carcinus*
745 *maenas* vary in relation to its diet on rocky versus fine-sediment shores of southwest New
746 Brunswick, Bay of Fundy, Canada?. Journal of Experimental Marine Biology and Ecology, 465,
747 121-129.

748 Yokoyama H, Tamaki A, Harada K, Shimoda K, Koyama K, and Ishihi Y (2005) Variability
749 of diet-tissue isotopic fractionation in estuarine macrobenthos. Marine Ecology Progress Series,
750 296, 115-128.

751 Yorio P, Suárez N, Kasinsky T, Pollicelli M, Ibarra C, and Gatto, A (2020) The introduced
752 green crab (*Carcinus maenas*) as a novel food resource for the opportunistic kelp gull (*Larus*
753 *dominicanus*) in Argentine Patagonia. Aquatic Invasions, 15(1), 140-159.

754 Young AM, and Elliott JA (2020) Life History and Population Dynamics of Green Crabs
755 (*Carcinus maenas*). Fishes, 5(1), 4.

Table 1. List of prey taxa at green crabs stomach content determinate by metabarcoding and visual analyzes. Frequency of occurrence (FO%) for metabarcoding (met) and visual inspection (vis) of each prey taxa are shown.

Phylum	Subphylum	Class	Subclass	Order/Subclass	Family/Suborder	Genus/Specie	FO (Met)	FO (Vis)
Mollusca							74.28	46.31
		Bivalvia					68.57	35.57
			Mytilida				17.14	35.57
					Mytilidae		17.14	35.57
						Aulacomya	2.86	-
						Brachidontes/ Aulacomya/ Mytilus	5.71	-
						Brachidontes/ Perumytilus	5.71	-
			Heterodonta				62.86	-
				Heterodonta			62.86	-
					Heterodonta			-
						Heterodonta Unid. 1	25.71	-
						Heterodonta Unid. 2	45.71	-
					Myoida		2.86	-
						Myoida Unid. 1	2.86	-
						Myoida Unid. 2	2.86	-
		Gastropoda					8.57	18.12
				Trochidae			8.57	12.75
					Tegulidae		8.57	12.75
						Tegula	8.57	12.75
				Siphonariida			-	4.03
					Siphonariidae		-	4.03
						<i>Siphonaria lessoni</i>	-	4.03
		Polyplacophora					-	0.67
			Chitonida				-	0.67
				Chaetopleuridae			-	0.67
					Chaetopleura		-	0.67
Annelida							22.87	16.78
		Clitellata					11.43	-
				Haplotaxida			11.43	-
					Naididae		11.43	-
						Nadididae Unid. 1	8.57	-
						Nadididae Unid. 2	2.86	-
					Haplotaxida Unid.		5.71	-

			Haplotaxida Unid. 1	2.86	-
			Haplotaxida Unid. 2	2.86	-
	Polychaeta			22.86	-
		Phyllodocida		14.28	4.70
		Nereididae			3.36
			Platynereis	5.71	-
			Nereidae Unid.	-	3.36
		Phyllodocidae		2.86	-
			Eulalia	2.86	-
		Polynoidae		-	0.67
			Polynoidae Unid.	-	0.67
		Glyceridae		-	0.67
			Glyceridae Unid.	-	0.67
		Palpata		2.86	-
		Sabellida		2.86	-
			Sabellida Unid.	2.86	-
		Eunicida		-	0.67
		Eunicidae		-	0.67
			Eunicidae Unid.	-	0.67
		Terebellida		2.86	-
		Terebellidae		2.86	-
			Terebellidae Unid.	2.86	-
		Other Polychaeta Unid.	Polychaeta Unid.	-	11.41
Nemertea				11.43	-
	Pilidiophora			11.43	-
		Heteronemertea		11.43	-
		Heteronemertea		11.43	-
			Heteronemertea Unid.	11.43	-
Arthropoda				94.28	20.80
	Crustacea			85.71	20.13
	Malacostraca			82.86	6.03
		Amphipoda		82.86	1.34
		Gammaridae		82.86	-
			Gammaridae Unid.	82.86	-
		Corophiidae		-	1.34
			<i>Monocorophium insidiosum</i>	-	1.34
		Decapoda		5.71	-
		Porcellanidae		5.71	-
			Pachycheles	5.71	-
		Isopoda		5.71	0.67
		Sphaeromatidae		5.71	-
			Exosphaeroma/	2.86	-

			Pseudosphaeroma		
			Sphaeroma	2.86	-
		Isopoda Unid.		-	0.67
	Euphausiacea			-	0.67
		Euphausiacea Unid.		-	0.67
			Euphausiacea Unid.	-	0.67
	Tanaidacea			-	2.68
		Tanaididae		-	2.68
			<i>Tanaïs dulongii</i>	-	2.68
	Cumacea			-	0.67
		Cumacea Unid.		-	0.67
			Cumacea Unid	-	0.67
Hexanauplia				2.86	-
	Copepoda			2.86	-
		Cyclopoida		2.86	-
			Oithonidae	2.86	-
			<i>Oithona similis</i>	2.86	-
Cirripedia				-	1.32
		Cirripedia Unid.		-	1.32
			Cirripedia Unid.	-	1.32
			Cirripedia Unid.	-	1.32
Other Crustacea Unid.			Crustacea Unid.	-	18.79
Insecta				54.28	0.67
		Diptera		54.28	0.67
			Chironomidae	54.28	0.67
			Chironomus	2.86	-
			Chironomidae Unid. 1	2.86	-
			Eukiefferiella	5.71	-
			Chironomidae Unid. 2	8.57	-
			Chironomidae Unid. 3	2.86	-
			Chironomidae Unid 4.	37.14	-
			Polypedilum	2.86	-
			Tanytarsus	2.86	-
			Chironomidae Unid. 5	31.43	-
			Chironomidae Unid. 6	-	0.67
Echinodermata				-	0.67
	Echinoidea			-	0.67
		Camarodonta		-	0.67
			Temnopleuridae	-	0.67
			<i>Pseudechinus magellanicus</i>	-	0.67
Cnidaria				2.86	-

	Hydrozoa		2.86	-
		Anthoathecata	2.86	-
		Pandeidae	2.86	-
		Amphinema	2.86	-
Porifera			14.28	-
	Demospongiae		14.28	-
		Demospongiae Unid.	14.28	-
Chordata			8.57	1.32
	Ascidacea		2.86	-
		Enterogona	2.86	-
		Enterogona Unid.	2.86	-
		Enterogona Unid 1.	2.86	-
	Actinopterygii		5.71	-
		Blenniiformes	2.86	-
		Blenniiformes Unid. 1	2.86	-
		Actinopterygii Unid.	5.71	-
		Actinopterygii Unid. 1	2.86	-
		Actinopterygii Unid. 2	5.71	-
	Fishes Unid.		-	1.32
Metazoa Unid.			5.71	-
		Metazoa Unid.	5.71	-
Orchrophyta			82.86	2.68
	Phaeophyceae		82.86	2.68
		Ectocarpales	48.57	-
		Acinetosporaceae	11.43	-
		Acinetosporaceae Unid. 1.	2.86	-
		Acinetosporaceae Unid. 2	8.57	-
		Chordariaceae	40.00	-
		Hecatonema	14.28	-
		Myrionema	8.57	-
		Microspongium	8.57	-
		Chordariaceae Unid. 1	2.86	-
		Chordariaceae Unid. 2	2.86	-
		Chordariaceae Unid. 3	2.86	-
		Chordariaceae Unid. 4	2.86	-
		Ectocarpaceae	17.14	-
		Ectocarpus Unid. 1	14.28	-

Florideophyceae		Ectocarpus Unid. 2	5.71	
		Scytosiphonaceae	14.28	-
		Scytosiphon	5.71	-
		Petalonia	5.71	-
		Planosiphon	2.86	-
		Ectocarpales Unid.	5.71	-
		Ectocarpales Unid. 1	2.86	-
		Ectocarpales Unid. 2	2.86	-
		Dictyotales	40.00	-
		Dictyotaceae	40.00	-
		Dictyota	40.00	-
		Ralfsiales	40.00	-
		Ralfsiales Unid.	40.00	-
		Ralfsiales Unid. 1	40.00	-
		Ralfsiales Unid. 2	40.00	-
		Sphacelariales	2.86	-
		Sphacelariales Unid.	2.86	-
		Sphacelariales Unid. 1	2.86	-
		Laminariales	34.28	-
		Alariaceae	34.28	-
		<i>Undaria pinnatifida</i>	34.28	-
			31.43	2.68
			28.57	-
		Ceramiales	22.86	-
		Rhodomelaceae	20.00	-
		Polysiphonia	2.86	-
		Streblocladia	2.86	
		Gredgaria	17.14	
		Ceramiales Unid.	2.86	-
		Ceramiales Unid.	2.86	-
		Corallinales	5.71	-
		Corallinaceae	2.86	-
		Corallina Unid.	2.86	-
	Corallinacea Unid.	2.86	-	
	Corallinacea Unid.	2.86	-	
Bangiophyceae			2.86	-
	Bangiales		2.86	-
	Bangiaceae		2.86	-
		Bangiales Unid.	2.86	-

				Bangiaceae Unid.	2.86	
Chlorophyta					2.86	0.67
	Ulvophyceae				2.86	-
		Ulvaes			2.86	-
			Ulvaceae		2.86	-
				Ulva	2.86	-
Other Algae Unid.				Algae Unid.	-	23.49

759 Figure Legends

760 **Figure 1.** Frequency of occurrence (FO) boxplots of mayor prey groups in green crab diet from
 761 Patagonia, Argentina for: A) visual and B) Metabarcoding analyses. The lower and upper hinges
 762 correspond to the first and third quartiles (the 25th and 75th percentiles). Points are outliers (data >
 763 $1.5 \times \text{IQR}$) and horizontal lines are medians. Annelida: “Ann”, Arthropoda: “Artr”, Mollusca: “Mol”,
 764 Chordata: “Chor”, Echinodermata: “Echin”, Cnidaria: “Cnid”, Porifera: “Por”, Nemertea: “Nem”,
 765 Chlorophyta: “Chlo”, Ochrophyta: “Orch”, Rhodophyta: “Rhod”, Unidentified: “Unid”.

766 **Figure 2. A)** Ordination of samples based on COI metabarcoding data (OTUs, NCBI-based
 767 assignments), depicted by sex. Distance measure: unweighted Unifrac. **B)** Distribution of OTU
 768 richness (number of different OTUs) by sex. Letters "a" and "b" indicate differences between sexes
 769 ($p < 0.05$; Kruskal-Wallis rank-sum test). **C)** Phylogenetic tree of OTU-representative COI
 770 sequences. The final, curated taxonomic assignment of each OTU is shown at the dendrogram tips.
 771 Occurrence by sex is depicted in circles (red: females, black: males). Local support values over
 772 1,000 resamplings are shown at the nodes.

773 **Figure 3. A)** Trophic position (TP) posterior estimates for female and male green crabs **B)** Alpha
 774 (α) posterior estimates for female and male green crabs at Nuevo Gulf, Patagonia Argentina. Means
 775 are indicated by dotted lines (males: blue, female: pink and overall: black).

776 **Figure 4. A)** $\delta^{15}\text{N}\text{‰}$ and $\delta^{13}\text{C}\text{‰}$ values for female and male green crabs. Ellipses represent the
 777 standard ellipses for each sex corrected for sample size. **B)** Standard ellipse area (‰^2) estimated by
 778 Bayesian methods (SEAb) for males and females green crabs.

779 **Figure 5.** Diagram representing the different techniques and their contribution to the analysis of
 780 trophic ecology (top left). R: Richness, QC: Quantitative Contribution, TL: Trophic Level, TP:
 781 Trophic Pathway. Line width represents the relative importance of each technique. The orange
 782 circle represents intra-population variability. The above diagram represents a simplified version of

783 the Atlantic Patagonian intertidal food webs showing the possible effects of the green crab invasion.
 784 The solid red lines represent a direct negative relationship (predation on resident species). The
 785 dotted red lines represent an indirect negative relationship to resident species (competition or
 786 parasite transfer). The solid green line represents a direct positive relationship to bird species (by
 787 food facilitation). The dotted green line represents a possible positive relationship to fish (by food
 788 facilitation). F: phytoplankton, M: macroalgae, D: detritus, Z: zooplankton, H: herbivores, C: native
 789 crabs as *Cyrtograptus angulatus* or *C. altimanus*, B: bivalves, T: *Trophon geversianus*, L: *Larus*
 790 *dominicanus*, P: fishes and GC: green crab *Carcinus maenas*. Photo credits: Antonela Leiva and
 791 Gregorio Bigatti.

792 **Figure 1**

794 **Figure 3**

795 **Figure 4**

796 **Figure 5**