Inflammation in Alzheimer's disease: do sex and APOE matter?

Paula Duarte-Guterman¹, Arianne Y. Albert², Amy M. Inkster^{3,4}, Cindy K. Barha⁵, Liisa A.M. Galea¹, on behalf of the Alzheimer's Disease Neuroimaging Initiative^δ

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of British Columbia, Vancouver, BC, Canada
2. Women's Health Research Institute of British Columbia, Vancouver, BC, Canada
3. Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada.
4. BC Children's Hospital Research Institute, Vancouver, BC, Canada
5. Djavad Mowafaghian Centre for Brain Health and Department of Physical Therapy, University of British Columbia, Vancouver, BC, Canada

Address all correspondence and requests for reprints to:

L. A. M. Galea, PhD
Djavad Mowafaghian Centre for Brain Health
2215 Wesbrook Mall
Vancouver, British Columbia
V6T 1Z3, Canada
E-mail: lgalea@psych.ubc.ca.

^δ Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Running title: Inflammation in AD

ABSTRACT

Alzheimer's disease (AD) disproportionately affects females with steeper cognitive decline and more neuropathology compared to males, which is exacerbated in females carrying the APOE ϵ 4 allele. The risk of developing AD is also higher in female APOE ϵ 4 carriers in earlier age groups (aged 65-75), and the progression from cognitively normal to mild cognitive impairment (MCI) and to AD may be influenced by sex. Inflammation is observed in AD and is related to aging, stress, and neuroplasticity, and although studies are scarce, sex differences are noted in inflammation. The objective of this study was to investigate underlying physiological inflammatory mechanisms that may help explain why there are sex differences in AD and APOE ϵ 4 carriers. We investigated, using the ADNI database, the effect of sex and APOE genotype (non-carriers or carriers of 1 and 2 APOE ϵ 4 alleles) and sex and diagnosis (cognitively normal (CN), MCI, AD) on CSF (N= 279) and plasma (N= 527) markers of stress and inflammation. We found CSF IL-16 and IL-8 levels were significantly lower in female non-carriers of APOE ϵ 4 alleles compared to males, whereas levels were similar between the sexes among carriers of APOE ϵ 4 alleles. Furthermore, females had on average higher levels of plasma CRP and ICAM1 but lower levels of CSF ICAM1, IL-8, IL-16, and IgA than males. Carrying APOE ϵ 4 alleles and diagnosis (MCI and AD) decreased plasma CRP in both sexes. Sex differences in inflammatory biomarkers support that the underlying physiological changes during aging differ by sex and tissue origin.

Keywords: Sex differences; Alzheimer's disease; APOE genotype; inflammation; cytokines.

INTRODUCTION

Alzheimer's disease (AD) is a neurodegenerative disease characterized by severe cognitive decline [1]. Risk factors for AD include modifiable risk factors such as sociocultural or lifestyle factors (e.g., education, marital status, exercise), chronic stress exposure [2], and medical conditions (diabetes, obesity, and cardiovascular disease) [3–5]. Non-modifiable lifetime risk factors for AD include age, female sex, and APOE genotype [6]. However, research on the effects of biological sex on risk for AD is equivocal and may depend on geographic location (reviewed in [4,7,8]). Nevertheless, females with AD show greater cognitive decline [9–11] and neuropathology compared to males (faster brain atrophy rates, neurofibrillary tangles; [10,12–15]). Intriguingly, the presence of APOEε4 alleles increases the risk to develop AD in females compared to males at an earlier age (aged 65-75; [16]), and accelerates neuropathology and cognitive decline more so in females than in males [10,11,14,17–19], indicating that the APOE genotype interacts with sex on various factors related to AD. However, there is limited research into the role of sex and its interaction with APOE genotype in the possible mechanisms underlying AD. Understanding why females in general and female APOEε4 carriers have a higher burden of the disease is important for the development of tailored treatments. Biomarkers are highly sought after to predict disease onset and progression and to understand the underlying mechanisms of diseases in order to develop or improve treatments.

Chronic low grade inflammation is a hallmark of AD, as evidenced by increased expression of proinflammatory cytokines in the brains of AD patients (not analyzed by sex), which can exacerbate AD pathology [20–22]. There is, however, increasing evidence that there are sex differences in immune responses in healthy adults with females mounting a stronger response compared to males after an acute challenge [23,24]. In response to an endotoxin, females have higher levels of pro-inflammatory plasma cytokines (TNF-α and IL-6) while males

have higher plasma levels of anti-inflammatory IL-10 [23,25]. In addition, aging affects the immune system differently in males and females, with females having higher genomic activity for adaptive cells and males having activity for monocytes and inflammation [26]. Although limited, there is evidence that sex differences in systemic inflammation are associated with greater AD pathology [27] but not cognitive decline in normal aging [28]. Specifically, higher C-reactive protein (CRP) levels in blood beginning in midlife are associated with higher brain amyloid levels later in life in healthy males, but not in healthy females [27]. To our knowledge, very few studies have stratified by sex and APOE genotype or sex and diagnosis of cognitive status on potential biomarkers of AD, including inflammation.

Sex differences in inflammatory biomarker systems may also differentially affect neuroplasticity [29,30], which is reduced in AD and correlates with cognitive decline [31,32]. In addition, peripheral cortisol, the main stress hormone in humans, is elevated in AD [33] and is associated with higher amyloid levels in the brain [34], a reduction in hippocampal volume, and cognitive impairment in older individuals [35] that may depend on MCI status [36]. Peripheral cortisol is also associated with elevated pro-inflammatory cytokines [23,37]. However, it is not known how sex differences in markers of inflammation (e.g., cytokines, immunoglobulins, CRP, intercellular adhesion molecule, ICAM1), and stress hormones (cortisol) may be related to sex differences in AD.

Using the ADNI database, we conducted exploratory analyses examining sex differences in CSF and plasma physiological biomarkers, inflammation and stress related, and how these may be affected by APOE genotype (non-carriers or carriers of APOE ϵ 4 alleles), and dementia status (cognitively healthy (CN), MCI, AD). We tested the hypothesis that females have higher

levels of inflammation and stress hormones compared to males and these levels are disproportionately affected by the presence of APOE ϵ 4 alleles and AD diagnosis.

METHODS

ADNI database

Data used in the preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). The ADNI was launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner, MD. The primary goal of ADNI has been to test whether serial magnetic resonance imaging (MRI), positron emission tomography (PET), other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD). For up-to-date information, see www.adni-info.org. Data used in this article were downloaded on or before Jan 16, 2019. Inclusion and exclusion criteria of participants [38,39] were the same for the two datasets analysed in the current study (biomarkers in CSF and plasma), and general procedures are detailed online (<http://adni.loni.usc.edu/methods/documents/>). Briefly, cognitively normal (CN) participants had normal memory function based on education-adjusted scores on the Wechsler Memory Scale Logical Memory II and a Clinical Dementia Rating (CDR) of 0. Amnesic late MCI (LMCI) participants had objective memory loss (measured by education-adjusted scores from Wechsler Memory Scale Logical Memory II), a CDR of 0.5, preserved daily activities, and absence of dementia. All AD participants met NINCDS/ADRDA Alzheimer's Criteria and a CDR of 0.5 or 1.0.

To address our research questions, we used two separate datasets from the ADNI database: CSF biomarkers and plasma biomarkers (Table 1). Although the datasets do not

overlap completely, within the plasma-CSF datasets there is an overlap of 85% (i.e., 85% of individuals with CSF biomarker data also had plasma levels of biomarkers). This is an exploratory study of these variables on sex by APOE genotype and sex by diagnosis and we discuss the limitation of these overlapping datasets below.

Statistical Methods: Inflammatory markers

We included all ADNI participants that had inflammatory markers measured in CSF (N = 279) and plasma (N = 527) listed in Tables 1. Data included in our analyses were: demographics (age, years of education, and ethnicity), baseline diagnosis (cognitively normal, CN; late MCI, LMCI; or AD), and number of APOE ϵ 4 alleles. We collapsed APOE genotype into two groups: (1) participants carrying any ϵ 4 alleles (homozygous ϵ 4/ ϵ 4 and heterozygous ϵ 4/-) and (2) participants with no ϵ 4 risk alleles (-/-). Plasma and CSF samples from the ADNI study were collected in CN, LMCI, and AD participants at baseline in the morning after an overnight fast. Processing, aliquoting and storage were performed according to the ADNI Biomarker Core Laboratory Standard Operating Procedures. Inflammatory markers were measured using a commercially available multiplex proteomic panel (Human Discovery Multi-Analyte Profile; Luminex xMAP) developed by Rules-Based Medicine (Austin, TX), that measures a variety of markers including cytokines, metabolic markers, and growth factors. We initially chose biomarkers available in plasma involved in inflammation and immune responses (cytokines, immunoglobulins, CRP, and ICAM1) and stress (cortisol; Table 2). We analysed the same biomarkers in CSF (however IgE and IL-18 are not available in CSF). The protocols used to quantify plasma and CSF analytes are described in Craig-Schapiro et al. [40] and Hu et al. [41]. We used the ADNI quality-controlled data for plasma and CSF provided by the ADNI

Consortium. For plasma IL-16, we removed one outlier that was more than two times lower than the 25th percentile in the plasma data. Sensitivity analysis with the outlier present suggested that it was disproportionately influencing the results.

We compared all available data for each study variable between the sexes using the Wilcoxon rank sum test for continuous variables and Fisher's exact test for categorical variables. Nonparametric tests are standard for comparing variables where the distribution is unknown or expected to be non-normal. We used general linear models to determine the relationships between (1) sex and APOE genotype (non-carriers or carriers of APOEε4 alleles) or (2) sex and baseline diagnosis as predictor variables, and biomarkers as dependent variables. Due to the limited sample size, we were not able to study sex, APOE genotype, and baseline diagnosis in one model. All models included age and education as covariates. Initially, all models included an interaction between sex and presence of APOEε4 alleles or sex and baseline diagnosis; if this interaction was not significant, it was removed from the model to estimate the main effects of sex and APOE genotype or diagnosis. Significance was based on the likelihood ratio test, and all P-values for comparisons of sex and either APOE genotype or diagnosis for all outcomes combined were corrected for multiple testing using the Benjamini-Hochberg false discovery rate method with the family-wise error rate set to 0.05 [42]. In total, three P-values per dependent variable were included in each set of models (interaction term and main effects of sex and APOE or diagnosis) resulting in 27 P-values corrected in CSF (9 dependent variables) and 33 P-values corrected in plasma (11 dependent variables; Supplementary Tables S1 to S4) for each of the two models (sex and APOE and sex and diagnosis). Significant interaction terms were followed up using pairwise simple-effects tests with Benjamini-Hochberg P-value correction. A subset of participants with CSF measurements had corresponding plasma measurements (N= 237 total, N=

88 females and N=149 males). For each biomarker, we calculated Pearson's correlation coefficients between CSF and plasma levels in males and females separately. We then compared these correlations using the Fisher r-to-Z transformation and Z-test using the method by Zou [43]. We report significance differences (adjusted $P \leq 0.05$) and trends (adjusted $P \leq 0.08$). All regression analyses were carried out in R v3.5.1 [44].

RESULTS

Demographic information

Table 1 gives a summary of the variables for the participants with: CSF biomarkers (Table 1A; N=279), plasma biomarkers (Table 1B; N=527). Given the differences in sample sizes, we performed demographic analyses on the two datasets. Females were younger than males in the CSF ($P < 0.01$) and plasma data set ($P = 0.051$). In the two datasets, females had fewer years of education than males ($P_s < 0.0001$). Thus, we used age and education level as covariates in the analyses. Although there were no sex differences in distribution of APOE ϵ 4 alleles in any of the two datasets (all $P_s > 0.4$), the proportion of participants in each of the diagnosis categories was marginally different for females and males in the CSF dataset ($P = 0.051$; Table 1 A) but not in the plasma dataset ($P > 0.1$; Table 1 B).

Sex and presence of APOE ϵ 4 alleles were associated with changes in inflammatory markers

Our first aim was to investigate whether sex and APOE genotype interact to influence inflammation using biomarkers, which we analysed separately in CSF and plasma (Supplementary Table S1 and S3, respectively). Caution should be noted as inflammatory signalling can differ depending on tissue examined [45,46].

For inflammatory markers measured in CSF, only IL-16 and IL-8 elicited a significant interaction between sex and APOE genotype ($P=0.016$ and $P=0.035$, respectively; Table 3). CSF IL-16 and IL-8 levels were significantly lower in females non-carriers of APOE ϵ 4 alleles compared to males (both P 's <0.001), whereas levels were similar between the sexes in carriers of APOE ϵ 4 alleles (P 's >0.9 ; Fig 1 A and B). Furthermore, in females with APOE ϵ 4 alleles, IL-16 was significantly higher than in non-APOE ϵ 4 female carriers ($P=0.050$), while a trend was observed in males ($P=0.062$). Whereas for IL-8, males with APOE ϵ 4 alleles had lower levels of IL-8 compared to males with no APOE ϵ 4 alleles ($P=0.014$) but there was no difference in females ($P>0.3$). Regardless of sex, CSF CRP levels were lower in carriers of APOE ϵ 4 alleles compared to non-carriers (main effect of genotype: $P=0.009$; Table 3; Fig 1 C). There was a trend for an increase in IL-6 receptor levels in APOE ϵ 4 carriers regardless of sex compared to non APOE ϵ 4 carriers (main effect of genotype: $P=0.071$; Table 3). Lastly females had significantly lower CSF levels of IgA and ICAM1 and a trend for lower CSF cortisol levels compared to males (main effect of sex: $P<0.001$; $P=0.009$, and $P=0.070$, respectively; Table 3). There were no other significant main or interaction effects on any other CSF biomarkers.

For biomarkers measured in plasma, there were no significant interactions between sex and APOE genotype (Supplementary Table S3). However, females had higher plasma CRP levels (main effect of sex: $P=0.048$; Fig 1 D) and ICAM1 (trend for a main effect of sex: $P=0.051$) compared to males and significantly lower levels of IL-18 (main effect of sex: $P=0.001$; Fig 1 E) and immunoglobulin E (IgE: main effect of sex: $P<0.001$; Fig 1 F) compared to males. Furthermore, plasma CRP decreased in carriers of APOE ϵ 4 alleles compared to non-carriers (main effect of genotype: $P<0.001$; Fig 1 D).

Sex and baseline diagnosis were associated with changes in inflammatory markers

We next tested whether sex and baseline diagnosis status (CN, LMCI, and AD) influenced CSF and plasma biomarkers of inflammation (Supplementary Table S2 and S4, respectively). There were no significant interactions between sex and diagnosis for any of the tested variables in CSF (Table 4 and Supplementary Table S2) or plasma (Supplementary Table S4). For CSF levels, females had significantly lower levels of IgA (main effect of sex: $P < 0.001$) and ICAM1 (main effect of sex: $P = 0.026$) and a trend for lower IL-16 levels (main effect of sex: $P = 0.055$) compared to males (Table 4; Fig 2 A-C), but we did not observe any significant main effects of diagnosis for any CSF variable.

In plasma, we found that females had lower levels of IgE (main effect of sex: $P < 0.001$) and IL-18 compared to males (main effect of sex: $P = 0.004$) and trends for females to have higher levels of ICAM1 (main effect of sex: $P = 0.056$) and CRP (main effect of sex: $P = 0.056$; Fig. 2 D) compared to males. In addition, we found diagnosis significantly influenced plasma cortisol (main effect of baseline diagnosis: $P = 0.01$) with lower levels in LMCI compared to AD ($P < 0.001$; Fig 2 E). We found trends for diagnosis to influence plasma IL-16, CRP, and CD 40 levels (main effect of diagnosis: $P = 0.054$, $P = 0.056$; $P = 0.067$). Plasma IL-16 ($P = 0.006$) and CRP ($P = 0.006$ and $P = 0.02$) levels were lower in LMCI and AD compared to CN (Fig 2 D and F). For plasma CD 40, levels were lower in LMCI compared to AD ($P = 0.01$; Supplementary Table S4). In summary, although we detected associations between sex and diagnosis and various biomarkers, we did not find evidence of a sex and diagnosis interaction on any variables examined.

Correlations between cerebrospinal and plasma levels of biomarkers were mostly positive

The results for inflammatory markers in plasma did not always match results in CSF (Supplementary Tables S2 and S4). We therefore investigated the relationship between plasma and CSF biomarkers in males and females (Table 5, Fig 3). Perhaps surprisingly, we found the majority of biomarkers were significantly positively correlated between plasma and CSF levels in both males and females. These significant positive correlations included CRP (males, $r=0.793$; females $r=0.860$; $P's < 0.0001$), IL-6 receptor (males, $r=0.459$; females $r=0.493$, $P's < 0.0001$), IgA (males, $r=0.705$; females $r=0.529$; $P's < 0.0001$), and cortisol in both sexes (males, $r=0.176$; females, $r=0.327$; $P=0.032$ and 0.002 , respectively). IL-16 was significantly correlated in females ($r=0.290$, $P=0.006$) but only a trend in males ($r=0.156$, $P=0.058$). Plasma and CSF levels of ICAM1 and CD 40 were positively correlated in males only ($r=0.231$, $P=0.005$ and $r=0.374$, $P < 0.0001$, respectively) whereas plasma and CSF IL-3 levels were negatively correlated in females only ($r=-0.246$, $P=0.021$; Fig 3). There were significant sex differences, favoring males, in the strength of correlation between the sexes for CD 40 ($P=0.01$) and IgA ($P=0.03$), with trends for sex differences, favouring males in ICAM1 ($P=0.06$) and favouring females in IL-3 ($P=0.06$).

DISCUSSION

In the present study using ADNI data from CN, LMCI, and AD participants we found interactions between sex and APOE genotype (but not between sex and diagnosis) on CSF and plasma levels of IL-8 and IL-16 (see Table 2 for summary of the results). CSF levels of IL-8 and IL-16 were on average lower in female APOE ϵ 4 non-carriers compared to males but similar levels were found between the sexes in APOE ϵ 4 allele carriers. Regardless of sex, the APOE ϵ 4 allele was associated with decreased levels of CSF and plasma CRP. Sex differences were seen

in inflammatory markers, regardless of diagnosis or genotype, as females had lower CSF cytokines (IL-16, IL-18), CSF ICAM1, CSF and plasma immunoglobulins (IgA, IgE), and plasma IL-18. However, tissue (CSF, plasma) mattered for results for certain inflammatory markers (ICAM1 and to a lesser extent CRP) as females had higher plasma CRP and ICAM1 compared to males, opposite to what was found in CSF. Despite these differences in outcomes between plasma and CSF biomarker analyses, plasma and CSF levels were positively correlated for cortisol, CRP, IL-6 receptor, IgA in both sexes, whereas IL-16, and IL-8 were correlated in females and CD 40 and ICAM1 were correlated in males, indicating good consistency between CSF and plasma levels of these biomarkers. Intriguingly, IL-3 stood out from all these biomarkers with a negative correlation between CSF and plasma levels in females only. Males exhibited significantly stronger correlations between plasma and CSF levels for CD 40 and IgA compared to females. Sex and APOE genotype differences in CSF and plasma inflammatory markers suggest differences in underlying physiology that may affect aging and the progression of AD and this should be considered in future studies. Researchers should be cautioned to use sex as a biological variable in all analyses.

Sex interacted with presence of APOEε4 alleles to affect levels of IL-8 and IL-16

In this study, we found that sex interacted with APOE genotype to influence CSF IL-16 and IL-8. CSF IL-16 and IL-8 levels were lower in females with no APOEε4 alleles compared to males, but no sex differences in these cytokine levels were detected in participants carrying APOEε4 alleles. Our results suggest that presence of APOEε4 alleles can modulate CSF (and potentially plasma) cytokine levels in a sex-dependent way. The APOE protein can regulate transcription in vitro [47] and APOE4, but not APOE3, increases levels of IL-6 and IL-8 in vitro

[48]. In the current study, we found that the sex differences in IL-16 and IL-8 levels disappeared in carriers of APOE ϵ 4 alleles. One possibility is that the APOE4 protein regulates cytokine levels differently in males and females. IL-16 has been implicated in AD [49] and plasma IL-16 levels decreased with diagnosis (in males and females; current study) and AD severity (analysis without regard to sex; [50]). On the other hand, levels of IL-8 were not affected by diagnosis, consistent with a meta-analysis of cytokines in AD [22]. It is unclear what the impact of regulation of CSF cytokine levels by sex and APOE ϵ 4 has on AD symptoms or pathology, however given that females with APOE ϵ 4 alleles are disproportionately affected by AD during certain ages [16,18], IL-16 and IL-8 levels are unlikely to be a mechanism for this effect as differences in sex by genotype were noticed in the absence not presence of APOE ϵ 4 alleles.

Females had higher CRP levels compared to males and CRP levels were lower in APOE ϵ 4 carriers

We found that plasma and CSF levels of CRP, a widely used inflammatory and cardiovascular marker [51,52], were independently affected by sex and APOE genotype. Females, regardless of diagnosis or APOE ϵ 4 alleles, had significantly higher plasma CRP relative to males, consistent with findings in healthy individuals [53]. Higher levels of peripheral CRP may suggest higher systemic inflammation in females, which is associated with an increased risk in all-cause dementia [54]. Higher levels of serum CRP are also associated with higher levels of serum estradiol in postmenopausal healthy females [55] which suggests that sex differences in CRP levels may be partly due to sex differences in estradiol levels or other sex hormones. A recent study using the ADNI database, found that low testosterone levels was associated with higher tau pathology especially among APOE ϵ 4 carriers, regardless of sex,

suggesting that testosterone maybe neuroprotective in both sexes [56]. In addition, we found that the presence of APOE ϵ 4 alleles decreased plasma and CSF CRP levels consistent with previous research in large population studies [57,58]. In our study, we also found a trend for lower levels of plasma CRP with LMCI and AD compared to CN. Recent meta-analyses did not find differences in peripheral levels of CRP in AD compared to healthy controls [59,60]. However, in participants with mild and moderate dementia only, serum CRP levels were lower compared to the cognitively healthy group [59]. In healthy individuals, higher levels of plasma CRP in midlife are associated with a higher amyloid burden later in life in males but not females [27]. However, despite this finding, higher systemic inflammation in midlife (including CRP) is associated with greater cognitive decline later in life in both sexes in healthy individuals [28]. It is important to acknowledge evidence that midlife obesity, but not later life obesity, is associated with an increased risk to develop dementia [61,62], which may be related to altered inflammation (e.g., cytokines and CRP) due to the accumulation of adipose tissue [63,64]. It is possible that sex differences in inflammation and/or obesity earlier in life have long-term effects on the transition to MCI and/or AD.

Females had lower cytokine and immunoglobulin levels compared to males

We found some biomarkers that were affected by sex, but not diagnosis or presence of APOE ϵ 4 alleles. For example, females had lower CSF levels of ICAM1 compared to males, regardless of APOE genotype or diagnosis, but, although a trend, the opposite effect was seen in plasma. In contrast, in healthy adults (18-55 years old), serum levels of ICAM1 are lower in females compared to males [65]. ICAM1 is a type of adhesion molecule associated with microvascular endothelial activation [66] and plasma ICAM1 levels (but not CSF levels; [67])

were higher in patients with AD [67–69]. Although in the present study we did not observe a significant effect of plasma ICAM1 with diagnosis, the unadjusted P-value was 0.063 with higher levels in LMCI and AD groups. It is intriguing that females have lower CSF levels of cytokines (IL-16, IL-18), ICAM1, and immunoglobulins (IgE and IgA) but higher plasma CRP and ICAM1 levels. Although neuroinflammation is associated with AD, it may be both a product and a driver of neurodegeneration, and it may have both beneficial and detrimental roles in AD [70,71]. In AD mouse models, inflammatory cytokines (e.g., IL1 β , IL-4, IL-6, IL-10, IFN γ , TNF α) can both increase amyloid beta deposition and reduce amyloid plaque pathology [72–80]. In transgenic mice, amyloid deposition is associated with low T-cell activation suggesting that the immune system is hypo-responsive to amyloid beta [81]. Thus, increases of inflammatory markers may not always be indicative of worse neuropathology or outcomes, but may be contributing to reductions in AD neuropathology. It is also possible that males and females have varying levels of beneficial vs detrimental immune responses which can differentially affect how the disease progresses between the sexes. Indeed, we found sex differences in the correlation between CSF and plasma biomarkers (CD 40, IgA, ICAM1, and IL-3), which suggests that plasma and CSF levels may be regulated differently in males and females.

Limitations

In this exploratory study, we used two separate ADNI datasets (CSF biomarkers and plasma biomarkers) with a large overlap of individuals (85%) but different sample sizes that resulted in differences in the demographics between the datasets and power across the datasets for the analyses conducted. Because of this, the proportion of APOE or diagnosis by sex could differ across these datasets. While the proportion of sex by APOE ϵ 4 carriers did not differ

substantially between the datasets, the proportion of participants in each of the diagnosis groups was not similar across datasets causing differences in statistical power to detect the interaction term of diagnosis and sex. In addition, in this cohort the proportion of participants in the different APOE ϵ 4 allele groups was correlated with diagnosis (Supplemental Table S5). Thus, a larger cohort is required to test how sex, APOE genotype, and diagnosis interact together in one model.

More generally, the ADNI cohort is not ethnically or socioeconomically diverse, being mostly composed of self-reported white (only 12 individuals were not-white) and highly educated individuals (average 15.69 years of education). As AD incidence, prevalence, and age of onset varies by ethnicity [82–84] and education [85], our conclusions may not apply to more ethnically and socially diverse populations. In addition to sex, it is possible the underlying mechanisms of AD are different depending on ethnicity. Additionally, other pathologies in these participants, such as cancer, cardiovascular disease, smoking status, or obesity may have influenced inflammatory markers and limited our interpretations.

CONCLUSION

The current study provides evidence that sex and presence of APOE ϵ 4 alleles are associated with CSF levels of the inflammatory markers IL-16 and IL-8. We found sex differences indicating that females had lower cytokine and immunoglobulin levels but higher plasma CRP and ICAM1 levels compared to males, although the direction of the ICAM1 finding was tissue-dependent. Together, our work suggests that that presence of APOE ϵ 4 alleles can affect cytokine levels differently in males and females and the underlying pathophysiology of aging and AD may be tissue- and sex-specific.

ACKNOWLEDGMENTS

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California. Funding for this study was provided by a Canadian Institutes of Health Research (CHIR) grant to LAMG (PJT-148662). PDG is funded by the Alzheimer's Association of the USA and Brain Canada with the financial support of Health Canada through the Brain Canada Research Fund (AARF-17-529705). The views expressed

herein do not necessarily represent the views of the Minister of Health or the Government of Canada. AMI is funded by a University of British Columbia Four Year Doctoral Fellowship and the CIHR Frederick Banting and Charles Best Masters Research Award. We thank Elizabeth Perez for help with data organization.

CONFLICT OF INTEREST

The authors have no conflict of interest to report.

REFERENCES

- [1] Alzheimer's Association (2017) 2017 Alzheimer's disease facts and figures. *Alzheimer's & Dementia* **13**, 325–373.
- [2] Caruso A, Nicoletti F, Mango D, Saidi A, Orlando R, Scaccianoce S (2018) Stress as risk factor for Alzheimer's disease. *Pharmacological Research* **132**, 130–134.
- [3] Baumgart M, Snyder HM, Carrillo MC, Fazio S, Kim H, Johns H (2015) Summary of the evidence on modifiable risk factors for cognitive decline and dementia: A population-based perspective. *Alzheimers Dement* **11**, 718–726.
- [4] Nebel RA, Aggarwal NT, Barnes LL, Gallagher A, Goldstein JM, Kantarci K, Mallampalli MP, Mormino EC, Scott L, Yu WH, Maki PM, Mielke MM (2018) Understanding the impact of sex and gender in Alzheimer's disease: A call to action. *Alzheimer's & Dementia* **14**, 1171–1183.
- [5] Xu W, Tan L, Wang H-F, Jiang T, Tan M-S, Tan L, Zhao Q-F, Li J-Q, Wang J, Yu J-T (2015) Meta-analysis of modifiable risk factors for Alzheimer's disease. *J Neurol Neurosurg Psychiatry* **86**, 1299–1306.
- [6] Riedel BC, Thompson PM, Brinton RD (2016) Age, APOE and sex: Triad of risk of Alzheimer's disease. *The Journal of Steroid Biochemistry and Molecular Biology* **160**, 134–147.
- [7] Laws KR, Irvine K, Gale TM (2018) Sex differences in Alzheimer's disease. *Current Opinion in Psychiatry* **31**, 133–139.
- [8] Mielke MM (2018) Sex and Gender Differences in Alzheimer's Disease Dementia. *Psychiatr Times* **35**, 14–17.
- [9] Irvine K, Laws KR, Gale TM, Kondel TK (2012) Greater cognitive deterioration in women than men with Alzheimer's disease: a meta analysis. *J Clin Exp Neuropsychol* **34**, 989–998.
- [10] Lin KA, Choudhury KR, Rathakrishnan BG, Marks DM, Petrella JR, Doraiswamy PM (2015) Marked gender differences in progression of mild cognitive impairment over 8 years. *Alzheimer's & Dementia: Translational Research & Clinical Interventions* **1**, 103–110.
- [11] Wang X, Zhou W, Ye T, Lin X, Zhang J, Initiative for ADN (2019) Sex Difference in the Association of APOE4 with Memory Decline in Mild Cognitive Impairment. *Journal of Alzheimer's Disease* **69**, 1161–1169.
- [12] Ardekani BA, Convit A, Bachman AH (2016) Analysis of the MIRIAD Data Shows Sex Differences in Hippocampal Atrophy Progression. *J. Alzheimers Dis.* **50**, 847–857.

- [13] Barnes LL, Wilson RS, Bienias JL, Schneider JA, Evans DA, Bennett DA (2005) Sex differences in the clinical manifestations of Alzheimer disease pathology. *Arch. Gen. Psychiatry* **62**, 685–691.
- [14] Holland D, Desikan RS, Dale AM, McEvoy LK, Alzheimer’s Disease Neuroimaging Initiative (2013) Higher rates of decline for women and apolipoprotein E epsilon4 carriers. *AJNR Am J Neuroradiol* **34**, 2287–2293.
- [15] Hua X, Hibar DP, Lee S, Toga AW, Jack CR, Weiner MW, Thompson PM (2010) Sex and age differences in atrophic rates: an ADNI study with n=1368 MRI scans. *Neurobiology of Aging* **31**, 1463–1480.
- [16] Neu SC, Pa J, Kukull W, Beekly D, Kuzma A, Gangadharan P, Wang L-S, Romero K, Arneric SP, Redolfi A, Orlandi D, Frisoni GB, Au R, Devine S, Auerbach S, Espinosa A, Boada M, Ruiz A, Johnson SC, Kosciuk R, Wang J-J, Hsu W-C, Chen Y-L, Toga AW (2017) Apolipoprotein E Genotype and Sex Risk Factors for Alzheimer Disease: A Meta-analysis. *JAMA Neurol.*
- [17] Beydoun MA, Boueiz A, Abougergi MS, Kitner-Triolo MH, Beydoun HA, Resnick SM, O’Brien R, Zonderman AB (2012) Sex differences in the association of the apolipoprotein E epsilon 4 allele with incidence of dementia, cognitive impairment, and decline. *Neurobiol. Aging* **33**, 720-731.e4.
- [18] Hohman TJ, Dumitrescu L, Barnes LL, Thambisetty M, Beecham G, Kunkle B, Gifford KA, Bush WS, Chibnik LB, Mukherjee S, Jager PLD, Kukull W, Crane PK, Resnick SM, Keene CD, Montine TJ, Schellenberg GD, Haines JL, Zetterberg H, Blennow K, Larson EB, Johnson SC, Albert M, Bennett DA, Schneider JA, Jefferson AL (2018) Sex-Specific Association of Apolipoprotein E With Cerebrospinal Fluid Levels of Tau. *JAMA Neurol* **75**, 989–998.
- [19] Koran MEI, Wagener M, Hohman TJ, Alzheimer’s Neuroimaging Initiative (2017) Sex differences in the association between AD biomarkers and cognitive decline. *Brain Imaging Behav* **11**, 205–213.
- [20] Heppner FL, Ransohoff RM, Becher B (2015) Immune attack: the role of inflammation in Alzheimer disease. *Nature Reviews Neuroscience* **16**, 358–372.
- [21] Kinney JW, Bemiller SM, Murtishaw AS, Leisgang AM, Salazar AM, Lamb BT (2018) Inflammation as a central mechanism in Alzheimer’s disease. *Alzheimer’s & Dementia: Translational Research & Clinical Interventions* **4**, 575–590.
- [22] Swardfager W, Lanctôt K, Rothenburg L, Wong A, Cappell J, Herrmann N (2010) A Meta-Analysis of Cytokines in Alzheimer’s Disease. *Biological Psychiatry* **68**, 930–941.
- [23] Engler H, Benson S, Wegner A, Spreitzer I, Schedlowski M, Elsenbruch S (2016) Men and women differ in inflammatory and neuroendocrine responses to endotoxin but not in the severity of sickness symptoms. *Brain, Behavior, and Immunity* **52**, 18–26.
- [24] Klein SL, Flanagan KL (2016) Sex differences in immune responses. *Nature Reviews Immunology* **16**, 626–638.
- [25] Wegner A, Benson S, Rebernig L, Spreitzer I, Jäger M, Schedlowski M, Elsenbruch S, Engler H (2017) Sex differences in the pro-inflammatory cytokine response to endotoxin unfold in vivo but not ex vivo in healthy humans. *Innate Immun* **23**, 432–439.
- [26] Márquez EJ, Chung C, Marches R, Rossi RJ, Nehar-Belaid D, Eroglu A, Mellert DJ, Kuchel GA, Banchereau J, Ucar D (2020) Sexual-dimorphism in human immune system aging. *Nature Communications* **11**, 751.
- [27] Walker KA, Windham BG, Brown IV CH, Knopman DS, Jack Jr CR, Mosley Jr TH, Selvin E, Wong DF, Hughes TM, Zhou Y, Gross AL, Gottesman RF (2018) The Association of Mid- and Late-Life Systemic Inflammation with Brain Amyloid Deposition: The ARIC-PET Study. *Journal of Alzheimer’s Disease* **66**, 1041–1052.
- [28] Walker KA, Gottesman RF, Wu A, Knopman DS, Gross AL, Mosley TH, Selvin E, Windham BG (2019) Systemic inflammation during midlife and cognitive change over 20 years. *Neurology* **92**, e1256–e1267.

- [29] Dantzer R (2018) Neuroimmune Interactions: From the Brain to the Immune System and Vice Versa. *Physiol. Rev.* **98**, 477–504.
- [30] de Miranda AS, Zhang C-J, Katsumoto A, Teixeira AL (2017) Hippocampal adult neurogenesis: Does the immune system matter? *Journal of the Neurological Sciences* **372**, 482–495.
- [31] Apostolova LG, Dutton RA, Dinov ID, Hayashi KM, Toga AW, Cummings JL, Thompson PM (2006) Conversion of mild cognitive impairment to Alzheimer disease predicted by hippocampal atrophy maps. *Arch. Neurol.* **63**, 693–699.
- [32] Petersen RC, Jack CR, Xu YC, Waring SC, O’Brien PC, Smith GE, Ivnik RJ, Tangalos EG, Boeve BF, Kokmen E (2000) Memory and MRI-based hippocampal volumes in aging and AD. *Neurology* **54**, 581–587.
- [33] Doecke JD, Laws SM, Faux NG, Wilson W, Burnham SC, Lam C-P, Mondal A, Bedo J, Bush AI, Brown B, Ruyck KD, Ellis KA, Fowler C, Gupta VB, Head R, Macaulay SL, Pertile K, Rowe CC, Rembach A, Rodrigues M, Rumble R, Szoek C, Taddei K, Taddei T, Trounson B, Ames D, Masters CL, Martins RN, Group for the ADNI and AIB and LR (2012) Blood-Based Protein Biomarkers for Diagnosis of Alzheimer Disease. *Arch Neurol* **69**, 1318–1325.
- [34] Toledo JB, Toledo E, Weiner MW, Jack CR, Jagust W, Lee VM-Y, Shaw LM, Trojanowski JQ (2012) Cardiovascular risk factors, cortisol, and amyloid- β deposition in Alzheimer’s Disease Neuroimaging Initiative. *Alzheimers Dement* **8**, 483–489.
- [35] Lupien SJ, de Leon M, de Santi S, Convit A, Tarshish C, Nair NP, Thakur M, McEwen BS, Hauger RL, Meaney MJ (1998) Cortisol levels during human aging predict hippocampal atrophy and memory deficits. *Nat. Neurosci.* **1**, 69–73.
- [36] Murphy KJ, Sheppard PAS, Troyer AK, Hampson E, Galea LAM (2019) Sex differences in cortisol and memory following acute social stress in amnesic mild cognitive impairment. *bioRxiv* 757484.
- [37] McEwen BS, Biron CA, Brunson KW, Bulloch K, Chambers WH, Dhabhar FS, Goldfarb RH, Kitson RP, Miller AH, Spencer RL, Weiss JM (1997) The role of adrenocorticoids as modulators of immune function in health and disease: neural, endocrine and immune interactions. *Brain Research Reviews* **23**, 79–133.
- [38] Aisen P, Petersen R, Donohue M, Gamst A, Raman R, Thomas R, Walter S, Trojanowski J, Shaw L, Beckett L, Jack C, Jagust W, Toga A, Saykin A, Morris J, Weiner M (2010) Clinical core of the Alzheimer’s Disease Neuroimaging Initiative: progress and plans. *Alzheimers Dement* **6**, 239–246.
- [39] Petersen RC, Aisen PS, Beckett LA, Donohue MC, Gamst AC, Harvey DJ, Jack CR, Jagust WJ, Shaw LM, Toga AW, Trojanowski JQ, Weiner MW (2010) Alzheimer’s Disease Neuroimaging Initiative (ADNI). *Neurology* **74**, 201–209.
- [40] Craig-Schapiro R, Kuhn M, Xiong C, Pickering EH, Liu J, Misko TP, Perrin RJ, Bales KR, Soares H, Fagan AM, Holtzman DM (2011) Multiplexed Immunoassay Panel Identifies Novel CSF Biomarkers for Alzheimer’s Disease Diagnosis and Prognosis. *PLOS ONE* **6**, e18850.
- [41] Hu WT, Holtzman DM, Fagan AM, Shaw LM, Perrin R, Arnold SE, Grossman M, Xiong C, Craig-Schapiro R, Clark CM, Pickering E, Kuhn M, Chen Y, Van Deerlin VM, McCluskey L, Elman L, Karlawish J, Chen-Plotkin A, Hurtig HI, Siderowf A, Swenson F, Lee VM-Y, Morris JC, Trojanowski JQ, Soares H, Alzheimer’s Disease Neuroimaging Initiative (2012) Plasma multianalyte profiling in mild cognitive impairment and Alzheimer disease. *Neurology* **79**, 897–905.
- [42] Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**, 289–300.
- [43] Zou GY (2007) Toward using confidence intervals to compare correlations. *Psychological Methods* **12**, 399–413.

- [44] R Core Team (2018) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria., R Foundation for Statistical Computing, Vienna, Austria.
- [45] Bettcher BM, Johnson SC, Fitch R, Casaletto KB, Heffernan KS, Asthana S, Zetterberg H, Blennow K, Carlsson CM, Neuhaus J, Bendlin BB, Kramer JH (2018) CSF and Plasma Levels of Inflammation Differentially Relate to CNS Markers of Alzheimer's Disease Pathology and Neuronal Damage. *J Alzheimers Dis* **62**, 385–397.
- [46] Setiawan E, Wilson AA, Mizrahi R, Rusjan PM, Miler L, Rajkowska G, Suridjan I, Kennedy JL, Rekkas PV, Houle S, Meyer JH (2015) Role of translocator protein density, a marker of neuroinflammation, in the brain during major depressive episodes. *JAMA Psychiatry* **72**, 268–275.
- [47] Theendakara V, Peters-Libeu CA, Bredesen DE, Rao RV (2018) Transcriptional Effects of ApoE4: Relevance to Alzheimer's Disease. *Molecular Neurobiology* **55**, 5243–5254.
- [48] Theendakara V, Peters-Libeu CA, Spilman P, Poksay KS, Bredesen DE, Rao RV (2016) Direct Transcriptional Effects of Apolipoprotein E. *The Journal of Neuroscience* **36**, 685–700.
- [49] Rosa MD, Dell'Ombra N, Zambito AM, Malaguarnera M, Nicoletti F, Malaguarnera L (2006) Chitotriosidase and inflammatory mediator levels in Alzheimer's disease and cerebrovascular dementia. *European Journal of Neuroscience* **23**, 2648–2656.
- [50] Motta M, Imbesi R, Di Rosa M, Stivala F, Malaguarnera L (2007) Altered plasma cytokine levels in Alzheimer's disease: Correlation with the disease progression. *Immunology Letters* **114**, 46–51.
- [51] Koenig W, Sund M, Fröhlich M, Fischer HG, Löwel H, Döring A, Hutchinson WL, Pepys MB (1999) C-Reactive protein, a sensitive marker of inflammation, predicts future risk of coronary heart disease in initially healthy middle-aged men: results from the MONICA (Monitoring Trends and Determinants in Cardiovascular Disease) Augsburg Cohort Study, 1984 to 1992. *Circulation* **99**, 237–242.
- [52] Ridker PM, Buring JE, Shih J, Matias M, Hennekens CH (1998) Prospective study of C-reactive protein and the risk of future cardiovascular events among apparently healthy women. *Circulation* **98**, 731–733.
- [53] Khera A, McGuire DK, Murphy SA, Stanek HG, Das SR, Vongpatanasin W, Wians FH, Grundy SM, de Lemos JA (2005) Race and gender differences in C-reactive protein levels. *J. Am. Coll. Cardiol.* **46**, 464–469.
- [54] Koyama A, O'Brien J, Weuve J, Blacker D, Metti AL, Yaffe K (2013) The role of peripheral inflammatory markers in dementia and Alzheimer's disease: a meta-analysis. *J. Gerontol. A Biol. Sci. Med. Sci.* **68**, 433–440.
- [55] Eldridge RC, Wentzensen N, Pfeiffer RM, Brinton LA, Hartge P, Guillemette C, Kemp TJ, Pinto LA, Trabert B (2020) Endogenous estradiol and inflammation biomarkers: potential interacting mechanisms of obesity-related disease. *Cancer Causes Control*.
- [56] Sundermann EE, Panizzon MS, Chen X, Andrews M, Galasko D, Banks SJ, for the Alzheimer's Disease Neuroimaging Initiative (2020) Sex differences in Alzheimer's-related Tau biomarkers and a mediating effect of testosterone. *Biology of Sex Differences* **11**, 33.
- [57] Hubacek JA, Peasey A, Pikhart H, Stavek P, Kubinova R, Marmot M, Bobak M (2010) APOE polymorphism and its effect on plasma C-reactive protein levels in a large general population sample. *Hum Immunol* **71**, 304–308.
- [58] Yun Y-W, Kweon S-S, Choi J-S, Rhee J-A, Lee Y-H, Nam H-S, Jeong S-K, Park K-S, Ryu S-Y, Choi S-W, Kim HN, Cauley JA, Shin M-H (2015) APOE Polymorphism Is Associated with C-reactive Protein Levels but Not with White Blood Cell Count: Dong-gu Study and Namwon Study. *J Korean Med Sci* **30**, 860–865.
- [59] Gong C, Wei D, Wang Y, Ma J, Yuan C, Zhang W, Yu G, Zhao Y (2016) A Meta-Analysis of C-Reactive Protein in Patients With Alzheimer's Disease. *Am J Alzheimers Dis Other Dement* **31**, 194–200.

- [60] Ng A, Tam WW, Zhang MW, Ho CS, Husain SF, McIntyre RS, Ho RC (2018) IL-1 β , IL-6, TNF- α and CRP in Elderly Patients with Depression or Alzheimer's disease: Systematic Review and Meta-Analysis. *Scientific Reports* **8**, 12050.
- [61] Albanese E, Launer LJ, Egger M, Prince MJ, Giannakopoulos P, Wolters FJ, Egan K (2017) Body mass index in midlife and dementia: Systematic review and meta-regression analysis of 589,649 men and women followed in longitudinal studies. *Alzheimers Dement (Amst)* **8**, 165–178.
- [62] Fitzpatrick AL, Kuller LH, Lopez OL, Diehr P, O'Meara ES, Longstreth WT, Luchsinger JA (2009) Midlife and Late-Life Obesity and the Risk of Dementia: Cardiovascular Health Study. *Arch Neurol* **66**, 336–342.
- [63] Nguyen JCD, Killcross AS, Jenkins TA (2014) Obesity and cognitive decline: role of inflammation and vascular changes. *Front. Neurosci.* **8**,.
- [64] Parimisetty A, Dorsemans A-C, Awada R, Ravanani P, Diotel N, Lefebvre d'Hellencourt C (2016) Secret talk between adipose tissue and central nervous system via secreted factors—an emerging frontier in the neurodegenerative research. *Journal of Neuroinflammation* **13**, 67.
- [65] Ponthieux A, Herbeth B, Drosch S, Lambert D, Visvikis S (2003) Age- and Sex-related Reference Values for Serum Adhesion Molecule Concentrations in Healthy Individuals: Intercellular Adhesion Molecule-1 and E-, P-, and L-Selectin. *Clinical Chemistry* **49**, 1544–1546.
- [66] Zenaro E, Piacentini G, Constantin G (2017) The blood-brain barrier in Alzheimer's disease. *Neurobiology of Disease* **107**, 41–56.
- [67] Nielsen HM, Londos E, Minthon L, Janciauskiene SM (2007) Soluble adhesion molecules and angiotensin-converting enzyme in dementia. *Neurobiol. Dis.* **26**, 27–35.
- [68] Huang C-W, Tsai M-H, Chen N-C, Chen W-H, Lu Y-T, Lui C-C, Chang Y-T, Chang W-N, Chang AYW, Chang C-C (2015) Clinical significance of circulating vascular cell adhesion molecule-1 to white matter disintegrity in Alzheimer's dementia. *Thrombosis and Haemostasis* **114**, 1230–1240.
- [69] Rentzos M, Michalopoulou M, Nikolaou C, Cambouri C, Rombos A, Dimitrakopoulos A, Kapaki E, Vassilopoulos D (2004) Serum Levels of Soluble Intercellular Adhesion Molecule-1 and Soluble Endothelial Leukocyte Adhesion Molecule-1 in Alzheimer's Disease. *J Geriatr Psychiatry Neurol* **17**, 225–231.
- [70] Heneka MT, Carson MJ, Khoury JE, Landreth GE, Brosseron F, Feinstein DL, Jacobs AH, Wyss-Coray T, Vitorica J, Ransohoff RM, Herrup K, Frautschy SA, Finsen B, Brown GC, Verkhratsky A, Yamanaka K, Koistinaho J, Latz E, Halle A, Petzold GC, Town T, Morgan D, Shinohara ML, Perry VH, Holmes C, Bazan NG, Brooks DJ, Hunot S, Joseph B, Deigendesch N, Garaschuk O, Boddeke E, Dinarello CA, Breitner JC, Cole GM, Golenbock DT, Kummer MP (2015) Neuroinflammation in Alzheimer's disease. *The Lancet Neurology* **14**, 388–405.
- [71] Walters A, Phillips E, Zheng R, Biju M, Kuruvilla T (2016) Evidence for neuroinflammation in Alzheimer's disease: Neuroinflammation in Alzheimer's. *Progress in Neurology and Psychiatry* **20**, 25–31.
- [72] Chakrabarty P, Li A, Ceballos-Diaz C, Eddy JA, Funk CC, Moore B, DiNunno N, Rosario AM, Cruz PE, Verbeeck C, Sacino A, Nix S, Janus C, Price ND, Das P, Golde TE (2015) IL-10 Alters Immunoproteostasis in APP Mice, Increasing Plaque Burden and Worsening Cognitive Behavior. *Neuron* **85**, 519–533.
- [73] Chakrabarty P, Tianbai L, Herring A, Ceballos-Diaz C, Das P, Golde TE (2012) Hippocampal expression of murine IL-4 results in exacerbation of amyloid deposition. *Mol Neurodegener* **7**, 36.
- [74] Chakrabarty P, Herring A, Ceballos-Diaz C, Das P, Golde TE (2011) Hippocampal expression of murine TNF α results in attenuation of amyloid deposition in vivo. *Molecular Neurodegeneration* **6**, 16.

- [75] Chakrabarty P, Ceballos-Diaz C, Beccard A, Janus C, Dickson D, Golde TE, Das P (2010) IFN-gamma promotes complement expression and attenuates amyloid plaque deposition in amyloid beta precursor protein transgenic mice. *J. Immunol.* **184**, 5333–5343.
- [76] Chakrabarty P, Jansen-West K, Beccard A, Ceballos-Diaz C, Levites Y, Verbeeck C, Zubair AC, Dickson D, Golde TE, Das P (2010) Massive gliosis induced by interleukin-6 suppresses Abeta deposition in vivo: evidence against inflammation as a driving force for amyloid deposition. *FASEB J.* **24**, 548–559.
- [77] Ghosh S, Wu MD, Shaftel SS, Kyrkanides S, LaFerla FM, Olschowka JA, O'Banion MK (2013) Sustained interleukin-1 β overexpression exacerbates tau pathology despite reduced amyloid burden in an Alzheimer's mouse model. *J. Neurosci.* **33**, 5053–5064.
- [78] Guillot-Sestier M-V, Doty KR, Gate D, Rodriguez J, Leung BP, Rezai-Zadeh K, Town T (2015) Il10 Deficiency Rebalances Innate Immunity to Mitigate Alzheimer-Like Pathology. *Neuron* **85**, 534–548.
- [79] Shaftel SS, Carlson TJ, Olschowka JA, Kyrkanides S, Matousek SB, O'Banion MK (2007) Chronic interleukin-1beta expression in mouse brain leads to leukocyte infiltration and neutrophil-independent blood brain barrier permeability without overt neurodegeneration. *J. Neurosci.* **27**, 9301–9309.
- [80] Town T, Laouar Y, Pittenger C, Mori T, Szekely CA, Tan J, Duman RS, Flavell RA (2008) Blocking TGF- β –Smad2/3 innate immune signaling mitigates Alzheimer-like pathology. *Nature Medicine* **14**, 681–687.
- [81] Ferretti MT, Merlini M, Späni C, Gericke C, Schweizer N, Enzmann G, Engelhardt B, Kulic L, Suter T, Nitsch RM (2016) T-cell brain infiltration and immature antigen-presenting cells in transgenic models of Alzheimer's disease-like cerebral amyloidosis. *Brain, Behavior, and Immunity* **54**, 211–225.
- [82] Fitten LJ, Ortiz F, Fairbanks L, Bartzokis G, Lu P, Klein E, Coppola G, Ringman J (2014) Younger age of dementia diagnosis in a Hispanic population in southern California. *Int J Geriatr Psychiatry* **29**, 586–593.
- [83] Mayeda ER, Glymour MM, Quesenberry CP, Whitmer RA (2016) Inequalities in dementia incidence between six racial and ethnic groups over 14 years. *Alzheimer's & Dementia* **12**, 216–224.
- [84] Steenland K, Goldstein FC, Levey A, Wharton W (2016) A Meta-Analysis of Alzheimer's Disease Incidence and Prevalence Comparing African-Americans and Caucasians. *J. Alzheimers Dis.* **50**, 71–76.
- [85] Sharp ES, Gatz M (2011) The Relationship between Education and Dementia An Updated Systematic Review. *Alzheimer Dis Assoc Disord* **25**, 289–304.

FIGURE CAPTIONS

Figure 1. Marginal mean ($\pm 95\%$ confidence interval) of **CSF levels** of A. IL-16 (pg/ml), B. IL-8 (pg/ml), C. C-reactive protein (CRP; $\mu\text{g/ml}$), and **plasma levels** of D. CRP (CRP; $\mu\text{g/ml}$), E. IL-18 (pg/ml), and F. IgE (ng/ml) in ADNI participants by sex and presence or absence of APOE $\epsilon 4$ alleles (none or 1 and 2 alleles).

Figure 2. Marginal mean ($\pm 95\%$ confidence interval) of **CSF levels** of A. IL-16 (pg/ml), B. IgA (mg/ml), C. Intercellular adhesion molecule (ICAM1; ng/ml), and **plasma levels** of D. C-reactive protein (CRP; $\mu\text{g/ml}$), E. cortisol (ng/ml), and F. IL-16 in ADNI participants by sex and diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; and AD, Alzheimer's disease).

Figure 3. Correlations between plasma and CSF levels of A. CD 40, B. ICAM1, C. IL-3, and D. IgA in males and females separately. CD 40 and ICAM1 were positively correlated in males while IL-3 was negatively correlated in females. IgA was more strongly correlated in males compared to females (see Table 5 for details).

Figure 1

CSF

A

B

C

Plasma

D

E

F

Figure 2

CSF

Plasma

Figure 3

Table 1. Demographic and clinical information for all ADNI participants subdivided by sex. Participants with measured biomarkers in (A) cerebrospinal fluid (CSF) and (B) plasma. We collapsed APOE genotype into two groups: (1) participants carrying any $\epsilon 4$ alleles (homozygous $\epsilon 4/\epsilon 4$ and heterozygous $\epsilon 4/-$) and (2) participants with no $\epsilon 4$ risk alleles ($-/-$). In the two subdata sets, females were significantly younger and had fewer years of education than males. In data set A (but not B), there was a trend for the proportion of female and male participants in each of the diagnosis to be different ($P=0.051$) with more females (27.5 % compared to 21.8%) diagnosed with AD, more females cognitively normal (32.1% compared to 22.9%), and fewer females diagnosed with LMCI compared to males (40.4% compared to 55.3%). The proportion of female and male participants carriers and non-carriers of APOE $\epsilon 4$ alleles was not significantly different in any of the two datasets analysed. 85% of individuals with CSF biomarkers (A) had also plasma biomarkers (B). CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease.

	A. CSF				B. Plasma			
	Total	Sex		P-value	Total	Sex		P-value
		Female	Male			Female	Male	
	No. 279	No. 109	No. 170		No. 527	No. 196	No. 330	
Age								
Mean (SD)	75.15 (± 6.86)	73.75 (± 6.69)	76.04 (± 6.83)	0.007	74.75 (± 7.40)	73.79 (± 7.63)	75.32 (± 7.21)	0.051
Education (years)								
Mean (SD)	15.69 (± 2.95)	14.68 (± 2.74)	16.34 (± 2.90)	< 0.0001	15.57 (± 3.04)	14.94 (± 2.89)	15.95 (± 3.07)	< 0.0001
Ethnicity								
White	267 (95.70%)	103 (94.50%)	164 (96.47%)	0.55	498 (94.68%)	186 (94.90%)	312 (94.55%)	0.27
Not White [†]	12 (4.30%)	6 (5.50%)	6 (3.53%)		28 (5.32%)	10 (5.10%)	18 (5.45%)	
Baseline diagnosis								
CN	74 (26.5%)	35 (32.1%)	39 (22.9%)	0.051	40 (7.6%)	19 (9.7%)	21 (6.4%)	0.16
LMCI	138 (49.5%)	44 (40.4%)	94 (55.3%)		378 (71.9%)	132 (67.3%)	246 (74.5%)	
AD	67 (24.0%)	30 (27.5%)	37 (21.8%)		108 (20.5%)	45 (23.0%)	63 (19.1%)	
APOE$\epsilon 4$ allele number								
0	134 (48.03%)	51 (46.79%)	83 (48.82%)	0.81	243 (46.20%)	90 (45.92%)	153 (46.36%)	0.93
1 or 2	145 (51.97%)	58 (53.21%)	87 (51.18%)		283 (53.80%)	106 (54.08%)	177 (53.64%)	
Cortisol (ng/mL)								
Mean (SD)	16.05 (± 6.04)	14.92 (± 6.01)	16.78 (± 5.96)	0.008	2.17 (± 0.13)	2.16 (± 0.13)	2.17 (± 0.13)	0.16
C reactive protein (ug/mL)								
Mean (SD)	-2.83 (± 0.56)	-2.77 (± 0.64)	-2.87 (± 0.51)	0.23	0.12 (± 0.54)	0.21 (± 0.55)	0.07 (± 0.52)	0.003
CD40 antigen (ng/mL)								
Mean (SD)	-0.65 (± 0.12)	-0.66 (± 0.10)	-0.64 (± 0.14)	0.12	-0.12 (± 0.13)	-0.12 (± 0.13)	-0.12 (± 0.14)	0.87
Interleukin 16 (pg/mL)								
Mean (SD)	0.91 (± 0.18)	0.87 (± 0.17)	0.94 (± 0.19)	0.004	2.55 (± 0.15)	2.54 (± 0.15)	2.55 (± 0.16)	0.34
Interleukin 3 (ng/mL)								
Mean (SD)	-2.22 (± 0.32)	-2.28 (± 0.29)	-2.17 (± 0.34)	0.001	-1.65 (± 0.29)	-1.65 (± 0.29)	-1.65 (± 0.30)	0.97
Interleukin 6 receptor (ng/mL)								
Mean (SD)	-0.01 (± 0.15)	-0.02 (± 0.14)	-0.00 (± 0.15)	0.30	1.46 (± 0.14)	1.48 (± 0.14)	.45 (± 0.13)	0.02
Interleukin 8 (pg/mL)								
Mean (SD)	1.68 (± 0.15)	1.64 (± 0.11)	1.70 (± 0.16)	0.001	1.02 (± 0.19)	1.02 (± 0.21)	1.01 (± 0.18)	0.1
Intercellular adhesion molecule 1 (ng/mL)								
Mean (SD)	0.96 (± 0.44)	0.83 (± 0.33)	1.04 (± 0.48)	0.0001	2.01 (± 0.15)	2.04 (± 0.14)	2.00 (± 0.15)	0.03
Immunoglobulin A (mg/mL)								
Mean (SD)	-2.54 (± 0.31)	-2.68 (± 0.26)	-2.45 (± 0.31)	< 0.0001	0.61 (± 0.23)	0.60 (± 0.23)	0.62 (± 0.22)	0.21

P-values are from Wilcoxon rank sum tests for continuous variables and Fisher's exact tests for categorical variables. [†]Includes self-reported Black, Asian, American Indian/Alaskan, and >1 ethnicity.

Table 2. List of biomarkers analysed in the current study with their main biological function and main finding in the CSF and plasma. Main effects of sex (sex difference), APOEε4 genotype (non-carriers or carriers), and diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease) and interaction between sex and APOEε4 genotype (sex * APOEε4 genotype) are shown. Significant effects are adjusted $P \leq 0.05$ and trends are adjusted $P \leq 0.08$. See results for details. n/a - not available

Biomarker	Biological function	Results in CSF	Results in Plasma
Cortisol	Stress hormone and inflammation	Sex difference (trend): ♀ < ♂	Diagnosis: LMCI < AD
Intercellular adhesion molecule 1	Immune response, immunoglobulin family	Sex difference: ♀ < ♂	Sex difference: ♀ > ♂ Sex difference: ♀ > ♂
C-reactive protein	Inflammation	APOEε4 genotype: non-carriers > carriers	APOEε4 genotype: non-carriers > carriers Diagnosis (trend): CN > LMCI and CN > AD
CD40 antigen	Immune and inflammatory responses		Diagnosis (trend): LMCI < AD
Interleukin 3	Immune and inflammatory responses		
Interleukin 6 receptor	Immune and inflammatory responses	APOEε4 genotype (trend): non-carriers < carriers	
Interleukin 8	Immune and inflammatory responses	Sex * APOEε4 genotype: non-carriers ♀ < ♂ carriers ♀ = ♂	
Interleukin 16	Immune and inflammatory responses	Sex * APOEε4 genotype: non-carriers ♀ < ♂ carriers ♀ = ♂	Diagnosis (trend): CN > LMCI CN > AD
Immunoglobulin A	Immune and inflammatory responses	Sex difference: ♀ < ♂	
Interleukin 18	Immune and inflammatory responses	n/a	Sex difference: ♀ < ♂
Immunoglobulin E	Immune and inflammatory responses	n/a	Sex difference: ♀ < ♂

Table 3. Linear regression results for models with sex and APOE genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles). P-values are for overall tests and are FDR-adjusted. Only shown are the models with significant associations (adjusted P≤0.05) and trends (adjusted P≤0.08). All model summaries are available in Supplementary Table S1.

<i>Predictors</i>	Cortisol Cortisol ng/ml			C Reactive Protein ug/ml			Interleukin 16 pg/ml			Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml		
	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>
(Intercept)	1.43 (-7.00 – 9.87)			-3.03 (-3.84 – -2.23)			0.42 (0.15 – 0.68)			-0.27 (-0.48 – -0.06)			1.33 (1.11 – 1.54)			-2.91 (-3.35 – -2.48)			-0.23 (-0.83 – 0.38)		
AGE (years)	0.22 (0.12 – 0.32)			0.01 (-0.00 – 0.02)			0.01 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.21 (-0.45 – 0.03)			-0.01 (-0.03 – 0.02)			-0.01 (-0.01 – 0.00)			-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	1.72 (0.25 – 3.19)	0.022	0.07	-0.11 (-0.25 – 0.03)	0.126	0.22	0.13 (0.07 – 0.19)			0.02 (-0.02 – 0.05)	0.372	0.5	0.09 (0.04 – 0.14)	<0.001		0.21 (0.14 – 0.29)	<0.001	<0.001	0.18 (0.07 – 0.28)	0.001	0.009
APOE4 1 or 2 alleles (ref = 0 alleles)	0.82 (-0.55 – 2.19)	0.241	0.35	-0.22 (-0.35 – -0.09)	0.001	0.009	0.08 (0.01 – 0.14)			0.04 (0.00 – 0.07)	0.025	0.071	0.03 (-0.02 – 0.09)	0.261		-0.00 (-0.07 – 0.07)	0.898	0.97	0.08 (-0.02 – 0.18)	0.128	0.22
Interaction: Male by 1 or 2 alleles							-0.13 (-0.22 – -0.05)	0.003	0.016				-0.09 (-0.16 – -0.02)	0.008	0.035						
Observations	279			279			279			279			279			279			279		
R ² / adjusted R ²	0.095 / 0.082			0.055 / 0.042			0.122 / 0.106			0.046 / 0.032			0.092 / 0.076			0.123 / 0.111			0.105 / 0.092		

Table 4. Linear regression results for models with sex and baseline diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease). Only shown are the models with significant associations (adjusted $P \leq 0.05$) and trends (adjusted $P \leq 0.08$). P-values are for overall tests and are FDR-adjusted. All model summaries are available in Supplementary Table S2. There were no significant interactions between diagnosis and sex.

<i>Predictors</i>	Interleukin 16 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml		
	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>
(Intercept)	0.51 (0.26 – 0.77)			-2.92 (-3.35 – -2.48)			-0.19 (-0.80 – 0.42)		
AGE (years)	0.01 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.01 (-0.01 – 0.00)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	0.06 (0.02 – 0.11)	0.007	0.055	0.21 (0.14 – 0.29)	<0.001	<0.001	0.17 (0.06 – 0.28)	0.002	0.026
Diagnosis (ref = CN)		0.4	0.61		0.99	0.99		0.54	0.67
LMCI	0.01 (-0.05 – 0.06)			0.00 (-0.08 – 0.09)			0.06 (-0.06 – 0.18)		
AD	-0.03 (-0.09 – 0.03)			-0.01 (-0.11 – 0.09)			0.02 (-0.12 – 0.16)		
Observations	279			279			279		
R ² / adjusted R ²	0.099 / 0.082			0.124 / 0.107			0.101 / 0.085		

Table 5. Pearson's correlations between plasma and CSF levels of the biomarkers analysed in the current study separately in males and females. Differences in the correlations were determined using confidence intervals. Significant correlations and differences between correlations are $P \leq 0.05$ and trends are $P \leq 0.08$

	Correlation (r) in Males n = 149	P-value	Correlation (r) in Females n = 88	P-value	Difference (95% CI)	P-value
Cortisol	0.176	0.032	0.327	0.002	-0.151 (-0.388-0.101)	0.24
C reactive protein	0.793	< 0.0001	0.860	< 0.0001	-0.067 (-0.149-0.019)	0.12
CD40 antigen	0.374	< 0.0001	0.016	0.88	0.358 (0.103-0.606)	0.01
IL-16	0.156	0.058	0.290	0.006	-0.134 (-0.376-0.121)	0.30
IL-3	0.001	0.989	-0.246	0.021	0.247 (-0.016-0.493)	0.06
IL-6 receptor	0.459	< 0.0001	0.493	< 0.0001	-0.034 (-0.232-0.179)	0.75
IL-8	0.138	0.093	0.287	0.007	-0.149 (-0.392-0.107)	0.25
IgA	0.705	< 0.0001	0.529	< 0.0001	0.176 (0.012-0.361)	0.03
ICAM1	0.231	0.005	-0.021	0.849	0.252 (-0.011-0.507)	0.06

Supplemental File

Inflammation in Alzheimer's disease: do sex and APOE matter?

Paula Duarte-Guterman¹, Arianne Y. Albert², Amy M. Inkster^{3,4}, Cindy K. Barha⁵, Liisa A.M. Galea¹, on behalf of the Alzheimer's Disease Neuroimaging Initiative^δ

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of British Columbia, Vancouver, BC, Canada
2. Women's Health Research Institute of British Columbia, Vancouver, BC, Canada
3. Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada.
4. BC Children's Hospital Research Institute, Vancouver, BC, Canada
5. Djavad Mowafaghian Centre for Brain Health and Department of Physical Therapy, University of British Columbia, Vancouver, BC, Canada

Address all correspondence and requests for reprints to:

L. A. M. Galea, PhD
 Djavad Mowafaghian Centre for Brain Health
 2215 Wesbrook Mall
 Vancouver, British Columbia
 V6T 1Z3, Canada
 E-mail: lgalea@psych.ubc.ca.

^δ Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Table S1. Linear regression results for all CSF variables investigated by sex and APOE genotype (non-carriers or carriers of APOEε4 alleles).

<i>Predictors</i>	Cortisol Cortisol ng/ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>
(Intercept)	1.43 (-7.00 – 9.87)			-3.03 (-3.84 – -2.23)			-1.10 (-1.27 – -0.93)			0.42 (0.15 – 0.68)			-3.03 (-3.49 – -2.57)		
AGE (years)	0.22 (0.12 – 0.32)			0.01 (-0.00 – 0.02)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.02)		
EDUCATION (years)	-0.21 (-0.45 – 0.03)			-0.01 (-0.03 – 0.02)			-0.00 (-0.01 – 0.00)			-0.01 (-0.01 – 0.00)			0.00 (-0.01 – 0.01)		
Male (ref = Female)	1.72 (0.25 – 3.19)	0.022	0.07	-0.11 (-0.25 – 0.03)	0.126	0.22	0.01 (-0.02 – 0.04)	0.432	0.55	0.13 (0.07 – 0.19)			0.08 (-0.00 – 0.16)	0.057	0.15
APOE4 1 or 2 alleles (ref = 0 alleles)	0.82 (-0.55 – 2.19)	0.241	0.35	-0.22 (-0.35 – -0.09)	0.001	0.009	0.01 (-0.02 – 0.04)	0.59	0.72	0.08 (0.01 – 0.14)			-0.04 (-0.12 – 0.03)	0.262	0.37
Interaction: Male by 1 or 2 alleles										-0.13 (-0.22 – -0.05)	0.003	0.016			
Observations	279			279			279			279			279		
R ² / adjusted R ²	0.095 / 0.082			0.055 / 0.042			0.125 / 0.112			0.122 / 0.106			0.076 / 0.063		

Table S1. Continued

<i>Predictors</i>	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml		
	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>	<i>Estimates (CI)</i>	<i>p</i>	<i>adjusted p</i>
(Intercept)	-0.27 (-0.48 – -0.06)			1.33 (1.11 – 1.54)			-2.91 (-3.35 – -2.48)			-0.23 (-0.83 – 0.38)		
AGE (years)	0.00 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	0.02 (-0.02 – 0.05)	0.372	0.5	0.09 (0.04 – 0.14)	<0.001		0.21 (0.14 – 0.29)	<0.001	<0.001	0.18 (0.07 – 0.28)	0.001	0.009
APOE4 1 or 2 alleles (ref = 0 alleles)	0.04 (0.00 – 0.07)	0.025	0.071	0.03 (-0.02 – 0.09)	0.261		-0.00 (-0.07 – 0.07)	0.898	0.97	0.08 (-0.02 – 0.18)	0.128	0.22
Interaction: Male by 1 or 2 alleles				-0.09 (-0.16 – -0.02)	0.008	0.035						
Observations	279			279			279			279		
R ² / adjusted R ²	0.046 / 0.032			0.092 / 0.076			0.123 / 0.111			0.105 / 0.092		

Table S2. Linear regression results for all CSF variables investigated by sex and baseline diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease). There were no significant interactions between diagnosis and sex.

Predictors	Cortisol Cortisol ng/ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	1.49 (-6.96 – 9.95)			-3.12 (-3.94 – -2.30)			-1.08 (-1.25 – -0.91)			0.51 (0.26 – 0.77)			-3.03 (-3.49 – -2.57)		
AGE (years)	0.22 (0.12 – 0.32)			0.01 (-0.00 – 0.02)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.02)		
EDUCATION (years)	-0.23 (-0.47 – 0.01)			-0.00 (-0.03 – 0.02)			-0.00 (-0.01 – 0.00)			-0.01 (-0.01 – 0.00)			0.00 (-0.01 – 0.01)		
Male (ref = Female)	1.55 (0.07 – 3.04)	0.041	0.18	-0.10 (-0.24 – 0.05)	0.19	0.42	0.01 (-0.02 – 0.04)	0.51	0.65	0.06 (0.02 – 0.11)	0.007	0.055	0.08 (-0.00 – 0.16)	0.052	0.18
Diagnosis (ref = CN)		0.24	0.49		0.15	0.36		0.067	0.2		0.4	0.61		0.5	0.65
LMCI	1.23 (-0.43 – 2.88)			-0.15 (-0.31 – 0.01)			0.01 (-0.03 – 0.04)			0.01 (-0.05 – 0.06)			-0.04 (-0.13 – 0.05)		
AD	0.10 (-1.83 – 2.02)			-0.15 (-0.34 – 0.03)			-0.03 (-0.07 – 0.00)			-0.03 (-0.09 – 0.03)			-0.06 (-0.17 – 0.04)		
Observations	279			279			279			279			279		
R ² / adjusted R ²	0.100 / 0.084			0.030 / 0.012			0.141 / 0.125			0.099 / 0.082			0.077 / 0.060		

Table S2. Continued

Predictors	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	-0.22 (-0.43 – -0.01)			1.35 (1.14 – 1.56)			-2.92 (-3.35 – -2.48)			-0.19 (-0.80 – 0.42)		
AGE (years)	0.00 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	0.02 (-0.02 – 0.05)	0.42	0.61	0.04 (0.01 – 0.08)	0.019	0.12	0.21 (0.14 – 0.29)	<0.001	<0.001	0.17 (0.06 – 0.28)	0.002	0.026
Diagnosis (ref = CN)		0.46	0.65		0.85	0.96		0.99	0.99		0.54	0.67
LMCI	0.01 (-0.04 – 0.05)			0.01 (-0.03 – 0.05)			0.00 (-0.08 – 0.09)			0.06 (-0.06 – 0.18)		
AD	-0.02 (-0.07 – 0.03)			0.01 (-0.03 – 0.06)			-0.01 (-0.11 – 0.09)			0.02 (-0.12 – 0.16)		
Observations	279			279			279			279		
R ² / adjusted R ²	0.034 / 0.016			0.062 / 0.045			0.124 / 0.107			0.101 / 0.085		

Table S3. Linear regression results for all plasma variables investigated by sex and APOE genotype (non-carriers or carriers of APOEε4 alleles).

Predictors	Cortisol Cortisol ng/ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	2.06 (1.92 – 2.19)			0.46 (-0.05 – 0.97)			-0.52 (-0.65 – -0.39)			2.42 (2.27 – 2.58)			-1.86 (-2.16 – -1.57)		
AGE (years)	0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.01)			0.01 (0.00 – 0.01)			0.00 (0.00 – 0.00)			0.00 (-0.00 – 0.00)		
EDUCATION (years)	-0.00 (-0.00 – 0.00)			-0.03 (-0.04 – -0.01)			-0.00 (-0.01 – 0.00)			-0.01 (-0.01 – -0.00)			0.01 (0.00 – 0.02)		
Male (ref = Female)	0.01 (-0.01 – 0.04)	0.26	0.57	-0.13 (-0.22 – -0.04)	0.006	0.048	-0.04 (-0.07 – -0.01)			0.01 (-0.02 – 0.04)	0.50	0.79	-0.02 (-0.07 – 0.04)	0.57	0.8
APOE4 1 or 2 alleles (ref = 0 alleles)	0.02 (-0.00 – 0.04)	0.11	0.29	-0.31 (-0.39 – -0.22)	<0.001	<0.001	-0.03 (-0.07 – 0.00)			-0.01 (-0.04 – 0.01)	0.31	0.66	0.02 (-0.03 – 0.07)	0.43	0.67
Interaction: Male by 1 or 2 alleles							0.05 (0.01 – 0.10)	0.022	0.11						
Observations	527			526			527			527			527		
R ² / adjusted R ²	0.013 / 0.005			0.123 / 0.117			0.126 / 0.118			0.042 / 0.033			0.011 / 0.004		

Table S3. Continued

Predictors	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml			Interleukin 18 pg/ml			Immunoglobulin E ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	1.43 (1.29 – 1.57)			0.79 (0.60 – 0.99)			0.48 (0.25 – 0.71)			1.89 (1.75 – 2.04)			2.42 (2.24 – 2.59)			1.42 (0.88 – 1.97)		
AGE (years)	0.00 (-0.00 – 0.00)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.00)			0.00 (0.00 – 0.00)			-0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.01)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			-0.00 (-0.01 – 0.01)			0.00 (-0.00 – 0.00)			-0.00 (-0.01 – 0.00)			0.01 (-0.01 – 0.02)		
Male (ref = Female)	-0.02 (-0.05 – -0.00)	0.049	0.14	-0.05 (-0.10 – 0.00)			0.02 (-0.02 – 0.06)	0.371	0.67	-0.04 (-0.06 – -0.01)	0.008	0.051	0.06 (0.03 – 0.09)	<0.001	0.001	0.24 (0.14 – 0.33)	<0.001	<0.001
APOE4 1 or 2 alleles (ref = 0 alleles)	0.00 (-0.02 – 0.03)	0.70	0.84	-0.08 (-0.13 – -0.02)			-0.02 (-0.06 – 0.02)	0.387	0.67	0.01 (-0.02 – 0.03)	0.586	0.80	-0.04 (-0.07 – -0.01)	0.019	0.10	0.01 (-0.09 – 0.10)	0.91	0.94
Interaction: Male by 1 or 2 alleles				0.07 (0.00 – 0.14)	0.046	0.14												
Observations	527			527			527			527			527			527		
R ² / adjusted R ²	0.011 / 0.004			0.036 / 0.026			0.019 / 0.012			0.008 / 0.000			0.038 / 0.031			0.052 / 0.045		

Table S4. Linear regression results for all plasma variables investigated by sex and baseline diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease). There were no significant interactions between diagnosis and sex.

	Cortisol ng/ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
Predictors	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	2.09 (1.95 – 2.22)			0.33 (-0.21 – 0.88)			-0.55 (-0.68 – -0.42)			2.48 (2.31 – 2.65)			-1.78 (-2.08 – -1.47)		
AGE (years)	0.00 (-0.00 – 0.00)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.01)			0.00 (0.00 – 0.00)			0.00 (-0.00 – 0.00)		
EDUCATION (years)	-0.00 (-0.00 – 0.00)			-0.02 (-0.04 – -0.01)			-0.00 (-0.00 – 0.00)			-0.01 (-0.01 – -0.00)			0.01 (0.00 – 0.02)		
Male (ref = Female)	0.02 (-0.01 – 0.04)	0.18	0.44	-0.12 (-0.21 – -0.03)	0.012	0.056	-0.01 (-0.03 – 0.01)	0.38	0.59	0.01 (-0.02 – 0.04)	0.39	0.57	-0.02 (-0.07 – 0.04)	0.56	0.75
Diagnosis (ref = CN)		0.0009	0.010		0.007	0.056		0.016	0.067		0.009	0.054		0.044	0.17
LMCI	-0.02 (-0.07 – 0.02)			-0.27 (-0.44 – -0.10)			-0.02 (-0.06 – 0.02)			-0.08 (-0.13 – -0.03)			-0.04 (-0.13 – 0.06)		
AD	0.03 (-0.02 – 0.08)			-0.24 (-0.43 – -0.05)			0.02 (-0.02 – 0.07)			-0.08 (-0.14 – -0.02)			-0.11 (-0.21 – -0.00)		
Observations	527			526			527			526			527		
R ² / adjusted R ²	0.034 / 0.024			0.061 / 0.052			0.131 / 0.123			0.048 / 0.039			0.022 / 0.013		

Table S4. Continued

	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml			Interleukin 18 pg/ml			Immunoglobulin E ng/ml		
Predictors	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	1.47 (1.33 – 1.61)			0.73 (0.54 – 0.93)			0.43 (0.20 – 0.67)			1.90 (1.75 – 2.05)			2.33 (2.15 – 2.51)			1.44 (0.89 – 2.00)		
AGE (years)	0.00 (-0.00 – 0.00)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.01)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			-0.00 (-0.01 – 0.01)			0.00 (-0.00 – 0.01)			-0.00 (-0.01 – 0.00)			0.01 (-0.01 – 0.02)		
Male (ref = Female)	-0.02 (-0.05 – 0.00)	0.063	0.19	-0.01 (-0.04 – 0.03)	0.65	0.79	0.02 (-0.02 – 0.06)	0.42	0.59	-0.03 (-0.06 – -0.01)	0.011	0.056	0.06 (0.03 – 0.09)	<0.001	0.004	0.24 (0.14 – 0.33)	<0.001	<0.001
Diagnosis (ref = CN)		0.34	0.56		0.11	0.30		0.72	0.82		0.063	0.19		0.32	0.56		0.88	0.94
LMCI	-0.03 (-0.08 – 0.01)			-0.04 (-0.10 – 0.02)			0.03 (-0.05 – 0.10)			-0.01 (-0.06 – 0.03)			0.04 (-0.01 – 0.10)			-0.00 (-0.18 – 0.17)		
AD	-0.03 (-0.08 – 0.02)			-0.00 (-0.07 – 0.07)			0.02 (-0.07 – 0.10)			0.02 (-0.03 – 0.08)			0.03 (-0.03 – 0.10)			-0.03 (-0.23 – 0.16)		
Observations	527			527			527			527			527			527		
R ² / adjusted R ²	0.015 / 0.006			0.029 / 0.020			0.008 / -0.002			0.029 / 0.019			0.032 / 0.023			0.053 / 0.043		

Table S5. Contingency table for the distribution of the APOE ϵ 4 alleles (0 or 1 and 2 ϵ 4 alleles) in each of the diagnosis groups (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease).

	CN	LMCI	AD
0	55 (74.32%)	60 (43.48%)	19(28.36%)
1 and 2	19 (25.68%)	78 (56.52%)	48 (71.64%)