

Regional gene expression patterns are associated with task-specific brain activation during reward and emotion processing measured with functional MRI

Arkadiusz Komorowski¹, Ramon Vidal², Aditya Singh³, Matej Murgas¹, Tonatiuh Pena-Centeno⁴, Gregor Gryglewski¹, Siegfried Kasper⁵, Jens Wiltfang^{6,7,8}, Rupert Lanzenberger^{1*}, Roberto Goya-Maldonado^{3*}

¹Department of Psychiatry and Psychotherapy, Division of General Psychiatry, Medical University of Vienna, Vienna, Austria.

²Max Delbrück Center for Molecular Medicine, Berlin, Germany.

³Laboratory of Systems Neuroscience and Imaging in Psychiatry (SNIPLab), Department of Psychiatry and Psychotherapy, University Medical Center Goettingen (UMG), Georg-August University, Von-Siebold-Str. 5, 37075 Goettingen, Germany.

⁴Bioinformatics Unit, Department for Epigenetics and Systems Medicine in Neurodegenerative Diseases, German Center for Neurodegenerative Diseases (DZNE), Von-Siebold-Str. 3a, 37075, Goettingen, Germany.

⁵Center for Brain Research, Medical University of Vienna, Vienna, Austria.

⁶Department of Psychiatry and Psychotherapy, University Medical Center Goettingen (UMG), Georg-August University, Von-Siebold-Str. 5, 37075 Goettingen, Germany.

⁷German Center for Neurodegenerative Diseases (DZNE), Von-Siebold-Str. 3a, 37075, Goettingen, Germany.

⁸Neurosciences and Signalling Group, Institute of Biomedicine (iBiMED), Department of Medical Sciences, University of Aveiro, Aveiro, Portugal.

***Correspondence to:**

Dr. Roberto Goya-Maldonado
Labor für Systemische Neurowissenschaften und Bildgebung in der Psychiatrie
Universitätsmedizin Göttingen
Von-Siebold-Str. 5, 37075 Goettingen, Germany
Tel: +49-551-39-22244
Email: roberto.goya@med.uni-goettingen.de

Prof. Rupert Lanzenberger
Department of Psychiatry and Psychotherapy, Medical University of Vienna
Waehringerguertel 18-20, 1090 Wien, Austria
Email: rupert.lanzenberger@meduniwien.ac.at

Abstract

The exploration of the relationship between gene expression profiles and neural response patterns known to be altered in major depressive disorder provides a unique opportunity to identify novel targets for diagnosis and therapy. Here, we estimated the spatial association between genome-wide transcriptome maps and brain activation patterns from functional magnetic resonance imaging (fMRI) with two established paradigms of great relevance for mood disorders. While task-specific neural responses during emotion processing were primarily associated with expression patterns of genes involved in cellular transport, reward processing was related to neuronal development, synapse regulation, as well as gene transcription. Multimodal integration of single-site and meta-analytic imaging data with risk genes associated with depression revealed a regional susceptibility of functional activity, modulated by master regulators TCF4 and MEF2C. The identification of multiple subordinate genes correlated with fMRI maps and their corresponding regulators presumably will reshape the prospects for neuroimaging genetics, especially for psychiatric disorders.

Introduction

Over the last decades, genetic and neuroimaging studies have significantly contributed to current knowledge about human neural functions. While single neuroscientific methods contributed to the comprehension of physiological processes as well as pathological alterations in psychiatric disorders, a multimodal integration of large-scale data has proven to be even more conducive for in-depth understanding, especially on the molecular scale^{1,2}. In particular, post-mortem gene expression data from the Allen Human Brain Atlas (AHBA) was repeatedly applied to investigate the relationship between the transcriptome and protein distribution³, brain connectivity⁴, or morphology⁵. These studies add up to numerous wide-ranging research findings combining topological mRNA expression with neuroimaging properties, partially offering toolboxes for an integrative data analysis^{6,7}. Different methods offer designated benefits and disadvantages, e.g. the precise temporal resolution of electroencephalography contrasts to its low spatial resolution⁸. In case of positron emission tomography, molecular specificity allows quantification of protein distributions in vivo, but availability of specific radioligands is limited⁹. The most prominent feature of the AHBA, however, is the availability of widespread gene expression across the whole brain. Regarding the measurement of brain activation during execution of specific paradigms, functional magnetic resonance imaging (fMRI) has evolved into one of the most popular neuroimaging techniques. Although a promising approach to contrast topological brain activation and gene expression patterns making use of the meta-analytic Neurosynth database was presented by Fox and colleagues¹⁰, their study lacked disease-related aspects. Other studies, by contrast, have successfully integrated post-mortem data and in vivo imaging findings, to assess the influence of regional gene expression on neurological or psychiatric disorders^{11,12}. While fMRI can depict neural correlates of specific psychological processes with high spatial resolution, measured signal patterns also appear susceptible to genotype variations¹³. In this context, the role of specific master regulators (MRs) modulating differentially expressed

genes that might influence regional blood oxygenation level dependent (BOLD) signaling has not been resolved yet.

From all psychiatric disorders, major depressive disorder (MDD) is now the leading cause of disability worldwide and a strong contributor to the overall global burden of disease, with increasing prevalence over the years¹⁴ as well as occurrence of subthreshold symptoms within the general population¹⁵. Granted that additive genetic effects attribute to approximately 9 % of the variation in liability of MDD¹⁶, regional variations in gene expression profiles may determine brain function within a continuum, spanning from a physiological to a more critical pathological state. In this regard, paradigms examining core depressive symptoms like impaired affect modulation or loss of interest and pleasure in common experiences are amongst the best-established within the realms of MDD. Furthermore, alterations of BOLD reactivity during processing of negatively valenced information as well as motivation- or incentive-based learning emphasize the characterization of neuropsychiatric disorders in terms of functions rather than diagnoses^{17,18}. Positive valence systems and systems for social processes thereby comprise prominent behavioral elements currently part of the Research Domain Criteria (RDoC) framework, which justifies the application of emotion and reward processing paradigms to investigate major domains of human functioning.

In this study, static gene expression patterns that correlated with task-specific brain activation were evaluated in regard to biological processes according to the Gene Ontology (GO) database¹⁹ and their relationship with MDD risk genes obtained from conventional Genome-wide association studies (GWAS)¹⁶. Potential influences of master regulating genes on functional neuroimaging properties were evaluated by means of single-site fMRI data and subsequently validated using meta-analytic activation maps from the Neurosynth online framework²⁰. Ultimately, we aimed to identify MRs implicated in depression that exert effects on individual genes associated with topological brain activation patterns during emotion and reward processing.

Results

Topological specificity of emotion and reward processing

In conjunction with known abnormalities in social interaction and reward responsiveness of patients with depressive disorders, fMRI activation within the social processes and positive valence systems domains of the RDoC framework was evaluated. We minimized unspecific signal variations related to visual, auditory, attentional and executive processing by contrasting brain activation elicited by the experimental condition with task control conditions in each participant. Sequentially, second level analyses provided specific activation patterns elicited by emotion and reward processing (full acquisition and analysis pipeline described in Methods section). The applied emotional face recognition task didn't focus attention on the emotional content of presented faces, but requested gender discrimination instead, provoking rather implicit emotional processing in limbic as well as non-limbic areas such as prefrontal cortices. When testing reward responsiveness, significant activations in dopaminergic brain regions were observed after acceptance of priorly conditioned stimuli, particularly in the mesolimbic reward system (Supplementary Table 1).

To expand the scope of results generated with single-site fMRI data, we obtained corresponding uniformity maps from the Neurosynth database, which inform about the consistency of functional brain activation for particular processes of emotion and reward circuits. Spatial activation clusters detected at our institution were thereby validated using matching statistical inference maps derived from 91 and 246 studies associated with the terms "fearful faces" and "rewards". Both meta-analytic maps, which can be obtained online from the Neurosynth framework, were representative of the expected neural correlates elicited by emotional face and reward tasks, respectively.

Associations between the transcriptome and functional brain activation

For each paradigm, correlation analyses showed spatial associations between functional brain activation and mRNA expression for a large number of individual genes (rankings of genes as well as corresponding Spearman's correlation coefficients for both datasets are reported in Supplementary Table 2). We performed all correlation analyses separately for cortical and subcortical regions, to prevent a possible bias that might arise from expression differences of the transcriptome between broad anatomical areas. Findings were highly specific for each paradigm, accounted for by the weak overlap of compiled correlation lists between the two psychological processes, i.e. emotion and reward processing (subcortex: $\rho_{\text{RRHO}} = -0.282$; cortex: $\rho_{\text{RRHO}} = 0.205$) (Supplementary Fig. 1). In contrast, high agreement of ranked mRNA-fMRI correlations between two different brain parcellation atlases was observed both for emotional face recognition (subcortex: $\rho_{\text{RRHO}} = 0.697$; cortex: $\rho_{\text{RRHO}} = 0.822$) and acceptance of monetary rewards (subcortex: $\rho_{\text{RRHO}} = 0.748$; cortex: $\rho_{\text{RRHO}} = 0.918$) (Fig. 1).

Regarding single-site data, brain activity patterns evoked by emotion processing correlated positively as well as negatively with whole-brain transcriptome maps. Region-wise correlations yielded similar results compared to voxel-wise analyses, ranging from $\rho = -0.739$ to $\rho = 0.865$ for subcortical and from $\rho = -0.449$ to $\rho = 0.431$ for cortical regions (Supplementary Fig. 2). Out of all spatial associations between gene expression and brain function the 10 highest positive correlating genes are listed in Table 1. In subcortical regions, MALL showed the strongest voxel-wise correlation ($\rho = 0.633$), while C10orf125 showed the highest region-wise correlation ($\rho = 0.865$). In the cortex, SPDYA yielded strongest voxel-wise ($\rho = 0.328$) and FOXN4 strongest region-wise ($\rho = 0.431$) correlation. FOXN4 also showed a high cortical ranking applying voxel-wise analysis ($\rho = 0.285$, 5th rank) (Fig. 2a,b).

Analogous to the emotion task, compiled ranked lists included genes spatially correlated with measured brain activity patterns related to the reward system, whereby the 10 highest positive correlating genes are listed in Table 2. Region-wise analyses yielded higher correlation coefficients than the voxel-wise approach with less prominent associations in the cortex ($\rho = -0.639$ to $\rho = 0.698$), compared to subcortical regions ($\rho = -0.788$ to $\rho = 0.81$). In the subcortex, MDK showed the strongest voxel-wise correlation of all genes ($\rho = 0.488$) and also a high region-wise correlation coefficient ($\rho = 0.803$, 3rd rank) (Fig. 2c,d). Comparing strongest voxel-wise vs. region-wise correlations in the cortex, 7 out of 10 genes were congruent (DUSP3, CA10, PIK3CD, HDAC9, LASS6, GRB14, OLFM3), indicating high agreement between both approaches (Supplementary Fig. 3). Overall, gene expression of DUSP3 showed strongest cortical correlations with reward processing both in the voxel-wise ($\rho = 0.548$) as well as the region-wise analysis ($\rho = 0.698$) (Supplementary Fig. 4).

Ontological analysis of task-specific biological processes

Considering both strong positive and negative mRNA-fMRI correlations, multiple genes associated with imaging data overlapped with specific gene sets listed in the GO knowledgebase (presented in detail in Supplementary Table 3 and Supplementary Fig. 5). Notably, a marked redundancy between GO terms was present within each paradigm, indicating rather task-specific associations of molecular programs with gene expression patterns throughout the human brain (Fig. 3). Considering neural responses during emotional face recognition in subcortical regions, significantly associated biological programs were related to cellular transport processes and mainly included genes that showed positive correlations between gene expression and imaging data. In line with the assumed relevance of molecular transduction for emotion regulation processes, the most significant overlap was observed for the GO term peptide transport. In total, 203 genes positively correlated with emotion processing were also present within this gene set ($p_{\text{corr}} = 0.013$), representing 7.9 %

of genes included within the GO term (GO:0015833). Functional brain activation in cortical regions yielded no significant overlap with biological processes. Analyzing neural responses during acceptance of monetary rewards in subcortical regions, associated GO terms mainly included genes with positive mRNA-fMRI correlations, predominantly comprising transcription processes. Highest significance was present for nucleic acid metabolic process (GO:0090304, $p_{\text{corr}} < 0.001$, 536 genes, 8.95 %), RNA metabolic process (GO:0016070, $p_{\text{corr}} < 0.001$, 486 genes, 8.89 %), as well as cellular macromolecule biosynthetic process (GO:0034645, $p_{\text{corr}} < 0.001$, 485 genes, 8.45 %). GO terms associated with genes negatively correlated with the acceptance of monetary rewards within subcortical regions mainly related to synaptic processes and neuronal development. Thereby, most significant GO terms were chemical synaptic transmission (GO:0007268, $p_{\text{corr}} < 0.001$, 54 genes, representing 6.19 % of genes included within this term) and modulation of chemical synaptic transmission (GO:0050804, $p_{\text{corr}} < 0.001$, 35 genes, 6.4 %). However, in the cortex only two overlapping GO terms were found for genes positively associated with reward processing (autophagy of mitochondrion, GO:0000422, $p_{\text{corr}} < 0.05$, 4 genes, 3.96 % and sensory perception of sound, GO:0007605, $p_{\text{corr}} = 0.04$, 4 genes, 2.34 %).

The role of risk genes implicated in depression

The relationship between neuronal activation patterns and 42 functional genes associated with MDD obtained from a pre-defined gene set was investigated to evaluate the superordinate role of these risk genes on specific BOLD activation. Single-site master regulator analysis of co-regulatory networks based on previously compiled ranked lists including strongest mRNA-fMRI correlations and revealed individual candidate MRs for each paradigm that potentially regulate genes spatially associated with imaging data. We found 4 regulators from the MDD risk gene set for emotion and 3 for reward processing in subcortical regions, with 79 and 89 possible targets, respectively ($p < 0.001$) (Table 3). Curiously, PAX6, LHX2, as well as

MEF2C were associated with negatively correlated genes for both paradigms, indicating a rather superordinate role of these MRs in MDD, regardless of cognitive system. Alternatively, TCF4 was identified both as a regulator for genes negatively correlated with emotion as well as genes positively correlated with reward processing. Overall, strongest regulation was found for genes negatively correlated with reward processing, whereby LHX2 and MEF2C coordinated more than half of the possible targets. Considering regulation for genes that showed positive spatial correlations with brain activation patterns, two MRs (SOX5 and TCF4) were identified for reward processing, while no significant regulators were found for genes positively correlated with emotion processing. While corresponding cortical analyses yielded no significant results, subcortical results were validated by means of an independent meta-analytical dataset from the Neurosynth framework. Identically, MEF2C emerged as regulator of genes negatively correlated with emotional face recognition as well as acceptance of monetary rewards (Fig. 4a), while TCF4 showed an inversed regulation of genes correlated with the emotion (negative association) and reward (positive association) paradigm (Fig. 4b). Both, single-site and meta-analytical results from gene set enrichment analysis (GSEA) complemented our findings from GO as well as master regulator analyses, showing an inversed aggregation of MDD risk genes for emotion and reward processing. Rather than focusing solely on ranks of single genes, by means of GSEA we assessed the role of the whole gene set associated with depressive disorders. Regarding emotion processing, risk genes were predominantly enriched within positively correlated genes (maximum enrichment score, ES: 2.881, $p = 0.43$) in single-site data, albeit not reaching statistical significance. In contrast, an enrichment of risk genes was present within negative correlations for reward processing (maximum ES: -0.2, $p = 0.527$). Within the Neurosynth dataset, the maximum subcortical ES yielded 0.275 ($p = 0.051$) for emotion and -0.243 for reward processing ($p = 0.65$), respectively (Fig. 4c). In the cortex, GSEA yielded identical enrichment of MDD risk genes compared to the subcortical analyses (Supplementary Fig. 6).

Discussion

Here, we applied a comprehensive and integrative methodological approach to investigate the relationship between regional gene expression patterns and macroscopic BOLD responses elicited by emotional face recognition and the acceptance of monetary rewards, under the assumption that strongly correlated genes would coincide with distinct biological programs and genes implicated in depressive disorders. Large-scale screening for spatial associations between mRNA expression and functional brain activation resulted in ranked lists of 18,686 genes positively and negatively correlated with BOLD signaling in healthy male and female subjects. Similar distributions of Spearman's correlation coefficients were present for emotion (ranging from $\rho = -0.739$ to $\rho = 0.865$ in subcortical and from $\rho = -0.449$ to $\rho = 0.431$ in cortical areas) and reward processing (from $\rho = -0.788$ to $\rho = 0.81$ in the subcortex and from $\rho = -0.639$ to $\rho = 0.698$ in the cortex). Considering strongest brain activation elicited by emotion processing in ventral striatum, amygdala, ventral tegmental area, fusiform gyri, insula and medial prefrontal cortex, in that order, it seems plausible that higher correlation levels were observed in subcortical regions, compared to the cortex. Similarly, subcortical activation was more prominent in the reward paradigm. Exploring the GO knowledgebase, we detected task-specific processes related to cellular transport as well as neuronal development, synapse regulation, and transcription for emotion and reward processing, respectively. Notably, associated ontologies were interrelated solely within each fMRI paradigm, thus indicating unique biological programs for both investigated RDoC domains. Focusing on systems for social processes and positive valence systems, a meta-analytic validation sample was obtained from the Neurosynth framework, comprising activation maps from over 90 fMRI studies. Based on correlations between imaging parameters and gene expression, we identified master regulators associated with depressive disorders and task-specific functional brain activation, TCF4 and MEF2C, in two independent datasets. While MEF2C showed a congruent regulation with negative associations related to emotion and reward processing,

TCF4 appeared simultaneously as a regulator for genes negatively correlated with the emotion task, but positively correlated with reward processing. Albeit not reaching statistical significance, the supplementary GWAS enrichment analysis suggested an inversed distribution of risk genes implicated in major depression, showing a rather positive association with imaging data for emotional face recognition and a negative association for the acceptance of monetary rewards.

Although the role of the transcriptome has been reviewed for several fMRI measures^{4,5}, individual genetic influences on emotional face recognition or adaptive reward-based decision-making have only been evaluated for the presence of single gene variants of functional proteins (i.e. CREB1), irrespective of topological distribution¹³. Particularly in MDD, additive genetic effects may attribute to individual differences in the phenotype and highlight the importance of large-scale data in systems medicine to resolve unsettled genetic influences on neuroimaging paradigms²¹. While over 322 million people worldwide suffer from depressive disorders, a number that increased by 18.4 % between 2005 and 2015¹⁴, a significant part of the population is also affected by subthreshold depressive symptoms, potentially originating from different levels of genetic susceptibility in relevant neuronal systems. Our findings strongly support the idea of a dimensional genetic background affecting brain regions involved in emotion and reward processing that continuously progresses from physiological to pathological states, recently highlighted within the much-noticed RDoC framework¹⁸. In line with the debilitating symptoms of depressive disorders, we investigated paradigms reflecting principal functions of both the reward responsiveness construct within the positive valence systems domain as well as the social communication construct, which is part of the systems for social processes domain. Hyper- as well as hypoactivations of brain regions involved in the integration of social information and a reduced reward sensitivity together with decision-making efficiency suggest a polygenic nature of depressive disorders

with distinct imaging features²²⁻²⁴. However, identifying the core set of risk genes is complicated due to widespread and disease-specific network interactions as well as modulatory MRs.

In this study, regulating genes associated with MDD included protein coding MEF2C (Myocyte Enhancer Factor 2C), which plays a role in neuronal development, as well as hippocampus-dependent learning and memory^{25,26}. Relevance for synapse regulation arises due to alternatively spliced transcript variants involved in neuronal processes, e.g. activated TLR4 signaling or the cAMP response element-binding protein (CREB) pathway. Besides depressive disorders, associations of this transcription activator with other psychiatric disorders like schizophrenia²⁷ or attention-deficit/hyperactivity disorder²⁸ have been reported. Most notably, significant dependence of striatal neuronal activation was described for an identified risk variant in the TMEM161B-MEF2C gene cluster during a reward task, endorsing well-known deficits of reward processing in MDD, observed as anhedonia²⁹. Besides, we affirm previously reported relationships between neuropsychiatric disorders and mutations of TCF4, which has been implicated not only in depression, but also in schizophrenia and autism^{30,31}. The transcription factor is mainly characterized by its regulatory role for the proliferation and differentiation of neuronal and glial progenitor cells³². Interestingly, the master regulatory role of TCF4 in schizophrenia was recently endorsed by Torshizi et al., based on the analysis of transcriptional networks in two independent datasets³³. Since gene variants exhibit their effects by uncountable molecular mechanisms, a closer investigation of genes strongly associated with imaging parameters, as well as the MRs reported in this study, will prospectively allow further statements about regional protein biosynthesis and allocation of resulting proteins to cellular compartments. For example, DNA-binding transcription factor FOXN4 (Forkhead Box N4), belonging to the Forkhead-box (FOX) superfamily, showed highest correlation with the emotional face recognition

paradigm in the cortex. FOX transcription factors are involved in regulatory biological processes and mutations in forkhead genes have been linked to developmental disorders in humans due to substitutions or frameshifts that disable or remove the DNA binding domain³⁴. The subtype FOXN4 thereby expresses developmental functions in neural and non-neural tissues, particularly during spinal neurogenesis by modulating a specific expression mosaic of other proneural factors³⁵. Further relevance for neural development was shown by Chen et al.³⁶, who demonstrated location of FOXN4 on neurons and astrocytes, as well as an increased expression after spinal cord injury lesions. Although associations with depressive or other neuropsychiatric disorders have not been published, the modulatory role of FOXN4 as a key transcriptional regulator during developmental processes demands further research, especially since the full set of its targets in the CNS are not known yet. Accordingly, the C10orf125 gene (Fucose Mutarotase, FUOM), expressed in the brain and other tissues, showed highest correlation with emotional face recognition in the subcortex. The corresponding gene transcript, fucose mutarotase, is an enzyme of the fucose-utilization pathway performing the interconversion between α -L-fucose and β -L-fucose on human cell surfaces. Hereof, besides one animal study demonstrating male-like sexual behavior in FUOM knock-out mice, presumably resulting from reduced fucosylation during neurodevelopment³⁷, further associations with pathological states in mammals have not been published for this gene. Regarding reward processing, the protein coding DUSP3 (Dual-specificity phosphatase 3) gene, member of the dual-specificity protein phosphatase subfamily, showed strongest correlation with measured fMRI data in the cortex. Members of these protein tyrosine phosphatases (PTPs) regulate the phosphorylation of the mitogen-activated protein (MAP) kinase signaling pathway and control cell signaling, especially in regard to cytoskeleton reorganization, apoptosis and RNA metabolism³⁸. DUSP3 shows a wide expression in different tissues as an opposing factor of protein tyrosine kinases (PTKs) and acts as a central mediator of cellular proliferation and differentiation. Whereas a role in neoplastic disorders,

pathologies related to immunology, angiogenesis as well as Parkinson's disease (PD) have been related to anomalous tyrosine phosphorylation^{39,40}, associations with psychiatric disorders have not been described. In subcortical regions, MDK (midkine) was among the highest correlating genes for the reward paradigm. MDK transcribes one of two growth factors from the heparin-binding cytokine family and plays a role during the differentiation of neurons⁴¹. An important role of this gene has been suggested within dopaminergic pathways, particularly facilitating neuroprotective effects in neurodegenerative disorders, drug-induced neurotoxicity in the striatum, or after neural injury⁴². Further, disease-related publications suggest accumulation of midkine in senile plaques and increased serum levels in patients with Alzheimer's disease⁴³, genetic variations associated with PD⁴⁴, and an influence on addictive behaviors⁴⁵. A possible therapeutic target of MDK was proposed for autoimmune diseases including multiple sclerosis⁴⁶, and more recently, associations of elevated serum levels of the neurotrophic factor have been found in autism spectrum disorder⁴⁷.

Despite advances and decreased costs of high-throughput gene expression profiling the necessity for large cohorts in genetic studies call for collaborations and integrative approaches. By including whole-brain expression patterns, our approach can be discriminated from previous neuroimaging studies solely investigating effects of individual disease-related SNPs or environmental factors. Still, a range of influencing factors hampers interpretation of spatial associations between gene expression and functional imaging data, like outdated annotation information or inaccurate sample assignment. Further, ubiquitous noise due to expression of genes with a low spatial dependence occurs, which has also been addressed by Gryglewski et al.⁴⁸. While mRNA levels don't necessarily reflect actual in vivo protein densities, signal alterations in fMRI studies are also impaired by confounding variables, such as network-architectures, structural and functional connectivity measures, or non-specific brain activation. To minimize BOLD signaling elicited by superimposed executive functions,

control conditions were implemented for each task, facilitating specificity of performed correlations for both paradigms. Noteworthy, meta-analytic brain activation maps eventually confirmed findings from single-site fMRI measurements. In contrast to a previous study¹⁰ that identified gene-cognition associations based on the Neurosynth framework, we increased spatial resolution and advanced probe selection of gene expression data by application of interpolated mRNA maps with continuous expression estimates throughout the whole brain. Surely, results of this study are limited due to the structure of applied data sets and reflect rather subtle genetic influences on psychological processes, disregarding the dynamic nature of short-term regulatory mechanisms, environmental factors or individual variations due to genetic ancestry. Inter-individual differences regarding effects of sex, age or genotype cannot be considered, when performing integrative analyses on the basis of the AHBA that originally derived expression values from only 6 post-mortem brains^{49,50}. Since female gene expression was acquired from solely one donor, sex-specific differences had to be neglected in this study. Predictions of in vivo gene expression are further limited due to locally and functionally regulated epigenetic and epitranscriptomic modifications that affect actual protein distribution⁵¹. Despite unbalanced sex distribution of the donor brains, the AHBA still offers the most comprehensive database for the investigation of human whole-brain gene expression, comprising nearly 20,000 genes. Within the framework of future studies, the vast potential of the AHBA might even be increased by re-assigning available mRNA probes to corresponding genes on the basis of the latest sequencing information to increase the number of specifically annotated genes. Additionally, harmonized data processing pipelines or methodological guidelines, instead of rather unique approaches to data integration and corresponding statistical measures, could increase comparability between studies⁵²⁻⁵⁴. Although a whole-brain proteome atlas including relevant genotypes would reflect actual protein expression more reliably, such database has not been published yet.

Since explanations of altered fMRI activations often neglect the spatial distribution of investigated genetic substrates, evaluating the role of gene expression in neuropsychiatric disorders requires an integration of high-resolution neural information from molecular sources for more precise interventions in the future. Our analyses highlight the advantages of a comprehensive approach to reveal genetic influences on functional brain imaging by integrating multimodal imaging and large-scale transcriptome data with sufficient power. While insignificant results of the performed GSEA are partly affected by the small number of risk genes implicated in depression, low enrichment of the risk gene set might also be caused by a more prominent role of individual MRs that modulate up- and downregulation of subordinate genes, including risk and non-risk genes. In general, the GWAS performed by Wray and colleagues¹⁶ is among the largest ever conducted in psychiatric genetics and provides a solid basis for further research about the genetic architecture of MDD. In this study, we performed an enrichment analysis with this GWAS data to explore imaging paradigms that specifically integrate core depressive symptoms in line with current RDoC, like dysfunction in social interaction and inability to properly experience reward.

While traditional imaging genomics strategies have solely investigated the effects of individual genotype variations, integration of whole-brain gene expression patterns provide superior information for the understanding of neuropsychiatric disorders. Here, biological programs related to cellular transport, neuronal development, synapse regulation, and transcription processes were specifically associated with the emotion and reward system, highlighting the role of topological gene expression for imaging paradigms. Further, identification of regulatory genes TCF4 and MEF2C endorses the investigation of commonly altered brain activation during emotional face recognition and the acceptance of monetary rewards in depressed individuals. Overall, our work exemplifies an integrative approach

including complementary information from multiscale data, which seems to be increasingly relevant in the big data era.

Materials and Methods

Participants

Healthy subjects were recruited from the university environment and gave written informed consent to study procedures previously approved by the Ethics Committee of the University Medical Center Göttingen. Included participants were aged between 22 and 52 years ($M = 40.5$, $SD = 14.37$), of Caucasian European ethnicity and were fluent in German language. Exclusion criteria comprised contraindications to MRI, past or present psychiatric, neurological, or medical disorders, consumption of psychotropic drugs as well as positive family history of psychiatric disorders. In total, a number of 26 men and 22 women completed two fMRI paradigms including reward processing and emotional face recognition. Excessive movement in any of the three translation (> 2 mm) or rotation ($> 2^\circ$) planes resulted in exclusion of 4 participants.

Functional brain imaging

Functional imaging data was acquired using a 3 T scanner (Siemens Magnetom TRIO, Siemens Healthcare, Erlangen, Germany) and a 32-channel head coil with a $2 \times 2 \times 2$ mm voxel size, TR 2500 ms, TE 33 ms, 70° flip angle, 10 % distance factor, FOV 256 mm and 60 slices with multiband factor of 3 for the acquisition of T2*-weighted images. Imaging data analysis was performed using Statistical Parameter Mapping (SPM12; Wellcome Department of Imaging Neuroscience, Institute of Neurology, London, UK) and Matlab R2015b (The Mathworks Inc., Natick, MA, USA). First, echo planar imaging (EPI) images were standardly preprocessed with slice time correction, realignment, and normalization into the MNI space, as well as smoothing with an $8 \times 8 \times 8$ mm FWHM Gaussian kernel.

Two different fMRI paradigms were analyzed for the same group of participants during two states. First, specific activation maps reflected brain activation during performance of the tasks. In contrast, corresponding control conditions represented non-specific hemodynamic

activity inherent to any task performance during fMRI measurement, caused by unspecific physiological activation, e.g. related to visual, auditory, attentional or motor functions.

Estimates of neural activity were initially computed with a general linear model (GLM) for each subject individually (first-level analysis) with nuisance movement parameters regressed as covariates-of-no-interest. Later, experimental and control conditions were evaluated at group level (second-level analysis) and resulting activation maps, which represented task-specific brain activation, were used for further analyses.

Emotional face recognition

The paradigm of implicit emotional face recognition contains two different contexts: human faces and geometric objects. Pictures of males and females with varying negative face expressions obtained from the Radboud database⁵⁵ were presented for 17 s, during which participants responded to the gender of the presented person with a button press. Thereby, perception of emotions was rather implicit, which has been shown to enhance the activation of emotional correlates⁵⁶. For the control condition, participants were instructed to respond analogously to the shape of an object, either an ellipse or a rectangle, positioned in the face area and made from scrambling original face trials. All trials were controlled for brightness, contrast and presented in a very similar composition. The activation patterns representing the experimental and control conditions were computed using first-level (single-subject) contrasts of the trials from emotional faces and object blocks, respectively. Resulting data were then used for second-level (group) analysis, as standardly performed for random effects model.

Reward processing

For this study, a previously established fMRI paradigm was implemented, which has been broadly used to investigate physiological and pathological reward mechanisms^{57,58}. Briefly, participants performed a modified delayed match to sample task, including two contexts

involving previously conditioned stimuli to monetary rewards: acceptance or rejection of rewards, i.e. pressing a button when squares are shown. Subjects were instructed they would receive 30 € for their participation, and that they were able to double this amount according to their task performance. As control trials, subjects responded with a button press to stimuli that required motor performance as well as attentional and memory resources, but were not conditioned to monetary reward. To compute the control condition, first-level (single-subject) contrasts of correctly matched sample trials within the same experimental block of reward trials were used. For experimental conditions, first-level experimental contrasts were calculated from brain activation elicited during acceptance of previously conditioned stimuli. At group-level, activations related to experimental and control trials were contrasted to obtain functional activation patterns.

Meta-analytic functional brain activation

Besides fMRI data obtained from participants performing two different tasks at our institution, we evaluated large-scale meta-analytical imaging data from the Neurosynth platform (<https://neurosynth.org/>), which provides probabilistic brain activation maps computed from an automated meta-analysis based on published fMRI studies. This online database combines text-mining and machine-learning techniques to generate statistical inference maps of currently 1,335 imaging terms from 14,371 fMRI studies including male and female participants²⁰. Within the framework of the Neurosynth database, particular psychological processes are labeled with terms of interest and represented by uniformity test maps. For this study, activation maps related to emotion (“fearful faces”) and reward (“rewards”) processing were downloaded in MNI152 2 mm space resolution to validate fMRI data obtained at our institution. Each uniformity test map provides z-scores, which reflect the proportion of studies that report cerebral activation at a given voxel. Thus, obtained

Neurosynth maps depicted specific brain areas that were consistently reported in studies investigating fMRI activation for emotion and reward processing, respectively.

In a first step, activation maps, with z-scores representing how brain regions are related to the chosen term, were generated from the Neurosynth online database. Since the online user interface only provides thresholded maps, we used the Neurosynth toolbox for python to create unthresholded maps that were further smoothed using 8 x 8 x 8 mm FWHM to match the kernel size of single-site data. Assessment of conformity between meta-analytic data and measured fMRI maps was performed qualitatively and quantitatively. Thereby, region-wise Spearman's correlation coefficients were calculated in cortical and subcortical regions, whereby the Neurosynth terms "fearful faces" and "rewards" matched well with functional activation related to recognition of negative faces and the acceptance of monetary rewards, respectively. Due to low fMRI activity, cerebellum was not considered for the comparison. In contrast to single-site fMRI data, meta-analytical information comprises rather positive values, due to the sparse reporting of brain regions showing negative activation in most neuroimaging studies. Hence, when processing unthresholded data from the uniformity test maps, mainly positive values determined the association analysis between single-site and Neurosynth data.

Whole-brain gene expression

The AHBA (www.brain-map.org) consists of microarray assessments from 3,702 brain tissue samples collected across 6 human donors (1 female, mean age = 42.5, SD = 13.4) derived from diverse regions of the brain, extensively described in the original publications^{49,50}. As delineated by other authors⁵⁴, multiple data processing steps such as gene annotation, probe selection, or sample assignment need to be considered to facilitate subsequent correlation analyses between gene expression and neuroimaging data. Also, by using common parcellation schemes expression levels in numerous brain regions devoid of tissue samples

remain indefinite, thus demanding the generation of whole brain transcriptome maps. To sufficiently meet proposed methodological requirements, gene expression data (\log_2 -values) from the AHBA were obtained according to by Gryglewski et al.⁴⁸ to compensate for inter-individual differences between mRNA probes and donor brains as well as sparse anatomical sampling, attributable to high efforts in tissue preparation and processing. Sex-specific differences were not considered due to low validity of female transcriptome data within the AHBA. Using Gaussian process regression, interpolated transcriptome maps comprising expression of 18,686 genes associated with Entrez Gene IDs at all cortical and subcortical structures were created (predicted transcriptome maps are available for download at www.meduniwien.ac.at/neuroimaging/mRNA.html). Nonetheless, for a certain number of remaining genes transcriptome maps have not been predicted sufficiently due to presence of insensitive probes or missing allocations to gene IDs.

Spatial correlation between gene expression and brain activity

Initially, gene lists ranking correlations between transcriptome maps and functional brain activation patterns elicited by selected paradigms were compiled, whereby the ranking of each gene depended on its correlation strength with the corresponding single-site BOLD activation map. After initial inspection, the cerebellum was excluded from further analysis, due to marginal activation during both fMRI paradigms. On the basis of known differences in gene expression between broad anatomical regions^{49,59} correlation analyses were assessed within cortical and subcortical regions separately (Supplementary Fig. 7). To conduct association analyses, all available transcriptome maps were aligned with group-averaged fMRI activation maps in MNI space for each paradigm. To account for partly non-symmetrical distribution of mRNA data and existence of outliers, Spearman's correlation coefficients were calculated between each gene-imaging pair (mRNA expression vs. fMRI activation). Main findings are reported for region-wise analyses, along with additional results for voxel-wise correlations

(total number of voxels was 129,817 in the cortex and 10,863 in subcortex; zero-values outside of the investigated area were excluded). Thereby, each statistical map was segmented into brain regions according to the Brainnetome atlas, which was selected for primary analyses, because it labels a sufficient number of subcortical ($n = 36$) and cortical ($n = 210$) regions-of-interest (ROIs)⁶⁰. A complementary analysis with fewer brain regions (12 subcortical and 78 cortical ROIs) was done using the automated anatomical labeling (AAL) brain atlas⁶¹, in order to evaluate influences of different parcellation methods. Both atlases were aligned with fMRI maps as well as transcriptome maps in MNI space using SPM12, while extraction of ROIs and correlation analyses were performed in MATLAB2018a (www.mathworks.com). Statistical significances of region-wise correlations were assessed by means of randomization tests, including 10,000 iterations. For each sampled permutation, mRNA values were randomly shuffled and correlated with fMRI data. Two-sided p-values were calculated as the proportion of sampled permutations where the absolute value was greater than the true correlation coefficient (non-shuffled data).

Identification of overlap between analyzed data set

To compare various sets of mRNA-fMRI correlations we used Rank-Rank Hypergeometric Overlap (RRHO) package (version 1.26.0) in R (<https://www.bioconductor.org/packages/release/bioc/html/RRHO.html>), which allows statistical testing of the extent of overlap between two ranked lists. RRHO determines the degree of differential expression observed in profiling experiments using the hypergeometric distribution. While originally applied for the comparison of gene expression profiles between different microarray platforms or types of model system, we used RRHO to compare genes that ranked according to relevant measures of differential information, in this case the correlation strength with fMRI data. Thereby, genes of two datasets were ranked according to their names and corresponding ranks tested for statistical overlap. We provide both a

graphical representation of the characteristics of analyzed datasets (corresponding p-values) as well as a statistical measure of overlap (ρ_{RRHO}). A high overlap implies that positively correlating genes of a given list show high ranks in the second list, while genes with a negative correlation are also negatively associated in the alternative list. Applying this method offered the advantage of using the whole continuum of previously correlated genes without the need to truncate the list by pre-defined thresholds for each combination of used datasets.

Analysis of biological processes

Making use of the GO knowledgebase^{19,62}, a comprehensive resource for computational analysis of large-scale data, we explored enriched biological processes that included genes with expression patterns highly correlated with each fMRI paradigm. Cytoscape plugin “ClueGO”⁶³ was used to compute GO enrichment (default parameters), comprising all listed terms at the time of analysis (18,361 biological process terms). Only findings of specialized biological processes (GO levels higher than 4) were reported, to attain more conclusive information about underlying genetic substrates and their functions within investigated emotion and reward systems. Compiled ranked lists originating from region-wise analyses were included for the investigation of gene ontology, due to more refined results. For each paradigm, the GO analysis was performed separately for positively and negatively correlated genes with cut-off values for correlation coefficients above $\rho = 0.5$ and below $\rho = -0.5$, which excluded non-spatially depending genes showing insufficient associations with fMRI data. Significance level was set to a p-value minimum of 0.05, after applying Bonferroni correction to the enriched GO terms. Considering potential overlaps across imaging paradigms, interrelations between all significant biological processes found in the GO analysis were quantified by means of Spearman’s correlation coefficients.

Association of risk genes implicated in major depression with functional imaging data

Topological expression patterns of genes implicated in major depression and their association with neuroimaging properties have not yet been investigated. Regarding genetic risks for depressive disorders, recently 44 genetic risk loci for major depression were identified in a genome-wide association meta-analysis by Wray et al.¹⁶, which included 135,458 cases and 344,901 controls. Based on this meta-analysis, we investigated if the MDD risk gene set contained MRs of previously compiled genes showing strong correlations between gene expression and imaging data. Thereby, the role of all 42 functional risk genes proposed by Wray et al. was evaluated for each fMRI paradigm (Supplementary Table 4).

By using the cytoscape plugin iRegulon⁶⁴, we performed master regulator analysis separately for all positive and negative mRNA-fMRI correlations above $\rho = 0.6$ and below $\rho = -0.6$, respectively. Rather low thresholds were set to ensure a sufficient number of evaluated genes in subcortical regions. In the cortex, insufficient data availability hampered analyses of co-regulatory networks, due to generally lower correlation coefficients of mRNA-fMRI associations compared to the subcortex. Parameters, such as enrichment score and significances, were used as default; distance from TSS was set to 500bp. We compared the master regulators on each predicted regulon with the genes associated with risk for major depression.

Additionally, potential enrichment of GWAS risk genes implicated in major depression was assessed by means of GSEA^{65,66} for each fMRI paradigm. Based on initially compiled gene lists that were ranked by correlation strengths with fMRI data, the position of each MDD risk gene was compared to the position of all other genes. Thereby, risk genes clustered either within positive or negative correlations with imaging data, depending on the investigated paradigm. In terms of the methodology, the GSEA enrichment score reflects the degree to which the analyzed MDD risk gene set is overrepresented within each ranked list. We tested, whether MDD risk genes were randomly distributed throughout each ranked list or primarily

found at the top (showing positive correlations with fMRI data) and bottom (showing negative correlations). We used the GSEA implementation in R available in the package `clusterProfiler`⁶⁷. Ranked correlation coefficients indicating the relationship between imaging and transcriptome data and corresponding p-values were defined as input factors for the enrichment analysis, which required a summarized biological value for each included gene. Genes ranked in higher positions contributed more to resulting enrichment p-values than lower ranked genes. Corresponding to a weighted Kolmogorov-Smirnov-like statistic, the ES was calculated by a stepwise increase or decrease of the total sum statistic of the ranked list, depending on the ranking of MDD risk genes, as described by Subramanian et al.⁶⁵. The maximum ES (with positive or negative values), representing the maximum deviation from zero, thereby indicated enrichment of positive or negative correlations, respectively.

References

- 1 Medland, S. E., Jahanshad, N., Neale, B. M. & Thompson, P. M. Whole-genome analyses of whole-brain data: working within an expanded search space. *Nature Neuroscience* **17**, 791-800 (2014).
- 2 Kitchen, R. R., Rozowsky, J. S., Gerstein, M. B., & Nairn, A. C. Decoding neuroproteomics: integrating the genome, transcriptome and functional anatomy. *Nature neuroscience* **17**, 1491-1499 (2014).
- 3 Komorowski, A. et al. Association of Protein Distribution and Gene Expression Revealed by PET and Post-Mortem Quantification in the Serotonergic System of the Human Brain. *Cerebral Cortex* **27**, 117-130 (2017).
- 4 Richiardi, J. et al. Correlated gene expression supports synchronous activity in brain networks. *Science* **348**, 1241-1244 (2015).
- 5 Shin, J. et al. Cell-specific gene-expression profiles and cortical thickness in the human brain. *Cerebral Cortex* **28**, 3267-3277 (2018).
- 6 Rizzo, G., Veronese, M., Expert, P., Turkheimer, F. E. & Bertoldo, A. MENGA: A New Comprehensive Tool for the Integration of Neuroimaging Data and the Allen Human Brain Transcriptome Atlas. *PLoS One* **11** (2016).
- 7 Unterholzner, J. et al. Topologically Guided Prioritization of Candidate Gene Transcripts Coexpressed with the 5-HT1A Receptor by Combining In Vivo PET and Allen Human Brain Atlas Data. *Cerebral Cortex* (2020).
- 8 Tulay, E. E., Metin, B., Tarhan, N. & Arkan, M. K. Multimodal Neuroimaging: Basic Concepts and Classification of Neuropsychiatric Diseases. *Clinical EEG and neuroscience* **50**, 20-33 (2018).
- 9 Jones, T. & Rabiner, E. A. The development, past achievements, and future directions of brain PET. *Journal of Cerebral Blood Flow & Metabolism* **32**, 1426-1454 (2012).
- 10 Fox, A. S., Chang, L. J., Gorgolewski, K. J. & Yarkoni, T. Bridging psychology and genetics using large-scale spatial analysis of neuroimaging and neurogenetic data. *bioRxiv* **012310** (2014).
- 11 Romme, I. A., de Reus, M. A., Ophoff, R. A., Kahn, R. S. & van den Heuvel, M. P. Connectome Disconnectivity and Cortical Gene Expression in Patients With Schizophrenia. *Biological psychiatry* **81**, 495-502 (2017).
- 12 Freeze, B., Acosta, D., Pandya, S., Zhao, Y. & Raj, A. Regional expression of genes mediating trans-synaptic alpha-synuclein transfer predicts regional atrophy in Parkinson disease. *Neuroimage Clin* **18**, 456-466 (2018).
- 13 Wolf, C. et al. CREB1 genotype modulates adaptive reward-based decisions in humans. *Cerebral Cortex* **26**, 2970-2981 (2015).
- 14 James, S. L. et al. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* **392**, 1789-1858 (2018).
- 15 Lewinsohn, P. M., Solomon, A., Seeley, J. R. & Zeiss, A. Clinical implications of "subthreshold" depressive symptoms. *Journal of Abnormal Psychology* **109**, 345-351 (2000).
- 16 Wray, N. R. et al. Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nature genetics* **50**, 668-681 (2018).
- 17 Foland-Ross, L. C. & Gotlib, I. H. Cognitive and neural aspects of information processing in major depressive disorder: an integrative perspective. *Front Psychol* **3**, 489 (2012).

- 18 Sanislow, C. A., Ferrante, M., Pacheco, J., Rudorfer, M. V. & Morris, S. E. Advancing translational research using NIMH research domain criteria and computational methods. *Neuron* **101**, 779-782 (2019).
- 19 The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing strong. *Nucleic acids research* **47**, D330-D338 (2018).
- 20 Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. Large-scale automated synthesis of human functional neuroimaging data. *Nat Methods* **8**, 665-670 (2011).
- 21 Fabbri, C. et al. Consensus paper of the WFSBP Task Force on Genetics: Genetics, epigenetics and gene expression markers of major depressive disorder and antidepressant response. *World J Biol Psychiatry* **18**, 5-28 (2017).
- 22 Knutson, B., Bhanji, J. P., Cooney, R. E., Atlas, L. Y. & Gotlib, I. H. Neural responses to monetary incentives in major depression. *Biological psychiatry* **63**, 686-692 (2008).
- 23 Ghaemi, S. N. & Vohringer, P. A. The heterogeneity of depression: an old debate renewed. *Acta psychiatrica Scandinavica* **124**, 497 (2011).
- 24 Luijten, M., Schellekens, A. F., Kühn, S., Machielse, M. W. & Sescousse, G. Disruption of reward processing in addiction: an image-based meta-analysis of functional magnetic resonance imaging studies. *JAMA Psychiatry* **74**, 387-398 (2017).
- 25 Paciorkowski, A. R. et al. MEF2C Haploinsufficiency features consistent hyperkinesia, variable epilepsy, and has a role in dorsal and ventral neuronal developmental pathways. *Neurogenetics* **14**, 99-111 (2013).
- 26 Barbosa, A. C. et al. MEF2C, a transcription factor that facilitates learning and memory by negative regulation of synapse numbers and function. *Proc. Natl Acad. Sci. USA* **105**, 9391-9396 (2008).
- 27 Mitchell, A. C. et al. MEF2C transcription factor is associated with the genetic and epigenetic risk architecture of schizophrenia and improves cognition in mice. *Molecular psychiatry* **23**, 123-132 (2017).
- 28 Shadrin, A. A. et al. Novel loci associated with attention-deficit/hyperactivity disorder are revealed by leveraging polygenic overlap with educational attainment. *J Am Acad Child Adolesc Psychiatry* **57**, 86-95 (2018).
- 29 Muench, C. et al. The major depressive disorder GWAS-supported variant rs10514299 in TMEM161B-MEF2C predicts putamen activation during reward processing in alcohol dependence. *Translational psychiatry* **8**, 1-10 (2018).
- 30 Li, H. et al. Disruption of TCF4 regulatory networks leads to abnormal cortical development and mental disabilities. *Molecular psychiatry* **24**, 1235-1246 (2019).
- 31 Amare, A. T. et al. Bivariate genome-wide association analyses of the broad depression phenotype combined with major depressive disorder, bipolar disorder or schizophrenia reveal eight novel genetic loci for depression. *Molecular psychiatry* (2019).
- 32 Ross, S. E., Greenberg, M. E. & Stiles, C. D. Basic helix-loop-helix factors in cortical development. *Neuron* **39**, 13-25 (2003).
- 33 Torshizi, A. D. et al. Deconvolution of transcriptional networks identifies TCF4 as a master regulator in schizophrenia. *Science Advances* **5** (2019).
- 34 Carlsson, P. & Mahlapuu, M. Forkhead transcription factors: Key players in development and metabolism. *Developmental Biology* **250**, 1-23 (2002).
- 35 Misra, K., Luo, H., Li, S., Matise, M. & Xiang, M. Asymmetric activation of Dll4-Notch signaling by Foxn4 and proneural factors activates BMP/TGF β signaling to specify V2b interneurons in the spinal cord. *Development* **141**, 187-198 (2014).
- 36 Chen, X., Yao, Y., Guan, J., Chen, X. & Zhang, F. Up-regulation of FoxN4 expression in adult spinal cord after injury. *Journal Of Molecular Neuroscience* **52**, 403-409 (2014).

37. Park, D., Choi, D., Lee, J., Lim, D. S. & Park, C. Male-like sexual behavior of female mouse lacking fucose mutarotase. *BMC Genet* **11**, 62 (2010).
38. Tonks, N. K. Protein tyrosine phosphatases - from housekeeping enzymes to master regulators of signal transduction. *The FEBS Journal* **280**, 346-378 (2013).
39. Russo, L. C., Farias, J. O., Ferruzo, P. Y. M., Monteiro, L. F. & Forti, F. L. Revisiting the roles of VHR/DUSP3 phosphatase in human diseases. *Clinics* **73** (2018).
40. Cohen, P. & Alessi, D. R. Kinase drug discovery—what’s next in the field?. *ACS chemical biology* **8**, 96-104 (2013).
41. F Alguacil, L. & Herradón, G. Midkine and pleiotrophin in the treatment of neurodegenerative diseases and drug addiction. *Recent patents on CNS drug discovery* **10**, 28-33 (2015).
42. Yoshida, Y., Sakakima, H., Matsuda, F. & Ikutomo, M. Midkine in repair of the injured nervous system. *British journal of pharmacology* **171**, 924-930 (2014).
43. Salama, R. H. et al. Increased midkine levels in sera from patients with Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry* **29**, 611-616 (2005).
44. Prediger, R. D. et al. Mice with genetic deletion of the heparin-binding growth factor midkine exhibit early preclinical features of Parkinson's disease. *Journal of neural transmission* **118**, 1215-1225 (2011).
45. Gramage, E. et al. Regulation of extinction of cocaine-induced place preference by midkine is related to a differential phosphorylation of peroxiredoxin 6 in dorsal striatum. *Behavioural brain research* **253**, 223-231 (2013).
46. Takeuchi, H. Midkine and multiple sclerosis. *British journal of pharmacology* **171**, 931-935 (2014).
47. Esnafoglu, E. & Cirrik, S. Increased serum midkine levels in autism spectrum disorder patients. *International Journal of Neuroscience* **128**, 677-681 (2018).
48. Gryglewski, G. et al. Spatial analysis and high resolution mapping of the human whole-brain transcriptome for integrative analysis in neuroimaging. *NeuroImage* **176**, 259-267 (2018).
49. Hawrylycz, M. J. et al. An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature* **489**, 391-399 (2012).
50. Shen, E. H., Overly, C. C. & Jones, A. R. The Allen Human Brain Atlas: comprehensive gene expression mapping of the human brain. *Trends in neurosciences* **35**, 711-714 (2012).
51. Maier, T., Güell, M. & Serrano, L. Correlation of mRNA and protein in complex biological samples. *FEBS letters* **583**, 3966-3973 (2009).
52. Kim, M. S. et al. A draft map of the human proteome. *Nature* **509**, 575-581 (2014).
53. Keil, J. M., Qalieh, A. & Kwan, K. Y. Brain Transcriptome Databases: A User's Guide. *Journal of Neuroscience* **38**, 2399-2412 (2018).
54. Arnatkevičiūtė, A., Fulcher, B. D. & Fornito, A. A practical guide to linking brain-wide gene expression and neuroimaging data. *NeuroImage* **189**, 353-367 (2019).
55. Langner, O. et al. Presentation and validation of the Radboud Faces Database. *Cognition & Emotion* **24**, 1377-1388 (2010).
56. Keightley, M. L. et al. An fMRI study investigating cognitive modulation of brain regions associated with emotional processing of visual stimuli. *Neuropsychologia* **41**, 585-596 (2003).
57. Goya-Maldonado, R. et al. Dissociating pathomechanisms of depression with fMRI: bottom-up or top-down dysfunctions of the reward system. *Eur Arch Psychiatry Clin Neurosci* **265**, 57-66 (2015).
58. Diekhof, E. K. & Gruber, O. When Desire Collides with Reason: Functional Interactions between Anteroventral Prefrontal Cortex and Nucleus Accumbens

- Underlie the Human Ability to Resist Impulsive Desires. *Journal of Neuroscience* **30**, 1488-1493 (2010).
- 59 Chen, L. et al. Analysis of Gene Expression Profiles in the Human Brain Stem, Cerebellum and Cerebral Cortex. *PLoS One* **11** (2016).
- 60 Fan, L. et al. The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture. *Cerebral Cortex* **26**, 3508-3526 (2016).
- 61 Tzourio-Mazoyer, N. et al. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage* **15**, 273-289 (2002).
- 62 Ashburner, M. et al. Gene Ontology: tool for the unification of biology. *Nature genetics* **25**, 25-29 (2000).
- 63 Bindea, G. et al. ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* **25**, 1091-1093 (2009).
- 64 Janky, R. et al. iRegulon: From a Gene List to a Gene Regulatory Network Using Large Motif and Track Collections. *PLoS computational biology* **10** (2014).
- 65 Subramanian, A. et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl Acad. Sci. USA* **102**, 15545-15550 (2005).
- 66 Mootha, V. K. et al. PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nature genetics* **34**, 267-273 (2003).
- 67 Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics: a journal of integrative biology* **16**, 284-287 (2012).

Acknowledgements

We thank the students of the Laboratory of Systems Neuroscience and Imaging in Psychiatry (SNIPLab) and the Neuroimaging Labs (NIL) for support. Parts of this study have been presented by AK at the 32nd ECNP Congress, 07-10 September 2019, Copenhagen, Denmark. With relevance to this work there is no conflict of interest to declare. AS and RGM are funded by the German Federal Ministry of Education and Research (Bundesministerium fuer Bildung und Forschung, BMBF: 01 ZX 1507, “PreNeSt - e:Med”). MM is funded by the Austrian Science Fund FWF DOC 33-B27. GG was recipient of a DOC-fellowship of the Austrian Academy of Sciences at the Department of Psychiatry and Psychotherapy, MUV. JW is supported by an Ilídio Pinho professorship, iBiMED (UID/BIM/04501/2013) and FCT project PTDC/DTP-PIC/5587/2014 at the University of Aveiro, Portugal.

Author contributions

RGM and RL conceived the study; AK and RGM designed the study, primarily interpreted results, and drafted the article; RGM, RV, AS, MM, TPC, and GG made significant contributions to data collection, including quality control, data processing and statistical analysis; SK and JW revised the study and contributed to the intellectual content. All authors have critically revised the article and approved it for publication.

Competing interests

SK received grants/research support, consulting fees and/or honoraria within the last three years from Angelini, AOP Orphan Pharmaceuticals AG, Celegne GmbH, Eli Lilly, Janssen-Cilag Pharma GmbH, KRKA-Pharma, Lundbeck A/S, Mundipharma, Neuraxpharm, Pfizer, Sanofi, Schwabe, Servier, Shire, Sumitomo Dainippon Pharma Co. Ltd. and Takeda. RL received travel grants and/or conference speaker honoraria within the last three years from Bruker BioSpin MR, Heel, and support from Siemens Healthcare regarding clinical research

using PET/MR; he is shareholder of BM Health GmbH since 2019. The remaining authors declare no competing interests.



Data availability

The functional imaging data generated during the current study are available from the corresponding author upon reasonable request. The correlation lists generated in this study, the associated biological processes, as well as the analyzed risk genes are included in the supplementary information files. All meta-analytical imaging datasets analyzed are available in the Neurosynth repository, <https://neurosynth.org/>, whole-brain transcriptome maps are available from the Neuroimaging Labs (NIL), www.meduniwien.ac.at/neuroimaging/mRNA.html, and gene ontology data are available from the Gene Ontology Consortium, <http://geneontology.org/>.

Code availability

No custom algorithm or software that are central to the conclusion have been used in this study. The codes that were included in this study are available from the corresponding authors upon reasonable request. To process data, licensed software as well as freely accessible webbased tools were used, whereby the authors presume further availability without restrictions on existing code or algorithm availability in the foreseeable future.

Figures









Fig. 1: Rank–rank hypergeometric overlap (RRHO) visual representation of measured imaging data applying automated anatomical labeling (AAL) vs. the Brainnetome atlas. Analyzing single-site data, region-wise RRHO comparing ranked lists including 18,686 genes indicated high agreement between the atlases. Genes with congruent correlation coefficients (either positive or negative) show higher statistical significance in the bottom left and top right corner. Comparisons of both parcellation methods were performed for emotional face recognition in the subcortex ($\rho_{\text{RRHO}} = 0.697$) and cortex ($\rho_{\text{RRHO}} = 0.822$), as well as for reward processing in subcortical ($\rho_{\text{RRHO}} = 0.748$) and cortical regions ($\rho_{\text{RRHO}} = 0.918$).

(a)




(b)





(c)






Fig. 2: Comparison of task-specific functional brain activation and mRNA expression of high-correlating genes in cortical and subcortical regions. **(a)** The scatter plots depict correlations between cortical mRNA levels of FOYN4 and single-site imaging data (emotional face recognition) for voxel-wise ($\rho = 0.285$; 129,817 voxels) and region-wise ($\rho = 0.431$; 210 regions, $p < 0.0001$) analyses. Each dot represents expression values and corresponding imaging parameters at target coordinates or within anatomical regions, respectively. **(b)** Cortical FOYN4 gene expression and brain activation patterns during emotion processing are visualized in MNI space based on whole-brain transcriptome maps (\log_2) and functional magnetic resonance imaging data (t-value). **(c)** Associations between subcortical mRNA levels of MDK and single-site imaging data (acceptance of monetary rewards) for voxel-wise ($\rho = 0.488$; 10,863 voxels) and region-wise ($\rho = 0.803$; 36 regions, $p < 0.0001$) analyses. **(d)** Subcortical MDK gene expression (\log_2) and brain activation patterns during reward processing (t-value).




Fig. 3: Heatmap depicting interrelations of associated biological processes listed within the gene ontology (GO) knowledgebase for emotion and reward processing. Overall, 5 (emotion, positive), 78 (reward, negative), and 27 (reward, positive) GO terms that overlapped with genes strongly correlated with single-site imaging data were compared in subcortical regions (GO terms of analyzed biological programs are provided in Supplementary Table 3). Results yielded marked redundancies for both paradigms (emotional face recognition: blue; acceptance of monetary rewards: red). Considering the multitude of possible associations of 18,686 genes with all listed GO terms, analyses were restricted to higher ontological hierarchies (GO levels above 4). Further, only reasonable correlations between imaging and gene expression data were analyzed, with a cut-off value of $\rho < -0.5$ and > 0.5 , respectively.





Fig. 4: Relationship between risk genes for depressive disorders and genes correlated with functional brain imaging. **(a)** Two master regulators associated with depressive disorders and corresponding subordinate genes correlated with emotion and reward processing present in single-site and meta-analytical data sets are presented. Graphical visualizations are based on associations between gene expression and single-site imaging data in subcortical regions ($p < 0.001$). The size of each circle corresponds to the absolute value of Spearman's correlation coefficient of the respective gene. MEF2C mainly regulates genes with expression patterns negatively correlated with emotion (16 targets, blue) and reward processing (51 targets, red). **(b)** In contrast, TCF4 inversely regulates genes showing negative associations with emotion (15 targets), but positive associations with reward processing (67 targets). **(c)** Gene set enrichment analysis revealed an inversed relationship between emotion and reward processing in the subcortex. Vertical lines on the x-axis represent positions of 42 functional risk genes within each ranked list including 18,686 genes, dashed lines mark the locations of the maximum enrichment score (ES) which draws a density line depicting the peak enrichment of risk genes. Analyzing data from the Neurosynth framework, ES yielded a positive value for emotion (0.275, $p = 0.051$) and a negative value for reward processing (-0.243, $p = 0.65$).

Tables

Table 1: Ranking of Spearman's correlation coefficients for genes with expression patterns showing highest positive associations with single-site imaging data (emotional face processing).

Subcortex				Cortex			
Voxel-wise correlations		Region-wise correlations		Voxel-wise correlations		Region-wise correlations	
rho	Gene name	rho	Gene name	rho	Gene name	rho	Gene name
0.633	MALL	0.865	C10orf125	0.328	SPDYA	0.431	FOXN4
0.616	HRASLS5	0.845	PTRH1	0.298	CCDC62	0.395	PIK3R6
0.614	FAT4	0.818	GHRLOS	0.296	PYGO2	0.390	RYBP
0.612	SCARA5	0.817	SLC24A4	0.289	CPZ	0.387	STC1
0.610	MESP1	0.817	AC022098.3	0.285	FOXN4	0.384	CCDC62
0.608	LINC00260	0.812	ZNF280C	0.283	FRMD3	0.379	FUBP1
0.604	SKAP2	0.810	FUT1	0.282	PHOX2B	0.378	SMYD1
0.602	SCPEP1	0.804	NLE1	0.280	ATXN10	0.356	MIA2
0.596	CRHBP	0.803	FBP1	0.276	XAGE3	0.356	TMPRSS4
0.594	RAB3GAP1	0.803	C16orf55	0.273	EGFL6	0.350	DLG3

Footnote: All listed region-wise correlation coefficients were significant in permutation tests ($p < 0.0001$). Genes marked in bold ranked within the 10 highest positively correlating genes in both voxel-wise and region-wise analyses.

Table 2: Ranking of Spearman’s correlation coefficients for genes with expression patterns showing highest positive associations with single-site imaging data (acceptance of monetary rewards).

Subcortex				Cortex			
Voxel-wise correlations		Region-wise correlations		Voxel-wise correlations		Region-wise correlations	
rho	Gene name	rho	Gene name	rho	Gene name	rho	Gene name
0.488	MDK	0.810	VMO1	0.548	DUSP3	0.698	DUSP3
0.481	HELLS	0.805	OSTM1	0.547	CA10	0.681	CA10
0.475	RBBP8	0.803	MDK	0.542	PIK3CD	0.680	PIK3CD
0.453	ATF1	0.798	KRT18P19	0.533	GRB14	0.653	HDAC9
0.451	KRT18P19	0.791	IMPACT	0.517	ASS1	0.640	LASS6
0.450	CD274	0.787	USP24	0.516	LASS6	0.639	CCNYL1
0.446	C8orf22	0.784	NEK1	0.505	HDAC9	0.637	GRB14
0.442	SALL4	0.782	RCBTB2	0.504	FBXL2	0.627	OLFM3
0.440	USP24	0.782	PCBD2	0.497	OLFM3	0.609	SHC1
0.439	SFRP5	0.781	CD99	0.492	TMEM150C	0.607	NT5DC2

Footnote: All listed region-wise correlation coefficients were significant in permutation tests ($p < 0.0001$). Genes marked in bold ranked within the 10 highest positively correlating genes in both voxel-wise and region-wise analyses.

Table 3: Master regulator analysis in subcortical regions for emotion and reward processing.

	Emotional face recognition		Acceptance of monetary rewards	
	Negative gene associations	Positive gene associations	Negative gene associations	Positive gene associations
Single-site imaging data	TCF4 (15/79)	---	PAX6 (20/89)	SOX5 (57/513)
	MEF2C (16/79)		LHX2 (43/89)	TCF4 (67/513)
	LHX2 (18/79)		MEF2C (51/89)	
	PAX6 (18/79)			
Meta-analytical imaging data	MEF2C (52/1332)	---	MEF2C (21/348)	TCF4 (87/208)
	TCF4 (649/1332)			

Footnote: Master regulators were evaluated in single-site as well as meta-analytical imaging datasets by means of co-regulatory networks built with iRegulon software ($p < 0.001$).

Ranked genes that showed highest correlation with brain activation maps and 42 functional risk genes associated with major depression were used as input data. Values in parenthesis represent the number of targets of each regulatory gene within the group and total number of possible targets.