

1 **Mutually Beneficial Symbiosis Between Human and Gut-Dominant *Bacteroides***
2 **Species Through Bacterial Assimilation of Host Mucosubstances**

3

4 Masahiro Sato¹, Kanta Kajikawa¹, Tomoya Kumon, Daisuke Watanabe, Ryuichi
5 Takase, and Wataru Hashimoto²

6

7 Laboratory of Basic and Applied Molecular Biotechnology, Division of Food Science
8 and Biotechnology, Graduate School of Agriculture, Kyoto University, Uji, Kyoto 611-
9 0011, Japan

10 ¹M.S. and K. K. contributed equally to this work.

11 ²To whom correspondence may be addressed. Wataru Hashimoto; Laboratory of Basic
12 and Applied Molecular Biotechnology, Division of Food Science and Biotechnology,
13 Graduate School of Agriculture, Kyoto University, Uji, Kyoto 611-0011, Japan; Tel.:
14 +81-774-38-3766; Email: hashimoto.wataru.8c@kyoto-u.ac.jp

15

16 Keywords: *Bacteroides* / glycosaminoglycans / microbiota / probiotics / short-chain
17 fatty acids

18

1 **Abstract (228/250 words)**

2 The composition of gut microbiota is influenced by the quantity and type of nutrients in
3 host. Even with some *Bacteroides* species being categorized as pathogens, *Bacteroides*
4 is one of the most dominant gut bacteria. Here we indicate the physiological
5 determinants of the species of *Bacteroides* for being dominant in human gut microbiota.
6 Each of the host extracellular mucosubstances including glycosaminoglycans (GAGs)
7 and mucin has grown human gut microbiota. In spite of the differences among initial
8 microbiota profiles, *Bacteroides* species dominated the community when GAG (e.g.,
9 chondroitin sulfate or hyaluronan) was used as a sole carbon source. In fact, GAGs and
10 the *Bacteroides* genes which are vital for the degradation of GAGs were commonly
11 detected in human feces. Mucin has encouraged the growth of *Bacteroides* and several
12 other genera. A comprehensive analysis on the degradation and assimilation of
13 mucosubstances by the genus *Bacteroides* using around 30 species has shown that most
14 species degrade and assimilate GAGs and mucin, showing that *Bacteroides* species can
15 survive even in the undernutrition condition including the fasting state. In the
16 assimilation of GAG or mucin, *Bacteroides* species significantly secreted essential
17 amino acids, γ -amino butyrate (GABA), and/or short-chain fatty acids which are needed
18 for human health. This is the first report as regards mutually beneficial interaction
19 between human and *Bacteroides* species via bacterial assimilation of host
20 mucosubstances and secretion of metabolites for host health promotion.

21

22 **Significance (120/120 words)**

23 The genus *Bacteroides* is one of the most dominant gut bacteria, although its beneficial
24 effects on human health have not been well understood. Here, we show modes of action

1 in human-*Bacteroides* interrelationship. Mucosubstances including GAGs and mucin
2 secreted by human host are abundant in gut for microbiota to grow well. *Bacteroides*
3 species are dominant in the community in the presence of GAGs, and provide human
4 host with a considerable amount of essential amino acids, γ -amino butyrate, and short-
5 chain fatty acids produced from mucosubstances. These results postulate mutually
6 beneficial symbiosis system between human and *Bacteroides* through bacterial
7 assimilation of host mucosubstances and secretion of metabolites for human body and
8 mental health promotion even in the undernutrition condition including the fasting state.

9

10 **Introduction**

11 In the human gut, there are 100 trillion bacteria communicating with each other and
12 forming a complex microbiota (1). The gut microbiota's composition is influenced by
13 different host diets and metabolizable nutrition for bacteria (2). Contrarily, gut bacteria
14 colonize mucosubstances including mucus layer mucin or extracellular matrix
15 glycosaminoglycan (GAG) made by the host to preserve the intestine (3). Mucin is
16 among the principal components of the mucus layer of the intestine and is comprised of
17 the main chain protein and side chain sugars (e.g., *N*-acetylglucosamine (GlcNAc), *N*-
18 acetylgalactosamine (GalNAc), galactose, fucose, sialic acid) (4). GAG is comprised of
19 a two-sugar component-repeated structure, uronate like iduronic acid or glucuronic acid
20 (GlcUA), and amino sugar including GlcNAc or GalNAc (5). In amino sugars, both
21 hydroxyl and amino groups are highly sulfated. GAG is categorized into chondroitin
22 sulfate (CS), hyaluronic acid (HA), heparin (HP), heparan sulfate, dermatan sulfate, and
23 keratan sulfate based on constituent sugar, glycoside bond manner, and the level of
24 sulfated groups (5).

1 Some human gut bacteria, for example, *Bacteroides*, *Clostridium*, *Enterococcus*, and
2 *Streptococcus* genera, break down GAGs that are supplied by the host intestine which
3 are not dependent on intake of meals (6-9). Our previous studies have exhibited
4 molecular machineries of depolymerization, import, degradation, and metabolism of
5 GAGs in the genus *Streptococcus* (10, 11) (*SI Appendix*, Fig. S1). Mainly among these
6 gut bacteria, *Bacteroides* species assimilate polysaccharides that are supplied by the
7 host and colonize the colon intestinal mucus layer as the largest group in humans in
8 spite of race (12-14), even though the genus *Bacteroides* is often classified as an
9 opportunistic pathogen (15). *Bacteroides thetaiotaomicron* assimilates both mucin O-
10 binding polysaccharides and GAGs including CS, HA, and HP (16-18).

11 On the other hand, short-chain fatty acids which are made by some gut bacteria
12 positively influence the host physiology, for example (19), (i) improve the gut
13 environment by decreasing the pH, (ii) adjust the host immune system, (iii) be the
14 energy source of the host, (iv) improve insulin resistance, and (v) decrease the risk of a
15 heart attack. Some species of *Bacteroides* can create short-chain fatty acids (20),
16 indicating their function as next-generation probiotics (15, 21, 22). For one to know the
17 reason why humans allow *Bacteroides* species to be dominant in gut microbiota and
18 why humans do not repress the bacterial species, this article deals with the mechanism
19 of symbiosis between humans and gut-dominant genus *Bacteroides* through
20 comprehensive analyses to explore *Bacteroides* species capability to degrade as well as
21 assimilate human mucosubstances (GAGs and mucin) and to dictate bacterial
22 metabolites during the human secretion assimilation.

23

24 **Results**

1 **GAG Is Available to Human Gut Microbiota.** Extracellular matrices under the mucus
2 layer which is comprised of mucin (23) are generally composed of GAGs, including CS,
3 HA, and HP. Since the mucus layer hinders bacteria from invading the host cells (24),
4 these bacteria can hardly utilize GAGs. On the contrary, human feces have components
5 from epithelial cell shedding (25), indicating that GAGs that are from the intestinal
6 tracts are available to the gut microbiota. Therefore, the concentration of GAGs,
7 especially those in sulfated forms, in feces was measured using the 1,9-dimethyl-
8 methylene blue (DMMB) method (26). The contents of GAGs in independent feces
9 from three volunteers A to C (two twenties and one fifties) were as follows: volunteer
10 A, 0.098% (w/w); volunteer B, 0.112%; and volunteer C, 0.033%. These values are
11 similar to those of GAGs that are part of bacterial media, indicating that there are plenty
12 GAGs in the human gut for microbiota to grow well. In fact, *B. thetaiotaomicron* has
13 shown a remarkable growth in the minimal medium with the use of 0.02% (w/v) GAG
14 (HA) as a sole carbon source (*SI Appendix*, Fig. S2).

15 ***Bacteroides* Species Are Dominant in the Community in the Presence of GAG.** To
16 explore the effects of carbon sources on human gut microbiota, fecal samples were
17 cultured for 48 hours in a minimal medium that contains glucose, mucin, or
18 representative GAG (i.e., chondroitin sulfate C (CSC), HA, or HP) as a sole carbon
19 source. Microbes in independent samples from three donors (one twenties, one thirties,
20 and one fifties) has exhibited vigorous growth in each medium (*SI Appendix*, Fig. S3),
21 indicating that microbes are present to degrade and/or assimilate GAG or mucin. The
22 collected microbiota samples before and after growth were subjected to 16S rDNA
23 amplicon sequence analysis (Fig. 1). Prior to cultivation, each sample had a wide
24 variety of bacteria (Fig. 1A): Donors A, B, and C comprised of *Bifidobacterium*,

1 *Prevotella*, and *Bacteroides* as the major genera, respectively. *Collinsella*, *Blautia*,
2 *Faecalibacterium*, and *Dialister* genera were also detected well. The frequency of
3 *Bacteroides* species ranged from 1.9 to 27.2%; among them, both *B. vulgatus* and *B.*
4 *dorei* were most commonly observed. Succeeding cultivation for 24 hours in the
5 presence of glucose, various genera such as *Bacteroides*, *Bifidobacterium*,
6 *Erysipelatoclostridium*, and *Escherichia* propagated, even though many other early
7 genera or species decayed. In a mucin-containing medium, *Collinsella* and *Blautia*
8 genera also increased or maintained the frequency. Contrarily, only the *Bacteroides*
9 species, especially *B. ovatus* and *B. xylanisolvens*, predominated the community (over
10 50% of total amplicons) in a GAG-containing medium in any donor sample. In view of
11 the low initial frequency of both *B. ovatus* and *B. xylanisolvens* (0.0–1.7% and 0.0–
12 0.6%, respectively), these species exhibited the best adaptive fitness to the used
13 experimental conditions with GAG. Altogether, our data indicate that carbon sources
14 are the key determinants in microbiota formation and maintenance.

15 Principal component analysis (PCA) was done with the use of 0- and 24-hour data of
16 averaged genus (or *Bacteroides* species) frequency (Fig. 1B). The first and second
17 principal components (PC1 and PC2) accounted for 35.0% and 23.3% of the variance,
18 respectively. Both the factor loading plot and the PCA score plot have shown that the
19 majority of *Bacteroides* species, such as *B. ovatus*, *B. xylanisolvens*, *B. uniformis*, *B.*
20 *finegoldii*, and *B. faecis*, formed a cluster with large negative PC1 values, corresponding
21 to the high frequency in CSC, HA, or HP-containing medium. Therefore, GAG may be
22 one of the reasons for the dominance of most *Bacteroides* species in gut microbiota.
23 The growth profiles of six major genera, exhibited from the growth curves of whole
24 microbiota and the genus frequency in the microbiota, further supported that

1 *Bacteroides* species prefer GAG than glucose or mucin (*SI Appendix*, Fig. S4),
2 although, based on Fig. 1B, both *B. vulgatus* and *B. dorei* were plotted apart from the
3 major *Bacteroides* group. The frequency of these two species was relatively high prior
4 to cultivation but was lesser when GAG was used as a carbon source. Mucin was the
5 only carbon source to grow the *Bacteroides* as well as five other major genera (*SI*
6 *Appendix*, Fig. S4).

7 **Most *Bacteroides* Degrade GAGs.** Since *Bacteroides* species prefer GAGs, their
8 GAG-degrading abilities were assessed with the use of CSC, HA, and HP. We have
9 earlier reported the degradation of GAGs by 6 *Bacteroides* species of 11 species tested
10 through the halo assay with the use of GAG minimal medium plates (9). The detection
11 of halo means there is a degradation of GAGs (27). Thus, to assess GAG degradation as
12 one of the common characteristics among *Bacteroides* species, the other 17 species
13 were also subjected to the halo assay (*SI Appendix*, Fig. S5), thus finding that *B. caccae*,
14 *B. faecis*, and *B. finegoldii* degrade CSC. Additionally, *B. caccae*, *B. faecis*, *B.*
15 *finegoldii*, *B. nordii*, and *B. xylanisolvens* were found to be novel HA-degrading
16 *Bacteroides*. Contrarily, the species tested here do not degrade HP, showing that
17 *Bacteroides* species prefer both CSC and HA containing 1,3-glycoside bond than HP
18 containing 1,4-glycoside bond.

19 We then conducted the halo assay for the confirmation of the GAG-degrading ability
20 of *Bacteroides* species under high nutrition conditions (Fig. 2). The nutrient-rich
21 medium (Gifu Anaerobic Medium: GAM) plate was used for the halo assay since some
22 *Bacteroides* species showed little growth in the GAG minimal plate, due to the lack of
23 essential components including cysteine which is used for a reducing agent. In addition
24 to 17 species used for the halo assay with GAG minimal plate, 8 other species were

1 investigated with the use of the GAM plate. Because there was no precipitation formed
2 in the HA-containing GAM plate, HA was not appropriate for this halo assay under
3 high-nutrition conditions. In addition to CSC-degrading *Bacteroides* species found on
4 the minimal medium plate, six species were found to degrade CSC: *B. cellulosityticus*,
5 *B. eggerthii*, *B. fluxus*, *B. gallinarum*, *B. massiliensis*, and *B. oleiciplenus*. However, in
6 the case of HP minimal medium, there were no *Bacteroides* species found to degrade
7 HP; the HP-containing GAM plate formed a halo, so that nine species exhibited HP-
8 degrading ability as follows: *B. clarus*, *B. coprosuis*, *B. eggerthii*, *B. faecis*, *B.*
9 *finegoldii*, *B. intestinalis*, *B. ovatus*, *B. stercoris*, and *B. thetaiotaomicron*. All in all, 18
10 of 28 *Bacteroides* species that were tested showed GAG (CSC, HA, and/or HP)-
11 degrading ability (*SI Appendix*, Table S1).

12 **Assimilation of GAGs by *Bacteroides* Species.** The assimilation of GAGs by 29
13 *Bacteroides* species was further investigated by monitoring the optical density at 600
14 nm (OD₆₀₀) of the bacterial culture in the liquid minimal medium such as CSC, HA, or
15 HP as a sole carbon source (Fig. 3a, *SI Appendix*, Fig. S6), resulting in the growth of 20
16 species in the presence of CSC as follows: *B. barnesiae*, *B. caccae*, *B. cellulosityticus*,
17 *B. clarus*, *B. coagulans*, *B. coprocola*, *B. coprophilus*, *B. eggerthii*, *B. faecis*, *B.*
18 *finegoldii*, *B. fluxus*, *B. gallinarum*, *B. intestinalis*, *B. massiliensis*, *B. nordii*, *B.*
19 *oleiciplenus*, *B. ovatus*, *B. stercoris*, *B. thetaiotaomicron*, and *B. xylanisolvens*. In the
20 case of HA, 15 species proliferated, that is, *B. barnesiae*, *B. caccae*, *B. clarus*, *B.*
21 *coagulans*, *B. coprocola*, *B. coprophilus*, *B. faecis*, *B. finegoldii*, *B. fluxus*, *B.*
22 *gallinarum*, *B. massiliensis*, *B. nordii*, *B. ovatus*, *B. thetaiotaomicron*, and *B.*
23 *xylanisolvens*. HP was assimilated by 13 species as follows: *B. clarus*, *B. eggerthii*, *B.*
24 *faecis*, *B. finegoldii*, *B. fluxus*, *B. gallinarum*, *B. intestinalis*, *B. massiliensis*, *B. nordii*,

1 *B. ovatus*, *B. stercoris*, *B. thetaiotaomicron*, and *B. xylanisolvens*. There is no species
2 that has shown growth without GAGs, showing that these species grew by assimilating
3 GAGs as a sole carbon source.

4 The time-dependent decrease of GAG concentration in the CSC minimal medium
5 was examined during the growth of *B. faecis* as a representative. In accordance to the
6 OD₆₀₀ increase of the bacterial culture broth, there was a concentration decrease of
7 CSC, and the polysaccharide was finally totally consumed (Fig. 3B). This result has
8 directly shown that CSC was assimilated by *B. faecis*. In total, 20 species of 29
9 *Bacteroides* species that were tested here assimilate GAGs (*SI Appendix*, Table S1).

10 In the majority of *Bacteroides* species, the growth profiles in human gut microbiota
11 (Fig. 1B) and in pure culture (Fig. 3A) were quite steady: *B. ovatus*, *B. xylanisolvens*, *B.*
12 *fingoldii*, and *B. faecis* in the major *Bacteroides* group showed high assimilation
13 ability, while *B. vulgatus* and *B. dorei* did not have GAG utilization. However, a major-
14 group species (*B. uniformis*) has no degradation or assimilation of GAG. Such species
15 could not feed by themselves on GAG but may grow in microbiota with the help of
16 GAG-degrading/ GAG-assimilating bacteria.

17 **Frequent Detection of GAG Lyase Gene in Human Gut Microbiota.** As a first step
18 of bacterial action on GAGs, bacterial GAG lyases are very important for
19 depolymerization of GAGs (*SI Appendix*, Fig. S1). Among these, HA/CS and heparan
20 sulfate lyases are categorized as Polysaccharide Lyase Families 8 (PL8) and PL12,
21 respectively, in the database CAZy (<http://www.cazy.org/>) (6, 16, 17). The primers to
22 amplify these lyase genes were designed by referring to those genes in four *Bacteroides*
23 species as follows: *B. cellulosilyticus*, *B. ovatus*, *B. thetaiotaomicron*, and *B.*
24 *xylanisolvens*. Family PL8 and PL12 GAG lyase genes were discovered from three

1 independent feces, and their frequency was 6.0%, 0.85%, and 3.2% in human gut
2 microbiota on the basis of the bacterial cells calculated from 16S rDNA amplification
3 (*SI Appendix*, Table S2). These values were reasonable based on the frequency of
4 *Bacteroides* species in fecal samples as described above.

5 **Assimilation of Mucin by *Bacteroides* Species.** Besides GAG, the assimilation of
6 mucin was assessed with the use of 28 *Bacteroides* species on the purified mucin
7 minimal medium. Although no *Bacteroides* species showed a significant growth on the
8 medium without mucin, 15 species assimilated mucin as follows: *B. caccae*, *B. clarus*,
9 *B. coprosuis*, *B. faecis*, *B. finegoldii*, *B. fragilis*, *B. gallinarum*, *B. helcogenes*, *B.*
10 *intestinalis*, *B. nordii*, *B. ovatus*, *B. stercoris*, *B. thetaiotaomicron*, *B. uniformis*, and *B.*
11 *xylanisolvans* (Fig. 3A, *SI Appendix*, Table S1). Among the mentioned species, six, *B.*
12 *clarus*, *B. coprosuis*, *B. faecis*, *B. gallinarum*, *B. helcogenes*, and *B. nordii*, were the
13 first ones to show mucin assimilation. Since commercially available mucin contained
14 GAGs, the amount of sulfated GAGs in the purified mucin was measured using the
15 DMMB method, resulting in the purified mucin containing 7–8% (w/v) of GAGs, which
16 corresponds to 0.02% (w/v) GAGs included in 0.25% (w/v) mucin medium which is
17 used for the growth assay. To remove the possibility that *Bacteroides* species
18 assimilated only GAG in the mucin minimal medium, *B. thetaiotaomicron* was cultured
19 in 0.25% (w/v) mucin or 0.02% (w/v) GAG minimal medium. As a consequence, there
20 was a higher growth level on mucin than that on GAG (*SI Appendix*, Fig. S2), showing
21 that the bacterial cells assimilated mucin.

22 ***Bacteroides* Species Utilize GAG or Mucin as a Sole Nitrogen Source.** Nitrogen is
23 one of the essential elements, despite its limited quantity in the gut environment.
24 Therefore, bacteria need to acquire nitrogen to be indigenous in the colon. Because

1 nitrogen is included in GAG or mucin, we tested whether *Bacteroides* species can use
2 HA or mucin as a sole nitrogen source (*SI Appendix*, Fig. S7). Three species, *B.*
3 *thetaiotaomicron*, *B. ovatus*, and *B. faecis*, have shown a remarkable growth even in
4 nitrogen-restricted GAG or mucin medium. In the meantime, these three species showed
5 no growth in nitrogen-restricted glucose medium, even though they grew well in the
6 presence of ammonium sulfate. *Bacteroides* species can, therefore, use GAG or mucin
7 as carbon and nitrogen sources. This feature is one of the reasons why *Bacteroides*
8 species have dominance in the gut environment.

9 **Metabolites from *Bacteroides* Species during Assimilation of Host Mucosubstances.**

10 Metabolites in the culture broth secreted by *B. thetaiotaomicron* or *B. ovatus* were
11 investigated in the GAG (HA) or mucin minimal medium, resulting in the detection of
12 19 kinds of standard amino acids, γ -amino butyrate (GABA), α -amino butyrate, and
13 ethanol amine (Fig. 4A). A huge amount of Ala was seen under all conditions, and
14 GABA, Glu, and Val were remarkably secreted when HA was assimilated. On the
15 contrary, when the bacterial cells assimilated mucin, there was an excretion of basic
16 amino acids, for example, Lys and Arg. Moreover, organic acids were also investigated.
17 Three *Bacteroides* species, *B. thetaiotaomicron*, *B. ovatus*, and *B. faecis*, grown in the
18 HA minimal medium secrete short-chain fatty acids including acetic and propionic acids
19 as well as lactic, succinic, and/or formic acids (Fig. 4B).

20

21 **Discussion**

22 Based on the above-described results, we have discussed the following: Since GAGs
23 and mucin are constantly supplied in human gut independent from nutrient intake by the
24 host, GAG or mucin assimilation abilities are necessary for survival of dominant

1 bacteria in the human gut, supporting the hypothesis that assimilation of GAG or mucin
2 is essential for gut microbiota dominance in the human gut. In fact, the abundance of
3 *Bacteroides* species in the gut microbiota has increased in female mice by oral
4 administration of chondroitin sulfates A and C (28). The gene expression involved in
5 the assimilation of GAGs is also upregulated in *Bacteroides* species in the presence of
6 GAGs (18, 29). As regards structure, GAG is simpler than mucin, so that GAG may be
7 degraded easier than mucin (3, 30). Indeed, more species have shown an assimilation
8 ability toward GAG rather than mucin (Fig. 3A, *SI Appendix*, Table S1).

9 Amino acids secreted by gut microbiota are used by the host as well as other gut
10 bacteria (31). *Bacteroides* species grown in GAG or mucin minimal medium secreted
11 most of the standard amino acids and organic acids such as short-chain fatty acids (Fig.
12 4), showing that the host human and other bacteria can assimilate these metabolites.
13 Additionally, *Bacteroides* species could provide important amino acids to host human
14 by converting host GAGs to those amino acids which are not dependent from nutrient
15 intake by the host. Even with the weight of gut microbiota of about 1.5 kg and
16 *Bacteroides* species being dominant species (~50% of total), the amount of secreted
17 amino acids is not negligible, even though amino acids are generally absorbed by the
18 small intestine and *Bacteroides* species are broadly distributed from the stomach to the
19 large intestine (32). The quantities of amino acids produced by *Bacteroides* species are
20 almost equal to the recommended amount for a person on a day by the WHO (*SI*
21 *Appendix*, Table S3), indicating that the human obtains a considerable amount of
22 nutrition from metabolites secreted by gut microbiota. In the recent time, *Bacteroides*
23 species produce much GABA, and negative correlation is apparent between their
24 abundance and depressive symptoms (33), showing that GABA which is produced by

1 human gut microbiota is as a signal molecule in the enteric nervous system (34). Thus,
2 besides nutritional effects, *Bacteroides* species producing GABA from the host GAG
3 may be a contributing factor to human mental health.

4 Furthermore, gut microbiota is involved in host immunity and/or energy source via
5 the secretion of organic acids such as short-chain fatty acids (19). For the most part,
6 short-chain fatty acids which are very important to the host are produced at 50–100 mM
7 in the colonic lumen by gut microbiota (35, 36). Based on the secretion level from the
8 mucosubstances (GAG and mucin) (Fig. 4B), *Bacteroides* species are the main
9 producers of short-chain fatty acids in the human gut. Since the intestine mucus of the
10 host is supplied inside of the gut which is independent from the nutrient uptake by the
11 host, *Bacteroides* species should supply physiologically essential molecules to the host
12 or other bacteria under fasting conditions. These results support the fact that
13 *Bacteroides* species are known as the next-generation probiotics (15, 22) and/or
14 pharmabiotics (34).

15 The overall results acquired here hypothesized that there is a mutually beneficial
16 relationship model between *Bacteroides* species and the human host (Fig. 4C). It is
17 essential for humans to secrete extracellular gut mucosubstances including GAGs and
18 mucin for the maintenance of cell structure/function and prevention of inflammation
19 due to pathogenic bacteria. Some gut bacteria convert these mucosubstances to essential
20 molecules for the host. This mutual beneficial relationship has probably been accepted
21 as a result of natural selection in the human gut. For humans, *Bacteroides* species in the
22 gut leads to the promotion of host health by providing amino acids and organic acids
23 such as short-chain fatty acids, so that the human immune system has not excluded
24 *Bacteroides* species from the gut. On the contrary, for the *Bacteroides* species, the

1 human gut is a well-organized place to survive due to the continuous supply of large
2 quantities of mucosubstances in which *Bacteroides* species can assimilate dominantly in
3 human gut microbiota. Human health is also necessary for *Bacteroides* species to keep
4 the residence place safe.

5 Finally, the mutually beneficial symbiosis between humans and gut-dominant
6 *Bacteroides* species is posited via bacterial assimilation of host mucosubstances and
7 secretion of metabolites for host health promotion.

8

9 **Materials and Methods**

10 **Materials.** HA and DMMB were purchased from Sigma-Aldrich. Mucin from porcine
11 stomach and CSC were acquired from Wako. HP was from Nacalai Tesque. The
12 nutrition-rich medium, GAM, was from Nissui Pharmaceutical Co. All other analytical
13 grade chemicals utilized in this study were available commercially. The feces were
14 kindly given to us as gifts by the Japanese volunteers. Informed consent was acquired
15 from all subjects, and experiments with the use of these feces were approved by the
16 Committee of Research Activity Promotion of Graduate School of Agriculture, Kyoto
17 University.

18 **Purification of GAG and Mucin.** Low-molecular-weight molecules contaminated in
19 each GAG reagent were removed by dialysis against pure water. Unless otherwise
20 stated, the percent concentration represents w/v. Mucin was dissolved at a 2% final
21 concentration in 20 mM potassium phosphate buffer (pH 7.8), which contains 0.1 M
22 NaCl. Several drops of toluene were placed in the mucin suspension, which is followed
23 by stirring for 1 hour at room temperature. The suspension was then adjusted to a pH of
24 7.2 with 2 M NaOH and stirred for 23 hours at room temperature. The supernatant was

1 collected after centrifugation (4°C, 10,000×g, 10 min). Cold ethanol was then added to
2 become 60% (v/v), for the precipitation of mucin to occur. Centrifugation was done
3 with the same condition, and the resultant pellet was then dissolved in 0.1 M NaCl. The
4 addition of ethanol followed by centrifugation was repeated twice. The last step was to
5 dissolve the pellet in pure water (80 mL) instead of 0.1 M NaCl. This solution was
6 dialyzed against pure water and then freeze-dried to obtain the purified mucin (37).

7 **Bacteria Strains.** A total of 30 *Bacteroides* species was utilized in this study. Among
8 them, 29 species were purchased from Japan Collection of Microorganisms (JCM), such
9 as *B. acidifaciens* JCM10556, *B. barnesiae* JCM13652, *B. caccae* JCM9498, *B.*
10 *cellulosilyticus* JCM15632, *B. clarus* JCM16067, *B. coagulans* JCM12528, *B.*
11 *coprocola* JCM12979, *B. coprophilus* JCM13818, *B. coprosuis* JCM13475, *B. dorei*
12 JCM13471, *B. eggerthii* JCM12986, *B. faechinchillae* JCM17102, *B. faecis* JCM16478,
13 *B. finegoldii* JCM13345, *B. fluxus* JCM16101, *B. fragilis* JCM11019, *B. gallinarum*
14 JCM 13658, *B. helcogenes* JCM6297, *B. intestinalis* JCM13265, *B. massiliensis*
15 JCM13223, *B. nordii* JCM12987, *B. oleiciplenus* JCM16102, *B. ovatus* JCM5824, *B.*
16 *paurosaccharolyticus* JCM15092, *B. salanitronis* JCM13657, *B. stercoris* JCM9496, *B.*
17 *thetaitaomicron* JCM5827, *B. uniformis* JCM5828, and *B. xylanisolvens* JCM15633.

18 In addition to this, *B. vulgatus* NBRC14291 was from the Biological Resource Center in
19 the National Institute of Technology and Evaluation (NBRC/NITE).

20 **16S rDNA Amplicon Sequence Analysis.** There is approximately 1 g of human fecal
21 sample dissolved in 10 mL of sterilized saline (0.9% NaCl). A 300-μL aliquot of this
22 suspension was inoculated into a test tube which contains 15 mL of assimilation
23 validation liquid medium [0.1% ammonium sulfate, 0.226% KH₂PO₄, 0.09% KH₂PO₄,
24 0.0004% FeSO₄(II), 0.09% NaCl, 0.0027% CaCl₂/2H₂O, 0.002% MgCl₂/6H₂O, 0.001%

1 MnCl₂/4H₂O, 0.001% CoCl₂, 0.0005% hemin, 0.00001% vitamin K₁, 0.00001%
2 ethanol, 0.0000005% vitamin B₁₂, and 0.04% L-cysteine, with or without 0.25%
3 glucose, dialyzed GAG (CSC, HA, or HP) or the purified mucin]. After being
4 anaerobically cultured at 37°C for 24 or 48 hours, cells acquired after centrifugation
5 were washed using the sterilized saline and immediately frozen in liquid nitrogen.

6 Both DNA extraction from human gut microbiota and 16S rDNA amplicon sequence
7 analysis were performed by TechnoSuruga Laboratory Co., based on a previously
8 reported method (38). In summary, the V3–V4 region of 16S rDNA was amplified with
9 the use of the 341F/R806 primer sets. Sequencing was conducted using a paired-end,
10 2×300-bp cycle run on a MiSeq sequencing system (Illumina) and MiSeq Reagent Kit
11 version 3 (600 cycle) chemistry. After the sequencing was done, image analysis, base
12 calling, and error estimation were done with the use of the Illumina Real-Time Analysis
13 software (version 1.17.28). Paired-end sequencing with read lengths of approximately
14 430 bp was done as well. Succeeding the demultiplexing, a clear overlap in the paired-
15 end reads was seen. This made paired reads be joined together with the fastq-join
16 program. Only reads that had quality value (QV) scores of ≥ 20 for more than 99% of
17 the sequence were extracted for supplemental analysis. Metagenome@KIN software
18 (World Fusion) was utilized for homology searching with the determined 16S rDNA
19 sequences, against the DB-BA13.0 microbial identification database (TechnoSuruga
20 Laboratory) (39, 40). Bacterial species were then identified based on the data from 97%
21 similarity cut-off with DB-BA13.0. All sequences have been deposited in the DNA
22 Data Bank of Japan (DDBJ) under the accession number DRA010273.

23 **GAG Quantitation.** Fecal samples (about 2 g) acquired from three volunteers were
24 dissolved in 50 mL of pure water followed by overnight rotation. Centrifugation and

1 filtration were done for insoluble residues to be removed. The reagent DMMB
2 determined the GAG content at a final concentration of 0.0016% in 0.304% glycine,
3 0.16% NaCl, and 0.057% (v/v) acetic acid as quantitative reagent (26). Standard curve
4 was made using a variety of CSC concentrations (0, 10, 20, 30, 40, and 50 mg/mL). The
5 absorbance at 525 nm (Abs_{525}) was measured after 80 μ L of samples and 800 μ L of
6 quantitative solution were mixed. To remove the effect of other contaminated materials
7 toward Abs_{525} in feces, Abs_{525} values of tenfold diluted fecal samples with pure water
8 were measured and subtracted from Abs_{525} values of DMMB when GAGs in feces were
9 quantified.

10 **Halo Assay for the Detection of GAG Degradation.** *Bacteroides* species were grown
11 in an anaerobic condition at 37°C overnight in 5 ml of liquid GAM (1% peptone, 0.3%
12 soy peptone, 1% proteose peptone, 1.35% digested serum powder, 0.5% yeast extract,
13 0.22% meat extract, 0.12% liver extract, 0.3% glucose, 0.25% KH_2PO_4 , 0.3% NaCl,
14 0.5% soluble starch, 0.03% L-cysteine hydrochloride, and 0.03% sodium thioglycolate)
15 as a preculture. The preculture broth (1 mL) was centrifuged at 9,700 \times g for 5 min, and
16 the resulting pellet (bacterial cells) was then washed with 1 mL of sterilized saline
17 (0.9% NaCl), which is followed by centrifugation (9,700 \times g, 5 min). The pellet was
18 suspended in saline to acquire $OD_{600} = 1$. The cell suspension (10 μ L) was seen on the
19 center of the halo minimal medium plate [0.1% yeast extract, 0.1% KH_2PO_4 , 0.1%
20 Na_2HPO_4 , 0.01% $MgSO_4/7H_2O$, 1% bovine serum albumin (BSA), 0.2% dialyzed GAG
21 (CSC, HA, or HP), and 1% agar]. The plate was incubated in an anaerobic condition at
22 37°C for 7 days. Afterward, 1 mL of 2 M acetic acid was spread on the plate to form
23 white precipitation of the BSA-GAG complex. In the case of GAG-degrading bacteria,
24 halo (clear zone) was seen on the plate due to the lack of GAG polymer (27).

1 For some *Bacteroides* species with a little growth in the halo minimal medium plate,
2 the nutrition-rich halo medium plate was also utilized. This plate has GAM to supply a
3 proper reducing agent for anaerobic bacteria as follows: 0.4% peptone, 0.12% soy
4 peptone, 0.4% proteose peptone, 0.54% digested serum powder, 0.2% yeast extract,
5 0.088% meat extract, 0.048% liver extract, 0.12% glucose, 0.1% KH₂PO₄, 0.12% NaCl,
6 0.2% soluble starch, 0.012% L-cysteine hydrochloride, 0.012% sodium thioglycolate,
7 1% BSA, and 0.2% dialyzed GAG (CSC or HP).

8 **Assimilation Assay.** To investigate the mucosubstance assimilation by *Bacteroides*
9 species, OD₆₀₀ of the culture broth was measured after their inoculation into a variety of
10 liquid media. *Bacteroides* species precultured in liquid GAM was centrifuged at
11 15,000×g for 10 min, and the resulting pellet (bacterial cells) was washed using
12 sterilized saline (0.9% NaCl). The bacterial cells were washed thrice and finally
13 suspended in saline to acquire OD₆₀₀ = 1. The cell suspension (300 μL) was inoculated
14 into 15 mL of the modified assimilation validation liquid medium (41) [0.1%
15 ammonium sulfate, 0.226% KH₂PO₄, 0.09% KH₂PO₄, 0.0004% FeSO₄ (II), 0.09%
16 sodium chloride, 0.0027% CaCl₂/2H₂O, 0.002% MgCl₂/6H₂O, 0.001% MnCl₂/4H₂O,
17 0.001% CoCl₂, 0.0005% hemin, 0.00001% vitamin K₁, 0.00001% ethanol, 0.0000005%
18 vitaminB₁₂, 0.04% L-cysteine, and 0.25% dialyzed GAG (CSC, HA, or HP) or the
19 purified mucin] and was further cultured at a temperature of 37°C with anaerobic
20 conditions. The negative control, validation liquid medium with no inoculation, was
21 also incubated to check the background of OD₆₀₀. For *B. faecis*, OD₆₀₀ of the bacterial
22 culture with CSC as a sole carbon source was monitored by measuring the CSC
23 concentration.

1 Three species, *B. faecis*, *B. ovatus*, and *B. thetaiotaomicron*, were grown in nitrogen-
2 restricted medium (assimilation validation liquid medium without ammonium sulfate)
3 with HA, mucin, or glucose present as a sole carbon source to check whether these
4 bacterial cells can use GAG or mucin as a sole nitrogen source. As indicated above,
5 bacterial growth was monitored by measuring OD₆₀₀ of the culture broth.

6 **Metagenomics of Gut Bacteria for GAG Degradation.** Metagenomics were
7 accomplished with the use of fecal samples from three volunteers. The analyses were
8 conducted by TechnoSuruga Laboratory Co. DNA primers, for the amplification of
9 GAG lyase genes, were designed by referring to family PL8 and PL12 lyase gene
10 sequences found in database CAZy (<http://www.cazy.org/>) as follows: PL8 forward,
11 CTSGAYGGDGCMAVAAAYATAGA; PL8 reverse,
12 TTTCCATCGGGAGWDCCRGCHAD; PL12 forward,
13 RAYTAYCCVGGWYTRGARAAAG; and PL12 reverse,
14 WCCAYTKATGRCGATGMADYTG. The total bacterial cell numbers were estimated
15 by real-time PCR with the use of primers which are specific to 16S rDNA.

16 **Metabolic Assay of *Bacteroides* Species through Assimilation of GAG or Mucin.**

17 Three species, *B. faecis*, *B. ovatus*, and *B. thetaiotaomicron*, were cultured in the
18 assimilation validation liquid medium that contained HA or mucin as a sole carbon
19 source. Organic acids in the bacterial culture broth for 48 hours were measured by
20 TechnoSuruga Laboratory Co. at 45°C of the column oven temperature and 0.8 mL/min
21 of flow rate with the use of 5 mM *p*-toluenesulfonic acid as a solvent by HPLC system
22 (Shimadzu) that is equipped with a column of Shim-pack SCR-102(H) and electrical
23 conductivity detector of CDD-10A. As regards *B. thetaiotaomicron* and *B. ovatus*, the
24 bacterial culture broth was also subjected to Amino Acid Analyzer L-8900 (Hitachi

1 High-Technologies) which is equipped with a column of #2622SC-PF and a detector of
2 photometer in the Global Facility Center of Hokkaido University. Free amino acids
3 were seen at wavelengths of 570 and 440 nm by the post-staining method with
4 ninhydrin. Before analyses, mucin was taken out with the use of ethanol precipitation.
5 The medium with no cells was also investigated as a background.

6

7 **References**

- 8 1. A. Suau *et al.*, Direct analysis of genes encoding 16S rRNA from complex
9 communities reveals many novel molecular species within the human gut. *Appl.*
10 *Environ. Microbiol.* **65**, 4799–4807 (1999).
- 11 2. S. A. Smits *et al.*, Seasonal cycling in the gut microbiome of the Hadza hunter-
12 gatherers of Tanzania. *Science* **357**, 802–806 (2017).
- 13 3. E. C. Martens, H. C. Chiang, J. I. Gordon, Mucosal glycan foraging enhances
14 fitness and transmission of a saccharolytic human gut bacterial symbiont. *Cell*
15 *Host Microbe* **4**, 447–457 (2008).
- 16 4. R. Bansil, B. S. Turner, The biology of mucus: Composition, synthesis and
17 organization. *Adv. Drug Deliv. Rev.* **124**, 3–15 (2018).
- 18 5. N. S. Gandhi, R. L. Mancera, The structure of glycosaminoglycans and their
19 interactions with proteins. *Chem. Biol. Drug Des.* **72**, 455–482 (2008).
- 20 6. A. Cartmell *et al.*, How members of the human gut microbiota overcome the
21 sulfation problem posed by glycosaminoglycans. *Proc. Natl. Acad. Sci. USA* **114**,
22 7037–7042 (2017).

- 1 7. B. Canard, T. Garnier, B. Saint-Joanis, S. T. Cole, Molecular genetic analysis of
2 the *nagH* gene encoding a hyaluronidase of *Clostridium perfringens*. *Mol. Gen.*
3 *Genet.* **243**, 215–224 (1994).
- 4 8. M. J. Jedrzejewski, Unveiling molecular mechanisms of bacterial surface proteins:
5 *Streptococcus pneumoniae* as a model organism for structural studies. *Cell. Mol.*
6 *Life Sci.* **64**, 2799–2822 (2007).
- 7 9. K. Kawai, R. Kamochi, S. Oiki, K. Murata, W. Hashimoto, Probiotics in human
8 gut microbiota can degrade host glycosaminoglycans. *Sci. Rep.* **8**, 10674 (2018).
- 9 10. Y. Maruyama *et al.*, Metabolic fate of unsaturated glucuronic/iduronic acids from
10 glycosaminoglycans: molecular identification and structure determination of
11 streptococcal isomerase and dehydrogenase. *J. Biol. Chem.* **290**, 6281–92 (2015).
- 12 11. S. Oiki *et al.*, Streptococcal phosphotransferase system imports unsaturated
13 hyaluronan disaccharide derived from host extracellular matrices. *PLoS One* **14**,
14 e0224753 (2019).
- 15 12. E. S. Friedman *et al.*, Microbes vs. chemistry in the origin of the anaerobic gut
16 lumen. *Proc. Natl. Acad. Sci. USA* **115**, 4170–4175 (2018).
- 17 13. N. L. Zitomersky *et al.*, Characterization of adherent *Bacteroidales* from intestinal
18 biopsies of children and young adults with inflammatory bowel disease. *PLoS*
19 *One* **8**, e63686 (2013).
- 20 14. F. Bäckhed *et al.*, Host-bacterial mutualism in the human intestine. *Science* **307**,
21 1915–1920 (2005).
- 22 15. F. Sun *et al.*, A potential species of next-generation probiotics? The dark and light
23 sides of *Bacteroides fragilis* in health. *Food Res. Int.* **126**, 108590 (2019).

- 1 16. N. A. Pudlo *et al.*, Symbiotic human gut bacteria with variable metabolic
2 priorities for host mucosal glycans. *mBio* **6**, e01282-15 (2015).
- 3 17. D. Ndeh *et al.*, Metabolism of multiple glycosaminoglycans by *Bacteroides*
4 *thetaiotaomicron* is orchestrated by a versatile core genetic locus. *Nat. Commun.*
5 **11**, 646 (2020).
- 6 18. V. Raghavan, E. C. Lowe, G. E. 2nd Townsend, D. N. Bolam, E. A. Groisman,
7 Tuning transcription of nutrient utilization genes to catabolic rate promotes
8 growth in a gut bacterium. *Mol. Microbiol.* **93**, 1010–1025 (2014).
- 9 19. B. T. Layden, A. R. Angueira, M. Brodsky, V. Durai, W. L. Jr. Lowe, Short chain
10 fatty acids and their receptors: new metabolic targets. *Transl. Res.* **161**, 131–140
11 (2013).
- 12 20. B. S. Ramakrishna, Role of the gut microbiota in human nutrition and
13 metabolism. *J. Gastroenterol. Hepatol.* **28**, 9–17 (2013).
- 14 21. A. G. Wexler, A. L. Goodman, An insider's perspective: *Bacteroides* as a window
15 into the microbiome. *Nat. Microbiol.* **2**, 17026 (2017).
- 16 22. H. Tan, Q. Zhai, W. Chen, Investigations of *Bacteroides* spp. towards next-
17 generation probiotics. *Food Res. Int.* **116**, 637–644 (2019).
- 18 23. M. P. Vélez, S. C. De Keersmaecker, J. Vanderleyden, Adherence factors of
19 *Lactobacillus* in the human gastrointestinal tract. *FEMS Microbiol. Lett.* **276**,
20 140–148 (2007).
- 21 24. M. E. Johansson, J. M. Larsson, G. C. Hansson, The two mucus layers of colon
22 are organized by the MUC2 mucin, whereas the outer layer is a legislator of host-
23 microbial interactions. *Proc. Natl. Acad. Sci. USA* **108**, 4659–4665 (2011).

- 1 25. C. Rose, A. Parker, B. Jefferson, E. Cartmell, The characterization of feces and
2 urine: A review of the literature to inform advanced treatment technology. *Crit.*
3 *Rev. Environ. Sci. Technol.* **45**, 1827–1879 (2015).
- 4 26. C. B. Whitley, M. D. Ridnour, K. A. Draper, C. M. Dutton, J. P. Neglia,
5 Diagnostic test for mucopolysaccharidosis. I. Direct method for quantifying
6 excessive urinary glycosaminoglycan excretion. *Clin. Chem.* **35**, 374–379 (1989).
- 7 27. R. F. Smith, N. P. Willett, Rapid plate method for screening hyaluronidase and
8 chondroitin sulfatase-producing microorganisms. *Appl. Microbiol.* **16**, 1434–1436
9 (1968).
- 10 28. Q. Shang *et al.*, Structural modulation of gut microbiota by chondroitin sulfate
11 and its oligosaccharide. *Int. J. Biol. Macromol.* **89**, 489–498 (2016).
- 12 29. E. C. Martens *et al.*, Recognition and degradation of plant cell wall
13 polysaccharides by two human gut symbionts. *PLoS Biol.* **9**, e1001221 (2011).
- 14 30. L. E. Tailford, E. H. Crost, D. Kavanaugh, N. Juge, Mucin glycan foraging in the
15 human gut microbiome. *Front. Genet.* **6**, 81 (2015).
- 16 31. R. Lin, W. Liu, M. Piao, H. Zhu, A review of the relationship between the gut
17 microbiota and amino acid metabolism. *Amino Acids* **49**, 2083–2090 (2017).
- 18 32. N. Zmora, *et al.*, Personalized gut mucosal colonization resistance to empiric
19 probiotics is associated with unique host and microbiome features. *Cell* **174**,
20 1388–1405 (2018).
- 21 33. P. Strandwitz *et al.*, GABA-modulating bacteria of the human gut microbiota.
22 *Nat. Microbiol.* **4**, 396–403 (2019).
- 23 34. E. Patterson *et al.*, Gut microbiota, the pharmabiotics they produce and host
24 health. *Proc. Nutr. Soc.* **73**, 477–489 (2014).

- 1 35. P. M. Smith *et al.*, The microbial metabolites, short-chain fatty acids, regulate
2 colonic T_{reg} cell homeostasis. *Science* **341**, 569–573 (2013).
- 3 36. J. H. Cummings, E. W. Pomare, W. J. Branch, C. P. Naylor, G. T. Macfarlane,
4 Short chain fatty acids in human large intestine, portal, hepatic and venous blood.
5 *Gut* **28**, 1221–1227 (1987).
- 6 37. R. S. Miller, L. C. Hoskins, Mucin degradation in human colon ecosystems. Fecal
7 population densities of mucin-degrading bacteria estimated by a “most probable
8 number” method. *Gastroenterology* **81**, 759–65 (1981).
- 9 38. S. Takahashi, J. Tomita, K. Nishioka, T. Hisada, M. Nishijima, Development of a
10 prokaryotic universal primer for simultaneous analysis of Bacteria and Archaea
11 using next-generation sequencing. *PLoS One* **9**, e105592 (2014).
- 12 39. T. Hisada, K. Endoh, K. Kuriki, Inter- and intra-individual variations in seasonal
13 and daily stabilities of the human gut microbiota in Japanese. *Arch. Microbiol.*
14 **197**, 919–34 (2015).
- 15 40. C. Kasai *et al.*, Comparison of the gut microbiota composition between obese and
16 non-obese individuals in a Japanese population, as analyzed by terminal
17 restriction fragment length polymorphism and next-generation sequencing. *BMC*
18 *Gastroenterol.* **15**, 100 (2015).
- 19 41. A. A. Salyers, J. R. Vercellotti, S. E. H. West, T. D. Wilkins, Fermentation of
20 mucin and plant polysaccharides by strains of *Bacteroides* from the human colon.
21 *Appl. Environ. Microbiol.* **33**, 319-322 (1977).

22

23 **Acknowledgements**

1 This work was supported in part by Grants-in-Aid for Scientific Research from the
2 Japan Society for the Promotion of Science (to W.H.), and Research Grant (to W.H.)
3 from Yakult Bio-science Foundation. The authors would like to thank Enago
4 (www.enago.com) for the English language review.

5

6 **Author contributions**

7 W.H. designed the study; M.S., K.K., T.K., R.T., and W.H. performed the experiments;
8 M.S., K.K., T.K., D.W., R.T., and W.H. analyzed the data; M.S., D.W., R.T., and W.H.
9 wrote the manuscript.

10

11 **Competing interests**

12 The authors declare no competing interests.

1 **Figure legends**


2 **Fig. 1.** Human gut microbiota changes in the presence of mucosubstances. (A) Genera
3 (or *Bacteroides* species) frequency profiles of the fecal samples cultivated in the
4 presence of a sole carbon source (Glc, mucin, HA, HP, or CSC). (B) Principal
5 component analysis of the microbiota profiles. The factor loading plot (*left*) and the
6 PCA score plot (*right*) are seen. For marker colors in the PCA score plot, see the
7 symbol legend in (A).

8
9 **Fig. 2.** Degradation of GAGs by *Bacteroides* species. The halo assay for GAG
10 degradation after 7-day incubation with *Bacteroides* species. CSC (*left*) or HP (*right*)
11 was part of the nutrient-rich medium plate.

12
13 **Fig. 3.** Assimilation of GAGs by *Bacteroides* species. (A) Growth level (OD₆₀₀) of
14 *Bacteroides* species in GAG (CSC, HA, or HP) or mucin minimal medium is shown
15 using the heat map with OD₆₀₀ from 0 (white) to 1.5 (black). (B) *Bacteroides faecis*
16 growth-dependent degradation of CSC. Closed and open circles represent OD₆₀₀ values
17 of the culture broth in the presence and absence of *B. faecis*, respectively. Closed and
18 open squares are the CSC concentrations of the culture broth in the presence and
19 absence of *B. faecis*, respectively. Error bars show standard deviations (n = 3).

20
21 **Fig. 4.** Mutually beneficial symbiosis between human and *Bacteroides* species. (A)
22 Secretion of amino acids from GAG (HA) and mucin by *Bacteroides* species. n.d., not
23 detected. (B) Secretion of organic acids from GAG (HA) and mucin by *Bacteroides*
24 species. n.d., not detected. (C) Symbiosis model. The human provides *Bacteroides*

- 1 species in the gut with extracellular mucosubstances (GAGs and mucin) for their
- 2 nutrition and residence place. *Bacteroides* species dominate gut microbiota by
- 3 assimilating host mucosubstances as a carbon/nitrogen source and secrete molecules
- 4 including amino acids and short-chain fatty acids which are essential in human health
- 5 promotion.
- 6



1

2

3 **Fig. 1.** Human gut microbiota changes in the presence of mucosubstances. (A) Genera

4 (or *Bacteroides* species) frequency profiles of the fecal samples cultivated in the


5 presence of a sole carbon source (Glc, mucin, HA, HP, or CSC). (B) Principal

6 component analysis of the microbiota profiles. The factor loading plot (*left*) and the

7 PCA score plot (*right*) are seen. For marker colors in the PCA score plot, see the

8 symbol legend in (A).

9




1

2

3 **Fig. 2.** Degradation of GAGs by *Bacteroides* species. The halo assay for GAG
4 degradation after 7-day incubation with *Bacteroides* species. CSC (*left*) or HP (*right*)
5 was part of the nutrient-rich medium plate.

6



1

2 **Fig. 3.** Assimilation of GAGs by *Bacteroides* species. (A) Growth level (OD₆₀₀) of
3 *Bacteroides* species in GAG (CSC, HA, or HP) or mucin minimal medium is shown
4 using the heat map with OD₆₀₀ from 0 (white) to 1.5 (black). (B) *Bacteroides faecis*
5 growth-dependent degradation of CSC. Closed and open circles represent OD₆₀₀ values
6 of the culture broth in the presence and absence of *B. faecis*, respectively. Closed and
7 open squares are the CSC concentrations of the culture broth in the presence and
8 absence of *B. faecis*, respectively. Error bars show standard deviations (n = 3).


1 **A**

	<i>B. thetaiotaomicron</i>		<i>B. ovatus</i>	
	GAG (μM)	Mucin (μM)	GAG (μM)	Mucin (μM)
Ala	636.8 \pm 32.9	20.8 \pm 7.8	362.8 \pm 188.2	17.2 \pm 14.7
Arg	n.d.	10.7 \pm 4.3	0.5 \pm 0.9	10.5 \pm 4.9
Asn	8.0 \pm 6.4	5.0 \pm 5.6	59.0 \pm 44.1	8.7 \pm 4.8
Asp	n.d.	1.0 \pm 0.3	14.0 \pm 21.8	0.5 \pm 0.6
Cys	n.d.	n.d.	n.d.	n.d.
Gln	0.2 \pm 0.1	0.3 \pm 0.3	2.0 \pm 3.5	0.4 \pm 0.7
Glu	41.9 \pm 20.1	10.6 \pm 4.1	98.3 \pm 58.5	9.0 \pm 2.2
Gly	14.8 \pm 0.7	6.9 \pm 2.4	19.8 \pm 7.3	6.7 \pm 1.0
His	n.d.	2.7 \pm 0.2	3.3 \pm 4.1	2.4 \pm 1.8
Ile	18.5 \pm 3.4	2.1 \pm 0.4	25.7 \pm 13.1	1.6 \pm 0.4
Leu	15.2 \pm 3.1	2.9 \pm 0.3	24.4 \pm 10.5	2.5 \pm 0.3
Lys	n.d.	19.7 \pm 21.4	2.2 \pm 3.4	19.0 \pm 20.3
Met	4.8 \pm 2.0	1.3 \pm 0.4	7.1 \pm 5.5	0.9 \pm 0.4
Phe	5.1 \pm 1.8	2.0 \pm 0.1	6.0 \pm 2.6	1.9 \pm 0.6
Pro	17.1 \pm 3.5	19.4 \pm 4.9	24.1 \pm 10.5	12.6 \pm 14.4
Ser	1.5 \pm 1.3	0.3 \pm 0.4	1.2 \pm 2.6	n.d.
Thr	5.0 \pm 0.5	0.2 \pm 0.2	5.4 \pm 1.8	0.2 \pm 0.1
Trp	n.d.	0.2 \pm 0.3	2.0 \pm 3.5	0.1 \pm 0.2
Tyr	3.5 \pm 0.6	1.1 \pm 0.0	4.4 \pm 2.6	1.2 \pm 0.5
Val	37.3 \pm 0.7	4.1 \pm 1.2	55.0 \pm 14.8	3.4 \pm 1.4
α -ABA	10.0 \pm 0.9	0.9 \pm 0.1	11.8 \pm 0.9	0.6 \pm 0.1
γ -ABA	207.9 \pm 84.4	n.d.	108.3 \pm 84.6	0.1 \pm 0.1
β -Ala	0.9 \pm 0.3	n.d.	2.1 \pm 0.6	n.d.
Citrulline	n.d.	0.2 \pm 0.2	0.2 \pm 0.3	0.1 \pm 0.2
EtOH NH ₂	9.3 \pm 1.1	2.7 \pm 0.9	9.6 \pm 3.1	2.5 \pm 3.2
Hypro	n.d.	n.d.	n.d.	2.1 \pm 3.7
Ornithine	n.d.	0.1 \pm 0.2	n.d.	n.d.
PEA	8.5 \pm 4.0	0.3 \pm 2.4	22.5 \pm 11.4	1.0 \pm 1.9
P-Ser	4.1 \pm 1.2	1.4 \pm 7.6	23.5 \pm 0.7	6.5 \pm 1.7
Sarcosine	n.d.	1.1 \pm 1.9	3.5 \pm 0.9	0.4 \pm 0.7
Urea	n.d.	55.7 \pm 28.3	113.7 \pm 80.7	51.4 \pm 4.8

2 **B**

	<i>B. thetaiotaomicron</i>		<i>B. ovatus</i>		<i>B. faecis</i>	
	GAG (mM)	Mucin (mM)	GAG (mM)	Mucin (mM)	GAG (mM)	Mucin (mM)
Succinic acid	1.08 \pm 0.59	0.05 \pm 0.00	0.86 \pm 0.28	n.d.	0.66 \pm 0.14	0.10 \pm 0.01
Lactic acid	3.63 \pm 1.21	n.d.	2.22 \pm 0.14	n.d.	2.46 \pm 0.90	0.09 \pm 0.01
Formic acid	3.04 \pm 2.10	n.d.	2.82 \pm 2.16	n.d.	2.12 \pm 1.38	0.27 \pm 0.02
Acetic acid	14.93 \pm 1.65	1.16 \pm 0.11	14.00 \pm 1.56	0.69 \pm 0.02	12.49 \pm 2.92	1.66 \pm 0.10
Propionic acid	1.42 \pm 0.55	0.51 \pm 0.04	1.66 \pm 0.47	0.22 \pm 0.01	1.86 \pm 0.61	1.02 \pm 0.06

3 **C**



4

5 **Fig. 4.** Mutually beneficial symbiosis between human and *Bacteroides* species. (A)
6 Secretion of amino acids from GAG (HA) and mucin by *Bacteroides* species. n.d., not
7 detected. (B) Secretion of organic acids from GAG (HA) and mucin by *Bacteroides*
8 species. n.d., not detected. (C) Symbiosis model. The human provides *Bacteroides*
9 species in the gut with extracellular mucosubstances (GAGs and mucin) for their
10 nutrition and residence place. *Bacteroides* species dominate gut microbiota by
11 assimilating host mucosubstances as a carbon/nitrogen source and secrete molecules
12 including amino acids and short-chain fatty acids which are essential in human health
13 promotion.