1 The wheat cytosolic glutamine synthetase *GSL1* modulates N assimilation and
2 spike development by characterizing CRISPR-edited mutants

3 Yazhou Wang^{1,2*}, Wan Teng^{1*#}, Yanpeng Wang¹, Xiang Ouyang^{1,3}, Xue, He¹,
 4 Xueqiang Zhao¹, Caixia Gao¹, Yiping Tong^{1,2}

5
 6 ¹The State Key Laboratory for Plant Cell and Chromosome Engineering, Institute of
 7 Genetics and Developmental Biology, The Innovative Academy of Seed Design,
 8 Chinese Academy of Sciences, Beijing 100101, China.

9 ²University of Chinese Academy of Sciences, Beijing 100049, China.

10 ³State Key Laboratory of Hybrid Rice, Hunan Hybrid Rice Research Center
 11 (HHRRC), Changsha 410125, China

12

13 ***These authors contributed equally to this work.**

14 **# Author for correspondence:** tengwan@genetics.ac.cn

15 **Phone:** +86 10 64807887

16

17 Eight figures, one table

18 Nine supplemental figures, four supplemental tables, one supplemental document

19 6483 words

20 **Highlight**

21 The wheat cytosolic glutamine synthetase *TaGS1.1* is important for N assimilation
 22 and remobilization, and is required for wheat adaptation to low-N stress and spike
 23 development.

24 **Running head :**

25 *TaGS1.1* functions in N use and spike growth

26 **Abstract**

27 Glutamine synthetase (GS) mediates the first step in the assimilation of inorganic
 28 nitrogen (N) into amino acids, however the function of GS encoding genes is not well
 29 understood in wheat (*Triticum aestivum*). We found that the cytosolic *TaGS1.1* was
 30 the major transcribed *GS1* gene and was up-regulated by low-N availability.
 31 CRISPR/Cas9 mediated genome editing was employed to develop two *gs1.1* mutants
 32 with mutated *TaGS1.1-6A*, *-6B*, and *-6D*. Both mutants had lower grains per spike and
 33 grain yield per plant than the wild type under both low-N and high-N conditions in
 34 field experiments. In a hydroponic culture treated with different N resources, the two
 35 mutants was more sensitive to low-N stress than the wild type, but showed similar
 36 sensitivity to high ammonium stress with the wild type. The growth deficiency and
 37 impaired spike development were associated with the imbalance of N metabolites in the
 38 mutant plants. During grain filling, *TaGS1.1* mutation reduced N translocation
 39 efficiency and delayed leaf N loss and grain N filling. Our results suggested that
 40 *TaGS1.1* is important for N assimilation and remobilization, and required for wheat
 41 adaptation to N-limited conditions and spike development.

42 **Key word:** Amino acid homeostasis, Glutamine synthetase, Grain number, Nitrogen
 43 assimilation, Nitrogen translocation, *Triticum aestivum*

44

45 **Introduction**

46 Wheat is one of the most important food crops, it alone provides more than 20% of
 47 the calories and protein for the world's population (Braun *et al.*, 2010; Tilman *et al.*,
 48 2011). Breeding and fertilizer application have greatly increased grain yield, and
 49 further yield increase is facing the challenges of a slow genetic gain in yield in recent
 50 years and efficient use of resources in wheat production (Hawkesford *et al.*, 2013).
 51 Early-season nitrogen (N) fertilizer is known to increase tiller/spike number, grain
 52 number per spike, whereas late-season N mainly increases the kernel weight and grain
 53 protein concentration (Otterson *et al.*, 2008; Peltonen, 1992, 1993). As such, efficient
 54 uptake and assimilation of N is critical for the formation of yield components, and it is
 55 important to understand the roles of N-use related genes in controlling wheat yields.

56 Glutamine synthetase (GS) /glutamate synthase (GOGAT) cycle is the first step
 57 in the assimilation of inorganic N onto carbon (C) skeletons for the production of
 58 glutamine (Gln) and glutamate (Glu). Gln and Glu can then be used to form aspartate
 59 (Asp) and asparagine (Asn) through the activity of aspartate aminotransferase (AAT)
 60 and asparagine synthetase (ASN) (Coruzzi, 2003). These four amino acids are then
 61 converted into all other amino acids and serve as major transport molecules of N
 62 between source and sink tissues (Coruzzi, 2003; Galili *et al.*, 2008). There are two GS
 63 isoforms in plants, the cytosolic isoform GS1 and chloroplastic isoform GS2. GS1
 64 isoenzymes assimilate ammonium derived from primary N uptake and various
 65 internal N recycling pathways (Coruzzi, 2003). GS1 is encoded by a small family of
 66 genes that are well conserved across plant species and are crucial for N assimilation
 67 and N recycling (Bernard and Habash, 2009). The critical roles of *GS1* genes in N
 68 assimilation have been well documented by analyzing N metabolites in the *GS1*
 69 mutants of Arabidopsis (*Arabidopsis thaliana*) (Konishi *et al.*, 2017; Konishi *et al.*,
 70 2018; Lothier *et al.*, 2011; Moison *et al.*, 2018), rice (*Oryza sativa*) (Funayama *et al.*,
 71 2013; Kusano *et al.*, 2020; Kusano *et al.*, 2011), and maize (*Zea mays*) (Canas *et al.*,
 72 2010; Martin *et al.*, 2006). The crucial roles of *GS1* genes in N remobilization also
 73 have been demonstrated by characterizing the *GS1* deficient mutants (Guan *et al.*,
 74 2015; Kamachi *et al.*, 1991; Masclaux-Daubresse *et al.*, 2010; Moison *et al.*, 2018;
 75 Yamaya and Kusano, 2014). *GS1* genes play non-overlapping roles in N use and plant
 76 growth. For example, the rice *OsGS1.2* is responsible for the primary assimilation of
 77 ammonium in roots, while *OsGS1.1* is important in the process of N remobilization in

senescing organs (Funayama *et al.*, 2013; Yamaya and Kusano, 2014). In line with their physiological role in N use, disruption of *OsGS1.2* greatly reduces active tiller number and hence panicle number at harvest; whereas loss-of-function mutation in *OsGS1.1* greatly inhibits rice growth, grain number per panicle, grain size and grain filling (Yamaya and Kusano, 2014). *OsGS1.1* and *OsGS1.2* are unable to compensate for the individual function of another (Tabuchi *et al.*, 2005; Yamaya and Kusano, 2014). The maize GS1 genes *ZmGln1.3* and *1.4* are specifically involved in the control of kernel number and kernel size, respectively (Martin, et al., 2006).

Physiological correlation and QTL mapping have revealed the importance of GS genes in N use and yield formation in wheat. GS activity is positively correlated with total N, chlorophyll, soluble protein, ammonium, and amino acids in flag leaves (Kichey *et al.*, 2006; Kichey *et al.*, 2007). N remobilization contributes to approximately 70% of grain N, and this contribution varies among wheat cultivars and closely related to leaf GS activity (Kichey *et al.*, 2006; Zhang *et al.*, 2017c). QTL mapping also has detected the co-localization between QTL for GS activity and N use and yield-related traits (Fontaine *et al.*, 2009; Habash *et al.*, 2007; Li *et al.*, 2015). As such, GS activity can be served as a marker to predict the N status of wheat, and GS1 genes are considered valuable in breeding with improved N use efficiency (NUE) and yield. For example, overexpression of a GS1 gene in wheat increased root growth, N uptake, and grain yield (Habash *et al.*, 2001). However, overexpression studies using GS1 to increase NUE have not yielded consistent results. To develop future strategies for the use of GS1 in increasing NUE, it is required to understand the pivotal role of GS1 in the maintenance of essential N flows and internal N sensing during critical stages of plant development (Thomsen *et al.*, 2014). Therefore, understanding the function of GS1 genes in mediating N use and yield performance will facilitate the use of GS1 genes in wheat breeding.

Common wheat has three GS1 genes, one GS2 gene, one NADH-GOGAT gene, and one Fd-GOGAT gene in each of the sub-genomes. Our previous studies have shown the essential roles of GS2 and NADH-GOGAT in mediating N use and plant growth in wheat (Hu *et al.*, 2018; Yang *et al.*, 2019; Zhang *et al.*, 2017a). Here we developed *gs1.1* mutants with mutations in *TaGS1.1-6A*, *-6B*, and *-6D* through genome editing. Investigation of the N use and growth-related traits revealed that *TaGS1.1* is critical for N assimilation, N remobilization, and adaptation to low-N

environments. This mutant has reduced grain number per spike and grain yield under both low-N and high-N conditions.

Materials and Methods

Plant materials

The winter wheat variety KN199 was used in this study. KN199 was commercially released in 2006, and was used to isolate *TaGS1.1* sequences and develop the *gs1.1* mutants.

Genome editing of TaGS1.1

One sgRNA target for *TaGS1.1* was designed on the conserved domains of all three genomes of wheat variety KN199. The activities of the sgRNA was evaluated by co-transforming the pJIT163-Ubi-Cas9 (Wang *et al.*, 2014) and TaU6-sgRNA (Shan *et al.*, 2013) plasmids into wheat protoplasts. Wheat protoplasts were isolated and transformed as previously described (Shan *et al.*, 2014). The DNAs of plasmids pJIT163-Ubi-Cas9 and pTaU6-sgRNA were simultaneously delivered into immature embryos of KN199 via particle bombardment, as previously described (Zhang *et al.*, 2017b). After bombardment, the embryos were cultured for plantlet regeneration on medium without selective agent. PCR-RE (PCR-restriction enzyme) assays and Sanger sequencing were used to identify wheat mutants in target regions, as described previously (Wang *et al.*, 2014).

Hydroponic culture

The hydroponic culture was conducted in a greenhouse under the following conditions: 20°C ± 1°C, 50% to 70% relative humidity, 300 μmol photons m⁻² s⁻¹ and a 16-h-day/8-h-night cycle. The germinated seedlings were transferred to the nutrient solution which was described previously (Ren *et al.*, 2012). Three treatments were used, the standard-N (SN), low-N (LN), and ammonium-N (AN) treatments which contained 1.0 mM NH₄NO₃, 0.1 mM NH₄NO₃, and 4 mM NH₄⁺, respectively. The nutrient solutions were refreshed every two days.

Field experiment

The field experiments were carried out in the 2017-2018 and 2018-2019 growing seasons in Hebei Province, China. Both field experiments consisted of two N

141 conditions, each of which had four biological replications. The high-N treatment was
 142 applied 18 g m⁻² N as urea, with 12 g m⁻² applied before sowing and 6 g m⁻² applied at
 143 the stem elongation stage. The low-N treatment was applied 3.45 g m⁻² N before
 144 sowing. The two N treatments were applied 6 g m⁻² P as calcium superphosphate
 145 before sowing. The seeds were sown in a 2-m-long row with a sowing density of 89
 146 seeds per m². In the 2017-2018 growing season, the seeds were sown in two rows for
 147 each genotype in each replicate, and four biological replications were set for each
 148 sampling time. At stem elongation, anthesis, 14 days post-anthesis (DPA), and 28
 149 DPA, the aerial parts of five representative plants were collected for dry weight and N
 150 analysis. At maturity, the aerial parts of 20 representative plants in each replication
 151 were harvested for dry weight, agronomic traits, and total N analysis. In the
 152 2018-2019 growing season, the seeds were sown in four rows for each genotype in
 153 each of the four biological replications. At maturity, at least 25 representative plants
 154 were harvested in each replication for the measurement of dry weight and agronomic
 155 traits. The photosynthetic parameters were measured at 14 DPA by using LI-6400
 156 Portable Photosynthesis System (LI-COR Biosciences, Lincoln, Nebraska USA). Five
 157 flag leaves in each replicate were measured in 9-11 am.

158 *Analysis of N metabolites*

159 The fresh samples stored at -80 °C were homogenized for the measurement of free
 160 nitrate and ammonium. The nitrate concentrations in plant tissues were quantified
 161 according to the methods described by Cataldo *et al.* (1975), and the free ammonium
 162 in plant tissues was determined by Berthelot reaction (Husted *et al.*, 2000). To
 163 measure free amino acids, the fresh samples were frozen dried and ground.
 164 Ultrasound-assisted extraction was performed for 30 min by adding 1 ml ultra-pure
 165 water to 20 mg of ground powder, and the mixture was centrifuged at 10, 000 g for 10
 166 min at 4 °C. The amino acids in the supernatant were derivatized before injection, and
 167 then the reaction products were separated and detected by HPLC with a BEH C₁₈
 168 sorbent (Waters Alliance e2695, Waters Corporation, Milford, MA). The dried
 169 samples were ground for total N analysis by the automated Kjeldahl method (Kjeltec
 170 TM 8400, Foss Analytical A/S, Demark).

171 *Quantitative Real-time PCR*

172 Total RNA extraction and real-time quantitative reverse transcription PCR (qRT-PCR)
173 were performed according to the methods of Yang et al. (2019). The primers for
174 qRT-PCR were detailed in [Supplemental Table S1](#). The gene expression levels were
175 normalized to the internal control of *TaActin*.

176 ***Western blot and GS activity assay***

177 The fresh plant samples stored at -80 °C were ground to a fine powder under liquid N
178 and then homogenized in an extraction buffer containing 50 mM Tris-HCl (pH 8.0), 2
179 mM MgCl₂, 2 mM DTT, and 0.4 M sucrose. The homogenate was centrifuged at 10,
180 000 g for 20 min two times at 4 °C. The supernatant fraction was used for western
181 blot and GS activity assay. The total protein concentration was determined by a
182 Bradford assay. Western blot analysis was performed using an antibody raised against
183 GS1 protein in rabbits (Abmart, Shanghai, China). The GS activity was determined by
184 using Glutamine Synthetase Detection Kit A047 (Nanjing Jiancheng Biotechnology).

185 ***Statistical analysis of data***

186 Statistically significant differences using SPSS17.0 for Windows (SPSS) were
187 computed based on Student's *t*-test.

188 **Results**

189 ***The TaGS1 genes differentially respond to N availability***

190 The published *GS* sequences in wheat were used to blast the reference sequence of
191 Chinese spring in the Ensembl Plants database (<http://plants.ensembl.org/index.html>).
192 We identified three *GS1* genes and one *GS2* gene in each sub-genome, and the *GS1*
193 genes were named according to their orthologous relation to the *GS1* genes rice
194 ([Supplemental Figure S1](#)). The Gene IDs and former names of *GS* genes were
195 presented in [Supplemental Table S2](#).

196 Gene expression analysis revealed that *TaGS1.1* had much higher transcript
197 abundance than *TaGS1.2* and *TaGS1.3* in roots and shoots of the wheat variety
198 KN199 grown under 1.0 mM NH₄NO₃ (SN), 0.1 mM NH₄NO₃ (LN) and 4.0 mM
199 NH₄⁺ (AN) conditions ([Fig. 1](#)). Compared with SN treatment, *TaGS1.1* was
200 up-regulated in roots by LN treatment, and down-regulated in shoots by AN treatment
201 ([Fig. 1A](#)); *TaGS1.2* was down-regulated by LN and AN treatments in roots, and by

LN treatment in shoots (Fig. 1B); *TaGSI.3* was down-regulated in both shoots and roots by LN treatment (Fig. 1C).

We then quantified the expression of *TaGSI* genes in different organs of KN199 grown under field conditions at 20 days post-anthesis (DPA). *TaGSI.1* was most strongly transcribed in leaf blades and sheaths, followed by roots, and the lowest in seeds; *TaGSI.2* was mainly expressed in leaf blades and sheaths, glumes and rachises; whereas *TaGSI.3* was presented at a very low level in all the investigated organs (Supplemental Figure S2A). In flag leaf blades, the expression of *TaGSI.1* displayed a substantial decline from stem elongation (Zadoks growth scale Z37) to 28 DPA, while that of *TaGSI.2* sharply increased from 14 DPA to 28 DPA (Supplemental Figure S2B).

Development of *TaGSI.1* knockout mutants

Considering the relatively high transcription abundance of *TaGSI.1* and the potential role of *TaGSI.1* in wheat adaptation to low-N availability, we then created *TaGSI.1* knockout mutants of KN199 by using transgenic free CRISPR/Cas9 mediated genome editing. We cloned the three homoeologous *TaGSI.1* genes (*GSI.1-6A*, *-6B*, and *-6D*) in KN199. A single-guide RNA (sgRNA) that matched perfectly with *TaGSI.1-6A* and *-6D* but had one mismatch with *TaGSI.1-6B* was designed (Fig. 2) in the fifth exon. We used the sgRNA sequence to blast with the Chinese spring reference genomic sequence, and the blast result showed that this sgRNA sequence could avoid off-target (Supplementary Document S1). The genome-editing obtained a mutant. In this mutant, *TaGSI.1-6A* and *-6D* each had homozygous mutation with a 12-bp deletion and a 4-bp deletion respectively, and *TaGSI.1-6B* had heterozygous mutations with a 4-bp deletion in mutation 1 and a 9-bp deletion in mutation 2 (Fig. 2). We grew the T₂ transgenic plants and WT under high-N and low-N conditions in the 2016-2017 growing season, and identified two kinds of homozygous mutants by PCR-RE (PCR-restriction enzyme) assays and Sanger sequencing. The *gsl.1-1* and *-2* had homozygous mutation 1 and 2 in *TaGSI.1-6B* respectively, and both mutants had homozygous mutations in *TaGSI.1-6A* and *-6B*. Preliminary investigation of agronomic traits showed that both mutants had lower plant height (PH), grain yield per plant (GY) and spike grain weight (SGW) than WT under both high-N and low-N conditions (Supplementary Figure S3A, B, D), but WT and the mutants showed similar spike number per plant (SNPP, Supplementary Figure S3C).

235

236 ***TaGS1.1 mutation decreases grain number and grain yield***

237 We evaluated the agronomic traits in two consecutive growing seasons from
238 2017-2019 under low-N and high-N conditions. The *gs1.1* mutants headed about two
239 days later than WT ([Supplemental Figure S4](#)). The WT and mutant plants did not
240 show a visible difference in leaf and spike color before 14 days post-anthesis (DPA,
241 [Fig. 3A, C](#)), but the leaves and spikes of the mutant plants looked greener than those
242 of WT after 25 DPA ([Fig. 3B, D](#)). The spikes of the mutant plants looked smaller than
243 those of WT ([Fig. 3C, D](#)). Compared with WT in the 2017-2018 growing season, both
244 of the *gs1.1* mutants had a lower GY under high-N and low-N conditions ([Fig. 3F](#)).
245 Investigation of yield components showed that the two *gs1.1* mutants had
246 significantly lower spike grain number (SGN) and spike grain weight (SGW) than
247 WT ([Fig. 3I, J](#)), but had similar SNPP and 1000-grain weight with WT ([Fig. 3G, H](#)).
248 We investigated the morphological traits of the main spikes and found that both of the
249 *gs1.1* mutant has shorter spike length and fewer spikelet number than WT ([Fig. 3K,](#)
250 [L](#)). We grew the WT and *gs1.1-1* mutant again in the 2018-2019 growing season, the
251 *gs1.1-1* mutant plants exhibited very similar phenotypes in agronomic traits with
252 those in the 2017-2018 growing season ([Supplemental Table S3](#)). These two growing
253 seasons clearly showed that the lower GY of the mutant plants mainly resulted from a
254 smaller spike size, which was reflected by spike length, spikelet number, and SGN.

255 We monitored the dry weight (DW) of different aerial organs in the main culm of
256 the WT and *gs1.1-1* mutant plants at stem elongation, anthesis, 14 DPA, 28DPA, and
257 maturity in the 2017-2018 growing season. All the investigated organs and the whole
258 culm displayed lower DW in the mutant than in WT under both high-N and low-N
259 conditions ([Supplemental Figure S5](#)). The relative difference between the WT and
260 *gs1.1* mutant was evaluated by using the DW ratio of the mutant over WT
261 (*gs1.1-1*/WT ratio). The lowest *gs1.1*/WT DW ratio was observed for the spike, the
262 youngest developing organ at stem elongation ([Supplemental Figure S5](#)). At stem
263 elongation, the *gs1.1*/WT DW ratio increased with the organ age ([Table 1](#)). These
264 results suggested that the inhibitory effect of disrupting *TaGS1.1* on organ DW
265 depended on the organ development stage, the younger the organ, the stronger the
266 inhibition.

267 ***Knockout of TaGS1.1 causes root and shoot growth deficiency in an N*** 268 ***resource-dependent manner***

269 Considering the potential role of *TaGS1.1* in mediating N assimilation, we then
270 investigate how *TaGS1.1* mutation affected root and shoot growth of wheat seedlings
271 supplied with different N resources. When the seedling were grown under SN, LN
272 and AN conditions, the *gs1.1-1* and *-2* mutants displayed shorter PH and leaf length,
273 and lower shoot dry weight (SDW), root dry weight (RDW) and root/shoot ratio (DW
274 R/S ration) than WT, in an N resource-dependent manner (Fig. 4). Among the three N
275 conditions, the mutants showed stronger phenotypes in SDW, RDW and DW R/S
276 ratio under LN conditions than under SN and AN conditions, indicating the role of
277 *TaGS1.1* in wheat adaptation to low-N availability.

279 ***Knockout of TaGS1.1 reduces GS1 protein abundance and GS activity***

280 To understand the mechanism underlying the growth deficiency of the *gs1.1* mutants,
281 we characterized the GS activity, N assimilation and remobilization of the *gs1.1-1*
282 mutant plants. The seedlings harvested in the hydroponic culture were used to
283 measure these traits. Analysis of gene expression found that the *gs1.1-1* mutant
284 exhibited much lower expression of *TaGS1.1-6B* and *-6D* in roots and shoots than WT
285 under both SN and LN conditions; whereas it showed a comparable expression of
286 *TaGS1.1-6A* with WT (Fig. 5A). It has been reported that a single bp deletion induced
287 frameshift and premature stop prevents mRNA accumulation and consequently results
288 in a low mRNA level of the mutated gene in tobacco (Voelker *et al.*, 1990). The
289 *gs1.1-1* mutant had a much lower GS1 protein level in roots and shoots than WT
290 under SN and LN conditions and lost the LN-induced GS1 protein increase which was
291 observed in WT (Fig. 5B, C). These results suggested that TaGS1.1 was the major
292 GS1 isoform in roots and shoots under both SN and LN conditions at the protein level
293 and majorly contributed to the LN-induced GS1 protein increase. Compared with WT,
294 the total GS activities of the mutant were significantly reduced in shoots under LN
295 conditions, and in roots under SN and LN conditions, whereas those in shoots and
296 roots under AN conditions were not significantly affected (Fig. 5D). These results
297 suggested that the mutant seedlings were deficient in GS activity in the presence of
298 nitrate.

299 In the hydroponic culture, the shoots of the *gs1.1-1* mutant plants had
300 significantly lower expression of *TaNR1* (nitrate reductase, [Supplemental Figure S6A](#)),
301 *TaNiR2* (nitrite reductase, [Supplemental Figure S6B](#)), *TaGS1.2* ([Supplemental Figure](#)
302 [S6C](#)), *TaNADH-GOGAT* ([Supplemental Figure S6E](#)) and *TaFd-GOGAT*
303 ([Supplemental Figure S6F](#)) than those of WT, depending on N supply level. In roots,
304 the mutant transcribed significantly fewer transcripts of *TaGS1.2* under SN conditions
305 ([Supplemental Figure S6C](#)) and *TaASN1* ([Supplemental Figure S6G](#)) than WT under
306 both SN and LN conditions. The *gs1.1-1* mutant and WT had similar mRNA levels of
307 *TaGS2* ([Supplemental Figure S6D](#)). We also analyzed the expression of *GS* and
308 *GOGAT* genes in flag leaves in the field experiment and found that knockout of
309 *TaGS1.1* reduced the expression of *TaGS1.1* and *TaFd-GOGAT*, but increased the
310 expression of *TaNADH-GOGAT* ([Supplemental Figure S7](#)).

311

312 ***The gs1.1-1 mutant displays an imbalance of N metabolites***

313 To understand the roles of *TaGS1.1* in mediating N assimilation, we measured N
314 metabolites in roots and shoots of wheat seedlings exposed to different N recourses as
315 described in [Fig. 4](#). For total N concentration (NC), the WT and *gs1.1-1* mutant only
316 differed in root NC under LN conditions, with the mutant having the lower root NC
317 ([Fig. 6A](#)). Compared with WT, the mutant accumulated much higher free NO_3^- in
318 shoots under SN conditions, and roots under SN and LN conditions ([Fig. 6B](#)); it also
319 contained significantly higher free NH_4^+ in shoots under LN conditions, and in roots
320 under LN and AN conditions ([Fig. 6C](#)).

321 The free amino acids were detected in both roots and shoots ([Supplemental Table](#)
322 [S4](#)). The concentrations of total free amino acids in shoots and roots of the mutant
323 were significantly reduced under SN and LN conditions but not under AN conditions,
324 as compared to WT ([Fig. 6D](#)). Under SN conditions, the mutant had significantly
325 lower Asn in shoots, and lower Gln, Asp, and Asn level in roots than WT, but had
326 significantly higher Glu level in shoots than WT; under LN conditions, the mutant had
327 significantly lower Glu, Gln and Asp level in shoots, and lower Glu level in roots than
328 WT, but had significantly higher Asp level in shoots than WT; under AN conditions,
329 the mutant had significantly lower Glu level in roots than WT ([Fig. 6E-H](#)). Besides
330 these four amino acids, the levels of many other amino acids were significantly

changed in the mutant in a tissue- and N resource-dependent manner, as compared to WT (Supplemental Table S4).

We also measured the free amino acids in the young spikes at stem elongation and seeds at maturity in the field experiment in the 2017-2018 growing season. In the young spikes at stem elongation stage, the concentrations of Asn, Gln, arginine (Arg), alanine (Ala) and proline (Pro) were significantly higher in the *gs1.1-1* mutant than in WT (Supplemental Figure S8A), those of Glu and Asp (Supplemental Figure S8A) and other amino acids (data not shown) did not show significant difference between the mutant and WT. In the seeds at maturity, the *gs1.1-1* mutant had significantly higher levels of Gln and Asn than WT, but had significantly lower level of Asp than WT (Supplemental Figure S8B).

Knockout of TaGS1.1 decreases flag leaf photosynthetic rate but delays flag leaf senescence

Since the *gs1.1-1* mutant exhibited growth deficiency, we then investigated if the knockout of *TaGS1.1* affected photosynthetic capacity flag leaves at 14 DPA in the 2017-2018 growing season. The mutant had a significantly lower net photosynthetic rate (P_n , Fig. 7A), but significantly higher stomatal conductance (G_s , Fig. 7B), intercellular CO₂ concentration (C_i , Fig. 7C); transpiration rate (Tr , Fig. 7D) than WT under both high-N and low-N conditions.

We also measured total N concentration (NC) in leaf blades, leaf sheaths, stems, spikes (with seeds removed), and grains at stem elongation, anthesis, 14 DPA, 28 DPA, and maturity. The strongest phenotype was observed in flag leaf blades (Supplemental Figure S9C and D), in which the *gs1.1-1* mutant had much higher NC than WT under both high-N and low-N conditions at 28 DPA, but had similar NC with WT at stem elongation, anthesis and maturity stage (Fig. 7E). The dynamic changes in SPAD values clearly showed that the flag leaf senescence was delayed in the mutant as compared to WT (Fig. 7F). These results indicated that *gs1.1* mutation delayed N loss in the flag blades and thereby flag leaf senescence. Similarly, the mutant also displayed a delay of N loss in the stems and middle leaf blades (top 2nd and 3rd leaves) as compared to WT (Supplemental Figure S9A, D). The significantly higher NC in the flag leaf blades and middle leaf blades of the *gs1.1-1* mutant than WT was observed at 14 and 28 DPA under low-N conditions, but was only observed

at 28 DPA under high-N conditions (Supplemental Figure S9C, D). The mutant had significantly higher grain NC than WT under both high-N and low-N conditions at maturity (Supplemental Figure S9G).

The gs1.1-1 mutant has an altered translocation efficiency of N and dry matter during grain filling

N translocation efficiency (NTE) and dry matter translocation efficiency (DMTE) of aerial organs in the main culm were calculated to reflect N and dry matter remobilization during grain filling by using the data in Supplemental Figure S5 and S9. The *gs1.1-1* mutant had significantly lower NTE in stems, spikes, leaf blades, and sheathes than WT, depending on N supply level (Fig. 8A). The *gs1.1-1* mutant exhibited significantly lower DMTE in stems, leaf sheathes, middle and bottom leaf blades than WT under both high-N and low-N conditions (Fig. 8C). We also calculated N harvest index (NHI) and harvest (HI) and found that the *gs1.1-1* mutant had lower NHI and HI than WT at 14 DPA and 28 DPA, but had similar NHI and HI with WT at maturity under both high-N and low-N conditions (Fig. 8B, D). These results suggested that disruption of *TaGS1.1* delayed grain N and dry matter filling, but not the final NHI and HI.

DISCUSSION

TaGS1.1 is the major GS1 isoform

The expression levels of the three *TaGS1* depended on N availability, organs, and developmental stages (Fig. 1, Supplemental Figure S5), suggesting their non-overlapping roles in N use. The sharp reduction of GS1 protein abundances in roots and shoots of the *gs1.1-1* mutant grown under SN and LN conditions indicated that TaGS1.1 is the major GS1 isoform in both roots and shoots in the presentation of nitrate (Fig. 5C). In line with this result, *TaGS1.1* was more abundantly transcribed than *TaGS1.2* and *TaGS1.3* in shoots and roots at the seedling stage (Fig. 1). A recent study reported that TaGS1.1 has much higher protein abundance than TaGS1.2 and 1.3 in roots and aerial parts (Wei et al., 2019). TaGS1.1 also majorly contributed to the increased GS1 protein level in roots and shoots by LN treatment (Fig. 5C), this result was associated with the up-regulation of *TaGS1.1* by LN treatment, as compared to SN treatment (Fig. 1A).

Although there was a sharp reduction in the GS1 protein level in the *gs1.1-1* mutant, the GS activity was moderately decreased in the mutant under SN and LN conditions, as compared to WT (Fig. 5D). Firstly, a possible explanation for this phenomenon was that the GS activity was measured as the sum of the GS1 and GS2 activities. Secondly, the *gs1.1-1* mutant had four amino acids deletion in TaGS1.1-6A and a frameshift mutation in TaGS1.1-6B and -6D (Fig. 2), the mutated TaGS1.1-6A may still have GS activity. Finally, TaGS1.2 and 1.3 had much higher $V_{\max}$ for Glu than TaGS1.1 (Wei et al., 2019), the higher $V_{\max}$ of TaGS1.2 and 1.3 may compensate the effect of low GS1 abundance on GS activity in the *gs1.1-1* mutant. Under AN conditions, the WT and *gs1.1-1* mutant had similar GS activity in both shoots and roots (Fig. 5D). This result was possibly related to the fact that a high level of ammonium greatly reduced TaGS1.1 protein abundances in shoots and roots (Wei et al., 2019). When the external NH_4^+ was greater than 2 mM, TaGS1.1 subunit abundance is difficultly detected in shoots and presents at a much lower level than TaGS1.2 in roots (Wei et al., 2019).

410 **TaGS1.1 is important for NH_4^+ assimilation**

Measurement of N metabolites revealed the crucial role of *TaGS1.1* in maintenance of internal inorganic N and amino acid homeostasis in wheat plants grown under different N conditions. The results from the seedlings grown in the hydroponic culture, and the young spikes and mature seeds from the field experiment showed that the effects of *TaGS1.1* mutation on N metabolites depended on tissue type and N resource. In the hydroponic culture treated with SN, LN and AN, the *gs1.1-1* mutant had a reduced level of many free amino acids and total free amino acids in shoots and roots under SN and LN conditions compared with WT (Fig. 6, Supplemental Table S4), suggesting the essential role of *TaGS1.1* for NH_4^+ assimilation in shoots and roots in the presence of nitrate. However, the *gs1.1-1* mutant had an increased free NH_4^+ level in both shoots and roots under LN conditions but not under SN conditions compared with WT (Fig. 6C), indicating that *TaGS1.1* was important for NH_4^+ assimilation and homeostasis in wheat plants under LN conditions. The importance of *TaGS1.1* for NH_4^+ assimilation under LN conditions was in line with the fact that *TaGS1.1* was up-regulated in roots by LN treatment (Fig. 1A) and majorly contributed to the LN-induced GS1 protein increase in shoots and roots (Fig. 2D). This importance was further supported by a recent study in which TaGS1.1 is found to be expressed in root

epidermis cells and displays a higher affinity for Glu and hydroxylamine than TaGS1.2 and TaGS1.3 (Wei et al., 2019). *OsGS1.1*, the rice orthologue of wheat *TaGS1.1* has also been reported its importance in N assimilation under low-ammonium conditions (Ishiyama *et al.*, 2004). Under AN conditions, the *gs1.1-1* mutant had a higher free NH_4^+ level and a lower level of Glu and seven other amino acids in roots than WT (Fig. 6C, E; Supplemental Table S4); however, only Ala level was significantly changed in the investigated N metabolites in shoots of the mutant compared with WT (Supplemental Table S4). These results suggested a role of *TaGS1.1* in NH_4^+ assimilation in roots, but not (or a limited role, if any) in shoots when NH_4^+ was used as the sole N resource. In contrast to the decreased Gln and Asn levels in shoots and roots of the hydroponically grown *gs1.1-1* seedlings in the presence of nitrate (SN and LN treatments), the field grown *gs1.1-1* plants had higher Gln and Asn in the young spikes and mature seeds than WT (Supplemental Figure S8A). As such, the role of TaGS1.1 in maintaining internal amino acid homeostasis depended on organ type or development stage.

The imbalance of N metabolites in the *gs1.1-1* mutant seemed the combination results of TaGS1.1 deficiency and the down-regulated expression of other genes involved in N assimilation (Supplemental Figure S6). Compared with the shoots of WT, for example, the increased free NO_3^- level was possibly associated with the down-regulated *TaNRI* in the *gs1.1-1* mutant under SN conditions (Supplemental Figure S6A). Knockout of *TaGS1.1* more substantially reduced the total free amino acid level in shoots and roots under SN conditions than under LN conditions compared with WT (Fig. 6D), this was possibly associated with the fact that the transcripts of *TaGS1.2* in shoots and roots of the mutant were significantly reduced under SN condition but not under LN conditions, as compared to WT (Supplemental Figure S6C). Among the detected free amino acids, the Asn level in roots under SN conditions was most sharply reduced (the lowest *gs1.1-1*/KN199 ratio, 0.19) by disrupting *TaGS1.1* (Supplemental Table S4). As ASN catalyzes the synthesis of Asn and Glu from Asp and Gln, the reduced *TaASN1* transcripts together with the reduced Asp and Gln level might contribute to the sharp reduction of Asn in the *gs1.1-1* roots under SN conditions (Fig. 6F, G; Supplemental Figure S6G).

The internal pools of amino acids within plants have been suggested to indicate N status of a plant, and are served as a signal to regulate gene expression (Miller *et al.*, 2008). Compared with SN treatment, LN treatment had a lower level in total NC and

most of the investigated N metabolites in both shoots and roots of WT (Fig. 6, Supplemental Table S4), indicating an N-deficient status of WT under LN conditions. The *gs1.1-1* mutant had a reduced level of many free amino acids and total free amino acids in shoots and roots under both SN and LN conditions (Fig. 6D, Supplemental Table S4). As such, the mutant might have a deficient N status under SN conditions and a more deficient N status under LN conditions compared with WT, and thereby displayed growth deficiency. In line with this claim, the responses of several N-assimilation genes to N availability were changed in the mutant. In shoots, the expression of *TaNRI*, *TaNiR2*, and *TaGS1.2* in WT was up-regulated by SN treatment as compared to LN treatment; however, these genes had significantly lower expression in the mutant than in WT under SN conditions (Supplemental Figure S6A-C). In roots, the expression of *TaGS1.2* and *TaASN1* in WT was up-regulated by SN treatment as compared to LN treatment; however, disrupting *TaGS1.2* impaired the SN up-regulated *TaGS1.1* expression, and significantly reduced *TaASN1* expression under SN and LN conditions (Supplemental Figure S6C, G).

477 ***TaGS1.1* functions in N remobilization**

Cytosolic GS1 isoforms are known to positively mediate N remobilization during leaf senescence (Moison *et al.*, 2018; Yamaya and Kusano, 2014). Our current study showed that knockout of *TaGS1.1* in the *gs1.1-1* mutant reduced N and dry matter translocation efficiency during grain filling (Fig. 8A, C), and delayed N loss and senescence of flag leaves (Fig. 7E, F). However, disruption of *TaGS1.1* did not significantly affect the final NHI and HI (Fig. 8B, D) and total N concentrations in the vegetative organs at maturity (Supplemental Figure S9). These phenomena were possibly associated with the dynamic changes of *TaGS1* transcripts. In flag leaves, the mRNA levels of *TaGS1.1* and *1.2* sharply decreased and increased after 14 DPA, respectively, and *TaGS1.1* had a higher mRNA level than *TaGS1.2* before 14 DPA, but a much lower level than *TaGS1.2* after 14 DPA (Supplemental Figure S2B). These results indicated that *TaGS1.1* may mainly function during relatively earlier grain filling, while *TaGS1.2* may mainly function during relatively later grain filling in modulating N and dry matter remobilization. Since knockout of *TaGS1.1* did not significantly alter the expression of *TaGS1.2* in flag leaves (Supplemental Figure S7B), it can be assumed that the functional *TaGS1.2* may partially compensate for the adverse effects of *TaGS1.1* deficiency on N and dry matter remobilization.

495 ***TaGS1.1 is required for wheat tolerance to low-N stress***

496 The three *GSI* genes showed a differential response to N availability (Fig. 1),
 497 suggesting their different roles in wheat adaptation to a fluctuating nutrient
 498 environment. In the hydroponic culture, the *gs1.1-1* and -2 mutant seedlings were
 499 more sensitive to low-N stress than WT (Fig. 4). At stem elongation, the young spike
 500 growth of the *gs1.1-1* mutant was more sensitive to low-N treatment than that of WT
 501 under field conditions (Table 1). The role of *TaGS1.1* in wheat adaptation to low-N
 502 stress was supported by its importance for NH_4^+ assimilation under N-limited
 503 conditions, which has been discussed in the context. The amino acids Glu, Gln, Asp,
 504 and Asn are the precursors of other amino acids in N assimilation, and Glu and Gln
 505 may serve as signals of organic N status (Gutierrez *et al.*, 2008). The reduced low-N
 506 tolerance of shoot growth in the *gs1.1-1* mutant was possibly associated with the fact
 507 that the mutant had a reduced level of Glu, Gln, and Asp in shoots under LN
 508 conditions but not under SN conditions, as compared to WT (Fig. 6E-G).

509 A high level of ammonium in the growth media is known to inhibit plant growth,
 510 as has been shown in our current study (Fig. 4). The WT and *gs1.1* plants did not
 511 differ in their sensitivity to high ammonium stress under our experimental conditions
 512 (Fig. 4). The underlying reason might be because that *TaGS1.1* had a limited role in
 513 NH_4^+ assimilation in shoots under AN conditions. The wheat *GS1.1* is orthologous to
 514 the rice *GS1.1/Gln1.1*. However, the rice *gs1.1* mutant exhibits a severe decrease in
 515 shoot growth and an imbalance in the levels of sugars, amino acids, and metabolites
 516 when ammonium is used as the sole N resource (Kusano *et al.*, 2011). Since nitrate
 517 and ammonium are, respectively, the major N resources for wheat and rice, it can be
 518 assumed that *TaGS1.1* and *OsGS1.1* are evolutionally adapted to nitrate and
 519 ammonium recourses, respectively.

520 ***TaGS1.1 Is Required for Spike Development***

521 We observed plant growth deficiency in the two *gs1.1* mutants at the seedling stage in
 522 the hydroponic culture (Fig. 4) and in the field experiments (Fig. 3, Supplemental
 523 Figure S3). In line with these results, the *gs1.1-1* mutant exhibited much lower *Pn* in
 524 flag leaves than WT under both high-N and low-N conditions at 14 DPA (Fig. 7A).
 525 The impaired plant growth and photosynthetic capacity were possibly caused by a
 526 deficiency in free amino acids and the imbalance of N metabolites but not by total N

deficiency in the mutant. In fact, the mutant had similar with, or even higher N concentrations in aerial parts than did WT (Supplemental Figure S9).

The field experiments showed that the lack of *TaGSI.1* reduced GY by inhibiting spike development, as both of the *gs1.1-1* and *-2* mutants had a shorter spike length, a fewer spikelets and grains per spike, and consequently a lower spike grain weight compared with WT (Fig. 3, Supplemental Figure S3, Supplemental Table S3). Monitoring aerial organ growth of the *gs1.1-1* and WT plants from stem elongation to mature revealed that the growth deficiency in the *gs1.1-1* mutant was most obvious in the young spikes at stem elongation stage (Table 1). This result together with the later heading date in the mutant (Supplemental Figure S4) indicated a delay in spike development in the mutant compared with WT. Calculation of the data in Fig. 3I and 3L showed that the *gs1.1-1* and *-2* mutant plants had fewer spikelets per spike and lower grains per spikelet (the ratio of grain number per spike over spikelet number per spike) than WT, indicating that both of spikelet development and floret development (or seed setting rate) were impaired in the mutants. Measurement of free amino acids in young spikes at stem elongation stage (corresponding to the floret development stage) revealed the over-accumulations of Gln, Asn, Arg, Ala and Pro in the *gs1.1-1* mutant plants (Supplemental Figure S8A), and these over-accumulations maybe associated with the impaired spike growth in the mutant. Maintenance of internal amino acid homeostasis has been shown the importance plant growth and development (Lu et al., 2018). Loss-of-function of *OsARG* encoding an arginine hydrolysis enzyme increases Arg accumulation in panicles and leads to small panicle and low seed-setting rate in rice (Ma et al., 2013). Exogenous application of a low level of Lys, Arg, Val, and Ala to nutrient solution promotes tiller bud outgrowth, but applying a high level of these amino acids inhibits tiller bud outgrowth in rice (Lu et al., 2018). Overexpression of an amino acid permease, *OsAAP5*, in rice transports more Lys, Arg, Val and Ala to the tiller base parts and reduces tiller bud outgrowth, while the opposite result occurs in the RNAi-mediated knockdown lines (Wang et al., 2019). These results suggest that an appropriate concentration range of free amino acids is required for tiller and spike growth. As such, it can be assumed that over accumulation of free amino acids such as Arg and Ala may inhibit spike growth in the *gs1.1-1* mutant.

In summary, the three *TaGSI* genes differentially responded to N availability. *TaGSI.1* is up-regulated by low-N and is the major *GSI* gene expressed in roots and

shoots at transcriptional and protein levels, supporting the importance of this gene in NH_4^+ assimilation and wheat adaptation to low-N stress. *TaGS1.1* has essential roles in remobilization, and its mutation delays leaf N loss and senescence during grain filling. Lack of *TaGS1.1* causes growth deficiency in an organ age-dependent manner and reduces yield by impairing spikelet and grain number. The current study and the reported literature showed that GS1.1 orthologues in wheat, rice, and maize were functionally diversified in adaptation to N resources and in mediating yield component formation. As such, strategies for the use of *GS1* genes in increasing yield and NUE should be optimized in different crops.

Acknowledgments

This research was supported by The National Key Research and Development Program of China (2016YFD0100706) and the Strategic Priority Research Program of the Chinese Academy of Sciences (Precision Seed Design and Breeding, Grant No. XDA24010202).

Author contributions

YZW and WT performed the experiments; YPW and CXG performed genome editing; WT and XO identified the *gs1.1* mutant; XH and XQZ assisted field experiments; WT and YPT wrote the article.

References

- Bernard SM, Habash DZ.** 2009. The importance of cytosolic glutamine synthetase in nitrogen assimilation and recycling. *New Phytologist* **182**, 608-620.
- Braun HJ, Atlin G, Payne T.** 2010. Multi-location testing as a tool to identify plant response to global climate change. In: Reynolds CK, ed. *Climate Change and Crop Production*. UK: CABI Climate Change Series, 20260-20264.
- Canas RA, Quillere I, Lea PJ, Hirel B.** 2010. Analysis of amino acid metabolism in the ear of maize mutants deficient in two cytosolic glutamine synthetase isoenzymes highlights the importance of asparagine for nitrogen translocation within sink organs. *Plant Biotechnology Journal* **8**, 966-978.
- Cataldo DA, Haroon M, Schrader LE, Youngs VL.** 1975. Rapid colorimetric determination of nitrate in plant-tissue by nitration of salicylic-acid. *Communications in Soil Science and Plant Analysis* **6**, 71-80.
- Coruzzi GM.** 2003. Primary N-assimilation into amino acids in Arabidopsis. *Arabidopsis Book* **2**, e0010.
- Fontaine JX, Ravel C, Pageau K, Heumez E, Dubois F, Hirel B, Le Gouis J.** 2009. A quantitative genetic study for elucidating the contribution of glutamine synthetase, glutamate dehydrogenase and other nitrogen-related physiological traits to the agronomic performance of common wheat. *Theoretical and Applied Genetics* **119**, 645-662.
- Funayama K, Kojima S, Tabuchi-Kobayashi M, Sawa Y, Nakayama Y, Hayakawa T, Yamaya T.** 2013. Cytosolic glutamine synthetase1;2 is responsible for the primary assimilation of ammonium in rice roots. *Plant and Cell Physiology* **54**, 934-943.
- Galili S, Amir R, Galili G.** 2008. Genetic engineering of amino acids in plants. *Advances in Plant Biochemistry and Molecular Biology*, 49-80.
- Guan M, Moller IS, Schjoerring JK.** 2015. Two cytosolic glutamine synthetase isoforms play specific roles for seed germination and seed yield structure in

Arabidopsis. *Journal of Experimental Botany* **66**, 203-212.

Gutierrez RA, Stokes TL, Thum K, Xu X, Obertello M, Katari MS, Tanurdzic M, Dean A, Nero DC, McClung CR, Coruzzi GM. 2008. Systems approach identifies an organic nitrogen-responsive gene network that is regulated by the master clock control gene CCA1. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 4939-4944.

Habash DZ, Bernard S, Schondelmaier J, Weyen J, Quarrie SA. 2007. The genetics of nitrogen use in hexaploid wheat: N utilisation, development and yield. *Theoretical and Applied Genetics* **114**, 403-419.

Habash DZ, Massiah AJ, Rong HL, Wallsgrove RM, Leigh RA. 2001. The role of cytosolic glutamine synthetase in wheat. *Annals of Applied Biology* **138**, 83-89.

Hawkesford MJ, Araus JL, Park R, Calderini D, Miralles D, Shen T, Zhang J, Parry MA. 2013. Prospects of doubling global wheat yields. *Food and Energy Security* **2**, 34-48.

Hu MY, Zhao XQ, Liu Q, Hong X, Zhang W, Zhang YJ, Sun LJ, Li H, Tong YP. 2018. Transgenic expression of plastidic glutamine synthetase increases nitrogen uptake and yield in wheat. *Plant Biotechnology Journal* **16**, 1858-1867.

Husted S, Hebborn CA, Mattsson M, Schjoerring JK. 2000. A critical experimental evaluation of methods for determination of NH_4^+ in plant tissue, xylem sap and apoplastic fluid. *Physiologia Plantarum* **109**, 167-179.

Ishiyama K, Inoue E, Tabuchi M, Yamaya T, Takahashi H. 2004. Biochemical background and compartmentalized functions of cytosolic glutamine synthetase for active ammonium assimilation in rice roots. *Plant and Cell Physiology* **45**, 1640-1647.

Kamachi K, Yamaya T, Mae T, Ojima K. 1991. A role for glutamine synthetase in the remobilization of leaf nitrogen during natural senescence in rice leaves. *Plant Physiology* **96**, 411-417.

Kichey T, Heumez E, Pocholle D, Pageau K, Vanacker H, Dubois F, Le Gouis J, Hirel B. 2006. Combined agronomic and physiological aspects of nitrogen

management in wheat highlight a central role for glutamine synthetase. *New Phytologist* **169**, 265-278.

Kichey T, Hirel B, Heumez E, Dubois F, Le Gouis J. 2007. In winter wheat (*Triticum aestivum* L.), post-anthesis nitrogen uptake and remobilisation to the grain correlates with agronomic traits and nitrogen physiological markers. *Field Crops Research* **102**, 22-32.

Konishi N, Ishiyama K, Beier MP, Inoue E, Kanno K, Yamaya T, Takahashi H, Kojima S. 2017. Contributions of two cytosolic glutamine synthetase isozymes to ammonium assimilation in *Arabidopsis* roots. *Journal of Experimental Botany* **68**, 610-625.

Konishi N, Saito M, Imagawa F, Kanno K, Yamaya T, Kojima S. 2018. Cytosolic glutamine synthetase isozymes play redundant roles in ammonium assimilation under low-ammonium conditions in roots of *Arabidopsis thaliana*. *Plant and Cell Physiology* **59**, 601-613.

Kusano M, Fukushima A, Tabuchi-Kobayashi M, Funayama K, Kojima S, Maruyama K, Yamamoto YY, Nishizawa T, Kobayashi M, Wakazaki M, Sato M, Toyooka K, Osanai-Kondo K, Utsumi Y, Seki M, Fukai C, Saito K, Yamaya T. 2020. Cytosolic GLUTAMINE SYNTHETASE1;1 modulates metabolism and chloroplast development in roots. *Plant Physiology* **182**, 1894-1909.

Kusano M, Tabuchi M, Fukushima A, Funayama K, Diaz C, Kobayashi M, Hayashi N, Tsuchiya YN, Takahashi H, Kamata A, Yamaya T, Saito K. 2011. Metabolomics data reveal a crucial role of cytosolic glutamine synthetase 1;1 in coordinating metabolic balance in rice. *The Plant Journal* **66**, 456-466.

Li HM, Liang H, Li Z, Tang ZX, Fu SL, Geng YY, Yan BJ, Ren ZL. 2015. Dynamic QTL analysis of protein content and glutamine synthetase activity in recombinant inbred wheat lines. *Genetics and Molecular Research* **14**, 8706-8715.

Lothier J, Gauffichon L, Sormani R, Lemaitre T, Azzopardi M, Morin H, Chardon F, Reisdorf-Cren M, Avise JC, Masclaux-Daubresse C. 2011. The

cytosolic glutamine synthetase *GLN1;2* plays a role in the control of plant growth and ammonium homeostasis in Arabidopsis rosettes when nitrate supply is not limiting. *Journal of Experimental Botany* **62**, 1375-1390.

Lu K, Wu B, Wang J, Zhu W, Nie H, Qian J, Huang W, Fang Z. 2018. Blocking amino acid transporter OsAAP3 improves grain yield by promoting outgrowth buds and increasing tiller number in rice. *Plant Biotechnology Journal* **16**, 1710-1722.

Ma X, Cheng Z, Qin R, Qiu Y, Heng Y, Yang H, Ren Y, Wang X, Bi J, Ma X, Zhang X, Wang J, Lei C, Guo X, Wang J, Wu F, Jiang L, Wang H, Wan J. 2013. OsARG encodes an arginase that plays critical roles in panicle development and grain production in rice. *The Plant Journal* **73**, 190-200.

Martin A, Lee J, Kichey T, Gerentes D, Zivy M, Tatout C, Dubois F, Balliau T, Valot B, Davanture M, Terce-Laforgue T, Quillere I, Coque M, Gallais A, Gonzalez-Moro MB, Bethencourt L, Habash DZ, Lea PJ, Charcosset A, Perez P, Murigneux A, Sakakibara H, Edwards KJ, Hirel B. 2006. Two cytosolic glutamine synthetase isoforms of maize are specifically involved in the control of grain production. *Plant Cell* **18**, 3252-3274.

Masclaux-Daubresse C, Daniel-Vedele F, Dechorgnat J, Chardon F, Gaufichon L, Suzuki A. 2010. Nitrogen uptake, assimilation and remobilization in plants: challenges for sustainable and productive agriculture. *Annals of Botany* **105**, 1141-1157.

Miller AJ, Fan XR, Shen QR, Smith SJ. 2008. Amino acids and nitrate as signals for the regulation of nitrogen acquisition. *Journal of Experimental Botany* **59**, 111-119.

Moison M, Marmagne A, Dinant S, Soulay F, Azzopardi M, Lothier J, Citerne S, Morin H, Legay N, Chardon F, Avise JC, Reisdorf-Cren M, Masclaux-Daubresse C. 2018. Three cytosolic glutamine synthetase isoforms localized in different-order veins act together for N remobilization and seed filling in Arabidopsis. *Journal of Experimental Botany* **69**, 4379-4393.

- Otteson BN, Mergoum M, Ransom JK, Schatz B.** 2008. Tiller contribution to spring wheat yield under varying seeding and nitrogen management. *Agronomy Journal* **100**, 406-413.
- Peltonen J.** 1992. Ear developmental stage used for timing supplemental nitrogen application to spring wheat. *Crop Science* **32**, 1029-1033.
- Peltonen J.** 1993. Grain-yield of high-protein and low-protein wheat cultivars as influenced by timing of nitrogen application during generative development. *Field Crops Research* **33**, 385-397.
- Ren YZ, He X, Liu DC, Li JJ, Zhao XQ, Li B, Tong YP, Zhang AM, Li ZS.** 2012. Major quantitative trait loci for seminal root morphology of wheat seedlings. *Molecular Breeding* **30**, 139-148.
- Shan QW, Wang YP, Li J, Gao CX.** 2014. Genome editing in rice and wheat using the CRISPR/Cas system. *Nature Protocols* **9**, 2395-2410.
- Shan QW, Wang YP, Li J, Zhang Y, Chen KL, Liang Z, Zhang K, Liu JX, Xi JJ, Qiu JL, Gao CX.** 2013. Targeted genome modification of crop plants using a CRISPR-Cas system. *Nature Biotechnology* **31**, 686-688.
- Tabuchi M, Sugiyama K, Ishiyama K, Inoue E, Sato T, Takahashi H, Yamaya T.** 2005. Severe reduction in growth rate and grain filling of rice mutants lacking *OsGSI;1*, a cytosolic glutamine synthetase1;1. *The Plant Journal* **42**, 641-651.
- Thomsen HC, Eriksson D, Moller IS, Schjoerring JK.** 2014. Cytosolic glutamine synthetase: a target for improvement of crop nitrogen use efficiency? *Trends in Plant Science* **19**, 656-663.
- Tilman D, Balzer C, Hill J, Befort BL.** 2011. Global food demand and the sustainable intensification of agriculture. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 20260-20264.
- Voelker TA, Moreno J, Chrispeels MJ.** 1990. Expression analysis of a pseudogene in transgenic tobacco: a frameshift mutation prevents mRNA accumulation. *Plant Cell* **2**, 255-261.

- Wang J, Wu B, Lu K, Wei Q, Qian J, Chen Y, Fang Z.** 2019. The amino acid permease 5 (*OsAAP5*) Regulates tiller number and grain yield in rice. *Plant Physiology* **180**, 1031-1045.
- Wang YP, Cheng X, Shan QW, Zhang Y, Liu JX, Gao CX, Qiu JL.** 2014. Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nature Biotechnology* **32**, 947-951.
- Wei YH, Wang XC, Zhang ZY, Xiong SP, Zhang YM, Wang LL, Meng XD, Zhang J, Ma XM.** 2019. How do three cytosolic glutamine synthetase isozymes of wheat perform N assimilation and translocation? bioRxiv 733857; doi:<https://doi.org/10.1101/733857>.
- Yamaya T, Kusano M.** 2014. Evidence supporting distinct functions of three cytosolic glutamine synthetases and two NADH-glutamate synthases in rice. *Journal of Experimental Botany* **65**, 5519-5525.
- Yang J, Wang M, Li W, He X, Teng W, Ma W, Zhao X, Hu M, Li H, Zhang Y, Tong Y.** 2019. Reducing expression of a nitrate-responsive bZIP transcription factor increases grain yield and N use in wheat. *Plant Biotechnology Journal* **17**, 1823-1833.
- Zhang W, Fan X, Gao Y, Liu L, Sun L, Su Q, Han J, Zhang N, Cui F, Ji J, Tong Y, Li J.** 2017a. Chromatin modification contributes to the expression divergence of three *TaGS2* homoeologs in hexaploid wheat. *Scientific Reports* **7**, 44677.
- Zhang YW, Bai Y, Wu GH, Zou SH, Chen YF, Gao CX, Tang DZ.** 2017b. Simultaneous modification of three homoeologs of *TaEDR1* by genome editing enhances powdery mildew resistance in wheat. *The Plant Journal* **91**, 714-724.
- Zhang ZY, Xiong SP, Wei YH, Meng XD, Wang XC, Ma XM.** 2017c. The role of glutamine synthetase isozymes in enhancing nitrogen use efficiency of N-efficient winter wheat. *Scientific Reports* **7**, 1000.

Table 1. Dry weight for different aerial organs of the main culm at stem elongation in the 2017-2018 growing season.

Organ	High-N treatment			Low-N treatment		
	KN199	<i>gs1.1-1</i>	<i>gs1.1-1</i> /KN199	KN199	<i>gs1.1-1</i>	<i>gs1.1-1</i> /KN199
Spike (mg)	77 ± 5	54 ± 6*	70.1%	54 ± 3	34 ± 6*	63.0%
Stem (mg)	413 ± 19	350 ± 17*	84.7%	396 ± 11	334 ± 29	84.3%
Leaf sheath (mg)	452 ± 24	372 ± 17*	82.3%	402 ± 14	343 ± 28	85.3%
Flag leaf blade (mg)	122 ± 5	106 ± 1*	86.9%	120 ± 12	93 ± 2	77.5%
Middle leaf blade (mg)	260 ± 5	213 ± 6**	81.9%	263 ± 10	211 ± 8**	80.2%
Bottom leaf blade (mg)	133 ± 7	123 ± 6	92.5%	136 ± 9	128 ± 14	94.1%

Data are mean ± SE of four replications. Asterisks indicate statistically significant differences between wild type and *gs1.1-1* mutant at $P < 0.05$ (*) and $P < 0.01$ (**).

Figure legend

Figure 1. The response of *TaGSI* genes to N availability. A-C, The expression of *TaGSI.1* (A), *TaGSI.2* (B), and *TaGSI.3* (C) in roots and shoots of wheat seedlings grown under 1.0 mM NH_4NO_3 (SN), 0.1 mM NH_4NO_3 (LN), and 4.0 mM NH_4^+ (AN) conditions. The germinated seedlings grown for 18 days under SN, LN, and AN conditions. The relative expression levels were normalized to the expression of *TaActin*. The data are expressed as mean $\pm$ SE of three replicates. Asterisks indicate the significant difference compared with SN treatment at $P < 0.05$ (*) and $P < 0.01$ (**) level.

Figure 2. Development of *TaGSI.1* mutant by CRISPR/Cas9-mediated genome editing. Target sites of single guide RNA (sgRNA) in the three *TaGSI.1* homoeologs, and CRISPR/Cas9-induced deletions in the three *TaGSI.1* genes in the *gs1.1* mutants are illustrated. The protospacer-adjacent motif (PAM) is highlighted in red, the single nucleotide polymorphism (SNP) between the three target sites is underlined, the *Bgl*III sites using in the PCR-RE are highlighted in bold font type .

Figure 3. Agronomic traits of the wild type and *gs1.1* mutants under low-N and high-N conditions in the 2017-2018 growing seasons. A and B, images of plants at 14 days post-anthesis (DPA) (A) and 28 DPA (B), bar = 10 cm; C and D, Spikes at 7 DPA (C) and 30 DPA (D), bar = 3 cm; E, Plant height; F, Grain yield per plant; G, Spike number per plant; H, 1000-grain weight; I, Spike grain number; J, Spike grain weight; K, Spike length; L, Spikelet number per main spike. Data are mean $\pm$ SE of four replications. Asterisks indicate statistically significant differences between wild type and *gs1.1* mutants at $P < 0.05$ (*) and $P < 0.01$ (**).

Figure 4. Root and shoot growth-related traits of the wild type and *gs1.1* mutants at seedling stage. The germinated seedlings were grown for 18 days under 1.0 mM NH_4NO_3 (SN), 0.1 mM NH_4NO_3 (LN) and 4.0 mM NH_4^+ (AN) conditions. A, Plant height; B, Shoot dry weight per plant (SDW); C, Tiller number per plant; D-F, Length of the 1st, 2nd and 3rd leaf; G, Root dry weight per plant (RDW); H, Root/Shoot dry weight ratio. Data are mean $\pm$ SE of four replications. Asterisks indicate statistically

618 significant differences between the wild type and *gs1.1* mutants at $P < 0.05$ (*) and P
619 < 0.01 (**).

620

621 **Figure 5.** GS activity of the *gs1.1-1* mutant. The germinated seedlings were grown for
622 18 days under 1.0 mM NH_4NO_3 (SN), 0.1 mM NH_4NO_3 (LN) and 4.0 mM NH_4^+ (AN)
623 conditions. A, Transcriptional expression of the three *TaGS1.1* homeologs in wheat
624 seedling of wild type (WT, KN199) and *gs1.1-1* mutant under SN and LN conditions,
625 the relative expression levels were normalized to the expression of *TaActin*; B and C,
626 Western blot (B) and relative protein abundance (C) of GS1, equal amounts of
627 proteins were loaded in each lane; D, GS activity. Data are represented as means $\pm$ SE
628 of four replicates. Asterisks indicate that the difference between the means of the
629 *gs1.1-1* mutant and KN199 was significant at the $P < 0.05$ (*) and $P < 0.01$ (**) level.

630

631 **Figure 6.** Concentration of N metabolites in roots and shoots of the wild type and
632 *gs1.1-1* mutant. The roots and shoots of wheat seedlings described in Fig. 4 were used
633 to measure the concentrations of total N, nitrate, ammonium, and free amino acids. A,
634 Total N concentration; B, Nitrate concentration; C, Ammonium concentration; D,
635 Concentration of total free amino acids; E, Glu concentration; F, Gln concentration; G,
636 Asp concentration; H, Asn concentration. Data are mean $\pm$ SE of four replications.
637 Asterisks indicate statistically significant differences between wild type and *gs1.1-1*
638 mutant at $P < 0.05$ (*) and $P < 0.01$ (**).

639

640 **Figure 7.** Photosynthetic parameters and N concentrations in flag leaves of the wild
641 type and *gs1.1-1* mutant in the 2017-2018 growing season. A, Net photosynthetic rate
642 (P_n); B, Stomatal conductance (G_s); C, Intercellular CO_2 concentration (C_i); D,
643 Transpiration rate (Tr); E, N concentrations in flag leaves; F, SPAD values of the flag
644 leaves. Data are mean $\pm$ SE of four replications. Asterisks indicate statistically
645 significant differences between wild type and *gs1.1-1* mutant at $P < 0.05$ (*) and $P <$
646 0.01 (**).

647

648 **Figure 8.** N and Dry matter translocation efficiencies in different aerial parts of the
649 main culm in the wild type and *gs1.1-1* mutant plants in the experiment of the
650 2017-2018 growing season. A, N translocation efficiency, which was expressed as (N

651 accumulation at anthesis – N accumulation at maturity) / N accumulation at anthesis ×
 652 100%; B, N harvest index; C, Dry matter translocation efficiency, which was
 653 expressed as (dry weight at anthesis – dry weight at maturity) / dry weight at anthesis
 654 × 100%. But the dry matter translocation efficiency for stem was calculated from the
 655 data at 14 DPA and maturity; D, Harvest index. DPA, days post-anthesis. Data are
 656 mean ± SE of four replications. Asterisks indicate statistically significant differences
 657 between wild type and *gs1.1-1* mutant at $P < 0.05$ (*) and $P < 0.01$ (**).

658 **Supplemental Data**

659 **Supplementary Document S1.** Off-target prediction

660

661 **Supplemental Figure S1.** Phylogenetic analysis of GS proteins in plants.

662

663 **Supplemental Figure S2.** Expression analysis of *GS1* genes in different organs of

664 wheat.

665

666 **Supplemental Figure S3.** Preliminary measurement of agronomic traits in the wild

667 type KN199 and the *gs1.1* mutants.

668

669 **Supplemental Figure S4.** The difference of heading time between the wild type and

670 *gs1.1-1* mutant plants.

671

672 **Supplemental Figure S5.** Dry weights of the aerial parts in the main culm at different

673 developmental stages in the 2017-2018 growing season.

674

675 **Supplemental Figure S6.** The expression of N assimilation genes in the WT and

676 *gs1.1-1* mutant seedlings.

677

678 **Supplemental Figure S7.** The expression of *GS* and *GOGAT* genes in flag leaves of

679 the WT and *gs1.1-1* mutant plants under high-N conditions in the field experiment of

680 the 2017-2018 growing season.

681

682 **Supplemental Figure S8.** Concentration of free amino acids in young spikes and

683 mature seeds in the field experiment of the 2017-2018 growing season.

684

685 **Supplemental Figure S9.** N concentrations in different aerial parts of the main culm

686 at different developmental stages in the field experiment in the 2017-2018 growing

687 season.

688

689 **Supplemental Table S1.** PCR primers used in this study.

690

691 **Supplemental Table S2.** Gene Ids and former names of *GS* genes in wheat.

692

693 **Supplemental Table S3.** Agronomic traits of the wild type and *gs1.1-1* mutant under
694 low-N and high-N conditions in the field experiment of the 2018-2019 growing
695 season.

696

697 **Supplemental Table S4.** Concentration of N metabolites in shoots and roots of the
698 wild type and *gs1.1-1* mutant grown under SN, LN, and AN conditions.

699

Figure 1. The response of *TaGS1* genes to N availability. A-C, The expression of *TaGS1.1* (A), *TaGS1.2* (B), and *TaGS1.3* (C) in roots and shoots of wheat seedlings grown under 1.0 mM NH_4NO_3 (SN), 0.1 mM NH_4NO_3 (LN), and 4.0 mM NH_4^+ (AN) conditions. The germinated seedlings grown for 18 days under SN, LN, and AN conditions. The relative expression levels were normalized to the expression of *TaActin*. The data are expressed as mean $\pm$ SE of three replicates. Asterisks indicate the significant difference compared with SN treatment at $P < 0.05$ (*) and $P < 0.01$ (**) level.

Figure 2. Development of *TaGS1.1* mutant by CRISPR/Cas9-mediated genome editing. Target sites of single guide RNA (sgRNA) in the three *TaGS1.1* homoeologs, and CRISPR/Cas9-induced deletions in the three *TaGS1.1* genes in the *gs1.1* mutants are illustrated. The protospacer-adjacent motif (PAM) is highlighted in red, the single nucleotide polymorphism (SNP) between the three target sites is underlined, the *Bg*/III sites using in the PCR-RE are highlighted in bold font type .

Figure 3. Agronomic traits of the wild type and *gs1.1* mutants under low N and high N conditions in the 2017-2018 growing seasons. A and B, images of plants at 14 days post-anthesis (DPA) (A) and 28 DPA (B), bar = 10 cm; C and D, Spikes at 7 DPA (C) and 30 DPA (D), bar = 3 cm; E, Plant height; F, Grain yield per plant; G, Spike number per plant; H, 1000-grain weight; I, Spike grain number; J, Spike grain weight; K, Spike length; L, Spikelet number per main spike. Data are mean $\pm$ SE of four replications. Asterisks indicate statistically significant differences between wild type and *gs1.1* mutants at $P < 0.05$ (*) and $P < 0.01$ (**).

Figure 4. Root and shoot growth-related traits of the wild type and *TaGSI.1* mutants at seedling stage. The germinated seedlings were grown for 18 days under 1.0 mM NH_4NO_3 (SN), 0.1 mM NH_4NO_3 (LN) and 4.0 mM NH_4^+ (AN) conditions. A, Plant height; B, Shoot dry weight per plant (SDW); C, Tiller number per plant; D-F, Length of the 1st, 2nd and 3rd leaf; G, Root dry weight per plant (RDW); H, Root/Shoot dry weight ratio. Data are mean $\pm$ SE of four replications. Asterisks indicate statistically significant differences between wild type and *gs1.1* mutants at $P < 0.05$ (*) and $P < 0.01$ (**).

Figure 5. GS activity of the *gs1.1-1* mutant. The germinated seedlings were grown for 18 days under 1.0 mM NH_4NO_3 (SN), 0.1 mM NH_4NO_3 (LN) and 4.0 mM NH_4^+ (AN) conditions. A, Transcriptional expression of the three *TaGS1.1* homeologs in wheat seedling of wild type (WT, KN199) and *gs1.1-1* mutant under SN and LN conditions, the relative expression levels were normalized to the expression of *TaActin*; B and C, Western blot (B) and relative protein abundance (C) of GS1, equal amounts of proteins were loaded in each lane; D, GS activity. Data are represented as means $\pm$ SE of four replicates. Asterisks indicate that the difference between the means of the *gs1.1-1* mutant and KN199 was significant at the $P < 0.05$ (*) and $P < 0.01$ (**) level.

Figure 6. Concentration of N metabolites in roots and shoots of the wild type and *gs1.1-1* mutant. The roots and shoots of wheat seedlings described in Fig. 4 were used to measure the concentrations of total N, nitrate, ammonium, and free amino acids. A, Total N concentration; B, Nitrate concentration; C, Ammonium concentration; D, Concentration of total free amino acids; E, Glu concentration; F, Gln concentration; G, Asp concentration; H, Asn concentration. Data are mean $\pm$ SE of four replications. Asterisks indicate statistically significant differences between wild type and *gs1.1-1* mutant at $P < 0.05$ (*) and $P < 0.01$ (**).

Figure 7. Photosynthetic parameters and N concentrations in flag leaves of the wild type and *gs1.1-1* mutant in the 2017-2018 growing season. A, Net photosynthetic rate (P_n); B, Stomatal conductance (G_s); C, Intercellular CO_2 concentration (C_i); D, Transpiration rate (Tr); E, N concentrations in flag leaves; F, SPAD values of the flag leaves. Data are mean $\pm$ SE of four replications. Asterisks indicate statistically significant differences between wild type and *gs1.1-1* mutant at $P < 0.05$ (*) and $P < 0.01$ (**).

Figure 8. N and Dry matter translocation efficiencies in different aerial parts of the main culm in the wild type and *gs1.1-1* mutant plants in the experiment of the 2017-2018 growing season. A, N translocation efficiency, which was expressed as (N accumulation at anthesis – N accumulation at maturity) / N accumulation at anthesis. B, N harvest index; C, Dry matter translocation efficiency, which was expressed as (dry weight at at anthesis – dry weight at maturity) / dry weight at at anthesis, but the dry matter translocation efficiency for stem was calculated from the data at 14 DPA and maturity. D, Harvest index. DPA, days post-anthesis. Data are mean $\pm$ SE of four replications. Asterisks indicate statistically significant differences between wild type and *gs1.1-1* mutant at $P < 0.05$ (*) and $P < 0.01$ (**).