Title: A single nuclei transcriptomic analysis of the Atlantic salmon gill through smoltification and seawater transfer

Authors: Alexander C. West¹, Yasutaka Mizoro², Shona H. Wood¹, Louise M. Ince³, Marianne Iversen¹, Even H. Jørgensen¹, Torfinn Nome⁴, Simen Rød Sandve⁴, Andrew S. I. Loudon² and David G. Hazlerigg¹

¹Arctic Chronobiology and Physiology, University of Tromsø, Framstredet 42, 9019 Tromsø, Norway

²Division of Diabetes, Endocrinology & Gastroenterology, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester, UK

³University of Geneva, Centre Médical Universitaire (CMU), Department of Pathology and Immunology, Switzerland

⁴Centre for Integrative Genetics, Department of Animal and Aquaculture Sciences, Norwegian University of Life Sciences, 1432 Ås, Norway

*Correspondence: david.hazlerigg@uit.no

Abstract

Anadromous salmonids begin life adapted to the freshwater environments of their natal streams before a developmental transition, known as smoltification, transforms them into marine-adapted fish. In the wild, the extending photoperiods of spring stimulates smoltification, typified by radical reprogramming of the gill from an ion-absorbing organ to ion-excreting organ. Prior work has highlighted the role of specialized “mitochondrion-rich”

cells in delivering this phenotype. However, transcriptomic studies identify thousands of smoltification-driven differentially regulated genes, indicating that smoltification causes a multifaceted, multicellular change; but direct evidence of this is lacking.

Here, we use single-nuclei RNAseq to characterize the Atlantic salmon gill during smoltification and seawater transfer. We identify 20 distinct clusters of nuclei, including known, but also novel gill cell types. These data allow us to isolate cluster-specific, smoltification-induced changes in gene expression. We also show how cellular make-up of the gill changes through smoltification. As expected, we noted an increase in the proportion of seawater mitochondrion-rich cells, however, we also identify a reduction of several immune-related cells. Overall, our results provide unrivaled detail of the cellular complexity in the gill and suggest that smoltification triggers unexpected immune reprogramming directly preceding seawater entry.

Keywords: Atlantic salmon, smoltification, photoperiod, seasonal, gill

Introduction

The Atlantic salmon migrates between fresh and seawater environments ¹. Atlantic salmon eggs hatch in freshwater streams where they develop for 1-4 years. On reaching a critical size threshold, young “parr” animals are sensitized by several weeks of winter photoperiod (day-lengths), after which long, summer-like photoperiods stimulates the parr to transform into a marine-adapted “smolt” fish ². This process, known as smoltification, drives divergent expression of endocrine factors that collectively deliver phenotypic remodeling, of length, weight, silvering, and in particular: gill physiology ¹.

The salmonid gill is a complex multifunctional organ, essential for gas exchange, nitrogenous waste excretion, pH balance and osmoregulation ³. It is also a major mucosal immune barrier harboring a dedicated lymphoid tissue ⁴. Structurally, the gills are arranged in symmetrical arches, each of which are populated by numerous filament structures, which are themselves densely flanked with lamellae. The gill is composed of seven major cell types ⁵. Pavement cells (PVCs) have an enlarged surface area on the apical membrane, and form the majority of the epithelium ⁶. Pillar cells (PCs), which are structural cells, define the blood spaces within the lamellae ⁷. Goblet cells (GCs) reside in the filament epithelium and excrete mucus ⁸. Non-differentiated progenitor cells (NDCs) colonize basal and intermediate layers of the gill epithelium ⁹. Chemosensory neuroepithelial cells (NECs) lie along the length of the efferent edge of the gills and are innervated by the central nervous system ¹⁰. Mitochondrion-rich cells (MRCs) and their adjacent accessory cells (ACs), finally, are located at the trough between two lamellae where they abundantly express the channels and pumps required to maintain the osmotic gradients between blood plasma and both fresh- and seawater ^{11–13}.

Under freshwater, Na⁺ ions are directly or indirectly exchanged for protons across the apical membrane then transported into the blood *via* the sodium potassium ATPase (NKA) on the basolateral membrane ^{14–16}. Cl⁻ ions, meanwhile, are exchanged or channeled across the apical membrane then enter the blood through an undefined channel ^{17–20}. Under saltwater, basolateral NKA generates a chemical and electrical gradient, motivating both loss of Cl⁻ ions *via* apical CFTR channels and paracellular escape of Na⁺ ions ^{13,21} (reviewed in ²²).

Smoltification converts the Atlantic salmon gill from a freshwater-adapted organ to a seawater-adapted organ. Rising cortisol and growth hormone along with falling prolactin propels smoltification. This change in endocrinology coincides with a switch in anatomical and

molecular phenotypes of MRCs and ACs, cell types which to date comprise the major focus of smoltification of gill physiology^{1,12,23}. Smoltification is, however, a complex developmental transition and the smolt gill phenotype likely extends far beyond changing MRC and AC cell phenotype. Therefore, to realize the complexities of smoltification-driven changes in gill physiology we adopt a new strategy, single-nuclei RNAseq, which provides transcriptional responses to smoltification and seawater transfer at individual nuclei-level resolution.

Results & Discussion

A single-nuclei survey of Atlantic salmon gill cells

We profiled 18,844 individual nuclei from eight Atlantic salmon gill samples from four distinct physiological states (Figure 1A). To define shared correlation structure across datasets we pooled replicate samples and integrated all four states using diagonalized canonical correlation analysis followed by L2 normalization. We next identified pairs of mutual nearest neighbors (MNNs) to identify anchors: cells that represent shared biological states across datasets. Anchors were then used to calculate “correction” vectors allowing all four states to be jointly analyzed as an integrated reference²⁴. Unsupervised graph clustering partitioned the nuclei into 20 clusters, which we visualized using a uniform manifold approximation and projection (UMAP) dimension reduction technique (Figure 1B).

Lists of co-expressed marker genes define individual clusters. We categorized individual clusters using gene ontology analysis of marker gene lists and unique expression of known marker genes. This approach allowed us to infer cell types including fresh- and seawater MRCs, ACs, neuroendothelial cells, goblet cells, non-differentiated cells, pillar cells, lymphatic endothelial cells and several types of blood cell. We also identified a novel population of fibrocyte-like cells, and several types of vascular- and endothelial-like cells that

partitioned across several clusters, together suggesting greater complexity in gill cytology that previously appreciated (Figure 1B).

We then defined expression signatures for each cell cluster. Our analysis re-identified several known marker genes, but also identifying several novel cell-type markers (Figure 1C). For example, the accessory cell signature included highly restricted expression of Slc26a6, an apical membrane $\text{Cl}^-/\text{HCO}_3^-$ exchanger, associated with gill function but until now misattributed to expression within MRCs^{25,26}. We were interested to note that the markers defining the erythrocyte population, including beta-globin, were expressed widely among all cell types. It is unclear exactly what role extra-erythroid haemoglobin plays in the gill, however, mammalian studies suggest that haemoglobin, in addition to its oxygen carrying capacity, may play an antimicrobial role²⁷. As a major mucosal immune barrier, this capacity may be pertinent to the gill²⁸.

Major changes in cell composition during smoltification

To understand how gene expression and cellular complexity changes within the gill during smoltification and seawater transfer we compared the snRNAseq profiles at different developmental points (Fig 2A, for confirmation of smolt status see²⁹). The abundance of six nuclei clusters changed dramatically (>3 fold change in percentage abundance) during smoltification (Figure 2B). SW MRCs increased in proportion steadily from T1-T4, consistent with previous descriptions of Atlantic salmon gill physiology. We also observed a marked increase in vascular cell number, with the major differences occurring between T2 and T3, suggesting that this vascular cell cluster proliferate in line with growth rates (Figure 2C). Interestingly, four immune-related nuclei clusters representing T cells, myeloid cells, dendritic cells and lymphatic endothelial cells fell dramatically during smoltification (Figure 2D).

Changes in cell abundance occurred with a similar profile in all immune-associated cell clusters, with consistent decline observed between T1-T3. In contrast, 24h SW transfer does not appear to affect immune-cell abundance directly. These results highlight the complex and dynamic changes in cellular composition that occur in the gill during smoltification.

Cell cluster-specific expression of smoltification-associated factors

Next, we wanted to identify cluster types where smoltification drives changes in local gene regulation. For statistical power, we cross-referenced our snRNAseq analysis with whole gill RNAseq analysis of T1-T3, identifying 9746 genes differentially regulated by smoltification (FDR <0.01). Pearson clustering of these genes resolved five major clusters that were associated with immune response, structural morphogenesis, autophagy, catabolism and mitochondrial respiration (Figure 3A). Within our analysis we identified a number of “classical” smoltification-related genes. As expected, CFTR was highest under constant light (LL), and was highly localized in expression to MRCs (Figure 3B). We also identified the reciprocal regulation of sodium-potassium ATPase subunits, specifically, suppression of NKAa1a and increase in NKAa1b¹². Inspection of cellular localization within our snRNAseq dataset showed that expression of these genes were, as anticipated, highest within the MRCs and ACs (Figure 3B).

Our previous work identified genes whose expression are predicated on exposure to several weeks of short-photoperiod exposure²⁹. In Atlantic salmon, these “winter-dependent” genes are analogous to vernalization dependent genes in *Arabidopsis*³⁰, where a dosage of exposure to a winter-like stimulus (in *Arabidopsis*, cold; in Atlantic salmon, short photoperiod) controls the presentation of a seasonal phenotype under summer-like stimulus (in *Arabidopsis*, warmth and long days; in Atlantic salmon, long photoperiod). Winter-dependent genes are therefore intrinsically linked to unidirectional smolt development, and may play a

mechanistic role in pre-adaptation of the gill for seawater migration. Surprisingly, canonical markers of smolt status, including the reciprocal expression of NKA subunits, are not winter dependent, meaning that their expression is passive to photoperiod rather than life history progression²⁹.

Using our RNAseq dataset we identified novel, winter dependent genes. Next, we mined our snRNAseq dataset to identify the cell clusters that express these factors (Figure S1). Of particular interest was *Cuzd1*, a gene associated with carcinogenesis, whose expression was tightly localized to non-differentiated cells ³¹. We also identified *Rhag*, an ammonium transporter thought to be erythrocyte specific in mammals, but here predominantly expressed in the vascular cell (VC 3) cluster ³²; and *Hg2a* (CD74) a ubiquitously expression multi-functional protein linked to immune defense ³³. Taken together our data show that smoltification engages all gill cells in diverse regulatory phenotypes.

Cell cluster-specific expression of seawater transfer-associated factors

Smoltification manifests when the Atlantic salmon smolts migrate downstream and arrive in the marine environment, thereby committing to an oceanic life ¹. To gain insight into this critical step we compared RNA profiles of gill samples between smolts in freshwater and 24h in seawater using whole gill RNAseq, identifying 144 induced and 107 suppressed genes (FDR<0.01). Gene ontologies showed that the induced gene cohort was significantly associated with keratinization, whereas the suppressed gene cohort related to immune function (Figure 4A). We cross-referenced our whole gill RNAseq data against our snRNAseq data to isolate cell-type specificity of gene expression. These data highlight the restricted expression of key up-regulated genes (Figure 4B). For example, we localize the expression of an enzyme involved in both ionic and acid/base balance, carbonic anhydrase, to MRCs ^{34,35}.

We also show that ATP-binding cassette sub-family A member 12 (Abca12), a gene important in epidermal lipid barrier formation ³⁶, is broadly expressed, but particularly concentrated in MRCs (SW), pavement, vascular, and non-differentiated cells. Interestingly, we show that a protein chaperone that helps regulate chromatin state, nucleoplasmin ^{37,38}, is expressed specifically in non-differentiated, vascular and pavement cells groups, suggesting that these cell types undergo a change in chromatin status under seawater exposure.

Conclusions

Our results bring insightful cellular resolution to the complexity of the Atlantic salmon gill and the compositional changes that occur during smoltification. Of particular interest was the suppression of immune cell types, which correlates with reduction in immune-related genes and suppression of immune function during smoltification ^{39–41}. These data are a puzzle. The marine environment is awash with parasites, bacteria and viruses to which the salmon is potentially vulnerable, so loss of immune function would make little sense. Future work should focus on why and how the immune system is affected in aquaculture. Conceivably these data point towards an adaptive immunological reprogramming that helps to avoid immune shock when the salmon transition between the distinctive pathogen complements of fresh- and seawater habitats ^{42,43}. Alternatively, artificial smolt production may drive abnormal immunosuppression. The constant light routinely used to stimulate smolts would profoundly undermine the immune defenses of mammals *via* disruption of the circadian clock ⁴⁴.

Our data also shows that smoltification-driven transcriptional regulation occurs not only in MRCs and ACs, but also in other distinctive cell types including pavement cells, vascular cells and non-differentiated cells. We anticipate that novel gene function within the context

of cell function will be a priority for future investigation, and will be assisted by the novel suite of marker genes which we present here.

References

1. Stefansson, S. O., Björnsson, B. T., Ebbesson, L. O. E. & McCormick, S. D. Smoltification. in *Fish Larval Physiology* 639–681 (2008).
2. Strand, J. E. T., Hazlerigg, D. & Jørgensen, E. H. Photoperiod revisited: is there a critical day length for triggering a complete parr–smolt transformation in Atlantic salmon *Salmo salar*? *J. Fish Biol.* **93**, 440–448 (2018).
3. Evans, D. H., Piermarini, P. M. & Choe, K. P. The multifunctional fish gill: Dominant site of gas exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous waste. *Physiol. Rev.* **85**, 97–177 (2005).
4. Koppang, E. O. *et al.* Salmonid T cells assemble in the thymus, spleen and in novel interbranchial lymphoid tissue. *J. Anat.* **217**, 728–739 (2010).
5. Wilson, J. M. & Laurent, P. Fish gill morphology: Inside out. *J. Exp. Zool.* **293**, 192–213 (2002).
6. Laurent, P. & Dunel-Erb, S. *The Pseudobranch: Morphology and Function. Fish Physiology* vol. 10 (1984).
7. Newstead, J. D. Fine structure of the respiratory lamellae of teleostean gills. *Zeitschrift für Zellforsch. und Mikroskopische Anat.* **79**, 396–428 (1967).
8. Rojo, M. C., Blánquez, M. J. & González, M. E. A histochemical study of the distribution

- 202 of lectin binding sites in the developing branchial area of the trout *Salmo trutta*. *J.*
203 *Anat.* **189**, 609–621 (1996).
- 204 9. Laurent, P., Dunel-Erb, S., Chevalier, C. & Lignon, J. Gill epithelial cells kinetics in a
205 freshwater teleost, *Oncorhynchus mykiss* during adaptation to ion-poor water and
206 hormonal treatments. *Fish Physiol. Biochem.* **13**, 353–370 (1994).
- 207 10. Dunel-Erb, S., Chevalier, C. & Laurent, P. Distribution of neuroepithelial cells and
208 neurons in the trout gill filament: comparison in spring and winter. *Can. J. Zool.* **72**,
209 1794–1799 (1994).
- 210 11. Hootman, S. R. & Philpott, C. W. Accessory cells in teleost branchial epithelium. *Am. J.*
211 *Physiol. - Regul. Integr. Comp. Physiol.* **7**, (1980).
- 212 12. McCormick, S. D., Regish, A. M., Christensen, A. K. & Björnsson, B. T. Differential
213 regulation of sodium – potassium pump isoforms during smolt development and
214 seawater exposure of Atlantic salmon. 1142–1151 (2013) doi:10.1242/jeb.080440.
- 215 13. Sardet, C., Pisam, M. & Maetz, J. The surface epithelium of teleostean fish gills. *J. Cell*
216 *Biol.* **80**, 96–117 (1979).
- 217 14. Edwards, S. L., Tse, C. M. & Toop, T. Immunolocalisation of NHE3-like
218 immunoreactivity in the gills of the rainbow trout (*Oncorhynchus mykiss*) and the
219 blue-throated wrasse (*Pseudolabrus tetrius*). *J. Anat.* **195**, 465–469 (1999).
- 220 15. Yan, J. J., Chou, M. Y., Kaneko, T. & Hwang, P. P. Gene expression of Na⁺/H⁺
221 exchanger in zebrafish H⁺-ATPase-rich cells during acclimation to low-Na⁺ and acidic
222 environments. *Am. J. Physiol. - Cell Physiol.* **293**, 1814–1823 (2007).
- 223 16. Hirata, T. *et al.* Mechanism of acid adaptation of a fish living in a pH 3.5 lake. *Am. J.*

- 224 *Physiol. - Regul. Integr. Comp. Physiol.* **284**, (2003).
- 225 17. Hiroi, J., McCormick, S. D., Ohtani-Kaneko, R. & Kaneko, T. Functional classification of
226 mitochondrion-rich cells in euryhaline Mozambique tilapia (*Oreochromis*
227 mossambicus) embryos, by means of triple immunofluorescence staining for Na⁺/K⁺-
228 ATPase, Na⁺/K⁺/2Cl⁻ cotransporter and CFTR anion channel. *J. Exp. Biol.* **208**, 2023–
229 2036 (2005).
- 230 18. Hiroi, J., Yasumasu, S., McCormick, S. D., Hwang, P. P. & Kaneko, T. Evidence for an
231 apical Na-Cl cotransporter involved in ion uptake in a teleost fish. *J. Exp. Biol.* **211**,
232 2584–2599 (2008).
- 233 19. Tang, C. H., Hwang, L. Y. & Lee, T. H. Chloride channel CLC-3 in gills of the euryhaline
234 teleost, *tetraodon nigroviridis*: Expression, localization and the possible role of
235 chloride absorption. *J. Exp. Biol.* **213**, 683–693 (2010).
- 236 20. Preest, M. R., Gonzalez, R. J. & Wilson, R. W. A pharmacological examination of Na⁺
237 and Cl⁻ transport in two species of freshwater fish. *Physiol. Biochem. Zool.* **78**, 259–
238 272 (2005).
- 239 21. Marshall, W. S., Lynch, E. M. & Cozzi, R. R. F. Redistribution of immunofluorescence of
240 CFTR anion channel and NKCC cotransporter in chloride cells during adaptation of the
241 killifish *Fundulus heteroclitus* to sea water. *J. Exp. Biol.* **205**, 1265–1273 (2002).
- 242 22. Evans, D. H. Freshwater fish gill ion transport: August Krogh to morpholinos and
243 microprobes. *Acta Physiol. (Oxf)*. **202**, 349–359 (2011).
- 244 23. Lorgen, M. *et al.* Functional divergence of type 2 deiodinase paralogs in the Atlantic
245 salmon. *Curr. Biol.* **25**, 936–941 (2015).

- 246 24. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-
247 1902.e21 (2019).
- 248 25. Leguen, I., Le Cam, A., Montfort, J., Peron, S. & Fautrel, A. Transcriptomic analysis of
249 trout gill ionocytes in fresh water and sea water using laser capture microdissection
250 combined with microarray analysis. *PLoS One* **10**, 1–22 (2015).
- 251 26. Boyle, D., Clifford, A. M., Orr, E., Chamot, D. & Goss, G. G. Mechanisms of Cl⁻ uptake in
252 rainbow trout: Cloning and expression of slc26a6, a prospective Cl⁻/HCO₃⁻ exchanger.
253 *Comp. Biochem. Physiol. -Part A Mol. Integr. Physiol.* **180**, 43–50 (2015).
- 254 27. Saha, D. *et al.* Hemoglobin Expression in Nonerythroid Cells: Novel or Ubiquitous? *Int.*
255 *J. Inflam.* **2014**, (2014).
- 256 28. Koppang, E. O., Kvellestad, A. & Fischer, U. *Mucosal Health in Aquaculture Fish*
257 *mucosal immunity: gill. Mucosal Health in Aquaculture* (Elsevier Inc., 2015).
258 doi:10.1016/B978-0-12-417186-2/00005-4.
- 259 29. Iversen, M. *et al.* RNA profiling identifies novel, photoperiodhistory dependent
260 markers associated with enhanced saltwater performance in juvenile Atlantic salmon.
261 *PLoS One* **15**, 1–21 (2020).
- 262 30. Song, J., Angel, A., Howard, M. & Dean, C. Vernalization - a cold-induced epigenetic
263 switch. *J. Cell Sci.* **125**, 3723–3731 (2012).
- 264 31. Liaskos, C., Rigopoulou, E. I., Orfanidou, T., Bogdanos, D. P. & Papandreou, C. N.
265 CUZD1 and Anti-CUZD1 Antibodies as Markers of Cancer and Inflammatory Bowel
266 Diseases. **2013**, (2013).
- 267 32. Tilley, L. *et al.* A new blood group system , RHAG : three antigens resulting from amino

- 268 acid substitutions in the Rh-associated glycoprotein. 151–159 (2010)
- 269 doi:10.1111/j.1423-0410.2009.01243.x.
- 270 33. Su, H., Na, N., Zhang, X. & Zhao, Y. The biological function and significance of CD74 in
- 271 immune diseases. *Inflamm. Res.* **66**, 209–216 (2017).
- 272 34. Houde, A. L. S. *et al.* Salmonid gene expression biomarkers indicative of physiological
- 273 responses to changes in salinity and temperature, but not dissolved oxygen. *J. Exp.*
- 274 *Biol.* **222**, (2019).
- 275 35. Zbanyszek, R. & Smith, L. S. Changes in carbonic anhydrase activity in coho salmon
- 276 smolts resulting from physical training and transfer into seawater. *Comp. Biochem.*
- 277 *Physiol. -- Part A Physiol.* **79**, 229–233 (1984).
- 278 36. Akiyama, M. The roles of ABCA12 in epidermal lipid barrier formation and
- 279 keratinocyte differentiation. *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids* **1841**, 435–
- 280 440 (2014).
- 281 37. Bouleau, A. *et al.* Maternally inherited npm2 mRNA is crucial for egg developmental
- 282 competence in zebrafish. *Biol. Reprod.* **91**, 1–9 (2014).
- 283 38. Chen, P. *et al.* Nucleoplasmin is a limiting component in the scaling of nuclear size
- 284 with cytoplasmic volume. *J. Cell Biol.* **218**, 4063–4078 (2019).
- 285 39. Jensen, I., Overrein, M. C., Fredriksen, B. N., Strandskog, G. & Seternes, T. Differences
- 286 in smolt status affect the resistance of Atlantic salmon (*Salmo salar* L.) against
- 287 infectious pancreatic necrosis, while vaccine-mediated protection is unaffected. *J. Fish*
- 288 *Dis.* **42**, 1271–1282 (2019).
- 289 40. Johansson, L. H., Timmerhaus, G., Afanasyev, S., Jørgensen, S. M. & Krasnov, A.

Smoltification and seawater transfer of Atlantic salmon (*Salmo salar* L.) is associated with systemic repression of the immune transcriptome. *Fish Shellfish Immunol.* **58**, 33–41 (2016).

41. Nuñez-Ortiz, N. *et al.* Atlantic salmon post-smolts adapted for a longer time to seawater develop an effective humoral and cellular immune response against Salmonid alphavirus. *Fish Shellfish Immunol.* **82**, 579–590 (2018).

42. Lee, S.-Y. & Eom, Y.-B. Analysis of Microbial Composition Associated with Freshwater and Seawater. *Biomed. Sci. Lett.* **22**, 150–159 (2016).

43. Wang, Y. *et al.* Comparison of the levels of bacterial diversity in freshwater, intertidal wetland, and marine sediments by using millions of illumina tags. *Appl. Environ. Microbiol.* **78**, 8264–8271 (2012).

44. Scheiermann, C., Gibbs, J., Ince, L. & Loudon, A. Clocking in to immunity. *Nature Reviews Immunology* vol. 18 423–437 (2018).

45. Matson, K. J. E. *et al.* Isolation of adult spinal cord nuclei for massively parallel single-nucleus RNA sequencing. *J. Vis. Exp.* **2018**, 1–12 (2018).

Material and Methods

Animal welfare statement

The Atlantic salmon smoltification experiment was conducted as part of the routine, smolt production at Kårvik havbruksstasjonen, approved by the Norwegian Animal Research Authority (NARA) for the maintenance of stock animals for experiments on salmonids. This is in accordance with Norwegian and European legislation on animal research.

Experimental Design

Atlantic salmon (*Salmo salar*, Aquagene commercial strain) were raised from hatching in freshwater, under continuous light (LL, > 200 lux at water surface) at ambient temperature (~10°C). Juvenile salmon were housed in 500 L circular tanks and fed continuously with pelleted salmon feed (Skretting, Stavanger, Norway). At seven months of age parr (mean weight 49.5g) were sampled for T1 (experiment start). Two days later remaining parr were equally distributed between two 100L circular tanks, and over the next seven days the photoperiod was incrementally reduced to a short photoperiod (SP, 8h light:16h darkness). T2 sampling occurred on experimental day 53 (44 days on SP), remaining parr were transferred back to LL on experimental day 60. T3 sampling occurred on experimental day 110 (50 days after return to LL), then a sub-cohort of fish were netted out and transferred to full strength seawater for 24h before the final T4 collection.

RNAseq Analysis

Gill samples were collected, RNA extracted and libraries prepared, sequenced and mapped as Iversen et al (2020). Raw counts were analysed using EdgeR (ver. 3.30.0), using R (ver. 4.0.2) and RStudio (ver. 1.1.456). An ANOVA-like test was used to identify differential expressed genes between T1-T3 samples. Clustering analysis was performed using Pearson correlation, and heatmap rendered using the R package pheatmap. An exact test was performed to identify differential expressed genes between T3 and T4.

Single nuclei RNAseq Analysis

Gills for single nuclei analysis were collected on dry ice and stored at -80°C. Duplicate samples were processed for T1-T4. Nuclei were released by detergent mechanical lysis, then samples were homogenized (30s) and nuclei isolated by sucrose gradient⁴⁵. Libraries were created

using Chromium Single Cell 3' GEM, Library & Gel Bead Kit v3 (10x technologies) by University of Manchester genomic technology core facility (UK). Raw data was converted to counts per cell using Cell Ranger (10x Technologies, ver. 3.1.0) and processed using NCBI annotations. The R package Seurat (ver. 3.1.5) was used to perform an integrated analysis using all snRNAseq data²⁴, further details in results and discussion. Raw data will be available following peer-reviewed publication.

Acknowledgements

The authors thank all of the animal staff at Kårvik havbruksstasjonen for their expert care of the research animals, and the University of Manchester Genomics Technology core facility (UK) for performing chromium 10x library preparation for snRNAseq. ACW is supported by the Tromsø forskningsstiftelse (TFS) grant awarded to DGH (TFS2016DH). The Sentinel North Transdisciplinary Research Program Université Laval and UiT awarded to DGH supports this work. SHW is supported a grant from the Tromsø forskningsstiftelse (TFS) starter grant TFS2016SW. Experimental costs were covered by HFSP grant "Evolution of seasonal timers" RGP0030 /2015 awarded to ASIL and DGH.

Author Contributions

Conceptualization, A.C.W., Y.M., E.H.J., A.S.I.L and D.G.H; Resources, A.C.W, Y.M and M.I.; Investigation, A.C.W., Y.M., M.I. E.H.J and D.G.H; Formal Analysis, A.C.W, Y.M., L.M.I., T.N and S.R.S; Visualization, A.C.W and S.H.W; Writing – Original Draft, A.C.W and S.H.W; Writing –

Review & Editing, All; Supervision, A.S.I.L and D.G.H; Project Administration, A.S.I.L and D.G.H;
Funding Acquisition, A.S.I.L and D.G.H.

Figure Legends

Figure 1. Single nuclei RNAseq analysis of Atlantic salmon gill tissue. A) Gill tissue processing.
B) UMAP plot of pooled cell data from 18844 cells representing eight samples from four
collection states. The plot indicates 20 separate cell clusters. C) Expression of marker genes in
20 cell clusters. From left to right: hierarchical relatedness of difference cell clusters; total cells
in each cluster; UMI number in each cell cluster; gene features in each cell cluster; violin plots
showing expression pattern of marker genes for each cluster. Abbreviations: ACs - accessory
cells, DCs - dendritic cells, ECs - epithelial cells, fib - fibrocytes, GCs - goblet cells, LECs -
lymphatic endothelial cells, MCs - myeloid cells, MRC - mitochondrion-rich cells, NDCs - non-
differentiated cells, PVCs, pavement cells, RBCs - red blood cells (erythrocytes), TCs - T cells,
VCs - vascular cells.

Figure 2. Comparative abundance of cell clusters at different sampling points. A) Experimental
design. Fish were kept in constant light (LL) from hatching then transferred to short
photoperiod (SP; 8L:16D) for 8 weeks before being returned to constant light (LL) for 8 weeks.
Finally the fish were transferred to sea water for 24h. Sample points are indicated T1-T4. B)
Subset of cell clusters from T2, T3 and T4 (green and red dots) overlaid on T1 cells (grey dots).
C) Increasing abundance of sea-water mitochondrion-rich cells (MRCs SW) and vascular cells
(VC 3) during smoltification C) Decreasing abundance of leukocytes and immune-associated
cells during smoltification.

Figure 3. Photoperiodic changes in gill gene expression and localized cell cluster expression.

A) Heat map representing 9746 genes differentially regulated (FDR <0.01) from T1-T3. Regulatory patterns for 5 major cluster are shown as amplitude index and 95% confidence limits. Major gene ontology terms for each cluster are shown. B) RNAseq data for “classical” smoltification-related genes and violin plots showing their cluster specific expression.

Figure 4. Sea-water transfer-associated changes in gill gene expression and localized cell cluster expression. A) Genes differentially regulated (FDR <0.01) by 24h seawater transfer. Major gene ontology terms for each cluster are shown. B) RNAseq data for sea-water transfer-related genes and violin plots showing their cluster specific expression.

Supplemental Figure 1. RNAseq data for winter-dependent genes and violin plots showing their cluster specific expression.

Figure 1.

Figure 2.

Figure 3.

B.

Abca12

Carbonic Anhydrase

Nucleoplasmin

Figure 4.