***Ageratina adenophora* and *Lantana camara* in Kailash Sacred Landscape, India: Current distribution and future climatic scenarios through modeling**

Authors:

Alka Chaudhary¹ <https://orcid.org/0000-0002-4789-6911>;
e-mail: chaudhary.alka43@gmail.com

Mriganka Shekhar Sarkar^{2,3} <https://orcid.org/0000-0001-8367-2774>;
e-mail: mriganka.sarkar@gbpihed.nic.in

Bhupendra Singh Adhikari^{1,*} <https://orcid.org/0000-0001-5632-0044>;
e-mail: adhikaribs@wii.gov.in

Gopal Singh Rawat¹ <https://orcid.org/0000-0002-5743-5035>;
e-mail: rawatg@wii.gov.in

* *Corresponding author*

Affiliations:

¹ Wildlife Institute of India, Dehradun, Uttarakhand 248001, India

² Department of Biological Sciences, Indian Institute of Science Education and Research Kolkata, West Bengal 741246, India

³ North-East Regional Centre, G. B. Pant National Institute of Himalayan Environment (NIHE), Itanagar, Arunachal Pradesh 791113, India

***Corresponding Author**

e-mail: adhikaribs@wii.gov.in

Abstract:

The Himalayan region is one of the global biodiversity hotspots. However, its biodiversity and ecosystems are threatened due to abiotic and biotic drivers. One of the major biotic threats to biodiversity in this region is the rapid spread of invasive alien species (IAS). Natural forests and grasslands are increasingly getting infested by IAS affecting regeneration of native species and decline in availability of bio-resources. Assessing the current status of IAS and prediction of their future spread would be vital for evolving specific species management interventions. Keeping this in view, we conducted an in-depth study on two IASs, viz., *Ageratina adenophora* and *Lantana camara* in the Indian part of Kailash Sacred Landscape (KSL), Western Himalaya. Intensive field surveys were conducted to collect the presence of *A. adenophora* ($n = 567$) and *L. camara* ($n = 120$) along an altitudinal gradient between 300 and 3000 m a.s.l. We performed Principle Component Analysis to nullify the multi-collinearity effects of the environmental predictors following *MaxEnt* species distribution model in the current and future climatic scenarios for both the species. All current and future model precision (i.e. Area Under the Curve; AUC) for both species was higher than 0.81. It is predicted that under the current rate of climate change and higher emission (i.e. RCP8.5 pathway), *A. adenophora* will spread 45.3% more than its current distribution and is likely to reach up to 3029 m a.s.l. Whereas, *L. camara* will spread 29.8% more than its current distribution range and likely to reach up to 3018 m a.s.l. Our results will help in future conservation planning and participatory management of forests and grasslands in the KSL–India.

Keywords: Invasive species; *Ageratina adenophora*; *Lantana camara*; Principal Component Analysis; *MaxEnt*; Kailash Sacred Landscape

Introduction

Invasive alien species (IAS) rank among the top three threats to global biodiversity, the other two being unsustainable harvest of various species from the wild and habitat degradation and loss (1). Invasive species, coupled with unsustainable resource use and climate change have seriously affected livelihoods of millions of people in south and south-east Asia (2). Rapid spread of invasive alien species affects natural habitats, regeneration potential of native species and affect productivity of forests and grassland habitats (3). Spread of alien species is particularly challenging in terrestrial ecosystems across the world (4, 5, 6). Changes in climatic conditions and land use practices favour introduction and spread of IAS in most parts of the world (7, 8, 9). Therefore, it is vital to assess the current extent and future scenarios of invasion for various IAS (10,11,12).

The Himalayan region, one of the global biodiversity hotspots, is quite vulnerable due to climate change and other abiotic and biotic drivers of change including rapid spread of IAS. Some of the most obnoxious IAS in this region include *Lantana camara*, *Mikania micrantha*, *Chromolaena odorata*, and *Ageratina eupatoria* (13). During the past few decades increasing anthropogenic pressures such as unabated linear infrastructure development, uncontrolled tourism, livestock grazing, agricultural expansion and extraction of non-timber forest products (NTFPs) have led to degradation of wildlife habitats and spread of a large number of invasive species (14). It is estimated that the Indian Himalayan region has over 600 IAS species and nearly 50% of them are said to have escaped from agriculture and accidental introduction (12). One of the major causes of spread of IAS in the Himalayan region is climate change (15). It has been observed that plant species of higher elevation are projected to shift higher, due to which a few IAS previously limited to the lower elevations are now shifting towards the higher altitudes in the Himalaya (15, 16).

As a signatory to Convention on Biological Diversity, India has set its National Target in alignment with Aichi Biodiversity Targets. Accordingly, India's Biodiversity Strategy and Action Plan aims to identify the priority species of IAS as well as their invasion pathways so that appropriate management strategies could be formulated. It is realized that a very few baseline studies have been conducted on the current extent and spread of even common species in the Himalayan region. Anthropogenic pressures coupled with climate change makes complex the future prediction of species spread. Several authors e.g., (15, 17-21); have opined that the IAS which were earlier limited to the lower areas are likely to shift towards the higher elevations in the Himalaya. This calls for validation of earlier models as well as multi-locational studies. In this paper, we present the findings of a case study on two IAS, viz., *A. adenophora* and *L. camara* at a landscape level within Indian part of Kailash Sacred Landscape, hereafter referred as Kailash Sacred Landscape (KSL) –India. Major objectives of the study were to predict potential current spread of these species in KSL–India and predict their invasion in future climate change scenarios.

Study Area

The study was carried out in KSL – India that is located between 29°26'35" to 30°35'13" N and 80°01'24" to 81°02'44" E. It is characterized by heterogeneous landscape, wide altitudinal range (ca. 800 to over 7000 m a.s.l.), diverse topography and rich biodiversity (17). This area is contiguous with far-western Nepal and Tibet Autonomous Region (TAR) of China. The study area is spread over 7,120 km² area (Figure 1).

KSL – India has the predominance of diverse natural ecosystems from grasslands to moist sub-tropical broadleaved to temperate oak forests, sub-alpine conifers, high altitude birch forests, while extensive alpine pastures occur in areas >3000 to 4000 m a.s.l beside agro-ecosystems in lower reaches (22-23). The diverse habitats cater to numerous indigenous flora and fauna. Forests in KSL – India fall into two categories - Protected Areas (PAs) and non-protected areas. KSL – India

includes only one legally designated PA (*i.e.* Askot Wildlife Sanctuary) which provides important ecosystem services for the region. However, the non-PA category of forests includes reserve forest and protected forests under the control of the State Forest Department of Uttarakhand managed for an extended period for production of timber and non-timber forest products (NTFPs). KSL – India also includes highly interspersed small forest fragments falling under two broad categories, *i.e.* civil/soyam forests and community forests are under the control of Revenue Department of the State Government. The latter category of community forests, for nearly nine decades or so, has been managed by the local communities and is referred to as ‘*Community Forests*’ or ‘*Van Panchayat Forests*’. This community forest serves as the primary source of livelihood to locals. The KSL – India has experienced a considerable change during recent decades in terms of land use and land cover, along with the development of infrastructure and patterns of natural resource used by the local communities (23- 24).

Methods

Species presence

After the extensive vegetation survey and consultation with experts, two most widely spread IAS, *i.e.* *Ageratina adenophora* (Asteraceae) and *Lantana camara* (Verbenaceae) were selected for the present study. *A. adenophora* is native of Central America (25) and distributed worldwide (26). *L. camara*, native to Tropical America (27) is considered to be one of the most troublesome invasive species worldwide (28). *A. Adenophora*, a perennial, herb produces a large number of tiny seeds which are dispersed by wind and human activities (29). Its ubiquitous property to invade diverse habitats, such as roadsides, riverbanks, forest edges, crop fields, wastelands, and rubbish dump edges makes it fit to invade various ecosystems. *L. camara* is one of the most widely distributed IAS in India (30), resistant to fire and can regrow if burnt (31). It reproduces primarily by seeds as well as through coppice and prefers to grow in degraded habitats.

Though, the species selected for the study are known to negatively affect the wildlife habitats and native vegetation, they are also known to have a few minor uses to local communities. Leaves of *A. adenophora* are used for cattle bedding (32), or its leaf paste is applied to cuts and wounds (33). The wood of *L. camara* is used as fuel (34), source of food for birds (35- 37) and nectar for butterflies and moths (38) as it flowers and fruits throughout the year.

Presence record

We recorded *A. adenophora* and *L. camara* presence between 300 and 2500 m a.s.l covering a wide range of habitats including agricultural lands, grasslands, fallow lands and forests, proximity to major road networks and naturally occurring drainages in different parts of the landscape during 2014 to 2017. A total of 567 and 120 presence locations for *A. adenophora* and *L. camara* respectively, were recorded using handheld Garmin N72 GPS. Of these, 80% locations were used for probability distribution modeling and the rest 20% were used to evaluate the model performance.

Selection of environmental and bioclimatic variables

We used two-time periods (“current” and “2050”) environmental data to model the *A. adenophora* and *L. camara* current and future distribution in KSL – India. A total of 22 environmental variables were used for probability distribution modeling in each time period. We retrieved standard 19 bioclimatic variables for WorldClim version 2 (39). These are the average for the years 1970-2000 with a spatial resolution of 30 seconds (~1 km²). These variables are considered as current bioclimatic variables. We also retrieved 19 bioclimatic variables with a spatial resolution of 30 seconds (~1 km²) for the year 2050 (average for 2041-2060). These are the climate projections as per the Intergovernmental Panel on Climate Change (IPCC) 5th assessment report from Global Climate Models (GCMs) for one Representative Concentration Pathways (RCPs). The data were downscaled and calibrated using WorldClim 1.4 as baseline 'current'

climate (39, 40). These bioclimatic data are also the most recent GCM climate projections that have already been considered in the IPCC fifth assessment report.

We also considered four major static topographic factors for both the “current” and “2050” time frame. These are elevation (ASTER, Digital elevation data from Advanced Space-borne Thermal Emission and Reflection Radiometer), slope, aspect, Euclidian distance from water channels and Euclidian distance from major roads. These variables were assumed to be constant across the entire study period. The details of all the variables used in this study are provided in Appendix 1.

Pair-wise Pearson correlations and Principal Component Analysis

We checked for multi co-linearity with pair-wise correlations for both “current” and “2050” environmental data sets in order to avoid spurious model calibrations (41). It is common a practice to retain only predictors with pair-wise correlation $< |0.7|$ (42). To nullify the correlation structure of the variables and inter-relationship among them, we implemented the Principal Component Analysis (43-44). The PCA reduces the number of orthogonal predictors each of which is the linear combination of 24 original environmental variables. All the exploratory variables were subjected to PCA to extract the factors of significant contribution using *prcomp* function and predicting function of R version 3.3.3 (45). Furthermore, we obtained 24 Principal Components (PCs) in raster format. For both “current” and “2050” data sets, the first six PCs accounted for 99% of the variability and were chosen for further multivariate species distribution modeling (Table 1).

MaxEnt species distribution modeling

Maximum Entropy (*MaxEnt*) distribution modeling was implied to identify the current and future invasion for both, *A. adenophora* and *L. camara* in KSL – India. *MaxEnt* is a machine learning algorithm which is widely used for species distribution modeling that uses presence-only data (46-

47) to predict potential distribution of a particular species in the current as well as future climatic scenarios (48-49).

The occurrence data from the field was collected for both *A. adenophora* ($n = 567$) and *L. camara* ($n = 120$). One thousand background points were drawn randomly across the KSL – India by considering pseudo-absence of both species. We generated four main *MaxEnt* models (two for each species) using the six PCA derived explanatory variables for both ‘current’ and ‘2050’ year.

We implemented k-fold cross-validation (50) to build each main *MaxEnt* model. Firstly, we used standard k-fold cross-validation in our randomly partitioned approach. In our k-fold cross-validation approach, the occurrence localities were divided randomly into 10 bins, where each bin constitutes an equal proportion of sample size (51-52). Then, each model was developed iteratively, using $(k - 1)$ bins for model calibration in each iteration, with the remaining fold retained for evaluation. This is repeated until all the folds were utilized at-least once for model evaluation. *MaxEnt* produces a model based on a series of ‘features’, such as a linear, hinge, quadratic, product, threshold and discrete; we used all except for discrete, as all of our explanatory variables were continuous (53). In the ‘feature’ functions, we set beta multiplier as 0.5 (medium) that affect the smoothness to the model output. We have opted for clamping function in each model as the use of the function is strongly suggested for projecting future species distributions by both Elith et al. (54) and Webber et al. (55).

As we set the models *via* k-fold cross validation ($k = 10$), for each dataset, we used the respective evaluation localities to calculate Area Under the Curve (AUC) of the Receiver Operation Characteristic (ROC), which is a measure of the overall discriminatory ability of the model or to precisely evaluate model performance by plotting its model sensitivity graph (56). We averaged those values from each sub-models of each main-model and represented the averages for each of

the four main *MaxEnt* models. The AUC score closer to 1 indicates that the model had accurately predicted the habitat, while a value ≤ 0.5 indicates low accuracy of the model (57).

Binary habitat invasion maps for the ‘current’ and future ‘2050’ were prepared by using the threshold of maximum training sensitivity plus the specificity logistic threshold. The threshold approach of ‘*Maximum training sensitivity + specificity*’ is a more reliable, restrictive and conservative approach to understand the potential distribution of a species. This particular threshold approach is one of the popular methods either for presence/absence or presence-only data (58). We used ‘*dismo*’ package in R (59) along with a source file developed by Feng et al. (60) to build all *MaxEnt* models as well as to evaluate their performance.

Invasion across the elevation gradient

To check the future spread of *A. adenophora* and *L. camara*, we extracted all elevational range values from a 30 m resolution digital elevation model (ASTER) using the threshold polygons generated from *MaxEnt* main models. Elevational gradients fall within the distribution ranges were re-classified into multiple classes considering 100 m a.s.l intervals. Area invaded by both species in each elevational class were compared between current and 2050 year using paired sample t - test. The distributions of elevational class vs. invaded areas of both *A. adenophora* and *L. camara* were plotted separately for the current year and 2050 to visually assess their spread pattern across the elevational classes.

Results

The pair-wise Pearson correlations were generally higher ($r > |0.70|$) for most of the variable (Figure 2). We were not able to use these variables directly in *MaxEnt* as they showed high correlation, and this avoids the overprediction of suitability in *MaxEnt* models. However, PCA

transformed the 24 environmental variables (each for ‘current’ and ‘2050’) into a composite set of 24 components that are orthogonal and un-correlated to each other. Six out of 24 components demonstrated the cumulative proportion of variance > 0.99 for both the ‘current’ and the year ‘2050’ (Table 1). The first six PCs each from the ‘current’ and ‘2050’ were represented in Appendices 2-3.

Both training and test area under ROC curve (AUC) values were > 0.81 for both species under the ‘current’ and future ‘2050’ climatic conditions (Figure 3, Table 2). The current occurrence locations of *A. adenophora* and *L. camara* where 95.06% and 92.5% respectively falls within their most climatically suitable current distribution ranges. The relative percent contribution of all PCs from both the ‘current’ and the year ‘2050’ are presented in Table 3, and all response curves associated with the above MaxEnt model predictions were given in Appendix 4.

The current and future distributions of the *A. adenophora* and *L. camara* along the elevational gradient of KSL – India are determined by the RCP 8.5 pathway projections for 2050 (Figure 4A-B).

The current distribution of *A. adenophora* ranges between 212 - 2616 m a.s.l, while *L. camara* distributes mostly below 2045 m a.s.l. The potential area that is currently vulnerable to invasion by *A. adenophora* and *L. camara* extends to 1403.7 km² and 1053.1 km², respectively. If the climate change continues to follow the higher emission RCP 8.5 pathway, by 2050, invasion of *A. adenophora* would reach up to 3029 m a.s.l, whereas, *L. camara* will invade up to 3018 m a.s.l. Both *A. adenophora* and *L. camara* will spread across the landscape occupying an area of 2039.7 km² and 1366.5 km², respectively. The area invasion by *A. adenophora* ($t = -7.61$, $df = 28$, $p \leq 0.05$) and *L. camara* ($t = -2.70$, $df = 28$, $p \leq 0.05$) will significantly increase by year 2050 across all designated elevational class (Figure 5).

Discussion

Plant invasion is the emerging area of science that impacts substantial economic and ecological imbalance by altering native plant community composition (61) and depleting native plant species diversity (62) thus affecting ecosystem processes (63-65). The lower and mid ecosystems in the mountains are more vulnerable to invasion by exotic plant species due to the altering climate change and increasing anthropogenic pressure (66). These plant species also exceeded their ranges and many of them have become abundant at higher elevations (67-68). The increasing incidence of invasion in the high-altitude ecosystems (9) poses a significant threat to the native biological diversity of the region and it is expected to expand in the future (21). The use of new and robust methods, especially combining PCA and *MaxEnt* species distribution modeling will reduce the chance of overpredicting species distribution map.

This demonstrated an increasing (both vertical and horizontal) trend of plant invasion in KSL – India. Altering climatic condition and mostly increased temperature have been providing favourable conditions for the invasion and spread of these species in the KSL-India. Joshi et al. (69) already demonstrated that *L. camara* occurrence was recorded widely in dry and exposed slope, forest fringes and deep forests throughout their study sites in the Indian butter tree (*Diploknema butyracea*) forest which is one of the economically, spiritually and culturally significant trees in the region with its great importance for local livelihood. Looking forward, *L. camara* might also affect the regeneration of such economically important trees in the area. Elevation was the most crucial factor in the current and future distribution. It appeared that in future both of these IAS would colonize more vigorously in the landscape where currently they are not present. This invasion of both species is likely to happen if climate change continues to follow the higher emission RCP 8.5 pathway. Furthermore, *L. camara* will spread within a similar altitudinal range where it is currently absent. According to Singh et al. (70), *A. adenophora* prefers

north-facing moist slopes along roadsides, forest fringe, cliffs, near water bodies, including agricultural and degraded lands and sub-tropical and warmer temperate landscapes. It also invades natural habitats, Deodar, Banj oak- Chir pine and mixed forests, degraded grassland areas and sometimes invading pure stands of pristine temperate broadleaved oak (*Quercus leucotrichophora*) forest representing climax vegetation between 1000 – 3500 m a.s.l. in KSL – India. Our study also showed a similar trend, as this species will reach 968 m a.s.l higher than its current distribution range.

Our predictive models also depicted high infestation and expansion in the present climatic scenario by *A. adenophora* and *L. camara* in the lower parts of KSL – India in western Himalaya. The MaxEnt model also predicted that the landscape provides more suitable conditions for the infestation of *A. adenophora* as compared to *L. camara* even at its current climatic condition. Both the species will expand their habitats with the increase in temperature and human activity in the landscape. Our study demonstrated similar results as compared to previous studies (71-73), but in a more robust and spatially explicit manner.

Our study has come up with current pattern of distribution and established future scenarios of invasion by two highly obnoxious IAS within KSL – India. The distribution maps generated in this exercise would be of much help in further conservation planning and participatory management of forest / grassland management. The findings of our study can be used in identification of vulnerable areas and important localities which are likely to be infested by these species. Habitat restoration and community-based monitoring at local scale involving Biodiversity Management Committee (BMC) would be one of the logical steps forward.

Acknowledgments

The authors are thankful to the Director of the Wildlife Institute of India for his support. The study was carried out under Kailash Sacred Landscape Conservation and Development Initiative (KSCDI) funded by ICIMOD, Nepal and National Mission on Himalayan Studies (NMHS) funded by Ministry of Environment, Forests and Climate Change, New Delhi. We would also like to acknowledge our gratitude to the local communities for their help, support, and cooperation during this study.

References

1. Early R, Bradley BA, Dukes JS, et al. Global threats from invasive alien species in the twenty-first century and national response capacities. *Nature communications*. 2016; 7(1): 1-9. <https://doi.org/10.1038/ncomms12485>.
2. IPBES. Summary for policy makers of the regional assessment report on biodiversity and ecosystem services for Asia and the Pacific of the Intergovernmental Science – Policy Platform on Biodiversity and Ecosystem Services. IPBES Secretariat. Bonn, Germany; 2018.p. 41.
3. Rai PK, Singh JS. Invasive alien plant species: Their impact on environment, ecosystem services and human health. *Ecological Indicators*. 2020; 111: p.106020. <https://doi.org/10.1016/j.ecolind.2019.106020>.
4. Dukes JS, Mooney HA. Does global change increase the success of biological invaders? *Trends in Ecology Evolution*. 1999; 14(4): 135–139. [https://doi.org/10.1016/S0169-5347\(98\)01554-7](https://doi.org/10.1016/S0169-5347(98)01554-7).
5. Mack RN, Simberloff D, Mark Lonsdale W, et al. Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological applications*. 2000; 10(3): 689–710. [https://doi.org/10.1890/1051-0761\(2000\)010\[0689:BICEGC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0689:BICEGC]2.0.CO;2).

6. Lin W, Zhou G, Cheng X, et al. Fast economic development accelerates biological invasions in China. *PLoS One*. 2007; 2(11): e1208. <https://doi.org/10.1371/journal.pone.0001208>.
7. Vila M, Weiner J. Are invasive plant species better competitors than native plant species? – evidence from pair-wise experiments. *Oikos*. 2004; 105(2): 229–238. <https://doi.org/10.1111/j.0030-1299.2004.12682.x>.
8. Hellmann JJ, Byers JE, Bierwagen BG, et al. Five potential consequences of climate change for invasive species. *Conservation biology*. 2008; 22(3): 534–543. <https://doi.org/10.1111/j.1523-1739.2008.00951.x>.
9. Thapa S, Chitale V, Rijal SJ, et al. Understanding the dynamics in distribution of invasive alien plant species under predicted climate change in Western Himalaya. *PloS One*. 2018; 13(4): e0195752. <https://doi.org/10.1371/journal.pone.0195752>.
10. Kohli RK, Dogra KS, Batish DR, et al. Impact of Invasive Plants on the Structure and Composition of Natural Vegetation of Northwestern Indian Himalayas. *Weed Technology*. 2004; 18(sp1): 1296–1301. [https://doi.org/10.1614/0890-037X\(2004\)018](https://doi.org/10.1614/0890-037X(2004)018) [1296: IOIPOT]2.0.CO;2.
11. Tylianakis JM, Didham RK, Bascompte J, et al. Global change and species interactions in terrestrial ecosystems. *Ecology Letters*. 2008; 11(12): 1351–1363. <https://doi.org/10.1111/j.1461-0248.2008.01250.x>.
12. Sekar, KC. Invasive alien plants of Indian Himalayan region—diversity and implication American. *Journal of Plant Sciences*. 2012; 3(2): 177. <https://doi.org/10.4236/ajps.2012.32021>.
13. Pathak R, Negi VS, Rawal RS, Bhatt ID. Alien plant invasion in the Indian Himalayan Region: state of knowledge and research priorities. *Biodiversity and Conservation*. 2019; 28: 3073–3102. <https://doi.org/10.1007/s10531-019-01829-1>.
14. Babu S, Love A, Babu CR. Ecological restoration of lantana-invaded landscapes in Corbett Tiger Reserve, India. *Ecological Restoration*. 2009; 27(4): 467–477. <https://doi.org/10.3368/er.27.4.467>.

15. Telwala Y, Brook BW, Manish K, et al. Climate-induced elevational range shifts and increase in plant species richness in a Himalayan biodiversity epicenter. PloS one. 2013; 8(2): e57103. <https://doi.org/10.1371/journal.pone.0057103>.
16. Shrestha BB, Joshi S, Bisht N, et al. Inventory and impact assessment of invasive alien plant species in Kailash Sacred Landscape: ICIMOD working paper 2018/2. ICIMOD Working Paper id: 12674, e Social Sciences; 2018. <http://lib.icimod.org/record/33744>.
17. Oli KP, Zomer R; (eds.) Kailash sacred landscape conservation initiative – Feasibility assessment report. Kathmandu: ICIMOD, Nepal; 2011. p. 1-92.
18. Taylor S, Kumar L, Reid N. Impacts of climate change and land-use on the potential distribution of an invasive weed: a case study of *Lantana camara* in Australia. Weed Research. 2012; 52(5): 391–401. <https://doi.org/10.1111/j.1365-3180.2012.00930.x>
19. Goncalves E, Herrera I, Duarte M, et al. Global invasion of *Lantana camara*: has the climatic niche been conserved across continents? PLoS One. 2014; 9(10): e111468. <https://doi.org/10.1371/journal.pone.0111468>.
20. Uddin K, Chaudhary S, Chettri N, et al. The changing land cover and fragmenting forest on the Roof of the World: A case study in Nepal's Kailash Sacred Landscape. Landscape and Urban Planning. 2015; 141: 1–10. <https://doi.org/10.1016/j.landurbplan.2015.04.003>.
21. Mungi NA, Coops NC, Ramesh K, et al. How global climate change and regional disturbance can expand the invasion risk? Case study of *Lantana camara* invasion in the Himalaya. Biological Invasions. 2018; 20(7): 1849–1863. <https://doi.org/10.1007/s10530-018-1666-7>.
22. Singh SP, Bassignana-Khadka I, Singh Karky B, et al. Climate change in the Hindu Kush-Himalayas: the state of current knowledge International Centre for Integrated Mountain Development (ICIMOD), Nepal; 2011. p. 1-88.
23. Singh G, Sarkar MS, Pandey A, et al. Quantifying Four Decades of Changes in Land Use and Land Cover in India's Kailash Sacred Landscape: Suggested Option for Priority Based Patch Level Future Forest Conservation. Journal of the Indian Society of Remote Sensing. 2018; 46(10): 1625-1635. <https://doi.org/10.1007/s12524-018-0817-8>.

24. Negi VS, Pathak R, Sekar KC, et al. Traditional knowledge and biodiversity conservation: a case study from Byans Valley in Kailash Sacred Landscape, India. *Journal of Environmental Planning and Management*. 2018; 61(10): 1722-1743. <https://doi.org/10.1080/09640568.2017.1371006>.
25. Datta A, Kühn I, Ahmad M, et al. Processes affecting altitudinal distribution of invasive *Ageratina adenophora* in western Himalaya: The role of local adaptation and the importance of different life-cycle stages. *PloS one*. 2017; 12(11): e0187708. <https://doi.org/10.1371/journal.pone.0187708>.
26. Cronk QC, Fuller JL. *Plant invaders: the threat to natural ecosystems*. London: Chapman and Hall; 1995. P. 241.
27. Holm LG, Plucknett DL, Pancho JV, et al. *The world's worst weeds: Distribution and biology*. University press of Hawaii; 1977. p. 609.
28. Kohli RK, Batish DR, Singh HP, et al. Status, invasiveness and environmental threats of three tropical American invasive weeds (*Parthenium hysterophorus* L, *Ageratum conyzoides* L, *Lantana camara* L.) in India. *Biological Invasions*. 2006; 8(7): 1501–1510. <https://doi.org/10.1007/s10530-005-5842-1>.
29. Sang W, Zhu L, Axmacher JC. Invasion pattern of *Eupatorium adenophorum* Spreng in southern China. *Biological Invasions*. 2010; 12(6): 1721–1730. <https://doi.org/10.1007/s10530-009-9584-3>.
30. Sundaram B, Krishnan S, Hiremath AJ, et al. Ecology and impacts of the invasive species, *Lantana camara*, in a social-ecological system in South India: perspectives from local knowledge. *Human ecology*. 2012; 40(6): 931–942. <https://doi.org/10.1007/s10745-012-9532-1>.
31. Hiremath AJ, Sundaram B. The fire-lantana cycle hypothesis in Indian forests. *Conservation and Society*. 2005; 3(1): 26-42. <http://eprints.atree.org/id/eprint/22>.
32. Shrestha TB. *Development ecology of the Arun river basin in Nepal Country report on biological diversity*, ICIMOD, Kathmandu, Nepal; 1989. p. 1-121.

33. Joseph J, Kharkongor P. A preliminary ethno-botanical survey in the Khasi and Jaintia hills, Meghalaya. In: S.K Jain (Ed.): Glimpses of Indian Ethnobotany. New Delhi: Oxford IBH Publishing Co; 1981. p. 115–123.
34. Sharma OP, Makkar HPS, Dawra RK (1988) A review of the noxious plant *Lantana camara*. Toxicon 26(11): 975–987. [https://doi.org/10.1016/0041-0101\(88\)90196-1](https://doi.org/10.1016/0041-0101(88)90196-1).
35. Corlett RT. Frugivory and seed dispersal by birds in Hong Kong shrub-land. Forktail 1998; 13: 23–27.
36. Bhatt D, Kumar A. Foraging ecology of Red-vented bulbul *Pycnonotus cafer* in Haridwar, India. Forktail. 2001; 17: 109–109.
37. Aruna R, Balasubramanian P. Fruiting phenology and avian frugivory on *Lantana camara* in a mixed dry deciduous forest, Western Ghats, Tamilnadu, India. International Journal of Fauna and Biological Studies. 2017; 4(4): 179–182.
38. Day MD, Wiley CJ, Playford J, et al. Lantana: current management status and future prospects. Australian Centre for International Agricultural Research. 2003; 5: 1–20.
39. Fick SE, Hijmans RJ. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology. 2017; 37(12): 4302–4315. <https://doi.org/10.1002/joc.5086>.
40. Hijmans RJ, Cameron SE, Parra JL, et al. Very high-resolution interpolated climate surfaces for global land areas. International Journal of Climatology: A Journal of the Royal Meteorological Society. 2005; 25(15): 1965–1978. <https://doi.org/10.1002/joc.1276>.
41. Guisan A, Thuiller W. Predicting species distribution: offering more than simple habitat models. Ecology Letters. 2005; 8(9): 993–1009. <https://doi.org/10.1111/j.1461-0248.2005.00792.x>.
42. Carvalheiro, L. G., Kunin, W. E., Keil, P., Aguirre-Gutiérrez, J., Ellis, W. N., Fox, R., ... & Van de Meutter, F. (2013). Species richness declines and biotic homogenisation have slowed down for NW-European pollinators and plants. *Ecology letters*, 16(7), 870-878.

43. Jolliffe I.T (Ed.) Principal Component Analysis (2nd Eds.) Aberdeen, UK: Springer; 2002. p. 488.
44. Cruz-Cárdenas, G., Villaseñor, J. L., López-Mata, L., Martínez-Meyer, E., & Ortiz, E. (2014). Selección de predictores ambientales para el modelado de la distribución de especies en Maxent. *Revista Chapingo. Serie ciencias forestales y del ambiente*, 20(2), 187-201.
45. Team R Core. R: A Language and Environment for Statistical Computing. Vienna, Austria. 2018 Jan 25. Available from: <https://www.R-project.org/>.
46. Phillips SJ, Anderson RP, Schapire RE. Maximum entropy modeling of species geographic distributions. *Ecological modelling*. 2006; 190(3–4): 231-259. <https://doi.org/10.1016/j.ecolmodel.2005.03.026>.
47. Elith J, Phillips SJ, Hastie T, et al. A statistical explanation of MaxEnt for ecologists. *Diversity and distributions*. 2011; 17(1): 43–57. <https://doi.org/10.1111/j.1472-4642.2010.00725.x>.
48. Loarie SR, Carter BE, Hayhoe K, et al. Climate change and the future of California's endemic flora. *PloS One*. 2008; 3(6): e2502. <https://doi.org/10.1371/journal.pone.0002502>.
49. Remya K, Ramachandran A, Jayakumar S. Predicting the current and future suitable habitat distribution of *Myristica dactyloides* Gaertn using MaxEnt model in the Eastern Ghats, India. *Ecological engineering*. 2015; 82: 184–188. <https://doi.org/10.1016/j.ecoleng.2015.04.053>.
50. Peterson AT, Soberón J, Pearson RG, et al. *Ecological niches and geographic distributions*. Princeton, UK: Princeton University Press. 2011. p. 328.
51. Boyce MS, Vernier PR, Nielsen SE, et al. Evaluating resource selection functions. *Ecological modelling*. 2002; 157(2-3): 281–300. [https://doi.org/10.1016/S0304-3800\(02\)00200-4](https://doi.org/10.1016/S0304-3800(02)00200-4).

52. Lehmann A, Overton JM, Leathwick JR. GRASP: generalized regression analysis and spatial prediction. *Ecological modelling*. 2002; 157(2-3): 189–207. [https://doi.org/10.1016/S0304-3800\(02\)00195-3](https://doi.org/10.1016/S0304-3800(02)00195-3).
53. Phillips SJ, Dudík M. Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography*. 2008; 31(2): 161–175. <https://doi.org/10.1111/j.0906-7590.2008.5203.x>.
54. Elith J, Kearney M, Phillips S. The art of modelling range-shifting species. *Methods in ecology and evolution*. 2010; 1(4): 330–342. <https://doi.org/10.1111/j.2041-210X.2010.00036.x>.
55. Webber BL, Yates CJ, Le Maitre DC, et al. Modelling horses for novel climate courses: insights from projecting potential distributions of native and alien Australian acacias with correlative and mechanistic models. *Diversity and Distributions*. 2011; 17(5): 978–1000. <https://doi.org/10.1111/j.1472-4642.2011.00811.x>.
56. Peterson AT, Papeş M, Soberón J. Rethinking receiver operating characteristic analysis applications in ecological niche modeling. *Ecological modelling*. 2008; 213(1): 63 –72. <https://doi.org/10.1016/j.ecolmodel.2007.11.008>.
57. Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology*. 1982; 143(1): 29–36. <https://doi.org/10.1148/radiology.143.1.7063747>.
58. Liu C, Berry PM, Dawson TP, et al. Selecting thresholds of occurrence in the prediction of species distributions. *Ecography*. 2005; 28(3): 385–393. <https://doi.org/10.1111/j.0906-7590.2005.03957.x>.
59. Hijmans RJ, Phillips S, Leathwick J, et al. Package ‘dismo’ Circles. 2017; 9(1): 1–68.
60. Feng X, Gebresenbet F, Walker C. Shifting from closed-source graphical-interface to open-source programming environment: a brief tutorial on running Maxent in R (No e3346v1). 2017; PeerJ Preprints. <https://doi.org/10.7287/peerj.preprints.3346v1>.

61. Gordon DR. Effects of invasive, non-indigenous plant species on ecosystem processes: lessons from Florida. *Ecological Applications*. 1998; 8(4): 975–989. [https://doi.org/10.1890/1051-0761\(1998\)008\[0975:EOINIP\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1998)008[0975:EOINIP]2.0.CO;2).
62. Hejda M, Pyšek, P Jarošík V. Impact of invasive plants on the species richness, diversity and composition of invaded communities. *Journal of Ecology*. 2009; 97(3): 393–403. <https://doi.org/10.1111/j.1365-2745.2009.01480.x>.
63. Weidenhamer, JD, Callaway RM. Direct and indirect effects of invasive plants on soil chemistry and ecosystem function. *Journal of Chemical Ecology*. 2010; 36(1): 59–69. <https://doi.org/10.1007/s10886-009-9735-0>.
64. Crooks JA. Characterizing ecosystem-level consequences of biological invasions: the role of ecosystem engineers. *Oikos*. 2002; 97(2): 153–166. <https://doi.org/10.1034/j.1600-0706.2002.970201.x>.
65. Dukes JS, Mooney HA. Disruption of ecosystem processes in western North America by invasive species. *Revista Chilena de Historia Natural*. 2004; 77(3): 411–437. <http://dx.doi.org/10.4067/S0716-078X2004000300003>.
66. Priyanka N, Joshi PK. A review of *Lantana camara* studies in India. *International Journal of Scientific and Research Publications*. 2013; 3(10): 1–11.
67. Alexander JM, Lembrechts JJ, Cavieres, LA, et al. Plant invasions into mountains and alpine ecosystems: current status and future challenges. *Alpine Botany*. 2016; 126(2): 89–103. <https://doi.org/10.1007/s00035-016-0172-8>.
68. Pauchard A, Kueffer C, Dietz H, et al. Ain't no mountain high enough: plant invasions reaching new elevations. *Frontiers in Ecology and the Environment*. 2009; 7(9): 479–486. <https://doi.org/10.1890/080072>.
69. Joshi NC, Chaudhary A, Rawat GS. Cheura (*Diploknema butyracea*) as a livelihood option for forest-dweller tribe (Van-Raji) of Pithoragarh, Uttarakhand, India. *ESSENCE – International Journal for Environmental Rehabilitation and Conservation IX*. 2018; 1: 134–141. <https://doi.org/10.31786/09756272.18.9.1.116>.

70. Singh G, Padalia H, Rai ID, et al. Spatial extent and conservation status of Banj oak (*Quercus leucotrichophora* A Camus) forests in Uttarakhand, Western Himalaya. *Tropical Ecology*. 2016; 57(2): 255-262.
71. Reese GC, Wilson KR, Hoeting JA, et al. Factors affecting species distribution predictions: a simulation modeling experiment. *Ecological Applications*. 2005; 15(2): 554–564. <https://doi.org/10.1890/03-5374>.
72. Hernandez PA, Graham CH, Master LL, et al. The effect of sample size and species characteristics on performance of different species distribution modeling methods. *Ecography*. 2006; 29(5): 773–785. <https://doi.org/10.1111/j.0906-7590.2006.04700.x>.
73. Wisz, MS, Hijmans, RJ, Li J, et al. Predicting Species Distributions Working Group: Effects of sample size on the performance of species distribution models. *Diversity and Distributions*. 2008; 14(5): 763–773. <https://doi.org/10.1111/j.1472-4642.2008.00482.x>.

Figure 1: Geographical location of the study area with sampling locations of *Lantana camara* and *Ageratina adenophora*.

Figure 3: Receiver operating characteristic (ROC) curve for the predicted distribution *MaxEnt* models averaged over the ten replicate runs: *Ageratina adenophora* (the “current” year – A; year 2050 – B) and *Lantana camara* (the “current” year – C; year 2050 – D).

Figure 4: Predicted distribution *MaxEnt* models averaged over the ten replicate runs. Orange and Red polygons represent maximum training sensitivity + specificity logistic threshold for the distribution of (A) *Ageratina adenophora* and (B) *Lantana camara* in the “current” and year 2050.

Figure 5: The predicted spread (in Km²) of *Ageratina adenophora* (A) and *Lantana camara* (B) in different elevational classes of KSL – India during the current year and the year 2050.

Table 1 Representation of the first seven principal components derived from all 24 original environmental variables along with the proportion of variance in environmental variables explained by each principal component and their cumulative proportion of variance.

Principal components	The “current” year		Year 2050	
	Proportion of Variance	Cumulative Proportion	Proportion of Variance	Cumulative Proportion
PC1	0.98890	0.98890	0.98880	0.98880
PC2	0.00810	0.99700	0.00810	0.99700
PC3	0.00276	0.99979	0.00277	0.99973
PC4	0.00013	0.99992	0.00015	0.99988
PC5	0.00007	0.99998	0.00008	0.99996
PC6	0.00001	0.99999	0.00003	0.99999
PC7	0.00000	1.00000	0.00000	1.00000

Table 2 The area under the receiver operating curve (AUC) score of MaxEnt models for both *Ageratina adenophora* and *Lantana Camara*.

Species	The “current” year		Year 2050	
	Test AUC	Training AUC	Test AUC	Training AUC
<i>Ageratina Adenophora</i>	0.81	0.82	0.81	0.82
<i>Lantana Camara</i>	0.93	0.93	0.92	0.93

Table 3 Relative contributions of each principal component are used in all the *MaxEnt* models.

	Contribution of <i>Ageratina adenophora</i> (%)		Contribution of <i>Lantana camara</i> (%)	
	The “current” year	Year 2050	The “current” year	Year 2050
PC1	83.00	86.10	62.8	65.2
PC2	00.00	00.10	0.1	0.2
PC3	01.10	02.40	9.1	15.6
PC4	04.60	04.60	2.1	1.1
PC5	07.30	05.40	16.7	15.8
PC6	04.00	01.50	9.1	2.2

Appendix 1 Description of variables of three time periods used to model *Ageratina adenophora* and *Lantana camara* distribution across the KSL-India.

Environmental variables		Data Source	
Bioclimatic variables			
Abbreviations	Description of variables	Current	2050
BIO1	Annual Mean Temperature	WorldClim version 2: average monthly climate data (average for 1970-2000)	WorldClim 1.4 downscaled (CMIP5) data (average for 2041-2060)
BIO2	Mean Diurnal Range (Mean of monthly (max temp - min temp))		
BIO3	Isothermality (BIO2/BIO7) (* 100)		
BIO4	Temperature Seasonality (standard deviation *100)		
BIO5	Max Temperature of Warmest Month		
BIO6	Min Temperature of Coldest Month		
BIO7	Temperature Annual Range (BIO5-BIO6)		
BIO8	Mean Temperature of Wettest Quarter		
BIO9	Mean Temperature of Driest Quarter		
BIO10	Mean Temperature of Warmest Quarter		
BIO11	Mean Temperature of Coldest Quarter		
BIO12	Annual Precipitation		
BIO13	Precipitation of Wettest Month		
BIO14	Precipitation of Driest Month		
BIO15	Precipitation Seasonality (Coefficient of Variation)		
BIO16	Precipitation of Wettest Quarter		
BIO17	Precipitation of Driest Quarter		
BIO18	Precipitation of Warmest Quarter		
BIO19	Precipitation of Coldest Quarter		
Topographic variables			
DEM	digital elevation model	Global Digital Elevation Model Version 2 (GDEM V2)	
EUCL_Water	euclidean distance from primary and secondary water bodies	OpenStreetMapData (http://openstreetmapdata.com)	
EUCL_Road	euclidean distance from road	OpenStreetMapData (http://openstreetmapdata.com)	
Slope	Angle of slope in degree	Derivative of digital elevation model	
Aspect	Slope faces in 3-dimensional space	Derivative of digital elevation model

Appendix 2 First six principal components as raster layers generated from the 24 environmental layers of current years.

Appendix 3 First six principal components as raster layers generated from the 24 environmental layers of 2050 years.

Appendix 4 These curves show how each PCs affects the MaxEnt prediction. The curves show how the predicted probability of presence changes as each PC is varied, keeping all other environmental variables at their average sample value. The curves show the mean response of the 10 replicates of each *MaxEnt* runs. (A-F): Response curves of *Ageratina adenophora* for current year, (G-L): Response curves of *Ageratina adenophora* for the year 2050, (M-R): Response curves of *Lantana camara* for current year, (S-X) Response curves of *Lantana camara* for the year 2050.