Ubiquitylome Analysis Reveals a Central Role for the Ubiquitin-Proteasome System in Plant Innate Immunity

Xiyu Ma^{1*}, Chao Zhang^{2*}, Do Young Kim^{3,4*}, Yanyan Huang¹, Ping He¹, Richard D. Vierstra^{3,5#}, and Libo Shan^{1,2#}

Author Affiliations:

¹Department of Biochemistry and Biophysics, Texas A&M University, College Station, TX 77843, USA

²Department of Plant Pathology and Microbiology, Texas A&M University, College Station, TX 77843, USA

³Department of Genetics, 425-G Henry Mall, University of Wisconsin-Madison, Madison, WI 53706, USA

⁴Advanced Bio Convergence Center, Pohang Technopark, Gyeong-Buk, South Korea, 37668

⁵Department of Biology, Washington University in St. Louis, St. Louis, MO 63130, USA

*These authors contribute equally.

#Corresponding authors:

Libo Shan

Tel: 979-845-8818; E-mail: lshan@tamu.edu

Richard D. Vierstra

Tel: 314-935-5058; E-mail: rdvierstra@wustl.edu

One-sentence summary: Proteome-wide catalogs of ubiquitylated proteins revealed a rapid engagement of the ubiquitin-proteasome system in Arabidopsis innate immunity.

Keywords: innate immunity, disease resistance, ubiquitylome, protein ubiquitylation, proteasome, mass spectrometry, Arabidopsis

Manuscript information: 47 pages and 8 figures.

ABSTRACT

Protein ubiquitylation profoundly expands proteome functionality and diversifies cellular signaling processes, with recent studies providing ample evidence for its importance to plant immunity. To gain a proteome-wide appreciation of ubiquitylome dynamics during immune recognition, we employed a two-step affinity enrichment protocol based on a 6His-tagged ubiquitin (Ub) variant coupled with high sensitivity mass spectrometry to identify *Arabidopsis* proteins rapidly ubiquitylated upon plant perception of the microbe-associated molecular pattern (MAMP) peptide flg22. The catalog from two-week-old seedlings treated for only 30 minutes with flg22 contained nearly 1,000 conjugates, 150 Ub footprints, and all seven types of Ub linkages, and included previously uncharacterized conjugates of immune components, such as RECEPTOR-LIKE KINASE 1 (RKL1) shown to negatively regulate plant immunity. *In vivo* ubiquitylation assays confirmed modification of several candidates upon immune elicitation, and revealed distinct modification patterns and dynamics for key immune components, including poly- and monoubiquitylation, as well as induced or reduced levels of ubiquitylation. Gene ontology and network analyses of the collection also uncovered rapid modification of the Ub-proteasome system itself, suggesting a critical auto-regulatory loop necessary for an effective MAMP-triggered immune response and subsequent disease resistance. Included targets were UBIQUITIN-CONJUGATING ENZYME 13 (UBC13) and proteasome component REGULATORY PARTICLE NON-ATPASE SUBUNIT 8b (RPN8b), whose subsequent biochemical and genetic analyses implied negative roles in immune elicitation. Collectively, our proteomic analyses further strengthened the connection between ubiquitylation and flg22-based immune signaling, identified novel components and pathways regulating plant immunity, and increased the database of ubiquitylated substrates in plants.

INTRODUCTION

Post-translational modifications (PTMs), such as phosphorylation, methylation, acetylation, glycosylation, myristoylation, and ubiquitylation diversify protein behaviors, thus increasing the functionality and regulation of the proteome. As one of the most abundant PTMs, ubiquitylation is involved in nearly all physiological and signaling processes in plants, including hormone perception, photomorphogenesis, circadian rhythms, self-incompatibility, and defense against biotic and abiotic challenges (Guerra and Callis, 2012; Trujillo and Shirasu, 2010; Vierstra, 2009; Zhou and Zeng, 2017). Attachment of the 76-amino acid ubiquitin (Ub) moiety occurs most commonly to accessible lysines (K) within the target via an isopeptide bond that links to the carboxyl (C)-terminal glycine of Ub.

This conjugation is typically mediated through a stepwise enzymatic cascade consisting of a Ub-activating enzyme (UBA, or E1), a Ub-conjugating enzyme (UBC, or E2), and finally, a Ub-protein ligase (or E3), the latter of which has greatly expanded and diversified in plants to recognize a plethora of substrates. Additional Ubs can become attached to one or more of the seven lysine residues (K6, K11, K27, K29, K31, K48, and K63) or the amino (N)-terminal amide nitrogen within previously attached Ubs, resulting in a myriad of poly-Ub chain architectures that provide additional functional diversity (Komander and Rape, 2012; Vierstra, 2009). The most common internal linkage involves lysine 48 (K48) whose poly-Ub chains mainly target proteins for breakdown by the 26S proteasome, a multisubunit proteolytic complex that recognizes the Ub moieties (Vierstra, 2009; Zhou and Zeng, 2017). Although not as well understood in plants (Paez Valencia et al., 2016; Romero-Barrios and Vert, 2018; Zhou and Zeng, 2017), other types of poly-Ub linkages, involving K6, K11, K27, K29, K33, or K63, and monoubiquitylation in which only a single Ub is attached, have also been connected to many intracellular events, including protein trafficking, signal transduction, protein-protein interactions, and protein degradation through autophagy (Komander and Rape, 2012).

In recent years, a myriad of plant defenses against a range of pathogens has become apparent. The front line of defense is to block pathogen entry by erecting physical barriers, such as wax layers, cell walls, cuticular lipids, cutin, and callose, which can be strengthened during infection (Thordal-Christensen, 2003). As a second line of defense, plants rely on innate immune responses to fend off infections internally, which are elicited upon recognition of pathogen- or microbe-associated molecular patterns (PAMPs/MAMPs) for pattern-triggered immunity (PTI), or pathogen-encoded effectors for effector-triggered immunity (ETI), which are encoded within the pathogen (Spoel and Dong, 2012; Zhou and Zhang, 2020). The host receptors known to detect these patterns/effectors are either receptor-like kinases (RLKs) or receptor-like proteins (RLPs) (Böhm et al., 2014; Couto and

Zipfel, 2016). The best understood is the RLK receptor FLAGELLIN SENSING 2 (FLS2) and its co-receptor BRASSINOSTEROID INSENSITIVE 1-ASSOCIATED KINASE 1 (BAK1) that recognize the bacterial 22-amino-acid flagellin epitope - flg22. Subsequently, PTI signaling is relayed through receptor-like cytoplasmic kinases (RLCKs), the mitogen-activated protein kinase (MAPK) cascade, and transcription factors, that trigger diverse immune responses, including the production of reactive oxygen species (ROS) and the stress hormone ethylene (Couto and Zipfel, 2016; Yu et al., 2017). Pathogen effectors are recognized directly or indirectly by NUCLEOTIDE-BINDING SITE LEUCINE-RICH REPEAT (NBS-LRR) proteins (NLRs), which often leads to localized cell death at the infection sites known as the hypersensitive response (HR) (Cui et al., 2015). The defense hormone salicylic acid (SA) is indispensable for NLR-triggered HR (Cui et al., 2015).

Protein ubiquitylation is also emerging as a dynamic process in both PTI and ETI signaling (Adams and Spoel, 2018; Cheng and Li, 2012; Zhou and Zeng, 2017). As examples, the MAMP receptor FLS2 is ubiquitylated upon ligand engagement by the E3s PLANT U-BOX 12 (PUB12) and PUB13, resulting in FLS2 degradation (Lu et al., 2011; Zhou et al., 2015), while PUB13 also directs the ubiquitylation and degradation of LYSIN MOTIF RECEPTOR KINASE 5 (LYK5), an RLK that recognizes the fungal cell wall polysaccharide chitin (Liao et al., 2017). The RLCK *BOTRYTIS*-INDUCED KINASE 1 (BIK1), which provides a convergent signaling hub downstream of multiple MAMP receptors, is strongly regulated by Ub addition (Ma et al., 2020; Wang et al., 2018). Two U-box E3s, PUB25, and PUB26 polyubiquitylate BIK1 to control its steady-state levels, while the RING-type E3 ligases, RING-H2 FINGER A3A (RHA3A) and RHA3B, monoubiquitylate BIK1 to direct its endocytosis and signaling activation (Ma et al., 2020; Wang et al., 2018). Additionally, the PUB22, PUB23, and PUB24 E3s work collectively to negatively regulate PTI responses through modification of EXOCYST SUBUNIT EXO-70 FAMILY PROTEIN B2 (EXO70B2), which promotes vesicle trafficking (Stegmann et al., 2012). Levels of several NLR proteins, including RESISTANT TO *PSEUDOMONAS SYRINGAE* 2 (RPS2) and SUPPRESSOR OF *npr1-1* CONSTITUTIVE 1 (SNC1), are also regulated by CONSTITUTIVE EXPRESSOR OF PR GENES 1 (CPR1), which is the target-recognition F-Box subunit within an SKP1-CULLIN1-F-BOX (SCF) E3 complex (Cheng et al., 2011; Gou et al., 2012). Despite these examples, a systematic investigation of ubiquitylation during plant immune activation is currently lacking.

Recent advance in proteomics has enabled high-throughput and comprehensive views of ubiquitylomes under specific physiological conditions. The first catalog was generated with yeast in which wild-type Ub was replaced by a 6His-tagged variant to enable enrichment of ubiquitylated proteins by nickel-nitrotrilotriacetic acid (Ni-NTA) affinity

chromatography under denaturing conditions, followed by liquid chromatography-mass spectrometric analysis (LC-MS) of the trypsinized preparations (Peng et al., 2003). Sarraco et al. (2009) then adopted this strategy for plants through the creation of a transgenic *Arabidopsis* line that constitutively expresses a hexa-6His-Ub concatamer which is post-translationally processed into functional 6His-Ub monomers. Subsequently, Ub-binding domains were exploited for the additional enrichment needed for more complex proteomes (Maor et al., 2007). Further improvements included Tandem-repeated Ub-Binding Entities (TUBEs) in which natural Ub-binding domains were oligomerized to increase enrichment and yield (Hjerpe et al., 2009). By combining TUBEs with the Ni-NTA affinity chromatography, stringent purifications were possible, which enabled the confident detection of over 1,000 ubiquitylated proteins in *Arabidopsis* (Aguilar-Hernandez et al., 2017; Kim et al., 2013). Notably, a number of Ub targets associated with disease resistance were identified.

Here, we employed this improved two-step TUBEs purification approach coupled with deep MS analysis of *Arabidopsis* seedlings before and shortly after exposure to flg22. The resulting ubiquitylome catalog revealed extensive proteome-wide changes soon after MAMP perception and uncovered a large number of ubiquitylated candidates potentially connected to plant innate immunity. We further observed by *in vivo* ubiquitylation assays distinct ubiquitylation patterns and dynamics for several immunity components, including key regulators that were modified by multiple ubiquitylation events. One target was the previously uncharacterized receptor protein kinase RECEPTOR-LIKE KINASE 1 (RKL1) that appears to negatively impact plant immunity. Interestingly, several Ub-proteasome pathway components were also confirmed as Ub targets upon PTI immune elicitation, including the E2 UBC13 and proteasome subunit REGULATORY PARTICLE NON-ATPASE SUBUNIT 8b (RPN8b). Taken together, our study identified novel connections between the Ub-proteasome system (UPS) and pathogen immunity as well as provided a valuable resource to further study this PTM during immune activation.

RESULTS

TUBEs enrichment of ubiquitylated *Arabidopsis* proteins upon flg22 treatment

To gain a global perspective of ubiquitylation during PTI elicitation, we employed a two-step TUBEs purification protocol to enrich for ubiquitylated proteins soon after exposure of two-week-old *A. thaliana* ecotype Col-0 seedlings to flg22 (Kim et al., 2013), which was based on a transgenic line harboring a synthetic *UBQ* gene encoding six 6His-Ub repeats expressed under the control of the Cauliflower mosaic virus (CaMV) 35S promoter (Sarraco et al., 2009). The polymer is processed by endogenous deubiquitylases (DUBs) to release tagged and fully functional Ub monomers (Isono and Nagel, 2014). In the first step, extracts from

hexa-6His-UBQ seedlings either untreated or treated for only 30 min to flg22 were prepared in a non-denaturing buffer containing the non-ionic detergent Triton X-100 and a protease inhibitor cocktail to minimize DUB activities that would disassemble Ub conjugates post homogenization, and then enriched for Ub-bearing species by TUBEs chromatography (Hjerpe et al., 2009; Raasi et al., 2004), using as the ligand four copies of the Ub-Associated Domain (UBA) sequence from human hHR23A interconnected through a short flexible linker, and fused with GLUTATHIONE S-TRANSFERASE (GST) (Figure 1A). Previous studies show that this GST-4X(UBA) fusion binds effectively to native Ub-conjugates bearing various chain topologies from *Arabidopsis*, thus allowing us to interrogate its near complete ubiquitylome (Aguilar-Hernandez et al., 2017; Kim et al., 2013). In the second step, we enriched for poly-His-containing proteins via Ni-NTA chromatography under strong denaturing conditions using buffers containing 7 M guanidine-HCl and 8 M urea for the application and washing steps, respectively, followed by elution with a buffer containing 8 M urea and 400 mM imidazole (Figure 1B).

As seen in Figure 1C and D, eluants from the TUBEs and Ni-NTA columns prepared with control and flg22-treated *hexa-6His-UBQ* seedlings had substantially more ubiquitylated species than those obtained from wild-type seedlings, as detected by immunoblot analysis with rabbit anti-Ub antibodies. They appeared particularly enriched in high molecular mass Ub conjugates, with limited amounts of free Ub or the Ub dimer that were easily seen in total seedling extracts (Figure 1D). The main contaminant was RIBULOSE-1,5-BISPHOSPHATE CARBOXYLASE/OXYGENASE (Rubisco), which is often recognized by antibodies generated in rabbits.

Identification of Ub-conjugates by LC-MS/MS

Ub-conjugates from the *hexa-6His-UBQ* plants were trypsinized, separated by reversed-phase nanoflow liquid chromatography (LC), and subjected to tandem MS (MS/MS) using a Velos LTQ mass spectrometer in the high energy collision mode (Figure 1B). Peptides were matched to the *Arabidopsis* ecotype Col-0 protein database (IPI database, version 3.85; <http://www.arabidopsis.org>) using SEQUEST, based on the identification of two or more different matching peptides with a $\leq 1\%$ false discovery rate (FDR), or if a single matching peptide of equivalent stringency was identified that harbored a canonical Ub footprint (Kim et al., 2013). This footprint was detected by the presence of a di-Gly remnant (Gly-Gly) from Ub isopeptide linked to a lysine within the peptide, which increased the monoisotopic mass of the residue by 114 Da as well as protected the site from trypsin cleavage (Peng et al., 2003; Saracco et al., 2009). To help eliminate contaminants that bind non-specifically to the TUBEs and/or Ni-NTA columns, a background LC-MS/MS database was also generated with

wild-type plants that underwent the same two-step affinity protocol; these proteins were subtracted from *hexa-6HIS-UBQ* datasets to obtain the final ubiquitylome catalog (Kim et al., 2013). In total, we generated MS/MS datasets from three biological replicates for control, untreated, and flg22-treated seedlings for final comparisons.

The summation of this MS analysis enabled the detection of 391 Ub conjugates in control samples (204, 210, 250 proteins for biological replicates 1, 2, 3, respectively) and 570 Ub conjugates in the flg22-treated samples (391, 400, 142 proteins for biological replicates 1, 2, 3, respectively) (Supplemental Table 1). Notably, 90 proteins in the control samples and 88 proteins in the flg22-treated samples were consistently identified in all three biological replicates, which likely represented a high abundance of ubiquitylated species. When combined, we uncovered 961 ubiquitylated proteins with high confidence; 690 were confined to either control or flg22-treated samples, whereas 271 proteins were present in both (Figure 2A). We detected 299 proteins unique to the flg22-treated samples versus 120 proteins unique to the control samples, implying that the PTI elicitation rapidly and markedly increases ubiquitylation in *Arabidopsis* (Figure 2A).

Gene ontology analyses of flg22-treated and -untreated ubiquitylomes

To gain a global perspective for how ubiquitylation is influenced by immune elicitation, we classified the identified targets in the control and flg22-treated ubiquitylomes based on their known or predicted functions within the Gene Ontology (GO) database using the DAVID Functional Annotation Tool (Huang et al., 2009). As shown in Figure 2B, the total number of significantly enriched GO terms (FDR <0.05) was larger for the flg22-treated ubiquitylome than for that for the control for the three main GO domains ('Biological Processes': 104 vs. 82, 'Molecular Functions': 59 vs. 45, and 'Cellular Components' 74 vs. 62), and were noticeably more dedicated to pathogen-related events. Enrichments included the GO terms 'defense response to fungus', and 'innate immune response', which were found only in the flg22-treated ubiquitylome, while the related terms 'defense response to bacterium', 'regulation of stomatal movement', 'defense response by callose deposition', and 'oxidation-reduction process' were more significantly enriched (Figure 2B, C).

As compared to the total proteome, we also saw significant enrichment for GO terms associated with 'Ub-dependent protein catabolic process' (Biological process), and 'Cul4-RING E3 Ub ligase' (Cellular components) in both control and flg22-treated ubiquitylomes, as were the GO terms related to stress, such as 'response to salt stress' and 'response to cold stress' (Figure 2B). Strikingly, this enrichment in Ub-related events was more robust in the flg22-treated versus control ubiquitylomes (3.7-fold vs. 2.4-fold for 'ubiquitin-dependent protein catabolic process' and 5.5-fold vs. 3.2-fold for 'Cul4-RING E3 Ub ligase',

respectively). A prevalence for UPS components in the flg22-treated samples was also evident for proteins associated with 'proteasomes' (11.4-fold vs. 9.2-fold, respectively; Figure 2C). Taken together, this non-targeted MS analysis directly connected rapid flg22-induced ubiquitylation to Arabidopsis proteins associated with immunity and the UPS itself.

Global ubiquitylation of immunity-related proteins

Among the identified Ub conjugates were many key immunity-related proteins (Table 1). To link these ubiquitylated species to immune signaling globally, we generated an interaction network using the Search Tool for Retrieval of Interacting Genes/Proteins (STRING) database (Szklarczyk et al., 2015), which revealed an interconnected web with defense-related proteins often situated at key hubs (Figure 3A, B). Examples included plasma membrane-resident leucine-rich repeat-RLKs (LRR-RLKs), such as RECEPTOR-LIKE KINASE 7 (RLK7) and SUPPRESSOR OF *BIR1-1*/EVERSHED (SOBIR1/EVR) that interact with multiple LRR-RLP receptors (Figure 3A, B). Whereas SOBIR1 serves as a common component in RLP-mediated plant immunity (Albert et al., 2015; Liebrand et al., 2014), RLK7 is a receptor for PAMP-INDUCED PEPTIDE 1 (PIP1), which acts as a damage-associated molecular pattern (DAMP) to amplify plant immune signaling (Hou et al., 2014). Additionally, LysM RLK1-INTERACTING KINASE 1 (LIK1), which was specific to the flg22-treated ubiquitylome, has been shown to interact with the fungal chitin receptor complex component CERK1 to positively regulate resistance against necrotrophic pathogens (Le et al., 2014).

Downstream of the immune receptor complexes are sequential phosphorylation events involving MAPK cascades driven by MAPK KINASE KINASES (MAPKK/MEKK), MAPK KINASES (MAPKK/MKK), and MAPKS (MPK) components (Meng and Zhang, 2013). Notably, MKK5, MPK3, and MPK11, which become phosphorylated upon flg22 perception, were detected here, with MKK5 and MPK3 only identified in the flg22-treated ubiquitylome (Figure 3B). Heterotrimeric G-proteins, consisting of G α , G β , and G γ subunits, act as molecular switches to mediate diverse signal transduction events (Urano et al., 2013). In particular, *ARABIDOPSIS* G PROTEIN β -SUBUNIT1 (AGB1), which associates with FLS2 and regulates BIK1 stability during plant PTI (Liang et al., 2016; Liu et al., 2013), was identified in the flg22-treated samples. Ubiquitylated forms of RECEPTOR FOR ACTIVATED C KINASE (RACK)-1B and RACK1C, which scaffolds and connects heterotrimeric G proteins with MAPK cascades to form a unique signaling pathway in plant immunity (Cheng et al., 2015), were identified in both control and flg22-treated ubiquitylomes, or only the flg22-treated ubiquitylome, respectively.

Protein trafficking, regulated by a cascade of factors such as small GTPases from Rab subfamilies, also plays important roles in plant immunity (Gu et al., 2017). RabE1D,

identified here in the flg22-treated ubiquitylome, is associated with Golgi and the plasma membrane and might regulate the apoplast secretion of anti-microbial proteins, such as PATHOGENESIS-RELATED PROTEIN 1 (PR1) that induce resistance against *Pseudomonas syringae* pv. *tomato* (Pst) DC3000 (Speth et al., 2009b). Likewise, the E3 KEEP ON GOING (KEG), which localizes to the trans-Golgi network/early endosome (TGN/EE) compartments and is essential for TGN/EE-mediated secretion of PR1 and papain-like Cys protease C14 (Gu and Innes, 2011) appears to be ubiquitylated by a flg22-induced event (Figure 3B).

ETI components were also identified as flg22-specific ubiquitylation substrates. Included were four uncharacterized NLR proteins (AT1G53350, AT1G63350, AT5G45240, and AT5G48770) along with ENHANCED DISEASE RESISTANCE 1 (EDS1), which is a nucleocytoplasmic lipase-like protein indispensable for multiple TOLL-INTERLEUKIN 1 RECEPTOR (TIR) domain NLR-mediated resistance events (Cui et al., 2015). Proteins implicated in long-distance resistance were identified in the flg22-treated ubiquitylome, such as the HEAT-repeat protein ILITHYIA (ILA) that encourages systemic acquired resistance (Monaghan et al., 2010). The peroxisome-localized glycol hydrolase PENETRATION2 (PEN2) was seen in all control and flg22-treated samples and thus considered to be a high abundance Ub target. PEN2, together with PEN1 and PEN3, actuate a cell wall-based defense against non-adapted pathogens by limiting pathogen entry (Figure 3B) (Clay et al., 2009; Collins et al., 2003; Fuchs et al., 2016). Similarly, the lignin biosynthetic enzymes, CINNAMYL ALCOHOL DEHYDROGENASE 4 (CAD4), identified in both control and flg22-treated ubiquitylomes, and CAD5 identified only in flg22-treated ubiquitylomes, likely protect against pathogen invasion by modifying the lignin content of cell walls (Tronchet et al., 2010). The callose synthase POWDERY MILDEW RESISTANT 4 (PMR4), which is required for wound- and/or pathogen-induced callose formation (Meyer et al., 2009; Wawrzynska et al., 2010), was also found in both datasets (Figure 3B). Taken together, our ubiquitylome profiles implicate Ub addition in multiple defense responses ranging from the preformed cell wall-based protection, to inducible PTI and ETI, and systemic defense (Figure 3B).

Distinct ubiquitylation patterns and dynamics of immunity-related proteins upon flg22 elicitation

To examine more closely the ubiquitylation dynamics (+/-flg22) of several immunity-related substrates, we exploited a previously established *in vivo* ubiquitylation system in which FLAG-tagged Ub (FLAG-Ub) is co-expressed with HA- or myc-tagged targets in wild-type Arabidopsis leaf protoplasts (Zhou et al., 2014). Ubiquitylated proteins are enriched by

immunoprecipitation with anti-FLAG antibody beads and then probed by immunoblot analysis with anti-HA or anti-myc antibodies to assay for possible ubiquitylated species.

As shown in Figure 4A, we successfully confirmed the ubiquitylation of MPK3, MKK5, SOBIR1, and EDS1 using this *in vivo* system combined with HA or myc-tagged versions. For MPK3-HA, multiple ubiquitylated species were evident in anti-FLAG immunoprecipitates with a dominant form seen at ~52 kDa. Being ~8 kDa bigger than that of unmodified MPK3-HA (~43.8 kDa), we consider it likely that it represents a monoubiquitylated form. Notably, ubiquitylation of MPK3-HA was transiently enhanced soon after flg22 treatment (10 min) but quickly returned to a basal level of modification at 30 min (Figure 4A), even though the levels of unmodified MPK3-HA remained steady for one hr of flg22 treatment (Figure 4A), suggesting that MPK3 ubiquitylation triggered by flg22 helps relay the flg22 signal. By contrast, modification of MKK5-myc, easily seen by a strong monoubiquitylated species, appeared to be constitutive, *i.e.*, its level was unchanged by flg22 elicitation (Figure 4B). For both SOBIR1-HA and EDS1-HA, poly-ubiquitylated species accumulated in the *in vivo* system. While, the smear of high molecular mass conjugates for SOBIR1 was strongly enhanced upon flg22 treatment, they were diminished for EDS1 (Figure 4C and 4D). Taken together, we confirmed the ubiquitylation of candidate proteins using this cell-based assay, which revealed surprisingly distinct ubiquitylation patterns and dynamics for the candidates upon immune elicitation with flg22.

Ubiquitylation of RKL1 and its role in plant immunity

The LRR III protein RECEPTOR-LIKE KINASE 1 (RKL1), identified here as a flg22-triggered ubiquitylation substrate, was previously shown to modulate the activity of the aquaporin PLASMA MEMBRANE INTRINSIC PROTEIN (PIP) family during osmotic and oxidative stress together with its close homolog RLK902 (Bellati et al., 2016). Intriguingly, while RKL1 was not yet connected to pathogen defense, RLK902 had been shown to transmit an immune signal by phosphorylating the RLCK BRASSINOSTEROID-SIGNALING KINASE 1 (BSK1) (Zhao et al., 2019). The RKL1 protein harbors a five-LEUCINE RICH REPEAT (LRR) extracellular domain, a transmembrane domain, and a cytosolic kinase domain (Figure 5A), and is mainly expressed in the stomata, hydathodes, and trichomes of young rosette leaves, and floral organ abscission zones based on transcriptomic analyses (Tarutani et al., 2004). When expressed in our leaf protoplast system with FLAG-Ub, we found that RKL1-HA was constitutively poly-ubiquitylated *in vivo*, despite our proteomic analyses of seedlings which found these species only in flg22-treated samples (Figure 5B).

To further characterize the role(s) of RKL1 in PTI, we assayed both MAMP-induced ROS production and bacteria disease spread, using a T-DNA insertion mutant compromising

RKL1 function available from the SALK insertion collection (Alonso et al., 2003). The *rkl1-1* mutant (*SALK_099094*) bears an insertion within the first exon (Tarutani et al., 2004), which was confirmed here by genomic PCR analysis using gene-specific and T-DNA border primers (Figure S1A). When compared to wild-type *Arabidopsis* Col-0 leaves, we found that homozygous *rkl1-1* leaves displayed enhanced ROS production upon flg22 treatment based on a chemiluminescence assay quantifying luminol oxidation. We also conducted an assay for disease spread by infiltrating *Arabidopsis* leaves with the bacterial pathogen *Pst* DC3000 and then measuring bacterial numbers 2 days afterwards. As shown in Figure 5D, the *rkl1-1* mutant displayed significantly less infection as compared to wild type, consistent with a robust defense (Figure 5D). Together, the data implicated for the first time that RKL1 is a negative regulator of plant immunity, which is possibly influenced by ubiquitylation.

Ubiquitylation of UPS components

As revealed above (Table 2; Figures 2 and 3), multiple UPS components were preferentially identified in flg22-treated ubiquitylome. Included were the E1 UBA1, the E2 isoforms UBC13, and UBC35 (UBC13A), the E3 components CULLIN 1 (CUL1), CUL4, and UBR1, components of the 26S proteasome, and the DUBs Ub-Specific Proteases (UBP)-13 and UBP24 and a predicted Ub-Carboxy-Terminal Hydrolase (UCH) (At4g24320) expected to release free Ub monomers from poly-Ub chains and poly-Ub translation products. UPL5 is part of the single polypeptide HECT E3 family. By contrast, CUL1, together with S-PHASE KINASE-ASSOCIATED PROTEIN 1 (SKP1), RING BOX 1 (RBX1), and one of possibly >700 F-box protein variants, assemble into SCF-type E3 complexes (Vierstra, 2009), while the CUL4 scaffolds DNA DAMAGE-BINDING (DDB)-type E3 complexes assembled with RBX1, and one of over 85 DWD box-containing DDB proteins (Vierstra, 2009). Previously, DDB1 was implicated in *PR* gene expression and resistance to *Agrobacterium* infection in tomato (Liu et al., 2012b), which might explain the connection between CUL4 and immune elicitation, while UPL5 is required for SA- and NONEXPRESSOR OF *PR* GENES 1 (NPR1)-mediated plant immunity (Furniss et al., 2018c). Our discovery of the DUBs UBP13, UBP24, and a UCH in the flg22-treated ubiquitylome implies that the release of Ub from targets is also a regulatory step in immune elicitation.

A common fate for ubiquitylated proteins is turnover by the 26S proteasome. This multisubunit particle consists of a 20S cylinder-shaped core protease (CP) that houses the peptidase activities, which is capped at one or both ends by the 19S regulatory particle (RP) that assist in substrate recognition, poly-Ub chain removal, unfolding, and finally translocation of the unfolded protein into to the CP for digestion (Marshall and Vierstra, 2018; Zhou and Zeng, 2017). The CP is assembled from four stacked heptameric rings of

related α (PAA-PAG) and β (PBA-PBG) subunits arranged in an $\alpha\beta\beta\alpha$ configuration. By contrast, RP is created by a ring of six RP ATPases (RPT1-RPT6) together with a collection of non-ATPase subunits (RPN1-3 and RPN5-14). The RPT ring provides the unfoldase activities that prepare substrates for CP import, RPN10, RPN13, and possible RPN14/SEM1 serves as Ub receptors, and RPN11 is one of the several proteasome-bound DUBs that help release the Ub moieties prior to proteolysis. Unexpectedly, we identified PAC1 (α 3), PBB1 (β 2), and PBE2 (β 5) within the CP, RPT3, and RPT6b within the RP ATPase ring, and RPN2b, RPN3a, RPN6, and RPN8a/b in the RP cap, as part of the flg22-induced ubiquitylome (Figure 2B; Tables 1 and 2). Additional RPN subunits were identified (RPT1a, RPT4b, RPT5b, RPN1a, RPN7, and RPN11) in the ubiquitylome from untreated samples as well. While this extensive modification of proteasomes is consistent with its ubiquitylation helping remove damaged particles via autophagy more generally (Marshall et al., 2015), and possibly as a target of during pathogen attack more specifically (Ustin et al., 2017), the preferential and rapid modification of some subunits after flg22 treatment also suggested a direct host role for this modification during the immune defense. As the selective modifications of several proteasome subunits occurred within 30 min of flg22 treatment and long before any impact on 26S proteasome synthesis, we propose that they reflect early defense signaling events.

Flg22 induces UBC13 ubiquitylation and its role in plant immunity

We further connected the ubiquitylation target UBC13 (AT3G46460) to flg22 perception by studying its ubiquitylation profile and impact on pathogen defense. (It should be mentioned that UBC13 is unrelated to the UBC35 and UBC36 subfamily in *Arabidopsis*, which is unfortunately also named UBC13A (At1g78870) and UBC13B (At1g16890), respectively (Wen et al., 2008).) As shown in Figure 6A, we confirmed that UBC13 is ubiquitylated using a UBC13-HA variant co-expressed in *Arabidopsis* leaf protoplasts with FLAG-Ub. Only a monoubiquitylated form was evident in the immunoprecipitates from both untreated and flg22-treated samples; intriguingly, its abundance was slightly elevated upon flg22 exposure. Whether this addition was directed by autoubiquitylation or by a second factor (E3?) is not currently known.

To examine whether UBC13 influences immune signaling, we examined two T-DNA insertion mutants *ubc13-1* (CS389049) and *ubc13-2* (CS389056) available from the GABI-Kat collection (Kleinboelting et al., 2012). Both mutants were predicted to harbor a T-DNA sequence within the 6th exon of *UBC13*, which was confirmed by genomic PCR with gene-specific and T-DNA border primers (Figure S1B, C). When compared to wild-type *Arabidopsis* Col-0 leaves, homozygous *ubc13-2* seedlings displayed enhanced ROS

production upon flg22 treatment, based on the chemiluminescence assay, and were significantly more resistant to infection spread by the pathogen *Pst* DC3000 (Figure 5D). Confirmation that the *ubc13* mutations were responsible was provided using complementation studies that attempted to rescue the *ubc13-2* phenotypes with a transgene expressing UBC13-HA. Both ROS production and infectivity by the *Pst* DC3000 pathogen was returned to wild-type levels in homozygous *UBC13-HA ubc13-2* plants (Figure 6B-D). Together, our results imply that UBC13 negatively regulates plant immunity possibly by increasing the ubiquitylation state of UBC13.

Flg22 induces polyubiquitylation of RPN8b to potentially impact plant immunity

Our discovery that RPN8 is rapidly ubiquitylated upon treating *Arabidopsis* with flg22 was particularly intriguing because of its role in RP assembly where it associates with the DUB RPN11 to form an RPN11-RPN8 hetero-dimeric subcomplex (Murata et al., 2009; Smalle and Vierstra, 2004). Two paralogs of RPN8 (RPN8a, and RPN8b) assemble into the 26S particle in *Arabidopsis*. While the RPN8a isoform was more highly expressed based on mRNA levels and more commonly detected within the particle by MS (Book et al., 2010), only RPN8b was identified here in flg22-treated ubiquitylome, suggesting that its modification has a selective effect on 26S proteasome structure/function. As shown in Figure 7A, we confirmed the ubiquitylation of RPN8b by co-expressing an HA-tagged variant with FLAG-Ub in our *Arabidopsis* protoplast ubiquitylation system. Polyubiquitylated species of RPN8b-HA were evident in the immunoprecipitates from both untreated and flg22-treated samples, but their abundance was markedly enhanced upon flg22 treatment (Figure 7A).

To test whether RPN8b contributes to immune signaling, we analyzed two T-DNA insertion mutants available from the SALK insertion collection (*rpn8b-1*, SALK_128568 and *rpn8b-2*, SALK_023568). The *rpn8b-1* allele was previously shown to harbor a T-DNA sequence within the 1st intron (Palm et al., 2019), whereas we found that the *rpn8b-2* allele harbored a T-DNA sequence within the 2nd intron, both of which were confirmed here by genomic PCR with gene-specific and T-DNA border primers (Figure S1D). When compared to wild-type *Arabidopsis* Col-0 leaves, homozygous *rpn8b-1* mutant leaves showed an increase in ROS production upon flg22 perception by the chemiluminescence assay, and enhanced disease resistance upon infiltrating leaves with *Pst* DC3000 (Figure 7B-D). Complementation studies then connected RPN8b to the phenotypes described here; transgenic plants expressing RPN8b-HA under the control of the CaMV 35S promoter in the *rpn8b-1* background had their levels of flg22-induced ROS production and disease resistance restored to those in wild type (Figure 7B-D).

Collectively, the data implicated RPN8b as a negative regulator of plant immunity possibly through a process that enhances its ubiquitylation state. Whether the RPN8a isoform is similarly ubiquitylated and involved in disease resistance is not yet known. However, we note that suppression of ROS production in *rpn8b-1* seedlings by introducing *RPN8b-HA* was even more striking for one of the rescued lines, suggesting that overexpression of RPN8b might provide a dominant effect possibly by swamping out the function(s) of RPN8a (Figure 7B-D).

Mapping ubiquitylation sites on Ub conjugates

One strategy to demonstrate the impact of ubiquitylation on protein function/stability is to identify the modified lysine within the target and then block this modification through lysine to arginine substitutions. Consequently, a catalog of such sites is valuable, which can be detected by MS identification of Ub footprints, *i.e.*, trypsin-protected lysines bearing an isopeptide linked di-glycine moiety (+114 kDa) (see Figure 8A). Directed queries of all of our MS datasets identified 150 ubiquitylation sites from 120 *Arabidopsis* proteins (Table 3), which adds to the expanding list generated by other reports (Aguilar-Hernandez et al., 2017; Kim et al., 2013; Maor et al., 2007; Saracco et al., 2009). Notable footprints within disease-related proteins identified here, some of which were previously unmapped, included K668 from CUL4, K301 from RPT6b, K502 and K507 from ARGONAUTE 7 (AGO7), K422 and K434 within the NLR proteins At1g63350 and At5g48770 respectively, K286 from UCH family protein At4g24320, and K736 from PHOSPHATIDYLINOSITOL-4-PHOSPHATE 5-KINASE (PIP5K2) (Table 3; Supplemental Table 2). Our datasets broadened the number of known Ub-attachment sites, especially within *Arabidopsis* proteins involved in flg22-triggered immune responses. Interestingly, noncanonical ubiquitylation sites detected in yeast or mammals, such as those involving serine, threonine, or cysteine residues (Iwai and Tokunaga, 2009; Okumoto et al., 2011; Shimizu et al., 2010) were not identified in here. Their absence is consistent with the previous ubiquitylome studies (Kim et al., 2013) and implies that these linkages are rarely, if at all, used by plants.

Analysis of poly-Ub linkages

Owing to the widespread observations that poly-Ub chains of various linkages confer unique functions to Ub (Komander and Rape, 2012; Vierstra, 2009), we examined how flg22 treatment might affect the abundances of these polymers, using the diagnostic Ub footprint peptides for quantification based on peptide spectral matches (Peng et al., 2003). While footprints on each of the seven internal lysines (K⁶, K¹¹, K²⁷, K²⁹, K³³, K⁴⁸, and K⁶³) were identified (Figure 8B), none were seen using the N-terminal methionine, which would have

signified linear Ub concatemers (Iwai and Tokunaga, 2009). K48-linked Ub-Ub connections were the most abundant and comprised ~30% of the total detected linkages in both control and flg22-treated samples (Figure 8C). The second most abundant linkage involved K63 and was followed in abundance by those involving K11, K29, K6, K33, and K27 (Figure 8B). Interestingly, flg22 exposure altered the relative abundance of poly-Ub linkages; the percentages of K6, K11, K33, and K48-linkages increased, while the percentages of K27, K29, and K63-linkages decreased (Figure 8B). The biological relevance of such changes awaits clarification.

DISCUSSION

Protein ubiquitylation is involved in nearly all aspects of eukaryotic biology, leading to a multitude of distinct signals within a variety of cellular and physiological contexts. In line with prior studies with yeast and mammals (Bennett et al., 2007; Ordureau et al., 2020), we show here and elsewhere (Aguilar-Hernandez et al., 2017) that dynamic and multifaceted changes in the plant ubiquitylome also occur. By employing a genetically encoded *hexa-6HIS-UBQ* variant coupled with an improved two-step enrichment protocol for ubiquitylated proteins, followed by deep LC-MS/MS analysis, we found profound and complex protein ubiquitylation patterns upon immune elicitation in *Arabidopsis*.

We propose that the use of two affinity-purification steps, with the second under strongly denaturing conditions, are essential to generate reliable catalogs, which resulted here in the detection of 961 possible substrates along with 150 ubiquitylation sites mapped onto 120 individual proteins. Among the 10 candidate proteins that we subsequently tested by *in vivo* conjugation assays, all were confirmed as ubiquitylated targets *in planta*. While it is possible that the ubiquitylated species seen in protoplasts were artifacts derived from ectopic expression, the fact that different types of linkages were seen (mono- versus polyubiquitylation) and that both increased and decreased responses were seen soon after flg22 elicitation argue otherwise.

We also identified Ub-Ub linkages involving all seven lysine residues based on the discovery of diagnostic footprint peptides, with those involving K48, K63, and K11 being the most abundant. Whereas K48 polyubiquitylation has been linked to 26S proteasome-mediated turnover, and K63 ubiquitylation has been connected to the endocytic internalization of plasma membrane proteins (Romero-Barrios and Vert, 2018), the functions of the other poly-Ub linkages remain largely unknown in *Arabidopsis* even though the total abundance of those linkages exceeds 40% of the total. In addition, we saw an increased number of ubiquitylated proteins (391 of control vs. 570 of flg22-treated) upon immune elicitation. Although the datasets were not derived from quantitative proteomic approaches,

a nearly 50% increase in the number of ubiquitylated proteins seen soon after immune elicitation, suggests a strong activation of the machinery that supports Ub transfer.

Notably, our flg22-induced datasets were obtained using seedlings treated for only 30 minutes, thus implicating ubiquitylation as an early step in immune signaling. Surprisingly, several resistance proteins known to be ubiquitylated such as FLS2, BRI1, and BIK1 were not identified here (Lu et al., 2011; Ma et al., 2020; Zhou et al., 2018), which might be explained by the nature of the tissues examined or the extraction method used, which possibly preferred cytoplasmic versus plasma membrane/organelle-associated proteins. Furthermore, we should also emphasize the TUBE affinity approach likely binds polyubiquitylated proteins better than monoubiquitylated proteins and thus reduced the latter in our catalogs. One strategy to avoid this complication is to enrich for Ub footprints directly from trypsin-digested cell lysates using an anti- K- ϵ -GG antibodies specific for the GG-remnant on ubiquitylated lysines (Udeshi et al., 2013). In fact, recent studies with such antibodies in rice discovered 1,376 possible ubiquitylated proteins and revealed strong changes in ubiquitylation after a three-hr exposure to flg22 or fungal chitin (Chen et al., 2018). The only complications were that designation of a protein as an Ub target was often based on a single Ub footprint peptide and the reliability of the anti-K- ϵ -GG antibodies. Clearly, further advancements in enrichment strategies and MS instrumentation should further expand the specificity, stringency, and depth of ubiquitylome during diverse physiological conditions.

Within the flg22-induced ubiquitylomes identified here, we confirmed several Ub targets by *in vivo* ubiquitylation assays in protoplasts. Surprisingly, we found distinct ubiquitylation patterns and dynamic temporal responses to flg22 treatment. For example, MKK5 and MPK3 were strongly monoubiquitylated while SOBIR1 and EDS1 were mainly polyubiquitylated. Ubiquitylation of RPN8b and MPK3 was increased by flg22, with MPK3 modification showing a transient effect. In contrast, ubiquitylation of EDS1 in the protoplasts was reduced in response to flg22 treatment. One of the more interesting substrates identified here was receptor protein kinase RKL1, which was polyubiquitylated, and found genetically to negatively regulate flg22-triggered ROS and immunity against bacterial *Pst* DC3000 infection. Intriguingly, RLK902, a close homolog of RKL1, was previously shown to positively regulate plant disease resistance (Zhao et al., 2019), suggesting that RKL1 and RLK902 counterbalance each other in immune signaling.

Ubiquitylation often intertwines with phosphorylation in regulating protein function (Hu and Sun, 2016; Reyes et al., 2011; Skubacz et al., 2016). A well-known example in plant disease resistance is BIK1, which is phosphorylated and subsequently monoubiquitylated upon flg22 perception (Ma et al., 2020). Here, we found that the phosphorylation (Asai et al.,

2002) target MPK3 is also ubiquitinated dynamically, and like phosphorylation, this modification was enhanced ~10 min after flg22 treatment, and then was reduced to basal levels after 30 min. The cognate E3 ligase for MPK3 is currently unknown; given that MPK3 phosphorylates the E3 ligase PUB22 and regulates its turnover (Furlan et al., 2017), it is possible that PUB22 directs MPK3 ubiquitylation thus creating an internal regulatory loop. As MPK3 remained unchanged despite the dynamics of its ubiquitylation after flg22 treatment, its ubiquitylation might not commit MPK3 to turnover but confer a reversible, non-proteolytic function. In line with some ubiquitylation events being reversible, we note the detection of several DUBs in our flg22-treated ubiquitylome datasets.

Pathway analysis of our ubiquitylome datasets upon MAMP perception revealed robust changes in the ubiquitylation status for numerous components involved in the UPS, including E1, E2, E3, and DUB family members. The *in vivo* ubiquitylation assays further confirmed that the E2 UBC13 in monoubiquitinated *in vivo* with the abundance of this modification increased upon flg22 treatment. While *ubc13* mutants morphologically resembled WT plants, they displayed an enhanced flg22-induced ROS burst and increased resistance against bacterial infection. Several E3s have also been implicated in immunity, including CPR1 (Cheng et al., 2011; Gou et al., 2012; Gou et al., 2009), PUB12/13 (Liao et al., 2017; Lu et al., 2011), PUB25/26 (Wang et al., 2018), PUB22/23/24 (Stegmann et al., 2012), and RHA3A/B (Ma et al., 2020). Although it is generally accepted that E3s, not E2s, are the key determinants in substrate selection, it is possible that UBC13 preferentially works with those E3s involved in immunity.

Proteasome components, including both the core and regulatory particles, are enriched in our ubiquitylome analysis, which is consistent with the observation in yeast (Peng et al., 2003), and the known role of ubiquitylation in directing proteasome turnover through autophagy (Marshall et al., 2015). Strikingly, several of the ubiquitylated subunits were only found in the flg22-treated ubiquitylome, implying an active regulation of the proteasome machinery upon flg22 perception. Previous studies showed that several regulatory particle subunits, such as RPN1a, RPN8a, and RPT2a, but not others, play a positive role in immunity (Yao et al., 2012). Unexpectedly, we found RPN8b negatively regulates the flg22-induced ROS burst and disease resistance against *Pst* DC3000 infection. As *Arabidopsis* also encodes a second isoform of RPN8 (RPN8a) that appears more dominant in providing this subunit (Book et al., 2010), one intriguing possibility is that the RPN8b isoform becomes assembled to a unique subset of proteasomes with functions more directed toward immune perception. The immunoproteasome assembled with unique isoforms of the CP subunits β 1, β 2, and β 5 could serve as an example of such non-redundancy (Finley, 2009). Clearly, further comparisons of mutants impacting both RPN8a

and RPN8b with respect to pathogen resistance, along with assays testing the ubiquitylation status of each is now necessary to explore this notion.

Acknowledgements

We thank the *Arabidopsis* Biological Resource Center (ABRC) for T-DNA insertion mutant seeds, and members of the laboratories of L.S., P.H. and R.D.V. for discussions and comments of the experiments. The work was supported by grants from the National Institutes of Health (NIH) (R01-GM092893) and National Science Foundation (NSF) (MCB-1906060) to P.H., NIH (R01-GM097247) and the Robert A. Welch Foundation (A-1795) grant to L.S., and a grant from the NIH (R01-GM124452) to R.D.V. C. Z. and Y.H. were partially supported by the China Scholarship Council (CSC).

Author Contributions

R.D.V., P.H., and L.S. conceived the project, designed experiments, and analyzed data. X.M., C.Z., D.K., and Y.H. performed experiments and analyzed data. X.M., C.Z., R.D.V., and L.S. wrote the manuscript with inputs from all co-authors.

Declaration of Interests

The authors declare no competing interests.

MATERIALS AND METHODS

Plant materials and growth conditions

Arabidopsis thaliana 6HIS-UBQ transgenic plants in the Col-0 background were described previously (Saracco et al., 2009). The *rkl1-1* (SALK_099094), *ubc13-1* (CS389049), *ubc13-2* (CS389056), *rpn8b-1* (SALK_128568), *rpn8b-2* (SALK_023568) T-DNA insertion lines were obtained from ABRC. *p35S::UBC13-HA* transgenic plants in the *ubc13-2* background and *p35S::RPN8b-HA* transgenic plants in the *rpn8b-1* background were generated in this study (see below for details). All *Arabidopsis* plants were grown in soil (Metro Mix 366, Sunshine LP5 or Sunshine LC1) in a growth chamber at 20-23 °C, 50% relative humidity and 75 $\mu\text{E m}^{-2}\text{s}^{-1}$ light with a 12-hr light/12-hr dark photoperiod for four weeks before pathogen infection assay, protoplast isolation, and ROS assays.

Plasmid construction and generation of transgenic plants

MPK3-HA, *MKK5-MYC*, and *FLAG-UBQ* in a plant gene expression vector *pHBT* used for protoplast assays were described previously (Asai et al., 2002; Lu et al., 2011). SOBIR1, RKL1, UBC13, RPN8b, and EDS1 tagged with HA in the *pHBT* vector were generated as

following: the gene was PCR amplified from total Col-0 cDNA with primers containing *Bam*HI (for *SOBIR1*, *RKL1*, *UBC13*, and *RPN8b*) or *Nco*I (for *EDS1*) at the 5' end and *Stu*I at the 3' end, followed by *Bam*HI/*Nco*I and *Stu*I digestion and ligation into the *pHBT* vector with a sequence encoding a HA tag at the 3' end. DNA fragments cloned into the *pHBT* vectors were confirmed as correct by Sanger sequencing. The *UBC13-HA* and *RPN8b-HA* transgenes were further shuttled into a *pCB302* vector by insertion into *Bam*HI and *Stu*I digestion sites. The *A. tumefaciens*-mediated floral dip was used to transform the above binary vectors into *ubc13-2* or *rpn8b-1* plants. Transgenic plants were selected by glufosinate-ammonium (Basta, 50 µg/ml) resistance. Multiple transgenic lines were analyzed by immunoblotting for protein expression. Two lines with a 3:1 segregation ratio for Basta resistance in the T3 generation were selected to obtain homozygous seeds for further studies.

Affinity purification of ubiquitylated proteins

Affinity purification of ubiquitylated proteins was described (Kim et al., 2013) with modifications. Seeds of *Arabidopsis* Col-0 and *6HIS-UBQ* transgenic plants were vernalized at 4 °C for three days and surface-sterilized with 75% ethanol for 10 min and 5% bleach for 10 min, and plated in 1/2-strength Murashige and Skoog (MS) medium containing 1% sucrose and 0.5% agar (100 seedlings per plate) at 23 °C, 75 µE m⁻²s⁻¹ light with a 12-hr light/12-hr dark photoperiod for 14 days. Approximately 5000 seedlings (50 plates) were used for each biological replicate which generated ~16-21 gram fresh weight of tissue. Seedlings were gently transferred into a Petri dish plate and incubated overnight in water. After removing the water by vacuum, seedlings were treated for 30 min with 100 nM flg22 or water as a control.

Seedlings were gently blot-dried on Kimwipes, frozen to liquid nitrogen temperatures, and homogenized in 0.5 g/mL of extraction buffer (EB; 50 mM Tris-HCl, pH 7.2, and 200 mM NaCl), containing 0.25% Triton X-100, 13 protease inhibitor cocktail (Roche, 1 tablet per 10 ml of EB), 2 mM phenylmethanesulfonyl fluoride, 10 mM 2-chloroacetamide, 10 mM sodium metasilicate, and 1 mM N-ethylmaleimide (freshly added). The homogenate was filtered through two layers of Miracloth and the supernatant was mixed with TUBEs beads as the ratio of 40 g of tissue to 1 mL of beads. After incubated for 6 hr at 4°C, the beads were collected in a 2 x 1-0cm chromatography column and washed three times with EB and three times with EB plus 1, 2, 4 M NaCl, and eluted in 10 mL of 7 M guanidine-HCl, 100 mM NaH₂PO₄, and 10 mM Tris-HCl (pH 8.0, 24 °C). The eluted conjugates were incubated with 500 µl of Ni-NTA beads for 12 hr at 4°C in the same buffer with the addition of 20 mM imidazole and 10 mM 2-chloroacetamide. The beads were collected and washed once in 6

M guanidine-HCl, 0.1% SDS, 100 mM NaH₂PO₄, and 10 mM Tris-HCl (pH 8.0), and once in urea buffer (UB; 8 M urea, 100 mM NaH₂PO₄, and 10 mM Tris-HCl (pH 8.0)) also containing 0.1% Triton X-100, twice in UB plus 20 mM imidazole, and three times in UB alone, and eluted by 1 ml of UB supplemented with 400 mM imidazole. The two-step purified conjugates were filtered by the Ultracel-10K filter (Millipore) and separated by SDS-PAGE followed by staining for protein with silver, or immunoblotting with anti-Ub (Sigma-Aldrich) or anti-His (Novagen) antibodies to detect Ub conjugates.

Mass spectrometry

Purified proteins (100 µl) or control samples were reduced with 10 mM DTT at room temperature for 1 hr, and alkylated in the dark in the presence of 30 mM 2-chloroacetamide at room temperature for a further 1 hr. Excess alkylating agent was quenched with 20 µL of 200 mM DTT for 10 min. Samples, ten-fold diluted with 25 mM ammonium bicarbonate, were digested for 12 hr at 37 °C with 2 mg of sequencing-grade trypsin (Promega), followed by a second incubation for 6 hr with an additional 2 mg of trypsin. Digestion products were desalted using a C18 solid-phase extraction pipette tip (SPEC PT C18, Varian), vacuum dried, and reconstituted in 10 µL of 0.1% formic acid and 5% acetonitrile in water for MS analysis.

Samples were analyzed by a nanoflow liquid chromatography system (nanoAcquity; Waters) combined with an electrospray ionization FT/ion-trap mass spectrophotometer (LTQ Orbitrap Velos; Thermo Fisher Scientific). The LC system employed a 100 × 365 -µm fused silica microcapillary column packed with 15 cm of 3-µm-diameter, 100-Å pore size, C18 beads (Magic C18; Bruker), with the emitter tip pulled to 2 mm using a laser puller (Sutter Instruments). Peptides were loaded onto the column for 30 min at a flow rate of 500 nL/min and eluted over 120 min at a 200 nL/min flow rate with a gradient of 2% to 30% acetonitrile in 0.1% formic acid. MS spectra were acquired in the FT orbitrap between 300 and 1500 mass-to-charge ratios (m/z) at a resolution of 60 000, followed by 10 MS/MS HCD scans of the 10 highest intensity parent ions at 42% relative collision energy and 7500 resolution, with a mass range starting at 100 mass-to-charge ratios. Dynamic exclusion was enabled with a repeat value of two for 30 sec and an exclusion window for 120 sec.

MS data analysis

The MS/MS data were searched using SEQUEST version 1.2 (ThermoFisher Scientific) against the *A. thaliana* ecotype Col-0 protein database (IPI database, version 3.85, containing 39,677 entries available within the *Arabidopsis* Information Resource [TAIR], <http://www.arabidopsis.org>) database. Masses for both precursor and fragment ions were

treated as monoisotopic. Met oxidation (+15.994915 Da), Cys carbamidomethylation (+57.021464 Da), the di-Gly Ub footprint indicating Ub addition to a lysine residue (+114.043 Da) were set as variable modifications. The database search allowed for up to two missed trypsin cleavages, and the ion mass tolerances were set to 10 ppm for precursor and 0.1 Da for HCD fragments. The data were filtered using a 1% FDR, and a minimum of two peptide matches was required or at least one peptide if it included a GGK Ub footprint.

GO analysis was evaluated by DAVID Functional Annotation Tool (Huang et al., 2009) using GO annotations in the TAIR GO database (<http://www.arabidopsis.org/>). Fold enrichments were calculated based on the frequency of proteins annotated to the term compared with their frequency in the proteome. The *p*-value combined with the FDR correction was used as criteria of significant enrichment for GO catalogs, whereas a *p*-value < 0.05 were considered to be enriched for GO terms. The GO annotation was classified based on the “biological processes,” “molecular functions,” and “cellular components” categories and visualized as a bubble plot by R Project 4.0.0 (<https://www.r-project.org/>) (Bonnot et al., 2019). Protein interaction networks were created by the STRING database version 11.0 (Szklarczyk et al., 2015), and visualized by Cytoscape version 3.8.0 (Shannon et al., 2003).

Detection of ROS production

Leaves from 4-5-week-old, soil-grown *Arabidopsis* plants were punched into 5-mm diameter discs. The discs were incubated in 100 µl of water with gentle shaking overnight, which was then replaced with 100 µl of reaction solution containing 50 µM luminol and 10 µg/ml horseradish peroxidase (Sigma-Aldrich) supplemented with or without 100 nM flg22. Luminescence was measured with a luminometer (GloMax®-Multi Detection System, Promega) with a setting of 1 min as the interval for 40 min. Detected values for ROS production were indicated as means of Relative Light Units (RLU).

Pathogen infection assays

Pseudomonas syringae pv. *tomato* (Pst) DC3000 was cultured for overnight at 28 °C in King's B medium supplemented with rifamycin (50 µg/ml). Cells were collected by centrifugation at 3,500 rpm, washed, and re-suspended to the density of 5 x 10⁵ cfu/ml in 10 mM MgCl₂. Leaves from four-week-old *Arabidopsis* plants were hand-inoculated with bacterial suspension using a needleless syringe. To measure *in planta* bacterial growth, three to four samples, each containing two 6-mm diameter leaf discs, were homogenized in 100 µl of water, and plated in serial dilutions on medium containing 1% tryptone, 1%

sucrose, 0.1% glutamic acid, 1.8% agar, and 25 µg/ml rifamycin. Plates were incubated at 28 °C and bacterial colony forming units (cfu) were counted after 2 days.

***In vivo* ubiquitylation assays**

Protoplast isolation and transient expression assay were as described previously (Zhou et al., 2014). Protoplasts were freshly isolated from leaves of four-week-old Col-0 plants and transfected at 2×10^5 cell/ml (500 µL) with various combinations of plasmids harboring genes encoding FLAG-UBQ (25 µL at 2 µg DNA/µL), and the HA or MYC-tagged proteins of interested (25 µL at 2 µg DNA /µL). The protoplasts were incubated for overnight at room temperature followed by a 2-hr treatment with DMSO alone or 2 µM MG132 dissolved in DMSO, and then a 2-hr exposure to 100 nM flg22. After homogenization in 300 µL of Immunoprecipitation (IP) buffer (150 mM NaCl, 50 mM Tris-HCl (pH 7.5), 5 mM Na₂EDTA, 1% Triton X-100, 2 mM Na₃VO₄, 2 mM NaF, 1 mM DTT, and 1:100 diluted protease inhibitor cocktail (Sigma-Aldrich), the FLAG-tagged proteins were immunoprecipitated by incubation of the extracts with 5 µL of anti-FLAG antibody agarose beads (Sigma-Aldrich) for 1 hr at 4 °C with gentle shaking. The anti-FLAG antibody beads were collected and washed three times with IP buffer (without cocktail) and once with 50 mM Tris-HCl (pH 7.5), and then incubated in 30 µL of hot SDS-PAGE sample buffer for 5 min. Twenty µL of the sample before adding the beads was used as the input control. The samples were separated by SDS-PAGE followed by electrophoretic transfer onto Immun-Blotpolyvinylidene difluoride (PVDF) membranes (Bio-Rad), and immunoblotting with appropriate antibodies. Antibodies against plant Ub (van Nocker et al., 1996) and 6His (Kim et al., 2013) were as described. Anti-FLAG, anti HA, and, anti-myc antibodies were purchased from Sigma-Aldrich (A8592), Roche (12013819001), and Santa Cruz (sc-40), respectively.

Statistical analyses

Data for quantification analyses are presented as mean ± standard error of the mean (s.e.m.). The statistical analyses were performed by Student's *t*-test or one-way analysis of variance (ANOVA) test (* *p*-value < 0.05, ** *p*-value < 0.01, *** *p*-value < 0.001). The number of replicates is shown in the figure legends.

FIGURE LEGENDS

Figure 1. Two-step affinity purification of ubiquitylated proteins from transgenic *Arabidopsis* expressing 6HIS-UBQ treated with or without flg22.

A. Diagram of the *p35S::hexa-6HIS-UBQ* transgene. A synthetic UBQ gene encoding six repeats of a Ub monomer N-terminally tagged with a 6His sequence followed by a glycine (G)-rich linker (MHHHHHHGSGGSA) was fused head-to-tail to form a single in-frame *hexa-6His-UBQ* transgene expressed under the control of the constitutive CaMV 35S promoter (Saracco et al., 2009).

B. Flow chart describing the proteomic approach to stringently identify Ub conjugates from *Arabidopsis p35S::hexa-6HIS-UBQ* seedlings. Ubiquitylated proteins were first enriched by the UBA-Ub affinity using GST-TUBEs beads under the native condition followed by Ni-NTA chromatography under the denaturing condition. The eluants were digested with trypsin and subjected to LC/ESI-MS/MS analysis.

C and D. Ubiquitylated proteins isolated from *Arabidopsis* plants expressing 6His-Ub by the two-step affinity purification. Ubiquitylated proteins enriched as outlined in (B) were subjected to SDS-PAGE and by either stained for protein with silver (C) or immunoblotted with anti-Ub antibodies (D). Samples from wild-type seedlings (WT) purified with both GST-TUBEs and Ni-NTA beads were included for comparison. FT and EL represent the flow-through and eluted fractions, respectively, from the Ub affinity (1) and Ni-NTA affinity columns (2). The total represents the total protein extracts before affinity purifications. The migration positions of the Ub dimers (Ub₂), and Ub conjugates (Ub-conj) are indicated by the brackets.

Figure 2. Gene ontology (GO) analysis of flg22-treated and untreated ubiquitylomes.

A. Venn diagram showing the overlap of the total ubiquitylated proteins isolated from plants with or without flg22 exposure.

B. Gene ontology (GO) enrichment of ubiquitylated proteins in control or flg22 treated ubiquitylomes. GO analysis in biological processes, molecular functions, and cellular components was performed using the DAVID Functional Annotation Tool (Huang da et al., 2009) and GO annotations from the TAIR GO database (<http://www.arabidopsis.org/>). The fold enrichment was calculated based on the frequency of proteins annotated to the term compared with their frequency in the total proteome. The dot size indicates the number of genes associated with each process and the dot color indicates the significance of the enrichment. FDR was based on the corrected *p*-values. The vertical grey dashed line represents a fold enrichment of 1. The total number of GO terms was listed under each GO

domain (biological processes, molecular functions, and cellular components). The selected terms related to stress and the UPS are shown.

C. Venn diagram showing the overlaps of ubiquitylated proteins with or without flg22 in the selected categories. Proteins were categorized into functional groups based on the GO annotations; those related to defense responses and the UPs are shown.

Figure 3. Protein interaction network of ubiquitylated proteins identified from 6HIS-UBQ seedlings treated with or without flg22.

A. Protein interaction networks were generated with the list of 391 (control) and 570 (flg22-treated) ubiquitylated proteins using the STRING database and visualized in Cytoscape. The node colors indicate proteins identified in control (yellow), flg22-treated (red), or both conditions (blue) of datasets. Clusters of the proteasome, ubiquitylation, and defense-related components are marked by dashed lines.

B. Map of proposed subcellular locations and functions of immunity-related proteins identified from the ubiquitylomes.

Figure 4. *In vivo* confirmation that several candidates are ubiquitylated in *Arabidopsis*.

Arabidopsis wild-type protoplasts were co-transfected with FLAG-tagged Ub (*FLAG-UBQ*) and HA- or myc-tagged substrate or a control vector (Ctrl) and incubated for 10 hr followed by treatment with 100 nM flg22 for the indicated. Protein extracts were immunoprecipitated with anti-FLAG agarose beads (IP: α -FLAG) and the ubiquitylated proteins were immunoblotted with anti-HA or anti-myc antibodies (top left) or anti-FLAG antibodies (bottom left). The input controls were shown by an anti-HA/myc (top right) or anti-FLAG (bottom right) immunoblots. **A.** Monoubiquitylation of MPK3 is enhanced upon flg22 treatment. Protoplasts were isolated at 0 to 30 min after flg22 exposure.

B. Polyubiquitylation of SOBIR1 *in vivo* is enhanced upon a 30 min treatment with flg22.

C, D. MKK5, and EDS1 were ubiquitylated *in vivo* following a 30-min treatment with flg22.

The above experiments were performed three times with similar results.

Figure 5. RKL1 is ubiquitylated and involved in PTI responses.

A. Diagram of the RKL1 protein. The signal peptide (SP), leucine-rich repeat (LRR), transmembrane (TM), and protein kinase (K) domains are shown. The amino acids demarcating each domain are shown.

B. *In vivo* ubiquitylation of RKL1 upon flg22 treatment. Protoplasts were co-transfected with *FLAG-UBQ* and HA-tagged RLK1 (*RKL1-HA*) or a control vector (Ctrl) and incubated for 10

hr followed by treatment with 100 nM flg22 for 30 min. Proteins were immunoprecipitated with anti-FLAG agarose beads (IP: α -FLAG) and the ubiquitylated proteins were detected by immunoblotting with anti-HA antibodies (top). The input controls were shown by an anti-HA immunoblot (bottom).

C. flg22-induced reactive oxygen species (ROS) burst is elevated in the *rk1-1* mutants. Leaf disks of four-week-old soil-grown *Arabidopsis* WT or *rk1-1* plants (SALK 099094) were treated with 100 nM flg22 or water as a control, and ROS production was monitored over time. The data are shown as mean $\pm$ s.e.m (n = 16) of relative fluorescence units (RLU) overlaid on the dot plot.

D. *rk1* mutants have elevated resistance against *Pst* DC3000. Leaf disks of four-week-old soil-grown *Arabidopsis* WT or *rk1-1* plants were hand-infiltrated with *Pst* DC3000 at $OD_{600}=5 \times 10^{-4}$ CFU/ml, and bacterial growth was counted at 0 or 3 dpi. The data are shown as mean $\pm$ s.e.m. overlaid on dot plot (n = 3 for 0 dpi; n = 4 for 3 dpi). The experiments were performed three times with similar results.

Figure 6. UBC13 is ubiquitylated and negatively regulates PTI responses.

A. *In vivo* ubiquitylation of UBC13 upon flg22 treatment. Protoplasts were co-transfected with *FLAG-UBQ* and HA-tagged UBC13 (*UBC13-HA*) and incubated for 10 hr followed by treatment with 100 nM flg22 for 30 min. Protein extracts were immunoprecipitated with anti-FLAG beads (IP: α -FLAG) and the ubiquitylated proteins were immunoblotted with anti-HA antibodies (top). The input control was shown by an anti-HA immunoblot (bottom).

B, C. UBC13 negatively regulates flg22-triggered ROS production. Leaf disks of four-week-old WT, *ubc13* (CS389049/*ubc13-1* & CS389056/*ubc13-2*), and plants harboring the *p35S::UBC13-HA* transgene in the *ubc13-2* mutant background were treated with 100 nM flg22. ROS production was monitored over time. The data are shown as mean $\pm$ s.e.m overlaid on dot plot (n =16) of RLU. The total photon count was shown in (**C**).

D. *ubc13* mutants have elevated resistance against *Pst* DC3000. Leaf disks of four-week-old soil-grown WT, *ubc13-1*, *ubc13-2*, and *p35S::UBC13-HA ubc13-2* (Line 1 and 2) plants were hand-infiltrated with *Pst* DC3000 at $OD_{600}=5 \times 10^{-4}$ CFU/ml, and bacterial growth was counted at 3 dpi. The data are shown as mean $\pm$ s.e.m overlaid on the dot plot (n = 3). The experiments were performed three times with similar results.

Figure 7. RPN8b is ubiquitylated and negatively regulates PTI responses.

A. *in vivo* ubiquitylation of RPN8b upon flg22 treatment. Protoplasts were co-transfected with *FLAG-UBQ* and HA-tagged RPN8b (*RPN8b-HA*) or a control vector (Ctrl) and incubated for 10 hr followed by treatment with 100 nM flg22 for 30 min. Protein extracts from

protoplasts were immunoprecipitated with anti-FLAG beads (IP: α -FLAG) and the ubiquitylated proteins were immunoblotted with anti-HA antibodies (top). The input control was shown by an anti-HA immunoblot (bottom).

B, C. RPN8b negatively regulates flg22-triggered ROS production. Leaf disks of four-week-old WT, *rpn8b-1* (SALK_128568), *rpn8b-2* (SALK_023568), and *rpn8b-1* plants harboring the *p35S::RPN8b-HA* transgene were treated with 100 nM flg22 or water as a control, and ROS production was monitored over time. The data are shown as mean $\pm$ s.e.m overlaid on the dot plot ($n \geq 13$) of RLU. The total photon count was shown in (C).

D. *rpn8b* mutants have elevated resistance against *Pst* DC3000. Leaf disks from four-week-old soil-grown WT, *rpn8b-1*, *rpn8b-2*, and *35S::RPN8b rpn8b-1* plants were hand-infiltrated with *Pst* DC3000 at $OD_{600}=5 \times 10^{-4}$ CFU/ml, and bacterial growth was counted at 3 dpi. The data are shown as mean $\pm$ s.e.m overlaid on dot plot ($n = 3$). The experiments were performed three times with similar results.

Figure 8. MS/MS mapping of Ub-Ub linkages.

A. Strategy for detecting ubiquitylation sites by MS/MS. After trypsin digestion, a diglycine remnant of Ub covalently appended to a lysine residue of a conjugated protein is detected by an increased mass of 114 kDa for the lysine residue, along with protection from trypsin cleavage after that residue. Amino acids are denoted by single-letter code.

B. The distribution of Ub-Ub linkages across the seven Ub lysines based on MS analysis from *Arabidopsis* transgenic plants carrying *6HIS-UBQ* seedlings treated with or without 100 nM flg22. Percentage of Ub footprints at each site were obtained from PSM counts of diagnostic footprint peptides.

Table 1. Ubiquitylation of immunity-related proteins identified from *6HIS-UBQ* seedlings treated with or without flg22.

Locus ID	Other names	Description	Category	Role in immunity	Condition	Ref.
AT4G34230	CAD5	Cinnamyl alcohol dehydrogenase 5	Preformed defense	Involves in lignin biosynthesis and disease resistance against <i>Pst</i> DC3000.	flg22	(Tronchet et al., 2010)
AT3G10340	PAL4	Phenylalanine ammonia-lyase 4	Preformed defense	Functions in the lignin and SA pathway in plant disease resistance.	flg22	(Chen et al., 2017; Huang et al., 2010)
AT5G36890	BGLU42	Beta glucosidase 42	Preformed defense	Promotes disease resistance against <i>B.cinerea</i> , <i>H.arabidopsidis</i> and <i>Pst</i> DC3000	flg22	(Stringlis et al., 2018; Zamioudis et al., 2014)
AT3G53420	PIP2A, PIP2;1	Plasma membrane intrinsic protein 2A	Preformed defense	Involves in stomatal closure triggered by abscisic acid (ABA) or flg22 for water and H ₂ O ₂ transport activities	flg22	(Rodrigues et al., 2017)
AT1G09970	RLK7	Leucine-rich repeat receptor-like kinase 7	RLK	Receptor for PIP1 defense peptide in PTI signaling	flg22	(Hou et al., 2014; Pitorre et al., 2010)
AT3G14840	LIK1	LysM RLK1-interacting kinase 1	RLK	Interacts with CERK1 and positively regulates resistance to necrotrophic pathogens.	flg22	(Le et al., 2014)
AT1G48480	RKL1	Receptor-like protein kinase 1	RLK	Characterized in this study	flg22	(Tarutani et al., 2004)
AT2G31880	SOBIR1,	Leucine-rich repeat	RLK	Interacts with multiple	flg22	(Albert et

	EVR	receptor-like protein		LRR-RLP and functions as a common component in RLP-mediated plant immunity		al., 2015; Liebrand et al., 2014)
AT1G53350	RPP8L2	RPP8 like protein 2; Resistance protein; CC-NBS-LRR	CNL	Belongs to CNL-D subclass, may involve in controlling virus infection	flg22	(Bakker et al., 2006; Mondragon-Palomino et al., 2017; Revers et al., 2003; Tan et al., 2007)
AT5G45240		Resistance protein; TIR-NBS-LRR	TNL	uncharacterized	flg22	
AT5G48770		Resistance protein; TIR-NBS-LRR	TNL	uncharacterized	flg22	
AT5G13530	KEG	RING E3 Ub-protein ligase; KEEP ON GOING	Signaling	Essential for the secretion of apoplast antimicrobial proteins.	flg22	(Gu and Innes, 2011)
AT2G46240	BAG6	BCL-2-associated athanogene 6	Signaling	Co-chaperone regulating diverse cellular pathways, such as programmed cell death, abiotic stress responses, and plant basal resistance	flg22	(Li et al., 2016)
AT2G21660	CCR2, GRP7	Cold, circadian rhythm, and RNA binding 2; glycine-rich RNA binding protein 7	Signaling	Associates with FLS2 mRNA and proteins; maintains the FLS2 protein level	flg22	(Nicaise et al., 2013)

AT4G34460	AGB1, ELK4	Heterotrimeric G-protein beta subunit; ERECTA-LIKE 4	Signaling	Complexes with FLS2 and regulates BIK1 stability	flg22	(Liang et al., 2016; Liu et al., 2013)
AT5G03520	RAB8C, RABE1D	Rab GTPase homolog 8C	Signaling	Involves in protein trafficking, interacts with AvrPto, and accumulates in response to bacterial infection	flg22	(Hou and Gao, 2017; Speth et al., 2009a)
AT3G21220	MEK5, MKK5	Mitogen-activated protein kinase kinase 5	Signaling	Activate MPK3 and MPK6 in PTI signaling.	flg22	(Meng and Zhang, 2013)
AT3G45640	MPK3	Mitogen-activated protein kinase 3	Signaling	Activated by MAMP treatment in PTI signaling	flg22	(Asai et al., 2002; Pecher et al., 2014; Su et al., 2018)
AT2G33340	MAC3B; PUB60	MOS4-associated complex 3B; Plant U-Box 60 E3 ligase	Signaling	Localizes to the nucleus and involves in plant innate immunity.	flg22	(Monaghan et al., 2009)
AT1G77740	PIP5K2	Phosphatidylinositol-4-phosphate 5-kinase	Signaling	Responsible for PI(4,5)P2 biosynthesis; inhibits fungal pathogen development and triggers disease resistance	Control	(Qin et al., 2020)
AT3G18130	RACK1C	Receptor for activated C kinase 1C	Signaling	scaffold proteins connecting heterotrimeric G proteins with a MAPK cascade in plant immunity	Control, flg22	(Cheng et al., 2015; Nakashima et al., 2008; Su et al., 2015)
AT3G48090	EDS1	Enhanced disease susceptibility 1; a lipase-like protein	Signaling	Required for multiple TNL-mediated resistance.	flg22	(Cui et al., 2015; Heidrich et

						al., 2011; Parker et al., 1996)
AT3G50480	HR4	Homolog of RPW8	Signaling	Involved in autoimmunity mediated by RPP7 oligomerization	flg22	(Li et al., 2020)
AT1G64790	ILA	ILITYHIA; HEAT repeat protein	SAR	Involved in immunity against bacterial infection, non-host resistance, and systemic acquired resistance	flg22	(Monaghan and Li, 2010)
AT2G44490	BGLU26, PEN2	Beta glucosidase 26, PENETRATION 2	Preformed defense	Required for callose deposition and glucosinolate activation in pathogen-induced resistance	Control, flg22	(Clay et al., 2009; Pastorczyk and Bednarek, 2016)
AT3G19450	CAD4	Cinnamyl alcohol dehydrogenase 4	Preformed defense	Involved in lignin biosynthesis and disease resistance to bacterial pathogens.	Control, flg22	(Kim et al., 2004; Tronchet et al., 2010)
AT4G03550	GSL5, PMR4	Callose synthase	Preformed defense	Required for callose formation induced by pathogen infections.	Control, flg22	(Jacobs et al., 2003; Wawrzynska et al., 2010)
AT1G63350		Resistance protein; CC-NBS-LRR	CNL	uncharacterized	Control, flg22	
AT1G01560	MPK11	MAP Kinase 11	Signaling	Activated upon flg22 treatment	Control, flg22	(Bethke et al., 2012; Eschen-Lippold et al., 2012)

AT4G23650	CPK3	Calcium-dependent protein kinase 3	Signaling	required for sphingolipid-induced cell death and viral infection	Control, flg22	(Lachaud et al., 2013; Perraki et al., 2017)
AT1G48630	RACK1B	Receptor for activated C kinase 1B	Signaling	scaffold proteins connecting heterotrimeric G proteins with a MAPK cascade in plant immunity	Control, flg22	(Cheng et al., 2015; Nakashima et al., 2008; Su et al., 2015)

908

909

Table 2. Examples of UBQ pathway components identified from ubiquitylomes

Locus ID	Other names	Description	Category	Role in immunity	Condition	Ref.
AT2G47110	UBQ6	Poly-UBQ gene	Ubiquitin	Not known.	flg22	
AT2G30110	UBA1	Ub-activating enzyme 1	E1	Involved in PTI and ETI.	flg22	(Furniss et al., 2018b; Goritschnig et al., 2007)
AT1G78870	UBC13A, UBC35	Ub-conjugating enzyme 35/13A	E2	Regulates the immune response mediated by Fen and other R proteins through Lys-63–linked ubiquitylation	flg22	(Mural et al., 2013; Wang et al., 2019)
AT3G46460	UBC13	Ub-conjugating enzyme 13		Characterized in this study.	flg22	
AT4G02570	AXR6, CUL1	Cullin-1, a component of SCF Ub E3 ligase complexes		SKP1-CULLIN1-CPR1 (SCF), regulates protein stability of SNC1 and RPS2 and autoimmunity	flg22	(Cheng et al., 2011)
AT5G46210	CUL4	CULLIN4, Ub E3 ligase		Required for <i>PR</i> gene expression and resistance to Agrobacterial infection.	flg22	(Liu et al., 2012a)
AT4G12570	UPL5	Ub E3 ligase UPL5	E3	Required for SA- and NPR1-mediated plant immunity	flg22	(Furniss et al., 2018a; Miao and Zentgraf, 2010; Zhou and Zeng,

						2017)
AT1G70320	UPL2	Ub E3 ligase 2		Not known.	Control, flg22	
AT4G30890	UBP24	Ub-specific protease 24	UBP	Negatively regulates abscisic acid signaling; a potential MPK3 interactor.	flg22	(Rayapura m et al., 2018; Zhao et al., 2016)
AT3G11910	UBP13	Ub-specific protease 13		Negatively regulates plant resistance against <i>Pst</i> DC3000 infections	flg22	(Ewan et al., 2011)
AT5G06600	UBP12	Ub-specific protease 12			Control, flg22	
AT4G24320		Ub carboxyl-terminal hydrolase family protein	UCH	Not known.	flg22	
AT3G61140	CSN1, FUS6	COP9 signalosome subunit 1	COP9	Not known.	flg22	
AT5G56280	CSN6A	COP9 signalosome complex subunit 6a		Not known.	flg22	
AT3G22110	PAC1	26S proteasome subunit alpha type-4	Proteasome component	Both the 26S proteasome core and regulatory particles were implicated to be involved in plant defense against pathogen infections.	flg22	(Adams and Spoel, 2018; DIELEN et al., 2010; Lee et al., 2006; Marino et al., 2012; Üstün et al., 2016; Yao et al.,
AT3G27430	PBB1	26S proteasome subunit beta type-7A			flg22	
AT3G26340	PBE2	26S proteasome subunit beta type-5B			flg22	

AT5G58290	RPT3	26S proteasome regulatory AAA-ATPase subunit			flg22	2012)
AT5G20000	RPT6b	26S proteasome regulatory AAA-ATPase subunit			flg22	
AT2G32730	RPN2a	26S proteasome regulatory subunit			flg22	
AT1G75990	RPN3b	26S proteasome regulatory subunit			flg22	
AT1G29150	RPN6	26S proteasome regulatory subunit			flg22	
AT3G11270	RPN8b	26S proteasome regulatory subunit			flg22	
AT1G53750	RPT1a	26S proteasome regulatory AAA-ATPase subunit			Control, flg22	
AT1G45000	RPT4b	26S proteasome regulatory AAA-ATPase subunit			Control, flg22	

AT2G20580	RPN1a	26S proteasome regulatory subunit			Control, flg22	
AT1G20200	RPN3a, EMB2719	26S proteasome regulatory subunit			Control, flg22	
AT4G24820	RPN7	26S proteasome regulatory subunit			Control, flg22	
AT5G05780	RPN8a	26S proteasome regulatory subunit			Control, flg22	
AT5G23540	RPN11	26S proteasome regulatory subunit			Control, flg22	
AT1G09100	RPT5b	26S proteasome regulatory AAA-ATPase subunit			Control	
AT1G04810	RPN2b	26S proteasome regulatory subunit			Control	
AT2G02560	CAND1, ETA2	Cullin-associated and neddylation dissociated 1	Ub component	Regulators of SCF (CPR1) complexes	flg22	(Huang et al., 2016)
AT3G20620		F-box family protein-related		Not known.	Control, flg22	

AT3G23633		F-box associated ubiquitylation effector family protein		Not known.	Control, flg22	
AT1G77710	Ufm1, CCP2	Ub-like, Ufm1		Not known.	Control	

910

911

Table 3. Examples of candidates with Ub-footprints identified from ubiquitylomes.

Locus ID	Other names	Description	Ub-footprint peptide	Ub attachment sites	Condition
AT3G62150	ABCB21, PGP21	ABC transporter B family member 21	SSLSRSLSK _(Ub) R	Lys677 of 1296	flg22
AT2G19590	ACO1	ACC oxidase 1	LSK _(Ub) LMCENLGLDQEDIMNA FSGPK _(Ub) GPAFGTK	Lys135 of 310 Lys156 of 310	+ Contr ol, flg22
AT1G60880	AGL56	AGAMOUS-LIKE-56	K _(Ub) ASQLCLLSPATQIAILAAP MTSK	Lys33 of 201	Contr ol
AT3G53230	CDC48B	Cell division cycle 48B	REK _(Ub) THGEVER	Lys319 of 815	flg22
AT1G07270	CDC6	Cell division control 6	K _(Ub) DATVGELNK _(Ub) LYLEICK	Lys427 of 505 Lys438 of 505	+ flg22
AT4G34530	CIB1	Transcription factor (cryptochrome-interacting)	MKFLQDLVPGCDK _(Ub) ITGK	Lys210 of 335	Contr ol
AT1G48260	CIPK17, SNRK3.21	SNF1-related kinase 3.21	TK _(Ub) IYMVLECVTGGDLFDR	Lys83 of 432	flg22
AT3G46790	CRR2	Chlororespiratory reduction 2	GVLYELETEEK _(Ub) ER	Lys582 of 657	Contr ol, flg22
AT5G06150	CYCB1;2	Cyclin B1;2	K _(Ub) PINGDNKVPALGPKR	Lys77 of 445	flg22
AT4G38170	FRS9	FAR1-related sequence 9	ELK _(Ub) ADYEATNSTPVMKTPS PMEK	Lys275 of 545	Contr ol
AT5G14040	MTP3, PHT3;1	Phosphate transporter 3;1	KTYSDLAGPEYTA _(Ub) YK	Lys172 of 375	flg22

AT5G25610	RD22	Responsive to dehydration 22	YHVRAVSTEVAK _(Ub) K	Lys287 of Contr 392 ol
AT5G62090	SLK2	SEUSS-like 2	YVRCLQISEVVSSMK _(Ub) DMID FCR	Lys537 of 816 flg22
AT3G23633		F-box associated ubiquitylation effector family protein	SLLVCMSSSLFHEEK _(Ub) K _(Ub) LA VCCNLNR	Lys123 + Contr Lys124 of ol, 157 flg22
AT1G63350		Resistance protein; CC-NBS-LRR family	K _(Ub) ENLIEYWICEEIIDGSEGID K	Lys422 of 898 flg22
AT1G77740	PIP5K2	Phosphatidylinositol-4-phosphate 5-kinase	SLQADPASISAVDPK _(Ub) LYSR	Lys736 of Contr 754 ol
AT3G26340	PBE2	26S proteasome subunit beta type-5B	K _(Ub) AVEMVK _(Ub) PAK	Lys47 + Lys53 of flg22 273
AT4G24320		Ub carboxyl-terminal hydrolase family protein	K _(Ub) TETDNVICLG EYLGFGMR FK	Lys286 of 395 flg22
AT5G20000	RPT6b	26S proteasome regulatory AAA-ATPase subunit	TMLELLNQLDGF EASNK _(Ub) IK VLMATNR	Lys301 of 419 flg22
AT5G46210	CUL4	CULLIN4, Ub E3 ligase	TLQSLACGK _(Ub) VRVLQK	Lys668 of 792 flg22
AT5G48770		Resistance protein; TIR-NBS-LRR family	DK _(Ub) TIFIRVACLFNGEPVSR	Lys434 of 1190 flg22

References

- Adams, E.H.G., and Spoel, S.H. (2018). The ubiquitin-proteasome system as a transcriptional regulator of plant immunity. *J Exp Bot* 69, 4529-4537.
- Aguilar-Hernandez, V., Kim, D.Y., Stankey, R.J., Scalf, M., Smith, L.M., and Vierstra, R.D. (2017). Mass Spectrometric Analyses Reveal a Central Role for Ubiquitylation in Remodeling the Arabidopsis Proteome during Photomorphogenesis. *Mol Plant* 10, 846-865.
- Albert, I., Böhm, H., Albert, M., Feiler, C.E., Imkamp, J., Wallmeroth, N., Brancato, C., Raaymakers, T.M., Oome, S., and Zhang, H. (2015). An RLP23–SOBIR1–BAK1 complex mediates NLP-triggered immunity. *Nature Plants* 1, 1-9.
- Alonso, J.M., Stepanova, A.N., Leisse, T.J., Kim, C.J., Chen, H., Shinn, P., Stevenson, D.K., Zimmerman, J., Barajas, P., Cheuk, R., *et al.* (2003). Genome-wide insertional mutagenesis of Arabidopsis thaliana. *Science* 301, 653-657.
- Asai, T., Tena, G., Plotnikova, J., Willmann, M.R., Chiu, W.L., Gomez-Gomez, L., Boller, T., Ausubel, F.M., and Sheen, J. (2002). MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature* 415, 977-983.
- Bakker, E.G., Toomajian, C., Kreitman, M., and Bergelson, J. (2006). A genome-wide survey of R gene polymorphisms in Arabidopsis. *Plant Cell* 18, 1803-1818.
- Bellati, J., Champeyroux, C., Hem, S., Rofidal, V., Krouk, G., Maurel, C., and Santoni, V. (2016). Novel Aquaporin Regulatory Mechanisms Revealed by Interactomics. *Mol Cell Proteomics* 15, 3473-3487.
- Bennett, E.J., Shaler, T.A., Woodman, B., Ryu, K.Y., Zaitseva, T.S., Becker, C.H., Bates, G.P., Schulman, H., and Kopito, R.R. (2007). Global changes to the ubiquitin system in Huntington's disease. *Nature* 448, 704-708.
- Bethke, G., Pecher, P., Eschen-Lippold, L., Tsuda, K., Katagiri, F., Glazebrook, J., Scheel, D., and Lee, J. (2012). Activation of the Arabidopsis thaliana mitogen-activated protein kinase MPK11 by the flagellin-derived elicitor peptide, flg22. *Mol Plant Microbe Interact* 25, 471-480.
- Böhm, H., Albert, I., Fan, L., Reinhard, A., and Nürnberger, T.J.C.o.i.p.b. (2014). Immune receptor complexes at the plant cell surface. 20, 47-54.
- Bonnot, T., Gillard, M., and Nagel, D. (2019). A Simple Protocol for Informative Visualization of Enriched Gene Ontology Terms. *Bio-Protocol* 9.
- Book, A.J., Gladman, N.P., Lee, S.S., Scalf, M., Smith, L.M., and Vierstra, R.D. (2010). Affinity purification of the Arabidopsis 26 S proteasome reveals a diverse array of plant proteolytic complexes. *The Journal of biological chemistry* 285, 25554-25569.
- Chen, X.L., Xie, X., Wu, L., Liu, C., Zeng, L., Zhou, X., Luo, F., Wang, G.L., and Liu, W. (2018). Proteomic Analysis of Ubiquitinated Proteins in Rice (*Oryza sativa*) After Treatment With Pathogen-Associated Molecular Pattern (PAMP) Elicitors. *Front Plant Sci* 9, 1064.
- Chen, Y., Li, F., Tian, L., Huang, M., Deng, R., Li, X., Chen, W., Wu, P., Li, M., and Jiang, H. (2017). The phenylalanine ammonia lyase gene LJPAL1 is involved in plant defense responses to pathogens and plays diverse roles in Lotus japonicus-rhizobium symbioses. *Molecular Plant-Microbe Interactions* 30, 739-753.
- Cheng, Y.T., and Li, X. (2012). Ubiquitination in NB-LRR-mediated immunity. *Curr Opin Plant Biol* 15, 392-399.
- Cheng, Y.T., Li, Y.Z., Huang, S.A., Huang, Y., Dong, X.N., Zhang, Y.L., and Li, X. (2011). Stability

of plant immune-receptor resistance proteins is controlled by SKP1-CULLIN1-F-box (SCF)-mediated protein degradation. *Proc Natl Acad Sci USA* 108, 14694-14699.

Cheng, Z., Li, J.F., Niu, Y., Zhang, X.C., Woody, O.Z., Xiong, Y., Djonovic, S., Millet, Y., Bush, J., McConkey, B.J., *et al.* (2015). Pathogen-secreted proteases activate a novel plant immune pathway. *Nature* 521, 213-216.

Clay, N.K., Adio, A.M., Denoux, C., Jander, G., and Ausubel, F.M. (2009). Glucosinolate metabolites required for an Arabidopsis innate immune response. *Science* 323, 95-101.

Collins, N.C., Thordal-Christensen, H., Lipka, V., Bau, S., Kombrink, E., Qiu, J.L., Huckelhoven, R., Stein, M., Freialdenhoven, A., Somerville, S.C., *et al.* (2003). SNARE-protein-mediated disease resistance at the plant cell wall. *Nature* 425, 973-977.

Couto, D., and Zipfel, C. (2016). Regulation of pattern recognition receptor signalling in plants. *Nat Rev Immunol* 16, 537-552.

Cui, H., Tsuda, K., and Parker, J.E. (2015). Effector-triggered immunity: from pathogen perception to robust defense. *Annual review of plant biology* 66, 487-511.

DIELEN, A.S., Badaoui, S., Candresse, T., and GERMAN-RETANA, S. (2010). The ubiquitin/26S proteasome system in plant-pathogen interactions: a never-ending hide-and-seek game. *Molecular plant pathology* 11, 293-308.

Eschen-Lippold, L., Bethke, G., Palm-Forster, M.A., Pecher, P., Bauer, N., Glazebrook, J., Scheel, D., and Lee, J. (2012). MPK11—a fourth elicitor-responsive mitogen-activated protein kinase in Arabidopsis thaliana. *Plant signaling & behavior* 7, 1203-1205.

Ewan, R., Pangestuti, R., Thornber, S., Craig, A., Carr, C., O'Donnell, L., Zhang, C., and Sadanandom, A. (2011). Deubiquitinating enzymes AtUBP12 and AtUBP13 and their tobacco homologue NtUBP12 are negative regulators of plant immunity. *New Phytol* 191, 92-106.

Finley, D. (2009). Recognition and processing of ubiquitin-protein conjugates by the proteasome. *Annual review of biochemistry* 78, 477-513.

Fuchs, R., Kopischke, M., Klapprodt, C., Hause, G., Meyer, A.J., Schwarzlander, M., Fricker, M.D., and Lipka, V. (2016). Immobilized Subpopulations of Leaf Epidermal Mitochondria Mediate PENETRATION2-Dependent Pathogen Entry Control in Arabidopsis. *Plant Cell* 28, 130-145.

Furlan, G., Nakagami, H., Eschen-Lippold, L., Jiang, X., Majovsky, P., Kowarschik, K., Hoehenwarter, W., Lee, J., and Trujillo, M. (2017). Changes in PUB22 Ubiquitination Modes Triggered by MITOGEN-ACTIVATED PROTEIN KINASE3 Dampen the Immune Response. *The Plant cell* 29, 726-745.

Furniss, J.J., Grey, H., Wang, Z., Nomoto, M., Jackson, L., Tada, Y., and Spoel, S.H. (2018a). Proteasome-associated HECT-type ubiquitin ligase activity is required for plant immunity. *PLoS Pathog* 14, e1007447.

Furniss, J.J., Grey, H., Wang, Z., Nomoto, M., Jackson, L., Tada, Y., and Spoel, S.H. (2018b). Proteasome-associated HECT-type ubiquitin ligase activity is required for plant immunity. *PLoS pathogens* 14, e1007447.

Furniss, J.J., Grey, H., Wang, Z., Nomoto, M., Jackson, L., Tada, Y., and Spoel, S.H.J.P.p. (2018c). Proteasome-associated HECT-type ubiquitin ligase activity is required for plant immunity. *14*, e1007447.

Goritschnig, S., Zhang, Y., and Li, X. (2007). The ubiquitin pathway is required for innate immunity in Arabidopsis. *The Plant Journal* 49, 540-551.

1001 Gou, M., Shi, Z., Zhu, Y., Bao, Z., Wang, G., and Hua, J. (2012). The F-box protein CPR1/CPR30
1002 negatively regulates R protein SNC1 accumulation. *Plant J* 69, 411-420.

1003 Gou, M., Su, N., Zheng, J., Huai, J., Wu, G., Zhao, J., He, J., Tang, D., Yang, S., and Wang,
1004 G.J.T.P.J. (2009). An F-box gene, CPR30, functions as a negative regulator of the defense response
1005 in Arabidopsis. 60, 757-770.

1006 Gu, Y., and Innes, R.W. (2011). The KEEP ON GOING protein of Arabidopsis recruits the
1007 ENHANCED DISEASE RESISTANCE1 protein to trans-Golgi network/early endosome vesicles.
1008 *Plant Physiol* 155, 1827-1838.

1009 Gu, Y., Zavaliev, R., and Dong, X.J.M.p. (2017). Membrane trafficking in plant immunity. 10,
1010 1026-1034.

1011 Guerra, D.D., and Callis, J. (2012). Ubiquitin on the move: the ubiquitin modification system plays
1012 diverse roles in the regulation of endoplasmic reticulum- and plasma membrane-localized proteins.
1013 *Plant Physiol* 160, 56-64.

1014 Heidrich, K., Wirthmueller, L., Tasset, C., Pouzet, C., Deslandes, L., and Parker, J.E. (2011).
1015 Arabidopsis EDS1 connects pathogen effector recognition to cell compartment-specific immune
1016 responses. *Science* 334, 1401-1404.

1017 Hjerpe, R., Aillet, F., Lopitz-Otsoa, F., Lang, V., England, P., and Rodriguez, M.S. (2009). Efficient
1018 protection and isolation of ubiquitylated proteins using tandem ubiquitin-binding entities. *EMBO*
1019 *reports* 10, 1250-1258.

1020 Hou, S., Wang, X., Chen, D., Yang, X., Wang, M., Turra, D., Di Pietro, A., and Zhang, W. (2014).
1021 The secreted peptide PIP1 amplifies immunity through receptor-like kinase 7. *PLoS pathogens* 10,
1022 e1004331.

1023 Hou, X., and Gao, Y. (2017). Investigation on the Interaction of Pseudomonas syringae Effector
1024 AvrPto with AtRabE1d GTPase. *Protein and peptide letters* 24, 661-667.

1025 Hu, H.B., and Sun, S.C. (2016). Ubiquitin signaling in immune responses. *Cell research* 26, 457-
1026 483.

1027 Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative analysis of
1028 large gene lists using DAVID bioinformatics resources. *Nature protocols* 4, 44-57.

1029 Huang, J., Gu, M., Lai, Z., Fan, B., Shi, K., Zhou, Y.H., Yu, J.Q., and Chen, Z. (2010). Functional
1030 analysis of the Arabidopsis PAL gene family in plant growth, development, and response to
1031 environmental stress. *Plant Physiol* 153, 1526-1538.

1032 Huang, S., Chen, X., Zhong, X., Li, M., Ao, K., Huang, J., and Li, X. (2016). Plant TRAF Proteins
1033 Regulate NLR Immune Receptor Turnover. *Cell Host Microbe* 20, 271.

1034 Isono, E., and Nagel, M.K. (2014). Deubiquitylating enzymes and their emerging role in plant
1035 biology. *Front Plant Sci* 5, 56.

1036 Iwai, K., and Tokunaga, F. (2009). Linear polyubiquitination: a new regulator of NF-kappaB
1037 activation. *EMBO reports* 10, 706-713.

1038 Jacobs, A.K., Lipka, V., Burton, R.A., Panstruga, R., Strizhov, N., Schulze-Lefert, P., and Fincher,
1039 G.B. (2003). An Arabidopsis callose synthase, GSL5, is required for wound and papillary callose
1040 formation. *The Plant Cell* 15, 2503-2513.

1041 Kim, D.Y., Scalf, M., Smith, L.M., and Vierstra, R.D. (2013). Advanced proteomic analyses yield a
1042 deep catalog of ubiquitylation targets in Arabidopsis. *The Plant cell* 25, 1523-1540.

1043 Kim, S.J., Kim, M.R., Bedgar, D.L., Moinuddin, S.G., Cardenas, C.L., Davin, L.B., Kang, C., and

1044 Lewis, N.G. (2004). Functional reclassification of the putative cinnamyl alcohol dehydrogenase
1045 multigene family in Arabidopsis. *Proc Natl Acad Sci U S A* *101*, 1455-1460.

1046 Kleinboelting, N., Huep, G., Kloetgen, A., Viehoveer, P., and Weisshaar, B. (2012). GABI-Kat
1047 SimpleSearch: new features of the Arabidopsis thaliana T-DNA mutant database. *Nucleic Acids Res*
1048 *40*, D1211-1215.

1049 Komander, D., and Rape, M. (2012). The ubiquitin code. *Annu Rev Biochem* *81*, 203-229.

1050 Lachaud, C., Prigent, E., Thuleau, P., Grat, S., Da Silva, D., Brière, C., Mazars, C., and Cotellet, V.
1051 (2013). 14-3-3-Regulated Ca²⁺-dependent protein kinase CPK3 is required for sphingolipid-
1052 induced cell death in Arabidopsis. *Cell Death & Differentiation* *20*, 209-217.

1053 Le, M.H., Cao, Y., Zhang, X.C., and Stacey, G. (2014). LIK1, a CERK1-interacting kinase,
1054 regulates plant immune responses in Arabidopsis. *PLoS One* *9*, e102245.

1055 Lee, B.-J., Kwon, S.J., Kim, S.-K., Kim, K.-J., Park, C.-J., Kim, Y.-J., Park, O.K., and Paek, K.-H.
1056 (2006). Functional study of hot pepper 26S proteasome subunit RPN7 induced by Tobacco mosaic
1057 virus from nuclear proteome analysis. *Biochemical and biophysical research communications* *351*,
1058 405-411.

1059 Li, L., Habring, A., Wang, K., and Weigel, D. (2020). Atypical Resistance Protein RPW8/HR
1060 Triggers Oligomerization of the NLR Immune Receptor RPP7 and Autoimmunity. *Cell Host*
1061 *Microbe* *27*, 405-417 e406.

1062 Li, Y., Kabbage, M., Liu, W., and Dickman, M.B. (2016). Aspartyl protease-mediated cleavage of
1063 BAG6 is necessary for autophagy and fungal resistance in plants. *The Plant Cell* *28*, 233-247.

1064 Liang, X., Ding, P., Lian, K., Wang, J., Ma, M., Li, L., Li, L., Li, M., Zhang, X., Chen, S., *et al.*
1065 (2016). Arabidopsis heterotrimeric G proteins regulate immunity by directly coupling to the FLS2
1066 receptor. *Elife* *5*, e13568.

1067 Liao, D., Cao, Y., Sun, X., Espinoza, C., Nguyen, C.T., Liang, Y., and Stacey, G. (2017).
1068 Arabidopsis E3 ubiquitin ligase PLANT U-BOX13 (PUB13) regulates chitin receptor LYSIN
1069 MOTIF RECEPTOR KINASE5 (LYK5) protein abundance. *The New phytologist* *214*, 1646-1656.

1070 Liebrand, T.W., van den Burg, H.A., and Joosten, M.H. (2014). Two for all: receptor-associated
1071 kinases SOBIR1 and BAK1. *Trends in plant science* *19*, 123-132.

1072 Liu, J., Ding, P., Sun, T., Nitta, Y., Dong, O., Huang, X., Yang, W., Li, X., Botella, J.R., and Zhang,
1073 Y. (2013). Heterotrimeric G proteins serve as a converging point in plant defense signaling activated
1074 by multiple receptor-like kinases. *Plant physiology* *161*, 2146-2158.

1075 Liu, J., Li, H., Miao, M., Tang, X., Giovannoni, J., Xiao, F., and Liu, Y. (2012a). The tomato UV-
1076 damaged DNA-binding protein-1 (DDB1) is implicated in pathogenesis-related (PR) gene
1077 expression and resistance to *Agrobacterium tumefaciens*. *Molecular plant pathology* *13*, 123-134.

1078 Liu, J., Li, H., Miao, M., Tang, X., Giovannoni, J., Xiao, F., and Liu, Y.J.M.p.p. (2012b). The
1079 tomato UV-damaged DNA-binding protein-1 (DDB1) is implicated in pathogenesis-related (PR)
1080 gene expression and resistance to *Agrobacterium tumefaciens*. *13*, 123-134.

1081 Lu, D., Lin, W., Gao, X., Wu, S., Cheng, C., Avila, J., Heese, A., Devarenne, T.P., He, P., and Shan,
1082 L. (2011). Direct ubiquitination of pattern recognition receptor FLS2 attenuates plant innate
1083 immunity. *Science* *332*, 1439-1442.

1084 Ma, X., Claus, L.A.N., Leslie, M.E., Tao, K., Wu, Z., Liu, J., Yu, X., Li, B., Zhou, J., Savatin, D.V.,
1085 *et al.* (2020). Ligand-induced monoubiquitination of BIK1 regulates plant immunity. *Nature* *581*,
1086 199-203.

1087 Maor, R., Jones, A., Nuhse, T.S., Studholme, D.J., Peck, S.C., and Shirasu, K. (2007).
1088 Multidimensional protein identification technology (MudPIT) analysis of ubiquitinated proteins in
1089 plants. *Mol Cell Proteomics* 6, 601-610.

1090 Marino, D., Peeters, N., and Rivas, S. (2012). Ubiquitination during plant immune signaling. *Plant*
1091 *physiology* 160, 15-27.

1092 Marshall, R.S., Li, F., Gemperline, D.C., Book, A.J., and Vierstra, R.D. (2015). Autophagic
1093 Degradation of the 26S Proteasome Is Mediated by the Dual ATG8/Ubiquitin Receptor RPN10 in
1094 Arabidopsis. *Molecular cell* 58, 1053-1066.

1095 Marshall, R.S., and Vierstra, R.D. (2018). Autophagy: The Master of Bulk and Selective Recycling.
1096 *Annual review of plant biology* 69, 173-208.

1097 Meng, X., and Zhang, S. (2013). MAPK cascades in plant disease resistance signaling. *Annual*
1098 *review of phytopathology* 51, 245-266.

1099 Meyer, D., Pajonk, S., Micali, C., O'Connell, R., and Schulze-Lefert, P.J.T.P.J. (2009). Extracellular
1100 transport and integration of plant secretory proteins into pathogen-induced cell wall compartments.
1101 *57*, 986-999.

1102 Miao, Y., and Zentgraf, U. (2010). A HECT E3 ubiquitin ligase negatively regulates Arabidopsis
1103 leaf senescence through degradation of the transcription factor WRKY53. *The Plant Journal* 63,
1104 179-188.

1105 Monaghan, J., and Li, X. (2010). The HEAT repeat protein ILITYHIA is required for plant
1106 immunity. *Plant and cell physiology* 51, 742-753.

1107 Monaghan, J., Li, X.J.P., and physiology, c. (2010). The HEAT repeat protein ILITYHIA is required
1108 for plant immunity. *51*, 742-753.

1109 Monaghan, J., Xu, F., Gao, M., Zhao, Q., Palma, K., Long, C., Chen, S., Zhang, Y., and Li, X.
1110 (2009). Two Prp19-like U-box proteins in the MOS4-associated complex play redundant roles in
1111 plant innate immunity. *PLoS Pathogens* 5.

1112 Mondragon-Palomino, M., Stam, R., John-Arputharaj, A., and Dresselhaus, T. (2017).
1113 Diversification of defensins and NLRs in Arabidopsis species by different evolutionary
1114 mechanisms. *BMC Evol Biol* 17, 255.

1115 Mural, R.V., Liu, Y., Rosebrock, T.R., Brady, J.J., Hamera, S., Connor, R.A., Martin, G.B., and
1116 Zeng, L. (2013). The tomato Fni3 lysine-63-specific ubiquitin-conjugating enzyme and suv
1117 ubiquitin E2 variant positively regulate plant immunity. *Plant Cell* 25, 3615-3631.

1118 Murata, S., Yashiroda, H., and Tanaka, K. (2009). Molecular mechanisms of proteasome assembly.
1119 *Nature reviews Molecular cell biology* 10, 104-115.

1120 Nakashima, A., Chen, L., Thao, N.P., Fujiwara, M., Wong, H.L., Kuwano, M., Umemura, K.,
1121 Shirasu, K., Kawasaki, T., and Shimamoto, K. (2008). RACK1 functions in rice innate immunity by
1122 interacting with the Rac1 immune complex. *Plant Cell* 20, 2265-2279.

1123 Nicaise, V., Joe, A., Jeong, B., Korneli, C., Boutrot, F., Wested, I., Staiger, D., Alfano, J., and Zipfel,
1124 C. (2013). Pseudomonas HopU1 affects interaction of plant immune receptor mRNAs to the RNA-
1125 binding protein GRP7. *EMBO J* 32, 701-712.

1126 Okumoto, K., Misono, S., Miyata, N., Matsumoto, Y., Mukai, S., and Fujiki, Y. (2011). Cysteine
1127 ubiquitination of PTS1 receptor Pex5p regulates Pex5p recycling. *Traffic* 12, 1067-1083.

1128 Ordureau, A., Paulo, J.A., Zhang, J., An, H., Swatek, K.N., Cannon, J.R., Wan, Q., Komander, D.,
1129 and Harper, J.W. (2020). Global Landscape and Dynamics of Parkin and USP30-Dependent

Ubiquitylomes in iNeurons during Mitophagic Signaling. *Molecular cell* 77, 1124-1142 e1110.

Paez Valencia, J., Goodman, K., and Otegui, M.S. (2016). Endocytosis and Endosomal Trafficking in Plants. *Annual review of plant biology* 67, 309-335.

Palm, D., Streit, D., Shanmugam, T., Weis, B.L., Ruprecht, M., Simm, S., and Schleiff, E. (2019). Plant-specific ribosome biogenesis factors in *Arabidopsis thaliana* with essential function in rRNA processing. *Nucleic Acids Res* 47, 1880-1895.

Parker, J.E., Holub, E.B., Frost, L.N., Falk, A., Gunn, N.D., and Daniels, M.J. (1996). Characterization of eds1, a mutation in *Arabidopsis* suppressing resistance to *Peronospora parasitica* specified by several different RPP genes. *The Plant Cell* 8, 2033-2046.

Pastorczyk, M., and Bednarek, P. (2016). The function of glucosinolates and related metabolites in plant innate immunity. In *Advances in Botanical Research* (Elsevier), pp. 171-198.

Pecher, P., Eschen-Lippold, L., Herklotz, S., Kuhle, K., Naumann, K., Bethke, G., Uhrig, J., Weyhe, M., Scheel, D., and Lee, J. (2014). The *Arabidopsis thaliana* mitogen-activated protein kinases MPK 3 and MPK 6 target a subclass of 'VQ-motif'-containing proteins to regulate immune responses. *New Phytologist* 203, 592-606.

Peng, J., Schwartz, D., Elias, J.E., Thoreen, C.C., Cheng, D., Marsischky, G., Roelofs, J., Finley, D., and Gygi, S.P. (2003). A proteomics approach to understanding protein ubiquitination. *Nat Biotechnol* 21, 921-926.

Perraki, A., Gronnier, J., Gouguet, P., Boudsocq, M., Deroubaix, A.-F., Simon, V., German-Retana, S., Zipfel, C., Bayer, E., and Mongrand, S. (2017). The plant calcium-dependent protein kinase CPK3 phosphorylates REM1.3 to restrict viral infection. *BioRxiv*, 205765.

Pitorre, D., Llauro, C., Jobet, E., Guillemot, J., Brizard, J.-P., Delseny, M., and Lasserre, E. (2010). RLK7, a leucine-rich repeat receptor-like kinase, is required for proper germination speed and tolerance to oxidative stress in *Arabidopsis thaliana*. *Planta* 232, 1339-1353.

Qin, L., Zhou, Z., Li, Q., Zhai, C., Liu, L., Quilichini, T.D., Gao, P., Kessler, S.A., Jaillais, Y., Datla, R., *et al.* (2020). Specific Recruitment of Phosphoinositide Species to the Plant-Pathogen Interfacial Membrane Underlies *Arabidopsis* Susceptibility to Fungal Infection. *Plant Cell* 32, 1665-1688.

Raasi, S., Orlov, I., Fleming, K.G., and Pickart, C.M. (2004). Binding of polyubiquitin chains to ubiquitin-associated (UBA) domains of HHR23A. *Journal of molecular biology* 341, 1367-1379.

Rayapuram, N., Bigeard, J., Alhoraibi, H., Bonhomme, L., Hesse, A.M., Vinh, J., Hirt, H., and Pflieger, D. (2018). Quantitative Phosphoproteomic Analysis Reveals Shared and Specific Targets of *Arabidopsis* Mitogen-Activated Protein Kinases (MAPKs) MPK3, MPK4, and MPK6. *Mol Cell Proteomics* 17, 61-80.

Revers, F., Guiraud, T., Houvenaghel, M.-C., Mauduit, T., Le Gall, O., and Candresse, T. (2003). Multiple resistance phenotypes to Lettuce mosaic virus among *Arabidopsis thaliana* accessions. *Molecular plant-microbe interactions* 16, 608-616.

Reyes, F.C., Buono, R., and Otegui, M.S. (2011). Plant endosomal trafficking pathways. *Current opinion in plant biology* 14, 666-673.

Rodrigues, O., Reshetnyak, G., Grondin, A., Saijo, Y., Leonhardt, N., Maurel, C., and Verdoucq, L. (2017). Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA- and pathogen-triggered stomatal closure. *Proceedings of the National Academy of Sciences* 114, 9200-9205.

Romero-Barrios, N., and Vert, G. (2018). Proteasome-independent functions of lysine-63

polyubiquitination in plants. *The New phytologist* *217*, 995-1011.

Saracco, S.A., Hansson, M., Scalf, M., Walker, J.M., Smith, L.M., and Vierstra, R.D. (2009). Tandem affinity purification and mass spectrometric analysis of ubiquitylated proteins in *Arabidopsis*. *Plant J* *59*, 344-358.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T.J.G.r. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *13*, 2498-2504.

Shimizu, Y., Okuda-Shimizu, Y., and Hendershot, L.M. (2010). Ubiquitylation of an ERAD substrate occurs on multiple types of amino acids. *Molecular cell* *40*, 917-926.

Skubacz, A., Daszkowska-Golec, A., and Szarejko, L. (2016). The Role and Regulation of ABI5 (ABA-Insensitive 5) in Plant Development, Abiotic Stress Responses and Phytohormone Crosstalk. *Front Plant Sci* *7*.

Smalle, J., and Vierstra, R.D. (2004). The ubiquitin 26S proteasome proteolytic pathway. *Annual review of plant biology* *55*, 555-590.

Speth, E.B., Imboden, L., Hauck, P., and He, S.Y. (2009a). Subcellular localization and functional analysis of the *Arabidopsis* GTPase RabE. *Plant physiology* *149*, 1824-1837.

Speth, E.B., Imboden, L., Hauck, P., and He, S.Y. (2009b). Subcellular Localization and Functional Analysis of the *Arabidopsis* GTPase RabE. *149*, 1824-1837.

Spoel, S.H., and Dong, X. (2012). How do plants achieve immunity? Defence without specialized immune cells. *Nature reviews Immunology* *12*, 89-100.

Stegmann, M., Anderson, R.G., Ichimura, K., Pecenkova, T., Reuter, P., Zarsky, V., McDowell, J.M., Shirasu, K., and Trujillo, M. (2012). The ubiquitin ligase PUB22 targets a subunit of the exocyst complex required for PAMP-triggered responses in *Arabidopsis*. *Plant Cell* *24*, 4703-4716.

Stringlis, I.A., Yu, K., Feussner, K., de Jonge, R., Van Bentum, S., Van Verk, M.C., Berendsen, R.L., Bakker, P.A., Feussner, I., and Pieterse, C.M. (2018). MYB72-dependent coumarin exudation shapes root microbiome assembly to promote plant health. *Proceedings of the National Academy of Sciences* *115*, E5213-E5222.

Su, J., Xu, J., and Zhang, S. (2015). RACK1, scaffolding a heterotrimeric G protein and a MAPK cascade. *Trends Plant Sci* *20*, 405-407.

Su, J., Yang, L., Zhu, Q., Wu, H., He, Y., Liu, Y., Xu, J., Jiang, D., and Zhang, S. (2018). Active photosynthetic inhibition mediated by MPK3/MPK6 is critical to effector-triggered immunity. *PLoS biology* *16*, e2004122.

Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J., Simonovic, M., Roth, A., Santos, A., Tsafou, K.P., *et al.* (2015). STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* *43*, D447-452.

Tan, X., Meyers, B.C., Kozik, A., West, M.A., Morgante, M., St Clair, D.A., Bent, A.F., and Michelson, R.W. (2007). Global expression analysis of nucleotide binding site-leucine rich repeat-encoding and related genes in *Arabidopsis*. *BMC Plant Biol* *7*, 56.

Tarutani, Y., Morimoto, T., Sasaki, A., Yasuda, M., Nakashita, H., Yoshida, S., Yamaguchi, I., and Suzuki, Y. (2004). Molecular characterization of two highly homologous receptor-like kinase genes, RLK902 and RKL1, in *Arabidopsis thaliana*. *Biosci Biotechnol Biochem* *68*, 1935-1941.

Thordal-Christensen, H. (2003). Fresh insights into processes of nonhost resistance. *Curr Opin Plant Biol* *6*, 351-357.

Tronchet, M., Balague, C., Kroj, T., Jouanin, L., and Roby, D. (2010). Cinnamyl alcohol dehydrogenases-C and D, key enzymes in lignin biosynthesis, play an essential role in disease resistance in Arabidopsis. *Molecular plant pathology* *11*, 83-92.

Trujillo, M., and Shirasu, K. (2010). Ubiquitination in plant immunity. *Current opinion in plant biology* *13*, 402-408.

Udeshi, N.D., Svinkina, T., Mertins, P., Kuhn, E., Mani, D.R., Qiao, J.W., and Carr, S.A. (2013). Refined Preparation and Use of Anti-diglycine Remnant (K-epsilon-GG) Antibody Enables Routine Quantification of 10,000s of Ubiquitination Sites in Single Proteomics Experiments. *Molecular & Cellular Proteomics* *12*, 825-831.

Urano, D., Chen, J.G., Botella, J.R., and Jones, A.M. (2013). Heterotrimeric G protein signalling in the plant kingdom. *Open Biol* *3*, 120186.

Üstün, S., Sheikh, A., Gimenez-Ibanez, S., Jones, A., Ntoukakis, V., and Börnke, F. (2016). The proteasome acts as a hub for plant immunity and is targeted by *Pseudomonas* type III effectors. *Plant Physiology* *172*, 1941-1958.

van Nocker, S., Walker, J.M., and Vierstra, R.D. (1996). The Arabidopsis thaliana UBC7/13/14 genes encode a family of multiubiquitin chain-forming E2 enzymes. *The Journal of biological chemistry* *271*, 12150-12158.

Vierstra, R.D. (2009). The ubiquitin-26S proteasome system at the nexus of plant biology. *Nat Rev Mol Cell Biol* *10*, 385-397.

Wang, J., Grubb, L.E., Wang, J., Liang, X., Li, L., Gao, C., Ma, M., Feng, F., Li, M., Li, L., *et al.* (2018). A Regulatory Module Controlling Homeostasis of a Plant Immune Kinase. *Molecular cell*.

Wang, L., Wen, R., Wang, J., Xiang, D., Wang, Q., Zang, Y., Wang, Z., Huang, S., Li, X., and Datla, R.J.N.P. (2019). Arabidopsis UBC 13 differentially regulates two programmed cell death pathways in responses to pathogen and low-temperature stress. *221*, 919-934.

Wawrzynska, A., Rodibaugh, N.L., and Innes, R.W. (2010). Synergistic activation of defense responses in Arabidopsis by simultaneous loss of the GSL5 callose synthase and the EDR1 protein kinase. *Mol Plant Microbe Interact* *23*, 578-584.

Wen, R., Torres-Acosta, J.A., Pastushok, L., Lai, X., Pelzer, L., Wang, H., and Xiao, W. (2008). Arabidopsis UEV1D promotes Lysine-63-linked polyubiquitination and is involved in DNA damage response. *The Plant cell* *20*, 213-227.

Yao, C., Wu, Y., Nie, H., and Tang, D. (2012). RPN1a, a 26S proteasome subunit, is required for innate immunity in Arabidopsis. *The Plant journal : for cell and molecular biology* *71*, 1015-1028.

Yu, X., Feng, B., He, P., and Shan, L. (2017). From Chaos to Harmony: Responses and Signaling upon Microbial Pattern Recognition. *Annual review of phytopathology* *55*, 109-137.

Zamioudis, C., Hanson, J., and Pieterse, C.M. (2014). β -Glucosidase BGLU 42 is a MYB 72-dependent key regulator of rhizobacteria-induced systemic resistance and modulates iron deficiency responses in Arabidopsis roots. *New Phytologist* *204*, 368-379.

Zhao, J., Zhou, H., Zhang, M., Gao, Y., Li, L., Gao, Y., Li, M., Yang, Y., Guo, Y., and Li, X. (2016). Ubiquitin-specific protease 24 negatively regulates abscisic acid signalling in Arabidopsis thaliana. *Plant, cell & environment* *39*, 427-440.

Zhao, Y., Wu, G., Shi, H., and Tang, D. (2019). RECEPTOR-LIKE KINASE 902 Associates with and Phosphorylates BRASSINOSTEROID-SIGNALING KINASE1 to Regulate Plant Immunity. *Mol Plant* *12*, 59-70.

1259 Zhou, B., and Zeng, L. (2017). Conventional and unconventional ubiquitination in plant immunity.
1260 *Molecular plant pathology* 18, 1313-1330.

1261 Zhou, J.-M., and Zhang, Y. (2020). Plant Immunity: Danger Perception and Signaling. *Cell*.

1262 Zhou, J., He, P., and Shan, L. (2014). Ubiquitination of plant immune receptors. *Methods Mol Biol*
1263 1209, 219-231.

1264 Zhou, J., Liu, D., Wang, P., Ma, X., Lin, W., Chen, S., Mishev, K., Lu, D., Kumar, R., Vanhoutte, I.,
1265 *et al.* (2018). Regulation of Arabidopsis brassinosteroid receptor BRI1 endocytosis and degradation
1266 by plant U-box PUB12/PUB13-mediated ubiquitination. *Proc Natl Acad Sci U S A* 115, E1906-
1267 E1915.

1268 Zhou, J., Lu, D., Xu, G., Finlayson, S.A., He, P., and Shan, L. (2015). The dominant negative ARM
1269 domain uncovers multiple functions of PUB13 in Arabidopsis immunity, flowering, and senescence.
1270 *J Exp Bot* 66, 3353-3366.

1271

Figure 1. Two-step affinity purification of ubiquitylated proteins from transgenic Arabidopsis expressing 6His-UBQ treated with or without flg22.

A. Diagram of the p35S::hexa-6His-UBQ transgene. A synthetic UBQ gene encoding six repeats of a Ub monomer N-terminally tagged with a 6His sequence followed by a glycine (G)-rich linker (MHHHHHHGGGGGSA) was fused head-to-tail to form a single in-frame hexa-6His-UBQ transgene expressed under the control of the constitutive CaMV 35S promoter (Saracco et al., 2009).

B. Flow chart describing the proteomic approach to stringently identify Ub conjugates from Arabidopsis p35S::hexa-6His-UBQ seedlings. Ubiquitylated proteins were first enriched by the UBA-Ub affinity using GST-TUBEs beads under the native condition followed by Ni-NTA chromatography under the denaturing condition. The eluants were digested with trypsin and subjected to LC/ESI-MS/MS analysis.

C and D. Ubiquitylated proteins isolated from Arabidopsis plants expressing 6His-Ub by the two-step affinity purification. Ubiquitylated proteins enriched as outlined in (B) were subjected to SDS-PAGE and by either stained for protein with silver (C) or immunoblotted with anti-Ub antibodies (D). Samples from wild-type seedlings (WT) purified with both GST-TUBEs and Ni-NTA beads were included for comparison. FT and EL represent the flow-through and eluted fractions, respectively, from the Ub affinity (1) and Ni-NTA affinity columns (2). The total represents the total protein extracts before affinity purifications. The migration positions of the Ub dimers (Ub₂), and Ub conjugates (Ub-conj) are indicated by the brackets.

Figure 2. Gene ontology (GO) analysis of flg22-treated and untreated ubiquitylomes.

A. Venn diagram showing the overlap of the total ubiquitylated proteins isolated from plants with or without flg22 exposure.

B. Gene ontology (GO) enrichment of ubiquitylated proteins in control or flg22 treated ubiquitylomes. GO analysis in biological processes, molecular functions, and cellular components was performed using the DAVID Functional Annotation Tool (Huang da et al., 2009) and GO annotations from the TAIR GO database (<http://www.arabidopsis.org/>). The fold enrichment was calculated based on the frequency of proteins annotated to the term compared with their frequency in the total proteome. The dot size indicates the number of genes associated with each process and the dot color indicates the significance of the enrichment. FDR was based on the corrected p-values. The vertical grey dashed line represents a fold enrichment of 1. The total number of GO terms was listed under each GO domain (biological processes, molecular functions, and cellular components). The selected terms related to stress and the UPS are shown.

C. Venn diagram showing the overlaps of ubiquitylated proteins with or without flg22 in the selected categories. Proteins were categorized into functional groups based on the GO annotations; those related to defense responses and the UPS are shown.

Figure 3. Protein interaction network of ubiquitylated proteins identified from 6HIS-UBQ seedlings treated with or without flg22.

A. Protein interaction networks were generated with the list of 391 (control) and 570 (flg22-treated) ubiquitylated proteins using the STRING database and visualized in Cytoscape. The node colors indicate proteins identified in control (yellow), flg22-treated (red), or both conditions (blue) of datasets. Clusters of the proteasome, ubiquitylation, and defense-related components are marked by dashed lines.

B. Map of proposed subcellular locations and functions of immunity-related proteins identified from the ubiquitylomes.

Figure 4. In vivo confirmation that several candidates are ubiquitylated in Arabidopsis.

Arabidopsis wild-type protoplasts were co-transfected with FLAG-tagged Ub (FLAG-UBQ) and HA- or myc-tagged substrate or a control vector (Ctrl) and incubated for 10 hr followed by treatment with 100 nM flg22 for the indicated. Protein extracts were immunoprecipitated with anti-FLAG agarose beads (IP: α -FLAG) and the ubiquitylated proteins were immunoblotted with anti-HA or anti-myc antibodies (top left) or anti-FLAG antibodies (bottom left). The input controls were shown by an anti-HA/myc (top right) or anti-FLAG (bottom right) immunoblots.

A. Monoubiquitylation of MPK3 is enhanced upon flg22 treatment. Protoplasts were isolated at 0 to 30 min after flg22 exposure.

B. Polyubiquitylation of SOBIR1 in vivo is enhanced upon a 30 min treatment with flg22.

C, D. MKK5, and EDS1 were ubiquitylated in vivo following a 30-min treatment with flg22.

The above experiments were performed three times with similar results.

Figure 5. RKL1 is ubiquitinated and involved in PTI responses.

A. Diagram of the RKL1 protein. The signal peptide (SP), leucine-rich repeat (LRR), transmembrane (TM), and protein kinase (K) domains are shown. The amino acids demarcating each domain are shown.

B. In vivo ubiquitylation of RKL1 upon flg22 treatment. Protoplasts were co-transfected with FLAG-UBQ and HA-tagged RKL1 (RKL1-HA) or a control vector (Ctrl) and incubated for 10 hr followed by treatment with 100 nM flg22 for 30 min. Proteins were immunoprecipitated with anti-FLAG agarose beads (IP: α-FLAG) and the ubiquitylated proteins were detected by immunoblotting with anti-HA antibodies (top). The input controls were shown by an anti-HA immunoblot (bottom).

C. flg22-induced reactive oxygen species (ROS) burst is elevated in the *rkl1-1* mutants. Leaf disks of four-week-old soil-grown Arabidopsis WT or *rkl1-1* plants (SALK 099094) were treated with 100 nM flg22 or water as a control, and ROS production was monitored over time. The data are shown as mean ± s.e.m. (n = 16) of relative fluorescence units (RLU) overlaid on the dot plot.

D. *rkl1* mutants have elevated resistance against Pst DC3000. Leaf disks of four-week-old soil-grown Arabidopsis WT or *rkl1-1* plants were hand-infiltrated with Pst DC3000 at OD₆₀₀=5×10⁻⁴ CFU/ml, and bacterial growth was counted at 0 or 3 dpi. The data are shown as mean ± s.e.m. overlaid on dot plot (n = 3 for 0 dpi; n = 4 for 3 dpi). The experiments were performed three times with similar results.

Figure 6. UBC13 is ubiquitylated and negatively regulates PTI responses.

A. In vivo ubiquitylation of UBC13 upon flg22 treatment. Protoplasts were co-transfected with FLAG-UBQ and HA-tagged UBC13 (UBC13-HA) and incubated for 10 hr followed by treatment with 100 nM flg22 for 30 min. Protein extracts were immunoprecipitated with anti-FLAG beads (IP: α-FLAG) and the ubiquitylated proteins were immunoblotted with anti-HA antibodies (top). The input control was shown by an anti-HA immunoblot (bottom).

B, C. UBC13 negatively regulates flg22-triggered ROS production. Leaf disks of four-week-old WT, *ubc13* (CS389049/*ubc13-1* & CS389056/*ubc13-2*), and plants harboring the p35S::UBC13-HA transgene in the *ubc13-2* mutant background were treated with 100 nM flg22. ROS production was monitored over time. The data are shown as mean $\pm$ s.e.m overlaid on dot plot ($n = 16$) of RLU. The total photon count was shown in (C).

D. *ubc13* mutants have elevated resistance against Pst DC3000. Leaf disks of four-week-old soil-grown WT, *ubc13-1*, *ubc13-2*, and p35S::UBC13-HA *ubc13-2* (Line 1 and 2) plants were hand-infiltrated with Pst DC3000 at OD600=5 $\times$ 10⁻⁴ CFU/ml, and bacterial growth was counted at 3 dpi. The data are shown as mean $\pm$ s.e.m overlaid on the dot plot ($n = 3$). The experiments were performed three times with similar results.

Figure 7. RPN8b is ubiquitinated and negatively regulates PTI responses.

A. in vivo ubiquitylation of RPN8b upon flg22 treatment. Protoplasts were co-transfected with FLAG-UBQ and HA-tagged RPN8b (RPN8b-HA) or a control vector (Ctrl) and incubated for 10 hr followed by treatment with 100 nM flg22 for 30 min. Protein extracts from protoplasts were immunoprecipitated with anti-FLAG beads (IP: α-FLAG) and the ubiquitinated proteins were immunoblotted with anti-HA antibodies (top). The input control was shown by an anti-HA immunoblot (bottom).

B, C. RPN8b negatively regulates flg22-triggered ROS production. Leaf disks of four-week-old WT, *rpn8b-1* (SALK_128568), *rpn8b-2* (SALK_023568), and *rpn8b-1* plants harboring the p35S::RPN8b-HA transgene were treated with 100 nM flg22 or water as a control, and ROS production was monitored over time. The data are shown as mean ± s.e.m overlaid on the dot plot ($n \geq 13$) of RLU. The total photon count was shown in (C).

D. *rpn8b* mutants have elevated resistance against Pst DC3000. Leaf disks from four-week-old soil-grown WT, *rpn8b-1*, *rpn8b-2*, and 35S::RPN8b *rpn8b-1* plants were hand-infiltrated with Pst DC3000 at OD600=5×10⁻⁴ CFU/ml, and bacterial growth was counted at 3 dpi. The data are shown as mean ± s.e.m overlaid on dot plot ($n = 3$). The experiments were performed three times with similar results.

Figure 8. MS/MS mapping of Ub-Ub linkages.

A. Strategy for detecting ubiquitylation sites by MS/MS. After trypsin digestion, a diglycine remnant of Ub covalently appended to a lysine residue of a conjugated protein is detected by an increased mass of 114 kDa for the lysine residue, along with protection from trypsin cleavage after that residue. Amino acids are denoted by single-letter code.

B. The distribution of Ub-Ub linkages across the seven Ub lysines based on MS analysis from Arabidopsis transgenic plants carrying 6HIS-UBQ seedlings treated with or without 100 nM flg22. Percentage of Ub footprints at each site were obtained from PSM counts of diagnostic footprint peptides.

Parsed Citations

- Adams, E.H.G., and Spoel, S.H. (2018). The ubiquitin-proteasome system as a transcriptional regulator of plant immunity. *J Exp Bot* 69, 4529-4537.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Aguilar-Hernandez, V., Kim, D.Y., Stankey, R.J., Scalf, M., Smith, L.M., and Vierstra, R.D. (2017). Mass Spectrometric Analyses Reveal a Central Role for Ubiquitylation in Remodeling the Arabidopsis Proteome during Photomorphogenesis. *Mol Plant* 10, 846-865.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Albert, I., Böhm, H., Albert, M., Feiler, C.E., Imkampe, J., Wallmeroth, N., Brancato, C., Raaymakers, T.M., Oome, S., and Zhang, H. (2015). An RLP23-SOBIR1-BAK1 complex mediates NLP-triggered immunity. *Nature Plants* 1, 1-9.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Alonso, J.M., Stepanova, A.N., Leisse, T.J., Kim, C.J., Chen, H., Shinn, P., Stevenson, D.K., Zimmerman, J., Barajas, P., Cheuk, R., et al. (2003). Genome-wide insertional mutagenesis of Arabidopsis thaliana. *Science* 301, 653-657.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Asai, T., Tena, G., Plotnikova, J., Willmann, M.R., Chiu, W.L., Gomez-Gomez, L., Boller, T., Ausubel, F.M., and Sheen, J. (2002). MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature* 415, 977-983.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bakker, E.G., Toomajian, C., Kreitman, M., and Bergelson, J. (2006). A genome-wide survey of R gene polymorphisms in Arabidopsis. *Plant Cell* 18, 1803-1818.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bellati, J., Champeyroux, C., Hem, S., Rofidal, V., Krouk, G., Maurel, C., and Santoni, V. (2016). Novel Aquaporin Regulatory Mechanisms Revealed by Interactomics. *Mol Cell Proteomics* 15, 3473-3487.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bennett, E.J., Shaler, T.A., Woodman, B., Ryu, K.Y., Zaitseva, T.S., Becker, C.H., Bates, G.P., Schulman, H., and Kopito, R.R. (2007). Global changes to the ubiquitin system in Huntington's disease. *Nature* 448, 704-708.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bethke, G., Pecher, P., Eschen-Lippold, L., Tsuda, K., Katagiri, F., Glazebrook, J., Scheel, D., and Lee, J. (2012). Activation of the Arabidopsis thaliana mitogen-activated protein kinase MPK11 by the flagellin-derived elicitor peptide, flg22. *Mol Plant Microbe Interact* 25, 471-480.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Böhm, H., Albert, I., Fan, L., Reinhard, A., and Nürnberger, T.J.C.o.i.p.b. (2014). Immune receptor complexes at the plant cell surface. *20*, 47-54.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bonnot, T., Gillard, M., and Nagel, D. (2019). A Simple Protocol for Informative Visualization of Enriched Gene Ontology Terms. *Bio-Protocol* 9.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Book, A.J., Gladman, N.P., Lee, S.S., Scalf, M., Smith, L.M., and Vierstra, R.D. (2010). Affinity purification of the Arabidopsis 26 S proteasome reveals a diverse array of plant proteolytic complexes. *The Journal of biological chemistry* 285, 25554-25569.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Chen, X.L., Xie, X., Wu, L., Liu, C., Zeng, L., Zhou, X., Luo, F., Wang, G.L., and Liu, W. (2018). Proteomic Analysis of Ubiquitinated Proteins in Rice (Oryza sativa) After Treatment With Pathogen-Associated Molecular Pattern (PAMP) Elicitors. *Front Plant Sci* 9, 1064.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Chen, Y., Li, F., Tian, L., Huang, M., Deng, R., Li, X., Chen, W., Wu, P., Li, M., and Jiang, H. (2017). The phenylalanine ammonia lyase gene LJPAL1 is involved in plant defense responses to pathogens and plays diverse roles in Lotus japonicus-rhizobium symbioses. *Molecular Plant-Microbe Interactions* 30, 739-753.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Cheng, Y.T., and Li, X. (2012). Ubiquitination in NB-LRR-mediated immunity. *Curr Opin Plant Biol* 15, 392-399.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Cheng, Y.T., Li, Y.Z., Huang, S.A., Huang, Y., Dong, X.N., Zhang, Y.L., and Li, X. (2011). Stability of plant immune-receptor resistance proteins is controlled by SKP1-CULLIN1-F-box (SCF)-mediated protein degradation. *P Natl Acad Sci USA* 108, 14694-14699.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Cheng, Z., Li, J.F., Niu, Y., Zhang, X.C., Woody, O.Z., Xiong, Y., Djonovic, S., Millet, Y., Bush, J., McConkey, B.J., et al. (2015). Pathogen-secreted proteases activate a novel plant immune pathway. *Nature* 521, 213-216.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Clay, N.K., Adio, A.M., Denoux, C., Jander, G., and Ausubel, F.M. (2009). Glucosinolate metabolites required for an Arabidopsis innate immune response. *Science* 323, 95-101.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Collins, N.C., Thordal-Christensen, H., Lipka, V., Bau, S., Kombrink, E., Qiu, J.L., Huckelhoven, R., Stein, M., Freialdenhoven, A., Somerville, S.C., et al. (2003). SNARE-protein-mediated disease resistance at the plant cell wall. *Nature* 425, 973-977.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Couto, D., and Zipfel, C. (2016). Regulation of pattern recognition receptor signalling in plants. *Nat Rev Immunol* 16, 537-552.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cui, H., Tsuda, K., and Parker, J.E. (2015). Effector-triggered immunity: from pathogen perception to robust defense. *Annual review of plant biology* 66, 487-511.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

DIELEN, A.S., Badaoui, S., Candresse, T., and GERMAN-RETANA, S. (2010). The ubiquitin/26S proteasome system in plant-pathogen interactions: a never-ending hide-and-seek game. *Molecular plant pathology* 11, 293-308.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Eschen-Lippold, L., Bethke, G., Palm-Forster, M.A., Pecher, P., Bauer, N., Glazebrook, J., Scheel, D., and Lee, J. (2012). MPK11-a fourth elicitor-responsive mitogen-activated protein kinase in *Arabidopsis thaliana*. *Plant signaling & behavior* 7, 1203-1205.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ewan, R., Pangestuti, R., Thornber, S., Craig, A., Carr, C., O'Donnell, L., Zhang, C., and Sadanandom, A. (2011). Deubiquitinating enzymes AtUBP12 and AtUBP13 and their tobacco homologue NtUBP12 are negative regulators of plant immunity. *New Phytol* 191, 92-106.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Finley, D. (2009). Recognition and processing of ubiquitin-protein conjugates by the proteasome. *Annual review of biochemistry* 78, 477-513.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fuchs, R., Kopischke, M., Klapprodt, C., Hause, G., Meyer, A.J., Schwarzlander, M., Fricker, M.D., and Lipka, V. (2016). Immobilized Subpopulations of Leaf Epidermal Mitochondria Mediate PENETRATION2-Dependent Pathogen Entry Control in *Arabidopsis*. *Plant Cell* 28, 130-145.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Furlan, G., Nakagami, H., Eschen-Lippold, L., Jiang, X., Majovsky, P., Kowarschik, K., Hoehenwarter, W., Lee, J., and Trujillo, M. (2017). Changes in PUB22 Ubiquitination Modes Triggered by MITOGEN-ACTIVATED PROTEIN KINASE3 Dampen the Immune Response. *The Plant cell* 29, 726-745.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Furniss, J.J., Grey, H., Wang, Z., Nomoto, M., Jackson, L., Tada, Y., and Spoel, S.H. (2018a). Proteasome-associated HECT-type ubiquitin ligase activity is required for plant immunity. *PLoS Pathog* 14, e1007447.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Furniss, J.J., Grey, H., Wang, Z., Nomoto, M., Jackson, L., Tada, Y., and Spoel, S.H. (2018b). Proteasome-associated HECT-type ubiquitin ligase activity is required for plant immunity. *PLoS pathogens* 14, e1007447.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Furniss, J.J., Grey, H., Wang, Z., Nomoto, M., Jackson, L., Tada, Y., and Spoel, S.H.J.P.p. (2018c). Proteasome-associated HECT-type ubiquitin ligase activity is required for plant immunity. 14, e1007447.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Goritschnig, S., Zhang, Y., and Li, X. (2007). The ubiquitin pathway is required for innate immunity in *Arabidopsis*. *The Plant Journal* 49, 540-551.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gou, M., Shi, Z., Zhu, Y., Bao, Z., Wang, G., and Hua, J. (2012). The F-box protein CPR1/CPR30 negatively regulates R protein SNC1 accumulation. *Plant J* 69, 411-420.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gou, M., Su, N., Zheng, J., Huai, J., Wu, G., Zhao, J., He, J., Tang, D., Yang, S., and Wang, G.J.T.P.J. (2009). An F-box gene, CPR30, functions as a negative regulator of the defense response in *Arabidopsis*. 60, 757-770.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gu, Y., and Innes, R.W. (2011). The KEEP ON GOING protein of *Arabidopsis* recruits the ENHANCED DISEASE RESISTANCE1 protein to trans-Golgi network/early endosome vesicles. *Plant Physiol* 155, 1827-1838.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gu, Y., Zavaliev, R., and Dong, X.J.M.p. (2017). Membrane trafficking in plant immunity. 10, 1026-1034.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Guerra, D.D., and Callis, J. (2012). Ubiquitin on the move: the ubiquitin modification system plays diverse roles in the regulation of endoplasmic reticulum- and plasma membrane-localized proteins. *Plant Physiol* 160, 56-64.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

- Heidrich, K., Wirthmueller, L., Tasset, C., Pouzet, C., Deslandes, L., and Parker, J.E. (2011). Arabidopsis EDS1 connects pathogen effector recognition to cell compartment-specific immune responses. *Science* 334, 1401-1404.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Hjerpe, R., Aillet, F., Lopitz-Otsoa, F., Lang, V., England, P., and Rodriguez, M.S. (2009). Efficient protection and isolation of ubiquitylated proteins using tandem ubiquitin-binding entities. *EMBO reports* 10, 1250-1258.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Hou, S., Wang, X., Chen, D., Yang, X., Wang, M., Turra, D., Di Pietro, A., and Zhang, W. (2014). The secreted peptide PIP1 amplifies immunity through receptor-like kinase 7. *PLoS pathogens* 10, e1004331.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Hou, X., and Gao, Y. (2017). Investigation on the Interaction of Pseudomonas syringae Effector AvrPto with AtRabE1d GTPase. *Protein and peptide letters* 24, 661-667.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Hu, H.B., and Sun, S.C. (2016). Ubiquitin signaling in immune responses. *Cell research* 26, 457-483.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature protocols* 4, 44-57.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Huang, J., Gu, M., Lai, Z., Fan, B., Shi, K., Zhou, Y.H., Yu, J.Q., and Chen, Z. (2010). Functional analysis of the Arabidopsis PAL gene family in plant growth, development, and response to environmental stress. *Plant Physiol* 153, 1526-1538.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Huang, S., Chen, X., Zhong, X., Li, M., Ao, K., Huang, J., and Li, X. (2016). Plant TRAF Proteins Regulate NLR Immune Receptor Turnover. *Cell Host Microbe* 20, 271.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Isono, E., and Nagel, M.K. (2014). Deubiquitylating enzymes and their emerging role in plant biology. *Front Plant Sci* 5, 56.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Iwai, K., and Tokunaga, F. (2009). Linear polyubiquitination: a new regulator of NF-kappaB activation. *EMBO reports* 10, 706-713.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Jacobs, A.K., Lipka, V., Burton, R.A., Panstruga, R., Strizhov, N., Schulze-Lefert, P., and Fincher, G.B. (2003). An Arabidopsis callose synthase, GSL5, is required for wound and papillary callose formation. *The Plant Cell* 15, 2503-2513.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Kim, D.Y., Scalf, M., Smith, L.M., and Vierstra, R.D. (2013). Advanced proteomic analyses yield a deep catalog of ubiquitylation targets in Arabidopsis. *The Plant cell* 25, 1523-1540.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Kim, S.J., Kim, M.R., Bedgar, D.L., Moinuddin, S.G., Cardenas, C.L., Davin, L.B., Kang, C., and Lewis, N.G. (2004). Functional reclassification of the putative cinnamyl alcohol dehydrogenase multigene family in Arabidopsis. *Proc Natl Acad Sci U S A* 101, 1455-1460.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Kleinboelting, N., Huep, G., Kloetgen, A., Viehoveer, P., and Weisshaar, B. (2012). GABI-Kat SimpleSearch: new features of the Arabidopsis thaliana T-DNA mutant database. *Nucleic Acids Res* 40, D1211-1215.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Komander, D., and Rape, M. (2012). The ubiquitin code. *Annu Rev Biochem* 81, 203-229.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Lachaud, C., Prigent, E., Thuleau, P., Grat, S., Da Silva, D., Brière, C., Mazars, C., and Cotelle, V. (2013). 14-3-3-Regulated Ca²⁺-dependent protein kinase CPK3 is required for sphingolipid-induced cell death in Arabidopsis. *Cell Death & Differentiation* 20, 209-217.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Le, M.H., Cao, Y., Zhang, X.C., and Stacey, G. (2014). LIK1, a CERK1-interacting kinase, regulates plant immune responses in Arabidopsis. *PLoS One* 9, e102245.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Lee, B.-J., Kwon, S.J., Kim, S.-K., Kim, K.-J., Park, C.-J., Kim, Y.-J., Park, O.K., and Paek, K.-H. (2006). Functional study of hot pepper 26S proteasome subunit RPN7 induced by Tobacco mosaic virus from nuclear proteome analysis. *Biochemical and biophysical research communications* 351, 405-411.
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Li, L., Habring, A., Wang, K., and Weigel, D. (2020). Atypical Resistance Protein RPW8/HR Triggers Oligomerization of the NLR Immune Receptor RPP7 and Autoimmunity. *Cell Host Microbe* 27, 405-417 e406.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li, Y., Kabbage, M., Liu, W., and Dickman, M.B. (2016). Aspartyl protease-mediated cleavage of BAG6 is necessary for autophagy and fungal resistance in plants. *The Plant Cell* 28, 233-247.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liang, X., Ding, P., Lian, K., Wang, J., Ma, M., Li, L., Li, L., Li, M., Zhang, X., Chen, S., et al. (2016). Arabidopsis heterotrimeric G proteins regulate immunity by directly coupling to the FLS2 receptor. *Elife* 5, e13568.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liao, D., Cao, Y., Sun, X., Espinoza, C., Nguyen, C.T., Liang, Y., and Stacey, G. (2017). Arabidopsis E3 ubiquitin ligase PLANT U-BOX13 (PUB13) regulates chitin receptor LYSIN MOTIF RECEPTOR KINASE5 (LYK5) protein abundance. *The New phytologist* 214, 1646-1656.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liebrand, T.W., van den Burg, H.A., and Joosten, M.H. (2014). Two for all: receptor-associated kinases SOBIR1 and BAK1. *Trends in plant science* 19, 123-132.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu, J., Ding, P., Sun, T., Nitta, Y., Dong, O., Huang, X., Yang, W., Li, X., Botella, J.R., and Zhang, Y. (2013). Heterotrimeric G proteins serve as a converging point in plant defense signaling activated by multiple receptor-like kinases. *Plant physiology* 161, 2146-2158.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu, J., Li, H., Miao, M., Tang, X., Giovannoni, J., Xiao, F., and Liu, Y. (2012a). The tomato UV-damaged DNA-binding protein-1 (DDB1) is implicated in pathogenesis-related (PR) gene expression and resistance to *Agrobacterium tumefaciens*. *Molecular plant pathology* 13, 123-134.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu, J., Li, H., Miao, M., Tang, X., Giovannoni, J., Xiao, F., and Liu, Y.J.M.p.p. (2012b). The tomato UV-damaged DNA-binding protein-1 (DDB1) is implicated in pathogenesis-related (PR) gene expression and resistance to *Agrobacterium tumefaciens*. 13, 123-134.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lu, D., Lin, W., Gao, X., Wu, S., Cheng, C., Avila, J., Heese, A., Devarenne, T.P., He, P., and Shan, L. (2011). Direct ubiquitination of pattern recognition receptor FLS2 attenuates plant innate immunity. *Science* 332, 1439-1442.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ma, X., Claus, L.A.N., Leslie, M.E., Tao, K., Wu, Z., Liu, J., Yu, X., Li, B., Zhou, J., Savatin, D.V., et al. (2020). Ligand-induced monoubiquitination of BIK1 regulates plant immunity. *Nature* 581, 199-203.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Maor, R., Jones, A., Nuhse, T.S., Studholme, D.J., Peck, S.C., and Shirasu, K. (2007). Multidimensional protein identification technology (MudPIT) analysis of ubiquitinated proteins in plants. *Mol Cell Proteomics* 6, 601-610.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Marino, D., Peeters, N., and Rivas, S. (2012). Ubiquitination during plant immune signaling. *Plant physiology* 160, 15-27.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Marshall, R.S., Li, F., Gemperline, D.C., Book, A.J., and Vierstra, R.D. (2015). Autophagic Degradation of the 26S Proteasome Is Mediated by the Dual ATG8/Ubiquitin Receptor RPN10 in Arabidopsis. *Molecular cell* 58, 1053-1066.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Marshall, R.S., and Vierstra, R.D. (2018). Autophagy: The Master of Bulk and Selective Recycling. *Annual review of plant biology* 69, 173-208.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Meng, X., and Zhang, S. (2013). MAPK cascades in plant disease resistance signaling. *Annual review of phytopathology* 51, 245-266.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Meyer, D., Pajonk, S., Micali, C., O'Connell, R., and Schulze-Lefert, P.J.T.P.J. (2009). Extracellular transport and integration of plant secretory proteins into pathogen-induced cell wall compartments. 57, 986-999.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Miao, Y., and Zentgraf, U. (2010). AHECT E3 ubiquitin ligase negatively regulates Arabidopsis leaf senescence through degradation of the transcription factor WRKY53. *The Plant Journal* 63, 179-188.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Monaghan, J., and Li, X. (2010). The HEAT repeat protein ILITYHIA is required for plant immunity. *Plant and cell physiology* 51, 742-753.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Monaghan, J., Li, X.J.P., and physiology, c. (2010). The HEAT repeat protein ILITYHIA is required for plant immunity. 51, 742-753.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Monaghan, J., Xu, F., Gao, M., Zhao, Q., Palma, K., Long, C., Chen, S., Zhang, Y., and Li, X. (2009). Two Prp19-like U-box proteins in the MOS4-associated complex play redundant roles in plant innate immunity. *PLoS Pathogens* 5.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mondragon-Palomino, M., Stam, R., John-Arputharaj, A., and Dresselhaus, T. (2017). Diversification of defensins and NLRs in Arabidopsis species by different evolutionary mechanisms. BMC Evol Biol 17, 255.

Google Scholar: [Author Only Title Only Author and Title](#)

Mural, R.V., Liu, Y., Rosebrock, T.R., Brady, J.J., Hamera, S., Connor, R.A., Martin, G.B., and Zeng, L. (2013). The tomato Fni3 lysine-63-specific ubiquitin-conjugating enzyme and suv ubiquitin E2 variant positively regulate plant immunity. Plant Cell 25, 3615-3631.

Google Scholar: [Author Only Title Only Author and Title](#)

Murata, S., Yashiroda, H., and Tanaka, K. (2009). Molecular mechanisms of proteasome assembly. Nature reviews Molecular cell biology 10, 104-115.

Google Scholar: [Author Only Title Only Author and Title](#)

Nakashima, A., Chen, L., Thao, N.P., Fujiwara, M., Wong, H.L., Kuwano, M., Umemura, K., Shirasu, K., Kawasaki, T., and Shimamoto, K. (2008). RACK1 functions in rice innate immunity by interacting with the Rac1 immune complex. Plant Cell 20, 2265-2279.

Google Scholar: [Author Only Title Only Author and Title](#)

Nicaise, V., Joe, A., Jeong, B., Korneli, C., Boutrot, F., Wested, I., Staiger, D., Alfano, J., and Zipfel, C. (2013). Pseudomonas HopU1 affects interaction of plant immune receptor mRNAs to the RNA-binding protein GRP7. EMBO J 32, 701-712.

Google Scholar: [Author Only Title Only Author and Title](#)

Okumoto, K., Misono, S., Miyata, N., Matsumoto, Y., Mukai, S., and Fujiki, Y. (2011). Cysteine ubiquitination of PTS1 receptor Pex5p regulates Pex5p recycling. Traffic 12, 1067-1083.

Google Scholar: [Author Only Title Only Author and Title](#)

Ordureau, A., Paulo, J.A., Zhang, J., An, H., Swatek, K.N., Cannon, J.R., Wan, Q., Komander, D., and Harper, J.W. (2020). Global Landscape and Dynamics of Parkin and USP30-Dependent Ubiquitylomes in iNeurons during Mitophagic Signaling. Molecular cell 77, 1124-1142 e1110.

Google Scholar: [Author Only Title Only Author and Title](#)

Paez Valencia, J., Goodman, K., and Otegui, M.S. (2016). Endocytosis and Endosomal Trafficking in Plants. Annual review of plant biology 67, 309-335.

Google Scholar: [Author Only Title Only Author and Title](#)

Palm, D., Streit, D., Shanmugam, T., Weis, B.L., Ruprecht, M., Simm, S., and Schleiff, E. (2019). Plant-specific ribosome biogenesis factors in Arabidopsis thaliana with essential function in rRNA processing. Nucleic Acids Res 47, 1880-1895.

Google Scholar: [Author Only Title Only Author and Title](#)

Parker, J.E., Holub, E.B., Frost, L.N., Falk, A., Gunn, N.D., and Daniels, M.J. (1996). Characterization of eds1, a mutation in Arabidopsis suppressing resistance to Peronospora parasitica specified by several different RPP genes. The Plant Cell 8, 2033-2046.

Google Scholar: [Author Only Title Only Author and Title](#)

Pastorczyk, M., and Bednarek, P. (2016). The function of glucosinolates and related metabolites in plant innate immunity. In Advances in Botanical Research (Elsevier), pp. 171-198.

Google Scholar: [Author Only Title Only Author and Title](#)

Pecher, P., Eschen-Lippold, L., Herklotz, S., Kuhle, K., Naumann, K., Bethke, G., Uhrig, J., Weyhe, M., Scheel, D., and Lee, J. (2014). The Arabidopsis thaliana mitogen-activated protein kinases MPK 3 and MPK 6 target a subclass of 'VQ-motif'-containing proteins to regulate immune responses. New Phytologist 203, 592-606.

Google Scholar: [Author Only Title Only Author and Title](#)

Peng, J., Schwartz, D., Elias, J.E., Thoreen, C.C., Cheng, D., Marsischky, G., Roelofs, J., Finley, D., and Gygi, S.P. (2003). A proteomics approach to understanding protein ubiquitination. Nat Biotechnol 21, 921-926.

Google Scholar: [Author Only Title Only Author and Title](#)

Perraki, A., Gronnier, J., Gouguet, P., Boudsocq, M., Deroubaix, A-F., Simon, V., German-Retana, S., Zipfel, C., Bayer, E., and Mongrand, S. (2017). The plant calcium-dependent protein kinase CPK3 phosphorylates REM1.3 to restrict viral infection. BioRxiv, 205765.

Google Scholar: [Author Only Title Only Author and Title](#)

Pitorre, D., Llauro, C., Jobet, E., Guilleminot, J., Brizard, J-P., Delseny, M., and Lasserre, E. (2010). RLK7, a leucine-rich repeat receptor-like kinase, is required for proper germination speed and tolerance to oxidative stress in Arabidopsis thaliana. Planta 232, 1339-1353.

Google Scholar: [Author Only Title Only Author and Title](#)

Qin, L., Zhou, Z., Li, Q., Zhai, C., Liu, L., Quilichini, T.D., Gao, P., Kessler, S.A., Jaillais, Y., Datla, R., et al. (2020). Specific Recruitment of Phosphoinositide Species to the Plant-Pathogen Interfacial Membrane Underlies Arabidopsis Susceptibility to Fungal Infection. Plant Cell 32, 1665-1688.

Google Scholar: [Author Only Title Only Author and Title](#)

Raasi, S., Orlov, I., Flerning, K.G., and Pickart, C.M. (2004). Binding of polyubiquitin chains to ubiquitin-associated (UBA) domains of HHR23A. Journal of molecular biology 341, 1367-1379.

Google Scholar: [Author Only Title Only Author and Title](#)

Rayapuram, N., Bigeard, J., Alhoraibi, H., Bonhomme, L., Hesse, A.M., Vinh, J., Hirt, H., and Pflieger, D. (2018). Quantitative Phosphoproteomic Analysis Reveals Shared and Specific Targets of Arabidopsis Mitogen-Activated Protein Kinases (MAPKs) MPK3, MPK4, and MPK6. *Mol Cell Proteomics* 17, 61-80.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Revers, F., Guiraud, T., Houvenaghel, M.-C., Mauduit, T., Le Gall, O., and Candresse, T. (2003). Multiple resistance phenotypes to Lettuce mosaic virus among Arabidopsis thaliana accessions. *Molecular plant-microbe interactions* 16, 608-616.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Reyes, F.C., Buono, R., and Otegui, M.S. (2011). Plant endosomal trafficking pathways. *Current opinion in plant biology* 14, 666-673.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rodrigues, O., Reshetnyak, G., Grondin, A., Saijo, Y., Leonhardt, N., Maurel, C., and Verdoucq, L. (2017). Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA- and pathogen-triggered stomatal closure. *Proceedings of the National Academy of Sciences* 114, 9200-9205.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Romero-Barrios, N., and Vert, G. (2018). Proteasome-independent functions of lysine-63 polyubiquitination in plants. *The New phytologist* 217, 995-1011.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Saracco, S.A., Hansson, M., Scafi, M., Walker, J.M., Smith, L.M., and Vierstra, R.D. (2009). Tandem affinity purification and mass spectrometric analysis of ubiquitylated proteins in Arabidopsis. *Plant J* 59, 344-358.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T.J.G.r. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *13*, 2498-2504.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shimizu, Y., Okuda-Shimizu, Y., and Hendershot, L.M. (2010). Ubiquitylation of an ERAD substrate occurs on multiple types of amino acids. *Molecular cell* 40, 917-926.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Skubacz, A., Daszkowska-Golec, A., and Szarejko, L. (2016). The Role and Regulation of ABI5 (ABA-Insensitive 5) in Plant Development, Abiotic Stress Responses and Phytohormone Crosstalk. *Front Plant Sci* 7.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Smalle, J., and Vierstra, R.D. (2004). The ubiquitin 26S proteasome proteolytic pathway. *Annual review of plant biology* 55, 555-590.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Speth, E.B., Imboden, L., Hauck, P., and He, S.Y. (2009a). Subcellular localization and functional analysis of the Arabidopsis GTPase RabE. *Plant physiology* 149, 1824-1837.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Speth, E.B., Imboden, L., Hauck, P., and He, S.Y. (2009b). Subcellular Localization and Functional Analysis of the Arabidopsis GTPase RabE. *149*, 1824-1837.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Spoel, S.H., and Dong, X. (2012). How do plants achieve immunity? Defence without specialized immune cells. *Nature reviews Immunology* 12, 89-100.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stegmann, M., Anderson, R.G., Ichimura, K., Pecenkova, T., Reuter, P., Zarsky, V., McDowell, J.M., Shirasu, K., and Trujillo, M. (2012). The ubiquitin ligase PUB22 targets a subunit of the exocyst complex required for PAMP-triggered responses in Arabidopsis. *Plant Cell* 24, 4703-4716.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stringlis, I.A., Yu, K., Feussner, K., de Jonge, R., Van Bentum, S., Van Verk, M.C., Berendsen, R.L., Bakker, P.A., Feussner, I., and Pieterse, C.M. (2018). MYB72-dependent coumarin exudation shapes root microbiome assembly to promote plant health. *Proceedings of the National Academy of Sciences* 115, E5213-E5222.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Su, J., Xu, J., and Zhang, S. (2015). RACK1, scaffolding a heterotrimeric G protein and a MAPK cascade. *Trends Plant Sci* 20, 405-407.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Su, J., Yang, L., Zhu, Q., Wu, H., He, Y., Liu, Y., Xu, J., Jiang, D., and Zhang, S. (2018). Active photosynthetic inhibition mediated by MPK3/MPK6 is critical to effector-triggered immunity. *PLoS biology* 16, e2004122.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J., Simonovic, M., Roth, A., Santos, A., Tsafou, K.P., et al. (2015). STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 43, D447-452.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tan, X., Meyers, B.C., Kozik, A., West, M.A., Morgante, M., St Clair, D.A., Bent, A.F., and Michelmore, R.W. (2007). Global expression

analysis of nucleotide binding site-leucine rich repeat-encoding and related genes in Arabidopsis. BMC Plant Biol 7, 56.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tarutani, Y., Morimoto, T., Sasaki, A., Yasuda, M., Nakashita, H., Yoshida, S., Yamaguchi, I., and Suzuki, Y. (2004). Molecular characterization of two highly homologous receptor-like kinase genes, RLK902 and RKL1, in Arabidopsis thaliana. Biosci Biotechnol Biochem 68, 1935-1941.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Thordal-Christensen, H. (2003). Fresh insights into processes of nonhost resistance. Curr Opin Plant Biol 6, 351-357.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tronchet, M., Balague, C., Kroj, T., Jouanin, L., and Roby, D. (2010). Cinnamyl alcohol dehydrogenases-C and D, key enzymes in lignin biosynthesis, play an essential role in disease resistance in Arabidopsis. Molecular plant pathology 11, 83-92.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Trujillo, M., and Shirasu, K. (2010). Ubiquitination in plant immunity. Current opinion in plant biology 13, 402-408.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Udeshi, N.D., Svinkina, T., Mertins, P., Kuhn, E., Mani, D.R., Qiao, J.W., and Carr, S.A. (2013). Refined Preparation and Use of Anti-diglycine Remnant (K-epsilon-GG) Antibody Enables Routine Quantification of 10,000s of Ubiquitination Sites in Single Proteomics Experiments. Molecular & Cellular Proteomics 12, 825-831.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Urano, D., Chen, J.G., Botella, J.R., and Jones, A.M. (2013). Heterotrimeric G protein signalling in the plant kingdom. Open Biol 3, 120186.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Üstün, S., Sheikh, A., Gimenez-Ibanez, S., Jones, A., Ntoukakis, V., and Börnke, F. (2016). The proteasome acts as a hub for plant immunity and is targeted by Pseudomonas type III effectors. Plant Physiology 172, 1941-1958.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

van Nocker, S., Walker, J.M., and Vierstra, R.D. (1996). The Arabidopsis thaliana UBC7/13/14 genes encode a family of multiubiquitin chain-forming E2 enzymes. The Journal of biological chemistry 271, 12150-12158.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vierstra, R.D. (2009). The ubiquitin-26S proteasome system at the nexus of plant biology. Nat Rev Mol Cell Biol 10, 385-397.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang, J., Grubb, L.E., Wang, J., Liang, X., Li, L., Gao, C., Ma, M., Feng, F., Li, M., Li, L., et al. (2018). A Regulatory Module Controlling Homeostasis of a Plant Immune Kinase. Molecular cell.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang, L., Wen, R., Wang, J., Xiang, D., Wang, Q., Zang, Y., Wang, Z., Huang, S., Li, X., and Datla, R.J.N.P. (2019). Arabidopsis UBC 13 differentially regulates two programmed cell death pathways in responses to pathogen and low-temperature stress. 221, 919-934.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wawrzynska, A., Rodibaugh, N.L., and Innes, R.W. (2010). Synergistic activation of defense responses in Arabidopsis by simultaneous loss of the GSL5 callose synthase and the EDR1 protein kinase. Mol Plant Microbe Interact 23, 578-584.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wen, R., Torres-Acosta, J.A., Pastushok, L., Lai, X., Pelzer, L., Wang, H., and Xiao, W. (2008). Arabidopsis UEV1D promotes Lysine-63-linked polyubiquitination and is involved in DNA damage response. The Plant cell 20, 213-227.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yao, C., Wu, Y., Nie, H., and Tang, D. (2012). RPN1a, a 26S proteasome subunit, is required for innate immunity in Arabidopsis. The Plant journal : for cell and molecular biology 71, 1015-1028.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yu, X., Feng, B., He, P., and Shan, L. (2017). From Chaos to Harmony: Responses and Signaling upon Microbial Pattern Recognition. Annual review of phytopathology 55, 109-137.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zamioudis, C., Hanson, J., and Pieterse, C.M. (2014). β -Glucosidase BGLU 42 is a MYB 72-dependent key regulator of rhizobacteria-induced systemic resistance and modulates iron deficiency responses in Arabidopsis roots. New Phytologist 204, 368-379.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhao, J., Zhou, H., Zhang, M., Gao, Y., Li, L., Gao, Y., Li, M., Yang, Y., Guo, Y., and Li, X. (2016). Ubiquitin-specific protease 24 negatively regulates abscisic acid signalling in Arabidopsis thaliana. Plant, cell & environment 39, 427-440.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhao, Y., Wu, G., Shi, H., and Tang, D. (2019). RECEPTOR-LIKE KINASE 902 Associates with and Phosphorylates BRASSINOSTEROID-SIGNALING KINASE1 to Regulate Plant Immunity. Mol Plant 12, 59-70.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhou, B., and Zeng, L. (2017). Conventional and unconventional ubiquitination in plant immunity. Molecular plant pathology 18, 1313-1330.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhou, J.-M., and Zhang, Y. (2020). Plant Immunity: Danger Perception and Signaling. Cell.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhou, J., He, P., and Shan, L. (2014). Ubiquitination of plant immune receptors. Methods Mol Biol 1209, 219-231.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhou, J., Liu, D., Wang, P., Ma, X., Lin, W., Chen, S., Mishev, K., Lu, D., Kumar, R., Vanhoutte, I., et al. (2018). Regulation of Arabidopsis brassinosteroid receptor BRI1 endocytosis and degradation by plant U-box PUB12/PUB13-mediated ubiquitination. Proc Natl Acad Sci U S A 115, E1906-E1915.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhou, J., Lu, D., Xu, G., Finlayson, S.A., He, P., and Shan, L. (2015). The dominant negative ARM domain uncovers multiple functions of PUB13 in Arabidopsis immunity, flowering, and senescence. J Exp Bot 66, 3353-3366.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)