1 ***Common variants contribute to intrinsic human brain functional networks***

2

3 **Running title: GWAS of intrinsic brain function**

4

5 Bingxin Zhao^{1,14}, Tengfei Li^{2,3,14}, Stephen M. Smith⁴, Di Xiong¹, Xifeng Wang¹, Yue Yang¹,
6 Tianyou Luo¹, Ziliang Zhu¹, Yue Shan¹, Nana Matoba^{5,6}, Quan Sun¹, Yuchen Yang⁵, Mads
7 E. Hauberg^{7,8,10,11}, Jaroslav Bendl⁷⁻⁹, John F. Fullard⁷⁻⁹, Panagiotis Roussos^{7-10,12}, Weili
8 Lin^{2,3}, Yun Li^{1,5,13}, Jason L. Stein^{5,6}, and Hongtu Zhu^{1,3*}

9

10 ¹Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

11 ²Department of Radiology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

12 ³Biomedical Research Imaging Center, School of Medicine, University of North Carolina at Chapel Hill,
13 Chapel Hill, NC, USA

14 ⁴Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical
15 Neurosciences, University of Oxford, Oxford, UK

16 ⁵Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

17 ⁶UNC Neuroscience Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

18 ⁷Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA

19 ⁸Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA

20 ⁹Department of Genetics and Genomic Science and Institute for Multiscale Biology, Icahn School of
21 Medicine at Mount Sinai, New York, NY, USA

22 ¹⁰iPSYCH, The Lundbeck Foundation Initiative for Integrative Psychiatric Research, Denmark

23 ¹¹Centre for Integrative Sequencing (iSEQ), Aarhus University, Aarhus, Denmark

24 ¹²Mental Illness Research, Education, and Clinical Center (VISN 2 South), James J. Peters VA Medical
25 Center, Bronx, NY, USA

26 ¹³Department of Computer Science, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

27 ¹⁴These authors contributed equally to this work.

28

29 ****Corresponding author:***

30 Hongtu Zhu

31 3105C McGavran-Greenberg Hall, 135 Dauer Drive, Chapel Hill, NC 27599.

32 E-mail address: htzhu@email.unc.edu Phone: (919) 966-7250

1 Abstract

2 The human brain remains active in the absence of explicit tasks and forms networks of
 3 correlated activity. Resting-state functional magnetic resonance imaging (rsfMRI)
 4 measures brain activity at rest, which has been linked with both cognitive and clinical
 5 outcomes. The genetic variants influencing human brain function are largely unknown.
 6 Here we utilized rsfMRI from 44,190 individuals of multiple ancestries (37,339 in the UK
 7 Biobank) to discover and validate the common genetic variants influencing intrinsic
 8 brain activity. We identified hundreds of novel genetic loci associated with intrinsic
 9 functional signatures ($P < 2.8 \times 10^{-11}$), including associations to the central executive,
 10 default mode, and salience networks involved in the triple network model of
 11 psychopathology. A number of intrinsic brain activity associated loci colocalized with
 12 brain disorder GWAS (e.g., Alzheimer's disease, Parkinson's disease, schizophrenia) and
 13 cognition, such as 19q13.32, 17q21.31, and 2p16.1. Particularly, we detected a
 14 colocalization between one (rs429358) of the two variants in the *APOE* $\epsilon 4$ locus and
 15 function of the default mode, central executive, attention, and visual networks. Genetic
 16 correlation analysis demonstrated shared genetic influences between brain function and
 17 brain structure in the same regions. We also detected significant genetic correlations
 18 with 26 other complex traits, such as ADHD, major depressive disorder, schizophrenia,
 19 intelligence, education, sleep, subjective well-being, and neuroticism. Common variants
 20 associated with intrinsic brain activity were enriched within regulatory element in brain
 21 tissues.

22
 23 **Keywords:** Amplitude; Functional connectivity; Intrinsic brain activity; GWAS;
 24 Resting-state fMRI; Triple network model; UK Biobank.

The human brain is a complex system where functional organization and communication between brain networks are necessary for behavior and cognition¹⁻⁴. The human brain remains active in the absence of explicit tasks or stimuli, resulting in an intrinsic functional architecture. Utilizing changes in blood oxygen level-dependent (BOLD) signal^{5,6}, resting-state functional magnetic resonance imaging⁷ (rsfMRI) captures spontaneous intrinsic brain activity⁸. Specifically, the spontaneous neural activity and non-neural physiological processes within each functional region are quantified by the amplitude of low frequency fluctuations (ALFF) in BOLD time series^{6,9,10}. Moreover, the inter-regional correlations in spontaneous neuronal variability are used to construct a functional connectivity matrix, which measures the magnitude of temporal synchrony between each pair of brain regions^{6,11}.

rsfMRI has led to the discovery of multiple resting-state networks (RSNs) present in neurotypical human brains, including the default mode, central executive (i.e., frontoparietal), attention, limbic, salience, somatomotor, and visual networks¹²⁻¹⁴. Among these RSNs, the central executive, default mode, and salience networks are three core neurocognitive networks that support efficient cognition¹⁵⁻¹⁷. Accumulating evidence suggests that the functional organization and dynamic interaction of these three networks underlie a wide range of mental disorders, resulting in the triple network model of psychopathology^{16,18}. Supporting this model, differences in RSNs have been detected in multiple neurological and psychiatric disorders¹⁹ relative to neurotypical controls, such as Alzheimer's disease²⁰, Parkinson's disease²¹, and major depressive disorder (MDD)²².

Twin and family studies have largely reported a low to moderate degree of genetic contributions to intrinsic brain activity²³⁻²⁹. For example, the family-based heritability estimates of major RSNs ranged from 20% to 40% in the Human Connectome Project (HCP)³⁰. In a previous study using about 8,000 UK Biobank (UKB) individuals³¹, the SNP heritability³² of amplitude and functional connectivity traits can be higher than 30%. Although there were multiple candidate gene studies for intrinsic brain activity (such as for *APOE*³³ and *KIBRA*³⁴), currently only one genome-wide association study (GWAS)³¹ has been successfully performed on rsfMRI²³ ($n \approx 8000$). This is likely due to both

insufficient sample size for GWAS discovery and weaker genetic effects on brain function than structure^{31,35-39}. It is also known that functional connectivity traits in rsfMRI are typically noisier than brain structural traits measured in other neuroimaging modalities. In addition, imaging batch effects⁴⁰ (e.g., image acquisition, processing procedures, and software) may cause additional technical variability in rsfMRI analyses⁴¹, making GWAS meta-analysis and independent replication particularly challenging. Therefore, genetic variants influencing intrinsic brain activity have remained largely undiscovered and their shared genetic influences with other complex traits and clinical outcomes are unknown.

To address these challenges, here we collected individual-level rsfMRI data from four independent studies, including the UK Biobank⁴², Adolescent Brain Cognitive Development (ABCD⁴³), Philadelphia Neurodevelopmental Cohort (PNC⁴⁴), and HCP⁴⁵. We harmonized rsfMRI processing procedures by following the unified UKB brain imaging pipeline^{9,46}. Functional brain regions and corresponding functional connectivity were characterized via spatial Independent Component Analysis (ICA)^{47,48} for 44,190 individuals from multiple ancestries, including 37,339 from UK Biobank. As in previous studies^{9,31,49}, two parcellations with different dimensionalities^{13,50} (25 and 100 regions, respectively) were separately applied in spatial ICA and we focused on the 76 (21 and 55, respectively) regions that had been previously confirmed to be non-artifactual⁹. Two group of neuroimaging phenotypes were then generated: the first group contains 76 (node) amplitude traits reflecting the regional spontaneous neuronal activity; and the second group includes 1,695 (i.e., $21 \times 20/2 + 55 \times 54/2$) (edge) functional connectivity traits that quantify the inter-regional co-activity, as well as 6 global functional connectivity measures summarizing all of the 1,695 pairwise functional connectivity traits³¹. These 1,777 traits were then used to explore the genetic architecture of intrinsic brain activity. To aid interpretation of GWAS results, the functional brain regions characterized in ICA were labelled by using the automated anatomical labeling (AAL) atlas⁵¹ and were mapped onto major functional networks defined in Yeo, et al.¹⁴ and Finn, et al.¹². Our GWAS results can be easily explored and downloaded through the Brain Imaging Genetics Knowledge Portal (BIG-KP) <https://bigkp.org>.

1 RESULTS

2 Genetics of the intrinsic brain functional architecture.

3 SNP heritability was estimated for the 1,777 intrinsic brain activity traits via GCTA⁵². The
4 mean heritability (h^2) estimate was 27.2% (range = (10%, 36.5%), standard error = 6.0%)
5 for the 76 amplitude traits, all of which remained significant after adjusting for multiple
6 comparisons by using the Benjamini-Hochberg procedure to control false discovery rate
7 (FDR) at 0.05 level (1,777 tests, **Fig. 1a** and **Supplementary Table 1**). Among the 1,701
8 functional connectivity traits, 1,230 had significant (again at 5% FDR) heritability with
9 estimates varying from 3% to 61% (mean = 9.6%, standard error = 5.8%). Ten functional
10 connectivity traits had heritability higher than 30%, including 4 global functional
11 connectivity measures (**Supplementary Fig. 1**) and 6 pairwise functional connectivity
12 traits (**Fig. 1b**). These most heritable traits were most related to the central executive,
13 default mode, and salience networks in the triple network model of psychopathology¹⁶.
14 To examine whether intrinsic brain activity within the triple network in general had
15 higher heritability, we classified the 76 amplitude traits into two categories 1) fully or
16 partially within the triple network and 2) outside the triple network. Correspondingly,
17 the 1,695 pairwise functional connectivity traits were classified into 1) within the triple
18 network, 2) outside the triple network, and 3) between the triple and non-triple
19 networks. We found that amplitude traits within the triple network had significantly
20 higher heritability than those outside the triple network (mean = 30.5% vs. 22.3%, $P =$
21 6.3×10^{-11} , two-sided Wilcoxon rank test) (**Fig. 1c**). Similarly, functional connectivity
22 traits within the triple network had higher heritability than interactions outside the
23 triple network or between the triple and non-triple networks (mean = 12.5% vs. 7%, $P =$
24 1.9×10^{-26}). These results indicate that the level of genetic control might be higher in
25 core neurocognitive networks. The range of heritability estimates was consistent with
26 previous results³¹, suggesting that common genetic variants had a low to moderate
27 degree of contributions to inter-individual variability of intrinsic brain activity. The
28 overall genetic effects on both amplitude and functional connectivity were lower than
29 those on brain structure. For example, the average heritability was reported to be 48.7%
30 for diffusion tensor imaging (DTI) traits of brain structural connectivity in white matter
31 tracts⁵³ and 40% for regional brain volumes measuring brain morphometry³⁷.
32 Nevertheless, as shown below, intrinsic brain activity may be more functionally relevant

with stronger genetic connections to brain disorders than brain structure, such as Alzheimer's disease.

Genome-wide association discovery was carried out for 1,777 intrinsic brain activity traits using UKB individuals of British ancestry ($n = 34,691$, Methods). The Manhattan and QQ plots can be found in the BIG-KP server. At the significance level 2.8×10^{-11} ($5 \times 10^{-8}/1,777$, i.e., the standard GWAS threshold, Bonferroni-adjusted for the 1,777 traits), FUMA⁵⁴ identified 264 lead independent variants (linkage disequilibrium [LD] $r^2 < 0.1$), and then characterized 606 significant locus-trait associations for 197 traits (75 amplitude and 122 functional connectivity (**Supplementary Tables 2-3, Supplementary Fig. 2**, Methods). The amplitude traits typically had multiple associated variants and a number of variants were widely related to the amplitude in different brain regions, such as rs429358 (nearest gene *APOE*), rs2274224 (*PLCE1*), and rs1133400 (*INPP5A*). In addition, rs2279829 (*ZIC4*), rs62158211 (*AC016745.1*), and rs115877304 (*NR2F1-AS1*) were associated with multiple functional connectivity traits. Global and pairwise functional connectivity traits that had at least 5 significant variants were again most related to the central executive, default mode, and salience networks (**Supplementary Fig. 3**). Of the 14 associated variants that had been identified in the previous GWAS³¹, 12 were in LD ($r^2 \geq 0.6$) with our significant variants, most of which were associated with amplitude traits. In summary, our analyses identify many novel variants associated with intrinsic functional signatures and illustrate the global genetic influences on functional connectivity across the whole brain. The degree of genetic control is higher in the central executive, default mode, and salience networks, whose cross-network interactions closely control multiple cognitive functions and affect major brain disorders¹⁸.

Replication and the effect of ancestry.

We aimed to replicate our results in UKB British GWAS using other independent datasets. First, we repeated GWAS on UKB individuals of White but Non-British ancestry (UKBW, $n = 1,970$) and three non-UKB European-ancestry cohorts, including ABCD European (ABCDE, $n = 3,821$), HCP ($n = 495$), and PNC ($n = 510$). We meta-analyzed the four European GWAS (total $n = 6,796$) and checked whether the locus-trait associations

detected in UKB British GWAS can be replicated. For the 606 significant associations, 101 (16.7%) passed the 8.2×10^{-5} (i.e., 0.05/606) Bonferroni significance level in this validation GWAS, and 599 (98.8%) were significant at FDR 5% level. Next, we performed GWAS on four non-European validation datasets: the UKB Asian (UKBA, $n = 446$), UKB Black (UKBBL, $n = 232$), ABCD Hispanic (ABCDH, $n = 768$), and ABCD African American (ABCDAA, $n = 1,257$). We meta-analyzed these four non-European GWAS (total $n = 2,703$) and found that 39 (6.4%) passed the Bonferroni significance level and 601 (99.2%) were significant at FDR 5% level. Some associations with rs3781658 (*ANO1*), rs7083220 (*PWWP2B*), rs9373978 (*FHL5*), rs11187838 (*PLCE1*), and rs35124509 (*EPHA3*) were replicated in both European and non-European datasets at the stringent Bonferroni significance level. Moreover, we performed a third meta-analysis to combine all of the eight validation datasets, after which the number of replicated associations moved up to 136 (22.4%) and 602 (99.3%) at Bonferroni and FDR significance levels, respectively. These results are summarized in **Supplementary Table 4**. Overall, our results suggest that the associated genetic loci discovered in UKB British GWAS have high generalizability in independent rsfMRI studies, despite the fact that these studies may use different imaging protocols/MRI scanners and recruit participants from different age groups. The strong homogeneity of GWAS results likely benefit, in part, from the consistent rsfMRI processing procedures that we applied to these datasets.

In addition, we utilized polygenic risk scores⁵⁵ (PRS) derived from UKB British GWAS for further evidence of replication (Methods). For the 197 traits that had significant variants, 168 had significant PRS in at least one of the four European validation GWAS datasets at FDR 5% level (197×4 tests, **Supplementary Table 5**), illustrating the significant out-of-sample prediction power of polygenic influences from our discovery GWAS results. The largest incremental R-squared (after adjusting the effects of age, sex, and ten genetic principal components) were observed on the 2nd, 3rd, 4th, and 6th global functional connectivity measures in UKBW and HCP datasets, which were larger than 5% (range = (5.1%, 5.7%), P range = (1.1×10^{-24} , 4×10^{-13})). To evaluate the consistency across ancestry, PRS was also constructed on the four non-European validation datasets. UKBA had the best validation performance among the four datasets, with 86 PRS being significant at FDR 5% level (197×4 tests, **Supplementary Table 6**).

1 The number of significant PRS was reduced to 59, 39, and 31 in ABCDH, ABCDA, and
2 UKBBL, respectively. In summary, these PRS results illustrate the overall consistency of
3 genetic effects in European cohorts and also show that there may be population specific
4 influences on brain function in other cohorts, though much smaller sample sizes and
5 difficulty in conducting cross ancestry PRS strongly limit the interpretability of these
6 analyses. More efforts are required to identify causal variants associated with functional
7 brain in global diverse populations and perform better cross-population PRS predictions.

8 9 **The shared genetic loci with brain-related complex traits and disorders.**

10 To evaluate the shared genetic influences between intrinsic brain activity and other
11 complex traits, we carried out association lookups for independent significant variants
12 (and their LD tags, i.e., variants with LD, $r^2 \geq 0.6$) detected in UKB British GWAS
13 (Methods). In the NHGRI-EBI GWAS catalog⁵⁶, our results tagged many variants reported
14 for a wide range of complex traits in different trait domains, such as neurological and
15 psychiatric disorders, cognitive performance, education, bone mineral density, sleep,
16 smoking/drinking, brain structure, and anthropometric traits. Below we highlighted
17 colocalizations in a few selected genomic regions.

18
19 The index variants rs429358 (*APOE*), rs34404554 (*TOMM40*), rs157582(*TOMM40*), and
20 rs157592 (*APOC1*) in the 19q13.32 region (**Fig. 2a, Supplementary Fig. 4**) had genetic
21 effects on the amplitude of many functional brain regions that were most in the default
22 mode, central executive (i.e., frontoparietal), attention, and visual networks. It is well
23 known that 19q13.32 is a risk locus of Alzheimer's disease and rs429358 is one of the
24 two variants in the *APOE* $\epsilon 4$ locus. In this region, we tagged variants associated with
25 dementia and decline in mental ability, including Alzheimer's disease⁵⁷⁻⁵⁹,
26 frontotemporal dementia⁶⁰, cerebral amyloid angiopathy⁶¹, cognitive decline⁶²,
27 cognitive impairment test score⁶³, as well as many biomarkers of Alzheimer's disease,
28 such as neurofibrillary tangles⁶¹, neuritic plaque⁶¹, cerebral amyloid deposition⁶⁴,
29 cerebrospinal fluid protein levels⁶³, and cortical amyloid beta load⁶⁵. Altered amplitude
30 activity has been widely reported in patients of cognitive impairment and Alzheimer's
31 disease^{66,67}. The brain degeneration related to Alzheimer's disease may begin in the
32 frontoparietal regions⁶⁸ and was associated with dysfunction of multiple RSNs,

1 especially the default mode network²⁰. Our findings suggest the shared genetic
2 influences between intrinsic neuronal activity and brain atrophy of Alzheimer's disease.

3

4 Next, the variant rs62061845 (*KANSL1*) in the 17q21.31 region (**Supplementary Fig. 5**)
5 was associated with functional connectivity over the inferior frontal, middle frontal,
6 superior frontal, middle temporal, and supplementary motor area regions in the default
7 mode and salience networks. Variants in LD with rs62061845 have been frequently
8 reported to be associated with Parkinson's disease studies⁶⁹⁻⁷³. As a system-level
9 progressive neurodegenerative disorder⁷⁴, Parkinson's disease not only leads to motor
10 abnormalities, but also has non-motor symptoms such as temporal perception
11 abnormalities⁷⁵ and impaired connectivity among frontal regions⁷⁶. Cognitive
12 dysfunction and disrupted coupling between default mode and salience networks were
13 commonly reported in Parkinson's disease¹⁷. In addition to Parkinson's disease, the
14 17q21.31 region was widely related to other complex traits, including neurological
15 disorders (e.g., Alzheimer's disease⁷⁷, corticobasal degeneration⁷⁸, progressive
16 supranuclear palsy⁷⁹), psychiatric disorders (e.g., autism spectrum disorder⁸⁰, depressive
17 symptoms⁸¹), educational attainment⁸², psychological traits (e.g., neuroticism⁸¹),
18 cognitive traits (cognitive ability⁸³), sleep⁸⁴, heel bone mineral density⁸⁵, alcohol use
19 disorder⁸⁶, subcortical brain volumes³⁸, cortical surface area and thickness³⁶, and white
20 matter microstructure⁵³.

21

22 In addition, the 2p16.1 (**Fig. 2b, Supplementary Fig. 6**) and 5q15 (**Supplementary Fig. 7**)
23 regions were mainly associated with interactions among the central executive, default
24 mode, and salience networks. We observed colocalizations with psychiatric disorders
25 (e.g., schizophrenia⁸⁷, MDD⁸⁸, depressive symptoms⁸⁹, autism spectrum disorder⁹⁰),
26 psychological traits (e.g., neuroticism⁸¹, well-being spectrum⁹¹), sleep⁹², cognitive traits
27 (e.g., intelligence⁹³), and educational attainment⁸². Dysregulated triple network
28 interactions were frequently reported in patients of schizophrenia⁹⁴, depression⁹⁵, and
29 autism spectrum disorder⁹⁶. Similarly, the 2q24.2 (**Supplementary Fig. 8**) and 10q26.13
30 (**Supplementary Fig. 9**) regions had genetic effects on functional connectivity traits
31 involved in the central executive, default mode, salience, and limbic networks. In these
32 two regions, our identified variants tagged those that have been implicated with

schizophrenia⁹⁷, educational attainment⁸², cognitive traits (e.g., cognitive ability⁸³), smoking/drinking (e.g., smoking status⁹⁸, alcohol consumption⁹⁹), hippocampus subfield volumes¹⁰⁰, and heel bone mineral density⁸⁵. We also observed colocalizations in some other genomic regions, such as in 2q14.1 region (**Fig. 2c, Supplementary Fig. 10**) with sleep traits (e.g., sleep duration⁸⁴, insomnia⁹²), in 3p11.1 (**Supplementary Fig. 11**) with cognitive traits (e.g., intelligence¹⁰¹, math ability⁸²), and in 5q14.3 (**Supplementary Fig. 12**) with cognitive traits⁸³ and educational attainment⁸². All of these results are summarized in **Supplementary Table 7**. In summary, intrinsic brain function has wide genetic links to a large number of brain-related complex traits and clinical outcomes, especially neurological and psychiatric disorders and cognitive traits. Integration of GWAS of brain function with these clinical outcomes may help to explain the underlying brain functional mechanisms leading to risk for these disorders.

Genetic correlations with brain structure, brain disorders, and cognition.

The intricate brain neuroanatomical structure is fundamental in supporting brain function. To explore whether genetically mediated brain structural changes were associated with brain function, we examined pairwise genetic correlations (gc) between 1,777 intrinsic brain activity traits and 315 brain structure traits via LDSC¹⁰² (Methods), including 100 regional brain volumes³⁷ and 215 DTI traits of brain structural connectivity in white matter tracts¹⁰³. There were 151 significant pairs between 94 intrinsic brain functional traits and 73 brain structural traits at FDR 5% level ($315 \times 1,777$ tests, $|gc|$ range = (0.22, 0.61), P range = (1.2×10^{-21} , 1.5×10^{-5}), **Supplementary Table 8**).

We found significant genetic correlations between regional brain volumes and functional connectivity strengths ($|gc|$ range = (0.22, 0.61), P range = (1.2×10^{-21} , 1.2×10^{-5}), **Supplementary Fig. 13**). Most of the observed correlations were related to higher order brain functional networks, particularly the attention, default mode, salience, and central executive networks. For example, the insula has been widely implicated to be associated with multiple functions, including but not limited to emotion, addiction, and cognition through extensive connections to neocortex, the limbic system, and amygdala¹⁰⁴. We observed genetic correlations between insula volumes and the connection strengths of multiple pairs of brain regions ($|gc|$ range = (0.22, 0.27), $P < 1.2$

1 $\times 10^{-5}$, **Figs. 3a-b**), which were largely in the default mode and central executive
2 networks, including the angular and the inferior and superior frontal regions. Similarly,
3 left inferior parietal lobule volume exhibited strong genetic correlations with
4 connectivity strengths over multiple pairs of brain regions that were known to be a part
5 of the default mode, visual, attention, and salience networks ($|gc|$ range = (0.34, 0.49),
6 $P < 9.7 \times 10^{-6}$, **Supplementary Fig. 14a**). Interestingly, however, the above identified
7 genetic correlations appeared to be more specific to the left but not right. The inferior
8 parietal has been implicated to be associated with language function and is connected
9 with the Broca's region via the superior longitudinal fasciculus (SLF)¹⁰⁵⁻¹⁰⁷. Considering
10 language processing is left-lateralized in about 95% of right-handers and 75% of
11 left-handers¹⁰⁸⁻¹¹¹, the observed associations of the left inferior parietal are consistent
12 with the results reported in the literature. In addition, we observed spatial
13 colocalizations between regional brain volumes and their genetically correlated
14 functional connectivity traits in multiple brain regions. For instance, left pericalcarine
15 volume was genetically correlated with the connectivity strengths among its
16 neighboring regions, such as the calcarine, superior occipital, cuneus, precuneus, and
17 lingual, which were largely in the visual, default mode, and central executive networks
18 (**Fig. 3c**). More spatial overlap/proximity examples included the associations between
19 right precuneus volume and functional connectivity pairs over the precuneus, angular,
20 inferior parietal, and middle temporal regions (**Supplementary Fig. 14b**); and the
21 associations between postcentral volumes and functional interactions among the
22 postcentral, inferior and superior parietal, supramarginal, and precuneus regions
23 (**Supplementary Fig. 14c**).

24
25 Significant genetic correlations were also observed between brain structural
26 connectivity and functional connectivity ($|gc|$ range = (0.25, 0.49), P range = (5.5×10^{-10} ,
27 1.3×10^{-5}), **Supplementary Fig. 15**). Many of the white matter tracts, in particular the
28 SLF and corpus callosum, manifested a strong genetic correlation with the interactions
29 of functional networks (**Fig. 4a**). These results provided genetic evidence on how these
30 distributed networks communicate across large distances. The SLF has been widely
31 documented connecting brain regions in temporal, parietal, and frontal lobes¹¹².
32 Functionally, SLF has been reported associated with a wide array of brain functions,

including working memory¹¹³, attention^{114,115}, and language functions^{116,117}. We observed significant genetic correlations between SLF and connectivity strengths over multiple pairs of brain regions including the frontal, parietal, and temporal regions ($|gc|$ range = (0.33, 0.49), $P < 2.4 \times 10^{-6}$, **Fig. 4b**). For example, a significant association between insula and temporal connection and SLF was observed. This finding is consistent with the well documented broad functions of insula, including attention and salience processes¹⁰⁴. Furthermore, parietal and frontal connections most likely reflected attention and executive control networks. Moreover, the splenium of corpus callosum (SCC) is located in the most posterior part of the corpus callosum and connects brain regions in the temporal, posterior parietal, and occipital lobes. Our results show that SCC was genetically associated with brain regions within the parietal lobe ($|gc|$ range = (0.34, 0.48), $P < 6.5 \times 10^{-6}$, **Fig. 4c**). In particular, multiple regions connected to the precuneus were observed, such as the inferior parietal, supramarginal, and occipital regions. The precuneus has been shown to connect multiple cortical and subcortical regions. Functionally, the precuneus is one of the critical areas of the default mode network and has also been implicated to be associated with attention as well as memory functions¹¹⁸. Our findings suggest that these connections may be genetically mediated by the SCC. Besides functional connectivity traits, amplitude traits also had significant genetic associations with regional brain volumes and white matter tracts (**Supplementary Figs. 16-17, Supplementary Note**). Overall, our results uncover the genetic links between intrinsic brain function networks and the associated structural substrates. As illustrated, a few pairs of the genetically correlated brain functional and structural traits show high congruity in spatial location and the involved functions. There has been growing interest to understand how brain topography interacts with brain functional networks¹¹⁹. To our knowledge, our results are the first to indicate that genetic changes in brain structure may also impact brain function.

Next, we examined the genetic correlations between 1,777 intrinsic brain activity traits and 30 other complex traits, mainly focusing on brain disorders and cognition (**Supplementary Table 9**). We found 176 significant pairs between 26 complex traits and 102 intrinsic brain activity traits at FDR 5% level ($30 \times 1,777$ tests, P range = (8.6×10^{-12} , 2.3×10^{-3}), **Supplementary Table 10**). Particularly, functional connectivity strengths

were genetically correlated with a few brain disorders, including attention deficit hyperactivity disorder (ADHD), schizophrenia (SCZ), major depressive disorder (MDD), and cross disorder (five major psychiatric disorders¹²⁰) ($|gc|$ range = (0.18, 0.37), $P < 1.2 \times 10^{-4}$, **Fig. 5a**). For example, we observed a significant genetic correlation between ADHD and functional interactions among the precentral, supplementary motor area, superior frontal, putamen, and caudate regions, which were largely in the attention, salience, motor, and subcortical-cerebellum networks (**Fig. 5b**). These brain regions have been widely implicated with ADHD in previous studies. ADHD patients have been observed to have stronger connectivity across the supplementary motor area, precentral, and superior frontal regions¹²¹. These regions are also associated with difficulties in performing some fine motor skills¹²². In addition, the putamen and caudate regions compose the dorsal striatum, one largest part of the basal ganglia, which is important in controlling motor functions^{123,124}. Moreover, significant genetic correlations were observed between SCZ and connection strengths over the precentral, postcentral, precuneus, frontal, and superior parietal regions (**Fig. 5c**); and between MDD and the interactions among the middle temporal, angular, and superior and middle frontal regions (**Fig. 5d, Supplementary Note**).

In addition, many genetic correlations were observed between functional connectivity and cognitive traits studied in previous GWAS, including intelligence, cognitive performance, general cognitive function, and numerical reasoning. For example, intelligence had genetic correlations with connection strengths over multiple brain regions ($|gc|$ range = (0.11, 0.34), $P < 1.8 \times 10^{-4}$, **Fig. 5e**). The strongest correlation located at the superior and middle frontal regions in the central executive and salience networks. It is known that the frontal lobe is associated with higher level cognitive skills, such as problem solving, thinking, planning, and organizing¹²⁵. Wang, et al. ¹²⁶ revealed a general intelligence network for logical-math, general intelligence, and linguistic skills, which widely included frontal, parietal, occipital, temporal, and limbic regions. Furthermore, significant genetic correlations were broadly observed on subjective well-being, education, neuroticism, sleep, risk tolerance, automobile speeding, manual occupation, BMI, high blood pressure, and behavioral factors (drinking and smoking)

(**Supplementary Figs. 18-19**). More details and interpretations can be found in **Supplementary Note**.

Gene-level association analysis and biological annotations.

Gene-level association was tested via MAGMA¹²⁷ (Methods), which detected 970 significant gene-trait associations ($P < 1.5 \times 10^{-9}$, adjusted for 1,777 phenotypes) for 123 genes (**Supplementary Fig. 20, Supplementary Table 11**). In addition, we applied FUMA⁵⁴ to map significant variants ($P < 2.8 \times 10^{-11}$) to genes via physical position, expression quantitative trait loci (eQTL) association, and 3D chromatin (Hi-C) interaction, which yielded 197 more associated genes that were not discovered in MAGMA (276 in total, **Supplementary Table 12**). For the 320 genes associated with intrinsic brain activity in either MAGMA or FUMA, 84 had been linked to white matter microstructure¹⁰³, 48 were reported to be associated with regional brain volumes³⁷, and 42 were related to both of them (**Supplementary Table 13**). These triple overlapped genes were also widely associated with other complex traits, such as Parkinson's disease, neuroticism, stroke, alopecia, handedness, and intelligence (**Supplementary Table 14**), providing more insights into the genetic overlaps among brain structure, brain function, and other brain-related traits. For example, *MAPT*, *NSF*, *WNT3*, and *LRRC37A3* were risk genes of Parkinson's disease, which were also associated with pallidum volumes³⁷, white matter microstructure¹⁰³, and intrinsic functional connectivity in central executive, default mode, and salience networks. These complementary neuroimaging traits had all been used to study the pathophysiology of Parkinson's disease¹²⁸⁻¹³⁰. Similarly, *CDKN2C* and *FAF1* were associated with ischemic stroke¹³¹ as well as multiple neuroimaging traits of brain structure and function. In addition, 4 of our intrinsic brain activity associated genes (*CALY*, *SLC47A1*, *CYP2C8*, and *CYP2C9*) were targets for 11 nervous system drugs¹³², such as 4 psycholeptics (ATC code: N05) to produce calming effects, 2 anti-depressants (N06A) to treat MDD and related conditions, 2 anti-migraine (N02C), and one anti-dementia (N06D) (**Supplementary Table 15**).

It is of particular interest to study the functional connectivity dysfunction in Alzheimer's disease and identify the overlapped genes^{20,133}. Our gene-level analysis replicated *APOE* and *SORL1*, which were frequently targeted in Alzheimer's disease-candidate gene

1 studies of functional connectivity^{23,134}. More importantly, we uncovered more
 2 overlapped genes between intrinsic brain activity and Alzheimer's disease, such as
 3 *PVRL2*, *TOMM40*, *APOC1*, *MAPK7*, *CLPTM1*, *HESX1*, *BCAR3*, *ANO3*, and *YAP1*
 4 (**Supplementary Table 16**). Interestingly, through the BIG-KP server, we found that
 5 these genes had much stronger associations with intrinsic brain function than brain
 6 structure. We also observed many pleiotropic genes associated with serum metabolite,
 7 low density lipoprotein cholesterol, high density lipoprotein cholesterol, triglyceride,
 8 type II diabetes mellitus, and blood protein measurements, all of which might be related
 9 to the Alzheimer's disease^{135,136}. These results expand the overview of the shared
 10 genetic components among metabolic dysfunction, blood biomarkers, brain function in
 11 Alzheimer's disease research, suggesting the potential value of integrating these traits in
 12 future studies.

13
 14 To identify the tissues and cell types in which genetic variation yields differences in brain
 15 functional connectivity, we performed partitioned heritability analyses¹³⁷ for tissue type
 16 and cell type specific regulatory elements¹³⁸ (Methods). We focused on the 10
 17 functional connectivity traits that had heritability higher than 30%. At FDR 5% level, the
 18 most significant enrichments of heritability were observed in active gene regulation
 19 regions of fetal brain tissues, neurospheres, and neuron/neuronal progenitor cultured
 20 cells (**Supplementary Fig. 21, Supplementary Table 17**). We also tried to further identify
 21 brain cell type specific enrichments using chromatin accessibility data of two main gross
 22 brain cell types¹³⁹ (i.e., neurons (NeuN+) and glia (NeuN-)) and multiple neuronal and
 23 glial cell subtypes, including oligodendrocyte (NeuN-/Sox10+), microglia, and astrocyte
 24 (NeuN-/Sox10-), as well as GABAergic (NeuN+/Sox6+) and glutamatergic neurons
 25 (NeuN+/Sox6-). Although enrichments were observed in some cell types, few of them
 26 remained significant after adjusting for multiple testing (**Supplementary Fig. 22,**
 27 **Supplementary Table 18**). Next, we performed MAGMA tissue-specific gene property¹²⁷
 28 analysis for 13 GTEx¹⁴⁰ (v8) brain tissues (Methods). We found that genes with higher
 29 expression levels in human brain tissues generally had stronger associations with
 30 intrinsic brain activity, particularly for tissues sampled from cerebellar hemisphere and
 31 cerebellum regions ($P < 1.9 \times 10^{-5}$, **Supplementary Fig. 23, Supplementary Table 19**).

32

Among the associated variants of intrinsic brain activity, a few resided in frequently interacting regions (FIREs) and topologically associating domain (TAD) boundaries in brain tissues^{141,142} (**Supplementary Table 20**). Partitioned heritability analysis also provided suggestive evidence of heritability enrichment in these FIREs and TAD boundaries (**Supplementary Fig. 24, Supplementary Table 21**). We performed additional gene mapping using 14 recent Hi-C datasets of brain tissue and cell types¹⁴¹⁻¹⁴⁵ (Methods). This Hi-C gene mapping prioritized 29 genes, 14 of which were not identified by the Hi-C analysis in FUMA⁵⁴ (**Supplementary Table 22**). Many of the newly mapped genes have been reported for brain-related disorders/conditions, sleep, and intelligence, including *APOE*, *HSPG2*, *APOC1*, *UFL1*, *NR2F1*, *NPM1*, *FAM172A*, *FADD*, *FHL5*, and *EPHA3*. Finally, MAGMA¹²⁷ gene-set analysis was performed to prioritize the enriched biological pathways (Methods). We found 59 significantly enriched gene sets after Bonferroni adjustment ($P < 1.8 \times 10^{-9}$, **Supplementary Table 23**). Multiple pathways related to nervous system were detected, such as “go neurogenesis” (GO: 0022008), “go neuron differentiation” (GO: 0030182), “go regulation of nervous system development” (GO: 0051960), “go regulation of neuron differentiation” (GO: 0045664), “go cell morphogenesis involved in neuron differentiation” (GO: 0048667), and “go neuron development” (GO: 0048666).

DISCUSSION

In the present study, we evaluated the influences of common variants on intrinsic brain functional architecture using harmonized rsfMRI data of 44,190 subjects from four independent studies. Genome-wide association analysis found hundreds of novel loci related to intrinsic brain activity in the UKB British cohort, which were successfully replicated in independent datasets. The interactions across core neurocognitive networks (central executive, default mode, and salience) in the triple network model had genetic links with cognition and multiple brain disorders. Shared genetic influences among functional, structural, and diffusion neuroimaging traits were also uncovered, showing that brain structure and function are intimately related. Gene-level analysis detected many overlapped genes between intrinsic brain activity and Alzheimer’s disease. We also detected a colocalization between one of the two variants in the *APOE* $\epsilon 4$ locus and function of the default mode, central executive, attention, and visual

networks, which may explain in part the functional mechanism underlying Alzheimer's risk. The enriched tissues and biological pathways were also prioritized in bioinformatic analyses. Compared to the previous study³¹ with about 8,000 subjects, this large-scale GWAS much improved our understanding of the genetic architecture of functional human brain.

Our study faces a few limitations. First, the samples in our discovery GWAS were mainly from European ancestry. In our PRS analysis, we illustrated a relatively poor replication of the European GWAS results within validation cohorts with non-European ancestry. The non-European GWAS was of small sample size, so population specific influences will be better understood when more data from global populations become available. Second, our study focused on the brain functional activity at rest. A recent study²⁸ had found that combining rsfMRI and task functional magnetic resonance imaging (tfMRI) may result in higher heritability estimates and potentially boost the GWAS power. Thus, future studies could model rsfMRI and tfMRI together to uncover more insights into the genetic influences on brain function. In addition, we applied ICA in this study, which was a popular approach to characterize the functionally connected brain⁶. It is also of great interest to evaluate the performance of other popular rsfMRI approaches (such as seed-based analysis) in these large-scale datasets. Finally, although we found genetic links between brain function and other complex traits, future work is needed to dissect the underlying mechanisms by which genetic variation leads to differences in brain activity. We expect that accumulating publicly available imaging genetics data resources will lead to a better understanding of specific genes involved in human brain structure function relationships and how variants can alter these relationships leading to risk for neuropsychiatric disorders.

URLs.

Brain Imaging Genetics Knowledge Portal (BIG-KP), <https://bigkp.org/>;
Brain Imaging GWAS Summary Statistics, <https://github.com/BIG-S2/GWAS>;
UKB Imaging Pipeline, https://git.fmrib.ox.ac.uk/falmagro/UK_biobank_pipeline_v_1;
PLINK, <https://www.cog-genomics.org/plink2/>;
GCTA & fastGWA, <http://cnsgenomics.com/software/gcta/>;

- 1 METAL, <https://genome.sph.umich.edu/wiki/METAL>;
- 2 FUMA, <http://fuma.ctglab.nl/>;
- 3 MGAMA, <https://ctg.cncr.nl/software/magma>;
- 4 LDSC, <https://github.com/bulik/ldsc/>;
- 5 FINDOR, <https://github.com/gkichaev/FINDOR>;
- 6 NHGRI-EBI GWAS Catalog, <https://www.ebi.ac.uk/gwas/home>;
- 7 The atlas of GWAS Summary Statistics, <http://atlas.ctglab.nl/>.

9 METHODS

10 Methods are available in the **Methods** section.

11 *Note: One supplementary information pdf file and one supplementary table zip file are*
 12 *available.*

14 ACKNOWLEDGEMENTS

15 This research was partially supported by U.S. NIH grants MH086633 (H.Z.) and
 16 MH116527 (TF.L.). We thank the individuals represented in the UK Biobank, ABCD, HCP,
 17 and PNC studies for their participation and the research teams for their work in
 18 collecting, processing and disseminating these datasets for analysis. We gratefully
 19 acknowledge all the studies and databases that made GWAS summary data available.
 20 This research has been conducted using the UK Biobank resource (application number
 21 22783), subject to a data transfer agreement. Part of the data used in the preparation of
 22 this article were obtained from the Adolescent Brain Cognitive Development (ABCD)
 23 Study (<https://abcdstudy.org>), held in the NIMH Data Archive (NDA). This is a multisite,
 24 longitudinal study designed to recruit more than 10,000 children age 9-10 and follow
 25 them over 10 years into early adulthood. The ABCD Study is supported by the National
 26 Institutes of Health and additional federal partners under award numbers
 27 U01DA041022, U01DA041028, U01DA041048, U01DA041089, U01DA041106,
 28 U01DA041117, U01DA041120, U01DA041134, U01DA041148, U01DA041156,
 29 U01DA041174, U24DA041123, U24DA041147, U01DA041093, and U01DA041025. A full
 30 list of supporters is available at <https://abcdstudy.org/federal-partners.html>. A listing of
 31 participating sites and a complete listing of the study investigators can be found at
 32 <https://abcdstudy.org/scientists/workgroups/>. ABCD consortium investigators designed

and implemented the study and/or provided data but did not necessarily participate in analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators. Support for the collection of the PNC datasets was provided by grant RC2MH089983 awarded to Raquel Gur and RC2MH089924 awarded to Hakon Hakonarson. All PNC subjects were recruited through the Center for Applied Genomics at The Children's Hospital in Philadelphia. HCP data were provided by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University.

AUTHOR CONTRIBUTIONS

B.Z., H.Z., J.L.S., S.M.S., and Y.L. designed the study. B.Z., T.F.L., D.X., X.W., Y.Y., T.Y.L., N.M., Q.S., Y.C.Y. analyzed the data. T.F. L., Z.Z., and Y.S. downloaded the datasets, processed rsfMRI data, and undertook quality controls. P.R., M.E.H., J.B., and J.F.F. analyzed brain cell chromatin accessibility data. B.Z. and H.Z. wrote the manuscript with feedback from all authors.

CORRESPONDENCE AND REQUESTS FOR MATERIALS should be addressed to H.Z.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

REFERENCES

1. Petersen, S.E. & Sporns, O. Brain networks and cognitive architectures. *Neuron* **88**, 207-219 (2015).
2. Park, H.-J. & Friston, K. Structural and functional brain networks: from connections to cognition. *Science* **342**(2013).
3. Medaglia, J.D., Lynall, M.-E. & Bassett, D.S. Cognitive network neuroscience. *Journal of cognitive neuroscience* **27**, 1471-1491 (2015).

- 1 4. van den Heuvel, M.P. & Sporns, O. Network hubs in the human brain. *Trends in*
2 *cognitive sciences* **17**, 683-696 (2013).
- 3 5. Fox, M.D. & Greicius, M. Clinical applications of resting state functional
4 connectivity. *Frontiers in systems neuroscience* **4**, 19 (2010).
- 5 6. Lv, H. *et al.* Resting-state functional MRI: everything that nonexperts have always
6 wanted to know. *American Journal of Neuroradiology* **39**, 1390-1399 (2018).
- 7 7. Biswal, B., Zerrin Yetkin, F., Haughton, V.M. & Hyde, J.S. Functional connectivity
8 in the motor cortex of resting human brain using echo-planar MRI. *Magnetic*
9 *resonance in medicine* **34**, 537-541 (1995).
- 10 8. Fox, M.D. & Raichle, M.E. Spontaneous fluctuations in brain activity observed
11 with functional magnetic resonance imaging. *Nature reviews neuroscience* **8**,
12 700-711 (2007).
- 13 9. Alfaro-Almagro, F. *et al.* Image processing and Quality Control for the first 10,000
14 brain imaging datasets from UK Biobank. *NeuroImage* **166**, 400-424 (2018).
- 15 10. Jia, X.-Z. *et al.* Percent amplitude of fluctuation: a simple measure for
16 resting-state fMRI signal at single voxel level. *Plos one* **15**, e0227021 (2020).
- 17 11. Hutchison, R.M. *et al.* Dynamic functional connectivity: promise, issues, and
18 interpretations. *Neuroimage* **80**, 360-378 (2013).
- 19 12. Finn, E.S. *et al.* Functional connectome fingerprinting: identifying individuals
20 using patterns of brain connectivity. *Nature neuroscience* **18**, 1664-1671 (2015).
- 21 13. Smith, S.M. *et al.* Correspondence of the brain's functional architecture during
22 activation and rest. *Proceedings of the national academy of sciences* **106**,
23 13040-13045 (2009).
- 24 14. Yeo, B.T. *et al.* The organization of the human cerebral cortex estimated by
25 intrinsic functional connectivity. *Journal of neurophysiology* (2011).
- 26 15. Menon, V. & Uddin, L.Q. Saliency, switching, attention and control: a network
27 model of insula function. *Brain Structure and Function* **214**, 655-667 (2010).
- 28 16. Menon, V. Large-scale brain networks and psychopathology: a unifying triple
29 network model. *Trends in cognitive sciences* **15**, 483-506 (2011).
- 30 17. Putcha, D., Ross, R.S., Cronin-Golomb, A., Janes, A.C. & Stern, C.E. Saliency and
31 default mode network coupling predicts cognition in aging and Parkinson's

- 1 disease. *Journal of the International Neuropsychological Society: JINS* **22**, 205
- 2 (2016).
- 3 18. Menon, V. The triple network model, insight, and large-scale brain organization
- 4 in autism. *Biological psychiatry* **84**, 236 (2018).
- 5 19. Lee, M.H., Smyser, C.D. & Shimony, J.S. Resting-state fMRI: a review of methods
- 6 and clinical applications. *American Journal of neuroradiology* **34**, 1866-1872
- 7 (2013).
- 8 20. Badhwar, A. *et al.* Resting-state network dysfunction in Alzheimer's disease: a
- 9 systematic review and meta-analysis. *Alzheimer's & Dementia: Diagnosis,*
- 10 *Assessment & Disease Monitoring* **8**, 73-85 (2017).
- 11 21. Wolters, A.F. *et al.* Resting-state fMRI in Parkinson's disease patients with
- 12 cognitive impairment: A meta-analysis. *Parkinsonism & Related Disorders* **62**,
- 13 16-27 (2019).
- 14 22. Mulders, P.C., van Eijndhoven, P.F., Schene, A.H., Beckmann, C.F. & Tendolkar, I.
- 15 Resting-state functional connectivity in major depressive disorder: a review.
- 16 *Neuroscience & Biobehavioral Reviews* **56**, 330-344 (2015).
- 17 23. Foo, H. *et al.* Genetic influence on ageing-related changes in resting-state brain
- 18 functional networks in healthy adults: a systematic review. *Neuroscience &*
- 19 *Biobehavioral Reviews* (2020).
- 20 24. Teeuw, J. *et al.* Genetic and environmental influences on functional connectivity
- 21 within and between canonical cortical resting-state networks throughout
- 22 adolescent development in boys and girls. *NeuroImage* **202**, 116073 (2019).
- 23 25. Meda, S.A. *et al.* Multivariate analysis reveals genetic associations of the resting
- 24 default mode network in psychotic bipolar disorder and schizophrenia.
- 25 *Proceedings of the National Academy of Sciences* **111**, E2066-E2075 (2014).
- 26 26. Duncan, L. *et al.* Analysis of polygenic risk score usage and performance in
- 27 diverse human populations. *Nature communications* **10**, 1-9 (2019).
- 28 27. Glahn, D. *et al.* Genetic control over the resting brain. *Proceedings of the*
- 29 *National Academy of Sciences* **107**, 1223-1228 (2010).
- 30 28. Elliott, M.L. *et al.* General functional connectivity: Shared features of
- 31 resting-state and task fMRI drive reliable and heritable individual differences in
- 32 functional brain networks. *NeuroImage* **189**, 516-532 (2019).

- 1 29. Ge, T., Holmes, A.J., Buckner, R.L., Smoller, J.W. & Sabuncu, M.R. Heritability
2 analysis with repeat measurements and its application to resting-state functional
3 connectivity. *Proceedings of the National Academy of Sciences* **114**, 5521-5526
4 (2017).
- 5 30. Adhikari, B.M. *et al.* Heritability estimates on resting state fMRI data using
6 ENIGMA analysis pipeline. (2018).
- 7 31. Elliott, L.T. *et al.* Genome-wide association studies of brain imaging phenotypes
8 in UK Biobank. *Nature* **562**, 210-216 (2018).
- 9 32. Yang, J., Zeng, J., Goddard, M.E., Wray, N.R. & Visscher, P.M. Concepts,
10 estimation and interpretation of SNP-based heritability. *Nature Genetics* **49**,
11 1304-1310 (2017).
- 12 33. Chiesa, P.A. *et al.* Revolution of resting-state functional neuroimaging genetics in
13 Alzheimer's disease. *Trends in neurosciences* **40**, 469-480 (2017).
- 14 34. Zhang, N. *et al.* APOE and KIBRA interactions on brain functional connectivity in
15 healthy young adults. *Cerebral Cortex* **27**, 4797-4805 (2017).
- 16 35. Satizabal, C.L. *et al.* Genetic architecture of subcortical brain structures in 38,851
17 individuals. *Nature genetics* **51**, 1624-1636 (2019).
- 18 36. Grasby, K.L. *et al.* The genetic architecture of the human cerebral cortex. *Science*
19 **367**(2020).
- 20 37. Zhao, B. *et al.* Genome-wide association analysis of 19,629 individuals identifies
21 variants influencing regional brain volumes and refines their genetic
22 co-architecture with cognitive and mental health traits. *Nature Genetics* **51**,
23 1637-1644 (2019).
- 24 38. Hibar, D.P. *et al.* Common genetic variants influence human subcortical brain
25 structures. *Nature* **520**, 224-229 (2015).
- 26 39. Stein, J.L. *et al.* Identification of common variants associated with human
27 hippocampal and intracranial volumes. *Nature Genetics* **44**, 552-561 (2012).
- 28 40. Smith, S.M. & Nichols, T.E. Statistical challenges in "big data" human
29 neuroimaging. *Neuron* **97**, 263-268 (2018).
- 30 41. Botvinik-Nezer, R. *et al.* Variability in the analysis of a single neuroimaging
31 dataset by many teams. *Nature*, 1-7 (2020).

- 1 42. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes
2 of a wide range of complex diseases of middle and old age. *PLoS Medicine* **12**,
3 e1001779 (2015).
- 4 43. Casey, B. *et al.* The adolescent brain cognitive development (ABCD) study:
5 imaging acquisition across 21 sites. *Developmental cognitive neuroscience* **32**,
6 43-54 (2018).
- 7 44. Satterthwaite, T.D. *et al.* Neuroimaging of the Philadelphia neurodevelopmental
8 cohort. *Neuroimage* **86**, 544-553 (2014).
- 9 45. Somerville, L.H. *et al.* The Lifespan Human Connectome Project in Development:
10 A large-scale study of brain connectivity development in 5–21 year olds.
11 *NeuroImage* **183**, 456-468 (2018).
- 12 46. Miller, K.L. *et al.* Multimodal population brain imaging in the UK Biobank
13 prospective epidemiological study. *Nature Neuroscience* **19**, 1523-1536 (2016).
- 14 47. Beckmann, C.F. & Smith, S.M. Probabilistic independent component analysis for
15 functional magnetic resonance imaging. *IEEE transactions on medical imaging*
16 **23**, 137-152 (2004).
- 17 48. Hyvarinen, A. Fast and robust fixed-point algorithms for independent component
18 analysis. *IEEE transactions on Neural Networks* **10**, 626-634 (1999).
- 19 49. Shen, X. *et al.* Resting-state connectivity and its association with cognitive
20 performance, educational attainment, and household income in the UK Biobank.
21 *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* **3**, 878-886
22 (2018).
- 23 50. Smith, S.M. *et al.* Resting-state fMRI in the human connectome project.
24 *Neuroimage* **80**, 144-168 (2013).
- 25 51. Rolls, E.T., Huang, C.-C., Lin, C.-P., Feng, J. & Joliot, M. Automated anatomical
26 labelling atlas 3. *NeuroImage* **206**, 116189 (2020).
- 27 52. Yang, J., Lee, S.H., Goddard, M.E. & Visscher, P.M. GCTA: a tool for genome-wide
28 complex trait analysis. *The American Journal of Human Genetics* **88**, 76-82
29 (2011).
- 30 53. Zhao, B. *et al.* Large-scale GWAS reveals genetic architecture of brain white
31 matter microstructure and genetic overlap with cognitive and mental health
32 traits (n = 17,706). *Molecular Psychiatry* (2019).

- 1 54. Watanabe, K., Taskesen, E., Bochoven, A. & Posthuma, D. Functional mapping
2 and annotation of genetic associations with FUMA. *Nature Communications* **8**,
3 1826 (2017).
- 4 55. Jansen, I.E. *et al.* Genome-wide meta-analysis identifies new loci and functional
5 pathways influencing Alzheimer's disease risk. *Nature genetics* **51**, 404-413
6 (2019).
- 7 56. Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide
8 association studies, targeted arrays and summary statistics 2019. *Nucleic Acids*
9 *Research* **47**, D1005-D1012 (2018).
- 10 57. Jun, G.R. *et al.* Transethnic genome-wide scan identifies novel Alzheimer's
11 disease loci. *Alzheimer's & Dementia* **13**, 727-738 (2017).
- 12 58. Nazarian, A., Yashin, A.I. & Kulminski, A.M. Genome-wide analysis of genetic
13 predisposition to Alzheimer's disease and related sex disparities. *Alzheimer's*
14 *research & therapy* **11**, 5 (2019).
- 15 59. Harold, D. *et al.* Genome-wide association study identifies variants at CLU and
16 PICALM associated with Alzheimer's disease. *Nature genetics* **41**, 1088 (2009).
- 17 60. Ferrari, R. *et al.* A genome-wide screening and SNPs-to-genes approach to
18 identify novel genetic risk factors associated with frontotemporal dementia.
19 *Neurobiology of aging* **36**, 2904. e13-2904. e26 (2015).
- 20 61. Beecham, G.W. *et al.* Genome-wide association meta-analysis of
21 neuropathologic features of Alzheimer's disease and related dementias. *PLoS*
22 *Genet* **10**, e1004606 (2014).
- 23 62. Davies, G. *et al.* A genome-wide association study implicates the APOE locus in
24 nonpathological cognitive ageing. *Molecular Psychiatry* **19**, 76-87 (2014).
- 25 63. Liu, C. *et al.* Genome-wide association and mechanistic studies indicate that
26 immune response contributes to Alzheimer's disease development. *Frontiers in*
27 *genetics* **9**, 410 (2018).
- 28 64. Yan, Q. *et al.* Genome-wide association study of brain amyloid deposition as
29 measured by Pittsburgh Compound-B (PiB)-PET imaging. *Molecular psychiatry*,
30 1-13 (2018).
- 31 65. Scelsi, M.A. *et al.* Genetic study of multimodal imaging Alzheimer's disease
32 progression score implicates novel loci. *Brain* **141**, 2167-2180 (2018).

- 1 66. Liang, P. *et al.* Altered amplitude of low-frequency fluctuations in early and late
2 mild cognitive impairment and Alzheimer's disease. *Current Alzheimer Research*
3 **11**, 389-398 (2014).
- 4 67. Liu, X. *et al.* Abnormal amplitude of low-frequency fluctuations of intrinsic brain
5 activity in Alzheimer's disease. *Journal of Alzheimer's Disease* **40**, 387-397 (2014).
- 6 68. Mattsson, N. *et al.* Emerging β -amyloid pathology and accelerated cortical
7 atrophy. *JAMA neurology* **71**, 725-734 (2014).
- 8 69. Edwards, T.L. *et al.* Genome-wide association study confirms SNPs in SNCA and
9 the MAPT region as common risk factors for Parkinson disease. *Annals of Human*
10 *Genetics* **74**, 97-109 (2010).
- 11 70. Pickrell, J.K. *et al.* Detection and interpretation of shared genetic influences on
12 42 human traits. *Nature genetics* **48**, 709 (2016).
- 13 71. Consortium, I.P.D.G. Imputation of sequence variants for identification of genetic
14 risks for Parkinson's disease: a meta-analysis of genome-wide association
15 studies. *The Lancet* **377**, 641-649 (2011).
- 16 72. Vacic, V. *et al.* Genome-wide mapping of IBD segments in an Ashkenazi PD
17 cohort identifies associated haplotypes. *Human molecular genetics* **23**,
18 4693-4702 (2014).
- 19 73. Pankratz, N. *et al.* Meta-analysis of Parkinson's disease: identification of a novel
20 locus, RIT2. *Annals of neurology* **71**, 370-384 (2012).
- 21 74. Caligiore, D. *et al.* Parkinson's disease as a system-level disorder. *npj Parkinson's*
22 *Disease* **2**, 1-9 (2016).
- 23 75. Bernardinis, M., Atashzar, S.F., Jog, M.S. & Patel, R.V. Differential Temporal
24 Perception Abilities in Parkinson's Disease Patients Based on Timing Magnitude.
25 *Scientific reports* **9**, 1-16 (2019).
- 26 76. Rowe, J. *et al.* Attention to action in Parkinson's disease: impaired effective
27 connectivity among frontal cortical regions. *Brain* **125**, 276-289 (2002).
- 28 77. Jun, G. *et al.* A novel Alzheimer disease locus located near the gene encoding tau
29 protein. *Molecular Psychiatry* **21**, 108-117 (2016).
- 30 78. Kouri, N. *et al.* Genome-wide association study of corticobasal degeneration
31 identifies risk variants shared with progressive supranuclear palsy. *Nature*
32 *communications* **6**, 7247 (2015).

- 1 79. Höglinger, G.U. *et al.* Identification of common variants influencing risk of the
2 tauopathy progressive supranuclear palsy. *Nature genetics* **43**, 699 (2011).
- 3 80. Grove, J. *et al.* Identification of common genetic risk variants for autism
4 spectrum disorder. *Nature genetics* **51**, 431 (2019).
- 5 81. Nagel, M. *et al.* Meta-analysis of genome-wide association studies for
6 neuroticism in 449,484 individuals identifies novel genetic loci and pathways.
7 *Nature Genetics* **50**, 920 (2018).
- 8 82. Lee, J.J. *et al.* Gene discovery and polygenic prediction from a genome-wide
9 association study of educational attainment in 1.1 million individuals. *Nature*
10 *Genetics* **50**, 1112–1121 (2018).
- 11 83. Davies, G. *et al.* Genome-wide association study of cognitive functions and
12 educational attainment in UK Biobank (N= 112 151). *Molecular Psychiatry* **21**,
13 758–767 (2016).
- 14 84. Dashti, H. *et al.* GWAS in 446,118 European adults identifies 78 genetic loci for
15 self-reported habitual sleep duration supported by accelerometer-derived
16 estimates. *bioRxiv*, 274977 (2018).
- 17 85. Morris, J.A. *et al.* An atlas of genetic influences on osteoporosis in humans and
18 mice. *Nature genetics* **51**, 258 (2019).
- 19 86. Sanchez-Roige, S. *et al.* Genome-wide association study meta-analysis of the
20 Alcohol Use Disorders Identification Test (AUDIT) in two population-based
21 cohorts. *American Journal of Psychiatry*, appi. ajp. 2018.18040369 (2018).
- 22 87. Pardiñas, A.F. *et al.* Common schizophrenia alleles are enriched in
23 mutation-intolerant genes and in regions under strong background selection.
24 *Nature Genetics* **50**, 381–389 (2018).
- 25 88. Hyde, C.L. *et al.* Identification of 15 genetic loci associated with risk of major
26 depression in individuals of European descent. *Nature genetics* **48**, 1031 (2016).
- 27 89. Howard, D.M. *et al.* Genome-wide meta-analysis of depression identifies 102
28 independent variants and highlights the importance of the prefrontal brain
29 regions. *Nature neuroscience* **22**, 343 (2019).
- 30 90. Anney, R.J. *et al.* Meta-analysis of GWAS of over 16,000 individuals with autism
31 spectrum disorder highlights a novel locus at 10q24. 32 and a significant overlap
32 with schizophrenia. *Molecular Autism* **8**, 21 (2017).

- 1 91. Baselmans, B.M. *et al.* Multivariate genome-wide analyses of the well-being
2 spectrum. *Nature genetics* **51**, 445-451 (2019).
- 3 92. Lane, J.M. *et al.* Biological and clinical insights from genetics of insomnia
4 symptoms. *Nature genetics* **51**, 387-393 (2019).
- 5 93. Hill, W. *et al.* A combined analysis of genetically correlated traits identifies 187
6 loci and a role for neurogenesis and myelination in intelligence. *Molecular*
7 *Psychiatry* **24**, 169–181 (2019).
- 8 94. Supekar, K., Cai, W., Krishnadas, R., Palaniyappan, L. & Menon, V. Dysregulated
9 brain dynamics in a triple-network saliency model of schizophrenia and its
10 relation to psychosis. *Biological psychiatry* **85**, 60-69 (2019).
- 11 95. Zheng, H. *et al.* The altered triple networks interaction in depression under
12 resting state based on graph theory. *BioMed research international* **2015**.
- 13 96. Uddin, L.Q. *et al.* Salience network–based classification and prediction of
14 symptom severity in children with autism. *JAMA psychiatry* **70**, 869-879 (2013).
- 15 97. Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated genetic loci.
16 *Nature* **511**, 421 (2014).
- 17 98. Kichaev, G. *et al.* Leveraging polygenic functional enrichment to improve GWAS
18 power. *The American Journal of Human Genetics* **104**, 65-75 (2019).
- 19 99. Liu, M. *et al.* Association studies of up to 1.2 million individuals yield new insights
20 into the genetic etiology of tobacco and alcohol use. *Nature genetics* **51**, 237-244
21 (2019).
- 22 100. van der Meer, D. *et al.* Brain scans from 21,297 individuals reveal the genetic
23 architecture of hippocampal subfield volumes. *Molecular Psychiatry*, in press.
24 (2018).
- 25 101. Savage, J.E. *et al.* Genome-wide association meta-analysis in 269,867 individuals
26 identifies new genetic and functional links to intelligence. *Nature Genetics* **50**,
27 912-919 (2018).
- 28 102. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases
29 and traits. *Nature Genetics* **47**, 1236-1241 (2015).
- 30 103. Zhao, B. *et al.* Common genetic variation influencing human white matter
31 microstructure. *bioRxiv* (2020).

- 1 104. Uddin, L., Nomi, J., Hébert-Seropian, B., Ghaziri, J. & Boucher, O. Structure and
2 Function of the Human Insula. *Journal of Clinical Neurophysiology: Official*
3 *Publication of the American Electroencephalographic Society* **34**, 300-306 (2017).
- 4 105. Binkofski, F.C., Klann, J. & Caspers, S. On the neuroanatomy and functional role
5 of the inferior parietal lobule and intraparietal sulcus. in *Neurobiology of*
6 *language* 35-47 (Elsevier, 2016).
- 7 106. Cappelletti, M., Lee, H.L., Freeman, E.D. & Price, C.J. The role of right and left
8 parietal lobes in the conceptual processing of numbers. *Journal of Cognitive*
9 *Neuroscience* **22**, 331-346 (2010).
- 10 107. Koenigs, M., Barbey, A.K., Postle, B.R. & Grafman, J. Superior parietal cortex is
11 critical for the manipulation of information in working memory. *Journal of*
12 *Neuroscience* **29**, 14980-14986 (2009).
- 13 108. Artemenko, C., Sitnikova, M.A., Soltanlou, M., Dresler, T. & Nuerk, H.-C.
14 functional lateralization of arithmetic processing in the intraparietal sulcus is
15 associated with handedness. *Scientific reports* **10**, 1-11 (2020).
- 16 109. Bethmann, A., Tempelmann, C., De Bleser, R., Scheich, H. & Brechmann, A.
17 Determining language laterality by fMRI and dichotic listening. *Brain research*
18 **1133**, 145-157 (2007).
- 19 110. Szaflarski, J.P. *et al.* Language lateralization in left-handed and ambidextrous
20 people: fMRI data. *Neurology* **59**, 238-244 (2002).
- 21 111. Knecht, S. *et al.* Handedness and hemispheric language dominance in healthy
22 humans. *Brain* **123**, 2512-2518 (2000).
- 23 112. Unger, S.E. *et al.* The superior longitudinal fasciculus in typically developing
24 children and adolescents: diffusion tensor imaging and neuropsychological
25 correlates. *Journal of Child Neurology* **30**, 9-20 (2015).
- 26 113. Vestergaard, M. *et al.* White matter microstructure in superior longitudinal
27 fasciculus associated with spatial working memory performance in children.
28 *Journal of Cognitive Neuroscience* **23**, 2135-2146 (2011).
- 29 114. Klarborg, B. *et al.* Sustained attention is associated with right superior
30 longitudinal fasciculus and superior parietal white matter microstructure in
31 children. *Human Brain Mapping* **34**, 3216-3232 (2013).

- 1 115. Hamilton, L.S. *et al.* Reduced white matter integrity in attention-deficit
2 hyperactivity disorder. *Neuroreport* **19**, 1705 (2008).
- 3 116. Rizio, A.A. & Diaz, M.T. Language, aging, and cognition: Frontal aslant tract and
4 superior longitudinal fasciculus contribute to working memory performance in
5 older adults. *Neuroreport* **27**, 689 (2016).
- 6 117. Madhavan, K.M., McQueeney, T., Howe, S.R., Shear, P. & Szaflarski, J. Superior
7 longitudinal fasciculus and language functioning in healthy aging. *Brain research*
8 **1562**, 11-22 (2014).
- 9 118. Cavanna, A.E. & Trimble, M.R. The precuneus: a review of its functional anatomy
10 and behavioural correlates. *Brain* **129**, 564-583 (2006).
- 11 119. Bijsterbosch, J.D., Beckmann, C.F., Woolrich, M.W., Smith, S.M. & Harrison, S.J.
12 The relationship between spatial configuration and functional connectivity of
13 brain regions revisited. *Elife* **8**, e44890 (2019).
- 14 120. Consortium, C.-D.G.o.t.P.G. Identification of risk loci with shared effects on five
15 major psychiatric disorders: a genome-wide analysis. *The Lancet* **381**, 1371-1379
16 (2013).
- 17 121. Barber, A.D. *et al.* Connectivity supporting attention in children with attention
18 deficit hyperactivity disorder. *Neuroimage: clinical* **7**, 68-81 (2015).
- 19 122. Mokobane, M., Pillay, B.J. & Meyer, A. Fine motor deficits and attention deficit
20 hyperactivity disorder in primary school children. *South African Journal of*
21 *Psychiatry* **25**(2019).
- 22 123. DeLong, M. *et al.* Role of basal ganglia in limb movements. *Human neurobiology*
23 **2**, 235-244 (1984).
- 24 124. Alexander, G.E. & Crutcher, M.D. Preparation for movement: neural
25 representations of intended direction in three motor areas of the monkey.
26 *Journal of neurophysiology* **64**, 133-150 (1990).
- 27 125. Gain, U. The cognitive function and the framework of the functional hierarchy.
28 *Applied Computing and Informatics* (2018).
- 29 126. Wang, L., Song, M., Jiang, T., Zhang, Y. & Yu, C. Regional homogeneity of the
30 resting-state brain activity correlates with individual intelligence. *Neuroscience*
31 *letters* **488**, 275-278 (2011).

- 1 127. de Leeuw, C.A., Mooij, J.M., Heskes, T. & Posthuma, D. MAGMA: generalized
2 gene-set analysis of GWAS data. *PLoS Computational Biology* **11**, e1004219
3 (2015).
- 4 128. Atkinson-Clement, C., Pinto, S., Eusebio, A. & Coulon, O. Diffusion tensor imaging
5 in Parkinson's disease: review and meta-analysis. *Neuroimage: Clinical* **16**,
6 98-110 (2017).
- 7 129. Muralidharan, A. *et al.* Physiological changes in the pallidum in a progressive
8 model of Parkinson's disease: Are oscillations enough? *Experimental neurology*
9 **279**, 187-196 (2016).
- 10 130. Criaud, M. *et al.* Contribution of insula in Parkinson's disease: A quantitative
11 meta-analysis study. *Human brain mapping* **37**, 1375-1392 (2016).
- 12 131. Cheng, Y.-C. *et al.* Genome-wide association analysis of young-onset stroke
13 identifies a locus on chromosome 10q25 near HABP2. *Stroke* **47**, 307-316 (2016).
- 14 132. Wang, Q. *et al.* A Bayesian framework that integrates multi-omics data and gene
15 networks predicts risk genes from schizophrenia GWAS data. *Nature*
16 *neuroscience* **22**, 691 (2019).
- 17 133. Axelrud, L.K. *et al.* Genetic risk for Alzheimer's disease and functional brain
18 connectivity in children and adolescents. *Neurobiology of aging* **82**, 10-17 (2019).
- 19 134. Karch, C.M. & Goate, A.M. Alzheimer's disease risk genes and mechanisms of
20 disease pathogenesis. *Biological psychiatry* **77**, 43-51 (2015).
- 21 135. Zhu, Z., Lin, Y., Li, X., Driver, J.A. & Liang, L. Shared genetic architecture between
22 metabolic traits and Alzheimer's disease: a large-scale genome-wide cross-trait
23 analysis. *Human genetics* **138**, 271-285 (2019).
- 24 136. Varma, V.R. *et al.* Brain and blood metabolite signatures of pathology and
25 progression in Alzheimer disease: A targeted metabolomics study. *PLoS medicine*
26 **15**, e1002482 (2018).
- 27 137. Finucane, H.K. *et al.* Partitioning heritability by functional annotation using
28 genome-wide association summary statistics. *Nature genetics* **47**, 1228-1235
29 (2015).
- 30 138. Kundaje, A. *et al.* Integrative analysis of 111 reference human epigenomes.
31 *Nature* **518**, 317 (2015).

139. Fullard, J.F. *et al.* An atlas of chromatin accessibility in the adult human brain. *Genome research* **28**, 1243-1252 (2018).
140. Aguet, F. *et al.* The GTEx Consortium atlas of genetic regulatory effects across human tissues. *BioRxiv*, 787903 (2019).
141. Schmitt, A.D. *et al.* A compendium of chromatin contact maps reveals spatially active regions in the human genome. *Cell Reports* **17**, 2042-2059 (2016).
142. Giusti-Rodriguez, P.M. & Sullivan, P.F. Using three-dimensional regulatory chromatin interactions from adult and fetal cortex to interpret genetic results for psychiatric disorders and cognitive traits. *BioRxiv*, 406330 (2019).
143. Jung, I. *et al.* A compendium of promoter-centered long-range chromatin interactions in the human genome. *Nature genetics*, 1-8 (2019).
144. Song, M. *et al.* Mapping cis-regulatory chromatin contacts in neural cells links neuropsychiatric disorder risk variants to target genes. *Nature genetics* **51**, 1252 (2019).
145. Song, M. *et al.* 3D Epigenomic Characterization Reveals Insights Into Gene Regulation and Lineage Specification During Corticogenesis. *bioRxiv* (2020).
146. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203-209 (2018).
147. Jiang, L. *et al.* A resource-efficient tool for mixed model association analysis of large-scale data. *Nature genetics* **51**, 1749 (2019).
148. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *The American Journal of Human Genetics* **81**, 559-575 (2007).
149. Willer, C.J., Li, Y. & Abecasis, G.R. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* **26**, 2190-2191 (2010).
150. Consortium, I.H. Integrating common and rare genetic variation in diverse human populations. *Nature* **467**, 52-58 (2010).
151. Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739-1740 (2011).

METHODS

1 Imaging phenotypes and datasets. The rsfMRI datasets were consistently processed
2 following the procedures in UK Biobank imaging pipeline⁹. Details about image
3 acquisition, preprocessing, and phenotype generation in each dataset can be found in
4 **Supplementary Note**. Following the previous study³¹, we generated two groups of
5 phenotypes, including 76 node amplitude traits reflecting the spontaneous neuronal
6 activity, and 1,695 pairwise functional connectivity traits quantifying co-activity for node
7 pairs, as well as 6 global functional connectivity measures to summarize all pairwise
8 functional connectivity. To aid interpretation of these phenotypes, the functional brain
9 regions characterized in ICA were labelled using the automated anatomical labeling
10 atlas⁵¹ (**Supplementary Table 24**) and were mapped onto major functional networks
11 defined in Yeo, et al. ¹⁴ and Finn, et al. ¹² (**Supplementary Figs. 25-26**). The assigned
12 location and functional networks are provided in **Supplementary Table 25**. Details of
13 our mapping procedures are provided in **Supplementary Note**. For each continuous
14 phenotype or covariate variable, values greater than five times the median absolute
15 deviation from the median value were removed. We analyzed the following nine
16 datasets separately: 1) the UKB discovery GWAS, which used data of individuals of
17 British ancestry¹⁴⁶ in the UKB study ($n = 34,691$); 2) four European validation GWAS: UKB
18 White but Non-British (UKBW, $n = 1,970$), ABCD European (ABCDE, $n = 3,821$), HCP ($n =$
19 495), and PNC ($n = 510$); 3) two non-European UKB validation GWAS: UKB Asian (UKBA,
20 $n = 446$) and UKB Black (UKBBL, $n = 232$); and 4) two non-European non-UKB validation
21 GWAS, including ABCD Hispanic (ABCDH, $n = 768$) and ABCD African American (ABCD A, n
22 $= 1,257$). See **Supplementary Table 26** for a summary of these datasets and
23 demographic information. The assignment of ancestry in UKB was based on
24 self-reported ethnicity (Data-Field 21000), which was verified in Bycroft, et al. ¹⁴⁶. The
25 ancestry in ABCD was assigned by combining the self-reported ethnicity and ancestry
26 inference results as in Zhao, et al. ¹⁰³.

27

28 GWAS discovery and validation. Details of genotyping and quality controls can be found
29 in **Supplementary Note**. SNP heritability was estimated by GCTA⁵² using all autosomal
30 SNPs in the UKB British cohort. We adjusted the effects of age (at imaging),
31 age-squared, sex, age-sex interaction, age-squared-sex interaction, imaging site, and the
32 top 40 genetic principle components (PCs). Genome-wide association analysis was

performed in linear mixed effect model using fastGWA¹⁴⁷, while adjusting the same set of covariates as in GCTA. GWAS were also separately performed via Plink¹⁴⁸ in the eight validation datasets, including UKBW, UKBBL, UKBA, ABCDA, ABCDH, ABCDE, HCP, and PNC, where the effects of age, age-squared, sex, imaging sites (if applicable), scanners (if applicable), age-sex interaction, age-squared-sex interaction, and top ten genetic PCs were adjusted.

To validate results in the UKB British discovery GWAS, meta-analysis was performed using the sample-size weighted approach via METAL¹⁴⁹. We examined whether the locus-level associations detected in the British GWAS can be replicated in the 1) meta-analyzed four European validation GWAS (UKBW, ABCDE, HCP, and PNC); 2) meta-analyzed four non-European validation GWAS (UKBBL, UKBA, ABCDA, and ABCDH); and 3) the combination of the above eight validation GWAS. Specifically, for each meta-analyzed GWAS, we checked and reported the smallest *P*-value among the variants within each associated locus identified in the UKB British discovery GWAS. Polygenic risk scores (PRS) were constructed on eight validation datasets using Plink. The BLUP effect sizes estimated from GCTA-GREML analysis in UKB British discovery GWAS were used as weights in PRS construction, which accounted for the LD structures. Ambiguous variants (i.e. variants with complementary alleles) were removed from analysis. We tried 17 *P*-value thresholds for variant selection according to their marginal *P*-values from fastGWA: 1, 0.8, 0.5, 0.4, 0.3, 0.2, 0.1, 0.08, 0.05, 0.02, 0.01, 1×10^{-3} , 1×10^{-4} , 1×10^{-5} , 1×10^{-6} , 1×10^{-7} , and 1×10^{-8} . The best prediction accuracy achieved by a single threshold was reported for each phenotype, which was measured by the additional phenotypic variation that can be explained by the polygenic profile (i.e., the incremental R-squared), while adjusting for the effects of age, sex, and top ten genetic PCs.

The shared loci and genetic correlation. The genomic loci associated with intrinsic brain activity traits were defined using FUMA (version 1.3.5e). We input UKB British discovery summary statistics after reweighting the *P*-values using functional information via FINDOR⁹⁸. To define the LD boundaries, FUMA identified independent significant variants, which were defined as variants with a *P*-value smaller than the predefined

threshold and were independent of other significant variants ($LD\ r^2 < 0.6$). FUMA then constructed LD blocks for these independent significant variants by tagging all variants in LD ($r^2 \geq 0.6$) with at least one independent significant variant and had a $MAF \geq 0.0005$. These variants included those from the 1000 Genomes reference panel that may not have been included in the GWAS. Moreover, within these significant variants, independent lead variants were identified as those that were independent from each other ($LD\ r^2 < 0.1$). If LD blocks of independent significant variants were close ($< 250\text{ kb}$ based on the closest boundary variants of LD blocks), they were merged into a single genomic locus. Thus, each genomic locus could contain multiple significant variants and lead variants. Independent significant variants and all the variants in LD with them ($r^2 \geq 0.6$) were searched by FUMA on the NHGRI-EBI GWAS catalog (version 2019-09-24) to look for previously reported associations ($P < 9 \times 10^{-6}$) with any traits. LDSC¹⁰² software (version 1.0.1) was used to estimate and test the pairwise genetic correlation. We used the pre-calculated LD scores provided by LDSC, which were computed using 1000 Genomes European data. We used HapMap3¹⁵⁰ variants and removed all variants in the major histocompatibility complex (MHC) region. The summary statistics of intrinsic brain activity traits were from the UKB British discovery GWAS and the resources of other summary statistics were provided in **Supplementary Table 9**.

Gene-level analysis and biological annotation. Gene-based association analysis was performed in UKB British participants for 18,796 protein-coding genes using MAGMA¹²⁷ (version 1.07). Default MAGMA settings were used with zero window size around each gene. We then carried out FUMA functional annotation and mapping analysis, in which variants were annotated with their biological functionality and then were linked to 35,808 candidate genes by a combination of positional, eQTL, and 3D chromatin interaction mappings. Brain-related tissues/cells were selected in all options and default values were used for all other parameters in FUMA. For the detected genes in MAGMA and FUMA, we performed lookups in the NHGRI-EBI GWAS catalog (version 2020-02-08) to explore their previously reported gene-trait associations. We performed heritability enrichment analysis via partitioned LDSC¹³⁷. Baseline models were adjusted when estimating and testing the enrichment scores for our tissue type and cell type specific annotations. Methods to analysis chromatin data of glial and neuronal cell subtypes can

be found in Zhao, et al.¹⁰³. We also performed gene property analysis for the 13 GTEx¹⁴⁰ v8 brain tissues via MAGMA. Specifically, we examined whether the tissue-specific gene expression levels can be linked to the strength of the gene-trait association. MAGMA was also used to explore the enriched biological pathways, in which we tested 500 curated gene sets and 9,996 Gene Ontology (GO) terms from the Molecular Signatures Database¹⁵¹ (MSigDB, version 7.0). Additional gene mapping was performed using 14 Hi-C datasets of brain tissue and cell types from five recent studies, including 1) the promoter capture Hi-C (PCHi-C) data of hippocampus and dorsolateral prefrontal cortex (DLPFC)¹⁴³; 2) the Hi-C data of hippocampus and DLPFC¹⁴¹; 3) the Hi-C data from fetal and adult cortices¹⁴², restricting to the high confidence interactions; 4) the PCHi-C data of primary astrocytes and three types of induced pluripotent stem cell (iPSC)-derived neurons¹⁴⁴ (cortical, hippocampal, and motor); and 5) proximity ligation assisted chromatin immunoprecipitation (PLAC-seq) data on sorted fetal neuron cells¹⁴⁵, including radial glial cells, intermediate progenitor cells, neurons, and interneurons. For interaction intensity cutoffs, we used 2 for the $-\log_{10}(P)$ used in datasets of Jung, et al.¹⁴³, 0.05 for the q -value in Schmitt, et al.¹⁴¹ and Giusti-Rodriguez and Sullivan¹⁴², 5 for the Chicago score in Song, et al.¹⁴⁴, and 0.01 for the FDR in Song, et al.¹⁴⁵.

Code availability

We made use of publicly available software and tools listed in URLs. Other codes used in our analyses are available upon reasonable request.

Data availability

Our GWAS summary statistics can be downloaded at <https://github.com/BIG-S2/GWAS>. The individual-level data used in the present study can be obtained from four publicly accessible data resources: UK Biobank (<http://www.ukbiobank.ac.uk/resources/>), ABCD (<https://abcdstudy.org/>), HCP (<https://www.humanconnectome.org/>), and PNC (<https://www.med.upenn.edu/bbl/philadelphianeurodevelopmentalcohort.html>). Our results can also be easily browsed through our knowledge portal <https://bigkp.org/>.

Figure 2: Selected genetic loci that associated with both rsfMRI traits of brain activity and other brain-related complex traits and disorders. We highlight local colocalization ($LD\ r^2 \geq 0.6$) in **a**) 19q13.32 (colocalized with Alzheimer's disease); **b**) 2p16.1 (with schizophrenia); and **c**) 2q14.1 (with sleep). For example, in 19q13.32, we observed colocalization between the amplitude of the precuneus region in the default mode and central executive networks with Alzheimer's disease. Location and functional network of the displayed three rsfMRI traits are illustrated on the bottom right. More examples of the shared genetic loci and the involved rsfMRI traits can be found in Supplementary Figures 4-12.

Figure 3: Selected pairwise genetic correlations between functional connectivity traits and regional brain volumes. **a)** The asterisks highlight significant associations after controlling the false discovery rate at 0.05 level. The left y-axis lists the location of functional connectivity traits, the right y-axis shows the associated functional networks, and the x-axis provides the name of regional brain volumes. The colors represent genetic correlations (rg). **b)** Location of the right insula and its neighboring brain regions whose functional connectivity strengths were genetically correlated with the right insula volume. The colors describe different brain regions. **c)** Location of the left pericalcarine and its neighboring brain regions whose functional connectivity strengths were genetically correlated with the left pericalcarine volume.

Figure 4: Selected pairwise genetic correlations between functional connectivity traits and fractional anisotropy (FA) of white matter tracts. a) The asterisks highlight significant associations after controlling the false discovery rate at 0.05 level. The left y-axis lists the location of functional connectivity traits, the right y-axis shows the associated functional networks, and the x-axis provides the name of white matter tracts. The colors represent genetic correlations (rg). **b)** Location of the SLF (left part) and its neighboring brain regions whose functional connectivity strengths were genetically correlated with the FA of SLF. The colors describe different brain regions. **c)** Location of the SCC (left part) and its neighboring brain regions whose functional connectivity strengths were genetically correlated with the FA of SCC.