Impaired M-current in KCNQ2 Encephalopathy Evokes Dyshomeostatic Modulation of Excitability

Dina Simkin^{1,2}, Timothy J. Searl², Brandon N. Piyevsky¹, Marc Forrest^{3,4}, Luis A. Williams⁵, Vaibhav Joshi⁵, Hongkang Zhang⁵, Steven J. Ryan⁵, Michael Schwake¹, Gabriella L. Robertson¹, Peter Penzes^{3,4}, Linda C. Laux⁶, Owen B. McManus⁵, Graham T. Dempsey⁵, John J. Millichap⁶, Alfred. L. George, Jr.^{2,*} and Evangelos Kiskinis^{1,3,*}

¹The Ken & Ruth Davee Department of Neurology, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

²Department of Pharmacology, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

³Department of Physiology, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

⁴Center for Autism and Neurodevelopment, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

⁵Q-State Biosciences, Cambridge, MA, USA

⁶Epilepsy Center and Division of Neurology, Departments of Pediatrics and Neurology, Ann & Robert H. Lurie Children's Hospital of Chicago, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

*Correspondence: al.george@northwestern.edu and evangelos.kiskinis@northwestern.edu

“The authors have withdrawn their manuscript whilst they perform additional experiments to test some of their conclusions further. Therefore, the authors do not wish this work to be cited as reference for the project. If you have any questions, please contact the corresponding authors.”