1 Germline DNA replication shapes the recombination landscape in 2 mammals

3 Florencia Pratto^{1*}, Kevin Brick^{1*}, Gang Cheng¹, Gabriel Lam¹, Jeffrey M. Cloutier¹, Daisy
4 Dahiya¹, Stephen R. Wellard², Philip W. Jordan², R. Daniel Camerini-Otero¹

5 ¹Genetics and Biochemistry Branch, NIDDK, NIH, Bethesda, Maryland, USA

6 ²Department of Biochemistry and Molecular Biology, Johns Hopkins University Bloomberg
7 School of Public Health, Baltimore, MD 21205

8 *These authors contributed equally

9 **Summary:**

10 Genetic recombination generates novel trait combinations and understanding how
11 recombination is distributed across the genome is key to modern genetics. The PRDM9 protein
12 defines recombination hotspots, however megabase-scale recombination patterning is
13 independent of PRDM9. The single round of DNA replication, which precedes recombination in
14 meiosis, may establish these patterns, therefore we devised a novel approach to study meiotic
15 replication that includes robust and sensitive mapping of replication origins. We find that meiotic
16 DNA replication is distinct; reduced origin firing slows replication in meiosis and a distinctive
17 replication pattern in human males underlies the sub-telomeric increase in recombination. We
18 detected a robust correlation between replication and both contemporary and ancestral
19 recombination and found that replication origin density coupled with chromosome size
20 determines the recombination potential of individual chromosomes. Our findings and methods
21 have far-reaching implications for understanding the mechanisms underlying DNA replication,
22 genetic recombination, and the landscape of mammalian germline variation.

Keywords

DNA replication, recombination, meiosis, germline, *in-silico* modelling, crossover, genome stability, human reproduction

Introduction

Sexual reproduction uses a specialized cell division called meiosis, in which a single round of DNA replication is followed by two cell divisions to create haploid gametes. Genetic recombination in meiosis assures faithful segregation of chromosomes and establishes patterns of genetic linkage and inheritance. Recombination is initiated by the formation of hundreds of programmed DNA double-strand breaks (DSBs). In mice and humans, DSBs are targeted by DNA sequence-specific binding of a meiosis-specific histone methyltransferase, PRDM9 (Baudat et al., 2010; Myers et al., 2010; Parvanov et al., 2010). Hundreds of PRDM9 alleles exist (Berg et al., 2010; Buard et al., 2014) that primarily differ in the DNA-binding domain. Thus, each variant may yield a distinct patterning of meiotic DSBs (Smagulova et al., 2016). Nonetheless, megabase-scale similarities between individuals demonstrate that a broad-scale, PRDM9-independent layer of control also shapes meiotic recombination (Davies et al., 2016; Myers et al., 2005; Smagulova et al., 2011). The clearest manifestation of this is seen in the elevation of meiotic DSBs in subtelomeric DNA of human males, independent of PRDM9 genotype (Pratto et al., 2014).

Since DNA replication determines genome structure (Klein et al., 2019) and immediately precedes DSB formation in meiosis, we hypothesized that DNA replication may drive the broad-scale regulation of recombination in mammals. Replication and recombination are correlated in yeasts (Borde et al., 2000; Murakami and Nurse, 2001) and in barley (Higgins et al., 2012). A mechanistic link has been demonstrated in baker's yeast, where a component of the DSB machinery is activated by passage of the replication fork (Murakami and Keeney, 2014). This results in a spatio-temporal coordination between replication and DSB formation.

All current knowledge of DNA replication in mammalian meiosis stems from classical papers using early molecular and cytological techniques (for review see (Chandley, 1986)). The paucity of studies of DNA replication in mammalian meiosis stems from the requirement to study meiosis *in-vivo*, in the context of a complex tissue (Handel et al., 2014). This contrasts with

cell-types that can be studied *in vitro*; indeed, mammalian DNA replication is exclusively studied in cell culture. This presents difficulties in adapting techniques for studying replication, which have been mostly designed and optimized for cell culture. To address this shortcoming, we devised the first method to map origins of replication in mammalian tissue (in this case, testis), developed a cell-type specific method to interrogate replication timing in meiotic S-phase and designed an *in-silico* modelling strategy to parameterize DNA replication *in-vivo*. This unique tripartite approach has generated the first comprehensive, parameterized description of DNA replication genome-wide in meiosis, or, for that matter, in any mammalian tissue. We found that DNA replication in the germline is distinct from replication in other cell types and that this unique replication patterning plays a direct role in shaping megabase-scale patterns of meiotic recombination and genome diversity.

Results

Highly specific replication origin mapping in mammalian testis

To identify origins of replication, we introduced an important adaptation to an existing method to sequence the RNA-primed short nascent leading strands (SNS) (Bielinsky and Gerbi, 1998) (Figure 1A). Briefly, RNA-primed leading strands are isolated by using lambda exonuclease to digest all DNA that lacks an RNA primer. Okazaki fragments, while also RNA-primed, are excluded by size selection. A key improvement upon SNS-Seq (Cayrou et al., 2012; Fu et al., 2014; Jodkowska et al., 2019; Picard et al., 2014) is that we directly sequence the captured single-stranded DNA (Khil et al., 2012) at origins. This retains the strand information of the nascent strand DNA (Origin-derived Single-Stranded DNA Sequencing; Ori-SSDS) (Figure 1A) and avoids biases resulting from random-priming and second-strand synthesis (Sequeira-Mendes et al., 2019). Characteristic bidirectional replication at origins manifests as a strand switch in sequenced reads. Distinct from SNS-Seq experiments, we can differentiate true origins from non-specific signals (Figure 1A-C, S1, S2) by imposing a requirement for leading-strand asymmetry at origins (see methods). Accounting for strand asymmetry substantially reduces the number of replication origins estimated by SNS-Seq experiments (40-50%; Figure S3; (Cayrou et al., 2015)) because local peaks within SNS-Seq signals have been spuriously defined as very closely-spaced replication origins (Figure S3).

81 We detected 10,932, 12,100 and 13,082 origins in replicate experiments from testes of
 82 individual male mice, demonstrating that our method is sufficiently sensitive to derive replication
 83 origin maps from individual animals. Prior to our experiments, the number of cells required to
 84 detect origins appeared to be prohibitive for *in-vivo* studies; for example, SNS-Seq typically
 85 starts from 10^9 exponentially growing cultured cells (Almeida et al., 2018). We used just 2×10^6
 86 replicating cells (Kojima et al., 2019) for Ori-SSDS; this is 3-5 fold fewer than methods that
 87 immunoprecipitate pre-replication complexes (i.e. ORC1/ORC2 ChIP-Seq (Miotto et al., 2016))
 88 and orders of magnitude fewer than are required for the sequencing of Okazaki fragments
 89 (Petryk et al., 2016). 94% of origins in the smallest set were found in at least one of the other
 90 two experiments (Figure S4) and origins unique to individual samples were weak (Figure 1D,
 91 S5). No origins were detected in control experiments, either using non-replicating tissue (sperm)
 92 or by hydrolyzing the leading strand RNA-primer (see above and methods) (Figure 1C, S1).
 93 94% of replication origins detected by Ori-SSDS in testis are found in SNS-based origin maps
 94 from other cell types (Mouse Embryonic Fibroblasts (MEFs) and Embryonic Stem (ES) cells;
 95 Figure S6). Furthermore, data from Okazaki-fragment sequencing (in activated B-cells) shows a
 96 switch in the polarity of Okazaki fragments around origin centers (Figure S6). These data
 97 strongly suggest that Ori-SSDS accurately captures origins of replication from tissue and that
 98 origins in testis mostly occur at sites that are also used as origins in other cell types.

99 Origin efficiency is a measure of the frequency with which an origin is used. We use the
 100 sequencing depth of “correctly” oriented Ori-SSDS read-pairs to infer origin efficiency (see
 101 methods). This efficiency varies ~100-fold and is highly correlated between replicates (Figure
 102 1D, S3; Spearman $R^2 = 0.65-0.74$). Despite the fragment asymmetry around the origins, there is
 103 considerable overlap of the Watson and Crick signals (Figure 1A, S1). Thus, while DNA
 104 replication originates at discrete loci, the initiation window spans several hundred base-pairs at
 105 the origin center. This contrasts with *S. cerevisiae*, where an 11 bp motif defines a more precise
 106 origin of replication (Broach et al., 1983; Stinchcomb et al., 1980). Together, these data show
 107 that Ori-SSDS can identify the origins of replication genome-wide from mammalian tissue.
 108 11,565 high-confidence origins, found in at least two of three replicates (Figure S4, S5), were
 109 used for subsequent analyses.

110 Origins identified from mouse testis are unevenly distributed in the genome (Figure S7A) and
 111 cluster in gene rich regions; 66% of origins occur within 1 Kb of a transcription start site (TSS)

(Figure S7B). DNA at the origin center is intrinsically flexible, flanked by relatively rigid DNA, and this may reflect a requirement of the ORC complex to bend DNA (Li et al., 2018) (Figure S7C). Origins are GC-rich, with elevated CpG, and to a lesser extent, GpC dinucleotides (Figure S7D). GC content, CpG density and GpC density are all positively correlated with origin efficiency (Figure S7E) implying that nucleotide content either directly or indirectly plays a role in origin firing. G-quadruplexes, a predicted secondary structure implicated in origin firing (Valton et al., 2014), are also strongly enriched at origin centers (Figure S7F). 74% of origins coincide with CpG islands (CGIs) in accessible chromatin (as measured by ATAC-Seq in spermatogonia - the germ-cells that precede meiosis (Maezawa et al., 2018); Figure 1E), yet only 53% of such sites are used for origin formation (Figure 1F). CGIs in inaccessible chromatin and open chromatin lacking a CGI are far less predictive of origin locations (Figure 1E,F). We did not identify any conserved sequence motif at replication origins, however the density of CpG dinucleotides alone can predict origin locations (Figure S7G). Indeed, CpG density is a better predictor of origin locations than binding sites are for transcription factors (Figure S7H). Many of these observations are consistent with properties of replication origins defined in cultured cells (Marchal et al., 2019; Miotto et al., 2016), and together suggest that replication in the germline initiates preferentially at CpG islands, near gene promoters, in accessible chromatin.

Broad replication initiation zones - up to 150 Kb and comprising ~7% of the genome - have been proposed as a major mode of replication initiation in humans (Petryk et al., 2016). Although most origins mapped by Ori-SSDS are narrow and discrete, 1,925 Ori-SSDS peak are wider than 10Kb, and may represent replication initiation zones (the expected width of the Ori-SSDS signal is approximately ± 3 Kb around the origin of replication; determined by the size selection step; see methods; Figure 1A; Figure S8A) To assess if putative initiation zones result from closely-spaced, but difficult-to-resolve discrete origins, we examined the Ori-SSDS signal around local peaks in CpG density (CpG peaks coincide with the center of isolated origins; Figure S8B). We found that within our defined initiation zones, most local CpG peaks exhibited origin-like Ori-SSDS strand asymmetry, implying that these sites are *de-facto* origins (Figure S8C). 620 zones contained multiple CpG peaks ($N_{\text{CpG Peaks}} = 1,645$), thus, closely-spaced and difficult-to-resolve origins account for a large number of initiation zones (Figure S8D-F). Among such zones is the extensively studied *HoxA* locus, where Ori-SSDS data suggests that ~19 origins of replication occur within just 110 Kb (Figure S8E). 816 zones contained a single CpG peak, and likely represent origins where we captured longer-than-expected nascent strands

144 (Figure S8G). The remaining 490 zones did not contain any CpG peak above our threshold and
 145 may represent more amorphous replication initiation regions. Alternatively, they may represent
 146 clusters of weak origins (Figure S8G,H). Overall, we found extensive evidence that clustered,
 147 yet discrete origins of replication can be mistaken for initiation “zones”. Previous experiments
 148 that identified “zones” lacked the resolution to detect this fine sub-structure.

149 **An unambiguous snapshot of replication in meiotic S-phase**

150 Stochastic origin firing and uneven origin density result in distinct earlier and later replicating
 151 parts of the genome. We inferred this replication timing (RT) in meiosis. Meiotic S-phase nuclei
 152 constitute just 1% of cells in adult testis (Kojima et al., 2019). Using a variant of fluorescence
 153 activated nuclei sorting (Lam et al., 2019), we isolated up to 2×10^5 meiotic S-phase nuclei at
 154 94% purity (inspection of wide-field images; Figure S9) from a single adult mouse (Figure 2A-C);
 155 this relied on a combination of DNA content, negative selection for a marker of non-meiotic cells
 156 (DMRT1) and positive selection for a meiotic protein (STRA8). A marker not used for sorting,
 157 but with expected meiotic expression (SYCP3) validated the purity of the meiotic population
 158 (Figure 2C). We then inferred meiotic RT from coverage imbalances in whole-genome
 159 sequencing of these nuclei (see methods; RT-Seq) (Koren et al., 2014). Replication origin
 160 density is high in early replicating regions and low in late-replicating DNA (Figure 2D). This
 161 yields the first snapshot of replication timing in mammalian meiosis. We also inferred RT from
 162 the other major replicating cell types in adult testis - undifferentiated and differentiating
 163 spermatogonia - and from published whole genome sequencing (WGS of other cell types (see
 164 methods; Figure S10). Meiotic RT profiles are highly correlated among replicates and, to a
 165 lesser extent, with RT from other cell types (Figure 2D-F, Figure S10). In particular, meiotic RT is
 166 notably similar to RT in the cells that immediately precede meiotic S-phase and to RT in other
 167 germ cells (PGCs, SSCs, ESCs). RT in all cell-types is highly correlated with GC-content
 168 (Spearman $R^2 = 0.27 - 0.40$) and with nuclear compartments (Spearman $R^2 = 0.57 -$
 169 0.70)(Figure 2G).

170 ***in-silico* modeling recapitulates replication timing from origin locations**

171 RT-Seq yields a static snapshot of replication and these static RT-Seq snapshots from different
 172 cell-types are often remarkably similar. Furthermore, the differences between RT-Seq profiles
 173 are difficult to interpret because RT-Seq can be strongly influenced by the average stage of

174 S-phase progression and by the relative synchrony of the population (Figure S11). We
175 hypothesized that subtle differences in the properties of replication among cell types could be
176 captured by *in-silico* modelling.

177 We built an *in silico* model of DNA replication that required experimentally defined origins as
178 input and that outputs a simulated RT profile. By examining the properties of replication that
179 produce a best-fit between simulated RT and experimental RT, we can understand the
180 parameters of replication that yield different RT profiles. Modeling parameters are the number of
181 active forks, whether to use origin efficiency estimates as a firing probability, and the total
182 duration of simulated S-phase (see methods). Replication fork speed is assumed constant
183 among cell types and throughout S-phase. After an initial round of simultaneous origin firing,
184 further origins fire only when two extant forks collide; this simulates the presence of factors that
185 limit the number of active origins (replisome ceiling), deemed important by previous simulations
186 of DNA replication (Gindin et al., 2014; Kelly and Callegari, 2019). Best fitting models were
187 obtained using a grid search for optimal model parameters (see methods).

188 We found that *in silico* modeling could accurately recapitulate experimental RT from both
189 meiotic (Figure 3A,B, Figure S12A) and non-meiotic S-phase cells (Figure 3B,C, Figure S12B).
190 In contrast, we could not obtain good-fitting models for non-replicating cells (2C) or when origin
191 locations were randomized (Ori(r); Figure 3B). Our success in modeling RT in all cell types
192 using testis-derived origins implies that a common set of origins are used during meiotic and
193 mitotic replication. Indeed in yeast, origins of replication are common to meiotic and mitotic cells
194 (Blitzblau et al., 2012; Wu and Nurse, 2014). Most testis-derived replication origins coincide with
195 CpG islands in open chromatin (ATAC-CGIs; Figure 1E,F), and these regions are broadly similar
196 across diverged cell types in mouse (Figure S13). RT can be modeled using ATAC-CGIs as a
197 proxy for origins, but these models are slightly worse than those using *de facto* origins (AS+CGI;
198 Figure 3B). Finally, RT-Sim does not explicitly distinguish between “early” and “late” firing
199 replication origins, suggesting that the paradigm of early and late origins is not required to
200 explain RT. Instead, “early replicating regions” and “late replicating regions” are defined by high
201 and low origin density, respectively. This is consistent with recent studies in other mammalian
202 cell types (Dileep and Gilbert, 2018; Gindin et al., 2014; Miotto et al., 2016; Takahashi et al.,
203 2019).

204 **Reduced origin firing slows DNA replication in meiosis**

205 To validate that RT-Sim can explain meaningful properties of replication, we examined model
 206 run-time of best-fitting models for very early-, early-, middle- and late- S-phase nuclei (Figure
 207 3D). Optimal model runtime - defined as the time at which simulated RT best fits experimental
 208 RT - should reflect the time a population has spent in S-phase. Indeed, optimal runtime got
 209 progressively longer from very early through late S-phase populations (Figure 3D),
 210 demonstrating that modeling yields meaningful numeric insights into the properties of DNA
 211 replication. Importantly, the differences between experimental RT in these populations do not
 212 substantially affect model estimates of total S-phase duration (Figure 3D). Accurate estimates of
 213 the duration of meiotic S-phase in mammals are notoriously difficult to obtain (Kofman-Alfaro
 214 and Chandley, 1970); in mice, estimates range from 14 hrs (Monesi, 1962) to 29 hrs (Ghosal
 215 and Mukherjee, 1971). Like in other organisms (Cha et al., 2000; Kofman-Alfaro and Chandley,
 216 1970; Lee and Amon, 2001), DNA replication in meiotic S-phase is thought to be longer than
 217 S-phase in non-meiotic cells (Kofman-Alfaro and Chandley, 1970). We find that for best-fitting
 218 models, median S-phase duration in meiosis is 1.4 - 1.8 fold longer than in spermatogonia - the
 219 cells that precede meiosis (Figure 3E, S14), despite having highly correlated
 220 experimentally-measured RT (Figure 2F). In *Saccharomyces cerevisiae*, the slow-down of DNA
 221 replication in meiosis may be partly to facilitate recombination, as knocking out *Spo11* - the
 222 protein that makes meiotic DSBs - reduces S-phase duration by 30% (Cha et al., 2000). Unlike
 223 in yeast, we found no reduction in meiotic S-phase duration in *Spo11*^{-/-} mice (Figure S14).

224 The slow-down we observe in meiotic S-phase results from the use of fewer replisomes;
 225 best-fitting models use just 0.24 - 0.28 replisomes Mb⁻¹ in meiosis (648 - 756 replisomes per
 226 haploid genome; 2,700 Mb genome), compared to 0.44 - 0.54 replisomes Mb⁻¹ in ES cells
 227 (1,200 - 1,500 replisomes per haploid genome) (Figure 3F). These estimates in ES cells are
 228 similar to the replisome count in cultured mouse C2C12 cells (imaging-based estimates; 0.46 -
 229 0.48 replisomes Mb⁻¹; (Chagin et al., 2016)). The decreased replisome density in meiosis
 230 mirrors findings in newt spermatocytes, where replication tracts in meiosis were notably longer
 231 than in mitotic cells (Callan, 1973). Although we don't explicitly model fork speed, universally
 232 altering fork speed without changing origin density cannot explain the presence of longer
 233 replication tracts in meiosis. Indeed, in other organisms, fork speed does not vary between
 234 meiosis and mitosis (Borde et al., 2000; Callan, 1973). By extrapolating from published

estimates of S-phase duration in intermediate-stage spermatogonia (12.5 hr (Monesi, 1962)), we estimate that meiotic S-phase in mice takes 21-24 hrs. Interestingly, the range of S-phase duration estimates for the best-fitting models of meiotic S phase is larger than that of the other populations (Figure 3E, S14). Given that the model simulates replication in single cells, this might indicate that there is a higher heterogeneity of RT profiles in meiosis vs mitosis. This is in line with the experimental data spread seen in yeast (Cha et al., 2000).

Together, these analyses suggest both that the origins captured by Ori-SSDS reflect meiotic replication initiation and that the same origins can yield tangibly different RT profiles if other properties of replication vary. This tripartite approach of origin mapping, RT-Seq and *in-silico* modeling, offers an alternative to classical cytogenetic approaches and yields an unprecedented description of the DNA replication landscape.

Direct coupling of replication with meiotic recombination

Programmed DSB formation in meiosis occurs after DNA replication. In *Saccharomyces cerevisiae*, passage of the replication fork favors meiotic DSB formation, likely through the phosphorylation of Mer2 (a key protein for DSB formation) (Murakami and Keeney, 2014). This interplay is completely unexplored in large mammalian genomes. In mice and humans, local DSB patterning is determined by the sequence-specific binding of PRDM9 (Baudat et al., 2010; Myers et al., 2010), yet at megabase scales, DSB density is highly correlated between individuals with different PRDM9 alleles (Davies et al., 2016; Smagulova et al., 2011). We therefore asked whether DNA replication underlies the megabase-scale control of DSB patterning.

Meiotic DSBs form through a well-documented cascade; PRDM9 tri-methylates H3K4 (Brick et al., 2012; Diagouraga et al., 2018) and/or H3K36 (Diagouraga et al., 2018; Powers et al., 2016) which appear to recruit the DSB complex. A DSB is made by the SPO11 protein, which is then released with a short oligonucleotide (Bergerat et al., 1997; Keeney et al., 1997; Neale et al., 2005). The 5' DNA is resected and the DMC1 protein loads on exposed ssDNA to facilitate homologous recombination (Bishop et al., 1992). DSBs ultimately repair as a crossover (CO) or as a non-crossover (NCO). H3K4me3 ChIP-Seq, SPO11-associated oligo mapping (Lange et al., 2016), DMC1 ChIP-single-stranded DNA sequencing (Brick et al., 2012) and repair outcome

264 mapping (COs/NCOs) provide differing quantitative readouts of intermediates in this cascade
265 (Figure 4A).

266 We found that all measures of recombination are enriched in the early replicating regions
267 (ERRs) (Figure 4). Nonetheless, informative differences are apparent. PRDM9-mediated
268 H3K4m3 and DSB frequency (SPO11-oligo) are similarly enriched relative to RT (Figure 4B;
269 yellow and magenta lines). Both profiles “flatten” in the earliest replicating DNA. This likely
270 reflects hotspot erosion (Boulton et al., 1997; Myers et al., 2010), a process by which strong
271 PRDM9 binding sites are purged from the genome (Figure 4C). Hotspot erosion is indicative of
272 ancestral recombination, reinforcing the strong association between RT and recombination in
273 mice. These findings also imply that the action of PRDM9 decouples replication and
274 recombination through the erosion of PRDM9 binding sites. In contrast to PRDM9-mediated
275 H3K4me3 and SPO11-oligo density, DSB repair intermediates (DMC1-SSDS, RPA-SSDS) are
276 relatively depleted in the very earliest replicating regions (Figure 4B,D, S15). SSDS captures
277 both the frequency (Lange et al., 2016) and the lifetime (Pratto et al., 2014) of DSB repair
278 intermediates. Since DSB frequency closely mirrors PRDM9-mediated H3K4me3, we conclude
279 that DSBs forming in early replicating regions are more rapidly repaired than those elsewhere.
280 In agreement with this hypothesis, the signature of rapid repair is no longer seen in
281 DMC1-SSDS from a mouse in which all meiotic DSBs remain unrepaired (*Hop2*^{-/-}; Figure S15).
282 The signature of rapid DSB repair in ERRs is also seen in mice that lack PRDM9 (*Prdm9*^{-/-};
283 Figure S15). Thus, this phenomenon is independent of the mechanisms that determine the local
284 patterning of meiotic DSBs.

285 Rapid DSB repair in meiosis has been proposed as a hallmark of crossover-biased DSB repair
286 (Brick et al., 2018a; Hinch et al., 2019). To simplify the study of repair outcomes we turned to a
287 mouse homozygous for a “humanized” PRDM9 allele; this allele of PRDM9 has a binding
288 preference not found naturally in mice, and therefore, has left no footprint of hotspot erosion in
289 the mouse genome. Indeed, in these mice, PRDM9-mediated H3K4me3 is linearly correlated
290 with RT in the earliest replicating DNA (Figure 4D; yellow line), while the DMC1-SSDS signal
291 still shows a relative depletion indicative of rapid repair. We found that all inter-homolog repair
292 products (COs and NCOs) were notably depleted in the earliest replicating DNA, where rapid
293 DSB repair is occurring (Figure 4E, S15). The “missing” repair outcomes likely result from DSBs
294 that use the sister chromatid as a repair template, as inter-sister repair products cannot be

detected (sister chromatids are genetically identical). Inter-sister DSB repair is generally disfavored in meiosis to assure crossover formation between homologs. Nonetheless, this inter-homolog bias appears to be gradually established, such that the earliest-forming DSBs can still repair from the sister chromatid (Joshi et al., 2015; Sandhu et al., 2020). Given that we find evidence for inter-sister repair of DSBs and given the fact that this type of repair is only favored at the earliest-forming DSBs in meiosis, the implication of our findings is that DSBs at the earliest replicating DNA are also formed early in meiosis. Other correlates of DNA replication, such as GC-content or chromatin structure lack any mechanism that would link them to early-forming DSBs. Therefore, the most parsimonious conclusion is that there exists a direct link between replication and recombination in male mouse meiosis.

Chromosome-scale regulation of recombination is coupled to DNA replication

Chromosome-scale regulation of recombination is a PRDM9-independent aspect of recombination patterning. For example, short chromosomes, by virtue of their size, are statistically less likely to receive a crossover-competent DSB, yet mechanisms exist to assure that they do (Duret and Galtier, 2009; Kaback et al., 1992; Murakami et al., 2020). In mouse males, we find that short chromosomes have a higher crossover density (Figure 5A), however much of the variance in crossover density remains unexplained. We hypothesized that DNA replication may play a role because origin density varies greatly among mouse chromosomes; for example, origin density on chromosome 13 (120 Mb; 327 origins) is three times lower than on chromosome 11 (122 Mb; 1,023 origins). Indeed, we found that origin density is positively correlated with crossover density (Figure S16). Origin density is a crude estimate of replication activity that does not account for stochastic origin firing or competition between chromosomes for limiting firing factors. We therefore derived a more integrated metric of replication from RT-Sim models that measures how quickly chromosomes are replicated relative to each other (replication speed; Figure 5B, S16). Alone, replication speed predicts crossover density substantially better than origin density (Figure 5C, S16) and when coupled with chromosome size in a multiple linear regression model can explain 90% of the per-chromosome variance in crossover density (Figure 5D, S16). Strikingly, this implies that the recombination potential of chromosomes is mostly established before DSB formation.

In yeast, compensation for chromosome size is achieved by increasing the DSB density on short chromosomes (Murakami et al., 2020), but such an increase is not so apparent in mice

(Lange et al., 2016). Whether differences in DNA replication modulate DSB formation or alter DSB repair outcomes remains to be explored.

Distinct sub-telomeric patterning of DNA replication in the human male germline

In human males, like in mice, meiotic DSBs exhibit megabase-scale correlations across the genome (Figure 6A). An overt manifestation of this phenotype is seen at the ends of human chromosomes, where both DSBs (Pratto et al., 2014) and crossovers (Coop et al., 2008) are grossly enriched in males, independent of PRDM9 genotype (Figure 6B).

We isolated S-phase nuclei expressing meiotic markers from human testes (Figure S17) and inferred replication timing. Strikingly, we found that sub-telomeric regions of human chromosomes replicate early and that therefore, germline RT is highly correlated with DSB hotspot density ($R^2 = 0.75$) (Figure 6B-C). This germline-specific RT patterning may explain why only a weak link was previously found between RT in a lymphoblastoid cell line (LCL) and meiotic recombination (Koren et al., 2012). The distal pattern of early replication in male meiosis appears highly distinct, as sub-telomeric DNA does not replicate notably early in any of the other cell types we studied (Figure 6C, S18). Intriguingly, GC-content is also elevated in sub-telomeric DNA (Figure 6C,D); thus, genomic GC-content is better correlated with RT in the germline than in other cell types (Figure 6D). Importantly, the earliest meiotic DSBs in human males are detected almost exclusively in sub-telomeric DNA (Pratto et al., 2014). Thus, early DNA replication in distal regions in the germline underlies the spatio-temporal patterning of meiotic recombination in human males.

Discussion

The single round of DNA replication that begins meiosis has remained an enigma, despite huge advances in our ability to interrogate nuclear processes. Aside from a cadre of classical papers in the 1970s, few studies have addressed the intricacies of meiotic DNA replication in mammals at a molecular level. We have addressed this shortcoming by developing a comprehensive framework for the study of DNA replication *in vivo* and have used this to generate the first major advances in the study of DNA replication in mouse and human meiosis in decades.

A major hurdle to overcome was to map origins of replication in the few S-phase cells that are found in the mammalian testis. Existing protocols to map origins are designed for the controlled

environment of cell culture, typically require tens to hundreds of millions of cells, and have not been performed in tissue. By adapting methods for nascent-strand sequencing to retain the directionality of captured strands, we generated a high-confidence map of ~11,500 origins of replication in the testis. This is also the first map of *in-vivo* replication origins in a mammal and paves the way for future studies in other cell types. Replication origins appear to be determined by the combination of open chromatin and CpG density and are unevenly distributed in the genome. CpG islands in open chromatin are broadly conserved across cell types, and indeed, we find that the origins identified in testis represent a core set of cell-type-agnostic replication origins. This does not negate the possibility of cell-type specific origins, as have been reported in other studies (Smith et al., 2016).

Despite using the same population of replication origins, we exposed fundamental differences in S-phase duration among cell-types through *in-silico* modelling. Importantly, experimental methods to infer S-phase duration are laborious and imprecise or are limited to cell culture systems (Grant et al., 2018). Meiotic S-phase has been suggested to be notably long in a variety of organisms, including animals, plants, and yeast (Bennett et al., 1972; Callan, 1973; Cha et al., 2000; Holm, 1977). Experimental estimates of S-phase duration in mice are 14-15 hr and 29-30 hr in Spermatogonia-B and Spermatocytes, respectively (Ghosal and Mukherjee, 1971; Monesi, 1962). In agreement, we found that meiotic S-phase in mice is approximately 1.8 times longer than the germ-cells that immediately precede meiosis. The similarity between *in-silico* and experimental estimates implies that the assumptions of our model are reasonable; this includes the assumption that replication fork speed is similar across cell-types. To accurately model replication, a limiting factor that caps the number of active replisomes is strictly required ((Gindin et al., 2014) & data not shown). We found that this replisome “ceiling” differs among cell types, and, in turn, modulates S-phase duration. Importantly, such estimates of the number of active replisomes per cell are extremely challenging to obtain experimentally (Chagin et al., 2016). What components of the replication machinery govern this “ceiling” remain unknown, but investigating the factors that slow down meiotic S-phase may help identify genes that modulate S-phase duration more generally. In yeast, the meiosis-specific cohesin Rec8p and DSB-forming protein Spo11p appear to play such a role. However, we found no evidence for a role of SPO11 in regulating S-phase duration in mouse meiosis. In addition to a lowered replisome “ceiling”, other, more complex mechanisms may contribute to slowing S-phase in meiosis. Our model only allows alteration of the global properties of replication, however

387 regional modifiers of replication fork speed such as replication slow zones (Cha and Kleckner,
388 2002) or common fragile sites (Smith et al., 2006) may impede fork progression differentially in
389 meiosis and other cell types.

390 Our tripartite approach to describe and parameterize DNA replication enabled us to ask if
391 replication influences recombination in mammalian meiosis. We found an extremely strong
392 positive correlation between multiple metrics of recombination and early replicating DNA. The
393 earliest measure we examined was the binding of the PRDM9 protein, where we found that the
394 preference for DSB formation in early replicating DNA was already established. One possibility
395 therefore is that opportunistic binding of PRDM9 to more accessible chromatin in early
396 replicated DNA establishes a link between replication and recombination. However, we found
397 that recombination was enriched in early replicating DNA even in the absence of functional
398 PRDM9. This implies that broad-scale patterning of recombination acts independent of the
399 factors that determine the local patterning of meiotic DSBs.

400 We found strong evidence for the rapid repair of meiotic DSBs in early replicating parts of the
401 genome. The rapid repair of DSBs is a correlate of crossover-biased resolution (Hinch et al.,
402 2019), however, we find that all interhomolog repair outcomes are depleted in these regions
403 suggesting increased use of the sister-chromatid as a repair template. In meiosis, DSB repair
404 with the sister chromatid is strongly disfavored. This assures that recombination occurs with the
405 homologous chromosome - a requirement for allelic exchange and accurate chromosome
406 segregation. At least in yeast, this inter-homolog bias is established gradually, and inter-sister
407 recombination is preferred only for the earliest DSBs that form in meiotic prophase (Joshi et al.,
408 2015; Sandhu et al., 2020). We therefore infer that the earliest DSBs occur in the earliest
409 replicating DNA, strongly implicating a mechanistic link between replication timing *per se* and
410 DSB formation. This would also imply that correlates of replication timing such as GC-content,
411 genome compartmentalization, heterochromatin, or gene density, which lack any temporal
412 component, may indirectly affect recombination by modifying replication patterns. In human
413 males, cytological evidence has shown that the earliest meiotic DSBs occur almost exclusively
414 in sub-telomeric DNA (Pratto et al., 2014). This coincides with the earliest replicating DNA,
415 substantiating the direct link between replication and recombination initiation. It remains
416 possible that there are two phenomena that modulate recombination at large-scales in the
417 genome; one that temporally favors early-forming DSBs in the wake of a passing replication fork

418 (like in yeast), and another that favors recombination in the resultant “permissive” chromatin
419 environment. Of course, since DNA replication establishes the 3D structure of the genome
420 (Klein et al., 2019), these two effects may be one-and-the-same.

421 It is well established that chromosome size partly determines per chromosome recombination
422 rates (Duret and Galtier, 2009; Kaback et al., 1992; Murakami et al., 2020). In addition, we
423 found that in mouse males, the per-chromosomes origin density has a similar predictive value to
424 chromosome size. Importantly, the predictive power of chromosome size and replication are
425 additive, which implies that these properties represent uncorrelated aspects of the mechanisms
426 that control per chromosome recombination rates. Together almost all of the per chromosome
427 variation in crossover density can be explained by these two properties, suggesting that
428 recombination will follow a deterministic path, established before DSBs are made. Short
429 chromosomes likely benefit from an elevated recombination rate to assure crossover formation,
430 however it is less intuitive to understand if or why elevated recombination on fast replicating
431 chromosomes would be beneficial. Fast-replicating chromosomes are origin rich, GC-rich,
432 replicate earlier, and have elevated gene density. A mechanism that links replication to
433 recombination would assure that such regions benefit from recombination to break linkage
434 blocks, generate diversity and purge deleterious mutations.

435 Any benefit of recombining in early replicating DNA could only manifest in the germline (where
436 meiotic recombination occurs) and intriguingly, in human males, germline DNA replication
437 mirrors GC content more closely than replication in other cell types. Thus, during development,
438 we propose that replication may follow a well-charted course, dictated by the underlying DNA. In
439 contrast, the commitment to differentiation in other cell types may render replication more
440 susceptible to epigenetic regulation (Hiratani et al., 2010). In being closely linked to GC content,
441 DNA replication may act as a “selectable trait” that influences the evolution of the genome. For
442 example, should a chromosome lack sufficient recombination capacity in meiosis, there would
443 be a strong selective advantage to increasing GC-content, leading to earlier replication and
444 hence, elevated DSB potential.

445 Errors in DNA replication, if unrepaired, can alter the genome, but will only be transmitted to the
446 next generation if they occur in the germline. Our finding that germline replication is substantially

different therefore has important implications for understanding the patterning of *de-novo* variation in the genome and its impact on population genetic structure.

Acknowledgments

We acknowledge all members of the Camerini-Otero lab for helpful discussion and comments. We also thank Galina Petukhova, Michael Lichten and Peggy Hsieh for critiquing the manuscript. We thank the Flow Cytometry Core of NHLBI, the NIDDK Laboratory of Animal Sciences Section and the NIDDK genomics core facility. This work utilized the computational resources of the NIH HPC Biowulf cluster (<http://hpc.nih.gov>). This work was funded by an NIGMS grant to P.W.J. (R01GM11755) and by the intramural program of the NIDDK (R.D.C.O.). We also thank the Washington Regional Transplant Community for their assistance in obtaining de-identified human testis donations for research.

Author Contributions

Conceptualization, F.P., K.B. and R.D.C.O.; Methodology, F.P., K.B., G.L. and G.C.; Investigation, F.P., K.B., G.L., D.D., J.C. and G.C.; Formal Analysis, F.P., K.B.; Software, K.B.; Visualization, F.P., K.B.; Writing - Original Draft, F.P. and K.B.; Writing - Review & Editing, F.P., K.B. and R.D.C.O.; Resources, S.R.W. and P.W.J.; Funding Acquisition, P.W.J. and R.D.C.O.; Supervision, R.D.C.O.

Declaration of Interests

The authors declare no competing interests.

Data & code availability

The sequencing data have been deposited at the GEO (accession number GSE148327, reviewer token XXXXXXXX). Custom pipelines and code are available at github: <https://github.com/kevbrick/prattoEtAlAnalyticPipeline.git>
<https://github.com/kevbrick/RTSeqPipeline.git>

471 <https://github.com/kevbrick/RTsim.git>

472 **Figures**

473 **Figure 1. Identification of replication origins**

474 **(A)** Schematic of the Ori-SSDS protocol to sequence nascent leading strands. **(B)** The
 475 Ori-SSDS signal in a typical 1 Mb region. Insets show the strand-specific Ori-SSDS signal at
 476 representative weak (inset 1) and strong (inset 3) origins and at a series of non-origin peaks
 477 (inset 2) that lack Ori-SSDS Watson-Crick asymmetry; red = Watson-strand coverage, blue =
 478 Crick-strand coverage). Coverage is calculated in 1 Kb windows with a 147 bp step. **(C)**
 479 Ori-SSDS signal at origins is reproducible but is lost in control experiments. The $\log_2$ ratio of
 480 Watson/Crick strand ssDNA fragments is shown, smoothed in 1 Kb windows. **(D)** Origin
 481 efficiency is highly correlated among replicates. Origin efficiency varies ~100-fold. Origins found

482 only in one replicate are colored purple and green, respectively. **(E)** Origins of replication occur
483 predominantly at CpG islands (CGIs) that coincide with open chromatin as measured by
484 ATAC-Seq (Maezawa et al., 2018). **(F)** Origins are only detected at about half of the CGIs at
485 ATAC-Seq peaks.

Figure 2. Replication timing in meiotic prophase I of male mice

(A-C) Meiotic S-phase and spermatogonia nuclei were isolated using fluorescence-activated nuclei sorting (Lam et al., 2019). (A) Replicating nuclei were isolated by gating 2-4C nuclei by DAPI content. (B) Meiotic S-phase nuclei (Meis) can be distinguished from S-phase nuclei of pre-meiotic germ cells using positive selection for nuclei expressing STRA8, and negative selection for DMRT1. This sorting paradigm can also distinguish three sub-types of germ cells (undifferentiated spermatogonia, SpgU; intermediate spermatogonia, Spgl; type-B spermatogonia, SpgB) (C) SYCP3 is a meiosis-specific protein that was not used for nuclei sorting. SYCP3 is elevated in the isolated meiotic nuclei relative to the other populations. Above-background SYCP3 is also seen in the SpgB population. These are the germ-cells that immediately precede meiotic entry. (D) Replication timing (RT; log₂ of normalized sequencing coverage (cov_n)/genome median) in meiotic S-phase nuclei correlates with origin locations in

498 testis. **(E)** RT in E14 embryonic stem cells. **(F)** RT is highly correlated across cell-types. RT was
 499 inferred from published whole genome sequencing data for Spermatogonial Stem Cells (SSC),
 500 Primordial Germ Cells (PGC), Embryonic Stem Cells (ESC) CD8⁺ cells (CD8), and activated
 501 B-cells (Bcell). Pre-processed RT data was obtained for Myoblast and Lymphoblastoid Cell
 502 Lines (LCL). Details of samples in Tables S1-3. **(G)** GC-content (in 10Kb windows; shown as
 503 $\log_2(\text{GC}/\text{meanGC})$; smoothed in 1Mb windows, 10Kb steps) and genome compartmentalization
 504 at zygonema (Patel et al., 2019) also correlate with RT. Hi-C track shows the eigenvector values
 505 for the first principal component of the Hi-C matrix (calculated in 100 kb windows, smoothed in
 506 1Mb windows, 10Kb steps). Active (A) and inactive (B) compartments are highlighted.

Figure 3. *In silico* modeling recapitulates RT from origin locations

(A) Best fitting RT-Sim model for Meiotic S-phase RT (MeiS(a)). Experimentally determined RT is shown as filled area. Simulated RT is the black line. Both simulated and experimental RT are normalized by mean and standard deviation (s.d.) ($\text{RT}_{\text{norm}} = (\text{RT} - \text{mean}(\text{RT})) / \text{s.d.}(\text{RT})$). **(B)** Fit scores for models built using RT from different cell types and different datasets as a proxy for origins. Mean of the top 0.015% of models is shown. Samples are described in detail in Tables S1-3. Briefly, RT is from: MeiS = Meiotic S-phase, Spg(B,I,U) = B-type, Intermediate and Undifferentiated Spermatogonia, ESC = Embryonic Stem Cells, SSC = Spermatogonial Stem Cells, PGC = Primordial Germ Cells, CD8 = CD8 cells, Myoblast = Myoblast cell line, LCL = Lymphoblastoid cell line, Bcell = Activated B-cells. (a,b,c) designate replicates. ESC(a) is from published whole genome sequencing data, ESC(b) is from cells grown by us. Ori = OriSSDS origins, Ori(r) = OriSSDS origins with randomized genomic location, AS+CGI = ATAC-Seq peak at a CGI, AS-CGI = ATAC-Seq peak not at a CGI, CGI-AS = CGI not at an ATAC-Seq peak. **(C)**

520 Best fitting RT-Sim model for RT in E14 ESCs (ESC(a)). **(D)** Sorting schematic for S-phase
 521 populations in lower panel. Populations have increasing DNA content from T1 to T4. The
 522 optimal simulation run-time for best-fitting models correlates with increasing DNA content.
 523 Nonetheless, the predicted S-phase duration is similar among all populations. **(E)** Total
 524 replication time for best-fitting models. Meiotic S-phase is notably slow. **(F)** Best fitting models
 525 imply that meiotic S-phase is slow because fewer replisomes fire than in other cell types.
 526 Activated B-cells appear to share this property of replication with meiotic S-phase.

Figure 4. Rapid DSB repair and elevated meiotic recombination in early-replicating regions

(A) Schematic of meiotic recombination. The colored lines in B-D reflect the colors in this schematic. PR9 = PRDM9; S11 = SPO11. (B,D) PRDM9-mediated H3K4m3 at DSB hotspots (yellow) is enriched in early replicating DNA in both wild-type B6 male mice ($B6^{wt/wt}$; B) and in mice with a humanized *PRDM9* allele ($B6^{h/h}$; C). DSB formation (SPO11-oligo mapping; pink) follows this Pre-DSB mark (B). The DMC1-SSDS (green) signal decays relative to H3K4me3 in the earliest replicating DNA. (C) B6 PRDM9 binding sites are depleted in early replicating DNA (measured by Affinity-Seq) in the $B6^{wt/wt}$ genome. This is indicative of hotspot erosion. (E) Inter-homolog repair products (crossovers + non-crossovers; (Li et al., 2019a)) are depleted in the earliest replicating DNA relative to DSB-associated H3K4m3 (Note that H3K4m3 are the same data as in D, plotted against RT in B6xCAST). Simulated RT from the T1 Meocyte population (Figure 3D) is used for B,C and D. Simulated RT from the Meiotic S-phase in B6xCAST F1 mice is used for E. Solid lines depict the LOESS smoothed signal $\pm$ standard error

541 (shaded). For all panels, dots represent the average signal from all autosomal bins in the
542 genome for each RT quantile ($N_{\text{bins}} = 250$).

Figure 5. DNA replication predicts per-chromosome crossover rates

(A) A linear regression model can predict crossover density from chromosome size (polynomial fit). Crossover density is calculated as total crossovers per Mb. Crossovers from (Yin et al., 2019). R^2 is strongly influenced by the single small outlier chromosome (chr19; rightmost). (B) Chromosomes replicate asynchronously. Chromosome 11 and 13 are highlighted because while they are a similar size, chromosome 11 replicates far earlier. The time required to replicate 75% of each chromosome is inferred from the best-fitting RT-Sim model (see also Figure S16). Boxplots indicate the range of times from individual modelled cells. (C) A linear regression model can predict crossover density from replication speed. (D) Per chromosome crossover

552 density can be accurately predicted from a linear model that combines replication speed and
553 chromosome size.

Figure 6. Sub-telomeric DNA replicates early in human male meiosis

(A) Extensive large-scale correlation in DSB density in human males with different PRDM9 genotypes (PRDM9^{A/A} homozygotes (A/A) and PRDM9^{C/L4} (C/L4) heterozygote). At fine scale (10 Kb), there is no correlation between DSB maps in the A/A and C/L4 individuals because

DSB hotspot locations differ. Comparing the maps at lower resolution reveals extensive correlation at megabase scales. **(B)** Meiotic DSBs are enriched in sub-telomeric regions in human males, independent of the *Prdm9* genotype. **(C)** Subtelomeric DNA replicates consistently early in human male meiosis but not in mitotic (RPE-1; Table S3) cells. Boxplots depict the range of replication timing values in 2 Mb regions across the genome; the median is shown as a grey bar and the box designates the interquartile range. The mean (red dot) $\pm$ 1 standard deviation (filled shadow) is also shown for each interval. GC content is also elevated in sub-telomeric DNA. **(D)** RT in the germline (all meiotic samples) correlates better with genomic GC content than RT in other cell types.

567 **Supplementary Figures**

569 **Figure S1. RNA hydrolysis abolishes the Ori-SSDS signal.**

570 RNA hydrolysis prior to lambda exonuclease treatment abolishes Watson/Crick asymmetry at
571 origins of replication. This is because RNA hydrolysis will remove the nascent strand RNA
572 primer and thus render nascent-strands susceptible to lambda exonuclease digestion (see
573 methods). Two replicate experiments were performed in parallel; WT_Rep4 & WT_Rep5 (see
574 methods) **(A,B,C)** The strand asymmetry of Ori-SSDS is lost following RNA hydrolysis (b,c

575 RNAh+ samples). Residual enrichment at origins, without Watson/Crick asymmetry is still seen.
576 This is likely because there is more DNA in the population at these sites as a result of DNA
577 replication. Upper panels in B,C depict the average Ori-SSDS signal at origins of replication for
578 Watson (red) and Crick (blue) strand reads.

Figure S2. Ori-SSDS signal at putative g-quadruplexes in replicating cells.

G-quadruplexes (G4s) are secondary structures that have been hypothesized to impede lambda exonuclease *in vitro* and may result in SNS-Seq peak artifacts (Foulk et al., 2015). Indeed, we detect small peaks at putative G4 sequences in Ori-SSDS data. Figures depict the average Ori-SSDS signal at origins of replication or predicted G-quadruplexes for Watson (red) and Crick (blue) strand reads. Ori-SSDS sequencing can distinguish G4-signals from origin-derived signals because G4s yield ssDNA on just one strand whereas reciprocal Watson/Crick asymmetry is seen around the center of origins. This differentiates Ori-SSDS from other SNS-Seq methods, which cannot distinguish G4s from origins. At origins that coincide with

588 putative G4s (Huppert and Balasubramanian, 2005) it is not possible to resolve these signals
 589 however, since the G4 signal is far weaker than that at origins, the contribution of G4s to true
 590 origin signal is a minor concern. Curiously, the signal at G4s is not seen in non-replicating tissue
 591 (sperm) or in samples where the RNA-primer is degraded. Thus, the G4-associated signal is
 592 replication-dependent and not simply the result of G4 resistance to lambda exonuclease. The
 593 stranded-ness of the signal is consistent with the capture of ligated Okazaki-fragments at the G4
 594 sites. This implies that G-quadruplexes are an impediment to fork passage *in-vivo*.

Figure S3. SNS-Seq peaks occur in clusters at individual origins of replication.

(A) Without knowledge of strand-asymmetry, SNS-Seq experiments have historically overestimated the number of origins in the genome. At this locus, Ori-SSDS reveals that 6 SNS-Seq origin calls likely represent a single origin of replication. Grey tracks represent three replicate SNS-Seq coverage tracks (Cayrou et al., 2015). Green boxes underneath the grey tracks represent SNS-Seq origin calls (from GEO: GSE68347_Initiation_Sites.bedGraph.gz; accession GSE68347; UCSC liftover was used to convert mouse mm9 to mm10 genome coordinates) (Cayrou et al., 2015). Ori-SSDS coverage from WT Rep 1 is shown. The green box under the Ori-SSDS track represents the single Ori-SSDS origin call in this region. This is validated by the characteristic Watson-Crick asymmetry. (B) We obtained SNS-Seq origin of replication peak calls (origins) for mouse (described in panel a.; (Cayrou et al., 2015)) and for human (from GEO; GSE134988_siNC-NS_peaks.bed.gz; accession GSE134988) (Long et al., 2020). For each dataset, origins were extended ± 1.5 Kb from the center-point and merged into origin "clusters"; clusters >10 Kb were discarded. We posit that each of the remaining clusters represents a single true origin of replication. Most SNS-Seq peaks occurred in a cluster. Just 37% and 29% occurred at isolated origins in mouse and human, respectively. (C) The number

611 of “origins of replication” from SNS-Seq studies is substantially reduced if clusters are
612 considered as single origins.

613 **Figure S4. Most origins are detected in multiple replicate experiments**

614 Proportionate venn diagram depicting the number of origins detected in each of three replicate
615 Ori-SSDS experiments and the overlaps between sets. Our consensus set of origins includes all
616 origins found in at least two Ori-SSDS experiments.

617 **Figure S5. Ori-SSDS gives a reproducible readout of origin efficiency**

618 Origin efficiency is a measure of the frequency at which each origin is used. This is inferred from
619 the sequencing read coverage in each experiment (see methods). Grey dots represent origins
620 found in both samples. Magenta and green dots represent origins found only in the x-axis
621 designated and y-axis designated samples, respectively. The top panels show all origins found
622 in either experiment for each pair-wise comparison. The bottom panels show only the
623 consensus set of 11,565 origins. R^2 is the squared Spearman correlation coefficient of log
624 transformed values. N indicates the number of origins for each pairwise comparison.

Figure S6. Ori-SSDS peaks coincide with replication measures from other experiments.

(A) Okazaki fragments are distributed asymmetrically around Ori-SSDS-defined origins of replication. Okazaki fragment sequencing (Ok-Seq) data in Embryonic Stem Cells (ESCs) are from (Petryk et al., 2018). The Okazaki fragment signal is far broader than the signal from Ori-SSDS because it captures all lagging-strand synthesis (for details, see (Petryk et al., 2016)); Ori-SSDS captures only RNA-primed leading strands at the origin of replication. Note that Ok-Seq exhibits the opposite polarity to the Ori-SSDS signal as expected from capture of lagging and leading strands, respectively. Data are shown in 50 Kb windows. Heatmaps depict the signal at individual Ori-SSDS defined origins. (B,C) Ok-Seq signal at subsets of origins of replication. For reference, the Ok-Seq signal at Ori-SSDS defined origins is shown in three panels (A,B,C). The heatmaps in panels B & C are scaled to reflect the number of origins in each subset. There are three genome-wide maps of origins of replication in mice. All three maps were generated using SNS-Seq. One map is derived from Mouse Embryonic Fibroblast (MEF)

638 cell culture (Almeida et al., 2018) and two from ESC culture (Almeida et al., 2018; Cayrou et al.,
639 2015). **(B)** Origins defined in SNS-Seq experiments that coincide with Ori-SSDS peaks exhibit
640 similar Ok-Seq asymmetry. Data for origins common to the two SNS-Seq experiments in ESCs
641 are also shown (ESC (x2)). **(C)** A large proportion of peaks detected in SNS-Seq experiments,
642 but not in Ori-SSDS appear to be false-positives as they lack Ok-Seq asymmetry. This is more
643 pronounced in the Almeida experiments than in the Cayrou experiment. **(D)** Most origins of
644 replication from Ori-SSDS in testis coincide with origins mapped in other cell types. Origins from
645 the three SNS-Seq experiments were combined to assess the overlap with “Any” origins defined
646 previously.

Figure S7. Origins of replication in mouse testis occur at GC-rich, accessible sites.

(A) Origins of replication are more closely spaced than expected in the genome. The expected distances between randomly chosen intervals is shown in grey. **(B)** Most origins occur near gene promoters; 66% occur within 1 Kbp of a transcription start site (TSS), far more than expected (random; grey line). **(C)** Replication origins coincide with local changes in DNA bendability (calculated from (Goodsell and Dickerson, 1994)). The mean (top) and standard deviation (bottom) of the DNA bendability at all origins is shown. Each point represents a 100 bp window. Solid line is a LOESS fit. **(D)** Origin centers are GC-rich and enriched in CpG and GpC dinucleotides. The $\log_2$ ratio of the central 500 bp compared to the region flanking this center is shown. **(E)** GC-content (GC%), CpG density and GpC density correlate positively with origin efficiency. **(F)** Putative G4-forming sequences are highly enriched at the center of origins. 68% of origins have a putative G4-forming sequence in the central $\pm$ 1 Kb. **(G)** Origin locations can be identified from CpG density. CpG dinucleotide density was evaluated in all 1 Kb regions in the genome using a 100 bp sliding window. Intervals at each CpG threshold were expanded $\pm$ 100 bp around the center and overlapping windows were merged. The centerpoint of each merged interval was defined as a CpG peak. CpG peaks with CpG content $\geq$ 7% overlap 6,233 origins (53% of total). 1,853 CpG peaks at this threshold do not occur at an origin (FP). **(H)** Origins are better predicted by CpG density than transcription factors (TFs) are by transcription factor binding sites (TFBS). Receiver operating curve (ROC) for predicting origin locations using CpG peaks (green) or for predicting peaks in ChIP-Seq datasets for 179 transcription factors (grey). For origins, at each CpG threshold (1% to 15% in 1% steps), peaks were called (as in G). All mouse TFs with ChIP-Seq data (in GTRD database; (Yevshin et al., 2017)) and with a designated TFBS in the HOCOMOCO database (Kulakovskiy et al., 2018) were used. SPRY-SARUS (Kulakovskiy et al., 2018) was used to identify TFBS motifs genome-wide with $p < 0.005$. For each TF, the score at which $\leq$ 12,000 peaks were found was used as a lower threshold (this is to assure a fairer comparison with origins). Motifs with scores less than this were discarded. TFs with $< 12,000$ peaks not used. Hits inside ChIP-Seq peaks were considered true positives.

675 **Figure S8. Deconvolution of closely spaced origins of replication**

676 **(A)** Origin of replication cumulative width distribution. **(B)** Peaks of CpG density coincide with
 677 origin centers. Only origins narrower than 6 Kb were considered. CpG dinucleotide density was
 678 evaluated in all 1 Kb regions in the genome using a 100 bp sliding window. Intervals with >5%
 679 CpGs were expanded ± 100 bp around the center and overlapping windows were merged. The
 680 centerpoint of each merged interval was defined as a CpG “peak”. **(C)** Almost all CpG peaks
 681 within broad initiation zones (origins >10 Kb) exhibit Ori-SSDS Crick-Watson asymmetry. This
 682 strongly implies that these are individual origins of replication. **(D)** Local CpG peak counts as a
 683 function of origin width. **(E,F)** Examples of a cluster of initiation zones. Peaks of CpG density
 684 coincide with Crick/Watson asymmetry at many loci, revealing extensive sub-structure. This
 685 locus likely contains many discrete origins of replication that are difficult to separate *in silico*.
 686 The Hox locus is shown in (E). **(G)** Two adjacent initiation zones with different properties. The
 687 zone on the left is 13 Kb wide, yet contains a single, clear CpG peak that coincides with
 688 Ori-SSDS asymmetry. This is likely a single replication origin. The zone on the right is also 13
 689 Kb, but contains no discernable CpG peak or origin. This may represent a diffuse initiation zone
 690 or a cluster of weak origins. **(H)** Four initiation zones with varying properties.

691 **Figure S9. High purity populations of sorted meiotic S-phase nuclei**

692 **(A)** Immunofluorescence microscopy images of sorted nuclei from the Meis population (same
693 population as in Fig. 2C; 40X magnification). Nuclei were concentrated in a small volume after
694 centrifugation. The nuclei suspension was then pipetted onto silane coated slides and mounted

695 with Vectashield. 49 nuclei are shown of which 46 are preleptotene nuclei (STRA8 positive,
 696 DMRT1 negative, weak and diffuse SYCP3 signal and DAPI morphology consistent with
 697 preleptotene cells (Bellve et al., 1977)). The magenta arrow points to a Sertoli cell nucleus and
 698 green arrowheads point to nuclei of unknown type. Nucleus types are indicated by a letter code
 699 superimposed on each cell of the DAPI-stained image (top-left). **(B)** Details of three
 700 preleptotene nuclei at higher magnification.

701 **Figure S10. RT-Seq yields similar RT profiles across varied cell-types**

702 RT for all cell types shown on Figure 2. RT is expressed as the $\log_2$ enrichment over genomic
 703 median coverage for all panels except for Myoblast and LCL data. These data were obtained in

704 processed form from the Replication Domain database. All panels depict chromosome 12.
705 Sample names are explained briefly in Figure 3 and in detail in Tables S1-3.

Figure S11. Experimentally derived RT varies as a function of the starting population

Four sub-populations of meiotic S-phase were isolated from early S (T1) through late S (T4). RT was inferred from WGS (see methods). The earliest and latest replicating regions exhibit substantial differences in RT across these four populations.

710 **A**

711 **B**

712 **Figure S12. Modelling recapitulates experimental RT genome-wide**

713 Experimental RT is shown as filled green (early replicating) and violet (late replicating) regions.
 714 The best-fitting RT-Sim predicted RT is shown as a solid black line. Per-chromosome Pearson
 715 correlation coefficients are shown. Both **(A)** Meiotic and **(B)** E14 ESC RT can be accurately
 716 modeled *in silico*. Both experimental and predicted RT are normalized by mean and standard
 717 deviation (s.d.); $RT_{norm} = (RT - \text{mean}(RT)) / \text{s.d.}(RT)$.

Figure S13. ATAC-Seq peaks at CpG islands are broadly conserved across diverse cell-types.

We called peaks from 8 published ATAC-Seq experiments in a variety of mouse cell types. The strongest 12,000 peaks were split into 4 groups by strength. We counted the overlap of each peak subset with the top 12,000 ATAC-Seq peaks from each other cell-type. The percentage of peaks from each subset (y-axis) found in other cell types (x-axis) is shown. Datasets used are as follows: Hepatocytes (Hep/cytes; SRA accessions: SRR6813698,SRR6813699)(Li et al., 2019b), Differentiated cKIT+ Spermatogonia (SpgD; SRA accessions: SRR5956508)(Maezawa et al., 2018), Undifferentiated THY+ Spermatogonia (SpgU; SRA accessions: SRR5956504)(Maezawa et al., 2018), Pachytene Spermatocytes (Pachy; SRA accessions: SRR5956512)(Maezawa et al., 2018), Mouse Embryonic Fibroblasts (MEF; SRA accessions: SRR7048429, SRR7048430), Hindlimb Muscle Cells E14.5 (Muscle (HL); SRA accessions: SRR8104383,SRR8104391)(Castro et al., 2019), Embryonic Stem Cells (ESC; SRA accessions: SRR7048437,SRR7048438), Embryonic Stem Cells - 6 Days (ESC (Day 6); SRA accessions: SRR7048433,SRR7048434).

733 **Figure S14. Meiotic S-phase duration is unaffected in *Spo11*^{-/-} male mice**

734 Meiotic S-phase nuclei, B-type spermatogonia and undifferentiated spermatogonia were
735 isolated from a *Spo11*^{-/-} male mouse testis (SPO11 in sample names) and a *wild-type* littermate
736 (wtLM in sample names). Details for each sample are in Tables S1-3. **(A)** 2-4C nuclei were

737 sorted and **(B)** gated by DMRT1 and STRA8 immunofluorescence signal. Gates were defined
 738 while sorting nuclei from the *Spo11*^{-/-} mouse and retained, unchanged for sorting nuclei from the
 739 littermate. **(C)** Summary statistics for best-fitting RT-Sim models for all samples. **(D)** Comparison
 740 of S-phase duration across cell types. The median S-phase duration of good models for each
 741 cell-type is compared to meiotic S-phase. **(E)** The “fit” of RT-Sim models for *Spo11*^{-/-} samples is
 742 comparable to the fit in other cell types. **(F)** The duration of meiotic S-phase is unaffected in
 743 *Spo11*^{-/-} mice relative to the *wild-type* littermate control. Notably, both mice exhibit slightly
 744 shorter S-phase duration compared to the three *wild-type* replicates. This is likely because for
 745 the *Spo11*^{-/-} and *wild-type* littermate, the sorting gates were defined in the *Spo11*^{-/-} mouse and
 746 left unchanged when sorting nuclei from the littermate. Cellularity differences from the loss of
 747 *Spo11* may result in the capture of a slightly different population than when sorting gates are
 748 defined using *wild-type*. **(G)** The firing rate is similarly unaffected.

749 **Figure S15. Rapid repair of DSBs in early replicating regions**

750 SSDS using DMC1 exhibits a dip in the earliest replicating DNA. This is mirrored by SSDS using

751 another ssDNA-binding protein, RPA. Thus, this dip does not simply reflect a property of DMC1

752 binding (two replicates shown). HOP2 is required for strand invasion and DSB repair. In *Hop2*^{-/-}

753 mice, DSBs are not repaired and therefore any effects resulting from repair dynamics should be

754 absent. We find that the dip in the DMC1-SSDS signal in the earliest replicating DNA is

755 abolished in *Hop2*^{-/-} mice and that the DMC1-SSDS curve now only reflects the rate of DSB

756 formation (like SPO11-oligo mapping; pink). Importantly, in mice lacking functional PRDM9, the

757 DMC1-SSDS signal also dips at the earliest replicating DNA. PRDM9 binding site erosion does

758 not affect PRDM9-independent DSB hotspots, therefore the dip is not as deep as in *wild-type*

759 mice, where both erosion and rapid repair play a role. Both crossovers (COs) and

760 Non-crossovers (NCO) are depleted in the earliest replicating DNA. For all figures, dots

761 represent the average signal from all autosomal bins for each RT quantile (N = 250). Simulated

762 RT from the T1 Meiocyte population (B6; Figure 3D) or simulated RT from a B6xCAST F1 hybrid

763 (B6xCAST) is used for RT estimates. The solid lines depict the LOESS smoothed signal ±

764 standard error (shaded). The dashed grey line is a projected linear correlation and the deviation
765 from a linear correlation was manually added as a black range bar. The phenomena contributing
766 to each dip are indicated by the colored circles (see key at top).

Figure S16. Replication speed differs among chromosomes and can predict crossover density.

(A) Some chromosomes replicate faster than others; for example, chromosome 11 and 13 are a similar size but chromosome 11 replicates far earlier. Notably, chromosome 11 has over 3-times

772 more replication origins. The time required to replicate 25% (top), 50% (middle) or 75% (bottom)
 773 of each chromosome is inferred from the best-fitting RT-Sim model. Boxplots indicate the range
 774 of times from 100 individual modelled cells. The median value for each chromosome is used as
 775 a measure of replication speed in subsequent panels. **(B)** We built simple regression models to
 776 predict crossover density from each individual property of chromosomes. DSBs are derived from
 777 DMC1-SSDS experiments in B6xCAS^T F1 mice (Smagulova et al., 2016). Crossovers in
 778 B6xCAS^T are taken from (Yin et al., 2019) and crossover density is calculated as total
 779 crossovers per Mb. Per-chromosome crossover density is best predicted by chromosome size
 780 (polynomial fit). This mirrors findings in other organisms (Murakami et al., 2020). **(C)** Next, we
 781 built a series of linear models of crossover density using chromosome size and each of the
 782 other chromosome properties. Replication speed and chromosome size together are a strong
 783 predictor of per-chromosome crossover density. The best combination (size + Rep.speed
 784 (75%)) yielded a 44% improvement on a model that used size alone and explained 90% of the
 785 variance in per-chromosome crossover density. Origin density and GC-content also improved
 786 the size-only models, but not as much as replication speed. **(D)** 94% of the variance in
 787 per-chromosome crossover density can be explained by adding DSB density to the best model
 788 from (C).

Figure S17. Isolation of human meiotic S-phase nuclei

(A-B) Human meiotic S-phase and spermatogonia nuclei are isolated using fluorescence-activated nuclei sorting (Lam et al., 2019). **(A)** Replicating nuclei were isolated by gating 2-4C nuclei by DAPI content. **(B)** Putative meiotic nuclei were isolated using a combination of DMRT1 and HORMAD1. Antibodies to the STRA8 protein (used for sorting mouse meiotic S-phase nuclei) did not work for human. Unlike in mouse, the DMRT1 protein is expressed in meiotic S-phase **(C-E)**. Thus, some spermatogonia may be present in the MeiS

796 population. **(C)** *t*-Distributed Stochastic Neighbor Embedding (t-SNE) projection of the single-cell
 797 transcription atlas of human testes (Guo et al., 2018). Each dot represents a single cell. Colors
 798 represent defined clusters. **(D)** Quantification of genes-of-interest in four cell clusters. Circles
 799 represent the relative abundance of transcripts. Circle size represents the fraction of cells
 800 showing expression. Depth of color represents average gene expression. **(E)** Single-cell
 801 expression patterns of genes-of-interest. Depth of red color indicates expression level (Red =
 802 high; Grey = low).

Figure S18. Early sub-telomeric replication is distinct to male meiosis

Consistently early replication in subtelomeric regions is only seen in the germline (green or blue) and not in any of the other cell types studied (grey). The pure meiotic S-phase population (green) exhibits the most pronounced signal. In the three samples which likely contain some spermatogonia (blue), early replication in subtelomeric DNA is seen, but less pronounced. This

808 could imply that sub-telomeric DNA does not replicate notably early in spermatogonia. Boxplots
 809 depict the range of replication timing values in 2 Mb regions across the genome; the median is
 810 shown as a grey bar and the box designates the interquartile range. The mean (red dot) $\pm$ 1
 811 standard deviation (filled shadow) is also shown for each interval. Details for each sample are in
 812 Table S1, S3.

813 **Methods**

814 **Tissue sources**

815 C57BL/6J (stock 000664) and CAST/Eij (000928) mice were obtained from The Jackson
816 Laboratory (Maine). Human testicular samples were obtained from a commercial source (Folio,
817 Ohio) that provides testicular biopsies from normal tumor adjacent tissue or from deidentified
818 donor testes from deceased individuals (obtained with the assistance of the Washington
819 Regional Transplant Community).

820 **Mapping origins of replication using short nascent-strand capture followed by** 821 **single-stranded DNA sequencing (Ori-SSDS)**

822 For the short nascent strands (SNS) preparation we adapted the protocol from (Picard et al.,
823 2014) to work with mouse tissue. Mice were euthanized and testes were retrieved. The tunica
824 albuginea was removed and both testes were suspended in 10 ml of DNAzol (Thermo-Fisher
825 catalog number 10503027) and transferred to a Dounce homogenizer. Tissue and cells were
826 disrupted with 5-7 strokes of a loose-fitting pestle. The DNA was precipitated with 5 ml of EtOH
827 and then spooled with a pipette tip and rinsed gently twice with EtOH 75% by sequential transfer
828 into 15 ml tubes. DNA was air-dried for 5 min and then resuspended in 500 µl of 1X TEN buffer
829 (Tris 10mM pH=8; EDTA 1mM; NaCl 100mM) + 80 U of RNaseOUT (Thermo-Fisher catalog
830 number 10777019) (40 U/µl) at 4°C for at least 24h.

831 DNA was denatured at 95°C for 5 min, chilled on ice for 5 min and size-fractionated on 5 - 30%
832 (w/v) linear sucrose gradients in TEN buffer in a Beckman Ultracentrifuge SW45 rotor for 18 h at
833 26,000 g at 20°C. 500 µl fractions were collected from the top of the gradient, precipitated by
834 adding 1 ml of EtOH 100 % and 50 µl of NaOAc 3M pH 5.2 and resuspended in 25 µl Tris Buffer
835 pH=8. Fractions were analyzed in a denaturing (50 mM NaOH, 1 mM EDTA) 1.2% agarose gel
836 and the ones containing ssDNA ranging in size from 800 to 2000 nt were pooled. The sample
837 was then heat-denatured for 5 min at 95°C and chilled on ice for 5 min and DNA was
838 phosphorylated by Polynucleotide Kinase in 150 µl reactions in PNK Buffer (100 U of PNK ; 1X
839 PNK buffer ; 1 mM ATP; 40U RNaseOUT). The reactions were incubated for 30 min at 37°C
840 and heat-inactivated at 75°C for 15 min. DNA was extracted with Phenol/Chloroform, EtOH
841 precipitated and resuspended in 50 µl Tris Buffer pH=8. Samples were heat-denatured again for
842 5 min and digested with lambda-exonuclease in 100 µl of custom lambda buffer (67 mM

glycine-KOH pH= 8.8, 2.5 mM MgCl₂, 50 µg/ml BSA) with 2.5 U to 10 U lambda-exonuclease/µg DNA (Thermo Scientific). 80U RNaseOUT, 5U of PNK and 1 mM ATP were added to the sample and incubated overnight at 37°C. Reactions were inactivated for 10 min at 75°C and DNA was purified with MinElute columns (QIAGEN), as indicated by the instruction manual. RNA primers were hydrolyzed by incubating the eluate for 30 min at 37°C in 0.25 N NaOH, followed by neutralization with acetic acid and purified again with MinElute columns. The sample was then sonicated in a Bioruptor UCD200 for 5 min 30 sec ON 30 sec OFF, “high” setting, and the library was prepared for SSDS as described (Brick et al., 2018b).

For one sample (wt_Rep2), separate libraries were made from three fractions from the sucrose gradient (Fraction 10 : 800-1,000 nt, Fraction 11 : 1,000-1,400 nt and Fraction 12 : 1,400-2,000 nt). Sequencing data were subsequently pooled.

To validate that the Ori-SSDS enrichment stems from RNA primed leading strands, we performed experiments where we hydrolyzed the RNA prior to lambda exonuclease treatment. This also necessitated changes to the above Ori-SSDS protocol as follows. Selected fractions from the sucrose gradient were split in two tubes. In one tube, leading strand RNA primers were hydrolyzed by incubating for 30 min at 37°C in 0.25 N NaOH, followed by neutralization with acetic acid. The other tube was incubated for 30 min at 37°C but in Tris Buffer pH=8. After neutralization, both tubes were heat denatured and spun through a Chromaspin TE-1000 size exclusion column (Clontech) to remove any small degradation products that could interfere with later steps. The eluate was then subjected to PNK treatment and subsequent steps outlined above. Two replicates were performed in parallel, with and without RNA hydrolysis (wt_Rep4, wt_Rep5).

Ori-SSDS in sperm was used as a negative control as there is no DNA replication. Sperm was retrieved from cauda and caput epididymides from 3 adult mice. The mix of somatic cells and sperm was resuspended in 0.1% SDS, 0.5% Triton X to selectively lyse somatic cells. Sperm was recovered by centrifugation and lysed in 6 M guanidinium, 30 mM sodium citrate (pH 7.0), 0.5% Sarkosyl, 0.20 mg/ml proteinase K and 0.3 M Beta-mercaptoethanol, and incubated at 55°C for 1h (Hossain et al., 1997). The DNA was precipitated with 2 volumes of isopropyl alcohol and then spooled with a pipette tip and rinsed gently twice with EtOH 75% by sequential transfer into 15 ml tubes. DNA was air-dried for 5 min and then resuspended in 500 µl of 1X

873 TEN buffer (Tris 10mM pH=8; EDTA 1mM; NaCl 100mM) + 80 U of RNaseOUT (Thermo-Fisher
874 catalog number 10777019) (40 U/μl) at 4°C for at least 24h.

875 **Identification of origins of replication**

876 Sequencing reads from Ori-SSDS experiments were aligned to the reference genome (mouse =
877 mm10; human = hg38) using bwa 0.7.12 and the single-stranded DNA pipeline (Brick et al.,
878 2018b; Khil et al., 2012). Only fragments derived unambiguously from ssDNA (ssDNA_type1)
879 were used in subsequent analyses. ssDNA fragments where either read1 or read2 had mapping
880 quality <30 were discarded. Fragments from the mitochondrial chromosome and fragments in
881 blacklisted regions of the genome were also discarded. Duplicate fragments were subsequently
882 discarded.

883 We used the MACS algorithm for peak calling. To assure that data derived from Ori-SSDS fit the
884 MACS peak model, we performed two operations: 1. The strand orientation of fragments was
885 reversed prior to peak calling. 2. The data was passed to MACS as ssDNA fragment BED files
886 (from SSDS pipeline). Peak calling was performed for each Ori-SSDS sample using MACS
887 2.1.2 and the following parameters : -g (hs or mm) -q 0.001 -extsize 2000 --nomodel
888 --nolambda.

889 The final origin set for mouse was defined by merging origin calls from wt_Rep1,2,3. Origin
890 intervals overlapping by ≥ 500 bp were merged. Origin efficiency was calculated in each sample
891 using a method analogous to that for DMC1-SSDS hotspot mapping (Brick et al., 2018b).
892 Briefly, each origin is re-centred to the midpoint of the Watson and Crick fragment distributions.
893 Signal is estimated as the Crick-strand signal to the left of the origin center + the Watson-strand
894 signal to the right of the origin center. The background is estimated by counting the remaining
895 fragments at the left and right edges of MACS-defined origins (15%); Watson-strand fragments
896 to the left of the origin center, Crick-strand fragments to the right. This background is then
897 extrapolated to the entire origin and subtracted from the signal. Following this process, origins
898 found in just one sample and origins that did not display the expected Crick/Watson asymmetry
899 (Figure 1A) in any sample were discarded.

900 **Whole genome sequencing for replication timing**

901 Nuclei from mouse or human testes were prepared as described in (Lam et al., 2019). To purify
902 the populations of interest, a combination of intranuclear markers were used (Table S1, Figure
903 2).

904 Mouse ESC were obtained from Jackson labs: 129X1/SvJ-PRX-129X1 #1 mES cells derived
905 from day 3.5 blastocysts of strain 129X1/SvJ (Stock Number 000691). The cells were cultured
906 as recommended by Jackson labs. Exponentially growing cells were cross-linked by adding
907 fresh 1% paraformaldehyde solution to the ES media for 10 minutes at room temperature. The
908 cells were washed twice with 1× phosphate-buffered saline and scraped off the plates, pelleted
909 and nuclei were prepared as described in (Lam et al., 2019). Replicating cells were sorted
910 based DNA content.

911 Pure nuclei populations were resuspended in 300 ul of lysis buffer (SDS 1%, 10mM EDTA,
912 50mM, Tris pH=8) and then sonicated in a Bioruptor UCD200 for 20 min 30 sec ON 30 sec OFF,
913 “high” setting. NaCl was added to a final concentration of 0.2M and the sample incubated
914 overnight at 65°C to reverse DNA-protein crosslinks. 5U of Proteinase K (NEB Cat. No.
915 P8107S) was added and the sample further incubated for 1h at 45°C. DNA was purified with
916 MinElute columns (QIAGEN), as indicated by the instruction manual. Whole genome
917 sequencing libraries were prepared from 100 ng of purified DNA using the KAPA Hyper Prep Kit
918 (Cat. No. KK8502) following the manufacturer's instructions.

919 **Inferring replication timing from whole genome sequencing (WGS) data**

920 The mem algorithm of bwa 0.7.12 (Li, 2013; Li and Durbin, 2009) was used to align each WGS
921 dataset to the reference genome (mm10 for mouse; hg38 for human). WGS from a
922 non-replicating population was used to calibrate the expected GC-biases from different
923 sequencing instruments (Illumina HiSeq 2000, Illumina HiSeq 2500, Illumina HiSeq X, Illumina
924 NextSeq 500). Briefly, these calibration files were generated by examining a subset of perfectly
925 mappable genomic loci. Each locus is a 101 bp region with the following properties:

- 926 1. Full mappability of all bases within the 101 bp window. To determine mappability, we
927 generated a fastq file of pseudoreads of the requisite length from the reference genome
928 fasta file. One pseudo-read was generated per bp; all bases were assigned a q-score of

60. Pseudo-reads were mapped to the reference genome using bwa mem 0.7.12 and the coverage at each base in the genome was assessed.

2. The window does not overlap with sequencing gaps (UCSC Gaps track), segmental duplications (UCSC SegDups track), high copy repeats (UCSC repeatmasker track) or otherwise blacklisted genomic regions (Amemiya et al., 2019; Kent et al., 2002).

The samples used to generate GC-correction files were the 2C_{MMC} and 2C_{HSC} samples (Table S1) for Hi-Seq X and Hi-Seq 2500, respectively. Samples used to generate GC-correction files for published data are detailed in Table S2 below.

The GC content (rounded to 1%) and WGS sequencing coverage of each 101 bp window was assessed. The median coverage for all windows of a given GC content was used as a correction factor. These correction factors vary substantially across the different sequencing platforms used. 2% of all qualifying 101 bp autosomal windows were used to calculate the GC correction coefficients.

For each RT-Seq experiment, the WGS coverage in 101-bp windows in the genome was calculated and corrected for GC biases using the correction coefficient for the appropriate sequencing platform. Replication timing was calculated as the log₂ ratio of corrected sequencing coverage at a locus to the genome-wide median (method was adapted from (Koren et al., 2014)). RT-Seq coverage was smoothed in 500 Kb windows with a 50 Kb step.

Inferring replication timing from published data

Whole genome sequencing data were obtained from the sources listed in Table S2. Sequencing data were aligned and replication timing profiles were inferred as described above.

Modelling DNA replication

DNA replication was modelled *in-silico* using custom R code (Brick, 2020). Replication was simulated in a haploid genome. Non-autosomal chromosomes were excluded. The initial wave of origin firing was assumed to be synchronous. *k* replication origins were selected at random and without replacement, using the origin efficiency as a selection weight. At each origin, a left-ward and a right-ward replication fork was established, and replication was then simulated bi-directionally outward from each origin in 10 Kb steps. After each step, we assessed which replication forks have collided with another fork and which forks reached the end of the

958 chromosome. These forks were terminated. For every two terminated forks, we allowed
 959 replication to begin from a new, randomly chosen origin (as above, weighted by origin
 960 efficiency). Origins that have previously fired and origins in regions that have already been
 961 replicated were not considered for selection. This iterative process continued until the genome
 962 was fully replicated or for a defined number of cycles. To simulate replication in a population,
 963 this simulation was performed in n haploid genomes and the results were combined.

964 **Hyperparameter search for replication models**

965 To assess the best-fitting parameters of DNA replication *in-silico*, we performed a
 966 grid-hyperparameter search. A hyperparameter search was performed for each pair-wise
 967 combination of RT and replication origins. We optimized for a scoring function defined as the
 968 Pearson correlation coefficient between the experimentally-determined replication timing and
 969 the *in-silico* simulated replication timing. A population of 250 replicating haploid genomes was
 970 used for the hyperparameter search. The grid-search generated an RT-Sim model for each
 971 combination of the following parameters:

- 972 a. Origin density; $R_d = 0.02$ to 0.4 in 0.01 steps; 0.45 to 0.6 in 0.05 steps
- 973 b. Modelling runtime; $N_{\text{epochs}} = 10$ to 7500 in steps of 10
- 974 c. Replicating cells in population; $P (\%) = 10, 30, 50, 70, 90$
- 975 d. Use origin efficiency; $E = \text{TRUE} / \text{FALSE}$ (if FALSE , all origins were equally weighted for
 976 “random” selection)

977 The fine-grained search yielded multiple models with similar scores, therefore in assessing
 978 results, the top $g\%$ of models were examined.

979 **Generating simulated replication timing**

980 The hyperparameters of the top-scoring RT-Sim model were used to generate an RT-Sim model
 981 from 500 haploid genomes and run it to completion. Simulated replication timing was then
 982 defined as the mean replication time across all simulations for each 10 Kb window in the
 983 genome.

984 Sources and processing of annotation data for the mouse genome

985 CpG islands in the mouse genome were obtained from the Irizarry lab at Harvard University,
986 USA (<http://www.haowulab.org/software/makeCGI/model-based-cpg-islands-mm9.txt>) (Wu et
987 al., 2010). UCSC liftover was used to convert mm9 to mm10 genome coordinates.

988 Meiotic DSB hotspots in C57BL6J mice were obtained from columns 1,2,3,6 of Supplementary
989 File 41586_2018_492_MOESM3_ESM.zip (Brick et al., 2018a). H3K4m3 ChIP-Seq signals at
990 DSB hotspots in C57BL6J mice (Baker et al., 2014) was obtained from columns 1,2,3,50 of
991 Supplementary File 41586_2018_492_MOESM3_ESM.zip (Brick et al., 2018a). Genome-wide
992 Spo11-associated oligo mapping data in C57BL6 mice (Lange et al., 2016) were obtained in
993 processed form from columns 1,2,3,55 of Supplementary File
994 41586_2018_492_MOESM3_ESM.zip (Brick et al., 2018a). DSB hotspots in *Prdm9*^{-/-} mice
995 (Brick et al., 2012) were obtained from GSM2664291 (Brick et al., 2018a). DSB hotspots from
996 mice homozygous for a humanized PRDM9 zinc-finger array (B6^{h/h}) were obtained from the
997 GEO (accession: GSM2049306; file: GSM2049306_dmc1hotspots_B6.PRDM9hh.txt.gz)
998 (Davies et al., 2016). B6^{h/h} hotspots overlapping hotspots from *Prdm9*^{-/-} were discarded.
999 H3K4m3 ChIP-Seq peaks in B6^{h/h} mice were obtained from the GEO (accession: GSM1904284;
1000 file GSM1904284_H3K4me3.ForceCalledValues.B6.PRDM9hh.txt) (Davies et al., 2016).
1001 Crossovers and non-crossovers defined by the humanized variant of mouse PRDM9 were
1002 obtained from supplementary file 3 (41467_2019_11675_MOESM6_ESM.csv) and 4
1003 (41467_2019_11675_MOESM7_ESM.csv), respectively, from (Li et al., 2019a). Affinity-Seq
1004 data describing the *in-vitro* binding preferences of the C57BL6J PRDM9 variant were obtained
1005 from the Sequence Read Archive (accession: SRR1976005) (Walker et al., 2015). Control data
1006 used for Affinity-Seq peak calling was obtained from the Sequence Read Archive (accession:
1007 SRR1976003 & SRR1976004) (Walker et al. 2015). Affinity-Seq and control reads were aligned
1008 to the reference mm10 genome using bwa mem 0.7.12. Affinity-Seq peak calling was performed
1009 using MACS 2.1.0.20150731 with the following arguments: -g mm -bw 1000 --keep-dup all
1010 --slocal 5000 -q 0.05. Affinity-Seq peak strength was determined as Affinity-Seq coverage less
1011 Control DNA coverage at Affinity-Seq peaks. Hi-C data from mouse zygotene cells was
1012 downloaded from the GEO (GEO accession: GSE122622; file:
1013 GSE122622_zygotene_overall.hic) (Patel et al., 2019). The Juicer tools 1.5.6 eigenvector
1014 program (Durand et al., 2016) was used to extract the first eigenvector from the .hic file, using
1015 the following arguments: KR, BP, binsize = 100000, -p. H3K9m2 (accession: SRR1975998) and

1016 H3K9m3 (accession: SRR1585300) ChIP-Seq data (Walker et al., 2015) from spermatocytes
1017 were obtained from the SRA. Reads were aligned to the reference mm10 genome using bwa
1018 mem 0.7.12.

1019 ATAC-Seq data were obtained from the published datasets described in Supplementary Figure
1020 9. Each dataset was aligned to the mouse mm10 reference genome using bwa mem 0.7.12.
1021 Peak calling was performed for each dataset using MACS 2.1.2 without a control and using
1022 default arguments. ATAC-Seq peaks coinciding with CpG-islands were defined for use on
1023 Supplementary Figure 9.

1024 **Mapping Okazaki-fragment sequencing data**

1025 Data from Okazaki-fragment sequencing in mouse ES-cells were obtained from SRA accession
1026 SRR7535256 (Petryk et al., 2018). Sequencing reads were aligned to the mm10 reference
1027 genome using the mem algorithm of bwa 0.7.12 (Li, 2013; Li and Durbin, 2009).

1028 **Obtaining published SNS-Seq replication origins data**

1029 Processed peak calls for MEFs and ESCs were obtained from GEO accession GSE99741
1030 (Almeida et al., 2018); MEFs: GSM2651107_MEFs_WT_repl_I_peaks.bigBed; ESCs:
1031 GSM2651111_mES_WT_repl_I_peaks.bigBed. Processed peak calls for ESCs from (Cayrou et
1032 al., 2015) were obtained from GEO accession GSE68347;
1033 GSE68347_Initiation_Sites.bedGraph.gz. The UCSC liftover tool was used to convert mm9 to
1034 mm10 coordinates.

1035 **Mapping DNA DSB hotspots from DMC1-SSDS experiments**

1036 A testicular biopsy was obtained from a commercial source (Folio, Ohio). Genotyping for
1037 PRDM9 alleles using Sanger sequencing (Pratto et al., 2014) revealed that this individual did
1038 not carry a copy of the most common PRDM9 variant in humans (PRDM9_A). Instead, this
1039 individual was heterozygous for the PRDM9_C and PRDM9_{L4} alleles (Berg et al., 2010). A
1040 DMC1-SSDS library was prepared using the protocol in (Brick et al., 2018b; Pratto et al., 2014).
1041 Published DMC1-SSDS data were also obtained for two human PRDM9_A-homozygous males
1042 (GEO accessions: GSM1447325 & GSM1447328) (Pratto et al., 2014). Sequencing reads were
1043 aligned to the human hg38 reference genome using bwa 0.7.12 and the single-stranded DNA

1044 pipeline (Brick et al., 2018b; Khil et al., 2012). Hotspots were called using the DMC1-SSDS DSB
1045 hotspot calling pipeline (Brick et al., 2018b).

1046 **Crossover and non-crossover data for humans**

1047 Crossover data for human males and females were obtained from the supplementary data files
1048 S1 and S2, respectively, from (Halldorsson et al., 2019). Non-crossover data were obtained
1049 from supplementary file S2 from (Halldorsson et al., 2016). NCOs from the CHIP-based
1050 approach and the sequencing-based approach were merged.

1051 **Data analysis tools and versions**

1052 BEDtools v.2.29.2 (Quinlan and Hall, 2010)

1053 BWA 0.7.12 (Li, 2013; Li and Durbin, 2009).

1054 DeepTools v.3.0.1 (Ramírez et al., 2014)

1055 Juicer v.1.5.6 (Durand et al., 2016)

1056 MACS v.2.1.0.20150731 (Zhang et al., 2008)

1057 MACS v.2.1.2 (Zhang et al., 2008)

1058 Nextflow v.20.01.0 (Di Tommaso et al., 2017)

1059 Picard v.2.9.2 (Broad Institute of MIT and Harvard, 2018)

1060 R v.3.6.0 (R Core Team, 2014)

1061 SAMtools v.1.9 (Li et al., 2009)

1062 SRA toolkit v.2.9.2 (Leinonen et al., 2011)

1063 UCSC toolkit v.385 (Kent et al., 2010)

References

- Almeida, R., Fernández-Justel, J.M., Santa-María, C., Cadoret, J.-C., Cano-Aroca, L., Lombraña, R., Herranz, G., Agresti, A., and Gómez, M. (2018). Chromatin conformation regulates the coordination between DNA replication and transcription. *Nat. Commun.* 9, 1590.
- Amemiya, H.M., Kundaje, A., and Boyle, A.P. (2019). The ENCODE Blacklist: Identification of Problematic Regions of the Genome. *Sci. Rep.* 9, 9354.
- Baker, C.L., Walker, M., Kajita, S., Petkov, P.M., and Paigen, K. (2014). PRDM9 binding organizes hotspot nucleosomes and limits Holliday junction migration. *Genome Res.* 24, 724–732.
- Baudat, F., Buard, J., Grey, C., Fledel-Alon, A., Ober, C., Przeworski, M., Coop, G., and de Massy, B. (2010). PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice. *Science* 327, 836–840.
- Bellve, A.R., Cavicchia, J.C., and Millette, C.F. (1977). Spermatogenic cells of the prepuberal mouse: isolation and morphological characterization. *J. Cell Biol.*
- Bennett, M.D., Smith, J.B., and Riley, R. (1972). The effects of polyploidy on meiotic duration and pollen development in cereal anthers. *Proceedings of the Royal Society of London. Series B. Biological Sciences* 181, 81–107.
- Berg, I.L., Neumann, R., Lam, K.-W.G., Sarbajna, S., Odenthal-Hesse, L., May, C.A., and Jeffreys, A.J. (2010). PRDM9 variation strongly influences recombination hot-spot activity and meiotic instability in humans. *Nat. Genet.* 42, 859–863.
- Bergerat, A., de Massy, B., Gadelle, D., Varoutas, P.C., Nicolas, A., and Forterre, P. (1997). An atypical topoisomerase II from Archaea with implications for meiotic recombination. *Nature* 386, 414–417.
- Bielinsky, A.K., and Gerbi, S.A. (1998). Discrete start sites for DNA synthesis in the yeast ARS1 origin. *Science* 279, 95–98.
- Bishop, D.K., Park, D., Xu, L., and Kleckner, N. (1992). DMC1: a meiosis-specific yeast homolog of E. coli recA required for recombination, synaptonemal complex formation, and cell cycle progression. *Cell* 69, 439–456.
- Blitzblau, H.G., Chan, C.S., Hochwagen, A., and Bell, S.P. (2012). Separation of DNA replication from the assembly of break-competent meiotic chromosomes. *PLoS Genet.* 8, e1002643.
- Borde, V., Goldman, A.S., and Lichten, M. (2000). Direct coupling between meiotic DNA replication and recombination initiation. *Science* 290, 806–809.
- Boulton, A., Myers, R.S., and Redfield, R.J. (1997). The hotspot conversion paradox and the

- 1098 evolution of meiotic recombination. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 8058–8063.
- 1099 Brick, K. (2020). Simulating replication timing in the mouse and human genomes (Github).
- 1100 Brick, K., Smagulova, F., Khil, P., Camerini-Otero, R.D., and Petukhova, G.V. (2012). Genetic
1101 recombination is directed away from functional genomic elements in mice. *Nature* **485**,
1102 642–645.
- 1103 Brick, K., Thibault-Sennett, S., Smagulova, F., Lam, K.-W.G., Pu, Y., Pratto, F., Camerini-Otero,
1104 R.D., and Petukhova, G.V. (2018a). Extensive sex differences at the initiation of genetic
1105 recombination. *Nature* **561**, 338–342.
- 1106 Brick, K., Pratto, F., Sun, C.-Y., Camerini-Otero, R.D., and Petukhova, G. (2018b). Analysis of
1107 Meiotic Double-Strand Break Initiation in Mammals. In *Methods in Enzymology*, (Academic
1108 Press), pp. 391–418.
- 1109 Broach, J.R., Li, Y.Y., Feldman, J., Jayaram, M., Abraham, J., Nasmyth, K.A., and Hicks, J.B.
1110 (1983). Localization and sequence analysis of yeast origins of DNA replication. *Cold Spring*
1111 *Harb. Symp. Quant. Biol.* **47 Pt 2**, 1165–1173.
- 1112 Broad Institute of MIT and Harvard (2018). Picard tools (Broad Institute of MIT and Harvard).
- 1113 Buard, J., Rivals, E., Dunoyer de Segonzac, D., Garres, C., Caminade, P., de Massy, B., and
1114 Boursot, P. (2014). Diversity of Prdm9 zinc finger array in wild mice unravels new facets of the
1115 evolutionary turnover of this coding minisatellite. *PLoS One* **9**, e85021.
- 1116 Callan, H.G. (1973). DNA Replication in the Chromosomes of Eukaryotes. In *Molecular*
1117 *Cytogenetics*, B.A. Hamkalo, and J. Papaconstantinou, eds. (Boston, MA: Springer New York),
1118 pp. 31–47.
- 1119 Castro, J.P.L., Yancoskie, M.N., Marchini, M., Belohlavy, S., Hiramatsu, L., Kučka, M., Beluch,
1120 W.H., Naumann, R., Skuplik, I., Cobb, J., et al. (2019). An integrative genomic analysis of the
1121 Longshanks selection experiment for longer limbs in mice. *eLife* **8**.
- 1122 Cayrou, C., Grégoire, D., Coulombe, P., Danis, E., and Méchali, M. (2012). Genome-scale
1123 identification of active DNA replication origins. *Methods* **57**, 158–164.
- 1124 Cayrou, C., Ballester, B., Peiffer, I., Fenouil, R., Coulombe, P., Andrau, J.-C., van Helden, J.,
1125 and Méchali, M. (2015). The chromatin environment shapes DNA replication origin organization
1126 and defines origin classes. *Genome Res.* **25**, 1873–1885.
- 1127 Cha, R.S., and Kleckner, N. (2002). ATR homolog Mec1 promotes fork progression, thus
1128 averting breaks in replication slow zones. *Science* **297**, 602–606.
- 1129 Cha, R.S., Weiner, B.M., Keeney, S., Dekker, J., and Kleckner, N. (2000). Progression of meiotic
1130 DNA replication is modulated by interchromosomal interaction proteins, negatively by Spo11p
1131 and positively by Rec8p. *Genes Dev.* **14**, 493–503.
- 1132 Chagin, V.O., Casas-Delucchi, C.S., Reinhart, M., Schermelleh, L., Markaki, Y., Maisei, A.,
1133 Bolius, J.J., Bensimon, A., Fillies, M., Domaing, P., et al. (2016). 4D Visualization of replication

1134 foci in mammalian cells corresponding to individual replicons. *Nat. Commun.* **7**, 11231.

1135 Chandley, A.C. (1986). A model for effective pairing and recombination at meiosis based on
1136 early replicating sites (R-bands) along chromosomes. *Hum. Genet.* **72**, 50–57.

1137 Davies, B., Hatton, E., Altemose, N., Hussin, J.G., Pratto, F., Zhang, G., Hinch, A.G., Moralli, D.,
1138 Biggs, D., Diaz, R., et al. (2016). Re-engineering the zinc fingers of PRDM9 reverses hybrid
1139 sterility in mice. *Nature* **530**, 171–176.

1140 Diagouraga, B., Clément, J.A.J., Duret, L., Kadlec, J., de Massy, B., and Baudat, F. (2018).
1141 PRDM9 Methyltransferase Activity Is Essential for Meiotic DNA Double-Strand Break Formation
1142 at Its Binding Sites. *Mol. Cell.*

1143 Dileep, V., and Gilbert, D.M. (2018). Single-cell replication profiling to measure stochastic
1144 variation in mammalian replication timing. *Nat. Commun.* **9**, 427.

1145 Di Tommaso, P., Chatzou, M., Floden, E.W., Barja, P.P., Palumbo, E., and Notredame, C.
1146 (2017). Nextflow enables reproducible computational workflows. *Nat. Biotechnol.* **35**, 316–319.

1147 Durand, N.C., Shamim, M.S., Machol, I., Rao, S.S.P., Huntley, M.H., Lander, E.S., and Aiden,
1148 E.L. (2016). Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C
1149 Experiments. *Cell Syst* **3**, 95–98.

1150 Duret, L., and Galtier, N. (2009). Biased gene conversion and the evolution of mammalian
1151 genomic landscapes. *Annu. Rev. Genomics Hum. Genet.* **10**, 285–311.

1152 Foulk, M.S., Urban, J.M., Casella, C., and Gerbi, S.A. (2015). Characterizing and controlling
1153 intrinsic biases of lambda exonuclease in nascent strand sequencing reveals phasing between
1154 nucleosomes and G-quadruplex motifs around a subset of human replication origins. *Genome*
1155 *Res.* **25**, 725–735.

1156 Fu, H., Besnard, E., Desprat, R., Ryan, M., Kahli, M., Lemaitre, J.-M., and Aladjem, M.I. (2014).
1157 Mapping replication origin sequences in eukaryotic chromosomes. *Curr. Protoc. Cell Biol.* **65**,
1158 22–20.

1159 Ghosal, S.K., and Mukherjee, B.B. (1971). THE CHRONOLOGY OF DNA SYNTHESIS,
1160 MEIOSIS AND SPERMOGENESIS IN THE MALE MOUSE AND GOLDEN HAMSTER. *Can. J.*
1161 *Genet. Cytol.* **13**, 672–682.

1162 Gindin, Y., Valenzuela, M.S., Aladjem, M.I., Meltzer, P.S., and Bilke, S. (2014). A chromatin
1163 structure-based model accurately predicts DNA replication timing in human cells. *Mol. Syst.*
1164 *Biol.* **10**, 722.

1165 Goodsell, D.S., and Dickerson, R.E. (1994). Bending and curvature calculations in B-DNA.
1166 *Nucleic Acids Res.* **22**, 5497–5503.

1167 Grant, G.D., Kedziora, K.M., Limas, J.C., Cook, J.G., and Purvis, J.E. (2018). Accurate
1168 delineation of cell cycle phase transitions in living cells with PIP-FUCCI. *Cell Cycle* **17**,
1169 2496–2516.

- 1170 Guo, J., Grow, E.J., Mlcochova, H., Maher, G.J., Lindskog, C., Nie, X., Guo, Y., Takei, Y., Yun,
1171 J., Cai, L., et al. (2018). The adult human testis transcriptional cell atlas. *Cell Res.* 28,
1172 1141–1157.
- 1173 Halldorsson, B.V., Hardarson, M.T., Kehr, B., Styrkarsdottir, U., Gylfason, A., Thorleifsson, G.,
1174 Zink, F., Jonasdottir, A., Jonasdottir, A., Sulem, P., et al. (2016). The rate of meiotic gene
1175 conversion varies by sex and age. *Nat. Genet.* 48, 1377–1384.
- 1176 Halldorsson, B.V., Palsson, G., Stefansson, O.A., Jonsson, H., Hardarson, M.T., Eggertsson,
1177 H.P., Gunnarsson, B., Oddsson, A., Halldorsson, G.H., Zink, F., et al. (2019). Characterizing
1178 mutagenic effects of recombination through a sequence-level genetic map. *Science* 363.
- 1179 Handel, M.A., Eppig, J.J., and Schimenti, J.C. (2014). Applying “Gold Standards” to
1180 In-Vitro-Derived Germ Cells. *Cell* 157, 1257–1261.
- 1181 Higgins, J.D., Perry, R.M., Barakate, A., Ramsay, L., Waugh, R., Halpin, C., Armstrong, S.J.,
1182 and Franklin, F.C.H. (2012). Spatiotemporal asymmetry of the meiotic program underlies the
1183 predominantly distal distribution of meiotic crossovers in barley. *Plant Cell* 24, 4096–4109.
- 1184 Hinch, A.G., Zhang, G., Becker, P.W., Moralli, D., Hinch, R., Davies, B., Bowden, R., and
1185 Donnelly, P. (2019). Factors influencing meiotic recombination revealed by whole-genome
1186 sequencing of single sperm. *Science* 363, eaau8861.
- 1187 Hiratani, I., Ryba, T., Itoh, M., Rathjen, J., Kulik, M., Papp, B., Fussner, E., Bazett-Jones, D.P.,
1188 Plath, K., Dalton, S., et al. (2010). Genome-wide dynamics of replication timing revealed by in
1189 vitro models of mouse embryogenesis. *Genome Res.* 20, 155–169.
- 1190 Holm, P.B. (1977). The premeiotic DNA replication of euchromatin and heterochromatin in *Lilium*
1191 *longiflorum* (Thunb.). *Carlsberg Res. Commun.* 42, 249–281.
- 1192 Hossain, A.M., Rizk, B., Behzadian, A., and Thorneycroft, I.H. (1997). Modified guanidinium
1193 thiocyanate method for human sperm DNA isolation. *Mol. Hum. Reprod.* 3, 953–956.
- 1194 Huppert, J.L., and Balasubramanian, S. (2005). Prevalence of quadruplexes in the human
1195 genome. *Nucleic Acids Res.* 33, 2908–2916.
- 1196 Jodkowska, K., Pancaldi, V., Almeida, R., Rigau, M., Graña-Castro, O., Fernández-Justel, J.M.,
1197 Rodríguez-Acebes, S., Rubio-Camarillo, M., Pau, E.C.S., Pisano, D., et al. (2019).
1198 Three-dimensional connectivity and chromatin environment mediate the activation efficiency of
1199 mammalian DNA replication origins.
- 1200 Joshi, N., Brown, M.S., Bishop, D.K., and Börner, G.V. (2015). Gradual implementation of the
1201 meiotic recombination program via checkpoint pathways controlled by global DSB levels. *Mol.*
1202 *Cell* 57, 797–811.
- 1203 Kaback, D.B., Guacci, V., Barber, D., and Mahon, J.W. (1992). Chromosome size-dependent
1204 control of meiotic recombination. *Science* 256, 228–232.
- 1205 Keeney, S., Giroux, C.N., and Kleckner, N. (1997). Meiosis-specific DNA double-strand breaks

- 1206 are catalyzed by Spo11, a member of a widely conserved protein family. *Cell* **88**, 375–384.
- 1207 Kelly, T., and Callegari, A.J. (2019). Dynamics of DNA replication in a eukaryotic cell. *Proc. Natl.*
1208 *Acad. Sci. U. S. A.* **116**, 4973–4982.
- 1209 Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M., and Haussler, D.
1210 (2002). The human genome browser at UCSC. *Genome Res.* **12**, 996–1006.
- 1211 Kent, W.J., Zweig, A.S., Barber, G., Hinrichs, A.S., and Karolchik, D. (2010). BigWig and
1212 BigBed: enabling browsing of large distributed datasets. *Bioinformatics* **26**, 2204–2207.
- 1213 Khil, P.P., Smagulova, F., Brick, K.M., Camerini-Otero, R.D., and Petukhova, G.V. (2012).
1214 Sensitive mapping of recombination hotspots using sequencing-based detection of ssDNA.
1215 *Genome Res.* **22**, gr – 130583.
- 1216 Klein, K.N., Zhao, P.A., Lyu, X., Bartlett, D.A., Singh, A., Tasan, I., Watts, L.P., Hiraga, S.-I.,
1217 Natsume, T., Zhou, X., et al. (2019). Replication timing maintains the global epigenetic state in
1218 human cells.
- 1219 Kofman-Alfaro, S., and Chandley, A.C. (1970). Meiosis in the male mouse. An autoradiographic
1220 investigation. *Chromosoma* **31**, 404–420.
- 1221 Kojima, M.L., de Rooij, D.G., and Page, D.C. (2019). Amplification of a broad transcriptional
1222 program by a common factor triggers the meiotic cell cycle in mice. *Elife* **8**.
- 1223 Koren, A., Polak, P., Nemesh, J., Michaelson, J.J., Sebat, J., Sunyaev, S.R., and McCarroll, S.A.
1224 (2012). Differential relationship of DNA replication timing to different forms of human mutation
1225 and variation. *Am. J. Hum. Genet.* **91**, 1033–1040.
- 1226 Koren, A., Handsaker, R.E., Kamitaki, N., Karlić, R., Ghosh, S., Polak, P., Eggan, K., and
1227 McCarroll, S.A. (2014). Genetic variation in human DNA replication timing. *Cell* **159**,
1228 1015–1026.
- 1229 Kulakovskiy, I.V., Vorontsov, I.E., Yevshin, I.S., Sharipov, R.N., Fedorova, A.D., Rumynskiy, E.I.,
1230 Medvedeva, Y.A., Magana-Mora, A., Bajic, V.B., Papatsenko, D.A., et al. (2018). HOCOMOCO:
1231 towards a complete collection of transcription factor binding models for human and mouse via
1232 large-scale ChIP-Seq analysis. *Nucleic Acids Res.* **46**, D252–D259.
- 1233 Lam, K.-W.G., Brick, K., Cheng, G., Pratto, F., and Camerini-Otero, R.D. (2019).
1234 Cell-type-specific genomics reveals histone modification dynamics in mammalian meiosis. *Nat.*
1235 *Commun.* **10**, 3821.
- 1236 Lange, J., Yamada, S., Tischfield, S.E., Pan, J., Kim, S., Zhu, X., Socci, N.D., Jasin, M., and
1237 Keeney, S. (2016). The Landscape of Mouse Meiotic Double-Strand Break Formation,
1238 Processing, and Repair. *Cell* **167**, 695–708.e16.
- 1239 Lee, B., and Amon, A. (2001). Meiosis: how to create a specialized cell cycle. *Curr. Opin. Cell*
1240 *Biol.* **13**, 770–777.
- 1241 Leinonen, R., Sugawara, H., Shumway, M., and International Nucleotide Sequence Database

- 1242 Collaboration (2011). The sequence read archive. *Nucleic Acids Res.* 39, D19–D21.
- 1243 Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
- 1244 Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
1245 transform. *Bioinformatics* 25, 1754–1760.
- 1246 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
1247 Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence
1248 Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078–2079.
- 1249 Li, N., Lam, W.H., Zhai, Y., Cheng, J., Cheng, E., Zhao, Y., Gao, N., and Tye, B.-K. (2018).
1250 Structure of the origin recognition complex bound to DNA replication origin. *Nature* 559,
1251 217–222.
- 1252 Li, R., Bitoun, E., Altemose, N., Davies, R.W., Davies, B., and Myers, S.R. (2019a). A
1253 high-resolution map of non-crossover events reveals impacts of genetic diversity on mammalian
1254 meiotic recombination. *Nat. Commun.* 10, 1–15.
- 1255 Li, W., Yang, L., He, Q., Hu, C., Zhu, L., Ma, X., Ma, X., Bao, S., Li, L., Chen, Y., et al. (2019b).
1256 A Homeostatic Arid1a-Dependent Permissive Chromatin State Licenses Hepatocyte
1257 Responsiveness to Liver-Injury-Associated YAP Signaling. *Cell Stem Cell* 25, 54–68.e5.
- 1258 Long, H., Zhang, L., Lv, M., Wen, Z., Zhang, W., Chen, X., Zhang, P., Li, T., Chang, L., Jin, C., et
1259 al. (2020). H2A.Z facilitates licensing and activation of early replication origins. *Nature* 577,
1260 576–581.
- 1261 Maezawa, S., Yukawa, M., Alavattam, K.G., Barski, A., and Namekawa, S.H. (2018). Dynamic
1262 reorganization of open chromatin underlies diverse transcriptomes during spermatogenesis.
1263 *Nucleic Acids Res.* 46, 593–608.
- 1264 Marchal, C., Sima, J., and Gilbert, D.M. (2019). Control of DNA replication timing in the 3D
1265 genome. *Nat. Rev. Mol. Cell Biol.* 20, 721–737.
- 1266 Miotto, B., Ji, Z., and Struhl, K. (2016). Selectivity of ORC binding sites and the relation to
1267 replication timing, fragile sites, and deletions in cancers. *Proc. Natl. Acad. Sci. U. S. A.* 113,
1268 E4810–E4819.
- 1269 Monesi, V. (1962). Autoradiographic study of DNA synthesis and the cell cycle in spermatogonia
1270 and spermatocytes of mouse testis using tritiated thymidine. *J. Cell Biol.* 14, 1–18.
- 1271 Murakami, H., and Keeney, S. (2014). Temporospatial coordination of meiotic DNA replication
1272 and recombination via DDK recruitment to replisomes. *Cell* 158, 861–873.
- 1273 Murakami, H., and Nurse, P. (2001). Regulation of premeiotic S phase and
1274 recombination-related double-strand DNA breaks during meiosis in fission yeast. *Nat. Genet.*
1275 28, 290–293.
- 1276 Murakami, H., Lam, I., Huang, P.-C., Song, J., van Overbeek, M., and Keeney, S. (2020).
1277 Multilayered mechanisms ensure that short chromosomes recombine in meiosis. *Nature* 582,

1278 124–128.

1279 Myers, S., Bottolo, L., Freeman, C., McVean, G., and Donnelly, P. (2005). A fine-scale map of
1280 recombination rates and hotspots across the human genome. *Science* 310, 321–324.

1281 Myers, S., Bowden, R., Tumian, A., Bontrop, R.E., Freeman, C., MacFie, T.S., McVean, G., and
1282 Donnelly, P. (2010). Drive against hotspot motifs in primates implicates the PRDM9 gene in
1283 meiotic recombination. *Science* 327, 876–879.

1284 Neale, M.J., Pan, J., and Keeney, S. (2005). Endonucleolytic processing of covalent
1285 protein-linked DNA double-strand breaks. *Nature* 436, 1053–1057.

1286 Parvanov, E.D., Petkov, P.M., and Paigen, K. (2010). Prdm9 controls activation of mammalian
1287 recombination hotspots. *Science* 327, 835.

1288 Patel, L., Kang, R., Rosenberg, S.C., Qiu, Y., Raviram, R., Chee, S., Hu, R., Ren, B., Cole, F.,
1289 and Corbett, K.D. (2019). Dynamic reorganization of the genome shapes the recombination
1290 landscape in meiotic prophase. *Nat. Struct. Mol. Biol.* 1.

1291 Petryk, N., Kahli, M., d'Aubenton-Carafa, Y., Jaszczyszyn, Y., Shen, Y., Silvain, M., Thermes, C.,
1292 Chen, C.-L., and Hyrien, O. (2016). Replication landscape of the human genome. *Nat.*
1293 *Commun.* 7, 10208.

1294 Petryk, N., Dalby, M., Wenger, A., Stromme, C.B., Strandsby, A., Andersson, R., and Groth, A.
1295 (2018). MCM2 promotes symmetric inheritance of modified histones during DNA replication.
1296 *Science* 361, 1389–1392.

1297 Picard, F., Cadoret, J.-C., Audit, B., Arneodo, A., Alberti, A., Battail, C., Duret, L., and Prioleau,
1298 M.-N. (2014). The spatiotemporal program of DNA replication is associated with specific
1299 combinations of chromatin marks in human cells. *PLoS Genet.* 10, e1004282.

1300 Powers, N.R., Parvanov, E.D., Baker, C.L., Walker, M., Petkov, P.M., and Paigen, K. (2016). The
1301 Meiotic Recombination Activator PRDM9 Trimethylates Both H3K36 and H3K4 at
1302 Recombination Hotspots In Vivo. *PLoS Genet.* 12, e1006146.

1303 Pratto, F., Brick, K., Khil, P., Smagulova, F., Petukhova, G.V., and Camerini-Otero, R.D. (2014).
1304 Recombination initiation maps of individual human genomes. *Science* 346, 1256442.

1305 Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
1306 features. *Bioinformatics* 26, 841–842.

1307 Ramírez, F., Dündar, F., Diehl, S., Grüning, B.A., and Manke, T. (2014). deepTools: a flexible
1308 platform for exploring deep-sequencing data. *Nucleic Acids Res.* 42, W187–W191.

1309 R Core Team (2014). R: A language and environment for statistical computing (R Foundation for
1310 Statistical Computing, Vienna, Austria.).

1311 Sandhu, R., Monge Neria, F., Monge Neria, J., Chen, X., Hollingsworth, N.M., and Börner, G.V.
1312 (2020). DNA Helicase Mph1FANCM Ensures Meiotic Recombination between Parental

- 1313 Chromosomes by Dissociating Precocious Displacement Loops. *Dev. Cell* 53, 458–472.e5.
- 1314 Sequeira-Mendes, J., Vergara, Z., Peiró, R., Morata, J., Aragüez, I., Costas, C.,
1315 Mendez-Giraldez, R., Casacuberta, J.M., Bastolla, U., and Gutierrez, C. (2019). Differences in
1316 firing efficiency, chromatin, and transcription underlie the developmental plasticity of the
1317 Arabidopsis DNA replication origins. *Genome Res.* 29, 784–797.
- 1318 Smagulova, F., Gregoret, I.V., Brick, K., Khil, P., Camerini-Otero, R.D., and Petukhova, G.V.
1319 (2011). Genome-wide analysis reveals novel molecular features of mouse recombination
1320 hotspots. *Nature* 472, 375–378.
- 1321 Smagulova, F., Brick, K., Pu, Y., Camerini-Otero, R.D., and Petukhova, G.V. (2016). The
1322 evolutionary turnover of recombination hot spots contributes to speciation in mice. *Genes Dev.*
1323 30, 266–280.
- 1324 Smith, D.I., Zhu, Y., McAvoy, S., and Kuhn, R. (2006). Common fragile sites, extremely large
1325 genes, neural development and cancer. *Cancer Lett.* 232, 48–57.
- 1326 Smith, O.K., Kim, R., Fu, H., Martin, M.M., Lin, C.M., Utani, K., Zhang, Y., Marks, A.B., Lalande,
1327 M., Chamberlain, S., et al. (2016). Distinct epigenetic features of differentiation-regulated
1328 replication origins. *Epigenetics Chromatin* 9, 18.
- 1329 Stinchcomb, D.T., Thomas, M., Kelly, J., Selker, E., and Davis, R.W. (1980). Eukaryotic DNA
1330 segments capable of autonomous replication in yeast. *Proc. Natl. Acad. Sci. U. S. A.* 77,
1331 4559–4563.
- 1332 Takahashi, S., Miura, H., Shibata, T., Nagao, K., Okumura, K., Ogata, M., Obuse, C.,
1333 Takebayashi, S.-I., and Hiratani, I. (2019). Genome-wide stability of the DNA replication
1334 program in single mammalian cells. *Nat. Genet.* 51, 529–540.
- 1335 Valton, A.-L., Hassan-Zadeh, V., Lema, I., Boggetto, N., Alberti, P., Saintomé, C., Riou, J.-F.,
1336 and Prioleau, M.-N. (2014). G4 motifs affect origin positioning and efficiency in two vertebrate
1337 replicators. *EMBO J.* 33, 732–746.
- 1338 Walker, M., Billings, T., Baker, C.L., Powers, N., Tian, H., Saxl, R.L., Choi, K., Hibbs, M.A.,
1339 Carter, G.W., Handel, M.A., et al. (2015). Affinity-seq detects genome-wide PRDM9 binding
1340 sites and reveals the impact of prior chromatin modifications on mammalian recombination
1341 hotspot usage. *Epigenetics Chromatin* 8, 31.
- 1342 Wu, P.-Y.J., and Nurse, P. (2014). Replication origin selection regulates the distribution of
1343 meiotic recombination. *Mol. Cell* 53, 655–662.
- 1344 Wu, H., Caffo, B., Jaffee, H.A., Irizarry, R.A., and Feinberg, A.P. (2010). Redefining CpG islands
1345 using hidden Markov models. *Biostatistics* 11, 499–514.
- 1346 Yevshin, I., Sharipov, R., Valeev, T., Kel, A., and Kolpakov, F. (2017). GTRD: a database of
1347 transcription factor binding sites identified by ChIP-seq experiments. *Nucleic Acids Res.* 45,
1348 D61–D67.
- 1349 Yin, Y., Jiang, Y., Lam, K.-W.G., Berletch, J.B., Disteche, C.M., Noble, W.S., Steemers, F.J.,

- 1350 Camerini-Otero, R.D., Adey, A.C., and Shendure, J. (2019). High-Throughput Single-Cell
 1351 Sequencing with Linear Amplification. *Mol. Cell*.
- 1352 Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C.,
 1353 Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS).
 1354 *Genome Biol.* 9, R137.