Dorsal-to-ventral imbalance in the superior longitudinal fasciculus mediates methylphenidate's effect on beta oscillations in ADHD

C. Mazzetti¹, C. G. Damatac^{1,3}, E. Sprooten^{1,3}, N. ter Huurne^{1,4}, J.K. Buitelaar^{1,3,4}, O. Jensen²

¹ Radboud University Nijmegen, Donders Institute for Brain, Cognition and Behaviour, Nijmegen, The Netherlands

² Centre for Human Brain Health, School of Psychology, University of Birmingham, Birmingham, UK;

³ Department of Cognitive Neuroscience, Radboudumc, Nijmegen, The Netherlands

⁴ Karakter Child and Adolescent Psychiatry University Centre, Nijmegen, The Netherlands

Corresponding author: Cecilia Mazzetti, Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen, P.O. Box 9101, NL-6500 HB Nijmegen, The Netherlands; phone: +31-24-36- 65059; email: c.mazzetti@donders.ru.nl

Short running title

Methylphenidate and beta oscillations in ADHD

Keywords

Brain oscillations, MEG, ADHD, DTI, Biomarker, Methylphenidate

Abstract

Background: While pharmacological treatment with Methylphenidate (MPH) is a first line intervention

for ADHD, its mechanisms of action have yet to be elucidated. In a previous MEG study, we demonstrated that MPH in ADHD normalizes beta depression in preparation to motor responses (1). We here seek to identify the white matter tracts that mediate MPH's effect on beta oscillations.

Methods: We implemented a double-blind placebo-controlled crossover design, where boys diagnosed with ADHD underwent behavioral and MEG measurements during a spatial attention task while on and off MPH. Results were compared with an age/IQ-matched typically developing (TD) group performing the same task. Estimates of white matter tracts were obtained through diffusion tensor imaging (DTI). Based on aprioristic selection model criteria, we sought to determine the fiber tracts associated with electrophysiological, behavioral and clinical features of attentional functions.

Results: We identified three main tracts: the anterior thalamic radiation (ATR), the Superior Longitudinal Fasciculus ('parietal endings') (SLFp) and Superior Longitudinal Fasciculus ('temporal endings') (SLFt). ADHD symptoms severity was associated with lower fractional anisotropy (FA) within the ATR. In addition, individuals with relatively higher FA in SLFp compared to SLFt showed faster and more accurate behavioral responses to MPH. Furthermore, the same parieto-temporal FA gradient explained the effects of MPH on beta modulation: subjects with ADHD exhibiting higher FA in SLFp compared to SLFt also displayed greater effects of MPH on beta power during response preparation.

Conclusions: Based on MPH's modulatory effects on striatal dopamine levels, our data suggest that the behavioral deficits and aberrant oscillatory modulations observed in ADHD depend on a structural connectivity imbalance within the SLF, caused by a diffusivity gradient in favor of temporal rather than parietal, fiber tracts.

Introduction

The neural mechanisms underlying selective attention processes are contingent upon a complex interaction of brain networks. Prior studies using electro- and magneto-encephalography (EEG/MEG) have provided insights into diverse oscillatory patterns indexing distractor suppression and appropriate target processing. These start prior to the display of the stimuli, when participants prepare to the response by specifically allocating attentional resources in order to meet task demands (2–5). One important function is the preparation to the motor response to a cued target (e.g., button press), which interacts with visual attentional allocation. The electrophysiological correlate underlying this behavioral component is visible as a suppression of oscillations in sensorimotor areas. Specifically, modulation of brain activity in the beta band (15 – 30Hz) has been associated with top-down control of motor preparation, reflecting an interaction between cognitive and motor functions, which has been interpreted as increased excitability of task-relevant brain areas (6,7). This motivates the investigation of brain oscillations during attention tasks in populations with attentional problems.

Attention deficit-hyperactivity disorder (ADHD) is a neurodevelopmental disorder characterized by age-inappropriate levels of inattention and hyperactivity-impulsivity (American Psychiatric Association, 2013). Aberrant modulation of oscillatory activity has indeed been observed in individuals identified with ADHD, both in adults (9–11) and pediatric populations (12–14), with the latter representing the most prominent line of research, given the predominance of the disorder in early life (15,16). Weaker suppression of β -band activity (13 – 30Hz) has been found in ADHD, prior to response preparation to a cued target (17), reflecting a lack of appropriate motor planning. Furthermore, recent evidence has observed weakened beta modulation during working memory encoding in ADHD (18) consistent with the view that modulation of beta oscillations reflect cognitive functions (19,20).

Previously, we showed that beta depression in preparation to responses to a cued target is reduced in children with ADHD as compared to matched TD peers (1). Additionally, in a double-blind randomized placebo-controlled design within the ADHD group we showed that MPH restores levels of beta depression in children with ADHD, by normalizing its values to the ones observed in the TD group. This effect is allegedly attributable to the effects of psychostimulants on catecholamines' levels in the midbrain (21). Specifically, MPH blocks the reuptake of norepinephrine and, particularly, dopamine at

the synaptic cleft (22–24) increasing their availability. A dopaminergic imbalance within networks mediated by the prefrontal cortex has indeed been proposed to underlie symptoms of attentional deficits and hyperactivity (25,26). Consistently, the modulatory action of dopamine is alleged to mediate the interaction between frontoparietal and default mode attentional networks (27).

So far, the link between beta oscillations and brain structure remains yet to be elucidated. This can be achieved by means of diffusion tensor imaging (DTI), which estimates the direction of diffusion of water molecules in the brain and is thought to reflect the underlying microstructural properties of white matter fiber tracts (28,29). Among the DTI-derived metrics is fractional anisotropy (FA), reflecting coherence of diffusion of water along the main tract direction (30,31) and, allegedly, the underlying tissue microstructure, such as integrity of myelin sheath, which impacts the overall mobility of water along axons (32). DTI studies in ADHD have so far pointed to reduced white matter integrity in widespread areas in the brain (33–35). Diffusion in the angular bundle of the cingulum correlates with hyperactivity-impulsivity symptomatology and associations of FA with ADHD are not uniformly distributed across white matter tracts (36). Importantly, the Superior Longitudinal Fasciculus, a white matter tract known to sustain spatial attention functions (38,39), has been consistently reported among the structural correlates of ADHD symptoms. In particular, lower integrity (FA) along this bundle has been associated with severity of ADHD symptoms and behavioral performance on cognitive tests (40–42). Taking advantage of the spatial resolution offered by MEG and the link to microstructural connectivity properties offered by DTI, this study aims to study the association between oscillatory and structural features in relation to attentional impairments in children with ADHD and typically developing (TD) children. We hence coupled the previously reported electrophysiological results (1) with an analysis of white matter microstructural properties in the same subjects.

Methods and Materials

A detailed explanation of participants' inclusion and exclusion criteria, attentional task, and study design has been previously described (1) and can be found in *Supplementary Materials*.

Participants

The study included 27 children diagnosed with ADHD and 27 typically developing (TD) male children. A total of 10 children (9 ADHD) withdrew from the experiment after at least one session, due to unwillingness to proceed and/or excessive complications during testing. As a result, structural and diffusion-weighted MRI scans and MEG data were acquired for a total of 49 (26 TD) and 46 participants (26 TD), respectively.

Experimental design

The study design is outlined in **Figure 1**. Overall, children in the ADHD group visited the lab three times, while children in the TD group twice. During the first intake-session, participants from groups underwent behavioral and IQ testing. For the ADHD group, a further in-depth intake was conducted to determine the medication dosage to be used during the task, based on operating procedures followed in prior studies (43). Based on the screening, a standardized dosage was chosen (either 10 or 15mg Methylphenidate immediate release; IR-MPH). Following the behavioral screenings, the MRI session took place for a duration of ~30mins.

During the second visit, both groups undertook the MEG testing. For the TD group, this constituted the last day of testing, while for the ADHD group, two MEG sessions were planned at two different visits, separated by at least one-week interval. The ADHD group performed the MEG task twice, i.e., under two conditions (MPH and placebo), according to a double-blind placebo controlled randomized design. Prior to each MEG session, a 24 hour treatment suspension allowed to control for withdrawal symptoms related to drug administration (rebound effect) (44). MEG testing began one hour after medication intake, allowing to reach on average moderate plasma concentration (C_{max}) of the drug along the experiment, which progressively increases and reaches its peak around the second hour post-intake (45). After completion of the MEG session, participants proceeded with their own regular stimulant.

The attention task

The task was presented as a child-friendly adaptation of a Posner's cueing paradigm for spatial orienting of attention, where a central cue (represented with a clown fish looking either at the left or at the right side of the screen) indicated the upcoming position of a target (a shark with an open mouth). Participants were asked to indicated via button press the position of the target, while ignoring the distractor on the opposite screen side (shark with mouth closed) (see **Figure 2**).

The experiment consisted of 360 trials, equally divided in 10 blocks, after which the participant was given the possibility to take a break and/or talk to the parents. Note that, for the ADHD group, the treatment order for the two MEG sessions was randomized across participants.

MEG data acquisition and analysis

Electromagnetic brain activity was recorded from the participants seated in a CTF 275-sensor whole-head MEG system with axial gradiometers (CTF MEG Systems, VSM MedTech Ltd.). Head position was monitored throughout the experiment via online head-localization software, allowing, if necessary, readjustment of the participant's position between blocks. Complete illustration of the steps followed for the analysis of power and beta oscillatory indices of interest, can be found in our previous paper (1) and are further described in the Supplementary Materials.

Importantly, in the above study, we report how the preparation to motor response to the cued target was accompanied by a desynchronization of beta oscillations in the MEG data at central sensors. For each subject, a Beta Preparation Index (PI(β)) was hence computed by considering the average beta band modulation over the sensors and time window of interest ($f = 15 - 30\text{Hz}$, $-1000 < t < 0 \text{ ms}$).

To estimate MPH modulatory effects exerted on beta preparation, we considered the difference in PI(β) between MPH drug and placebo in the ADHD sample, referred to as $\Delta\text{PI}(\beta)$:

$$\Delta\text{PI}(\beta)_s = \frac{\text{PI}(\beta)_{s,\text{Placebo}} - \text{PI}(\beta)_{s,\text{MPH}}}{\text{PI}(\beta)_{s,\text{Placebo}} + \text{PI}(\beta)_{s,\text{MPH}}} \quad \text{Eq.(1)}$$

Based on Eq.1, the larger the $\Delta PI(\beta)$ the higher the stronger in the beta depression following MPH intake.

Structural data analysis

MRI and DTI data were acquired via a 3T MAGNETOM Skyra MR scanner (Siemens AG, Healthcare Sector, Erlangen, Germany) with a product 32-channel head coil. The protocol included a T1-weighted MRI scan for anatomical reference and analysis and diffusion-weighted MRI scans for performing fiber tractography.

Anatomical and DTI images were analyzed in FreeSurfer 6.0 (<http://surfer.nmr.mgh.harvard.edu/>). The TRACULA toolbox (Tracts Constrained by Underlying Anatomy; Yendiki et al., 2011) was implemented for preprocessing of DWI images and for subsequent delineation of 18 major white matter tracts (8 bilateral and 2 interhemispheric): corticospinal tract (CST), inferior longitudinal fasciculus (ILF), uncinate fasciculus (UNC), anterior thalamic radiations (ATR), cingulum-cingulate gyrus bundle (CCG), cingulum-angular bundle (CAB), superior longitudinal fasciculus-parietal terminations (SLFP), superior longitudinal fasciculus-temporal terminations (SLFT), corpus callosum forceps major and minor (Fmaj, Fmin). TRACULA allows the automated reconstruction of major white matter pathways based on the global probabilistic approach described in (47), and further extends it by incorporating anatomical knowledge in the prior probability function: each resulting segmented tract is the best fit not only given the observed diffusion data within each subject, but also given its similarity to the known tract anatomy in relation to grey matter segmentations from FreeSurfer.

Preprocessing steps included eddy current compensation, head motion correction, intra-subject registration (individual DWI to individual T1), inter-subject registration (individual T1 affine registration to the MNI template space), creation of cortical masks (parcellation via probabilistic information estimated from a manually labeled training set based on Desikan-Killiany Atlas, **Figure 3**) and white-matter masks (based on (44)), tensor fitting for extraction of tensor-based measures,

computation of anatomical priors for white-matter pathways reconstruction (e.g., diffusion eigenvectors, eigenvalues and FA for each voxel). Next, *bedpostx* was applied with a two-fiber, ball-and-stick model to estimate the distributions of the diffusion parameters and create the input for probabilistic tractography (49).

See **Figure 4** for an example of TRACULA's output in a single healthy TD participant, showing the posterior distribution for all the white matter pathways included in the segmentation pipeline. Each participant's scan was then registered to standard MNI space for group-level analyses. A pairwise correlation matrix for bilateral structures is presented in **Figure 5**, showing that no negative association was present between tracts' FA values.

All statistical analyses were performed in MATLAB2019a. For all analyses, the main metric of interest was mean FA within the tracts.

Model selection of white matter ROIs

In order to determine the relationship between mean FA along the white matter tracts segmented, as well as electrophysiological and behavioural measures, we implemented a general linear model (GLM) specifying the FA values of the white matter structures as regressors for the prediction of each index of interest.

Consistent with the methods implemented in previous work (50), and given the relative heterogeneity of current results in the field (33,51,52), we implemented a data driven strategy aimed at selecting the optimal set of or regressors to be included in the model.

We started by focusing on the behavioral performance to constrain the optimal model, hence setting IESs as dependent variable. Next, we considered all possible combinations from 2 to 5 regressors reflecting the FA values of the bilateral tracts segmented with TRACULA: SLFt, SLFp, ATR, CAB, CCG, ILF (in addition to the model including all 6 regressors, referred to as the 'full model'). We hence separately considered the general linear mixed models (maximum likelihood estimation) derived from all possible combinations of n regressors, i.e., including either 2, 3, 4 or 5 regressors (i.e. ROIs), by

computing all possible unique permutations of n regressors from the subset defined. This resulted in a set of models for each of the 4 possibilities using 2, 3, 4 or 5 regressors. Next, for each of these subsets, we derived the model associated with the lowest *Akaike Information Criterion* (AIC), *Bayesian Information Criterion* (BIC), highest *log likelihood* (where the winning model would be the one associated with at least two highest criteria compared to the others in the same model subset). These values have been commonly used in model selection to identify the best predictor subsets for a statistical model (53). Upon selection, we ended up with the five ‘best’ models, representative of each of the four model options described above.

In a final step, we identified the ‘winning model’ among the 4 selected ones (based on the same criteria as the previous selection) and compared it with the following full model based on 6 regressors:

$$IES \sim b_0 + (b_1FA_{SLFT} + b_2FA_{SLFP} + b_3FA_{ATR} + b_4FA_{CAB} + b_5FA_{CCG} + b_6FA_{ILF}) * b_7Group + \varepsilon \quad mdl(1)$$

Where *Group* is a categorical variable describing the subjects’ group (ADHD vs TD).

We then tested how the winning model could account for ADHD symptoms (ADHD-RS), beta modulation ($PI(\beta)$) and modulation of drug related beta oscillatory response ($\Delta PI(\beta)$).

Results

While a total of 22 ADHD and 27 TD children underwent the MR session, only a smaller group of 18 ADHD and 26 TD successfully completed both MEG sessions (see Fig. 2 for the task). In the following section, we report the results of the latter group, where we quantified the association between white matter tracts microstructure, behavioral performance and MEG-derived measures. The bigger sample was considered for the association between diffusion weighted imaging (DWI) results and ADHD symptoms score.

Behavioral performance

Behavioral performance in the ADHD group significantly improved following MPH administration ($t_{(17)}=2.49, p=.023$), as reflected by lower inverse efficiency scores (IESs: accuracy/reaction time). No significant difference in IES was found between TD and ADHD: the TD group did not perform better when compared to the ADHD group in the Placebo ($t_{(42)}=-.22, p=.827$) nor in the MPH condition ($t_{(42)}=.77, p=.445$).

Beta desynchronization in preparation to response to the cued target

As depicted in **Figure 6**, ADHD subjects in the placebo conditions exhibited lower overall beta depression, which was restored following MPH intake. This was observed as a diminished $PI(\beta)$ s with values closer to those observed in the TD group (see Supplementary Methods and Materials). We will here focus on the association between beta modulation and white matter microstructural diffusion properties.

Definition of white matter structures of interest: model building strategy

Following the step-wise model selection criteria described in the *Materials and Methods* section, we identified a model with 3 regressors as the best fit (AIC= -67.34 and BIC=-53.09, Log Likelihood=42.67, Adjusted $R^2=.19$) which included the regressors FA_{SLFt} ($p=7\times 10^{-4}$), FA_{SLFp} ($p=.003$), and FA_{ATR} ($p=.068$). A 3D rendering of the three white matters tracts selected (SLFT, SLFP and ATR) is illustrated in **Figure 7** and they were considered as ROIs for later linear mixed model analyses.

We first enquired whether FA along these tracts differed between groups. We hence used a 2-way unbalanced ANOVA with factors ‘*tract*’ (ATR, SLFp, SLFt) and ‘*group*’ (‘TD’, ‘ADHD’) in relation to FA values. While the results yield a main effect of *group* ($p=.009$), showing a group difference in FA across all tracts, and *tract* ($p=4.8\times 10^{-13}$), reflecting a difference in FA between tracts, no significant interaction *group by tract* emerged ($p=.453$), indicating that the pattern of FA across tracts was similar across groups.

Functional anisotropy of the Superior Longitudinal Fasciculus relates to behavioral performance

We considered task performance in relation to the three identified tracts of the winning model. Beta coefficients and adjusted response plots (i.e., showing the response as a function of one predictor, averaged over the others) derived from the model are shown in **Figure 8**: SLFt and SLFp were associated with beta coefficients of opposite sign: while higher FA values in the SLFt were associated with a higher IES ($p = 7 \times 10^{-4}$), i.e. worse performance (**Figure 8B**), a higher FA in the SLFp accounted for lower IES ($p = .003$), i.e. better behavioral performance (**Figure 8C**). A significant interaction term furthermore emerged for FA_{SLFt} and FA_{SLFp} with *Group* ($p = .002$ and $p = .023$, associated with a negative and positive coefficient, respectively). A negative interaction term of FA_{SLFt} with *Group* indicates that the effect of FA_{SLFt} on IES for the TD group was relatively weaker than for the ADHD group. *Vice versa*, a positive interaction term of FA_{SLFp} with *Group*, denotes that the effect of FA_{SLFp} for the TD group was relatively stronger than for ADHD.

Given the strong association between the two regressors FA_{SLFt} and FA_{SLFp} ($r = .86$, $p = 2 \times 10^{-4}$; see correlation matrix in **Figure 5**) and based on the high degree of overlap between the two reconstructed tracts (as they overlapped anteriorly; **Figure 4,7**) more analysis was required to determine how SLFp and SLFt related to performance. We hence computed a measure describing the parietal-to-temporal SLF imbalance, according to:

$$FA_{SLF(p-t)} = \frac{FA_{SLFp} - FA_{SLFt}}{FA_{SLFp} + FA_{SLFt}} \quad \text{Eq.(3)}$$

As a result, a given subject would display a specific degree of parietal-to-temporal diffusivity (i.e. imbalance) along the SLF tract: A higher value of the imbalance reflects a stronger diffusivity at parietal locations along the tract, and a lower value along the gradient reflected a stronger diffusivity at temporal locations.

We therefore considered a model specifically incorporating the parietal-temporal imbalance:

$$IES \sim b_0 + (b_1 FA_{SLF(p-t)} + b_2 FA_{ATR}) * b_3 Group + \varepsilon \quad mdl(1.a)$$

Figure 8D shows the association between IES and $FA_{SLF(p-t)}$. As a corollary of Eq.(3), the negative partial association ($p=.001$) shows that subjects with a higher parietal than temporal FA along the SLF tract, were also the ones with a better behavioral performance. Also in this model, a significant interaction ($p=.006$) emerged between tract*Group, which reflected that the effect of $FA_{SLF(p-t)}$ on IES was stronger for the TD group compared to the ADHD.

Fractional anisotropy of the ATR predicts ADHD symptoms

In order to assess whether the FA values of the white matter ROIs were related to symptoms (as measured by ADHD-Rating Scale; ADHD-RS), we considered the full sample of participants. We included symptoms across both the TD and the ADHD (placebo condition) group, hence embracing the notion that ADHD symptomatology derives from a ‘spectrum’, rather than from a dichotomous distinction with TD peers. The following model was then applied:

$$ADHD - RS \sim b_0 + b_1 FA_{SLFT} + b_2 FA_{SLFP} + b_3 FA_{ATR} + \varepsilon \quad mdl(2)$$

The resulting model was significant ($R^2=.22$, $p=.01$, **Figure 9A**), and FA_{ATR} was associated with a significant partial coefficient ($p=.007$) showing a negative relationship with ADHD symptoms (**Figure 9B, 9C**): higher FA in the ATR predicted an overall lower ADHD symptomatology. FA_{SLFP} and FA_{SLFT} did not show a significant partial correlation with symptoms score ($p=.494$ and $p=.309$, respectively).

Parietotemporal gradient along the SLF reflects methylphenidate’s effect on preparatory beta depression

Finally, we sought to investigate the relationship between FA values in the selected ROIs, and the patterns of MPH associated beta depression ($PI(\beta)$). As described in (1), MPH intake normalized aberrant beta depression in ADHD, initially lower as compared to controls. In a first model, we aimed at identifying whether a combination of FA values in the selected ROIs accounted for the degree to

which subjects were able to suppress their somatosensory beta power in preparation to a motor response (PI(β)). To this end, we considered PI(β)s for TD subjects and ADHD subjects in the placebo condition as dependent variable in the following model:

$$PI(\beta) \sim b_0 + (b_1 FA_{SLFt} + b_2 FA_{SLFp} + b_3 FA_{ATR}) * b_4 Group + \varepsilon \quad \text{mdl(3)}$$

The resulting overall model was not significant ($p=.762$), neither were the main or the interaction effects in the regressors (FA_{SLFt} : $p=.890$; FA_{SLFp} : $p=.985$; FA_{ATR} : $p=.754$, $FA_{SLFt} * Group$: $p=.934$, $FA_{SLFp} * Group$: $p=.689$, $FA_{ATR} * Group$: $p=.655$).

Next, given the effects of MPH on modulations of the beta oscillations within the ADHD group, we enquired whether, instead, a linear combination of the ROIs' FA properties, could explain the *changes* in beta depression following medication intake ($\Delta PI(\beta)$). We hence implemented the following model, consistently with the principles abovementioned:

$$\Delta PI(\beta) \sim b_0 + (b_1 FA_{SLFT} + b_2 FA_{SLFP} + b_3 FA_{ATR}) * b_4 mg_{MPH} + \varepsilon \quad \text{mdl(4)}$$

Here, we added the term mg_{MPH} as a categorical variable to control for the dosage of MPH administered prior to the task (15/10mg). The resulting model was significant with $R^2=.83$ and $p=.003$ (**Figure 10A**). Analyses of main effects showed a significant negative association between FA_{SLFt} and $\Delta PI(\beta)$ (b_1 : $p=.001$) (**Figure 10B**), and a positive association between FA_{SLFp} and $\Delta PI(\beta)$ (b_2 : $p=1 \times 10^{-4}$) (**Figure 10C**), while no main effects for FA_{ATR} and mg_{MPH} were found ($p=.910$ and $p=.693$, respectively).

The main effects in the model denoted a bigger MPH effect for subjects displaying higher FA in the SLFp, while subjects displaying higher FA in the SLFt were the ones whose sensorimotor beta was less affected by MPH administration.

Although the model produced significant interaction terms between FA_{SLFt} and FA_{SLFp} with mg_{MPH} ($p=.001$ and $p=.002$, respectively), we did not pursue any post-hoc examination of such effect, given a

301 very low ratio between 10/15mg dosage in the sample (0.38) would not produce statistically reliable
302 results.

303 According to the same principle which led to mdl(1.a), we merged the two regressors FA_{SLFp} and FA_{SLFt}
304 into the gradient denoted as $FA_{SLF(p-t)}$, and considered an equivalent to mdl(4) as follows:

$$\Delta PI(\beta) \sim b_0 + (b_1 FA_{SLFT(p-t)} + b_2 FA_{ATR}) * b_3 Group + \varepsilon \quad \text{mdl(4.a)}$$

305 In **Figure 10D** we present an alternative and equivalent representation of the plot in **Figure 10B and**
306 **C**, where the linear association between $\Delta PI(\beta)$ and $FA_{SLF(p-t)}$ is presented. Here, we can observe a
307 significant positive partial association ($p=7 \times 10^{-4}$) which indicates that subjects with a higher parietal
308 compared to temporal FA along the SLF tract, were also the ones whose beta depression was more
309 affected (enhanced) by MPH.

Discussion

In the current study, We employed DTI to estimate microstructural properties of main bilateral fasciculi and explored their role in mediating behavior and beta power modulation associated with ADHD symptomatology.

We first showed that, in both groups, lower values of fractional anisotropy within the ATR were related to ADHD symptom severity and that parieto-to-temporal FA-imbalance within the SLF accounted for behavioral performance in the attentional task. Importantly, in the ADHD group, the same SLF gradient was predictive of the effects of MPH on beta power modulation.

The ATR originates from anterior and medial nuclei of the thalamus and radiates along the anterior thalamic peduncle and the anterior limb of the internal capsule to reach the frontal cortex (54,55). The thalamocortical feedback loop is crucial in conscious processing, and its role in attention is to provide a functional link between otherwise structurally segregated cortical areas, supporting different aspects of attentional selection and working memory (56,57). Prior morphological analyses have shown that ADHD is associated with altered shape and volume of thalamic nuclei, whose connections within several cortical regions, particularly frontal areas, seem disrupted in affected children (58,59). On the other hand, there is still controversy over the validity of thalamic volume as anatomical correlate of the disorder, given that case-control volumetric differences in this area were not confirmed in a recent mega-analysis (60). However, that mega-analysis did provide evidence for a different influence of age on thalamic volumes between clinical and non-clinical samples. The thalamus is a rather complex structure composed of a set of cytoarchitectonically segregated nuclei, each providing specific thalamocortical signals and relying on partially independent neural circuitry to mediate different cognitive functions. Hence, a more valid investigation of thalamic involvement in the pathophysiology of ADHD should take into account an specific analysis of such diverse nuclei (61), so far still absent in the ADHD literature. Arguably, currently the most convincing effects with regard to thalamus role in ADHD, are found in the context of connectivity studies, pointing to the importance of fronto-striatal circuits and their disruption in association to the symptomatology (62). Here, we corroborated these

findings by embracing a different approach: the association between thalamocortical diffusivity and symptom severity we report, exploits the inherent heterogeneity in FA values among subjects, by linking this variance to behavioral symptomatology along the broader clinical spectrum. In other words, our results speak to the importance of thalamocortical connections in the interaction between attention and premotor functions which, if anomalous, are at the basis of clinical behavioral symptoms (as observed in ADHD).

We postulate that the strong interconnections between the thalamus and striatal regions, among which different basal ganglia nuclei, is one of the core variables into consideration when aiming at identifying structural correlates of ADHD. Reinforcing this finding, the TD group in our study displayed stronger anisotropic diffusivity along the ATR as compared to the ADHD group. Such notion further highlights the role of the basal ganglia and their morphological and volumetric differences as potential predictors of the disorder (60,63–68).

The second main and novel finding related the parieto-to-temporal FA-imbalance within the SLF with behavioral performance in the attentional task (in both groups): a higher FA in the parietal-SLF compared to the temporal-SLF predicted faster and more accurate responses in both ADHD and controls. Furthermore, in the ADHD sample, the same gradient explained the effects of MPH on beta modulation: individuals displaying higher FA in the parietal than the temporal SLF were also the ones whose beta power during response preparation increased more with MPH. FA along the SLF likely reflects the functions of the frontoparietal control network (FPCN). This network is one of the core anatomical components providing the basis of flexible attentional adaptations to different task demands (69). While the above results linking FA imbalance within the SLF to behavior and beta oscillations are seemingly orthogonal to the previous finding relating FA_{ATR} to symptoms severity, the former are not independent from thalamo-frontal influences. Indeed, the dopaminergic regulation of the prefrontal cortex and the striatum has been proposed to mediate the interaction between the different attentional systems (27,70,71), some of which structurally rely on the SLF.

Dopaminergic availability is modulated by MPH (21,24,72), which blocks the reuptake of the neurotransmitter, hence allegedly increasing the functional interactions within attentional networks. Given these premises, it is not surprising that diffusivity along the superior longitudinal fasciculus favors the effects of MPH: a stronger connectivity, as indexed by anisotropy along the tract, promotes communication between frontoparietal areas, which is further maximized by the stimulant's action. Crucially, we found that an imbalance of FA in favor of parietal rather than temporal regions is associated with stronger effects of medication on beta modulation. Important nodes of the FPCN are found in dorsolateral prefrontal cortex, frontal eye fields and intraparietal sulcus (73), regions that are connected by the dorsal fibers of the SLF (74). Previous studies have already proposed that the mechanisms of action of stimulant medication in ADHD are strongly reflected by its activity on frontoparietal regions (18,75), whose under-activation is one of the neural correlates of the disorder (76–78).

It is important to consider the link between dopaminergic coordination of attentional networks and increased beta modulation. Evidence from Parkinson's Disease (PD) patients has provided a theoretical framework according to which dopamine levels within the basal ganglia-cortical loop have a direct influence on beta oscillations (79,80). Anomalous beta oscillations are highly correlated with PD pathology (81), with evidence from in vivo recordings and mathematical models suggesting they originate from dopamine degeneration in cortical projections to the striatum (82–84). Coupled with findings of increased striatal dopamine transporter (thus lower dopaminergic availability) in ADHD (85–88), we here propose an attentional network which relies on the dopaminergic input from thalamo-striatal regions to the prefrontal cortex, which in turn mediates the activity along the FPCN. Higher FA along frontoparietal tracts tends to reflect increased connectivity and is being investigated as indirect tool to infer dopaminergic functions in the pathological brain (89–91). Relatively stronger parietal FA may thus underly a facilitatory mechanisms for MPH action in the brain, as revealed by a stronger increase in preparatory beta desynchronization in children displaying higher parietal-to-temporal FA gradient.

Although limited in its sample size, this study offers important new insights in the potential of multimodal imaging to investigate and identify the sources of attention performance in the brain: by coupling evidence from electrophysiological measures and information about white matter integrity, we are able to investigate the origin of aberrant brain activity observed in ADHD and improve our understanding of the mechanisms of action of stimulant medication. Furthermore, it might encourage future research on the issue, clarifying whether FA represents a predictor of MPH responsiveness or it is rather one of the neural targets of medication, as could potentially be inferred by longitudinal research.

Disclosures

Jan K Buitelaar has been in the past 3 years a consultant to / member of advisory board of / and/or speaker for Janssen Cilag BV, Eli Lilly, Medice, Takeda/Shire, Roche, and Servier. He is not an employee of any of these companies, and not a stock shareholder of any of these companies. He has no other financial or material support, including expert testimony, patents, royalties. The remaining authors declare no competing and financial interests.

Acknowledgements

The authors gratefully acknowledge the support of the Marie-Curie ITN grant ChildBrain (grant number 641652).

References

1. Mazzetti C, ter Huurne N, Buitelaar JK, Jensen O. Methylphenidate normalizes aberrant beta oscillations and reduces alpha power during retention in children with ADHD. *bioRxiv* [Internet]. 2020 Jan 1;2020.03.13.990929. Available from: <http://biorxiv.org/content/early/2020/03/15/2020.03.13.990929.abstract>
2. van Ede F, Niklaus M, Nobre AC. Temporal Expectations Guide Dynamic Prioritization in Visual Working Memory through Attenuated α Oscillations. *J Neurosci*. 2017;37(2):437–45.
3. Tzagarakis C, West S, Pellizzer G. Brain oscillatory activity during motor preparation: Effect of directional uncertainty on beta, but not alpha, frequency band. *Front Neurosci*. 2015;9(JUN):1–13.
4. Herring JD, Thut G, Jensen O, Bergmann TO. Attention modulates TMS-locked alpha oscillations in the visual cortex. *J Neurosci*. 2015;35(43):14435–47.
5. van Ede F, de Lange FP, Maris E. Attentional Cues Affect Accuracy and Reaction Time via Different Cognitive and Neural Processes. *J Neurosci* [Internet]. 2012;32(30):10408–12. Available from: <http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.1337-12.2012>
6. Berger C, Müller-Godeffroy J, Marx I, Reis O, Buchmann J, Dück A. Methylphenidate promotes the interaction between motor cortex facilitation and attention in healthy adults: A combined study using event-related potentials and transcranial magnetic stimulation. *Brain Behav*. 2018;8(12):1–15.
7. Doyle LMF, Yarrow K, Brown P. Lateralization of event-related beta desynchronization in the EEG during pre-cued reaction time tasks. *Clin Neurophysiol*. 2005;116(8):1879–88.
8. American Psychiatric Association. Diagnostic and statistical manual of mental disorders : DSM-5. Association AP, Force APAD-5 T, editors. Arlington, VA: American Psychiatric Association; 2013.
9. ter Huurne N, Onnink M, Kan C, Buitelaar J, Jensen O. Diminished modulation of preparatory sensorimotor mu rhythm predicts attention-deficit / hyperactivity disorder severity. *Psychol Med*. 2018;(2017):1947–56.
10. ter Huurne N, Onnink M, Kan C, Franke B, Buitelaar J, Jensen O. Behavioral consequences of aberrant alpha lateralization in attention-deficit/hyperactivity disorder. *Biol Psychiatry* [Internet]. 2013;74(3):227–33. Available from: <http://dx.doi.org/10.1016/j.biopsych.2013.02.001>
11. ter Huurne N, Lozano-Soldevilla D, Onnink M, Kan C, Buitelaar J, Jensen O. Diminished modulation of preparatory sensorimotor mu rhythm predicts attention-deficit/hyperactivity disorder severity. *Psychol Med*. 2017;47(11):1947–56.
12. Arns M, Vollebregt MA, Palmer D, Spooner C, Gordon E, Kohn M, et al. Electroencephalographic biomarkers as predictors of methylphenidate response in attention-deficit/hyperactivity disorder. *Eur Neuropsychopharmacol* [Internet]. 2018;28(8):881–91. Available from: <https://doi.org/10.1016/j.euroneuro.2018.06.002>

13. Vollebregt MA, Zumer JM, ter Huurne N, Buitelaar JK, Jensen O. Posterior alpha oscillations reflect attentional problems in boys with Attention Deficit Hyperactivity Disorder. *Clin Neurophysiol* [Internet]. 2016;127(5):2182–91. Available from: <http://dx.doi.org/10.1016/j.clinph.2016.01.021>
14. Lenartowicz A, Mazaheri A, Jensen O, Loo SK. Aberrant Modulation of Brain Oscillatory Activity and Attentional Impairment in Attention-Deficit/Hyperactivity Disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging* [Internet]. 2018;3(1):19–29. Available from: <https://doi.org/10.1016/j.bpsc.2017.09.009>
15. Scahill L, Schwab-Stone M. Epidemiology of Adhd in School-Age Children. *Child Adolesc Psychiatr Clin N Am* [Internet]. 2000;9(3):541–55. Available from: <http://www.sciencedirect.com/science/article/pii/S1056499318301068>
16. Polanczyk G, Caspi A, Houts R, Kollins SH, Rohde LA, Moffitt TE. Implications of Extending the ADHD Age-of-Onset Criterion to Age 12: Results from a Prospectively Studied Birth Cohort. *J Am Acad Child Adolesc Psychiatry* [Internet]. 2010;49(3):210–6. Available from: <http://dx.doi.org/10.1016/j.jaac.2009.12.014>
17. Mazaheri A, Fassbender C, Coffey-Corina S, Hartanto TA, Schweitzer JB, Mangun GR. Differential oscillatory electroencephalogram between attention-deficit/ hyperactivity disorder subtypes and typically developing adolescents. *Biol Psychiatry* [Internet]. 2014;76(5):422–9. Available from: <http://dx.doi.org/10.1016/j.biopsych.2013.08.023>
18. Zammit N, Muscat R. Beta band oscillatory deficits during working memory encoding in adolescents with attention-deficit hyperactive disorder. *Eur J Neurosci*. 2019;(February):1–16.
19. Richter CG, Coppola R, Bressler SL. Top-down beta oscillatory signaling conveys behavioral context in early visual cortex. *Sci Rep* [Internet]. 2018;8(1):1–12. Available from: <http://dx.doi.org/10.1038/s41598-018-25267-1>
20. Engel AK, Fries P. Beta-band oscillations--signalling the status quo? *Curr Opin Neurobiol* [Internet]. 2010;20(2):156–65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20359884>
21. Faraone S V. The pharmacology of amphetamine and methylphenidate: Relevance to the neurobiology of attention-deficit/hyperactivity disorder and other psychiatric comorbidities. *Neurosci Biobehav Rev* [Internet]. 2018;87(January):255–70. Available from: <https://doi.org/10.1016/j.neubiorev.2018.02.001>
22. Tang B, Dafny N. Dorsal raphe neuronal activities are modulated by methylphenidate. *J Neural Transm*. 2013;120(5):721–31.
23. Hodgkins P, Shaw M, Coghill D, Hechtman L. Amphetamine and methylphenidate medications for attention-deficit/ hyperactivity disorder: Complementary treatment options. *Eur Child Adolesc Psychiatry*. 2012;21(9):477–92.
24. Devilbiss DM, Kelley AE, Schmeichel B, Andrzejewski ME, Berridge CW, Hamilton C, et al. Methylphenidate Preferentially Increases Catecholamine Neurotransmission within the Prefrontal Cortex at Low Doses that Enhance Cognitive Function. *Biol Psychiatry*. 2006;60(10):1111–20.
25. Jarczok TA, Haase R, Bluschke A, Thiemann U, Bender S. Bereitschaftspotential and lateralized readiness potential in children with attention deficit hyperactivity disorder: altered motor system activation and effects of methylphenidate. *Eur Neuropsychopharmacol*

- 465 [Internet]. 2019;29(8):960–70. Available from: <https://doi.org/10.1016/j.euroneuro.2019.05.003>
- 466 26. Zhong P, Liu W, Yan Z. Aberrant regulation of synchronous network activity by the attention-deficit/hyperactivity disorder-
467 associated human dopamine D4 receptor variant D4.7 in the prefrontal cortex. *J Physiol*. 2016;594(1):135–47.
- 468 27. Dang LC, O’Neil JP, Jagust WJ. Dopamine supports coupling of attention-related networks. *J Neurosci*. 2012;32(28):9582–7.
- 469 28. Alexander AL, Lee JE, Lazar M, Field AS. Diffusion tensor imaging of the brain. *Neurotherapeutics* [Internet]. 2007 Jul;4(3):316–
470 29. Available from: <https://www.sciencedirect.com/science/article/pii/S1933721307000955>
- 471 29. Chang EH, Argyelan M, Aggarwal M, Chandon TSS, Karlsgodt KH, Mori S, et al. The role of myelination in measures of white
472 matter integrity: Combination of diffusion tensor imaging and two-photon microscopy of CLARITY intact brains. *Neuroimage*
473 [Internet]. 2017;147(June 2016):253–61. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2016.11.068>
- 474 30. Le Bihan D, Mangin J-F, Poupon C, Clark CA, Pappata S, Molko N, et al. Diffusion tensor imaging: Concepts and applications. *J*
475 *Magn Reson Imaging* [Internet]. 2001 Apr;13(4):534–46. Available from: <http://doi.wiley.com/10.1002/jmri.1076>
- 476 31. Assaf Y, Pasternak O. Diffusion tensor imaging (DTI)-based white matter mapping in brain research: A review. *J Mol Neurosci*.
477 2008;34(1):51–61.
- 478 32. Beaulieu C. The basis of anisotropic water diffusion in the nervous system - A technical review. *NMR Biomed*. 2002;15(7–8):435–
479 55.
- 480 33. Onnink AMH, Zwiers MP, Hoogman M, Mostert JC, Dammers J, Kan CC, et al. Deviant white matter structure in adults with
481 attention-deficit/hyperactivity disorder points to aberrant myelination and affects neuropsychological performance. *Prog Neuro-*
482 *Psychopharmacology Biol Psychiatry* [Internet]. 2015;63:14–22. Available from: <http://dx.doi.org/10.1016/j.pnpbp.2015.04.008>
- 483 34. Lei D, Ma J, Du X, Shen G, Jin X, Gong Q. Microstructural abnormalities in the combined and inattentive subtypes of attention
484 deficit hyperactivity disorder: a diffusion tensor imaging study. *Sci Rep* [Internet]. 2014;4:6875. Available from:
485 <http://www.nature.com/srep/2014/141103/srep06875/full/srep06875.html>
- 486 35. van Ewijk H, Heslenfeld DJ, Zwiers MP, Buitelaar JK, Oosterlaan J. Diffusion tensor imaging in attention deficit/hyperactivity
487 disorder: A systematic review and meta-analysis. *Neurosci Biobehav Rev* [Internet]. 2012;36(4):1093–106. Available from:
488 <http://dx.doi.org/10.1016/j.neubiorev.2012.01.003>
- 489 36. Damatac CG, Chauvin RJM, Zwiers MP, van Rooij D, Akkermans SEA, Naaijen J, et al. White matter microstructure in attention-
490 deficit/hyperactivity disorder: a systematic tractography study in 654 individuals. *bioRxiv* [Internet]. 2020;787713. Available from:
491 <http://biorxiv.org/content/early/2020/03/18/787713.abstract>
- 492 37. Aoki Y, Cortese S, Castellanos FX. Research Review: Diffusion tensor imaging studies of attention-deficit/hyperactivity disorder:
493 meta-analyses and reflections on head motion. *J Child Psychol Psychiatry Allied Discip*. 2018;59(3):193–202.
- 494 38. de Schotten MT, Dell’Acqua F, Forkel SJ, Simmons A, Vergani F, Murphy DGM, et al. A lateralized brain network for visuospatial

495 attention. *Nat Neurosci* [Internet]. 2011;14(10):1245–6. Available from: <http://www.nature.com/doi/10.1038/nn.2905>

496 39. Marshall TR, Bergmann TO, Jensen O. Frontoparietal Structural Connectivity Mediates the Top-Down Control of Neuronal
497 Synchronization Associated with Selective Attention. *PLoS Biol*. 2015;13(10):1–17.

498 40. Wolfers T, Onnink AMH, Zwiers MP, Arias-Vasquez A, Hoogman M, Mostert JC, et al. Lower white matter microstructure in the
499 superior longitudinal fasciculus is associated with increased response time variability in adults with attentiondeficit/hyperactivity
500 disorder. *J Psychiatry Neurosci*. 2015;40(5):344–51.

501 41. Witt ST, Stevens MC. Relationship between white matter microstructure abnormalities and ADHD symptomatology in adolescents.
502 *Psychiatry Res - Neuroimaging* [Internet]. 2015;232(2):168–74. Available from:
503 <http://dx.doi.org/10.1016/j.psychresns.2015.02.009>

504 42. Konrad A, Dielentheis TF, El Masri D, Bayerl M, Fehr C, Gesierich T, et al. Disturbed structural connectivity is related to inattention
505 and impulsivity in adult attention deficit hyperactivity disorder. *Eur J Neurosci*. 2010;31(5):912–9.

506 43. Linssen AMW, Sambeth A, Vuurman EFPM, Riedel WJ. Cognitive effects of methylphenidate in healthy volunteers: A review of
507 single dose studies. *Int J Neuropsychopharmacol*. 2014;17(6):961–77.

508 44. Carlson GA, Kelly KL. Stimulant Rebound: How Common Is It and What Does It Mean? *J Child Adolesc Psychopharmacol*.
509 2003;13(2):137–42.

510 45. Quinn D, Bode T, Reiz JL, Donnelly GAE, Darke AC. Single-dose pharmacokinetics of multilayer-release methylphenidate and
511 immediate-release methylphenidate in children with attention-deficit/ hyperactivity disorder. *J Clin Pharmacol*. 2007;47(6):760–6.

512 46. Yendiki A, Panneck P, Srinivasan P, Stevens A, Zöllei L, Augustinack J, et al. Automated probabilistic reconstruction of white-
513 matter pathways in health and disease using an atlas of the underlying anatomy. *Front Neuroinform*. 2011;5(October):1–12.

514 47. Jbabdi S, Woolrich MW, Andersson JLR, Behrens TEJ. A Bayesian framework for global tractography. *Neuroimage* [Internet].
515 2007;37(1):116–29. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2007.04.039>

516 48. Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, et al. Whole Brain Segmentation: Automated Labeling of
517 Neuroanatomical Structures in the Human Brain. *Neuron*. 2002;33(3):341–55.

518 49. Behrens TEJ, Berg HJ, Jbabdi S, Rushworth MFS, Woolrich MW. Probabilistic diffusion tractography with multiple fibre
519 orientations: What can we gain? *Neuroimage*. 2007;34(1):144–55.

520 50. Mazzetti C, Staudigl T, Marshall TR, Zumer JM, Fallon SJ, Jensen O. Hemispheric asymmetry of globus pallidus relates to alpha
521 modulation in reward-related attentional tasks. *J Neurosci* [Internet]. 2019 Oct 2;610–9. Available from:
522 <http://www.jneurosci.org/content/early/2019/10/02/JNEUROSCI.0610-19.2019.abstract>

523 51. Beare R, Adamson C, Bellgrove MA, Vilgis V, Vance A, Seal ML, et al. Altered structural connectivity in ADHD: a network based
524 analysis. *Brain Imaging Behav*. 2017;11(3):846–58.

- 525 52. Damatac CG, Zwiers M, Chauvin R, Rooij D Van, Sophie EA. Hyperactivity-impulsivity in ADHD is associated with white matter
526 microstructure in the cingulum ' s angular bundle. 2019;1–17.
- 527 53. Ward EJ. A review and comparison of four commonly used Bayesian and maximum likelihood model selection tools. *Ecol Modell.*
528 2008;211(1–2):1–10.
- 529 54. Jones EG. Thalamic circuitry and thalamocortical synchrony. *Philos Trans R Soc B Biol Sci.* 2002;357(1428):1659–73.
- 530 55. George K, M.Das J. Neuroanatomy, Thalamocortical Radiations. *StatPearls Publ [Internet].* 2019;January. Available from:
531 <https://www.ncbi.nlm.nih.gov/books/NBK546699/>
- 532 56. Halassa MM, Kastner S. Thalamic functions in distributed cognitive control. *Nat Neurosci [Internet].* 2017;20(12):1669–79.
533 Available from: <http://dx.doi.org/10.1038/s41593-017-0020-1>
- 534 57. Sherman SM. Thalamus plays a central role in ongoing cortical functioning. *Nat Neurosci.* 2016;19(4):533–41.
- 535 58. Xia S, Li X, Kimball AE, Kelly MS, Lesser I, Branch C. Thalamic shape and connectivity abnormalities in children with attention-
536 deficit/hyperactivity disorder. *Psychiatry Res - Neuroimaging [Internet].* 2012;204(2–3):161–7. Available from:
537 <http://dx.doi.org/10.1016/j.psychres.2012.04.011>
- 538 59. Svatkova A, Nestrasil I, Rudser K, Goldenring Fine J, Bledsoe J, Semrud-Clikeman M. Unique white matter microstructural patterns
539 in ADHD presentations—a diffusion tensor imaging study. *Hum Brain Mapp.* 2016;37(9):3323–36.
- 540 60. Hoogman M, Bralten J, Hibar DP, Mennes M, Zwiers MP, Schweren LSJ, et al. Subcortical brain volume differences in participants
541 with attention deficit hyperactivity disorder in children and adults: a cross-sectional mega-analysis. *The Lancet Psychiatry.*
542 2017;4(4):310–9.
- 543 61. Battistella G, Najdenovska E, Maeder P, Ghazaleh N, Daducci A, Thiran JP, et al. Robust thalamic nuclei segmentation method
544 based on local diffusion magnetic resonance properties. *Brain Struct Funct.* 2017;222(5):2203–16.
- 545 62. Cupertino RB, Soheili-Nezhad S, Grevet EH, Bandeira CE, Picon FA, Tavares ME, et al. Reduced fronto-striatal volume in ADHD
546 in two cohorts across the lifespan. *bioRxiv [Internet].* 2019;i(51):790204. Available from:
547 <https://www.biorxiv.org/content/10.1101/790204v1>
- 548 63. Paclt I, Přibilová N, Kollárová P, Kohoutová M, Dezortová M, Hájek M, et al. Reverse asymmetry and changes in brain structural
549 volume of the basal ganglia in ADHD, developmental changes and the impact of stimulant medications. *Neuroendocrinol Lett.*
550 2016;37(1):29–32.
- 551 64. Aylward EH, Reiss AL, Reader MJ, Singer HS, Brown JE, Denckla MB. Basal ganglia volumes in children with attention-deficit
552 hyperactivity disorder. *J Child Neurol [Internet].* 1996;11(2):112–5. Available from:
553 http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=8881987
- 554 65. Qiu A, Crocetti D, Adler M, Mahone EM, Denckla MB, Miller MI, et al. Basal ganglia volume and shape in children with attention

555 deficit hyperactivity disorder. *Am J Psychiatry*. 2009;166(1):74–82.

556 66. Sethi A, Evelyn-Rahr E, Dowell N, Jain S, Voon V, Critchley HD, et al. Magnetization transfer imaging identifies basal ganglia
557 abnormalities in adult ADHD that are invisible to conventional T1 weighted voxel-based morphometry. *NeuroImage Clin* [Internet].
558 2017;15(March):8–14. Available from: <https://doi.org/10.1016/j.nicl.2017.03.012>

559 67. Von Rhein D, Oldehinkel M, Beckmann CF, Oosterlaan J, Heslenfeld D, Hartman CA, et al. Aberrant local striatal functional
560 connectivity in attention-deficit/hyperactivity disorder. *J Child Psychol Psychiatry Allied Discip*. 2016;57(6):697–705.

561 68. Oldehinkel M, Beckmann CF, Pruim RHR, Van Oort ESB, Franke B, Hartman CA, et al. Attention-Deficit/Hyperactivity Disorder
562 Symptoms Coincide with Altered Striatal Connectivity. *Biol Psychiatry Cogn Neurosci Neuroimaging* [Internet]. 2016;1(4):353–
563 63. Available from: <http://dx.doi.org/10.1016/j.bpsc.2016.03.008>

564 69. Dixon ML, De La Vega A, Mills C, Andrews-Hanna J, Spreng RN, Cole MW, et al. Heterogeneity within the frontoparietal control
565 network and its relationship to the default and dorsal attention networks. *Proc Natl Acad Sci U S A*. 2018;115(7):E1598–607.

566 70. Jenni NL, Larkin JD, Floresco SB. Prefrontal dopamine D1 and D2 receptors regulate dissociable aspects of decision making via
567 distinct ventral striatal and amygdalar circuits. *J Neurosci*. 2017;37(26):6200–13.

568 71. Tomasi D, Wang GJ, Volkow ND. Association between striatal dopamine D2/D3 receptors and brain activation during visual
569 attention: effects of sleep deprivation. *Transl Psychiatry*. 2016;6(5):e828.

570 72. Arnsten AFT. Stimulants: Therapeutic actions in ADHD. *Neuropsychopharmacology*. 2006;31(11):2376–83.

571 73. Xuan B, Mackie MA, Spagna A, Wu T, Tian Y, Hof PR, et al. The activation of interactive attentional networks. *Neuroimage*
572 [Internet]. 2016;129:308–19. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2016.01.017>

573 74. Makris N, Kennedy DN, McNerney S, Sorensen AG, Wang R, Caviness VS, et al. Segmentation of subcomponents within the
574 superior longitudinal fascicle in humans: A quantitative, in vivo, DT-MRI study. *Cereb Cortex*. 2005;15(6):854–69.

575 75. Kowalczyk OS, Cubillo AI, Smith A, Barrett N, Giampietro V, Brammer M, et al. Methylphenidate and atomoxetine normalise
576 fronto-parietal underactivation during sustained attention in ADHD adolescents. *Eur Neuropsychopharmacol* [Internet].
577 2019;29(10):1102–16. Available from: <https://doi.org/10.1016/j.euroneuro.2019.07.139>

578 76. Rubia K, Halari R, Cubillo A, Smith AB, Mohammad AM, Brammer M, et al. Methylphenidate normalizes fronto-striatal
579 underactivation during interference inhibition in medication-naïve boys with attention-deficit hyperactivity disorder.
580 *Neuropsychopharmacology* [Internet]. 2011;36(8):1575–86. Available from: <http://dx.doi.org/10.1038/npp.2011.30>

581 77. Chiang HL, Chen YJ, Lo YC, Tseng WYI, Gau SSF. Altered white matter tract property related to impaired focused attention,
582 sustained attention, cognitive impulsivity and vigilance in attention-deficit/hyperactivity disorder. *J Psychiatry Neurosci*.
583 2015;40(5):325–35.

584 78. Cai W, Griffiths K, Korgaonkar MS, Williams LM, Menon V. Inhibition-related modulation of salience and frontoparietal networks

- predicts cognitive control ability and inattention symptoms in children with ADHD. *Mol Psychiatry*. 2019;
79. Jenkinson N, Brown P. New insights into the relationship between dopamine, beta oscillations and motor function. *Trends Neurosci* [Internet]. 2011;34(12):611–8. Available from: <http://dx.doi.org/10.1016/j.tins.2011.09.003>
80. Moran RJ, Mallet N, Litvak V, Dolan RJ, Magill PJ, Friston KJ, et al. Alterations in brain connectivity underlying beta oscillations in parkinsonism. *PLoS Comput Biol*. 2011;7(8).
81. Cole SR, van der Meij R, Peterson EJ, de Hemptinne C, Starr PA, Voytek B. Nonsinusoidal beta oscillations reflect cortical pathophysiology in parkinson's disease. *J Neurosci*. 2017;37(18):4830–40.
82. McCarthy MM, Moore-Kochlacs C, Gu X, Boyden ES, Han X, Kopell N. Striatal origin of the pathologic beta oscillations in Parkinson's disease. *Proc Natl Acad Sci U S A*. 2011;108(28):11620–5.
83. Pavlides A, Hogan SJ, Bogacz R. Computational Models Describing Possible Mechanisms for Generation of Excessive Beta Oscillations in Parkinson's Disease. *PLoS Comput Biol*. 2015;11(12):1–29.
84. Wang DD, de Hemptinne C, Miocinovic S, Ostrem JL, Galifianakis NB, Luciano MS, et al. Pallidal deep-brain stimulation disrupts pallidal beta oscillations and coherence with primary motor cortex in Parkinson's disease. *J Neurosci*. 2018;38(19):4556–68.
85. Krause KH, Dresel SH, Krause J, Kung HF, Tatsch K. Increased striatal dopamine transporter in adult patients with attention deficit hyperactivity disorder: Effects of methylphenidate as measured by single photon emission computed tomography. *Neurosci Lett*. 2000;285(2):107–10.
86. Jucaite A, Fernell E, Halldin C, Forssberg H, Farde L. Reduced midbrain dopamine transporter binding in male adolescents with attention-deficit/hyperactivity disorder: Association between striatal dopamine markers and motor hyperactivity. *Biol Psychiatry*. 2005;57(3):229–38.
87. Tai YC, Chi MH, Chu CL, Chiu NT, Yao WJ, Chen PS, et al. Availability of Striatal Dopamine Transporter in Healthy Individuals With and Without a Family History of ADHD. *J Atten Disord*. 2019;23(7):665–70.
88. Hesse S, Ballaschke O, Barthel H, Sabri O. Dopamine transporter imaging in adult patients with attention-deficit/hyperactivity disorder. *Psychiatry Res - Neuroimaging* [Internet]. 2009;171(2):120–8. Available from: <http://dx.doi.org/10.1016/j.psychresns.2008.01.002>
89. Lenfeldt N, Eriksson J, Åström B, Forsgren L, Mo SJ. Fractional Anisotropy and Mean Diffusion as Measures of Dopaminergic Function in Parkinson's Disease: Challenging Results. *J Parkinsons Dis* [Internet]. 2017 Feb 7;7(1):129–42. Available from: <https://www.medra.org/servlet/aliasResolver?alias=iospress&doi=10.3233/JPD-161011>
90. Lorio S, Sambataro F, Bertolino A, Draganski B, Dukart J. The Combination of DAT-SPECT, Structural and Diffusion MRI Predicts Clinical Progression in Parkinson's Disease. *Front Aging Neurosci*. 2019;11(March):1–13.
91. Shimony JS, Rutlin J, Karimi M, Tian L, Snyder AZ, Loftin SK, et al. Validation of diffusion tensor imaging measures of

615 nigrostriatal neurons in macaques. PLoS One [Internet]. 2018 Sep 5;13(9):e0202201. Available from:
616 <https://doi.org/10.1371/journal.pone.0202201>

617

618

Figure 1. Experimental study design. ^a Case Report Form and Behavioural test (line bisection task, WISCIII *Vocabulary* and *Block design* subscales, ADHD rating scale, CBCL); ^b Psychiatric intake (basic medical screening and dosage determination); ^c Polhemus digitizer; ^d Medication intake (1 hour prior to the beginning of experimental task); Participants in the ADHD and Controls group visited the laboratory three and two times, respectively. During the first visit (day I) the psychiatric assessment took place, which determined participants' suitability for the study. The same day, the dummy MR took place, followed by the actual MR scan. The second day (day II) both groups performed the attentional task while electromagnetic activity was recorded with the MEG. The ADHD group performed the task twice (day II and day III), once upon administration of MPH, and once upon administration of a placebo pill. This was done according to a randomized crossover design, half of the participants received the MPH on day II,

619 Figures

Figure 2. Attentional task. (Adapted from Mazzetti et al., 2020). The paradigm consisted in a child-friendly adaptation of a Posner cueing paradigm for the study of spatial orienting of attention. Each trial (370 in total) began with the presentation of a fish in the middle of the screen, serving as fixation cross. An eye tracker ensured that the children kept proper fixation throughout the whole trial (as trials were stopped in case the subject performed a saccade). A cue was then presented for 200ms, represented by the fish looking either at the left or right side of the screen (cue side equally distributed across trials). After a preparation interval jittered in the range 1100 – 1500ms, the stimuli were then presented for 300ms, on the two sides of the screen. The child was asked to respond, via button press, indicating the position of the target (shark with an open mouth), while ignoring the distractor on the other side (shark with mouth closed). A positive feedback (happy fish) was then presented if a correct answer was provided within the response interval (1100ms). In case of wrong or no response, a negative feedback was presented (fish bone).

Figure 3- 3D rendering of cortical parcellation based on Desikan-Killiany atlas in one sample TD subject. A cortical parcellation was generated prior to tract estimation, which are combined with prior distributions on the neighboring anatomical structures of each pathway and subcortical segmentation to constrain the tractography solutions (obviating the need for user interaction thus automating the process).

Figure 4 – 3D isosurface rendering and 2D orthographic view of tract reconstructions in one sample subject obtained using TRACULA. Visualization of the probability distributions of all white-matter pathways simultaneously overlaid on 4D brain mask. All 18 tracts are displayed at 20% of their maximum threshold

Figure 5 – Correlation matrix of FA across bilateral white matter structures shows that no negative association subsists between segmented tracts.

632

Figure 6. Beta modulation indices in the three conditions (adapted from (1)). Topographic plot (left) and respective time frequency representations (TFRs) (right panel) of power modulation (β -MI) for the typically developing group (TD) (A), ADHD_{MPH} group (B) and ADHD_{PLA} group (C). Red dots superimposed on the topographies denote sensors of interest as defined in Figure 4A. Notably, beta preparation is stronger in the TD group, while progressively decreases in the ADHD_{MPH} group, being weakest in the ADHD_{PLA} group.

Figure 7 – 3D rendering of white matter ROI. Visualization of the probability distributions of the ROIs identified as tracts of interest according to the model selection approach: SLFp, SLFt and ATR.

Figure 8 – FA in SLFp and SLFt predict behavioral performance in the task. A. The bar plot displays the beta coefficients associated with the linear mixed model mdl(1), where mean FA values within the identified tracts of interest are set as explanatory variables for behavioral performance, as indexed by the IES. Error bars indicate standard error of the mean. The adjusted response plots in B and C show, respectively, the behavioral performance (IES) as a function of the FASLft and, FASLfp, while averaging over other regressors in the model in (A). D. Adjusted response plot displaying the association between IES and parietotemporal gradient SLF(p-t), averaged over the residual regressors (mdl(1.a)). Positive SLF(p-t) values indicate higher FA along parietal as compared to temporal endings within the SLF.

Figure 9 – FA in ATR predicts ADHD symptoms severity in all subjects. A. Bar plot shows beta coefficients associated with mdl(2), where mean FA values along the tracts of interest are defined as predictors for ADHD-RS symptoms score in all subjects. FAATR is associated with a significant partial regression coefficient ($p=0.007$). Scatter plots in B and C show, respectively, adjusted response plot averaged over the other predictors and the added variable plot (partial regression), of symptoms as a function of FAATR. Lower diffusivity along the ATR corresponded to higher average ADHD symptomatology along the spectrum.

635

Figure 10 – FA in SLFp and SLFt predict MPH effects on β depression in the ADHD group. A. Bar plot shows beta regression coefficients associated with mdl(4), where mean FA values along the tracts of interest are defined as predictors for changes in beta modulation due to medication intake ($\Delta PI(\beta)$). According to Eq.(2), a higher $\Delta PI(\beta)$ value for a given ADHD subject reflects bigger changes in depression of beta oscillations following medication intake. B. Adjusted response plot showing $\Delta PI(\beta)$ as a function of FA_{SLFt} . A significant beta coefficient ($p=.001$) reveals that FA_{SLFt} predicted lower changes in β power depression due to MPH. C. Adjusted response plot showing $\Delta PI(\beta)$ as a function of FA_{SLFp} . Higher FA_{SLFp} corresponded to stronger β depression changes following MPH intake ($p=1 \times 10^{-4}$). D. Adjusted response plot illustrating $\Delta PI(\beta)$ as a function of parieto-temporal SLF gradient according to mdl(4.a). Higher SLF(p-t), for a given ADHD subject, reflected higher diffusivity at parietal endings, as compared to temporal endings, of the SLF. The significant positive relationship ($p=7 \times 10^{-4}$) suggests that the gradient of parieto-temporal diffusivity in the SLF is predictive of the effects of MPH on β depression.