Changes in biochemistry and histochemical characteristics during somatic embryogenesis in *Ormosia henryi* Prain

Wu Gaoyin, Wei Xiaoli*, Wang Xiao, Liang Xian, Wei Yi

College of Forestry/Institute for Forest Resources & Environment of Guizhou, Guizhou University, Guiyang 550025, Guizhou, People's Republic of China.

*Corresponding authors: gdwxl-69@126.com; wugaoyin1234@163.com

Author contributions:

Wei X.L conceived the project, designed and supervised the experiments. Wu G.Y., Wang X., Liang X and Wei Y performed the experiments. Wu G.Y and Wang X analyzed data and wrote the manuscript. Wei X.L edited and revised this manuscript.

One-sentence summary:

Somatic embryogenesis and abnormal callus tissues formation of *Ormosia henryi* Prain were associated with energy substances, reactive oxygen species, enzyme activities and endogenous hormones, as well as histochemical characteristics.

Funding information:

This study was funded by the National Natural Science Foundation of China (31460193) and the High Level Innovative Talents Training Programme of Guizhou Province ([2016]5661).

ABSTRACT

Mature embryos were used as an explant for embryogenic callus (EC) induction, and then EC was further developed to form somatic embryos during somatic embryogenesis (SE) of *Ormosia henryi* Prain; however, some mature embryos could induced non-embryogenic callus (NEC), browning callus (BC) or snowflake callus (SC). These phenomena might be related to the biochemical and histochemical differences during somatic embryo induction. The present study was conducted to analyze the biochemical events and histochemical changes at different SE stages in *O. henryi*. The contents of soluble sugar, starch, soluble protein, H₂O₂, and endogenous hormones and the activities of polyphenoloxidase (PPO), superoxide dismutase (SOD), ascorbate peroxidase (APX), peroxidase (POD) and catalase (CAT) were measured at different SE stages, such as EC, globular embryo (GE), and cotyledon embryo (CE), and in abnormal tissue, such as NEC, BC, and SC. The results showed that the contents of soluble sugar and starch; the activities of PPO, SOD, APX and POD; and the ratios of indole-3-acetic acid/abscisic acid (IAA/ABA), IAA/gibberellins (IAA/GAs), auxin /GAs (AUX/GAs), and AUX/ABA decreased gradually at different SE stages. In contrast, the contents of soluble protein, H₂O₂, all endogenous hormones gradually increased. However, CAT activity and the ratios of IAA/cytokinins (IAA/CKs), AUX/CKs, ABA/CKs, and GAs/CKs first increased and then decreased. The high contents of GAs and ABA, high ratios of ABA/CKs and GAs/CKs and low ratios of IAA/ABA, IAA/GAs, AUX/GAs and AUX/ABA were responsible for the inability of the callus to form EC. The low enzyme activities, low contents of energy substances and H₂O₂ were related to NEC formation. The high contents of soluble sugar, H₂O₂, AUX, CKs and PPO activity and the low content of soluble protein were the basic causes of BC formation. The high-energy substances contents and low activities of SOD and POD facilitated SC formation . Histochemical observation showed that starch granule staining gradually lightened with SE development, but protein granules were darkly stained. Compared with EC, starch and protein granules were stained darker in SC, and lighter in NEC and BC. These results showed that energy substances were the material basis of SE, which affected enzyme

activities, regulated reactive oxygen species (ROS) metabolism, and thus regulated the morphogenesis and development of somatic embryos. In addition, the contents and ratios of endogenous hormones affected the dedifferentiation, dedifferentiation and embryogenesis of somatic cells. To induce EC from mature embryos and further develop their formation into somatic embryos, it is necessary to adjust the energy supply and hormone ratio in the medium.

Keywords: *Ormosia henryi*, Somatic embryogenesis, Biochemical events, Histochemical observation.

INTRODUCTION

Somatic embryogenesis (SE) is an efficient approach for the clonal propagation of plants (Correia et al., 2011), Its developmental process resembles zygotic embryo development (Leljak-Levanic et al., 2004), which is an ideal material for studying morphology, physiology, biochemistry and molecular biology in embryogenesis and development (Morel et al., 2014). Since the late 1950s, Steward (1958) found that carrot root cells could be regenerated through SE in vitro, and the SE for many plants has been studied. *O. henryi* is a precious tree species in China and has great economic and ecological potential as a raw material for making high-grade furniture and handicrafts; some factors restrict the scale propagation of *O. henryi*, such as scarce wild resources, biennial fruiting, dormancy caused by the hardness seed coat and pests that eat the seeds (Wei et al., 2014). Therefore, it is very important to research asexual propagation for *O. henryi*. Our previous research has established the SE protocols of *O. henryi* (Wu et al., 2020), but the underlying mechanism is not well understood. During the process of callus subculture, callus is prone to browning and eventually necrosis, and its transformation into SC is a common phenomenon, seriously

hindering the further development of callus. Manivannan et al.(2015) and Roowi et al.(2010) reported that the morphogenesis and external morphological characteristics of SE were regulated by complex biochemistry and molecules inside the cell tissues. Therefore, it is very important to understand its physiological mechanism in the regulation of this process.

Sugar is an indispensable energy substance and carbon skeleton for embryonic development (Bartos et al., 2018), and it can regulate cell osmotic pressure (Iraqi and Tremblay, 2001), protect cell membrane integrity (Bartos et al., 2018), and determine the shape and developmental fate of cell (Baluska et al., 2003). Proteins affect embryo development (Cangahuala-Inocente et al., 2014), embryo morphogenesis (Cangahuala-Inocente et al., 2014), embryo maturation (Jiménez, 2001) and cell signal transduction (Cangahuala-Inocente et al., 2014). Previous studies have confirmed that the changes in their contents could be used as a marker of different SE stages (Jiménez, 2001; Cangahuala-Inocente et al., 2014).

ROS, such as $\cdot\text{OH}$, H_2O_2 , and O_2^- continuously occur in mitochondria, chloroplasts and peroxisomes of plant cells as byproducts of various metabolic pathways of respiration and photosynthesis (Libik et al., 2005; Zhang et al., 2010). H_2O_2 plays the role of a secondary messenger in cell signaling transduction, affects gene expression (Libik et al., 2005) and stimulates embryonic cell formation (Cui et al., 1999). However, in vitro culture of explants provides a stressful environment, which can stimulate the overproduction of ROS (Fatima et al., 2011), and interfere with intracellular redox homeostasis, normal physiological activities and cell metabolism (Cui et al., 1999). Plants have an antioxidant system to remove excess ROS to maintain the balance of ROS contents in cells (Manivannan et al., 2015). In recent years, changes in the activities of SOD, CAT, POD and APX had been shown to be highly correlated with the ROS elimination at different SE stages (Cui et al., 1999; Libik et al., 2005).

Exogenous hormones play an irreplaceable role in SE induction, and their levels affect the balance of endogenous hormones in plants, regulate plant growth and morphogenesis (Guo et al., 2017), and even determine the developmental fate of cells

(Grzyb et al., 2018). Auxin triggers SE induction (Nic-Can and Loyola-Vargas, 2016), and its asymmetric distribution plays key roles in the polar transport of auxin, stem cell formation (Su and Zhang, 2009) and SE induction (Choi et al., 2001). Like auxin, cytokinin is a key regulatory factor for SE (Jiménez, 2005), and the balance between cytokinin and auxin determines the dedifferentiation and redifferentiation states of cells. In addition, ABA and GAs play a positive role in SE maturation and germination (Zhang et al., 2010; Kim et al., 2019). Although endogenous hormones have made some achievements in SE induction and regulation, their regulatory mechanism is still not well understood.

Biochemical events during SE in *O. henryi* were determined to gain a better understanding of the role of the energy substance, ROS, enzyme activities and endogenous hormones in triggering the embryogenic pathway, and they could serve as markers in the determination of different stages of somatic embryo development. Moreover, the histochemical changes in the accumulation and distribution of starch and protein granules were examined at different SE stages, and the changes in starch and protein contents were assessed at different SE stages based on the quantitative and qualitative analysis.

RESULTS

Energy substance

The contents of soluble sugar, starch and protein were significantly different at different SE stages (EC -GE -CE) (Fig .2). The contents of soluble sugar and starch decreased gradually with SE development, and the soluble protein contents increased, which indicated that carbohydrate substances were consumed by somatic embryo growth and development while the increase in protein content offered material conditions for the maturation and germination of somatic embryos. Compared with EC, the contents of soluble sugar, starch and protein were lower in NEC, which

indicated that energy substances were the material basis for EC transformation to the somatic embryo; in contrast, the contents of soluble sugar, starch and protein were significantly higher in SC than EC, indicating that the high-energy substance contents facilitated SC formation; however, the soluble protein content was lowest in BC, which indicated that BC activity was low, and might be a cause of its final necrosis.

H₂O₂ contents and enzyme activities

The H₂O₂ content and activities of PPO, SOD, APX, POD, and CAT were significantly different at different SE stages (Fig. 3). H₂O₂ content increased gradually, whereas the activities of PPO, SOD, APX and POD decreased gradually with SE development. CAT activity, however, was the highest in GE. It is worth noting that APX activity decreased sharply in GE and CE, which was 15% and 23% of EC, respectively. There were significant differences in APX and CAT activities between EC and GE. Among them, APX activity in EC was 570% higher than in GE, and CAT activity was 89% higher in GE than EC.

An increase in contents by 36% (H₂O₂) and activities by 63% (PPO), 13% (SOD), 64% (APX), 48% (POD), and 115% (CAT) were noticed in EC compared with NEC, which indicated that the high H₂O₂ contents and the high enzyme activities could promote SE. The H₂O₂ content and PPO activity were highest in BC, which might be the fundamental reason for callus turning into BC. Compared with EC, SOD and POD activities of SC were significantly decreased, which indicated that the low activities of SOD and POD affected SC formation.

Endogenous hormones

The contents of AUX, CKs, GAs and ABA were significantly different at different SE stages (Fig. 4). Their total contents showed an increasing trend. Except for AUX total contents, which were lowest in GE, the total contents of all endogenous hormones

were lowest in EC and highest in CE. These findings indicated that a relatively high AUX content and low contents of CKs, GAs and ABA were conducive to EC induction, a high AUX content was the premise for somatic embryo development and polarity establishment, a high CKs content regulated the differentiation of somatic embryos in the latter stage, and high contents of GAs and ABA promoted the maturation and germination of somatic embryos in the latter stage.

Compared with EC, the contents of AUX and CKs were similar in NEC and SC, and IAA contents in AUX and DZ contents in CKs were higher in EC; the contents of AUX and CKs in BC were significantly higher than those in EC. In addition, the contents of ME-IAA and tZ were lower in EC than NEC, BC and SC. These results showed that NEC formation was related to the low IAA content; the callus with high contents of AUX and CKs was not conducive to EC formation; while the callus with low contents of IAA and DZ, and high tZ content could easily form SC. Moreover, the contents of GAs and ABA in NEC, BC and SC were more than 6 times that of EC, which indicated that the high contents of GAs and ABA inhibited EC induction.

Endogenous hormones ratios

The interaction between endogenous hormones can be expressed by their ratios to evaluate the balance of endogenous hormones at different SE stages. The ratios of IAA/ABA, IAA/GAs, AUX/GAs and AUX/ABA decreased gradually. The ratios of IAA/CKs, AUX/CKs, ABA/CKs and GAs/CKs first increased and then decreased. The ratios of IAA/CKs and AUX/CKs were the lowest in CE, and the ratios of ABA/CKs, GAs/CKs were lowest in EC. These results indicated that the high ratios of IAA/ABA, IAA/GAs, AUX/GAs, and AUX/ABA and low ratios of ABA/CKs and GAs/CKs were conducive to EC induction at an early stage of SE. The low ratios of IAA/CKs and AUX/CKs promoted maturation and germination of the somatic embryo.

Compared with EC, the ratios of IAA/ABA, IAA/GAs, AUX/GAs, and AUX/ABA decreased significantly in NEC, BC, and SC, and the ratios of ABA/CKs and GAs/CKs were higher. In addition, the ABA/GAs ratio was higher in NEC and

lower in BC. These findings indicated that the low ratios of IAA/ABA, IAA/GAs, AUX/GAs, and AUX/ABA and the high ratios of ABA/CKs and GAs/CKs were responsible for the inability of the callus to form EC; however, an ABA/GAs ratio that was too high or too low ABA/GAs ratio was not conducive to SE induction.

Histochemical studies

Starch granules

As shown in Fig.5, starch granules were mainly distributed in the cell wall and intercellular space, followed by an uneven distribution in cytoplasm and reduced levels in the nucleus. These findings showed that carbohydrates were the main components of the carbon skeleton, and cytoplasm was the main site of carbohydrate synthesis.

At different SE stages, the starch granules staining was dark in the EC, and starch granule staining of EC masses (Fig. 5-c) was especially darker than in the surrounding tissues, which indicated that EC masses stored sufficient nutrients for the growth and development of SE. The starch granule staining gradually lightened with SE development, but the meristem staining (Fig. 5f-i) was dark, which confirmed that the meristem was active and might provide the best explanation for the strong embryogenic response of the shoot apical meristem (SAM).

Compared with EC, the starch granule staining was lightest in BC, followed by NEC, but it was darker in SC. These findings indicated that the abundant energy substances were the basis for callus formation of EC, but the callus with excessive energy substances could easily form SC.

Protein granules

As shown in Fig.6, the protein granule staining was relatively light, which was most obvious in the nucleus, followed by the cytoplasm, which indicated that proteins in

the nucleus were most abundant. This feature was related to the need for a large number of proteins and enzymes for DNA replication and transcription in the nucleus.

With SE development, the protein grain staining became gradually darkened (Fig. 6d-h), with the darkest staining observed in CE. More protein grains accumulated in the meristems, which indicated their strong activity and vigorous metabolism.

Compared with EC, the protein granules staining were light in NEC and BC, and with greater numbers of vacuolar cells, the numbers of nuclei in the cell were reduced; however, it was darker in SC. These results indicated that the callus with the low protein content and activity affected the ability of the callus to form EC, but the callus with the high protein content could easily form SC.

DISCUSSION

Energy substances

C and N participate in the metabolic energy, carbon skeletons and biochemical signaling during plant growth and development (Bartos et al., 2018), and they are closely related to the growth and development of SE and morphogenesis (Cangahuala-Inocente et al., 2014). In this study, the contents of soluble sugar and starch gradually decreased with SE development, similarly to findings reported in *Acca sellowiana* (Cangahuala-Inocente et al., 2014) and spruce (Hazubska-Przybyl et al., 2016). This decrease was related to the consumption of a large amount of energy for the growth and development of SE. However, the protein content increased with SE development, similarly to findings reported in *Eucalyptus globulus* (Pinto et al., 2010), *Coffea arabica* (Bartos et al., 2018), *Catharanthus roseus* (Fatima et al., 2011) and *Cordyline australis* (Warchoř et al., 2015). These phenomena could be related to the synthesis of late embryogenesis abundant protein (LEA) and reserve proteins, which might play an important role in embryonic development and protect embryos against dehydration (Gomes et al., 2014); however, the protein content in late stages of SE were decreased in *Acca sellowiana* (Cangahuala-Inocente et al., 2014) and *Persian walnut* (Jariteh et al., 2015).

EC induction was the basis of SE. The contents of soluble sugar, starch and protein were higher in EC than NEC, similarly to findings reported in banana (Wang et al., 2013), alfalfa (Martin et al., 2000) and *Catharanthus roseus* (Fatima et al., 2011), because it was difficult or impossible for NEC to develop into the somatic embryo. These findings showed that the accumulation of energy substances was the material basis for EC transformation into the somatic embryo. Compared with EC, the contents of starch and protein were lower in BC, which indicated that BC activity was low and thus affected the synthesis of starch and protein, which may explain the final necrosis of BC. In contrast, the contents of soluble sugar, starch and protein increased significantly in SC, but SC could not further develop into somatic embryos, which confirmed that the accumulation of energy substances affected the cell activity and morphogenesis. These results indicated that too low or too high contents of energy substances were not conducive to the further growth of callus. A previous study confirmed that the sucrose concentration affected carbon metabolism in EC, and medium supplemented with different sucrose concentrations was needed to reduce the frequency of SC, although the frequency of SC (9%) was significantly reduced at a low sucrose concentration (20g/L), the SE induction rate (19%) also reduced.

H₂O₂ contents and enzyme activities

The H₂O₂ content increased in the late stage of SE, and similar results were reported in wolfberry (Cui et al., 1999) and Begonia (Manivannan et al., 2015), which indicated that ROS metabolism played a decisive role in cell differentiation and development (Cui et al., 1999). The activities of SOD and APX decreased in GE and CE. Fatima et al.(2011) reported similar results in *Catharanthus roseus*; APX activity significantly decreased, which might be related to the high activities of POD and CAT in GE and CE, which consumed excessive H₂O₂, leading to the decrease in APX activity. A similar phenomenon was reported in *Citrus reticulata* (Liu, 2016).

H₂O₂ content and the activities of PPO, SOD, APX, POD and CAT in the EC of *O. henryi* were higher than those in NEC, which indicated that the callus with the low H₂O₂ content and low enzyme activities were not sufficient to induce SE. The H₂O₂

accumulation could trigger the stress response, play a role as a secondary messenger involved in cell signaling and promote SE development. Similar results have been reported in *Albizia lebbeck* (Saeed and Shahzad, 2015) and *Mesembryanthemum crystallinum* (Libik et al., 2005). Compared with EC, higher H₂O₂ contents, higher PPO activity and lower antioxidant enzyme activities were detected in BC, which caused BC to be in a state of high oxidative stress. Wichers et al.(1985) reported that PPO activity increased in cell culture of *Mucuna pruriens* during cell senescence, which might represent the main reasons for callus transformation into BC and even necrosis. To reduce the BC frequency, medium supplemented with anti-browning agents, such as polyvinylpyrrolidone, ascorbic acid and activated carbon, reduced the frequency of callus transformation into BC, but the results were not significant differences (data not shown). Compared with EC, SOD and POD activities decreased significantly in SC, and therefore, we speculated that callus with low SOD and POD activities could easily form SC. Fatima et al.(2011) reported that POD was a marker involved in cell morphogenesis.

Endogenous hormones

Endogenous hormone levels regulate the dedifferentiation, redifferentiation and embryogenesis of somatic cells at different SE stages (Grzyb et al., 2018). At different SE stages of *O. henryi*, IAA content increased significantly in CE, which was consistent with the results of SE maturation of *Norway Spruce* (Vondrakova et al., 2018) and larch (Aderkas et al., 2001), indicating that auxin is related to the establishment of embryo polarity (Vondrakova et al., 2018). Auxin is required for SE maturation to mediate SAM by establishing the homeodomain transcription factor WUSCHEL (WUS), which is essential for stem cell initiation and meristem maintenance (Lee and Clark, 2015). Similarly, CKs contents increased significantly in CE, which is consistent with a report on SE maturation and germination in spruce (Vondrakova et al., 2018). Recent advances have shown that CKs are essential factors that regulate shoot initiation via WUS expression (Cheng et al., 2013), and the CKs response factors ARABIDOPSIS RESPONSE REGULATORS (ARRs) directly

activate WUS expression and thus initiate SAM formation (Zhang et al., 2017; Liu et al., 2018). Some studies had shown that high GAs levels can inhibit SE induction, but GAs have a positive effect on the maturation and germination of embryos (Kim et al., 2019), as also confirmed in our study. ABA content was also the highest in CE, indicating that ABA could promote the maturation and differentiation of embryos. Vondrakova et al.(2018) found that ABA content was highest in the SE maturation stage of spruce. Cheng(2016) reported that the ABA content increased significantly after EC formation in cotton, and the genes related to ABA synthesis and signal transduction were also differentially expressed in cotton. The ABA2, NCED, PP2C, ABF, PYR/PYL and ABA receptor family were upregulated in the embryo.

We found that the IAA content was higher in EC than NEC, which was consistent with the results obtained for *Cyathea delgadii* (Grzyb et al., 2017). Su and Zhang(2009) reported that auxin gradients must be established for EC induction, and expressive of WUS gene was correlated with auxin gradients and initiated the polar localization of PIN1 in EC. These findings indicated that endogenous auxin had important significance for WUS gene and SE induction. The contents of tZ and cZ in NEC were higher than in EC, similarly to findings reported in *Prunus persica* (Pérez-Jiménez et al., 2013), this result was also supported by the research of Fraga et al.(2016), who found that the decrease in the Z level was related to the embryogenic potential. Compared with EC, BC with high contents of AUX and CKs was not conducive to EC formation; while the callus with low contents of IAA and DZ, and high tZ content could easily form SC. In addition, the contents of GAs and ABA in NEC, BC and SC were more than 6 times that of EC, which indicated that the high contents of GAs and ABA inhibited EC induction. Similar results had shown that the higher ABA content in NEC was also found in *Hevea brasiliensis* (Etienne et al., 1993) and *Cyathea delgadii* (Grzyb et al., 2017), while Pérez-Jiménez et al.(2013) found no significant difference in ABA content between NEC and EC in *Prunus persica*. In contrast, Nakagawa et al.(2001) found a higher ABA level in EC in *Cucumis melo*. These difference may depend on the sensitivity of different species to ABA and the difference in time and space. The content difference of these endogenous hormones in

different SE stages and different callus can provide a reference for SE induction and the application of exogenous hormone.

Endogenous hormone ratios

The ratios of endogenous hormones contents can be used as a physiological index to regulate the dedifferentiation and redifferentiation of the cell (Grzyb et al., 2018), and evaluate embryogenesis (Pérez-Jiménez et al., 2013). At different SE stages of *O. henryi*, the high ratios of IAA/ABA, IAA/GAs, AUX/GAs, and AUX/ABA and low ratios of ABA/CKs and GAs/CKs were favorable for EC induction; however, the ratios of IAA/CKs and AUX/CKs were lower in CE and promoted SE maturation and differentiation, which indicated that low AUX and high CKs contents were needed for SE maturation and differentiation, consistent with the actual theory.

The IAA/ABA ratio was significantly higher in EC than NEC, and similar results were found in *Corylus avellana* (Centeno et al., 1997); In contrast, the ABA/CKs ratio was significantly higher in NEC than EC, and similar results have been reported by Grzyb et al. (2017). Guo et al.(2018) reported that the ABA/GAs ratio can reflect the growth and dormancy of plant, as well as determine the developmental pattern of the embryo or late stages of the embryo (Zheng et al., 2013). Our study showed that the callus with low ratios of IAA/ABA, IAA/GAs, AUX/GAs, and AUX/ABA, and high ratios of ABA/CKs and GAs/CKs could not form EC, and an ABA/GAs ratio that was too high or too low was not conducive to SE induction.

Histochemical studies

Starch and protein are active metabolites in cell growth and morphogenesis. The starch granule staining gradually lightened with SE development of *O. henryi*, and the protein grain staining gradually darkened. Similar findings have been reported in *Picea sitchensis* SE (Liu, 2009). The changes at different SE stages can be used as a biochemical marker in *O. henryi* and provide important information for SE induction.

Compared with EC, the starch and protein granules staining were lighter in NEC.

Similar results have been reported in *Picea sitchensis* (Liu, 2009) and *Albizia lebbek* (Saeed and Shahzad, 2015); similarly, the starch and protein granules staining were lighter in BC and darker in SC. These phenomena were consistent with the results of our quantitative analysis, which indicated that contents of energy substances that were too low or too high were not conducive to callus development.

In summary, SE was a complex physiological process, and a single or two indexes was insufficient to evaluate the SE mechanism in *O. henryi*. The physiological indexes interacted with each other, and the results showed that energy substances were the material basis of SE, which affected the enzyme activities, regulated ROS metabolism, and thus regulated the morphogenesis and development of somatic embryos. In addition, the contents and ratios of endogenous hormones affected the dedifferentiation, dedifferentiation and embryogenesis of somatic cells. The formation of NEC was related to low contents of energy substances. H_2O_2 and low enzyme activities; the high contents of soluble sugar, H_2O_2 , AUX, CKs, and high PPO activity and low content of soluble protein were the basic causes of BC formation; the high-energy substance contents and low activities of SOD and POD could easily form SC. The high contents of Gas and ABA, high ratios of ABA/CKs and GAs/CKs, and low ratios of IAA/ABA, IAA/GAs, AUX/GAs, and AUX/ABA were explanations for the inability of the callus to form EC. Based on the above conclusion, the EC induction rate and frequency of EC transformation into a somatic embryo in *O. henryi* might be regulated and improved by adjusting the external conditions, such as the medium composition and culture environment, especially the application of exogenous hormone types and concentrations.

MATERIALS AND METHODS

Plant materials

Mature seeds were collected from a 120–150-year-old *O. henryi* tree growing in Mengguan town, Guiyang city (latitude: 26°14'23"N, longitude: 106°25'12"W) on Nov 2017. The mature seeds were treated with concentrated H_2SO_4 for 1 h, washed

with tap water for 30 min, and disinfected in 75% alcohol for 1 min and then in 2% NaClO for 8 min, followed by five rinses in sterile distilled water. Because the seed coat is hard, the mature seeds were soaked in sterile water for 24 h to make them swell and soften, which was conducive to stripping the mature embryos. Mature embryos were used as the explant for SE induction (Wu et al., 2020).

The mature embryos were inoculated into B5 medium (Gamborg et al., 1968) supplemented with 30 g/L sucrose, 500 mg/L glutamine, 2.5 g/L phytigel, 0.2 mg/L 6-benzylaminopurine (BA) and 2.0 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D) for callus induction. The pH values of the medium were adjusted to 5.90 ± 0.1 before autoclaving 121°C for 20 min. After 6 weeks, callus were transferred into B5 medium supplemented with 0.5 mg/L kinetin (KT), 1.0 mg/L 2,4-D for callus proliferation induction; after 4 weeks, NEC (Fig. 1-a), BC (Fig. 1-b), SC (Fig. 1-c) and EC (Fig. 1-d) were collected. GE (Fig. 1-e) was obtained after EC continued to be cultured in the above medium for 2 weeks in the dark at $25 \pm 2^{\circ}\text{C}$. CE (Fig. 1-f) was obtained after GE was cultured in B5 differentiation medium supplemented with 0.5 mg/L thidiazuron (TDZ) and 0.2 mg/L naphthaleneacetic acid (NAA) for 60 days.

The different tissues were collected, the culture medium of tissue surface was cleaned with deionized water. The tissues were placed in 5 ml centrifuge tubes, quickly frozen in liquid nitrogen, and then transferred to -80°C freezer for storage.

Biochemical analyses

The contents of energy substances, ROS, endogenous hormones and enzyme activities in different tissues, such as NEC, BC, SC, EC, GE and CE, were determined. Fresh tissues of 0.2-0.5g were obtained to detect and analyze biochemical events, with 3-5 replications performed per treatment.

Energy substances

The contents of soluble sugar and starch were determined using the sulfuric

acid-anthrone method (Liu et al., 2016), and the soluble protein content was determined using Coomassie brilliant blue (Bradford, 1976).

H₂O₂ contents and enzyme activities

H₂O₂ was determined according to the method of Ferguson et al. (1983) method. SOD activity was determined using the nitroblue tetrazolium method (Dhindsa et al., 1981). PPO activity was determined by pyrocatechol colorimetry (Liu et al., 2016). POD activity was measured using the guaiacol colorimetric method (Türkan et al., 2005). The determination of APX activity was based on the method of Yoshiyuki and Kozi (1981), and CAT activity was estimated according to Aebi(1984).

Endogenous hormones

The contents of endogenous hormones, such as AUX (indole-3-acetic acid [IAA], methyl indole-3-acetate [ME-IAA], 3-indolebutyric acid [IBA], indole-3-carboxylic acid [ICA], indole-3-carboxaldehyde [ICAld]), CKs (N6-isopentenyladenine [IP], trans-zeatin [tZ], cis-zeatin [cZ], dihydrozeatin [DZ]), GAs (gibberellin A1 [GA1], GA3, GA4, GA7, GA9, GA15, GA19, GA20, GA24, GA53) and ABA, were measured using the Metware (<http://www.metware.cn/>) method in different tissues. The tissues were removed from -80 °C freezer and ground into a powder with a grinder (MM 400, Retsch) (30 Hz, 1 min); then, 50 mg of the powdered tissues was added as an internal standard of the appropriate amount and extracted with methanol: water: formic acid = 15:4:1 (v: v: v). The extracts were concentrated and then redissolved in 100 µl 80% methanol aqueous solution and filtered with 0.22 µm PTFE filter membranes. They were then placed in sample bottles and analyzed using liquid chromatography-tandem mass spectrometry (LC-MS/MS). The peak area of each peak represented the relative content of the corresponding hormone. Finally, the qualitative and quantitative analysis results were obtained for all sample hormones.

463

464 **Histochemical studies**

465 The different tissues were fixed in FAA (formalin: acetic acid: 50% alcohol, 1:1:18)
 466 for 24 h. The tissue was processed with an alcohol-xylol series and embedded in
 467 paraffin. Sections (8 μ m) were cut on a rotary microtome (Leica, Germany) and
 468 dewaxed with xylol. Starch granules were stained with Schiff reagent (Liu, 2009), and
 469 protein granules with naphthol yellow S reagent (Liu, 2009). Finally, the sections
 470 were mounted with neutral balsam and examined under a DM3000 light microscope
 471 (Leica, Germany).

472

473 **Statistical analysis**

474 The results were expressed as the mean $\pm$ standard error for 3-4 replications per
 475 treatment. They were analyzed by one-way ANOVA followed by Tukey's test ($p <$
 476 0.05) with SPSS (version 18).

477

478 **Supplemental Data**

479 **Supplemental Energy substance**

480 **Supplemental H₂O₂ contents and enzyme activities**

481 **Supplemental hormones levels**

482

483

484 **Acknowledgments**

485 We are very grateful to the members of the research group for their help and

interesting scientific discussions. This work has benefited from the facilities and expertise of Metware (<http://www.metware.cn/>) and the experimental condition of the Institute for Forest Resources & Environment of Guizhou.

Tables

Table 1. Ratios of phytohormones during somatic embryogenesis in *O. henryi*.

Data are the means ($\pm$ SD) of three replicates. The use of the same lowercase letters indicates that the values were not significantly different according to Tukey's test ($p < 0.05$). CKs represent the sum of cZ, tZ, DZ and IP contents. GAs represent the sum of GA4, GA15 and GA19 contents. AUX represent the sum of IAA, ICAld and ME-IAA contents.

Phytohormone ratios	NEC	EC	BC	SC	GE	CE
IAA/ABA	0.16 $\pm$ 0.02b	3.16 $\pm$ 0.58a	0.57 $\pm$ 0.1b	0.3 $\pm$ 0.05b	0.74 $\pm$ 0.2b	0.86 $\pm$ 0.1b
IAA/CKs	1.24 $\pm$ 0.09a	2.38 $\pm$ 0.04a	1.74 $\pm$ 0.1a	1.3 $\pm$ 0.3a	4 $\pm$ 2a	0.88 $\pm$ 0.2a
IAA/GAs	0.36 $\pm$ 0.04c	2.94 $\pm$ 0.17a	0.36 $\pm$ 0.02c	0.33 $\pm$ 0.03c	0.6 $\pm$ 0.08bc	1.01 $\pm$ 0.13b
AUX/CKs	8.16 $\pm$ 0.44a	8.22 $\pm$ 0.26a	9.89 $\pm$ 1.08a	7.89 $\pm$ 1.51a	10.9 $\pm$ 5.26a	2.72 $\pm$ 0.43a
AUX/GAs	2.38 $\pm$ 0.62b	10.15 $\pm$ 0.88a	2 $\pm$ 0.06b	1.93 $\pm$ 0.21b	1.71 $\pm$ 0.33b	3.19 $\pm$ 0.76b
AUX/ABA	1.07 $\pm$ 0.14b	10.93 $\pm$ 3.52a	3.16 $\pm$ 0.61b	1.53 $\pm$ 0.21b	2.02 $\pm$ 0.63b	2.7 $\pm$ 0.35b
ABA/CKs	7.69 $\pm$ 0.68a	0.81 $\pm$ 0.16b	3.3 $\pm$ 0.6ab	5.2 $\pm$ 1.1ab	5.5 $\pm$ 2.7ab	1 $\pm$ 0.1b
ABA/GAs	2.29 $\pm$ 0.5a	0.99 $\pm$ 0.16b	0.65 $\pm$ 0.07b	1.3 $\pm$ 0.2ab	0.9 $\pm$ 0.1b	1.2 $\pm$ 0.2ab
GAs/CKs	3.55 $\pm$ 0.43a	0.81 $\pm$ 0.04a	4.93 $\pm$ 0.49a	4.02 $\pm$ 0.59a	6.95 $\pm$ 3.33a	0.88 $\pm$ 0.18a

Figure Legends

Fig .1 Somatic embryogenesis in *O. henryi*.

(a-d) The mature embryos were inoculated into B5 medium supplemented with 30 g/L sucrose, 500 mg/L glutamine, 2.5 g/L phytigel, 0.2 mg/L 6-benzylaminopurine (BA) and 2.0 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D) for callus induction. After 6 weeks, callus were transferred into B5 medium supplemented with 0.5 mg/L kinetin (KT), 1.0 mg/L 2,4-D for proliferation induction. After 4 weeks, non-embryogenic

callus (a), browning callus (b), snowflake callus (c), and embryogenic callus (d) were obtained.

(e) Globular embryo was obtained after embryogenic callus continued to be cultured in B5 medium supplemented with 0.5 mg/L kinetin (KT), 1.0 mg/L 2,4-D for 6 weeks in the dark at $25 \pm 2^\circ\text{C}$.

(f) Cotyledon embryo was obtained after globular embryo was cultured in B5 differentiation medium supplemented with 0.5 mg/L thidiazuron (TDZ) and 0.2 mg/L naphthaleneacetic acid (NAA) for 60 days.

Fig. 2 Contents of soluble sugar, starch and protein during somatic embryogenesis in *O. henryi*.

(I) Soluble sugar contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

(II) Starch contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

(III) Protein contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

Fig. 3 H_2O_2 content and enzymatic activity during somatic embryogenesis in *O. henryi*.

(I) H₂O₂ contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

(II) Polyphenoloxidase activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of four independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

(III) Superoxide dismutase activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

(IV) Ascorbate peroxidase activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of four independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

(V) POD activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

(VI) CAT activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of four independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

Fig. 4 Endogenous hormones contents during somatic embryogenesis in *O. henryi*.

(I) Auxin (IAA, ICAld and ME-IAA) contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC.

(II) Cytokinin (cZ, tZ, DZ and IP) contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC.

(III) Gibberellin acid (GA4, GA15 and GA19) contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC.

(IV) Absciscic acid (ABA) contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Data are the means $\pm$ standard error of three replicates. The use of the same lowercase letters indicates that the values were not significantly different according to Tukey's test ($p < 0.05$).

Fig. 5 Starch grains staining during somatic embryogenesis in *O. henryi*.

Starch granules were mainly distributed in cell wall and intercellular space, followed by uneven distribution in cytoplasm, less in nucleus, and then the meristem staining was dark. (a) Non-embryogenic callus. (b) Embryogenic callus. (c) Embryogenic callus. (d) Browning callus. (e) Snowflake callus. (f) Globular embryo. (g) Heart-shaped embryo. (h) Prophase of CE. (i) Cotyledon embryo.

Fig. 6 Protein grains staining during somatic embryogenesis in *O. henryi*.

Protein granules staining was relatively light, which was the most obvious in nucleus,

followed by cytoplasm, and then there were more protein grains accumulated in the meristems. **(a)** Non-embryogenic callus. **(b)** Embryogenic callus. **(c)** Browning callus. **(d)** Snowflake callus. **(e)** Globular embryo. **(f)** Heart-shaped embryo. **(g)** Prophase of cotyledon embryo. **(h)** Cotyledon embryo.

LITERATURE CITED

- Aderkas PV, Lelu MA, Label P** (2001) Plant growth regulator levels during maturation of larch somatic embryos. *Plant Physiol Bioch* **39**: 495-502
- Aebi H** (1984) Methods in Enzymology. Oxygen Radicals in Biological Systems. Academic Press, Orlando, pp 121-126
- Baluska F, Samaj J, Wojtaszek P, Volkmann D, Menzel D** (2003) Cytoskeleton-plasma membrane-cell wall continuum in plants. Emerging links revisited. *Plant Physiol* **133**: 482-491
- Bartos PMC, Gomes HT, Velho Do Amaral LI, Teixeira JB, Scherwinski-Pereira JE** (2018) Biochemical events during somatic embryogenesis in *Coffea arabica* L. *3 Biotech* **8**
- Bradford MM** (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248-254
- Cangahuala-Inocente GC, Silveira V, Caprestano CA, Floh EIS, Guerra MP** (2014) Dynamics of physiological and biochemical changes during somatic embryogenesis of *Acacia sellowiana*. *In Vitro Cell Dev-Pl* **50**: 166-175
- Centeno ML, Rodriguez R, Berros B, Rodriguez A** (1997) Endogenous hormonal content and somatic embryogenic capacity of *Corylus avellana* L. cotyledons. *Plant Cell Rep* **17**: 139-144
- Cheng WH** (2016) Research on physiological and molecular mechanisms of somatic embryogenesis in cotton (*Gossypium hirsutum* L.). Dissertation, Shihezi University
- Cheng ZJ, Wang L, Sun W, Zhang Y, Zhou C, Su YH, Li W, Sun TT, Zhao XY, Li XG, Cheng Y, Zhao Y, Xie Q, Zhang XS** (2013) Pattern of Auxin and Cytokinin Responses for Shoot Meristem Induction Results from the Regulation of Cytokinin Biosynthesis by AUXIN RESPONSE FACTOR3. *Plant Physiol* **161**: 240-251
- Choi YE, Katsumi M, Sano H** (2001) Triiodobenzoic acid, an auxin polar transport inhibitor, suppresses somatic embryo formation and postembryonic shoot/root development in *Eleutherococcus senticosus*. *Plant Ence* **160**: 1183-1190
- Correia S, Lopes ML, Canhoto JM** (2011) Somatic embryogenesis induction system for cloning an adult *Cyphomandra betacea* (Cav.) Sendt. (tamarillo). *Trees-Struct Funct* **25**: 1009-1020
- Cui K, Xing G, Liu X, Xing G, Wang Y** (1999) Effect of hydrogen peroxide on somatic embryogenesis of *Lycium barbarum* L. *Plant Sci* **146**: 16
- Dhindsa RS, Plumbdhindsa PL, Thorpe TA** (1981) Leaf senescence: correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J Exp Bot* **1**: 93-101
- Fraga HP, Vieira LD, Puttkammer CC, Dos Santos HP, Garighan JA, Guerra MP** (2016) Glutathione and abscisic acid supplementation influences somatic embryo maturation and hormone endogenous levels during somatic embryogenesis in *Podocarpus lambertii* Klotzsch ex Endl. *Plant*

632 Sci **253**: 98-106

633 **Etienne, H., Sotta, B., Montoro, P., Miginiac, E.** (1993) Relations between exogenous growth
634 regulators and endogenous indole-3-acetic acid and abscisic acid in the expression of somatic
635 embryogenesis in *Hevea brasiliensis* (Muell. Arg.). Plant Sci **88**: 91–96

636 **Steward FC** (1958) Growth and organized development of cultured cells. III. Interpretations of the
637 growth from free cell to carrot plan. Am J Bot **45**:709-713

638 **Fatima S, Mujib A, Samaj J** (2011) Anti-oxidant enzyme responses during in vitro embryogenesis in
639 *Catharanthus roseus*. J Hort Sci Biotech **86**: 569-574

640 **Ferguson IB, Watkins CB, Harman JE** (1983) Inhibition by calcium of senescence of detached
641 cucumber cotyledons: effect on ethylene and hydroperoxide production. Plant Physiol **71**: 182-186

642 **Gamborg OL, Miller RA, Ojima K** (1968) Nutrient requirements of suspension cultures of soybean
643 root cells. Exp Cell Res **50**: 151-158

644 **Gomes HT, Cunha Bartos PM, Silva CO, Velho Do Amaral LI, Scherwinski-Pereira JE** (2014)
645 Comparative biochemical profiling during the stages of acquisition and development of somatic
646 embryogenesis in African oil palm (*Elaeis guineensis* Jacq.). Plant Growth Regul **74**: 199-208

647 **Grzyb M, Kalandyk A, Mikula A** (2018) Effect of TIBA, fluridone and salicylic acid on somatic
648 embryogenesis and endogenous hormone and sugar contents in the tree fern *Cyathea delgadii* Sternb.
649 Acta Physiol Plant **40**

650 **Grzyb M, Kalandyk A, Waligorski P, Mikula A** (2017) The content of endogenous hormones and
651 sugars in the process of early somatic embryogenesis in the tree fern *Cyathea delgadii* Sternb. Plant
652 Cell Tiss Org Cult **129**: 387-397

653 **Guo B, He W, Zhao Y, Wu Y, Fu Y, Guo J, Wei Y** (2017) Changes in endogenous hormones and
654 H₂O₂ burst during shoot organogenesis in TDZ-treated *Saussurea involucre* explants. Plant Cell
655 Tiss Org Cult **128**: 1-8

656 Guo YQ, Huang DB, Chang XJ, Zhu C, Li XZ, Lai ZX (2018) Somatic embryogenesis and the changes
657 of endogenous hormones in *Camellia sinensis* Tieguanyin. Chinese Journal of Applied and
658 Environmental Biology **24**: 824-832

659 **Hazubska-Przybyl T, Kalembe EM, Ratajczak E, Bojarczuk K** (2016) Effects of abscisic acid and
660 an osmoticum on the maturation, starch accumulation and germination of *Picea* spp. somatic
661 embryos. Acta Physiol Plant **38**

662 **Iraqi D, Tremblay FM** (2001) The role of sucrose during maturation of black spruce (*Picea mariana*)
663 and white spruce (*Picea glauca*) somatic embryos. Physiol Plantarum **111**: 381-388

664 **Jariteh M, Ebrahimzadeh H, Niknam V, Mirmasoumi M, Vahdati K** (2015) Developmental
665 changes of protein, proline and some antioxidant enzymes activities in somatic and zygotic embryos
666 of Persian walnut (*Juglans regia* L.). Plant Cell Tiss Org Cult **122**: 101-115

667 **Jiménez MV**(2001) Regulation of in vitro somatic embryogenesis with emphasis on to the role of
668 endogenous hormones. Brazilian Journal of Plant Physiology **13**: 196-223

669 **Jiménez VM** (2005) Involvement of plant hormones and plant growth regulators on in vitro somatic
670 embryogenesis. Plant Growth Regul **47**: 91-110

671 **Kim DH, Kang KW, Sivanesan I** (2019) Influence of auxins on somatic embryogenesis in *Haworthia*
672 *retusa* Duval. Biologia **74**: 25-33

673 **Lee C, Clark SE** (2015) A WUSCHEL-Independent Stem Cell Specification Pathway Is Repressed by
674 PHB, PHV and CNA in Arabidopsis. Plos One **10**: e0126006

675 **Leljak-Levanic D, Bauer N, Mihaljevic S, Jelaska S** (2004) Somatic embryogenesis in pumpkin

(*Cucurbita pepo* L.): control of somatic embryo development by nitrogen compounds. J Plant Physiol **161**: 229-236

Libik M, Konieczny R, Pater B, Slesak I, Miszalski Z (2005) Differences in the activities of some antioxidant enzymes and in H₂O₂ content during rhizogenesis and somatic embryogenesis in callus cultures of the ice plant. Plant Cell Rep **23**: 834-841

Liu CQ (2009) The study of somatic embryogenesis induction and Histocytology in *Hippophae rhamnoides* subsp. sinensis Rousi and *Sequoia sempervirens* (Lamb.) Endl. China Environmental Science Press, Beijing, pp 93-136.

Liu P, Li MJ, Ding YF (2016) Plant physiology experiment. Science Press, Beijing, pp 50-78

Liu H, Zhang H, Dong YX, Hao YJ, Zhang XS (2018) DNA METHYLTRANSFERASE1-mediated shoot regeneration is regulated by cytokinin-induced cell cycle in Arabidopsis. New Phytol **217**: 219-232

Liu YQ (2016) Preliminary study on the mechanism of somatic embryogenesis of ponka (*Citrus reticulata* blanco). Dissertation, Yangzhou university

Manivannan A, Jana S, Soundararajan P, Ko CH, Jeong BR (2015) Antioxidant enzymes metabolism and cellular differentiation during the developmental stages of somatic embryogenesis in *Torilis japonica* (Houtt.) DC. Plant Omics **8**: 461-471

Martin, Cuadrado, Guerra, Gallego, Hita, Martin, Dorado, Villalobos (2000) Differences in the contents of total sugars, reducing sugars, starch and sucrose in embryogenic and non-embryogenic calli from *Medicago arborea* L. Plant Sci **154**: 143-151

Morel A, Teyssier C, Trontin JFO, Eliá Ová KI, Pe Ek BI, Beaufour M, Morabito D, Boizot N, Metté CL, Belal-Bessai L (2014) Early molecular events involved in *Pinus pinaster* Ait. somatic embryo development under reduced water availability: transcriptomic and proteomic analyses. Physiol Plantarum **152**: 184-201

Nakagawa H, Saijyo T, Yamauchi N, Shigyo M, Kako S, Ito A (2001) Effects of sugars and abscisic acid on somatic embryogenesis from melon (*Cucumis melo* L.) expanded cotyledon. Sci Hortic-Amsterdam **90**: 85-92

Nic-Can GI, Loyola-Vargas VM (2016) The role of the auxins during somatic embryogenesis. In: Loyola-Vargas VM, Ochoa-Alejo N (eds) Somatic embryogenesis: fundamental aspects and applications. Springer International Publishing, Switzerland, pp 171-182

Pérez-Jiménez M, Cantero-Navarro E, Acosta M, Cos-Terrer J (2013) Relationships between endogenous hormonal content and direct somatic embryogenesis in *Prunus persica* L. Batsch cotyledons. Plant Growth Regul **71**: 219-224

Pinto G, Silva S, Neves L, Araujo C, Santos C (2010) Histocytological changes and reserve accumulation during somatic embryogenesis in *Eucalyptus globulus*. Trees-Struct Funct **24**: 763-769

Roowi SH, Ho C, Alwee SSRS, Abdullah MO, Napis S (2010) Isolation and characterization of differentially expressed transcripts from the suspension cells of oil palm (*Elaeis guineensis* Jacq.) in response to different concentration of auxins. Mol Biotechnol **46**: 1-19

Saeed T, Shahzad A (2015) High frequency plant regeneration in Indian Siris via cyclic somatic embryogenesis with biochemical, histological and SEM investigations. Ind Crop Prod **76**: 623-637

Su YH, Zhang XS (2009) Auxin gradients trigger de novo formation of stem cells during somatic embryogenesis. Plant Signal Behav **4**: 574-576

Türkan S, Bor M, Zdemir F, Koca H (2005) Differential responses of lipid peroxidation and antioxidants in the leaves of drought-tolerant *P. acutifolius* Gray and drought-sensitive *P. vulgaris* L.

720 subjected to polyethylene glycol mediated water stress. *Plant Sci* **168**: 231

721 **Vondrakova Z, Dobrev PI, Pesek B, Fischerova L, Vagner M, Motyka V** (2018) Profiles of
 722 endogenous phytohormones over the course of norway spruce somatic embryogenesis. *Front Plant*
 723 *Sci* **9**: 1283

724 **Wang X, Shi L, Lin G, Pan X, Chen H, Wu X, Takac T, Samaj J, Xu C** (2013) A systematic
 725 comparison of embryogenic and non-embryogenic cells of banana (*Musa* spp. AAA): Ultrastructural,
 726 biochemical and cell wall component analyses. *Sci Hortic-Amsterdam* **159**: 178-185

727 **Warchol M, Skrzypek E, Kusibab T, Dubert F** (2015) Induction of somatic embryogenesis and
 728 biochemical characterization of *Cordyline australis* (G. Forst.) Endl. 'Red Star' callus. *Sci*
 729 *Hortic-Amsterdam* **192**: 338-345

730 **Wei XL, Meng XS, Deng Z** (2014) Relation between being endangered and seed reproductive ecology
 731 of a rare species *Ormosia henryi*. *Seed*, **33**: 82-86

732 **Wichers HJ, Malingre TM, Huizing HJ** (1985) Induction of phenoloxidase in cell suspension
 733 cultures of *Mucuna pruriens* L. : Effects on accumulation of L-3,4-dihydroxyphenylalanine and
 734 biotransformation capacity. *Planta* **165**: 264-268

735 **Wu GY, Wei XL, Wang X, Wei Y** (2020) Induction of somatic embryogenesis in different explants
 736 from *Ormosia henryi* Prain. *Plant Cell Tiss Org Cult* **142**: 229-240

737 **Yoshiyuki N, Kozi A** (1981) Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in
 738 spinach chloroplasts. *Plant Cell Physiol* **22**: 867-880

739 **Zhang S, Han S, Yang W, Wei H, Zhang M, Qi L** (2010) Changes in H₂O₂ content and antioxidant
 740 enzyme gene expression during the somatic embryogenesis of *Larix leptolepis*. *Plant Cell Tiss Org*
 741 *Cult* **100**: 21-29

742 **Zhang T, Lian H, Zhou C, Xu L, Jiao Y, Wang J** (2017) A Two-Step Model for de Novo Activation
 743 of WUSCHEL during Plant Shoot Regeneration. *Plant Cell* **29**: 1073-1087

744 **Zheng Q, Zheng Y, Perry SE** (2013) Decreased GmAGL15 expression and reduced ethylene
 745 synthesis may contribute to reduced somatic embryogenesis in a poorly embryogenic cultivar of
 746 *Glycine max*. *Plant Signal Behav* **8**: e25422

Fig .1 Somatic embryogenesis in *O. henryi*.

(a-d) The mature embryos were inoculated into B5 medium supplemented with 30 g/L sucrose, 500 mg/L glutamine, 2.5 g/L phytigel, 0.2 mg/L 6-benzylaminopurine (BA) and 2.0 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D) for callus induction. After 6 weeks, callus were transferred into B5 medium supplemented with 0.5 mg/L kinetin (KT), 1.0 mg/L 2,4-D for proliferation induction. After 4 weeks, non-embryogenic callus **(a)**, browning callus **(b)**, snowflake callus **(c)**, and embryogenic callus **(d)** were obtained. **(e)** Globular embryo was obtained after embryogenic callus continued to be cultured in B5 medium supplemented with 0.5 mg/L kinetin (KT), 1.0 mg/L 2,4-D for 6 weeks in the dark at $25 \pm 2^\circ\text{C}$. **(f)** Cotyledon embryo was obtained after globular embryo was cultured in B5 differentiation medium supplemented with 0.5 mg/L thidiazuron (TDZ) and 0.2 mg/L naphthaleneacetic acid (NAA) for 60 days.

Fig. 2 Contents of soluble sugar, starch and protein during somatic embryogenesis in *O. henryi*.

(I) Soluble sugar contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean±standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$). **(II)** Starch contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean±standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$). **(III)** Protein contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean±standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

Fig. 3 H₂O₂ content and enzymatic activity during somatic embryogenesis in *O. henryi*.

(I) H₂O₂ contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean ± standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$). **(II)** Polyphenoloxidase activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean ± standard error of four independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$). **(III)** Superoxide dismutase activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean ± standard error, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$). **(IV)** Ascorbate peroxidase activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean ± standard error of four independent experiments, bars denoted by the same

letter are not significantly different using the Tukey's test ($p < 0.05$). **(V)** POD activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of three independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$). **(VI)** CAT activity at different SE stages, such as EC, GE, and CE and in abnormal tissues, such as NEC, BC, and SC. Bars represent the mean $\pm$ standard error of four independent experiments, bars denoted by the same letter are not significantly different using the Tukey's test ($p < 0.05$).

Fig. 4 Endogenous hormones contents during somatic embryogenesis in *O. henryi*.

(I) Auxin (IAA, ICAld and ME-IAA) contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. **(II)** Cytokinin (cZ, tZ, DZ and IP) contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. **(III)** Gibberellin acid (GA4, GA15 and GA19) contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. **(IV)** Absciscic acid (ABA) contents at different SE stages, such as EC, GE, and CE, and in abnormal tissues, such as NEC, BC, and SC. Data are the means $\pm$ standard error of three replicates. The use of the same lowercase letters indicates that the values were not significantly different according to Tukey's test ($p < 0.05$).

Fig. 5 Starch grains staining during somatic embryogenesis in *O. henryi*.

Starch granules were mainly distributed in cell wall and intercellular space, followed by uneven distribution in cytoplasm, less in nucleus, and then the meristem staining was dark. **(a)** Non-embryogenic callus. **(b)** Embryogenic callus. **(c)** Embryogenic callus. **(d)** Browning callus. **(e)** Snowflake callus. **(f)** Globular embryo. **(g)** Heart-shaped embryo. **(h)** Prophase of CE. **(i)** Cotyledon embryo.

Fig. 6 Protein grains staining during somatic embryogenesis in *O. henryi*.

Protein granules staining was relatively light, which was the most obvious in nucleus, followed by cytoplasm, and then there were more protein grains accumulated in the meristems. **(a)** Non-embryogenic callus. **(b)** Embryogenic callus. **(c)** Browning callus. **(d)** Snowflake callus. **(e)** Globular embryo. **(f)** Heart-shaped embryo. **(g)** Prophase of cotyledon embryo. **(h)** Cotyledon embryo.

Parsed Citations

Aderkas PV, Lelu MA, Label P (2001) Plant growth regulator levels during maturation of larch somatic embryos. *Plant Physiol Bioch* 39: 495-502

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Aebi H (1984) *Methods in Enzymology. Oxygen Radicals in Biological Systems*. Academic Press, Orlando, pp 121-126

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Baluska F, Samaj J, Wojtaszek P, Volkmann D, Menzel D (2003) Cytoskeleton-plasma membrane-cell wall continuum in plants. *Emerging links revisited*. *Plant Physiol* 133: 482-491

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bartos PMC, Gomes HT, Velho Do Amaral LI, Teixeira JB, Scherwinski-Pereira JE (2018) Biochemical events during somatic embryogenesis in *Coffea arabica* L. *3 Biotech* 8

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248-254

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cangahuala-Inocente GC, Silveira V, Caprestano CA, Floh EIS, Guerra MP (2014) Dynamics of physiological and biochemical changes during somatic embryogenesis of *Acca sellowiana*. *In Vitro Cell Dev-PI* 50: 166-175

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Centeno ML, Rodriguez R, Berros B, Rodriguez A (1997) Endogenous hormonal content and somatic embryogenic capacity of *Corylus avellana* L. cotyledons. *Plant Cell Rep* 17: 139-144

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cheng WH (2016) *Research on physiological and molecular mechanisms of somatic embryogenesis in cotton (Gossypium hirsutum L.)*. Dissertation, Shihezi University

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cheng ZJ, Wang L, Sun W, Zhang Y, Zhou C, Su YH, Li W, Sun TT, Zhao XY, Li XG, Cheng Y, Zhao Y, Xie Q, Zhang XS (2013) Pattern of Auxin and Cytokinin Responses for Shoot Meristem Induction Results from the Regulation of Cytokinin Biosynthesis by AUXIN RESPONSE FACTOR3. *Plant Physiol* 161: 240-251

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Choi YE, Katsumi M, Sano H (2001) Triiodobenzoic acid, an auxin polar transport inhibitor, suppresses somatic embryo formation and postembryonic shoot/root development in *Eleutherococcus senticosus*. *Plant Ence* 160: 1183-1190

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Correia S, Lopes ML, Canhoto JM (2011) Somatic embryogenesis induction system for cloning an adult *Cyphomandra betacea* (Cav.) Sendt. (tamarillo). *Trees-Struct Funct* 25: 1009-1020

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cui K, Xing G, Liu X, Xing G, Wang Y (1999) Effect of hydrogen peroxide on somatic embryogenesis of *Lycium barbarum* L. *Plant Sci* 146: 16

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dhindsa RS, Plumbdhindsa PL, Thorpe TA (1981) Leaf senescence: correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J Exp Bot* 1: 93-101

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fraga HP, Vieira LD, Puttkammer CC, Dos Santos HP, Garighan JA, Guerra MP (2016) Glutathione and abscisic acid supplementation influences somatic embryo maturation and hormone endogenous levels during somatic embryogenesis in *Podocarpus lambertii* Klotzsch ex Endl. *Plant Sci* 253: 98-106

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Etienne, H., Sotta, B., Montoro, P., Miginiac, E. (1993) Relations between exogenous growth regulators and endogenous indole-3-acetic acid and abscisic acid in the expression of somatic embryogenesis in *Hevea brasiliensis* (Muell. Arg.). *Plant Sci* 88: 91-96

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Steward FC (1958) Growth and organized development of cultured cells. III. Interpretations of the growth from free cell to carrot plan. *Am J Bot* 45:709-713

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fatima S, Mujib A, Samaj J (2011) Anti-oxidant enzyme responses during in vitro embryogenesis in *Catharanthus roseus*. *J Hortic Sci Biotech* 86: 569-574

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ferguson IB, Watkins CB, Harman JE (1983) Inhibition by calcium of senescence of detached cucumber cotyledons: effect on ethylene and hydroperoxide production. *Plant Physiol* 71: 182-186

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gamborg OL, Miller RA, Ojima K (1968) Nutrient requirements of suspension cultures of soybean root cells. Exp Cell Res 50: 151-158

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gomes HT, Cunha Bartos PM, Silva CO, Velho Do Amaral LI, Scherwinski-Pereira JE (2014) Comparative biochemical profiling during the stages of acquisition and development of somatic embryogenesis in African oil palm (*Elaeis guineensis* Jacq.). Plant Growth Regul 74: 199-208

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Grzyb M, Kalandyk A, Mikula A (2018) Effect of TIBA, fluridone and salicylic acid on somatic embryogenesis and endogenous hormone and sugar contents in the tree fern *Cyathea delgadii* Sternb. Acta Physiol Plant 40

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Grzyb M, Kalandyk A, Waligorski P, Mikula A (2017) The content of endogenous hormones and sugars in the process of early somatic embryogenesis in the tree fern *Cyathea delgadii* Sternb. Plant Cell Tiss Org Cult 129: 387-397

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Guo B, He W, Zhao Y, Wu Y, Fu Y, Guo J, Wei Y (2017) Changes in endogenous hormones and H₂O₂ burst during shoot organogenesis in TDZ-treated *Saussurea involucre* explants. Plant Cell Tiss Org Cult 128: 1-8

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Guo YQ, Huang DB, Chang XJ, Zhu C, Li XZ, Lai ZX (2018) Somatic embryogenesis and the changes of endogenous hormones in *Camellia sinensis* Tieguanyin. Chinese Journal of Applied and Environmental Biology 24: 824-832

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hazubska-Przybyl T, Kalemba EM, Ratajczak E, Bojarczuk K (2016) Effects of abscisic acid and an osmoticum on the maturation, starch accumulation and germination of *Picea* spp. somatic embryos. Acta Physiol Plant 38

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Iraqi D, Tremblay FM (2001) The role of sucrose during maturation of black spruce (*Picea mariana*) and white spruce (*Picea glauca*) somatic embryos. Physiol Plantarum 111: 381-388

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jariteh M, Ebrahimzadeh H, Niknam V, Mirmasoumi M, Vahdati K (2015) Developmental changes of protein, proline and some antioxidant enzymes activities in somatic and zygotic embryos of Persian walnut (*Juglans regia* L.). Plant Cell Tiss Org Cult 122: 101-115

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jiménez MV (2001) Regulation of in vitro somatic embryogenesis with emphasis on to the role of endogenous hormones. Brazilian Journal of Plant Physiology 13: 196-223

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jiménez VM (2005) Involvement of plant hormones and plant growth regulators on in vitro somatic embryogenesis. Plant Growth Regul 47: 91-110

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kim DH, Kang KW, Sivanesan I (2019) Influence of auxins on somatic embryogenesis in *Haworthia retusa* Duval. Biologia 74: 25-33

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee C, Clark SE (2015) A WUSCHEL-Independent Stem Cell Specification Pathway Is Repressed by PHB, PHV and CNA in *Arabidopsis*. Plos One 10: e0126006

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Leljak-Levanic D, Bauer N, Mihaljevic S, Jelaska S (2004) Somatic embryogenesis in pumpkin (*Cucurbita pepo* L.): control of somatic embryo development by nitrogen compounds. J Plant Physiol 161: 229-236

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Libik M, Konieczny R, Pater B, Slesak I, Miszalski Z (2005) Differences in the activities of some antioxidant enzymes and in H₂O₂ content during rhizogenesis and somatic embryogenesis in callus cultures of the ice plant. Plant Cell Rep 23: 834-841

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu CQ (2009) The study of somatic embryogenesis induction and Histocytology in *Hippophae rhamnoides* subsp. *sinensis* Rousi and *Sequoia sempervirens* (Lamb.) Endl. China Environmental Science Press, Beijing, pp 93-136.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu P, Li MJ, Ding YF (2016) Plant physiology experiment. Science Press, Beijing, pp 50-78

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu H, Zhang H, Dong YX, Hao YJ, Zhang XS (2018) DNA METHYLTRANSFERASE1-mediated shoot regeneration is regulated by cytokinin-induced cell cycle in *Arabidopsis*. New Phytol 217: 219-232

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu YQ (2016) Preliminary study on the mechanism of somatic embryogenesis of ponka (*Citrus reticulata* blanco). Dissertation,

Manivannan A, Jana S, Soundararajan P, Ko CH, Jeong BR (2015) Antioxidant enzymes metabolism and cellular differentiation during the developmental stages of somatic embryogenesis in *Torilis japonica* (Houtt.) DC. Plant Omics 8: 461-471

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Martin, Cuadrado, Guerra, Gallego, Hita, Martin, Dorado, Villalobos (2000) Differences in the contents of total sugars, reducing sugars, starch and sucrose in embryogenic and non-embryogenic calli from *Medicago arborea* L. Plant Sci 154: 143-151

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Morel A, Teyssier C, Trontin JFO, Eliá Ová KI, Pe Ek BI, Beaufour M, Morabito D, Boizot N, Metté CL, Belal-Bessai L (2014) Early molecular events involved in *Pinus pinaster* Ait. somatic embryo development under reduced water availability: transcriptomic and proteomic analyses. Physiol Plantarum 152: 184-201

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nakagawa H, Saijyo T, Yamauchi N, Shigyo M, Kako S, Ito A (2001) Effects of sugars and abscisic acid on somatic embryogenesis from melon (*Cucumis melo* L.) expanded cotyledon. Sci Hortic-Amsterdam 90: 85-92

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nic-Can GI, Loyola-Vargas VM (2016) The role of the auxins during somatic embryogenesis. In: Loyola-Vargas VM, Ochoa-Alejo N (eds) Somatic embryogenesis: fundamental aspects and applications. Springer International Publishing, Switzerland, pp 171-182

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pérez-Jiménez M, Cantero-Navarro E, Acosta M, Cos-Terrer J (2013) Relationships between endogenous hormonal content and direct somatic embryogenesis in *Prunus persica* L. Batsch cotyledons. Plant Growth Regul 71: 219-224

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pinto G, Silva S, Neves L, Araujo C, Santos C (2010) Histocytological changes and reserve accumulation during somatic embryogenesis in *Eucalyptus globulus*. Trees-Struct Funct 24: 763-769

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Roowi SH, Ho C, Alwee SSRS, Abdullah MO, Napis S (2010) Isolation and characterization of differentially expressed transcripts from the suspension cells of oil palm (*Elaeis guineensis* Jacq.) in response to different concentration of auxins. Mol Biotechnol 46: 1-19

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Saeed T, Shahzad A (2015) High frequency plant regeneration in Indian Siris via cyclic somatic embryogenesis with biochemical, histological and SEM investigations. Ind Crop Prod 76: 623-637

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Su YH, Zhang XS (2009) Auxin gradients trigger de novo formation of stem cells during somatic embryogenesis. Plant Signal Behav 4: 574-576

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Türkan S, Bor M, Zdemir F, Koca H (2005) Differential responses of lipid peroxidation and antioxidants in the leaves of drought-tolerant *P. acutifolius* Gray and drought-sensitive *P. vulgaris* L. subjected to polyethylene glycol mediated water stress. Plant Sci 168: 231

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vondrakova Z, Dobrev PI, Pesek B, Fischerova L, Vagner M, Motyka V (2018) Profiles of endogenous phytohormones over the course of norway spruce somatic embryogenesis. Front Plant Sci 9: 1283

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang X, Shi L, Lin G, Pan X, Chen H, Wu X, Takac T, Samaj J, Xu C (2013) A systematic comparison of embryogenic and non-embryogenic cells of banana (*Musa* spp. AAA): Ultrastructural, biochemical and cell wall component analyses. Sci Hortic-Amsterdam 159: 178-185

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Warchol M, Skrzypek E, Kusibab T, Dubert F (2015) Induction of somatic embryogenesis and biochemical characterization of *Cordyline australis* (G. Forst.) Endl. 'Red Star' callus. Sci Hortic-Amsterdam 192: 338-345

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wei XL, Meng XS, Deng Z (2014) Relation between being endangered and seed reproductive ecology of a rare species *Ormosia henryi*. Seed, 33: 82-86

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wichers HJ, Malingre TM, Huizing HJ (1985) Induction of phenoloxidase in cell suspension cultures of *Mucuna pruriens* L. : Effects on accumulation of L-3,4-dihydroxyphenylalanine and biotransformation capacity. Planta 165: 264-268

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wu GY, Wei XL, Wang X, Wei Y (2020) Induction of somatic embryogenesis in different explants from *Ormosia henryi* Prain. Plant Cell Tiss Org Cult 142: 229-240

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yoshiyuki N, Kozi A (1981) Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. Plant Cell Physiol 22: 867-880

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang S, Han S, Yang W, Wei H, Zhang M, Qi L (2010) Changes in H₂O₂ content and antioxidant enzyme gene expression during the somatic embryogenesis of *Larix leptolepis*. Plant Cell Tiss Org Cult 100: 21-29

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang T, Lian H, Zhou C, Xu L, Jiao Y, Wang J (2017) A Two-Step Model for de Novo Activation of WUSCHEL during Plant Shoot Regeneration. Plant Cell 29: 1073-1087

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zheng Q, Zheng Y, Perry SE (2013) Decreased GmAGL15 expression and reduced ethylene synthesis may contribute to reduced somatic embryogenesis in a poorly embryogenic cultivar of *Glycine max*. Plant Signal Behav 8: e25422

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)