

NDRG1 expression is an independent prognostic factor in inflammatory breast cancer

Emilly S Villodre^{1,5}, Yun Gong^{2,5}, Lei Huo^{2,5}, Esther C Yoon², Naoto T Ueno^{1,5}, Wendy A Woodward³,
Debu Tripathy¹, Juhee Song⁴, Bisrat G Debeb^{1,5*}

Departments of ¹Breast Medical Oncology, ²Pathology, ³Radiation Oncology, ⁴Biostatistics and ⁵Morgan Welch Inflammatory Breast Cancer Clinic and Research Program, The University of Texas MD Anderson Cancer Center, Houston TX

*Corresponding author: Bisrat G. Debeb, DVM PhD, Department of Breast Medical Oncology, Section of Translational Breast Cancer Research, The Morgan Welch Inflammatory Breast Cancer Research Program and Clinic, The University of Texas MD Anderson Cancer Center

6565 MD Anderson Blvd, Houston, TX 77030 (Z12.5044). Office phone: 713-792-0696; Lab phone: 713-795-5041; Email: bgdebeb@mdanderson.org

ABSTRACT

NDRG1 is widely described as a metastasis suppressor in breast cancer. However, we found that NDRG1 is critical in promoting tumorigenesis and brain metastasis in mouse models of inflammatory breast cancer (IBC), a rare but highly aggressive form of breast cancer. We hypothesized that NDRG1 is a prognostic marker associated with poor outcome in patients with IBC. Microarray gene expression data from the IBC Consortium dataset were analyzed to compare *NDRG1* expression between IBC and non-IBC tumors and among breast cancer subtypes. NDRG1 levels in tissue microarrays from 64 IBC patients were evaluated by immunohistochemical staining with anti-NDRG1 primary antibody (32 NDRG1-low [\leq median], 32 NDRG1-high [$>$ median]). Overall and disease-free survival (OS and DSS) were analyzed with Kaplan–Meier curves and log-rank test. *NDRG1* mRNA expression was higher in IBC than in non-IBC tumors ($p=0.007$), and in more aggressive HER2+ and basal-like vs luminal IBC subtypes ($p<0.0001$). Univariate analysis showed NDRG1 expression, tumor grade, disease stage, estrogen receptor (ER) status, and receipt of adjuvant radiation to be associated with OS and DSS. NDRG1-high patients had poorer 10-year OS and DSS than NDRG1-low patients (OS, 19% vs 45%, $p=0.0278$; DSS, 22% vs 52%, $p=0.0139$). On multivariable analysis, NDRG1 independently predicted OS (hazard ratio [HR]=2.034, $p=0.0274$) and DSS (HR=2.287, $p=0.0174$). NDRG1-high ER-negative tumors had worse outcomes OS, $p=0.0003$; DSS, $p=0.0003$; and NDRG1-high tumors that received adjuvant radiation treatment had poor outcomes (OS, $p=0.0088$; DSS, $p=0.0093$). NDRG1 correlated positively with aggressive tumor characteristics in IBC and was a significant independent prognostic factor for DSS and OSS in IBC patients. Targeting NDRG1 may represent a novel strategy for improving clinical outcomes for patients with IBC.

Inflammatory breast cancer (IBC) is one of the most aggressive forms of breast cancer. Although rare, accounting for only 1%-4% of newly diagnosed breast cancer cases, it is responsible for a disproportionately high 10% of breast cancer-related deaths in the United States^{1,2}. IBC has a unique biology characterized by rapid proliferation and metastasis; indeed, almost all patients have lymph node involvement and more than 33% of patients with IBC present with distant metastasis at the time of diagnosis^{3,4}. Even with multimodality treatment approaches that include systemic chemotherapy, surgery, and radiation therapy, prognosis for patients with IBC is worse than for non-IBC patients (overall survival [OS] rates 40% versus 63% at 5 years)⁵⁻⁷. This may be due in part to 70% of IBC patients presenting with aggressive subtypes of HER2+ or triple-negative breast cancer (TNBC), compared with 40% of non-IBC tumors⁸. Efforts have been undertaken to identify molecular markers and therapeutic targets distinct to IBC and have identified important targets and pathways including EGFR, E-cadherin, eIFG4I, RhoC, and TIG1/AXL⁹⁻¹³. However, no IBC-specific molecular signature or target has been identified thus far, and effective targeted therapies for this disease remain limited.

N-myc downstream regulated gene 1 (NDRG1) is a stress response protein involved in hypoxia, cell growth, lipid metabolism, and resistance to chemotherapy¹⁴⁻¹⁹. NDRG1 is widely known as a metastasis suppressor in breast cancer, acting mainly by suppressing migration and invasion of breast cancer cells²⁰⁻²². However, we and others have shown NDRG1 to be a tumor promoter in aggressive breast cancer²³⁻²⁵. Nagai and colleagues also showed that high expression of NDRG1 was associated with aggressive breast cancer behaviors, including advanced stage at presentation and high-grade tumors and that NDRG1 was independently associated with poor survival outcome²⁶. However, the expression of NDRG1 and its clinical importance in IBC remains unknown.

Herein, we examined the expression of NDRG1 by using immunohistochemical staining of a tissue microarray (TMA) composed of samples from IBC patients and evaluated the expression of NDRG1 and its correlation with survival outcomes. We also assessed the association between NDRG1 expression and outcome stratified by known prognostic factors. Our findings showed that high expression of NDRG1 in IBC tumors was an independent predictor of worse OS and disease-specific survival (DSS).

RESULTS

The IBC Consortium dataset consists of microarray gene expression profiles of IBC and non-IBC tumors from three institutions²⁷. With this dataset, we analyzed *NDRG1* expression in patients with IBC and non-IBC and observed that *NDRG1* was expressed at higher levels in patients with IBC relative to non-IBC ($p=0.0287$; Supplementary Figure 1a). Considering the breast cancer subtypes within IBC tumors, *NDRG1* expression was significantly increased in more aggressive basal-like and HER2+ subtypes than in luminal subtypes of IBC ($p<0.0001$; Supplementary Figure 1b). We further found that *NDRG1* expression was significantly higher in patients with TNBC versus non-TNBC tumors in both IBC and non-IBC patients (Supplementary Figure 1c), supporting that *NDRG1* is positively associated with aggressive tumor characteristics. To determine whether *NDRG1* protein expression is associated with outcome in IBC, immunohistochemical staining was performed on TMAs from 64 patients with primary IBC who were treated between 1991 and 2004 at The University of Texas MD Anderson Cancer Center. The average age of these patients was 50 years (range 23-75 years). Eighty three percent of patients were stage III, 80% high grade and 62% were ER-negative tumors, and 67% of these patients received adjuvant radiation. The median follow-up time for the patients studied was 11.7 years and the median OS time was 3.7 years. *NDRG1* staining was predominantly cytoplasmic/membranous. Representative images of *NDRG1*-low and -high tumors are shown in Figure 1.

Table 1 summarizes patient characteristics based on *NDRG1* expression status, *NDRG1* expression was associated with negative HER2 status ($p=0.0077$). Univariate analysis (Table 2) showed that *NDRG1* expression (hazard ratio [HR]=2.1, $p=0.0150$), tumor grade (HR=2.4, $p=0.0463$), disease stage (HR=4.6, $p=0.0011$), hormone therapy (HR=0.3, $p=0.0026$), ER status (HR=0.4, $p=0.0098$), and adjuvant radiation therapy (HR=0.538, $p=0.0434$) were associated with OS. The same variables were also associated with DSS (Table 2).

The Kaplan-Meier method was used to evaluate the association of *NDRG1* expression and survival over time. Patients with *NDRG1*-low tumors experienced better actuarial 10-year OS

($p=0.0129$, Figure 2a) and DSS ($p=0.0134$, Figure 2b). Patients with NDRG1-high tumors showed significantly lower 10-year OS and DSS rates than patients with NDRG1-low (OS, 19% vs. 45%, $p=0.0278$; DSS, 22% vs 52%, $p=0.0139$). The median OS and DSS times were shorter for NDRG1-high patients (OS 2.5 years; DSS, 3.1 years) than for NDRG1-low patients (OS, 5.9 years; DSS, 10.7 years). Multivariable model predictors of OS and DSS included NDRG1 expression, ER status, disease stage, and receipt of adjuvant radiation (Table 3). Tumor grade was identified as being associated with OS and DSS at the univariate level but not at the multivariable level. NDRG1-high expression was a strong independent predictor of OS (HR=2.034, 95% confidence interval [CI]=1.081–3.822, $p=0.0274$) and DSS (HR=2.287, 95% CI=1.157–4.522, $p=0.0174$).

ER status was also an important prognostic factor for OS and DSS for patients with IBC; patients with ER-negative tumors had worse OS ($p=0.0077$) and DSS ($p=0.01$) relative to patients with ER-positive IBC tumors (Figure 3a-b). Multivariable analysis showed ER status to be an independent factor associated with OS (HR=0.307, 95% CI=0.145–0.650, $p=0.002$) and DSS (HR=0.307, 95% CI=0.147–0.642, $p=0.0017$) (Table 3). Interestingly, NDRG1-high and ER-negative tumors were associated with the worst clinical outcomes for patients with IBC (OS, $p=0.0003$; DSS, $p=0.0003$; Figure 3c-d). Survival outcomes of ER-positive patients were not affected by NDRG1 expression (Figure 3c-d). Analysis of median OS and DSS times highlights the importance of stratifying patients for both variables: patients with ER-negative tumors had a median 2.2 years for both OS and DSS, whereas those with ER-negative / NDRG1-high tumors had a median 1.6 years, and ER-negative / NDRG1-low tumors had medians of 3.2 years OS and 4.6 years DSS (Figure 3e-f).

Disease stage was another prognostic variable for OS (HR=3.212, 95% CI=1.171–8.814, $p=0.0234$) and DSS (HR=3.844, 95% CI=1.362–10.854, $p=0.011$). Kaplan-Meier analysis showed that patients with stage III IBC had better outcomes than did patients with stage IV tumors (OS, $p=0.0003$; DSS, $p<0.0001$) (Figure 4a-b). Further stratification of patients with stage III disease according to NDRG1 expression status showed a significant difference in outcomes, wherein patients with stage III NDRG1-high tumors had worse OS ($p=0.045$) and DSS ($p=0.0239$) than did patients with stage III NDRG1-low tumors (Figure 4c-d). We could not perform similar analyses of stage IV tumors owing to

small patient numbers. Interestingly, the median OS times for patients with stage III tumors differed considerably by NDRG1 expression level, being 9.1 years in NDRG1-low tumors to 4.6 years for NDRG1-high stage III tumors (Figure 4e). Similarly, the median DSS times were 4.9 years for stage III NDRG1-low tumors and 10.7 years for stage III NDRG1-high tumors (Figure 4f).

Receipt of adjuvant radiation was also an independent variable related to DSS (HR=0.510, 95% CI=0.261–0.994, $p=0.0481$) (Table 3). Patients who received adjuvant radiation had better survival outcomes than those who did not (OS, $p=0.0403$; DSS, $p=0.0137$) (Figure 5a-b). Among patients who received adjuvant radiation, those with NDRG1-high tumors showed poorer outcomes than those with NDRG1-low tumors (OS, $p=0.0088$; DSS, $p=0.0128$, Figure 5c-d). Among patients who did not receive adjuvant radiation therapy, NDRG1 expression did not correlate with survival outcomes (Figure 5e-f). The median survival times for all patients who received adjuvant radiation was 3.7 years for OS and 4.6 years for DSS. Stratification of radiation-treated patients by NDRG1 again showed distinct differences in survival time, with medians of 3.1 years for both OS and DSS for NDRG1-high tumors versus not achieved for NDRG1-low tumors (Figure 5g-h).

As expected, patients with lower tumor grades (I-II) had better outcomes than those with high-grade tumors (OS, $p=0.0399$; DSS, $p=0.0386$; Figure 6a-b). Despite the small number of low-grade tumors, we observed a significant difference in OS ($p=0.0363$) and DSS ($p=0.0210$) after stratifying for NDRG1-high versus NDRG1-low expression; patients with low-grade tumors and NDRG1-low expression had better outcomes than patients with NDRG1-high expression (Figure 6c-d). Outcomes may have been worse for patients with high-grade tumors and NDRG1-high expression relative to those with NDRG1-low expression, but those apparent differences were not statistically significant (OS, $p=0.0765$; DSS, $p=0.0699$) (Figure 6d-e). Low-grade, NDRG1-high tumors were associated with shorter survival, with median survival times of 4.3 years versus not achieved for low-grade, NDRG1-low tumors (Figure 6g-h).

DISCUSSION

IBC remains a relatively poorly defined disease that lacks specific therapeutic targets and prognostic biomarkers; the molecular characterization of IBC could advance our understanding of its unique biology and provide opportunities that could be translated into novel therapeutic strategies to improve clinical outcomes. Herein, we report that *NDRG1* protein expression was an independent predictor of poor survival outcomes for patients with IBC. In subset analyses, we report that *NDRG1*-high expression in patients with ER-negative, stage III tumors, and patients who received adjuvant radiation had worse outcomes than did patients with *NDRG1*-low tumors. Our results suggest that IBC patients could be stratified not only by known prognostic markers but also by biological determinants such as *NDRG1* expression status. Further, we found that *NDRG1* mRNA was overexpressed significantly in IBC relative to non-IBC, and in HER2+, basal-like, and TNBC aggressive breast cancer subtypes compared with luminal-like IBC tumors. These findings support our contention that *NDRG1* is associated with aggressive tumor features in IBC.

NDRG1 is a stress response gene that is highly activated and expressed in hypoxia and resistance to chemotherapy. Its function in breast cancer is widely described as a tumor and metastasis suppressor, acting mainly through inhibition of migration and invasion of cancer cells^{20-22,28}. The induction of *NDRG1* was shown in a mouse mammary tumor model to suppress metastasis by modulating WNT pathway signaling²¹. Chiang *et al.* also described how silencing *NDRG1* expression in MCF-7 cells led to increased proliferation and invasiveness of those breast cancer cells²⁸. Paradoxically, other studies showed that *NDRG1* may function as an oncogene or a prognostic biomarker in aggressive forms of breast cancer^{19,23,24,26}. Mao *et al.* found that *NDRG1* could be used as a marker for invasive breast cancer, observing that *NDRG1* expression was significantly higher in invasive breast cancer versus matched non-tumor tissues, and its levels were associated with progression from breast atypia to carcinoma. They also observed a correlation between advanced tumor stage and high *NDRG1* expression²³. Nagai *et al.* found associations between high *NDRG1* expression and worse DSS and OS in a cohort of 600 patients: the 10-year OS rate was 35% for *NDRG1*-high versus 67% for *NDRG1*-low, and *NDRG1* was an independent prognostic factor for both OS and DSS. Moreover, *NDRG1* was expressed at higher levels in stage III and IV breast cancer and

in grade 3 tumors²⁶. More recently, a study by Sevinsky and colleagues demonstrated that NDRG1 promotes breast cancer aggressiveness by altering lipid metabolism¹⁹. These observations are supported by our previous work showing that NDRG1 promotes tumorigenesis and brain metastasis in mouse models of aggressive breast cancer²⁵.

Expression of the ER is a well-known prognostic and predictive factor, and ER status is essential in the choice of treatment strategy. Patients with ER-positive breast cancer benefit from the use of hormonal therapy and have better OS than do patients with ER-negative disease, and this improvement is independent of disease stage and tumor grade²⁹⁻³¹. Our results in the present study confirmed that ER status was, indeed, an independent factor related to both OS and DSS, with ER-negative IBC patients exhibiting worse clinical outcomes. ER-negative status is associated with aggressive growth and shorter survival. Interestingly, in our study stratification of ER-negative patients by NDRG1 expression level showed significant differences in survival outcomes: ER-negative, NDRG1-high tumors were associated with worse outcomes than ER-negative, NDRG1-low tumors. However, no such difference was observed in ER-positive tumors stratified by NDRG1 expression. Our findings indicate that the clinical outcome of patients with ER-negative IBC can be stratified further based on NDRG1 expression status.

Adjuvant radiation therapy is an important part of breast cancer treatment and is known to improve breast cancer-specific survival and reduce tumor recurrence³²⁻³⁴. In the current study, we found that receipt of adjuvant radiation for IBC tumors independently correlated with improved breast cancer-specific survival. We also showed that patients who received adjuvant radiation and had NDRG1 low-expressing tumors had better clinical outcomes than did those with NDRG1-high-expressing radiation-treated tumors. These hypothesis-generating findings suggest that the role of NDRG1 in local failure in breast cancer patients and radiation resistance warrants further investigation. Previous studies of rectal cancer cells have shown that NDRG1 is one of the top highly upregulated genes in response to ionizing radiation, and that depleting it could be a promising strategy to sensitize cells to radiotherapy³⁵. Many studies have been conducted to develop a “radiosensitivity signature” to stratify patients according to

benefit from adjuvant radiation treatment³⁶⁻³⁸. However, no such molecular signature for radiation response has yet been identified.

In conclusion, NDRG1 expression was an independent prognostic factor for worse survival outcomes in patients with IBC, and together with other important prognostic factors, such as ER status and disease stage, can be used to further stratify prognostic outcome or treatment response. Because NDRG1 expression levels are generally high in IBC, targeting NDRG1 may provide a novel therapeutic strategy to improve outcome for patients with IBC.

MATERIALS AND METHODS

IBC tumor microarrays and immunohistochemical staining

This study was approved by institutional review board of The University of Texas MD Anderson Cancer Center. Details of disease diagnosis, preoperative and postoperative treatments, biomarker studies (including ER, PR, and HER2 status), and TMA construction with post-neoadjuvant residual tumors are reported elsewhere³⁹. Immunohistochemical staining of TMAs was done with a monoclonal antibody against NDRG1 (1:5000, #9485, Cell Signal) that was previously validated⁴⁰. NDRG1 staining was evaluated by percentage (0%–100%) and intensity (weak, moderate and strong) of invasive tumor cells showing cytoplasmic and/or membranous staining. NDRG1 H-score was calculated by multiplying the percentage with intensity, and the NDRG1 H-score median was used as cutoff, wherein 32 patients were grouped as NDRG1-low (\leq median), and 32 as NDRG1-high ($>$ median). Representative images of NDRG1-low and NDRG1-high tumors are shown in Figure 1.

Analysis of IBC Consortium dataset

We analyzed microarray gene expression data from the multi-institutional IBC Consortium dataset, which contains 137 IBC and 252 non-IBC tumors, to compare *NDRG1* mRNA expression between IBC and non-IBC tumors and among the molecular subtypes of breast cancer²⁷.

Statistical Analysis

Patient characteristics were summarized by NDRG1 value (Low [\leq median] vs. High [$>$ median]) and compared between patients with NDRG1-low and patients with NDRG1-high tumors. Two-sample *t* tests or Wilcoxon rank-sum tests were used for the comparison of continuous variables. Chi-square tests or Fisher's exact tests were used for the comparison of categorical variables. OS was defined as the interval from diagnosis to death, and DSS as the interval from diagnosis to death from breast cancer. Those patients without an event (death or breast cancer death) were censored at last follow-up. Kaplan-Meier curves and log-rank tests were used to compare survival distributions. Univariate and multivariate Cox proportional hazards regression models were used to compare OS (and DSS) between NDRG1-low and -high groups, adjusting for other covariates. The proportional hazards assumption was checked by scaled Schoenfeld residual plots and correlation between the scaled Schoenfeld residuals and survival time. *P* values of <0.05 indicated a statistically significant difference. SAS 9.4 (SAS institute INC, Cary, NC) was used for data analysis.

For analyses of the IBC Consortium dataset, patients were stratified as having NDRG1-high-expressing or NDRG1-low-expressing tumors according to the median *NDRG1* mRNA expression levels in the tumor samples. Mann-Whitney tests were used to compare the two categories. Black lines in each group indicate median \pm SD. GraphPad software (GraphPad Prism 8, La Jolla, CA) was used. *P* values of <0.05 were considered to indicate significant differences.

ACKNOWLEDGEMENTS

We thank Christine F. Wogan for scientific editing of the manuscript, and Carol M. Johnston from the Division of Surgery Histology Core at UT MD Anderson for her help with immunohistochemical staining.

Funding

This study was supported in part by the following grants: Susan G. Komen Career Catalyst Research grant (CCR16377813 to BGD), American Cancer Society Research Scholar grant (RSG-19-126-01 to BGD), and the State of Texas Rare and Aggressive Breast Cancer Research Program.

AUTHOR CONTRIBUTIONS

E.S.V. and B.G.D. conceived and designed the project. E.S.V. performed experiments, analyzed data and interpreted the results. Y.G. provided TMA of IBC patient samples. Y.G. L.H and E.C.Y. analyzed the TMA and interpreted results. J.S. provided statistical analysis support. N.T.U, W.A.W, and D.T. provided resources and contributed to revision of the manuscript. E.S.V. and B.G.D. wrote and edited the manuscript with input from all other authors.

COMPETING INTERESTS STATEMENT

The authors declare no conflict of interest.

REFERENCES

- 1 Chang, S., Parker, S. L., Pham, T., Buzdar, A. U. & Hursting, S. D. Inflammatory Breast Carcinoma Incidence and Survival. *Cancer* **82**, 2366-2372 (1998).
- 2 Hance, K. W., Anderson, W. F., Devesa, S. S., Young, H. A. & Levine, P. H. Trends in inflammatory breast carcinoma incidence and survival: the surveillance, epidemiology, and end results program at the National Cancer Institute. *J Natl Cancer Inst* **97**, 966-975 (2005).
- 3 Dirix, L. Y., Van Dam, P., Prove, A. & Vermeulen, P. B. Inflammatory breast cancer: current understanding. *Curr Opin Oncol* **18**, 563-571 (2006).
- 4 Wang, Z. *et al.* Pattern of distant metastases in inflammatory breast cancer - A large-cohort retrospective study. *J Cancer* **11**, 292-300 (2020).
- 5 Cristofanilli, M. *et al.* Inflammatory breast cancer (IBC) and patterns of recurrence: understanding the biology of a unique disease. *Cancer* **110**, 1436-1444 (2007).
- 6 Robertson, F. M. *et al.* Inflammatory breast cancer: the disease, the biology, the treatment. *CA Cancer J Clin* **60**, 351-375 (2010).
- 7 Fouad, T. M. *et al.* Overall survival differences between patients with inflammatory and noninflammatory breast cancer presenting with distant metastasis at diagnosis. *Breast Cancer Res Treat* **152**, 407-416 (2015).
- 8 Lim, B., Woodward, W. A., Wang, X., Reuben, J. M. & Ueno, N. T. Inflammatory breast cancer biology: the tumour microenvironment is key. *Nat Rev Cancer* **18**, 485-499 (2018).
- 9 Zhang, D. *et al.* Epidermal growth factor receptor tyrosine kinase inhibitor reverses mesenchymal to epithelial phenotype and inhibits metastasis in inflammatory breast cancer. *Clin Cancer Res* **15**, 6639-6648 (2009).
- 10 Kleer, C. G., van Golen, K. L., Braun, T. & Merajver, S. D. Persistent E-cadherin expression in inflammatory breast cancer. *Mod Pathol* **14**, 458-464 (2001).
- 11 van Golen, K. L., Wu, Z. F., Qiao, X. T., Bao, L. W. & Merajver, S. D. RhoC GTPase, a novel transforming oncogene for human mammary epithelial cells that partially recapitulates the inflammatory breast cancer phenotype. *Cancer Res* **60**, 5832-5838 (2000).
- 12 Costa, R. *et al.* Developmental therapeutics for inflammatory breast cancer: Biology and translational directions. *Oncotarget* **8**, 12417-12432 (2017).
- 13 Wang, X. *et al.* Targeting Signaling Pathways in Inflammatory Breast Cancer. *Cancers (Basel)* **12**, 2479 (2020).
- 14 Cangul, H. Hypoxia upregulates the expression of the NDRG1 gene leading to its overexpression in various human cancers. *BMC Genet* **5**, 27 (2004).

- 15 Lee, J. C. *et al.* The Iron Chelator, Dp44mT, Effectively Inhibits Human Oral Squamous Cell Carcinoma Cell
Growth in Vitro and in Vivo. *Int J Mol Sci* **17**, 1435 (2016).
- 16 Fotovati, A. *et al.* N-myc downstream-regulated gene 1 (NDRG1) a differentiation marker of human
breast cancer. *Pathol Oncol Res* **17**, 525-533 (2011).
- 17 Chen, Z. *et al.* The iron chelators Dp44mT and DFO inhibit TGF-beta-induced epithelial-mesenchymal
transition via up-regulation of N-Myc downstream-regulated gene 1 (NDRG1). *J Biol Chem* **287**, 17016-
17028 (2012).
- 18 Kovacevic, Z., Chikhani, S., Lui, G. Y., Sivagurunathan, S. & Richardson, D. R. The iron-regulated
metastasis suppressor NDRG1 targets NEDD4L, PTEN, and SMAD4 and inhibits the PI3K and Ras signaling
pathways. *Antioxid Redox Signal* **18**, 874-887 (2013).
- 19 Sevinsky, C. J. *et al.* NDRG1 regulates neutral lipid metabolism in breast cancer cells. *Breast Cancer Res*
20, 55 (2018).
- 20 Bandyopadhyay, S. *et al.* Role of the putative tumor metastasis suppressor gene Drg-1 in breast cancer
progression. *Oncogene* **23**, 5675-5681 (2004).
- 21 Liu, W. *et al.* N-myc downstream regulated gene 1 modulates Wnt-beta-catenin signalling and
pleiotropically suppresses metastasis. *EMBO Mol Med* **4**, 93-108 (2012).
- 22 Godbole, M. *et al.* Up-regulation of the kinase gene SGK1 by progesterone activates the AP-1-NDRG1
axis in both PR-positive and -negative breast cancer cells. *J Biol Chem* **293**, 19263-19276 (2018).
- 23 Mao, X. Y. *et al.* Increased N-myc downstream-regulated gene 1 expression is associated with breast
atypia-to-carcinoma progression. *Tumour Biol* **32**, 1271-1276 (2011).
- 24 Li, E. Y. *et al.* Aryl Hydrocarbon Receptor Activates NDRG1 Transcription under Hypoxia in Breast Cancer
Cells. *Sci Rep* **6**, 20808 (2016).
- 25 Villodre, E. S. *et al.* NdrG1-egfr axis in inflammatory breast cancer tumorigenesis and brain metastasis
[abstract]. In: *Proceedings of the 2019 San Antonio Breast Cancer Symposium; 2019 Dec 10-14; San
Antonio, TX. Philadelphia (PA): AACR Cancer Res* **80**, supplementary 4 (2020).
- 26 Nagai, M. A. *et al.* Prognostic value of NDRG1 and SPARC protein expression in breast cancer patients.
Breast Cancer Res Treat **126**, 1-14 (2011).
- 27 Van Laere, S. J. *et al.* Uncovering the molecular secrets of inflammatory breast cancer biology: an
integrated analysis of three distinct affymetrix gene expression datasets. *Clin Cancer Res* **19**, 4685-4696
(2013).
- 28 Chiang, K. C. *et al.* WNT-1 inducible signaling pathway protein-1 enhances growth and tumorigenesis in
human breast cancer. *Sci Rep* **5**, 8686 (2015).
- 29 Yip, C. H. & Rhodes, A. Estrogen and progesterone receptors in breast cancer. *Future Oncol* **10**, 2293-
2301 (2014).
- 30 Grann, V. R. *et al.* Hormone receptor status and survival in a population-based cohort of patients with
breast carcinoma. *Cancer* **103**, 2241-2251 (2005).
- 31 Munoz, D. F. & Plevritis, S. K. Estimating Breast Cancer Survival by Molecular Subtype in the Absence of
Screening and Adjuvant Treatment. *Med Decis Making* **38**, 32S-43S (2018).
- 32 Daugherty, E. C., Daugherty, M. R., Bogart, J. A. & Shapiro, A. Adjuvant Radiation Improves Survival in
Older Women Following Breast-Conserving Surgery for Estrogen Receptor-Negative Breast Cancer. *Clin
Breast Cancer* **16**, 500-506 e502 (2016).
- 33 Speers, C. & Pierce, L. J. Postoperative Radiotherapy After Breast-Conserving Surgery for Early-Stage
Breast Cancer: A Review. *JAMA Oncol* **2**, 1075-1082 (2016).
- 34 Ohri, N. & Haffty, B. G. The evolution of adjuvant radiation therapy for early-stage and locally advanced
breast cancer. *Breast J* **26**, 59-64 (2020).
- 35 Kim, S. C., Shin, Y. K., Kim, Y. A., Jang, S. G. & Ku, J. L. Identification of genes inducing resistance to
ionizing radiation in human rectal cancer cell lines: re-sensitization of radio-resistant rectal cancer cells
through down regulating NDRG1. *BMC Cancer* **18**, 594 (2018).
- 36 Servant, N. *et al.* Search for a gene expression signature of breast cancer local recurrence in young
women. *Clin Cancer Res* **18**, 1704-1715 (2012).

- 37 Tramm, T. *et al.* Development and validation of a gene profile predicting benefit of postmastectomy radiotherapy in patients with high-risk breast cancer: a study of gene expression in the DBCG82bc cohort. *Clin Cancer Res* **20**, 5272-5280 (2014).
- 38 Speers, C. *et al.* Development and Validation of a Novel Radiosensitivity Signature in Human Breast Cancer. *Clin Cancer Res* **21**, 3667-3677 (2015).
- 39 Gong, Y. *et al.* Polycomb group protein EZH2 is frequently expressed in inflammatory breast cancer and is predictive of worse clinical outcome. *Cancer* **117**, 5476-5484 (2011).
- 40 Zeng, L. *et al.* Prognostic value of biomarkers EpCAM and alphaB-crystallin associated with lymphatic metastasis in breast cancer by iTRAQ analysis. *BMC Cancer* **19**, 831 (2019).

Figure 1




Figure 1. Immunohistochemical staining of NDRG1 in IBC tumors. Representative images of NDRG1 immunostaining of **(a)** an NDRG1-low IBC tumor and **(b)** an NDRG1-high IBC tumor.

Table 1. Clinical and pathologic characteristics of tumor samples from patients with IBC according to NDRG1 expression.

Covariate	Level	NDRG1-low (n=32)	NDRG1-high (n=32)	p-value
Age		51.5 ± 12.1	48.6 ± 12	0.3861
Race	Non-White	8 (25%)	5 (16%)	0.5356
	White	24 (75%)	27 (84%)	
Histologic type	Others	2 (6%)	4 (13%)	0.6719
	Ductal	30 (94%)	28 (87%)	
Grade	1-2	6 (19%)	7 (22%)	0.7560
	3	26 (81%)	25 (78%)	
Lymphovascular invasion	No	4 (14%)	3 (10%)	0.7065
	Yes	25 (86%)	27 (90%)	
Stage	III	31 (97%)	27 (84%)	0.1961
	IV	1 (3%)	5 (16%)	
Estrogen receptor	No	21 (66%)	19 (61%)	0.7209
	Yes	11 (34%)	12 (39%)	
Progesterone receptor	No	24 (75%)	19 (61%)	0.2425
	Yes	8 (25%)	12 (39%)	
HER2	No	12 (37%)	22 (71%)	0.0077
	Yes	20 (63%)	9 (29%)	
Triple-negative breast cancer	No	27 (84%)	20 (64%)	0.0879
	Yes	5 (16%)	11 (36%)	
Adjuvant radiation	No	10 (31%)	11 (34%)	0.7901
	Yes	22 (69%)	21 (66%)	

Table 2. Univariate Cox regression analysis on overall survival and disease-specific survival among patients with IBC.

Covariate	Level	Overall survival			Disease-specific survival		
		HR	95% CI	p-value	HR	95% CI	p-value
Age	1 Unit Change	1.007	0.980 - 1.035	0.6080	1.000	0.971 - 1.029	0.9904
NDRG1	Low	1.000			1.000		
	High	2.107	1.155 - 3.842	0.0150	2.354	1.235 - 4.485	0.0092
Race	Non-white	1.000			1.000		
	White	1.823	0.769 - 4.321	0.1724	1.940	0.757 - 4.973	0.1676
Histologic type	Others	1.000			1.000		
	Ductal	1.027	0.368 - 2.870	0.9591	1.153	0.410 - 3.243	0.7874
Grade	1-2	1.000			1.000		
	3	2.404	1.014 - 5.698	0.0463	2.612	1.020 - 6.688	0.0453
Lymphovascular invasion	No	1.000			1.000		
	Yes	1.684	0.601 - 4.717	0.3216	1.493	0.529 - 4.213	0.4486
Stage	III	1.000			1.000		
	IV	4.638	1.847 - 11.647	0.0011	5.485	2.138 - 14.069	0.0004
Estrogen receptor	No	1.000			1.000		
	Yes	0.414	0.212 - 0.808	0.0098	0.426	0.211 - 0.860	0.0173
Progesterone receptor	No	1.000			1.000		
	Yes	0.699	0.359 - 1.361	0.2919	0.816	0.412 - 1.616	0.5598
HER2	No	1.000			1.000		
	Yes	0.748	0.409 - 1.366	0.3445	0.602	0.313 - 1.161	0.1300
Triple-negative breast cancer	No	1.000			1.000		
	Yes	1.557	0.810 - 2.992	0.1844	1.692	0.852 - 3.358	0.1328
Adjuvant radiation	No	1.000			1.000		
	Yes	0.538	0.295 - 0.982	0.0434	0.486	0.259 - 0.914	0.0252




Figure 2. NDRG1 is a predictor of poor outcome in patients with IBC. Kaplan-Meier analysis showed that patients whose tumors had NDRG1-high expression had **(a)** worse overall survival and **(b)** worse disease-specific survival than did patients whose tumors had NDRG1-low expression.

Table 3. Multivariate Cox regression analysis on overall survival and disease-specific survival among patients with IBC.

Covariate	Level	Overall survival			Disease-specific survival		
		HR	95% CI	p-value	HR	95% CI	p-value
NDRG1	Low	1.000			1.000		
	High	2.034	1.081 - 3.822	0.0274	2.287	1.157 - 4.522	0.0174
Estrogen receptor	No	1.000			1.000		
	Yes	0.313	0.156 - 0.628	0.0011	0.307	0.147 - 0.642	0.0017
Stage	III	1.000			1.000		
	IV	3.212	1.171 - 8.814	0.0234	3.844	1.362 - 10.854	0.0110
Adjuvant radiation	No	1.000			1.000		
	Yes	0.562	0.298 - 1.058	0.0741	0.510	0.261 - 0.994	0.0481

Figure 3




Figure 3. Overall survival and disease-specific survival in patients with IBC stratified by ER status and NDRG1 expression. Patients with estrogen receptor (ER)-negative tumors had **(a)** worse overall survival and **(b)** worse disease-specific survival versus patients with ER-positive tumors. **(c, d)** Stratification of patients by ER and NDRG1 expression status in terms of overall survival and disease-specific survival. Log-rank tests were used to obtain p values. **(e, f)** Median overall survival and disease-specific survival times, in years, for patients stratified by ER status and NDRG1 expression.

Figure 4




Figure 4. Overall survival and disease-specific survival in patients with IBC stratified by disease stage and NDRG1 expression. Patients with stage III IBC had better (a) overall survival and (b) disease-specific survival than did patients with stage IV IBC. (c, d) Patients with stage III IBC stratified by NDRG1 expression in terms of overall survival and disease-specific survival. Log-rank tests were used to obtain p values. (e, f) Median overall survival and disease-specific survival times, in years, for patients stratified by disease stage and NDRG1 expression.

Figure 5




Figure 5. High NDRG1 expression correlated with worse outcomes among patients who received adjuvant radiation therapy. IBC patients who received adjuvant radiation had better **(a)** overall survival **(b)** and disease-specific survival than did patients who did not receive radiation. **(c, d)** Patients who received adjuvant radiation treatment stratified by NDRG1 expression in terms of overall survival and disease-specific survival. **(e, f)** Patients who did not receive adjuvant radiation stratified by NDRG1 expression in terms of overall survival and disease-specific survival. Log-rank tests were used to obtain p values. **(g, h)** Median overall survival and disease-specific survival times, in years, for patients stratified by NDRG1 expression and adjuvant radiation treatment status.

Figure 6




Figure 6. Overall survival and disease-specific survival in patients with IBC stratified by tumor

grade and NDRG1 expression. Patients with IBC and low-grade cancer had better **(a)** overall survival and **(b)** disease-specific survival than did patients with grade III disease. **(c, d)** Patients with low-grade IBC stratified by NDRG1 expression in terms of overall survival and disease-specific survival. **(e, f)** Patients with high-grade IBC stratified by NDRG1 expression in terms of overall survival and disease-specific survival. Log-rank tests were used to obtain *p* values. **(g, h)** Median overall survival and disease-specific survival times, in years, for patients stratified by NDRG1 expression and tumor grade.