Novel taxa of Acidobacteriota involved in seafloor sulfur cycling

Mathias Flieder^a, Joy Buongiorno^{c,#}, Craig W. Herbold^a, Bela Hausmann^{a,d,e}, Thomas Rattei^f, Karen G. Lloyd^c, Alexander Loy^{a,b,d,*}, Kenneth Wasmund^{a,b,g,*}

^aDivision of Microbial Ecology, Centre for Microbiology and Environmental Systems Science, University of Vienna, Vienna, Austria.

^bAustrian Polar Research Institute, Vienna, Austria.

^cDepartment of Microbiology, University of Tennessee, Knoxville, USA.

^dJoint Microbiome Facility of the Medical University of Vienna and the University of Vienna, Vienna, Austria.

^eDepartment of Laboratory Medicine, Medical University of Vienna, Vienna, Austria.

^fDivision of Computational Systems Biology, Centre for Microbiology and Environmental Systems Science, University of Vienna, Vienna, Austria.

^gCenter for Microbial Communities, Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark.

[#]Current affiliation: Division of Natural Sciences, Maryville College, Maryville, Tennessee USA.

*Corresponding authors: Kenneth Wasmund (kwasmund@gmail.com) and Alexander Loy (alexander.loy@univie.ac.at)

Abstract

Acidobacteriota are widespread and often abundant in marine sediments, yet their metabolic and ecological properties are poorly understood. Here, we examined metabolisms and distributions of Acidobacteriota in marine sediments of Svalbard by functional predictions from metagenome-assembled genomes (MAGs), amplicon sequencing of 16S rRNA and dissimilatory sulfite reductase (*dsrB*) genes and transcripts, and gene expression analyses of tetrathionate-amended microcosms. Acidobacteriota were the second most abundant *dsrB*-harboring (averaging 13%) phylum after Desulfobacterota in Svalbard sediments, and represented 4% of *dsrB* transcripts on average. We propose two new Acidobacteriota genera, *Candidatus* Sulfomarinibacter (class Thermoanaerobaculia, 'sub-division 23') and *Ca. Polarisedimenticola* ('sub-division 22'), with distinct genetic properties that may explain their distributions in biogeochemically distinct fjord sediments. *Ca. Sulfomarinibacter* encodes flexible respiratory routes, with potential for oxygen, nitrous oxide, metal-oxide, tetrathionate, sulfur and sulfite/sulfate respiration, and possibly sulfur disproportionation. Potential nutrients and energy include cellulose, proteins, cyanophycin, hydrogen and acetate. A *Ca. Polarisedimenticola* MAG encodes enzymes to degrade proteins, and to reduce oxygen, nitrate, sulfur/polysulfide and metal-oxides. 16S rRNA gene and transcript profiling showed *Ca. Sulfomarinibacter* members were relatively abundant and transcriptionally active in sulfidic fjord sediments, while *Ca. Polarisedimenticola* members were more relatively abundant in metal-rich fjord sediments. Overall, we reveal various physiological features of uncultured marine Acidobacteriota that indicate fundamental roles in seafloor biogeochemical cycling.

41 Introduction

42 Bacteria of the phylum *Acidobacteriota* (also known as 'Acidobacteria') are highly diverse and
 43 inhabit a vast array of environments on Earth, yet the properties of various *Acidobacteriota* lineages
 44 remain poorly understood [1–6]. Knowledge regarding the functions and ecology of *Acidobacteriota* is
 45 biased to isolates and genomes obtained from soils, where they are especially prevalent and often
 46 dominate microbial communities [3, 4]. Soil-derived *Acidobacteriota* are generally known as aerobic
 47 heterotrophs that utilize various carbohydrates including polysaccharides like chitin or cellulose [3, 7,
 48 8]. Some *Acidobacteriota* known from other environments have unique physiological properties, such
 49 as the ability to reduce iron [9], perform phototrophy [9, 10], or exhibit thermophilic lifestyles [11].
 50 Members of *Acidobacteriota* sub-divisions 1 and 3 from peatland and permafrost soils have the
 51 potential to dissimilate inorganic and/or organic sulfur compounds [2, 12]. In comparison to terrestrial
 52 *Acidobacteriota*, even less is known about *Acidobacteriota* in marine systems.

53 *Acidobacteriota* 16S rRNA genes or genomes are frequently detected in marine environments
 54 including ocean waters, marine sponges, hydrothermal vents, or sediments [13–17]. Studies of 16S
 55 rRNA genes in marine sediments showed that *Acidobacteriota* are widespread and reach relative
 56 abundances in amplicon libraries of up to 23% [18–23]. This suggests they play important roles in
 57 microbial community functioning and biogeochemical processes, although our knowledge regarding
 58 their specific roles in sediments remains limited. A recent stable isotope probing study showed some
 59 *Acidobacteriota* in deep-sea sediments are capable of fixing nitrogen [24]. *Acidobacteriota* were also
 60 shown to be active, by incorporation of isotopically-labelled tracer into their DNA, under sulfidic
 61 conditions in incubations with estuarine sediment [6]. One novel *Acidobacteriota* metagenome-
 62 assembled genome (MAG) (*Candidatus* Guanabacteria) had genes for the CO dehydrogenase/CO-
 63 methylating acetyl-CoA synthase complex and heterodisulfide reductases, indicating a possible
 64 anaerobic lifestyle [25].

65 Marine sediments are a massive global habitat for microorganisms [26], with cell densities of
 66 microorganisms average up to 10^9 cells per cm^3 in surface sediments of organic-rich sediments [27].
 67 Substantial amounts of organic matter are processed in marine sediments, which makes them a critical
 68 component of marine and global biogeochemical cycles [28]. Marine sediments are often stratified with
 69 respect to redox states, whereby oxygen is typically depleted within millimetres to centimetres below

the surface at sites where organic inputs are relatively high [29]. Vast expanses of sediments are therefore anoxic, and many microorganisms survive via anaerobic lifestyles, such as fermentation, or respiration of nitrate, metals, sulfate or CO₂. Sulfate is abundant in sediments and is used by sulfite/sulfate-reducing microorganisms (SRMs) as an electron acceptor for anaerobic respiration. Sulfate reduction is estimated to facilitate approximately 29% of organic matter degradation in marine sediments globally [26, 28]. The sulfur cycle is therefore a major driver of microbial life and biogeochemical cycling in the seafloor, so understanding the microorganisms that catalyze sulfur cycling is of great importance.

Because sulfate reduction is a major process in marine sediments, the activities, distributions and diversity of SRMs have been relatively well studied [28, 30, 31]. Members of the Desulfobacterota (formerly ‘Deltaproteobacteria’) are known as abundant SRMs in marine sediments, playing key roles in anaerobic food webs by utilizing fermentation products released by primary degraders of organic matter [32–34]. They are also represented by various isolates, and many have been subject to genomic and physiological studies [35]. Surveys of functional marker genes for sulfite/sulfate reducers in marine sediments, i.e., of dissimilatory sulfite reductases (*dsrAB*), have repeatedly shown that *dsrAB* from the phylum Desulfobacterota are typically the dominant *dsrAB*-harbouring group in marine sediments, but importantly, that several other lineages of *dsrAB*-harbouring uncultivated organisms are also abundant and prevalent [36]. Recently, some *dsrAB* sequences in marine sediments have been inferred to belong to Acidobacteriota [6, 37], although nothing is known about the metabolic properties or the sulfur dissimilating pathways of the organisms that harbour these genes. Identifying and understanding these undescribed *dsrAB*-harbouring microorganisms is therefore critical for understanding the microbial groups that drive sulfur cycling in marine sediments.

In this study, we aimed to gain insights into the metabolic potential of uncultured Acidobacteriota in marine sediments. We therefore recovered metagenome-assembled genomes (MAGs) from abundant Acidobacteriota populations present in marine fjord sediments of Svalbard, and predicted their metabolic features. Focus was placed on MAGs from the Thermoanaerobaculia, which represent a newly described lineage of *dsrAB*-harbouring organisms that may be important sulfur cycling bacteria in marine sediments. These analyses were complemented with comparative genomics, incubation experiments, transcript analysis, and analyses of Acidobacteriota distributions in Svalbard

sediments, together revealing they may play various roles in sedimentary biogeochemical cycles, and that they are a prominent group of sulfur-dissimilating organisms.

Materials and Methods

Sample collection and microcosms

Marine sediments were collected from Smeerenburgfjorden, Kongsfjorden and Van Keulenfjorden, Svalbard, Norway, in July 2016 and/or June 2017 with the vessel 'MS Farm'. Extensive biogeochemical data for these sites is available from previous studies [38–42]. Maps of sample locations are presented in Michaud *et al.* 2020. From Smeerenburgfjorden, samples were taken from three stations: station GK (79°38.49N, 11°20.96E), station J (79°42.83N, 11°05.10E) and station GN (79°45.01N, 11°05.99E). Samples from Van Keulenfjorden were taken from sites AC (77°32.260'N, 15°39.434'E) and AB (77°35.249'N, 15°05.121'E). A sample was also taken from Kongsfjorden station F (78°55.075' N, 12°15.929' E) [43]. For molecular biological analyses, samples were taken with HAPS [44] or Rumohr corers [45]. Details of core subsampling procedures and microcosm incubations are provided in the Supplementary information. Additional samples for non-quantitative microscopy were taken from tidal flat sediments of Aveiro Lagoon, Portugal (40°34.14N 8°45.10W) in July 2019, and from Kristineberg station near Fiskebäckskil, Sweden (58°24.95N, 11°44.50E) in October 2019, with further details provided in the Supplementary information.

Nucleic acid extractions and reverse-transcription

For amplicon-based analyses, DNA and RNA was extracted from the sediment core samples (~500 µl) and microcosm samples (~250 µl) using the RNeasy PowerSoil Total RNA Kit (Qiagen) according to the manufacturer's instructions. Additionally, a phenol/chloroform based extraction method was used to extract nucleic acids from sediment samples from station J sampled in July 2016 (Supplementary information). Eluted nucleic acids were stored in molecular biology grade water at –80°C. Aliquots for DNA-based analyses were used as eluted, while aliquots for RNA-based analyses were DNase-treated using the TURBO DNA-free™ kit (Thermo Fisher), followed by reverse transcription of the RNA to cDNA using the RevertAid First Strand cDNA Synthesis Kit (Thermo Fisher) according to the manufacturer's instructions. To test if any DNA remained in the RNA samples after the DNase digestion step, control samples were processed as above except the RevertAid M-MuLV

Reverse Transcriptase was excluded. These controls were checked for DNA by PCR using 16S rRNA gene targeting primers (described below).

Sediment samples from 2016 were used for metagenome sequencing. DNA was extracted by the Vienna group from 3–5 mL of sediment from varying depths or microcosms derived from station J, Smeerenburgfjorden, and 18 centimeters below seafloor (cmbsf) from station AC of Van Keulenfjorden (Supp. Table 1) using the DNeasy PowerSoil Kit (Qiagen) according to the manufacturer's protocol. DNA was also extracted by the Knoxville group from 2 grams of sample spanning 0–5 cmbsf from site AB of Van Keulenfjorden and site F of Kongsfjorden (Supp. Table 1), using the RNeasy PowerSoil Kit (Qiagen) with DNA elution following the manufacturer's protocol.

Metagenome sequencing and genome binning

DNA libraries were prepared (detailed in Supplementary information) and sequenced using 2×150 bp paired-end mode on an Illumina HiSeq 3000 instrument at the Biomedical Sequencing Facility (BSF), Vienna. Metagenomic libraries were generated from the combined extracts from the first 5 cm (spanning 0 to 5 cm downcore) in sites AB and F in the Center for Environmental Biotechnology, Knoxville, using Illumina HiSeq, 2×250 bp in paired-end mode [43]. Sequencing output summaries are provided in Supp. Table 1.

Sequence reads were quality filtered, trimmed, and normalized as described in the Supplementary information. Processed reads from each sample were assembled separately using IDBA-UD (version 1.1.1) [46] with default settings and the following options: --min_contig 500 --pre_correction. Reads from site F (Kongsfjorden) were assembled via metaSPAdes (version 3.11) [47] with kmer sizes set to 21, 33, 55, 77, 99, and 127 to find the best assembly. All other samples were assembled using metaSPAdes on the KBase server [48] with the default parameters and following options: minimum contig length of 1000 bp, and kmer sizes of 21, 33, and 55. All samples were also assembled using Megahit [49] on the KBase server using default parameters.

Coverage profiles of assembled unbinned contigs were acquired by mapping trimmed reads (not normalised) to assemblies using BWA [50] and SAMtools [51]. Contigs from each assembly were then binned into metagenome-assembled genomes (MAGs) using MetaBat2 (using each binning strategy) (version 2.12.1) [52], CONCOCT (version 0.4.1) [53] and MaxBin2 (version 2.2.4). MAG collections derived from each binning strategy, from all respective assemblies, were then aggregated

using DasTool (version 1.1.0) (Supp. Fig. 1A) [54]. Finally, all MAGs were dereplicated using dRep (version 1.4.3) [55], with the options: an average nucleotide identity (ANI) of 98% was used as cut-off to dereplicate MAGs from the secondary ANI comparison [56], and MAGs >50% complete and <10% contamination were retained. Estimations of completeness and degree of contamination of MAGs were obtained by CheckM (version 1.0.7) [57]. Read mapping to compare relative abundances of read recruitment to MAGs was performed using BMap [58], with the default settings and 'minid' of 0.99 for the minimum identity threshold. Taxonomic affiliations of MAGs were determined with GTDB-Tk [59]. ANI comparisons of MAGs were obtained using JSpeciesWS server based on BLASTN ('ANiB') [60] and ANIcalculator [61].

Gene annotations and in silico analyses of inferred proteins

Calling of genes and annotations were performed via RAST [62]. Functions of predicted proteins of interest were manually checked after searches with BLASTP [63] against the NCBI-nr and SWISS-PROT databases [64] (>30% identity), and the Conserved Domain Database (CDD) [65] (default expect value of 0.01). Functional predictions for proteins were also evaluated using literature searches and the MetaCyc database [66]. Methods for further annotations and protein sequence analyses are described in the Supplementary information.

MiSeq amplicon sequencing

For amplification of bacterial and archaeal 16S rRNA genes or transcripts (cDNA) from Smeerenburgfjorden sediments, the primers 515F (5'-GTGYCAGCMGCCGCGGTAA-3') [67] and 806R (5'-GGACTACNVGGGTWTCTAAT-3') [68] including a 5'-head sequence for 2-step PCR barcoding [69], were used (further details in Supplementary information). Slight variants of these PCR primers 515F and 806R [70] for 16S rRNA genes were used in amplicon sequencing profiling of sediments from Van Keulenfjorden in a previous study, although a standard 'one-step PCR' approach was used [40]. Amplicon pools were extracted from the raw sequencing data using the FASTQ workflow in BaseSpace (Illumina) with default parameters. Demultiplexing was performed with the python package demultiplex (Laros JFJ, github.com/jfjaros/demultiplex) allowing one mismatch for barcodes and two mismatches for linkers and primers. DADA2 [71] was used for demultiplexing amplicon sequencing variants (ASVs) using a previously described standard protocol [72]. FASTQ reads 1 and 2 were trimmed at 220 nt and 150 nt with allowed expected errors of 2. Taxonomy was assigned to 16S rRNA gene/transcript

sequences based on SILVA taxonomy (release 138) using the naïve Bayesian classification method as implemented in *mothur* [73]. Amplicon sequence datasets were analyzed with the Rhea pipeline [74] implemented in R (<https://www.r-project.org/>).

Primers DSR-1762Fmix and DSR-2107Rmix, including a 5'-head sequence for barcoding, were used for amplification of *dsrB*-genes or -transcripts (cDNA) [75] (further details in Supplementary information). Raw reads were then processed as previously described [69, 75], into *dsrB* operational taxonomic units (OTUs) with >99% identity. Classification of DsrB sequences was performed using a combined phylogenetic and naïve Bayesian classification approach as previously described [75].

Quantitative reverse-transcription PCR

RT-qPCR assays targeting the octaheme cytochrome tetrathionate reductase (*otr*) and *dsrB* genes of MAG AM3-C were performed using the newly-designed primers TetraC-C-F (5'-CACCACGACCTGTCTCGG-3') and TetraC-C-R (5'-CCCCCTGGAGTTCTTGGT-3'), and Acido-dsrB-F (5'-GGAGAACTATGGGAAGTGGG-3') and Acido-dsrB-R (5'-GTTGAGGCAGCACGCGTA-3'). Primers 1329-B-F (5'-AACCTTTGGGCGATTTCTCG-3') and 1329-B-R (5'-GAGAGAGTGGCAACGTGAAC-3') targeting the DNA-directed RNA polymerase alpha subunit gene of MAG AM3-C were used to examine expression of a housekeeping gene. Details of RT-qPCR assay conditions are presented in the Supplementary information.

Phylogenetic analyses

A phylogenomic maximum-likelihood tree was created using the IQ-TREE web-server with automatic substitution model selection and ultra-fast bootstrapping (1000×) [76] using an alignment of concatenated protein sequences derived from single copy marker genes retrieved from CheckM [57]. The tree was visualized with iTol [77]. Phylogenetic analysis of 16S rRNA was performed in ARB [78] using the SILVA database release 138 [79], and *dsrAB* sequences were also analysed using ARB using previously described database [36, 75] (Supplementary information). Phylogenetic analyses of all other protein sequences were performed using the IQ-TREE web-server with automatic substitution model selection and ultra-fast bootstrapping (1000×) [76]. For the Complex-Iron-Sulfur-Molybdoenzyme (CISM) tree, query protein sequences were added to a pre-computer alignment of CISM protein sequences [80], using MAFFT using the 'add full length sequences' option (--add) [81]. All other proteins sequence alignments were made *de novo* with MUSCLE [82] within Mega6 [83].

Catalyzed reporter deposition-fluorescence in situ hybridisation (CARD-FISH)

Sediment samples from Svalbard, Portugal and Sweden were fixed with 4% formaldehyde for 3 hrs on ice and stored in PBS:ethanol (1:1) at -20°C using standard procedures [84]. Cells were extracted from sediments using Nycodenz density gradients (Supplementary information). Hybridisations were performed using the 5'-horseradish peroxidase-labeled (HRP) probe Acido-Sva-34-HRP (5'-GACTTATGTCATTGAGGACTCATGCGG-3') and unlabelled helper probes (5'-GGATAGCCTCGGGAAACCGAGGGTAA-3') and (5'-TGAGGGGAAAGGCGGGG-3'), or with HoAc1402-HRP (5'-CTTTCGTGATGTGACGGG-3') with competitor compHoAc1402 (5'-CTTTCGTGACGTGACGGG-3') [85]. Further details of hybridisation methods and probes are provided in the Supplementary information.

Sequence and MAG accessions

Metagenomic sequence reads from Van Keulenfjorden and Kongsfjorden samples are available under NCBI-Genbank Bioproject PRJNA493859. Metagenomic sequence reads, and 16S rRNA gene and *dsrB* sequence reads from Smeerenberfjorden samples are available under NCBI-Genbank Bioproject PRJNA623111. Metagenome-assembled genomes are available under NCBI-Genbank Bioproject PRJNA623111, with Biosample accessions SAMN15691661-SAMN15691666.

Results

Recovery of novel Acidobacteriota genomes from marine sediments

Metagenomic sequencing and genome binning was performed from DNA extracted and sequenced from sediments originating from three fjords from Svalbard, Norway (Supp. Table 1). Our genome binning strategy based on multiple assemblies and multiple binning algorithms recovered more MAGs with higher completeness, as compared to applying multiple binning approaches based on single assembly approaches (Supp. Fig. 1A and Supp. Fig. 1B). From the dereplicated MAGs ($n=97$), four represented populations of the phylum Acidobacteriota and were chosen for in-depth analyses.

Phylogenomic analyses showed three MAGs (AM1, AM2 and AM3-A) affiliated with GTDB family 'FEB-10' of the class Thermoanaerobaculia ('sub-division 23') (Fig. 1). We included two additional MAGs in our analyses, i.e., AM3-B and AM3-C, that were highly similar to the AM3-A MAG (>98% ANI), but were classified as redundant during MAG dereplication. They encoded enzymes of

interest, and were more complete than MAG AM3-A (Table 1). Comparisons of ANI values suggested these MAGs represent three distinct species (<95% ANI) (Supp. Table. 2) [86], all from a novel genus for which we propose the name *Candidatus Sulfomarinibacter*. The MAG AM3-C represents the type species *Ca. Sulfomarinibacter kjeldsenii* (Supp. Table 3). MAG AM4 represents the type species of another novel genus affiliated with the GTDB class 'Mor1' ('sub-division 22') (Fig. 1), and for which we propose the name *Ca. Polarisedimenticola svalbardensis* (Table 1 and Supp. Table 3).

Marine Acidobacteriota encode the full dissimilatory sulfate reduction pathway

Together, the gene content of the *Ca. Sulfomarinibacter* MAGs suggests they encode a complete canonical dissimilatory sulfate reduction pathway (Fig. 2 and Supp. Table 4). This includes enzymes required for sulfate activation to APS (Sat) and reduction of APS to sulfite (AprAB, QmoABC), and further reduction of sulfite to sulfide (DsrAB, DsrC, DsrMKJOP, DsrN) (Supp. Table 4). Acidobacteriota *dsr* were also found on scaffolds (up to 20 kb) that were not binned into MAGs, yet had highly similar genes and therefore derive from closely related populations, e.g., >99% *dsrB* nucleotide identity (Fig. 3 and Supp. Table 4). The unbinned acidobacteriotal contig 'ThM_scaffold_807' harboured all *dsr* on one contig (Fig. 3). The predicted DsrC had two conserved cysteine residues critical for respiratory functioning (Supp. Fig. 2) [87]. Similar to Acidobacteriota MAGs from peatlands and permafrost [2, 12], the marine Acidobacteriota encoded both DsrL and DsrD proteins. DsrL acts as a NAD(P)H:acceptor oxidoreductase for DsrAB [88], while the function of DsrD has not been proven, it is possibly a transcriptional regulator [89]. The DsrL sequences were phylogenetically related to group 'DsrL-2' from *Desulfurella amilsii*, peatland Acidobacteriota and other subsurface bacteria, and were phylogenetically distinct from group 'DsrL-1' of sulfur-oxidizing aerobes (Supp. Fig. 3A) [136]. The DsrL had conserved YRR-motifs in the NAD(P)H substrate-binding domains that are present in the DsrL-2 group, and absent in DsrL-1 of sulfur-oxidizing aerobes (Supp. Fig. 3B) [136].

The DsrAB sequences from the novel Acidobacteriota MAGs and unbinned metagenomic contigs are affiliated with the 'Uncultured family-level DsrAB lineage 9' within the 'Environmental supercluster 1', which is part of the 'reductive, bacterial-type DsrAB branch' in the DsrAB tree [36] (Fig. 4). Sequences of 'lineage 9' are primarily derived from marine sediments [36]. This lineage is closely related to the 'Uncultured family-level lineage 8' that harbours DsrAB sequences from peatland and permafrost derived Acidobacteriota of subdivisions 1 and 3 [2, 12]. We also identified several 'lineage

9' *dsrA* and/or *dsrB* sequences in Acidobacteriota MAGs from public databases that derived from marine or groundwater environments (Fig. 4). Herein, we refer to this clade as the 'Thermoanaerobaculia Dsr lineage'.

Marine Acidobacteriota use tetrathionate and potentially also other sulfur cycle intermediates

Several *Ca. Sulfomarinibacter* MAGs encoded c-type cytochromes annotated as octaheme tetrathionate reductases (Otr), which was supported by phylogenetic analysis (Supp. Fig. 4) [90]. The Otr were predicted to be periplasmic and may enable respiration with tetrathionate, a sulfur compound of intermediate oxidation state ('sulfur cycle intermediate' (SCI)) [31] (Fig. 2). Transcription of *otr* in Svalbard sediment microcosms with or without tetrathionate additions was analysed by RT-qPCR analysis of mRNA of *otr* of *Ca. Sulfomarinibacter* MAG AM3-C. This showed *otr* was upregulated (1.8-fold) at day 1 although not significantly, and was significantly upregulated ($p < 0.0488$) at day 8 (36-fold). The transcription of *dsrB* appeared lower at both days in tetrathionate-amended microcosms (0.48-0.63-fold), although not significantly (Fig. 5).

'YTD gene clusters' encoding sulfur-trafficking rhodanase-like proteins [91] were identified among *Ca. Sulfomarinibacter* MAGs. Genes for YedE-related permease-like proteins, a DsrE2-like protein, a rhodanase-domain containing sulfur carrier TusA, and two conserved hypothetical proteins were present (Supp. Table 4). The TusA sulfurtransferase had conserved Cys-Pro-X-Pro sulfane sulfur-binding domains (Supp. Fig. 6A). The TusA were phylogenetically most closely related to various TusA from anaerobic Desulfobacterota that are capable of reducing and/or disproportionating inorganic sulfur compounds such as elemental sulfur, sulfite and/or thiosulfate (Supp. Fig. 6B). Together, this suggested *Ca. Sulfomarinibacter* are capable of internal trafficking of sulfur, and may use it to reduce and/or disproportionate inorganic sulfur compounds of intermediate redox states.

The marine Acidobacteriota MAGs encoded several Complex-Iron-Sulfur-Molybdoenzyme (CISM) enzymes that may catalyse redox reactions of sulfur compounds. The *Ca. Sulfomarinibacter* MAG AM3-A encoded a putative tetrathionate reductase (TtrA) (Supp. Fig. 6), and also had an adjacent TtrB (FeS protein) encoded. A *ttrC* encoding a membrane anchor was missing, although the *ttrAB* were situated on the end of the contig and therefore *ttrC* may have been present in DNA that either was not sequenced or was not binned. The Ttr complex may provide an additional means to reduce tetrathionate.

Ca. P. svalbardensis MAG AM4 had genes for a CISM subunit A enzyme that phylogenetically affiliated with the polysulfide/thiosulfate reductase clade ('Psr') (Supp. Fig. 6). Subunits for PsrABC were encoded in a gene cluster, where the terminal reductase PsrA had a TAT-leader peptide for export from the cytoplasm, PsrB had FeS domains for electron transfer between PsrA and PsrC, and the PsrC subunit was predicted to be membrane-bound. This suggested a periplasm location and that the complex may play a role in respiration of sulfur/polysulfide or thiosulfate. Selenite reductases (SrrA) also phylogenetically affiliate with the polysulfide/thiosulfate reductase clade, but conserved rhodonase-like proteins encoded in the gene neighbourhood of SrrA are thought to be indicative of selenite-reducing organisms [92], but were absent near *psrABC* in MAG AM4.

Ca. P. svalbardensis MAG AM4 also harboured a gene cluster encoding four subunits of a sulfhydrogenase complex (Supp. Table 4). Similar to the characterized sulfhydrogenase from *Pyrococcus furiosus*, this included two NiFe hydrogenase subunits, as well as two subunits of anaerobic sulfite reductases [93–95]. These complexes can use elemental sulfur or polysulfides as electron sinks when available [95], or act in reverse as hydrogen-evolving hydrogenases during fermentative growth [96].

Marine Acidobacteriota may respire additional electron acceptors including metals

All MAGs had gene clusters encoding multi-heme c-type cytochromes with predicted periplasmic or extracellular locations, as well as associated predicted β -barrel proteins (Supp. Table 4). In known metal-reducing and/or -oxidizing bacteria, extracellular and periplasmic cytochromes insert into outer-membrane traversing β -barrel proteins, and transfer electrons through the complexes to/from metals [97, 98]. These gene clusters were syntenous among the MAGs and *Thermoanaerobaculum aquaticum* (Supp. Fig. 7A), a related hot spring-derived isolate that can anaerobically reduce iron- and manganese-oxides [11]. We therefore propose these cytochromes are likely candidates for facilitating the reduction of metal-oxides by *Thermoanaerobaculum aquaticum*, because no other predicted extracellular cytochromes are encoded. We therefore also propose the similar cytochromes in our marine MAGs may also perform this function.

The *Ca. P. svalbardensis* MAG AM4 encoded two additional cytochrome c proteins with similarity to metal-reducing outer-membrane cytochromes (OmcS) from known metal-reducing bacteria, i.e., various *Desulfuromonadia* (formerly *Desulfuromonadales*) such as *Geobacter* and

Geopsychrobacter spp. (Supp. Table 5) [99, 100]. These cytochromes had six heme-binding sites like characterised OmcS, and were also clustered among genes for predicted periplasmic cytochromes and β -barrel proteins (Supp. Fig.7B). They could therefore also potentially exchange electrons with metal oxides (or other insoluble substrates such as humic-like substances, or other cells).

The marine Acidobacteriota MAGs also encoded the potential to reduce oxygen (Fig. 2), nitrous oxide (Supp. Fig. 8), organohalides (Supp. Fig. 9), nitrate, and arsenate (Supp. Fig. 6, Supp. Table 4, and further detailed in Supplementary information).

Additional energy conserving mechanisms among marine Acidobacteriota

Electron bifurcating heterodisulfide reductase complexes were only encoded in *Ca. Sulfomarinibacter* MAGs (Supp. Table 4). These complexes enable flavin-based redox balancing and formation of low-potential electron carriers (i.e., ferredoxin and/or flavodoxin), and are common among strict anaerobes [101, 102]. A high-molecular-weight cytochrome c3-type protein and a predicted periplasmic location was encoded in *Ca. Sulfomarinibacter* MAG AM1 (Supp. Table 4). These typically act as periplasmic redox hubs to link electron flows between the periplasm and cytoplasm in SRM [103]. All Acidobacteriota MAGs recovered in this study encoded NADH-ubiquinone oxidoreductase (Nuo) complexes required for energy conservation via respiration (Supp. Table 4). Genes for additional sodium-dependent Nuo complexes were also present (Supp. Table 4). Apart from the potential for respiration, some Acidobacteriota MAGs from both *Ca. Sulfomarinibacter* and *Ca. P. svalbardensis* MAG AM4 encoded acetate kinase and phosphate acetyltransferase for fermentation via acetogenesis, or which may act in reverse to facilitate acetate consumption (Supp. Table 4).

Marine Acidobacteriota use diverse nutrient and electron sources

The *Ca. Sulfomarinibacter* AM3 MAGs encoded predicted cellulase A enzymes with signal peptides for export from the cytoplasm (Fig. 2). They were phylogenetically affiliated with cellulase A from various anaerobic degraders of cellulose and/or plant-derived polysaccharides (Supp. Fig. 10). A cellobiose phosphorylase was encoded in *Ca. S. kjeldsenii* MAG AM3-C, and had relatively high amino acid identity (63%) to a characterized cellobiose phosphorylase from *Thermotoga neapolitana* [104]. These enzymes catalyse phosphorolysis of cellobiose to α -D-glucose 1-phosphate (G1P) and D-glucose, thereby saving an ATP before entering glycolysis, and are typically used by anaerobic cellulose-degraders [105]. This suggests these organisms have the capacity to anaerobically degrade

cellulose, a derivative of cellulose, or a structurally similar compound. Overall, the marine *Ca. Sulfomarinibacter* MAGs encoded few genes for glycoside hydrolases or other carbohydrate active enzymes, i.e., 0.47-0.75% of protein encoding genes encoded glycoside hydrolases (further detailed in Supplementary information) (Supp. Table 6). The *Ca. P. svalbardensis* MAG AM4 also encoded few glycoside hydrolases (0.54% of protein encoding genes), with none predicted to be exported to the extracellular environment, and a single endo-1,4-beta-xylanase predicted to be periplasmic (Supp. Table 4).

Genes for cyanophycinases among *Ca. Sulfomarinibacter* MAGs indicated they may utilize the storage compound cyanophycin as a nutrient (Supp. Table 4). The cyanophycinases had Secretion-signal peptides (Sec-) for export from the cytoplasm, indicating they act on an external substrate and not an internally stored compound. Accordingly, no genes for cyanophycin synthetases were found. An isoaspartyl dipeptidase was encoded in *Ca. S. kjeldsenii* MAG AM3-C, which may enable utilization of the products released by the cyanophycinase, i.e., a dipeptide of aspartate and arginine (Supp. Table 4). The capacity to catabolically degrade aspartate and arginine was also encoded (Supp. Table 4).

The *Ca. Sulfomarinibacter* MAG AM3-C may degrade extracellular proteins using two predicted secreted proteases, as well as adjacently encoded peptidases predicted to be membrane-bound (Supp. Table 4). The *Ca. P. svalbardensis* MAG AM4 harboured numerous genes for proteases/peptidases ($n=7$) that were predicted to be secreted, strongly indicating these bacteria use proteins as nutrients (Supp. Table 4).

Membrane-bound NiFe uptake-hydrogenases were encoded by both *Ca. Sulfomarinibacter* and *Ca. P. svalbardensis* MAGs (Supp. Table 4). These may be important for oxidizing environmental hydrogen. The *Ca. Sulfomarinibacter* MAGs encoded 'type-1c' NiFe hydrogenases typically found in obligate anaerobes and that are thought to be oxygen sensitive (Supp. Fig. 11) [106]. The *Ca. P. svalbardensis* MAG AM4 encoded a 'type-1d' NiFe hydrogenase, which are typically found in aerobes and facultative anaerobes (Supp. Fig. 11) [106]. Inspection of best BLASTP hits from the NCBI-nr database to the *Ca. Sulfomarinibacter* NiFe hydrogenase sequences identified various sequences previously shown to be expressed in tidal flat sediments [107]. Formate dehydrogenases encoded among MAGs of both *Ca. Sulfomarinibacter* and *Ca. P. svalbardensis* also suggested formate may be used as an electron donor (Supp. Table 4).

Adaptations to marine environments

Comparative genomics with seven *dsr*-harbouring Acidobacteriota MAGs from peatland soil [2] suggested the marine Acidobacteriota encoded unique adaptations to marine settings (Supplementary information) (Supp. Fig. 12). These included various predicted transporters/symporters and pumps for ions (e.g., sodium and potassium) and metals/metalloids (e.g., zinc and arsenic) that were unique to the marine MAGs. Genes for a sodium-translocating NADH-quinone oxidoreductase complex, which are used by various marine microorganisms to support respiration and cellular homeostasis [108], were only present in marine MAGs. Symporters for the osmolytes proline, glutamate and glycine, were also only present in marine MAGs.

Acidobacteriota are abundant, active and diverse in marine sediments

Amplicon sequencing of 16S rRNA genes revealed Acidobacteriota had an average relative abundance of $4.5 \pm 2.2\%$ in Smeerenbergfjorden sediments (Supp. Fig. 13 and 14), which have high sulfate reduction rates (reaching around $100 \text{ nmol SO}_4^{2-} \text{ cm}^{-3} \text{ d}^{-1}$ around 5 cmbsf) [38, 41, 109]. Thermoanaerobaculia-affiliated sequences were the most dominant of any Acidobacteriota, and reached the most abundant (11%) genus-level clade of *Bacteria* at 31 cmbsf in Station J (2016). The same clade was on average the fourth most abundant genus-level clade in the same core (averaged $4.5 \pm 2.8\%$). 16S rRNA transcripts of Acidobacteriota were below 0.5% relative abundances in the surface sediments (0–1 cmbsf) of Smeerenbergfjorden cores (Supp. Fig. 14). At station GK, Acidobacteriota 16S rRNA transcripts reached 6% relative abundance at 15 cmbsf (Supp. Fig. 14). We also examined Acidobacteriota 16S rRNA genes from metal-rich Van Keulenfjorden sediments from a previously published study [40]. This showed *Ca. Polarisedimenticola* related sequences were the most prominent Acidobacteriota, reaching 1.5%, and averaging $1.1 \pm 0.21\%$ of communities in four cores (Supp. Fig. 14). Members of the Thermoanaerobaculia were in much lower abundances ($0.3 \pm 0.2\%$ average overall), although they reached 1.1% in deeper sections of core AB. Mapping of metagenomic reads to the Acidobacteriota MAGs supported the general distribution trends from 16S rRNA amplicon analyses, i.e., that Thermoanaerobaculia were abundant in Smeerenbergfjorden sediments and *Ca. Polarisedimenticola* were more abundant in Van Keulenfjorden sediments (Supp. Table 7, and further detailed in Supplementary information).

Phylogenetic analysis of 16S rRNA genes from Smeerenbergfjorden sediment (Supp. Fig. 15) and examination of Acidobacteriota 16S rRNA sequences in the SILVA database (Supp. Fig. 16) revealed diverse Acidobacteriota sequences from marine sediments. It also revealed that Thermoanaerobaculia (sub-division 23) and Ca. Polarisedimenticola (sub-division 22) sequences are the most prominent Acidobacteriota lineages in marine sediments in general (further detailed in Supplementary information).

Sequencing of *dsrB* genes and transcripts from Smeerenburgfjord sediments revealed the Acidobacteriota *dsrB* averaged 13±6.6% of all *dsrB* (DNA-derived) sequences, and 4±2 % of *dsrB*-transcripts (cDNA-derived) (Supp. Fig. 17). Acidobacteriota *dsrB* sequences were the second most abundant group after Desulfobacterota *dsrB*, which dominated the sediments and averaged 75±6% in relative abundance (Supp. Fig. 17). Acidobacteriota *dsrB* reached a maximum of 19% at station GK and 31% at station J. The most abundant Acidobacteriota *dsrB*-OTU-17 was 100% identical (over 321 nucleotides) to *dsrB* from Ca. Sulfomarinibacter AM3-B MAG (Fig. 4). Amplicon-derived DsrB sequences that affiliated with the DsrB from marine Acidobacteriota MAGs were phylogenetically diverse and spread through-out the 'Thermoanaerobaculia Dsr clade' (Fig. 4).

Description of novel Acidobacteriota Candidatus taxa

Based on their unique phylogeny, predicted metabolic properties, CARD-FISH visualized cells of Thermoanaerobaculia (thin rods present in three different sites, see Supp. Fig. 18) and relatively complete MAGs, we propose the following new *Candidatus* taxa of Acidobacteriota (Supp. Table 3):

class Thermoanaerobaculia (sub-division 23)
 order Thermoanaerobaculales
 fam. nov. Sulfomarinibacteraceae (GTDB family FEB-10)
 gen. nov. Ca. Sulfomarinibacter
 sp. nov. Ca. Sulfomarinibacter kjeldsenii sp. nov. MAG AM3-C
 Ca. Sulfomarinibacter sp. MAG AM1
 Ca. Sulfomarinibacter sp. MAG AM2
 class nov. Ca. Polarisedimenticola (GTDB class Mor1, sub-division 22)
 ord nov. Ca. Polarisedimenticolales (GTDB order Mor1)
 fam. nov. Ca. Polarisedimenticolaceae (GTDB family Mor1)
 gen. nov. Ca. Polarisedimenticola
 sp. nov. Ca. Polarisedimenticola svalbardensis MAG AM4

447

448 Discussion

449 This study provides the first insights into the genomes and metabolic potential of abundant
450 *Thermoanaerobaculia* from marine sediments, and new insights into the metabolisms of *Ca.*
451 *Polarisedimenticola* (sub-division 22, or Mor1). Most notably, we revealed that MAGs from both of the
452 major lineages of Acidobacteriota from marine sediments have capabilities to dissimilate various
453 inorganic sulfur compounds.

454 Genes for the full dissimilatory sulfate reduction pathway provided the first direct link between
455 genomes of marine sediment Acidobacteriota and DsrAB sequences of the previously undescribed
456 'Uncultured family-level lineage 9' clade (here named 'Thermoanaerobaculia Dsr lineage'). In addition
457 to being abundant and actively transcribed in Svalbard sediments as shown here, *dsrB* sequences of
458 this lineage often constitute a prominent fraction of *dsrB*-harbouring communities in various sediments,
459 e.g., making around 8–14% of *dsrB* sequences from Aarhus Bay sediments [110, 111], around 5% of
460 *dsrB* sequences in Baltic Sea sediments [112], and up to 15-25% of sequences in sediments from
461 various cores from the Greenland coast [37]. Together, this indicates Acidobacteriota are a widespread
462 and prominent group of inorganic sulfur-dissimilating microorganisms in marine sediments.

463 While enzymes of the dissimilatory sulfate reduction pathway are widely used for anaerobic
464 reduction of sulfite/sulfate [35], some organisms can use them in reverse for the oxidation of reduced
465 sulfur compounds [113], or for disproportionation of sulfur compounds [114, 115]. Because no enzymes
466 are currently known that distinguish these different metabolisms, discerning sulfur metabolisms based
467 on genomic data requires careful interpretation [114, 115]. For instance, the *Ca. Sulfomarinibacter*
468 MAGs encoded DsrL, which was previously thought to be exclusively found in sulfur-oxidizing bacteria
469 [88]. However, recent work showed DsrL can function in a reductive manner in biochemical assays [88]
470 [136], and was highly expressed during reductive sulfur- and thiosulfate-respiration by *Desulfurella*
471 *amilsii* [88, 116]. The DsrL of *Ca. Sulfomarinibacter* contained putative NADP(H)-binding domain
472 structures that may enable coupling of NADPH as electron donor to sulfite reduction [136], as well as
473 phylogenetic relatedness with DsrL of *Desulfurella amilsii*. Together, this indicates the DsrL of *Ca.*
474 *Sulfomarinibacter* has potential to facilitate a reductive pathway.

The *Ca. Sulfomarinibacter* MAGs encoded rhodanase-like TusA and DsrE2, which act as sulfur-
trafficking proteins in reverse-Dsr harbouring sulfur-oxidizing bacteria, i.e., they help deliver sulfur to
DsrABC for oxidation [117]. Interestingly, the 'YTD gene clusters' that encode these enzymes are also
common in genomes of anaerobic elemental sulfur-reducing and/or -disproportionating bacteria that
have Dsr, and are suggested to be genetic indicators for disproportionation potential among these
anaerobes [91]. The TusA proteins from *Ca. Sulfomarinibacter* were most closely related to TusA from
various anaerobic sulfur-reducing and -disproportionating Desulfobacteriota (Supp. Fig. 6). This
suggested *Ca. Sulfomarinibacter* could reduce and/or disproportionate elemental sulfur, or possibly
other sulfur compounds that can be trafficked by TusA, like thiosulfate [118]. Indeed, the ability to
disproportionate sulfur compounds is common among sulfate-reducing Desulfobacteriota [119].
Elemental sulfur is often the most abundant sulfur cycle intermediate (SCI) in marine sediments [120],
and was measured in sediments from Smeerenbergfjorden up to 0.15 wt % of total sulfur [39]. Overall,
the gene content of *Ca. Sulfomarinibacter* MAGs indicated flexible dissimilatory sulfur metabolisms that
may be dictated by and/or switch under different biogeochemical and redox conditions.

Results indicated *Ca. Sulfomarinibacter* likely use the dissimilatory sulfate reduction pathway in
a reductive direction in most depths of the sediments studied. Firstly, Acidobacteriota were relatively
abundant and expressed *dsrB* in deeper (>15-75 cmbsf), strictly anoxic sediment layers of
Smeerenbergfjorden. These sediments lack electron acceptors that could sustain these abundant
populations growing via biological oxidation of sulfides, i.e., oxygen, nitrate or oxidized metals [31,
121]. In Station J sediments, oxygen and nitrate are depleted within millimetres-to-centimetres of the
surface [122, 123], and sulfide oxidation facilitated by Fe(III) is negligible [41]. An alternative possibility
is that cryptic biogeochemical cycling could sustain sulfide oxidation, i.e., fast consumption and
production of low concentrations of sulfides and oxidants [124]. Nevertheless, it remains unproven
whether biological sulfide oxidation occurs in deep sediments that lack measurable concentrations of
required oxidants [28]. On the other hand, the relative abundances of Acidobacteriota peaked in
subsurface zones around 5 cmbsf in Station J sediments, where sulfate reduction rates also peak [41,
109]. In another study, Acidobacteriota 16S rRNA gene relative abundances were also highly
correlated with sulfate reduction rates in sediments from Greenland [37]. These associations therefore
point toward an active role in the reduction and/or disproportionation of sulfur compounds of various
oxidation states by *Ca. Sulfomarinibacter* in marine sediments.

Our results also suggested marine Acidobacteriota have potential to reduce various inorganic sulfur compounds independent of the Dsr pathway. Our tetrathionate-amended microcosm experiment suggested that *Ca. Sulfomarinibacter* use tetrathionate as an electron sink via cytochromes, which supports the roles of these enzymes in tetrathionate reduction within *in situ*-like conditions. This is noteworthy because these enzymes were only previously shown to perform this function during biochemical assays [125], i.e., their utilization under *in situ*-like conditions was unknown. The ability to utilize SCI, e.g., tetrathionate or elemental sulfur/polysulfides/thiosulfate, could be important in sediment zones where SCI might be generated from sulfides reacting with available oxidants [41].

The *Ca. Sulfomarinibacter* MAGs indicated they could respire oxygen using terminal cbb3- or aa3-type cytochromes, although we speculate these may instead be used for defence against oxygen because they encoded many characteristics of obligate anaerobes (Supplementary discussion). We also hypothesize that the different redox metabolisms of the two predominant Acidobacteriota groups in Svalbard sediments, i.e., the *Ca. Sulfomarinibacter* and *Ca. Polarisodimenticola*, may explain their different abundances among fjords with different biogeochemical properties (Supplementary discussion). That is, the *Ca. Sulfomarinibacter* may be adapted to low redox environments, and are thus more abundant in the reduced (visibly black), sulfidic subsurface sediments of Smeerenburgfjorden. In comparison, the *Ca. P. svalbardensis* MAG had additional genes to utilize high-potential electron acceptors such as oxygen, nitrate and oxidized metals, and may be better adapted to the more high redox, metal-rich sediments of Van Keulenfjorden (visibly reddish-orange).

If members of the *Ca. Sulfomarinibacter* are indeed SRM, a question arises regarding how they co-exist with dominant sulfate-reducing Desulfobacterota populations, as both apparently use hydrogen, acetate or formate as substrates. However, we identified genes for use of several organic substrates that may enable *Ca. Sulfomarinibacter* to occupy a distinct nutrient niche. Complex carbohydrates such as cellulose (or structurally similar compounds) could be used. Carbohydrates are not used by most known isolated Desulfobacterota SRM [35]. Plant-derived molecules could stem from terrestrial run-off, which is a major source of organic carbon to arctic sediments [126, 127] and to coastal marine systems in general [128]. Additionally, various marine algae are known to produce cellulose [129]. The predicted ability to utilize cyanophycin could also facilitate a unique nutrient niche. Cyanophycin is a multi-L-arginyl-poly-L-aspartic acid, commonly produced by cyanobacteria as a

storage compound [130, 131]. Indeed, few organisms are known to use cyanophycin anaerobically [132], and no anaerobes are known from marine sediments.

Ca. Polarisedimenticola svalbardensis appeared to have a high propensity for the degradation of proteins, which was indicated by a suite of predicted secreted peptidases. A related *Mor1* *Acidobacteriota* genome (GCA_001664505.1) (Fig. 1) was recovered as a bacterial co-inhabitant of a cyanobacterial enrichment culture from seawater, suggesting it used organic material/necromass from the primary-producing cyanobacterium [133]. *Ca. Polarisedimenticola* may therefore contribute to protein degradation in marine sediments, where proteinaceous organics comprise a large proportion (~10%) of available organic matter [134].

In summary, the genome-encoded dissimilatory sulfur metabolisms and the high abundances and activity of *Ca. Sulfomarinibacter* in the sulfidic zones of Svalbard sediments, suggested these novel *Acidobacteriota* of the class *Thermoanaerobaculia* (sub-division 23) are important players in the biogeochemical sulfur cycles of the sediments. Our data also indicated that *Ca. Sulfomarinibacter* thrive largely via anaerobic metabolisms with the capability to use various other electron acceptors with different redox potentials, including biogeochemically relevant metal-oxides. Additionally, we show that *Ca. Polarisedimenticola svalbardensis*, a member of a different class of *Acidobacteriota* (sub-division 22), has the genetic potential for protein degradation and for metabolisms driven by high redox potential electron acceptors such as oxygen, nitrate and metal-oxides.

References

1. Kuske CR, Ticknor LO, Miller ME, Dunbar JM, Davis JA, Barns SM, et al. Comparison of Soil Bacterial Communities in Rhizospheres of Three Plant Species and the Interspaces in an Arid Grassland. *Applied and Environmental Microbiology* 2002. , **68**: 1854–1863
2. Hausmann B, Pelikan C, Herbold CW, Köstlbacher S, Albertsen M, Eichorst SA, et al. Peatland Acidobacteria with a dissimilatory sulfur metabolism. *ISME J* 2018; **12**: 1729–1742.
3. Kielak AM, Barreto CC, Kowalchuk GA, van Veen JA, Kuramae EE. The Ecology of Acidobacteria: Moving beyond Genes and Genomes. *Front Microbiol* 2016; **7**: 744.
4. Eichorst SA, Trojan D, Roux S, Herbold C, Rattei T, Woebken D. Genomic insights into the Acidobacteria reveal strategies for their success in terrestrial environments. *Environ Microbiol* 2018; **20**: 1041–1063.
5. LaPara TM, Nakatsu CH, Pantea L, Alleman JE. Phylogenetic analysis of bacterial communities

- in mesophilic and thermophilic bioreactors treating pharmaceutical wastewater. *Appl Environ Microbiol* 2000; **66**: 3951–3959.
6. Coskun ÖK, Özen V, Wankel SD, Orsi WD. Quantifying population-specific growth in benthic bacterial communities under low oxygen using H₂¹⁸O. *ISME J* 2019; **13**: 1546–1559.
7. Ward NL, Challacombe JF, Janssen PH, Henrissat B, Coutinho PM, Wu M, et al. Three genomes from the phylum Acidobacteria provide insight into the lifestyles of these microorganisms in soils. *Appl Environ Microbiol* 2009; **75**: 2046–2056.
8. Challacombe JF, Eichorst SA, Hauser L, Land M, Xie G, Kuske CR. Biological consequences of ancient gene acquisition and duplication in the large genome of *Candidatus Solibacter usitatus* Ellin6076. *PLoS One* 2011; **6**: e24882.
9. Coates JD, Ellis DJ, Gaw CV, Lovley DR. *Geothrix fermentans* gen. nov., sp. nov., a novel Fe(III)-reducing bacterium from a hydrocarbon-contaminated aquifer. *Int J Syst Evol Microbiol* 1999; **49**: 1615–1622.
10. Garcia Costas AM, Liu Z, Tomsho LP, Schuster SC, Ward DM, Bryant DA. Complete genome of *Candidatus Chloracidobacterium thermophilum*, a chlorophyll-based photoheterotroph belonging to the phylum Acidobacteria. *Environ Microbiol* 2012; **14**: 177–190.
11. Losey NA, Stevenson BS, Busse H-J, Sinninghe Damsté JS, Rijpstra WIC, Rudd S, et al. *Thermoanaerobaculum aquaticum* gen. nov., sp. nov., the first cultivated member of Acidobacteria subdivision 23, isolated from a hot spring. *Int J Syst Evol Microbiol* 2013; **63**: 4149–4157.
12. Woodcroft BJ, Singleton CM, Boyd JA, Evans PN, Emerson JB, Zayed AAF, et al. Genome-centric view of carbon processing in thawing permafrost. *Nature* 2018; **560**: 49–54.
13. Fukunaga Y, Kurahashi M, Yanagi K, Yokota A, Harayama S. *Acanthopleuribacter pedis* gen. nov., sp. nov., a marine bacterium isolated from a chiton, and description of Acanthopleuribacteraceae fam. nov., Acanthopleuribacterales ord. nov., Holophagaceae fam. nov., Holophagales ord. nov. and Holophagae classis nov. in the phylum ‘Acidobacteria’. *Int J Syst Evol Microbiol* 2008; **58**: 2597–2601
14. Yilmaz P, Yarza P, Rapp JZ, Glöckner FO. Expanding the World of Marine Bacterial and Archaeal Clades. *Front Microbiol* 2015; **6**: 1524.
15. Quaiser A, Zivanovic Y, Moreira D, López-García P. Comparative metagenomics of bathypelagic plankton and bottom sediment from the Sea of Marmara. *ISME J* 2011; **5**: 285–304.
16. O’Connor-Sánchez A, Rivera-Domínguez AJ, Santos-Briones C de L, López-Aguilar LK, Peña-Ramírez YJ, Prieto-Davo A. Acidobacteria appear to dominate the microbiome of two sympatric Caribbean Sponges and one Zoanthid. *Biol Res* 2014; **47**: 67.
17. Zhou Z, Liu Y, Xu W, Pan J, Luo Z-H, Li M. Genome- and Community-Level Interaction Insights

into Carbon Utilization and Element Cycling Functions of Hydrothermarchaeota in Hydrothermal Sediment. *mSystems* 2020; **5**

18. Polymenakou PN, Lampadariou N, Mandalakis M, Tselepides A. Phylogenetic diversity of sediment bacteria from the southern Cretan margin, Eastern Mediterranean Sea. *Syst Appl Microbiol* 2009; **32**: 17–26.
19. Kielak AM, van Veen JA, Kowalchuk GA. Comparative analysis of acidobacterial genomic fragments from terrestrial and aquatic metagenomic libraries, with emphasis on acidobacteria subdivision 6. *Appl Environ Microbiol* 2010; **76**: 6769–6777.
20. Orcutt BN, Sylvan JB, Knab NJ, Edwards KJ. Microbial ecology of the dark ocean above, at, and below the seafloor. *Microbiol Mol Biol Rev* 2011; **75**: 361–422.
21. Wang Y, Sheng H-F, He Y, Wu J-Y, Jiang Y-X, Tam NF-Y, et al. Comparison of the levels of bacterial diversity in freshwater, intertidal wetland, and marine sediments by using millions of illumina tags. *Appl Environ Microbiol* 2012; **78**: 8264–8271.
22. Choi H, Koh H-W, Kim H, Chae J-C, Park S-J. Microbial Community Composition in the Marine Sediments of Jeju Island: Next-Generation Sequencing Surveys. *J Microbiol Biotechnol* 2016; **26**: 883–890.
23. Conte A, Papale M, Amalfitano S, Mikkonen A, Rizzo C, De Domenico E, et al. Bacterial community structure along the subtidal sandy sediment belt of a high Arctic fjord (Kongsfjorden, Svalbard Islands). *Sci Total Environ* 2018; **619**: 203–211.
24. Kapili BJ, Barnett SE, Buckley DH, Dekas AE. Evidence for phylogenetically and catabolically diverse active diazotrophs in deep-sea sediment. *ISME J* 2020; **14**: 971–983
25. Tschoeke DA, Coutinho FH, Leomil L, Cavalcanti G, Silva BS, Garcia GD, et al. New bacterial and archaeal lineages discovered in organic rich sediments of a large tropical Bay. *Mar Genomics* 2020; 100789.
26. Bowles MW, Mogollón JM, Kasten S, Zabel M, Hinrichs K-U. Global rates of marine sulfate reduction and implications for sub-sea-floor metabolic activities. *Science* 2014; **344**: 889–891.
27. Parkes RJ, John Parkes R, Cragg BA, Wellsbury P. Recent studies on bacterial populations and processes in subseafloor sediments: A review. *Hydrogeology Journal* . 2000. , **8**: 11–28
28. Jørgensen BB, Findlay AJ, Pellerin A. The Biogeochemical Sulfur Cycle of Marine Sediments. *Front Microbiol* 2019; **10**.
29. Revsbech NP, Barker Jorgensen B, Blackburn TH. Oxygen in the Sea Bottom Measured with a Microelectrode. *Science* 1980; **207**: 1355.
30. Anantharaman K, Hausmann B, Jungbluth SP, Kantor RS, Lavy A, Warren LA, et al. Expanded diversity of microbial groups that shape the dissimilatory sulfur cycle. *ISME J* 2018; **12**: 1715–1728.
31. Wasmund K, Mußmann M, Loy A. The life sulfuric: microbial ecology of sulfur cycling in marine

sediments. *Environ Microbiol Rep* 2017; **9**: 323–344.

32. Jørgensen BB. Mineralization of organic matter in the sea bed—the role of sulphate reduction. *Nature* 1982; **296**: 643–645.

33. Müller AL, Pelikan C, de Rezende JR, Wasmund K, Putz M, Glombitza C, et al. Bacterial interactions during sequential degradation of cyanobacterial necromass in a sulfidic arctic marine sediment. *Environ Microbiol* 2018; **20**: 2927–2940.

34. Finke N, Vandieken V, Jørgensen BB. Acetate, lactate, propionate, and isobutyrate as electron donors for iron and sulfate reduction in Arctic marine sediments, Svalbard. *FEMS Microbiol Ecol* 2007; **59**: 10–22

35. Rabus R, Venceslau SS, Wöhlbrand L, Voordouw G, Wall JD, Pereira IAC. A Post-Genomic View of the Ecophysiology, Catabolism and Biotechnological Relevance of Sulphate-Reducing Prokaryotes. *Adv Microb Physiol* 2015; **66**: 55–321.

36. Müller AL, Kjeldsen KU, Rattei T, Pester M, Loy A. Phylogenetic and environmental diversity of DsrAB-type dissimilatory (bi)sulfite reductases. *ISME J* 2015; **9**: 1152–1165.

37. Pelikan C, Jaussi M, Wasmund K, Seidenkrantz M-S, Pearce C, Kuzyk ZZA, et al. Glacial Runoff Promotes Deep Burial of Sulfur Cycling-Associated Microorganisms in Marine Sediments. *Front Microbiol* 2019; **10**: 2558.

38. Wehrmann LM, Formolo MJ, Owens JD, Raiswell R, Ferdeman TG, Riedinger N, et al. Iron and manganese speciation and cycling in glacially influenced high-latitude fjord sediments (West Spitsbergen, Svalbard): Evidence for a benthic recycling-transport mechanism. *Geochim Cosmochim Acta* 2014; **141**: 628–655.

39. Wehrmann LM, Riedinger N, Brunner B, Kamysny A, Hubert CRJ, Herbert LC, et al. Iron-controlled oxidative sulfur cycling recorded in the distribution and isotopic composition of sulfur species in glacially influenced fjord sediments of west Svalbard. *Chem Geol* 2017; **466**: 678–695.

40. Buongiorno J, Herbert LC, Wehrmann LM, Michaud AB, Laufer K, Røy H, et al. Complex Microbial Communities Drive Iron and Sulfur Cycling in Arctic Fjord Sediments. *Appl Environ Microbiol* 2019; **85**.

41. Michaud AB, Laufer K, Findlay A, Pellerin A, Antler G, Turchyn AV, et al. Glacial influence on the iron and sulfur cycles in Arctic fjord sediments (Svalbard). *Geochim Cosmochim Acta* 2020.

42. Jørgensen BB, Laufer K, Michaud AB, Wehrmann LM. Biogeochemistry and microbiology of high Arctic marine sediment ecosystems—Case study of Svalbard fjords. *Limnol Oceanogr* 2020; **30**: 85.

43. Buongiorno J, Sipes K, Wasmund K, Loy A, Lloyd KG. Woeseiales transcriptional response to shallow burial in Arctic fjord surface sediment. *PLoS One* 2020; **15**: e0234839

44. Kannevorff E, Nicolaisen W. The ‘Haps’ a frame-supported bottom corer. *Ophelia* 1972; **10**:

- 119–128.
45. Meischner D, Others. A light-weight, high-momentum gravity corer for subaqueous sediments. 1974.
46. Peng Y, Leung HCM, Yiu SM, Chin FYL. IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* 2012; **28**: 1420–1428
47. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile metagenomic assembler. *Genome Res* 2017; **27**: 824–834.
48. Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, et al. KBase: The United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol* 2018; **36**: 566–569.
49. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 2015; **31**: 1674–1676
50. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 2009; **25**: 1754–1760.
51. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 2009; **25**: 2078–2079.
52. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ* 2019; **7**: e7359.
53. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, et al. Binning metagenomic contigs by coverage and composition. *Nat Methods* 2014; **11**: 1144–1146.
54. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nature Microbiology* 2018; **3**: 836–843
55. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic comparisons that enables improved genome recovery from metagenomes through de-replication. *ISME J* 2017; **11**: 2864–2868.
56. Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, et al. Thousands of microbial genomes shed light on interconnected biogeochemical processes in an aquifer system. *Nature Communications* 2016; **7**
57. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 2015; **25**: 1043–1055.
58. Bushnell B, Rood J, Singer E. BBMerge--accurate paired shotgun read merging via overlap. *PLoS One* 2017; **12**.

59. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* 2019.
60. Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J. JSpeciesWS: a web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 2016; **32**: 929–931.
61. Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavrommatis K, Kyrpides NC, et al. Microbial species delineation using whole genome sequences. *Nucleic Acids Res* 2015; **43**: 6761–6771.
62. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 2008; **9**: 75.
63. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997; **25**: 3389–3402.
64. Boeckmann B, Bairoch A, Apweiler R, Blatter M-C, Estreicher A, Gasteiger E, et al. The SWISS-PROT protein knowledgebase and its supplement TrEMBL in 2003. *Nucleic Acids Res* 2003; **31**: 365–370.
65. Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, et al. CDD/SPARCLE: functional classification of proteins via subfamily domain architectures. *Nucleic Acids Res* 2017; **45**: D200–D203.
66. Caspi R, Altman T, Billington R, Dreher K, Foerster H, Fulcher CA, et al. The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of Pathway/Genome Databases. *Nucleic Acids Res* 2014; **42**: D459–71.
67. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environ Microbiol* 2016; **18**: 1403–1414.
68. Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquat Microb Ecol* 2015; **75**: 129–137.
69. Herbold CW, Pelikan C, Kuzyk O, Hausmann B, Angel R, Berry D, et al. A flexible and economical barcoding approach for highly multiplexed amplicon sequencing of diverse target genes. *Front Microbiol* 2015; **6**: 731.
70. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc Natl Acad Sci USA* 2011; **108**: 4516–4522.
71. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* 2016; **13**: 581–583.

72. Callahan BJ, Sankaran K, Fukuyama JA, McMurdie PJ, Holmes SP. Bioconductor Workflow for Microbiome Data Analysis: from raw reads to community analyses. *F1000Res* 2016; **5**: 1492.
73. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 2009; **75**: 7537–7541.
74. Lagkouvardos I, Fischer S, Kumar N, Clavel T. Rhea: a transparent and modular R pipeline for microbial profiling based on 16S rRNA gene amplicons. *PeerJ* 2017; **5**: e2836
75. Pelikan C, Herbold CW, Hausmann B, Müller AL, Pester M, Loy A. Diversity analysis of sulfite- and sulfate-reducing microorganisms by multiplex dsrA and dsrB amplicon sequencing using new primers and mock community-optimized bioinformatics. *Environ Microbiol* 2016; **18**: 2994–3009.
76. Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res* 2016; **44**: W232–5.
77. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res* 2016; **44**: W242–5.
78. Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadukumar, et al. ARB: a software environment for sequence data. *Nucleic Acids Res* 2004; **32**: 1363–1371.
79. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* 2013; **41**: D590–6.
80. Duval S, Ducluzeau A-L, Nitschke W, Schoepp-Cothenet B. Enzyme phylogenies as markers for the oxidation state of the environment: the case of respiratory arsenate reductase and related enzymes. *BMC Evol Biol* 2008; **8**: 206.
81. Katoh K, Misawa K, Kuma K-I, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 2002; **30**: 3059–3066.
82. Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* 2004; **5**: 113.
83. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* 2013; **30**: 2725–2729.
84. Wendeberg A. Fluorescence in situ hybridization for the identification of environmental microbes. *Cold Spring Harb Protoc* 2010; **2010**: db.prot5366.
85. Juretschko S, Loy A, Lehner A, Wagner M. The microbial community composition of a nitrifying-denitrifying activated sludge from an industrial sewage treatment plant analyzed by the full-cycle rRNA approach. *Syst Appl Microbiol* 2002; **25**: 84–99.
86. Barco RA, Garrity GM, Scott JJ, Amend JP, Nealson KH, Emerson D. A Genus Definition for Bacteria and Archaea Based on a Standard Genome Relatedness Index. *MBio* 2020; **11**.

87. Santos AA, Venceslau SS, Grein F, Leavitt WD, Dahl C, Johnston DT, et al. A protein trisulfide couples dissimilatory sulfate reduction to energy conservation. *Science* 2015; **350**: 1541–1545.
88. Löffler M, Feldhues J, Venceslau SS, Kammler L, Grein F, Pereira IAC, et al. DsrL mediates electron transfer between NADH and rDsrAB in *Allochromatium vinosum*. *Environ Microbiol* 2019.
89. Mizuno N, Voordouw G, Miki K, Sarai A, Higuchi Y. Crystal Structure of Dissimilatory Sulfite Reductase D (DsrD) Protein—Possible Interaction with B- and Z-DNA by Its Winged-Helix Motif. *Structure* 2003; **11**: 1133–1140.
90. Mowat CG, Rothery E, Miles CS, McIver L, Doherty MK, Drewette K, et al. Octaheme tetrathionate reductase is a respiratory enzyme with novel heme ligation. *Nat Struct Mol Biol* 2004; **11**: 1023–1024.
91. Umezawa K, Kojima H, Kato Y, Fukui M. Disproportionation of inorganic sulfur compounds by a novel autotrophic bacterium belonging to Nitrospirota. *Syst Appl Microbiol* 2020; **43**: 126110.
92. Wells M, McGarry J, Gaye MM, Basu P, Oremland RS, Stolz JF. Respiratory Selenite Reductase from *Bacillus selenitireducens* Strain MLS10. *J Bacteriol* 2019; **201**.
93. Ma K, Schicho RN, Kelly RM, Adams MW. Hydrogenase of the hyperthermophile *Pyrococcus furiosus* is an elemental sulfur reductase or sulfhydrogenase: evidence for a sulfur-reducing hydrogenase ancestor. *Proc Natl Acad Sci USA* 1993; **90**: 5341–5344.
94. Bryant FO, Adams MW. Characterization of hydrogenase from the hyperthermophilic archaeobacterium, *Pyrococcus furiosus*. *J Biol Chem* 1989; **264**: 5070–5079.
95. Ma K, Weiss R, Adams MW. Characterization of hydrogenase II from the hyperthermophilic archaeon *Pyrococcus furiosus* and assessment of its role in sulfur reduction. *J Bacteriol* 2000; **182**: 1864–1871.
96. van Haaster DJ, Silva PJ, Hagedoorn P-L, Jongejan JA, Hagen WR. Reinvestigation of the steady-state kinetics and physiological function of the soluble NiFe-hydrogenase I of *Pyrococcus furiosus*. *J Bacteriol* 2008; **190**: 1584–1587.
97. Edwards MJ, White GF, Lockwood CW, Lawes MC, Martel A, Harris G, et al. Structural modeling of an outer membrane electron conduit from a metal-reducing bacterium suggests electron transfer via periplasmic redox partners. *J Biol Chem* 2018; **293**: 8103–8112.
98. Edwards MJ, White GF, Butt JN, Richardson DJ, Clarke TA. The Crystal Structure of a Biological Insulated Transmembrane Molecular Wire. *Cell* 2020.
99. Mehta T, Coppi MV, Childers SE, Lovley DR. Outer membrane c-type cytochromes required for Fe(III) and Mn(IV) oxide reduction in *Geobacter sulfurreducens*. *Appl Environ Microbiol* 2005; **71**: 8634–8641.
100. Tang H-Y, Holmes DE, Ueki T, Palacios PA, Lovley DR. Iron Corrosion via Direct Metal-Microbe Electron Transfer. *MBio* 2019; **10**.

101. Yan Z, Wang M, Ferry JG. A Ferredoxin- and F420H₂-Dependent, Electron-Bifurcating, Heterodisulfide Reductase with Homologs in the Domains Bacteria and Archaea. *MBio* 2017; **8**.
102. Buckel W, Thauer RK. Flavin-Based Electron Bifurcation, A New Mechanism of Biological Energy Coupling. *Chem Rev* 2018; **118**: 3862–3886.
103. Pereira IAC, Ramos AR, Grein F, Marques MC, da Silva SM, Venceslau SS. A comparative genomic analysis of energy metabolism in sulfate reducing bacteria and archaea. *Front Microbiol* 2011; **2**: 69.
104. Yernool DA, McCarthy JK, Eveleigh DE, Bok JD. Cloning and characterization of the glucooligosaccharide catabolic pathway beta-glucan glucohydrolase and cellobiose phosphorylase in the marine hyperthermophile *Thermotoga neapolitana*. *J Bacteriol* 2000; **182**: 5172–5179.
105. Kim SK, Himmel ME, Bomble YJ. Expression of a cellobiose phosphorylase from *Thermotoga maritima* in *Caldicellulosiruptor bescii* improves the phosphorolytic pathway and results in a dramatic increase in cellulolytic activity. *Appl Environ Microbiol* 2018; **84**:e02348-17
106. Søndergaard D, Pedersen CNS, Greening C. HydDB: A web tool for hydrogenase classification and analysis. *Sci Rep* 2016; **6**: 34212.
107. Dykstra S, Pjevac P, Ovanesov K, Musmann M. Evidence for H₂ consumption by uncultured Desulfobacterales in coastal sediments. *Environ Microbiol* 2018; **20**: 450–461.
108. Verkhovsky MI, Bogachev AV. Sodium-translocating NADH:quinone oxidoreductase as a redox-driven ion pump. *Biochim Biophys Acta* 2010; **1797**: 738–746.
109. Vandieken V, Finke N, Jørgensen BB. Pathways of carbon oxidation in an Arctic fjord sediment (Svalbard) and isolation of psychrophilic and psychrotolerant Fe(III)-reducing bacteria. *Mar Ecol Prog Ser* 2006; **322**: 29–41.
110. Jochum LM, Chen X, Lever MA, Loy A, Jørgensen BB, Schramm A, et al. Depth Distribution and Assembly of Sulfate-Reducing Microbial Communities in Marine Sediments of Aarhus Bay. *Appl Environ Microbiol* 2017; **83**.
111. Petro C, Zäncker B, Starnawski P, Jochum LM, Ferdelman TG, Jørgensen BB, et al. Marine Deep Biosphere Microbial Communities Assemble in Near-Surface Sediments in Aarhus Bay. *Front Microbiol* 2019; **10**: 758.
112. Marshall IPG, Ren G, Jaussi M, Lomstein BA, Jørgensen BB, Røy H, et al. Environmental filtering determines family-level structure of sulfate-reducing microbial communities in subsurface marine sediments. *ISME J* 2019; **13**: 1920–1932.
113. Loy A, Duller S, Baranyi C, Musmann M, Ott J, Sharon I, et al. Reverse dissimilatory sulfite reductase as phylogenetic marker for a subgroup of sulfur-oxidizing prokaryotes. *Environ Microbiol* 2009; **11**: 289–299.
114. Thorup C, Schramm A, Findlay AJ, Finster KW, Schreiber L. Disguised as a Sulfate Reducer:

- Growth of the Deltaproteobacterium *Desulfurivibrio alkaliphilus* by Sulfide Oxidation with Nitrate. *mBio* 2017; **8**
115. Kjeldsen KU, Schreiber L, Thorup CA, Boesen T, Bjerg JT, Yang T, et al. On the evolution and physiology of cable bacteria. *Proc Natl Acad Sci USA* 2019; **116**: 19116–19125.
116. Florentino AP, Pereira IAC, Boeren S, van den Born M, Stams AJM, Sánchez-Andrea I. Insight into the sulfur metabolism of *Desulfurella amilsii* by differential proteomics. *Environ Microbiol* 2019; **21**: 209–225.
117. Dahl C. Cytoplasmic sulfur trafficking in sulfur-oxidizing prokaryotes. *IUBMB Life* 2015; **67**: 268–274.
118. Liu L-J, Stockdreher Y, Koch T, Sun S-T, Fan Z, Josten M, et al. Thiosulfate transfer mediated by DsrE/TusA homologs from acidothermophilic sulfur-oxidizing archaeon *Metallosphaera cuprina*. *J Biol Chem* 2014; **289**: 26949–26959.
119. Slobodkin AI, Slobodkina GB. Diversity of Sulfur-Disproportionating Microorganisms. *Microbiology* 2019; **88**: 509–522.
120. Zopfi J, Ferdelman TG, Fossing H. Distribution and fate of sulfur intermediates-sulfite, tetrathionate, thiosulfate, and elemental sulfur-in marine sediments. *Special Papers-Geological Society of America* 2004; 97–116.
121. Henkel JV, Dellwig O, Pollehne F, Herlemann DPR, Leipe T, Schulz-Vogt HN. A bacterial isolate from the Black Sea oxidizes sulfide with manganese(IV) oxide. *Proc Natl Acad Sci USA* 2019; **116**: 12153–12155.
122. Jørgensen BB, Dunker R, Grünke S, Røy H. Filamentous sulfur bacteria, *Beggiatoa* spp., in arctic marine sediments (Svalbard, 79°N). *FEMS Microbiol Ecol* 2010; **73**: 500–513.
123. Canion A, Overholt WA, Kostka JE, Huettel M, Lavik G, Kuypers MMM. Temperature response of denitrification and anaerobic ammonium oxidation rates and microbial community structure in Arctic fjord sediments. *Environ Microbiol* 2014; **16**: 3331–3344.
124. Kappler A, Bryce C. Cryptic biogeochemical cycles: unravelling hidden redox reactions. *Environ Microbiol* 2017; **19**: 842–846
125. Buckley A, MacGregor B, Teske A. Identification, Expression and Activity of Candidate Nitrite Reductases From Orange Beggiatoaceae, Guaymas Basin. *Front Microbiol* 2019; **10**: 644.
126. Benner R, Benitez-Nelson B, Kaiser K, Amon RMW. Export of young terrigenous dissolved organic carbon from rivers to the Arctic Ocean. *Geophys Res Lett* 2004; **31**.
127. Opsahl S, Benner R, Amon RMW. Major flux of terrigenous dissolved organic matter through the Arctic Ocean. *Limnol Oceanogr* 1999; **44**: 2017–2023.
128. Burdige DJ. Burial of terrestrial organic matter in marine sediments: A re-assessment. *Global Biogeochem Cycles* 2005; **19**.

129. Lakshmi DS, Trivedi N, Reddy CRK. Synthesis and characterization of seaweed cellulose derived carboxymethyl cellulose. *Carbohydr Polym* 2017; **157**: 1604–1610.
130. Simon RD. Cyanophycin Granules from the Blue-Green Alga *Anabaena cylindrica*: A Reserve Material Consisting of Copolymers of Aspartic Acid and Arginine. *Proc Natl Acad Sci USA* 1971; **68**: 265–267.
131. Simon RD, Weathers P. Determination of the structure of the novel polypeptide containing aspartic acid and arginine which is found in Cyanobacteria. *Biochim Biophys Acta* 1976; **420**: 165–176.
132. Obst M, Krug A, Luftmann H, Steinbüchel A. Degradation of cyanophycin by *Sedimentibacter hongkongensis* strain KI and *Citrobacter amalonaticus* strain G Isolated from an anaerobic bacterial consortium. *Appl Environ Microbiol* 2005; **71**: 3642–3652.
133. Cummings SL, Barbé D, Leao TF, Korobeynikov A, Engene N, Glukhov E, et al. A novel uncultured heterotrophic bacterial associate of the cyanobacterium *Moorea producens* JHB. *BMC Microbiol* 2016; **16**: 198.
134. Burdige DJ. Preservation of organic matter in marine sediments: controls, mechanisms, and an imbalance in sediment organic carbon budgets? *Chem Rev* 2007; **107**: 467–485.
135. Rasigraf O, Helmond NAGM, Frank J, Lenstra WK, Egger M, Slomp CP, et al. Microbial community composition and functional potential in Bothnian Sea sediments is linked to Fe and S dynamics and the quality of organic matter. *Limnol Oceanogr* 2020; **65**: e00169.
136. Löffler, Wallerang and Dahl. The Iron-Sulfur Flavoprotein DsrL as NAD(P)H:acceptor Oxidoreductase in Oxidative and Reductive Dissimilatory Sulfur Metabolism. *Front Microbiol* 2020; in press. doi: 10.3389/fmicb.2020.578209.

Acknowledgements

This research was supported by the Austrian Science Fund (FWF grants P29426 to KW and P25111-B22 to AL) and the MetaBac Research Platform of the University of Vienna. We thank Captain Stig Henningsen of MS Farm during Svalbard expeditions. We particularly thank Bo Barker Jørgensen, Alexander Michaud and Susann Henkel for organising the 2016 and 2017 Svalbard expeditions, and all members of Svalbard expeditions for help with sample collection, especially Claus Pelikan. We thank the Biomedical Sequencing Facility (BSF) Vienna for sequencing of metagenome samples, the Joint Microbiome Facility (JMF) of the Medical University of Vienna and the University of Vienna, and Microsynth for sequencing amplicons. We specifically thank Jasmin Schwarz, Gudrun Kohl and Petra Pjevac from the JMF for assisting with amplicon sequencing. We thank Marc Mussman and Stefan

920 Dyksma for providing the hydrogenase database. We are grateful to Bernhard Schink for help with
921 Latin naming of taxa.

922 Contributions

923 KW and AL conceived the study. KW, JB, KGL and AL collected samples. MF and KW performed
924 microcosm experiments. MF performed DNA/RNA extractions and PCRs for amplicon sequencing. KW,
925 MF and JB performed bioinformatic analyses of metagenomic and amplicon data, with support from
926 CH, BH and TR. MF and BH analysed amplicon sequencing data. MF designed and performed RT-
927 qPCR experiments. KW and MF interpreted genomic data. MF and KW performed CARD-FISH. MF,
928 KW and AL wrote the manuscript, with contributions from all authors.

929 Conflict of interest

930 The authors declare no conflicts of interest.

931 Correspondence

932 Correspondence to Kenneth Wasmund or Alexander Loy.

933 Figure captions

934 **Figure 1. Phylogenomic analysis reveals novel Acidobacteriota taxa in marine sediments.** Maximum-
935 likelihood tree of concatenated protein sequences from MAGs and genomes. Single marker genes
936 were retrieved with CheckM. Highlighted in blue are MAGs obtained in this study. Highlighted in purple
937 are *dsrAB*-containing MAGs obtained from the NCBI database from the class Thermoanaerobaculia.
938 The genus *Ca. Acidiflorens* is represented by the most complete MAG (GCA_003166525.1) from the
939 corresponding study [12]. Our phylogenomic analysis showed that one MAG that was previously
940 assigned to *Ca. Aminicenantes* (GCA_004524955.1), recovered from the Bothnian Sea [135], is
941 affiliated with the newly proposed family *Ca. Sulfomarinibacteraceae*. Black dots indicate *dsrAB*-
942 containing genomes/MAGs. Bootstrap values with >90% are indicated with filled black circles on nodes.
943 *Nitrospina gracilis* 3/211 (GCA_000341545.2) was used as an outgroup. The scale bar represents 10%
944 sequence divergence.

945 **Figure 2. Metabolic models of (A) *Ca. Sulfomarinibacter kjeldsenii* MAG AM3-C and (B) *Ca.***
946 ***Polarisedimenticola svalbardensis* MAG AM4 suggest different fundamental niches of the two species**
947 **in marine sediments.** GH = glycoside hydrolase, RDH = reductive dehalogenase homologous enzyme,
948 Ack = acetate kinase, Pta = phosphotransacetylase, PFL = pyruvate-formate lyase, FDH = formate
949 dehydrogenase, Hdr = heterodisulfide reductase, NUO = NADH dehydrogenase, Otr = Tetrathionate
950 reductase, NosZ = nitrous oxide reductase, Sat = sulfate adenylyltransferase, Apr = adenylylsulfate
951 reductase, Qmo = quinone-interacting membrane oxidoreductase complex, Dsr = dissimilatory sulfate
952 reductase, Nap = Periplasmic nitrate reductase, Psr = polysulfide reductase, Sdh = Sulfhydrogenase
953 complex, TusA = sulfur carrier protein.

Figure 3. Gene organization of the *dsr* gene cluster in Acidobacteriota. Scaffold names in blue were retrieved from this study. Scaffold names in purple were derived from best BLASTP hits to sequences from this study. *Ca. Sulfotellmatomonas gaucii* SbA5 was retrieved from Hausmann *et al.* 2018. Green: *dsr*, dark red: other genes, and orange: hypothetical genes. Shaded blue lines indicate degree of sequence similarity as determined by tBLASTx within EasyFig.

Figure 4. DsrAB uncultured lineage 9 in the DsrAB tree represents members of the Acidobacteriota class Thermoanaerobaculia (sub-division 23). Blue leaves in the DsrAB tree represent MAGs or contigs identified in this study. Red leaves represent the most abundant acidobacteriotal amplicon-derived DsrB sequences identified in this study. Purple leaves represent sequences from MAGs retrieved from public databases. The DsrAB sequences were added to the consensus tree from Müller *et al.* 2015 in ARB. SD, pertaining to 'sub-divisions' of Acidobacteriota. The scale bar represents 10% sequence divergence.

Figure 5. Box plots depicting the expression of *otr* and *dsrB* relative to a house-keeping gene (DNA-directed RNA polymerase, alpha subunit) from *Ca. Sulfomarinibacter kjeldsenii* MAG AM3-C during microcosm experiments with amendments of tetrathionate versus no-amendment controls. Relative expression was determined by rt-qPCR. Expression of *otr* was significantly higher at day 8 ($p=0.0488$) as determined using a two-tailed T-test, and is indicated by an asterisk. Center lines indicate medians; box limits indicate 25th and 75th percentiles as determined by R software; and whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles.

Supplementary Figure 1. A) Outline of the metagenomic binning strategy. B) Plot of completeness of MAGs (CheckM). Comparisons are derived from binning from single assemblies (IDBA or MetaSpades or Megahit), versus binning from multiple assemblies of each sample (outlined in panel A).

Supplementary Figure 2. Alignment of dissimilatory DsrC cysteine motifs. Sub-section (C-terminus) of alignment of DsrC proteins, showing two conserved cysteine residues (dark purple) that are present in dissimilatory versions of the enzymes.

Supplementary Figure 3. A) Phylogenetic tree of DsrL proteins. The sequences from MAGs recovered in this study are highlighted in blue. Other Acidobacteriota DsrL are highlighted in purple. Bootstrap values >50% are presented on nodes as black-filled circles. The scale bar represents 20% sequence divergence. **B) Alignment of DsrL proteins.** A subsection of the whole DsrL alignment is shown to highlight YRR amino acids for putative NAD(P)H-binding domain.

Supplementary Figure 4. Phylogenetic tree of multiheme cytochrome protein sequences. Sequences from MAGs recovered in this study are highlighted in blue. Sequences from other Acidobacteriota are highlighted in purple. Reference sequences were retrieved from Kern *et al.*, 2011, and from best BLASTP hits to our MAG-derived sequences. Functional assignments are labelled at the end of each leaf label. NrfA = respiratory cytochrome c nitrite reductase, Onr = octaheme cytochrome c nitrite reductase, Hao/Hzo = octaheme hydroxylamine oxidoreductase/hydrazine oxidoreductase, MccA = cytochrome c sulfite reductase, and Otr = octaheme tetrathionate reductase. Genbank accessions are presented in parentheses. The scale bar represents 50% sequence divergence.

Supplementary Figure 5. A) Phylogenetic tree of TusA proteins. The sequences from MAGs recovered in this study are highlighted in blue. The orange branch indicates TusA proteins from anaerobic organisms known to reduce or disproportionate sulfur cycle intermediates and that had TusA related to the Acidobacteriota TusA. Descriptions of sulfur metabolisms related to reduction or disproportionation of sulfur cycle intermediates are presented in parenthesis for TusA related to TusA from MAGs recovered in this study. Bootstrap values >50% are presented on nodes as black-filled circles. The scale bar represents 20% sequence divergence. **B) Alignment of TusA proteins** from marine Acidobacteriota showing Cys-Pro-X-Pro sulfane sulfur-binding domains.

Supplementary Figure 6. Phylogenetic tree of complex iron-sulfur molybdoenzyme (CISM) family proteins. The sequences from the MAGs recovered in this study are highlighted in blue. Reference sequences were obtained from Duval *et al.*, 2008, as well as selected additional sequences. Bootstrap values >90% are presented on nodes as black-filled circles. The scale bar represents 50% sequence divergence.

Supplementary Figure 7. A) Schematic of gene organisation and synteny of extracellular cytochrome-rich genomic loci among Acidobacteriota MAGs (AM1, AM3-C and AM4) and *Thermoanaerobaculum*

aquaticum (T. aq.). B) Schematic of gene organisation of genomic loci encoding OmcS-like proteins in MAG AM4. Shaded blue lines indicate degree of sequence similarity as determined by tblastx within EasyFig (Sullivan et al., 2011). Subcellular location predictions and number of heme-binding sites (CXXCH) are indicated in parentheses. SEC-peptides for Sec secretion systems were searched in proteins with 'unknown' location predictions using PRED-TAT (Bagos et al., 2011).

Supplementary Figure 8. Phylogenetic tree of nitrous oxide reductases (NosZ). Sequences from MAGs recovered in this study are highlighted in blue. The NosZ from MAG AM1 was omitted due to short sequence length, although it was most similar to the NosZ from AM3-B and AM3-C (>90% amino acid identity from 190 amino acids). Sequences from other Acidobacteriota are highlighted in purple. Clade of 'type I NosZ' = blue, and clade of 'type II NosZ' = red. Reference sequences were retrieved from the top 50 best BLASTP hits to the NosZ from MAG AM3-C were included. Genbank accessions are presented in parentheses. Black circles on nodes represent bootstraps values >90%. The scale bar represents 20% sequence divergence.

Supplementary Figure 9. Phylogenetic tree of reductive dehalogenase homolog A (RdhA) proteins. The sequence from the MAG recovered in this study are highlighted in blue. Reference sequences were obtained from Hug et al., 2013, and the top 10 best BLASTP hits to the RdhA from MAG AM3-C were also included. The RdhA sequence of MAG AM1 was not included due to the truncated protein sequence, although it was most similar to the RdhA from MAG AM3-C (>87% amino acid identity from 120 amino acids). The scale bar represents 50% sequence divergence.

Supplementary Figure 10. Phylogenetic tree of cellulase A-like proteins. Sequences from MAGs recovered in this study are highlighted in blue. Sequences from other Acidobacteriota are highlighted in purple. Sequences from genera or species known to perform cellulose degradation are highlighted in green. Reference sequences were retrieved from the top 50 best BLASTP hits to the cellulase A from MAG AM3-C. Genbank accessions are presented in parentheses. The tree was rooted with the cellulase A of *Bacillus subtilis*. Black circles on nodes represent bootstraps values >90%. The scale bar represents 20% sequence divergence.

Supplementary Figure 11. Phylogenetic tree of [NiFe]-hydrogenase large subunit proteins. The sequences from the MAGs recovered in this study are highlighted in dark blue. Sequences from other Acidobacteriota are highlighted in purple. Sequences from PCR-derived amplicons from tidal flat sediments (Dyksma et al., 2018) are highlighted in light blue. Reference sequences were derived from best BLASTP hits from NCBI-nr database. Hydrogenase 'types' were determined using HydDB (Søndergaard et al., 2016). Black circles on nodes represent bootstraps values >90%. The scale bar represents 20% sequence divergence.

Supplementary Figure 12. Comparisons of COG classifications of proteins representing unique ortholog groups (OGs) from marine versus terrestrial dsr-harboursing Acidobacteriota. OGs unique to each group of genomes were determined using OrthoFinder (Emms and Kelly 2019). Proteins were compared from the six MAGs recovered in this study, versus proteins from the seven MAGs recovered by Hausmann et al., 2018. Letters in parenthesis represent standard COG codes.

Supplementary Figure 13. Microbial community composition of Smeerenburgfjorden sediments. Relative abundance of 16S rRNA ASVs is derived from amplicon sequencing of the 16S rRNA gene. Taxa that are less abundant than 1% or are unclassified are shown in grey. Depth is shown in centimeters below seafloor (cmbfs) for A) Station J, B) Station GK, and C) Station GN.

Supplementary Figure 14. Relative abundances of 16S rRNA and DsrB genes and transcripts for Svalbard sediments. A) Acidobacteriota, B) Thermoanaerobaculia, C) *Ca. Polarisedimenticola* (SD-22), D) *Ca. Sulfomarinibacter* ASV-2257, E) Thermoanaerobaculia DsrB (uncultured family-level lineage 9). Smeerenburgfjorden stations GK, J and GN were sampled in June 2017, and J16 was sampled in July 2016. Replicate cores from Van Keulenfjorden stations AB and AC are derived from Buongiorno et al., 2019.

Supplementary Figure 15. Phylogenetic tree of 16S rRNA gene sequences. Red leaves are from all acidobacteriotal amplicon-derived sequences (ASVs) retrieved in this study from Smeerenburgfjorden, Svalbard. The red dot denotes the most abundant ASV in the dataset. The orange leaf represents the 16S rRNA sequence recovered from an acidobacteriotal metagenome-assembled genome (this study). Blue leaves represent sequences derived from marine environments and present in the SILVA database v138. Green leaves represent cultivated *Acidobacteriota*. SD = 'sub-division', and are numbered as per SILVA database (v138). The tree was built as a consensus of three maximum-likelihood methods (see Materials and Methods). The scale bar represents 10% sequence divergence.

Supplementary Figure 16. Sankey diagram of taxonomic breakdown of marine sediment derived *Acidobacteriota* 16S rRNA genes from the SILVA database (v138 NR).

Supplementary Figure 17. Community compositions of *dsrB*-harbouring microorganisms in Svalbard sediments. **A)** Compositions determined from *dsrB*-gene (DNA) amplicon sequencing. **B)** Compositions determined by *dsrB*-transcript (cDNA) amplicon sequencing. Groups with relative abundances <1% are grouped as 'other'.

Supplementary Figure 18. CARD-FISH images of *Acidobacteriota* from marine sediments. Sediment locations and probes used are listed above panels A-E. Panels with blue cells are DAPI stained, panels with green cells are CARD-FISH hybridised cells from corresponding fields of view. White scale bars represent 10 µm.

bioRxiv preprint doi: <https://doi.org/10.1101/2020.10.01.322446>; this version posted October 1, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

MAG	MAG (bp)	Estimated genome size (bp)	GC (%)	Complete- ness (%)	Contam- ination (%)	Strain hetero- geneity	Number of contigs	GenBank accession
<i>Ca</i> . Sulfomarinibacter MAG AM1	2,426,940	3,362,810	63.3	72.17	2.94	40	666	JACXVY0000000000
<i>Ca</i> . Sulfomarinibacter MAG AM2	1,835,749	3,146,099	63	58.35	7.69	43	828	JACXVZ0000000000
<i>Ca</i> . Sulfomarinibacter MAG AM3-A	1,779,173	2,206,863	60.9	80.62	1.1	0	127	JACXWA0000000000
<i>Ca</i> . Sulfomarinibacter MAG AM3-B	3,394,205	3,824,456	60.7	88.75	4.27	20	533	JACXWB0000000000
<i>Ca</i> . Sulfomarinibacter kjeldsenii MAG AM3-C	3,921,116	4,316,434	60.9	91.17	3.42	20	783	JACXWC0000000000
<i>Ca</i> . Polarisedimenticola svalbardensis MAG AM4	3,685,148	3,887,504	62.2	95.73	5.98	0	180	JACXWD0000000000

Figure 1. Phylogenomic analysis reveals novel Acidobacteriota taxa in marine sediments. Maximum-likelihood tree of concatenated protein sequences from MAGs and genomes. Single marker genes were retrieved with CheckM. Highlighted in blue are MAGs obtained in this study. Highlighted in purple are *dsrAB*-containing MAGs obtained from the NCBI database from the class Thermoanaerobaculia. The genus *Ca. Acidiflorens* is represented by the most complete MAG (GCA_003166525.1) from the corresponding study [12]. Our phylogenomic analysis showed that one MAG that was previously assigned to *Ca. Aminicenantes* (GCA_004524955.1), recovered from the Bothnian Sea [130], is affiliated with the newly proposed family *Ca. Sulfomarinibacteraceae*. Black dots indicate *dsrAB*-containing genomes/MAGs. Bootstrap values with >90% are indicated with filled black circles on nodes. *Nitrospina gracilis* 3/211 (GCA 000341545.2) was used as an outgroup. The scale bar represents 10% sequence divergence.

A. Candidatus Sulfomarinibacter kjeldsenii MAG AM3-C

Figure 3. Gene organization of the *dsr* gene cluster in Acidobacteriota. Scaffold names in blue were retrieved from this study. Scaffold names in purple were derived from best BLASTP hits to sequences from this study. *Ca. Sulfotellmatomonas gaucii* SbA5 was retrieved from Hausmann et al. 2018. Green: *dsr*, dark red: other genes, and orange: hypothetical genes. Shaded blue lines indicate degree of sequence similarity as determined by tBLASTx within EasyFig.

Figure 4. DsrAB uncultured lineage 9 in the DsrAB tree represents members of the Acidobacteriota class Thermoanaerobaculia (sub-division 23). Blue leaves in the DsrAB tree represent MAGs or contigs identified in this study. Red leaves represent the most abundant acidobacteriotal amplicon-derived DsrB sequences identified in this study. Purple leaves represent sequences from MAGs retrieved from public databases. The DsrAB sequences were added to the consensus tree from Müller et al. 2015 in ARB. SD, pertaining to 'sub-divisions' of Acidobacteriota. The scale bar represents 10% sequence divergence.

Figure 5. Box plots depicting the expression of *otr* and *dsrB* relative to a house-keeping gene (DNA-directed RNA polymerase, alpha subunit) from *Ca. Sulfomarinibacter kjeldsenii* MAG AM3-C during microcosm experiments with amendments of tetrathionate versus no-amendment controls. Relative expression was determined by RT-qPCR. Expression of *otr* was significantly higher at day 8 ($p=0.0488$) as determined using a two-tailed T-test, and is indicated by an asterisk. Center lines indicate medians; box limits indicate 25th and 75th percentiles; and whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles.