Title: The *Phaeodactylum tricornutum* Diaminopimelate Decarboxylase was Acquired via Horizontal Gene Transfer from Bacteria and Displays Substrate Promiscuity

Vincent A. Bielinski,^{a*} John K. Brunson,^{a,b*} Agnidipta Ghosh,^c Mark A. Moosburner,^{a,d} Erin A. Garza,^a Zoltan Fussy,^{a,e} Jing Bai,^a Shaun M.K. McKinnie,^f Bradley S. Moore,^{b,g} Andrew E. Allen,^{a,b} Steven C. Almo,^c and Christopher L. Dupont^{a,1}

^a J. Craig Venter Institute, La Jolla CA 92037

^b Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, UCSD, La Jolla CA 92093

^c Department of Biochemistry, Albert Einstein College of Medicine, NY 10461

^d Integrative Oceanography Division, Scripps Institution of Oceanography, University of California San Diego, La Jolla CA 92037

^e Faculty of Science, Charles University, BIOCEV, Vestec, Czech Republic

^f Department of Chemistry and Biochemistry, University of California, Santa Cruz, Santa Cruz, CA, 95064

^g Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California, San Diego, La Jolla, CA 92093

*These authors contributed equally to this work

¹ Address correspondence to cdupont@jcvl.org

Running title: A Novel DAPDC Displays Substrate Promiscuity

The author(s) responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Christopher L. Dupont (cdupont@jcvl.org)

ABSTRACT

Diatoms are predicted to synthesize certain amino acids within the chloroplast, including L-lysine via a diaminopimelate-dependent pathway. Herein, we report that the model diatom, *Phaeodactylum tricornutum*, possesses a chimeric lysine biosynthetic pathway, which coalesces bacterial and plant genes, and is terminated by a chloroplast-localized diaminopimelate decarboxylase (DAPDC, *PtLYSA*). We show that while RNAi ablation of *PtLYSA* is either synthetically lethal or concomitant with a slower growth rate, Cas9-mediated mutagenesis of *PtLYSA* results in recovery of heterozygous cells lines, suggesting that *PtLYSA* is an essential gene. Previously characterized DAPDCs are unique within the PLP-dependent decarboxylases where catalysis occurs at the D-stereocenter of the substrate and display a strict stereochemical preference for a (D,L)- or *meso*-substrate and not the D,D- or L,L-isomers of diaminopimelate (DAP) to synthesize L-lysine. Using decarboxylation assays and differential scanning calorimetry analyses, we validate that *PtLYSA* is a *bona fide* DAPDC and uncover its unexpected stereopromiscuous behavior in substrate specificity. The crystal structure of *PtLYSA* confirms the enzyme is an obligate homodimer in which both protomers reciprocally participate in the active site. The structure underscores features unique to the *PtLYSA* clan of DAPDC and provides structural insight into the determinants responsible for the substrate-promiscuity observed in *PtLYSA*.

INTRODUCTION

Lysine plays a variety of important roles in biology across the tree of life. Besides its role as an essential amino acid required for protein synthesis, L-lysine can also be degraded into the lipid building block acetyl-CoA via lysine catabolism pathways in some organisms, including *E. coli*, *Arabidopsis* and humans (Knorr et al., 2018; Zhu et al., 2004; Chang, 1978). In Gram-positive *Staphylococcus* strains, L-lysine is found as a structural component of the peptidoglycan layer and the biosynthetic enzymes required for peptidoglycan production are involved in β -lactam resistance (De Lencastre et al., 1999). Two main routes for the *de novo* production of L-lysine exist in nature, although new and unique pathways are still being discovered in bacteria (Price et al., 2018). Higher fungi, euglenids and some bacteria produce L-lysine from homocitrate via the α -amino adipate pathway (Xu et al., 2006; Kosuge et al., 1998). Land plants and most bacteria employ four biochemically distinct variations of the diaminopimelate (DAP) pathway to synthesize L-lysine from an aspartic acid precursor (Pratelli and Pilot, 2014; Kirma et al., 2012; Jander and Joshji, 2009; Hudson et al., 2005). In plants, the pathway includes an aminotransferase activity that produces L,L-DAP as an intermediate. L,L-DAP undergoes a subsequent epimerization to *meso*-DAP (D,L-DAP), the substrate for the final enzymatic step, which is catalyzed by diaminopimelate decarboxylase (DAPDC) to produce L-lysine (Bukhari and Taylor, 1971; White and Kelly, 1965; Dewey and Work, 1952).

DAPDC proteins are unique in the amino acid decarboxylase family as the only members to carry out decarboxylation exclusively at the D-stereocenter of the substrate. All DAPDC enzymes characterized thus far display a strict substrate preference for *meso*-DAP over other DAP stereoisomers and function as homodimeric complexes (Peverelli et al., 2016, Peverelli and Perugini, 2015, Griffin et al., 2012; Hu et al., 2008, Hutton et al., 2007). Elimination of CO₂ from *meso*-DAP by DAPDC requires the cofactor pyridoxal-5'-phosphate (PLP), which forms a Schiff base with a conserved lysine residue (K54 in *E. coli* LYSA, UNIPROT ID P00861) in the active site of the enzyme (Ray et al., 2002). PLP facilitates the decarboxylation of *meso*-DAP substrate via formation of a Schiff base involving the α -amino group of the *meso*-DAP D-stereocenter, while distal residues in the DAPDC catalytic pocket interact with the L-stereocenter of *meso*-DAP (Hu et al., 2008, Gokulan et al., 2003).

The structures of DAPDC orthologs reveal that both protomers of the dimer contribute to form the active sites in which a conserved arginine from the EPGR motif directs preference for *meso*-DAP (Crowther et al., 2019; Son and Kim, 2018; Weyand et al., 2009; Hu et al., 2008; Gokulan et al., 2003; Ray et al., 2002). These structures evince that DAPDC orthologs have a variable and flexible active site loop that shields the active site ligands from bulk solvent during catalysis (Hu et al., 2008).

In photosynthetic eukaryotes studied thus far, it appears the terminal steps of lysine biosynthesis occur exclusively within the chloroplast (Hudson, 2006; Perl, 1992; Wallsgrove, 1980). It has been postulated that lysine biosynthetic gene clusters from the cyanobacterial lineage were obtained via the primary plastid endosymbiotic event that resulted in the emergence of Archaeplastida that encompass glaucophytes, red and green algae and land plants (Reyes-Prieto, 2012; Kidron, 2007; Velasco, 2002). Diatoms are ubiquitous ocean phytoplankton and major contributors to the carbon cycle (Armbrust, 2009), which were derived from a secondary endosymbiosis event and thrive in upwelling-induced, nutrient-rich conditions making them the basis for the world's shortest and most energy-efficient food webs. Similar to land plants and green algae, diatoms are capable of *de novo* synthesis of essential amino acids, including L-lysine via a predicted aminotransferase-dependent DAP pathway (Figure 1). Genomic reconstruction of pathways in diatoms suggests many amino acid biosynthetic genes are present in multiple copies and function within organelles, while a subset of biosynthetic genes is present as single copies (Bromke, 2013; Bowler et al., 2008).

Within the genome of the model diatom *Phaeodactylum tricornutum*, the predicted lysine biosynthesis genes exist as single copies, and the corresponding proteins of the last four steps of the pathway predicted to localize within the chloroplast via canonical ASAF-type motifs and N-terminal chloroplast targeting peptides (Kilian and Kroth, 2005; Apt et al., 2002). In this report, we describe the phylogenetic origins of lysine biosynthetic machinery in diatoms. In addition, we report subcellular localization and biochemical activity of the enzyme involved in the terminal step of lysine biosynthetic pathway in *P. tricornutum*, and reveal that the enzyme (Phatr3_J21592, UNIPROT ID B7G3A2; *Pt*LYSA hereafter) is essential and harbors unexpected promiscuity in substrate utilization. In order to gain insight into this unique behavior, we determined the crystal structure of *Pt*LYSA, and defined the determinants responsible for substrate promiscuity in diatoms.

146

147

148

RESULTS

Phylogenetic Analysis

The DAP biosynthetic pathway in diatoms appears to be a hybrid of components from plants and bacteria potentially acquired via horizontal gene transfer (HGT), a process previously described in sequenced diatom genomes (Bowler, 2008). For the five enzymes catalyzing the synthesis of L-lysine from L-aspartate 4-semialdehyde (Table 1), orthologs in sequenced genomes and transcriptomes were identified, aligned, and examined using a maximum likelihood analysis (Figure 2). In parallel, target peptide analyses were conducted to predict subcellular localization (Table 1). The gene *PtDAPA* (Phatr3_J11151) encodes a putative dihydrodipicolinate synthase (DDPS) homolog in *P. tricornutum* and cannot be confidently localized based on target peptide analysis (Table 1). Our analyses reveal that *PtDAPA* has orthologs in other stramenopile genomes; however, the closest ancestors are in gammaproteobacteria (Figure 2A). For the three intermediate steps in the pathway, the diatom proteins (carried out by *PtDAPF*, *PtDAPL* and *PtDAPB*) also have orthologs in the stramenopiles, the closest ancestors of which are found in plant or cyanobacterial genomes (*DAPL* in Figure 2B, *DAPF* and *DAPB* in Supplemental Figure 1). Each of these proteins also contain putative N-terminal chloroplast targeting sequences (Gruber and Kroth, 2017; Apt et al., 2002).

The terminal step of the DAP pathway in *P. tricornutum* is predicated to be catalyzed by *PtLYSA* (Phatr3_J21592), which has orthologs throughout the stramenopile lineage; however, our analyses strongly suggest that the stramenopile lineage acquired the gene through horizontal gene transfer from Bacteria, specifically the PVC (Planctomycete-Verruca-Chlamydia) supergroup (Figure 2C). The diatom sequences for the enzymes in the pathway form monophyletic clades in phylogenetic inferences, suggesting that these enzymes are conserved through the entire lineage. In addition, our analyses reveal that the long branch length of the diatom and *Bolidophyceae* *LYSA* clade is associated with the insertion of a highly conserved 9 amino-acid segment into the active site loop of the predicted protein (Figure 2D). In this work we focused on experimentally validating the terminal step of this chimeric lysine biosynthetic pathway for the following reasons: 1) bioinformatic annotations of the *P. tricornutum* genome suggest the presence of three putative organic acid decarboxylase genes (Bowler et al., 2008) however the *LYSA* enzymes in diatoms

has not been biochemically validated thus far, **2)** the presence of the chloroplast targeting sequence in the *PtLYSA* protein suggesting a similar localization of the pathway observed in plants, and **3)** inclusion of a highly conserved substitution in a critical substrate-binding residue and a unique insertion in the active site loop, while maintaining its ancestral link to bacteria in the PVC supergroup.

Determination of *PtLYSA* Subcellular Localization

The subcellular localization of *PtLYSA* in *P. tricornutum* was determined using transgenic and exconjugant diatoms expressing C-terminal fusions of *PtLYSA* with either YFP or mTurquoise2 (mT2). Transgenic diatoms expressing *PtLYSA*-YFP fusions driven by *pFCPB* promoter were generated by particle bombardment (Apt et al., 1996). Confocal microscopy showed that on occasion the YFP signal formed distinct punctate distributions within the chloroplast (Figure 3A), while in other transgenic strains the YFP signal was located throughout the chloroplast. Exconjugants expressing similar *PtLYSA*-mT2 protein fusions on stable episomes were also generated, which provides a more controlled heterologous expression platform (Karas et al., 2015). Confocal microscopy of *Phaeodactylum* expressing a C-terminal mT2-fused version of *PtLYSA* revealed a distinct regional overlap of the mT2 fluorescence with the autofluorescence of chlorophyll in the chloroplast (Figure 3C). We observed that exconjugant cells with constitutive expression of *PtLYSA*-mT2 showed signs of cell stress, grew poorly and adopted an ovoid morphology. We chose to employ the nitrate inducible pNR promoter (Chu et al., 2016; Poulsen and Korger, 2005) to repress *PtLYSA*-mT2 expression during conjugation and strain selection and inducing expression after colony isolation. Exconjugants were generated and selected using ammonium as the nitrogen source and upon induction of cultures with nitrate, the cell morphology of these cultures quickly changed from pennate to ovoid and the cells began aggregating. In comparison, cultures expressing a cytosolic version of mT2 had little to no ovoid cells and grew to a higher cell density.

***PtLYSA* RNAi and CRISPR**

Reported transcriptomic analysis of the lysine pathway genes in *P. tricornutum* reveals the expression patterns of the associated genes are diel-regulated, with the aminotransferase *PtDAPL* showing the greatest change in expression over the

course of the day (Smith et al., 2016; Supplemental Figure 2). In order to determine if the putative *PtLYSA* protein is necessary for growth, we performed RNAi-driven gene knockdown, as well as TALEN and CRISPR-Cas9 gene mutagenesis experiments. For the RNAi-based gene knockdown, overlapping 250-bp and 400-bp intragenic fragments of *PtLYSA* were generated and cloned into a receiver vector as described (De Riso et al., 2008), with the exception that Gibson assembly (Gibson et al., 2008) was used instead of classical restriction enzyme dependent cloning. The receiver vector contains a transcriptional unit conferring phleomycin resistance and a cloning site with either the *pFCPB* or *pNR* upstream of the cognate terminator sequence. All attempts at attaining transformants of RNAi driven by *pFCPB* failed to yield colonies, despite the inclusion of a non-lethal positive control targeted to the *P. tricornutum* urease gene (Phatr3_J29702) that yielded transformants. We were able to attain RNAi-expressing transgenic lines where the hairpin loop for *PtLYSA* is driven by *pNR* only if the transformation is plated with ammonia as a sole nitrogen source. These lines exhibit reduced growth rates and *PtLYSA* protein content (Supplemental Figure 3) when grown on nitrate (RNAi-induced) instead of ammonia (RNAi-silenced).

Gene ablation vectors were made with the TALEN system described for *P. tricornutum* (Weyman et al., 2015); while colonies were obtained for the positive control urease knockout lines, none could be recovered for cell lines transformed with vectors targeting *PtLYSA*. Subsequent attempts were carried out using episome-delivered CRISPR-Cas9 methodology that includes multiplexed small-guide RNAs (sgRNAs, Moosburner et al., 2020). Two sgRNAs were delivered with Cas9 together or individually to *Phaeodactylum* using bacterial conjugation (Diner et al., 2016; Karas et al., 2015). The first attempt to knockout *PtLYSA* used a Cas9-fusion episome that harbors both sgRNAs, g21592-1 and g21592-2. Initially, two transformation attempts were made to produce transgenic *P. tricornutum* lines. Each time, less than 20 colonies were obtained on the plate, even after transformation optimization. On the third attempt, the selection plate growth media was supplemented with L-lysine (40 $\mu\text{g mL}^{-1}$), which produced over 100 colonies on the selection plates.

A total of 24 colonies from the third conjugation attempt were analyzed for Cas9-induced indels by Sanger sequencing and TIDE analysis. The expected mutagenesis genotype was an 18-bp deletion between the two sgRNA cleavage sites. Manual curation of the sequences did not produce any obvious knock-out

mutations but had a signature of Cas9 activity at the expected cut site(s). TIDE analysis revealed that 6 of 24 cell lines were heterozygous, or heterozygous where one allele contained an 18-bp deletion and the other allele retained a wild-type sequence (Figure 4). This observation was surprising since an 18-bp deletion should not knock-out the protein but could result in reduced enzyme activity or protein misfolding by removing 6 amino acids within the open-reading frame. Other attempts to produce a biallelic mutant cell line included re-streaking colonies to produce monoclonal cell lines and growing the heterozygous cell lines on increasing concentrations of L-lysine to force a biallelic mutation to occur. Nonetheless, the only genotype that could be produced using both sgRNAs to cut out an 18-bp sequence was heterozygous.

Subsequently, each sgRNA was delivered to *Phaeodactylum* individually and selected using growth media supplemented with L-lysine. 24 colonies were picked and analyzed for genotypes as described previously. Colonies that contained g21592-2 could not produce a transgenic genotype and were all wild-type at the *PtLYSA* locus. As for g21592-1, 2 of 24 colonies showed the effects of Cas9 activity after manual sequence curation and TIDE analysis. Since the induced mutations would be random indels, the colonies were re-streaked and monoclonal cell lines were picked again. Of 24 colonies, 18 had Cas9 activity at the g21592-1 cut site. TIDE analysis revealed that one third (8/24) of colonies were clearly heterozygous (Figure 4). Three colonies had the genotype of a 12-bp deletion plus wild-type and five colonies had the genotype of a 10-bp deletion plus wild-type. The other colonies had either a wild-type genotype (4), a mixed genotype without a wild-type signature (5), or a mix of indel and wild-type (2) (Supplemental Figure 4). In the 10-bp deletion plus wild-type heterozygous cell lines, a deletion of a 10-bp sequence would result in a deleterious frame-shift by introducing a premature stop codon in one of the two alleles. Therefore, one functional allele is sufficient to retain L-lysine biosynthesis in the chloroplast; however, using two sgRNAs, a biallelic mutant cell line of *PtLYSA* could not be obtained.

Biochemical Analysis of *PtLYSA*

Primary sequence analysis indicated that *PtLYSA* is homologous to an organic acid decarboxylase, likely a DAPDC, a well-studied and structurally characterized enzyme family (Hu et al., 2008; Gokulan et al., 2003; Ray et al., 2002). The *P.*

tricornutum genome is predicted to harbor genes for putative DAP, ornithine and arginine decarboxylases (Smith et al., 2016; Bowler et al., 2008). To biochemically validate *PtLYSA* as a DAPDC, we cloned full-length and varying N-terminal deletion constructs into a pBAD-driven bacterial expression vector, fusing them with a C-terminal 6xHis-Flag affinity tag (Savitsky et al., 2010). To screen for expression and solubility at small scale, the resultant plasmids were transformed into *E. coli* BL21 cells (Supplemental Figure 5). Overnight induction with 0.5% L-arabinose at 30 °C yielded several soluble constructs, with increased solubility at 18 °C. Judging by the level of expression and solubility, we chose to purify the $\Delta 36$ -*PtLYSA*-F481L variant protein in large quantities using immobilized metal (Ni) affinity chromatography (IMAC) and tested the purified enzyme for *in vitro* activity towards multiple substrates including ornithine and all three DAP stereoisomers (Figure 5, Supplemental Figure 6; Methods). The resolution of substrates and products of $\Delta 36$ -*PtLYSA*-F481L was achieved through pre-column derivatization with 1-fluoro-2,4-dinitrophenyl-5-L-alanine amide (L-FDAA), known as Marfey's reagent (Bhushan and Brückner, 2004).

Derivatization of DAP and lysine with Marfey's reagent allows for chromatographic separation of stereoisomers followed by UV-detection (340 nm) and was used for reaction assay analysis by HPLC (Methods). We found that the purified *PtLYSA* enzyme produced L-lysine from *meso*-DAP and displayed no activity towards ornithine (Supplemental Figure 6), confirming biochemical function of *PtLYSA* as a DAPDC. When the purified $\Delta 36$ -*PtLYSA*-F481L protein was challenged with three different DAP stereoisomers, to our surprise, we observed that the enzyme is capable of producing D-lysine from the D,D-DAP and L-lysine from L,L-DAP. Both the *meso*-DAP and D,D-DAP reactions go to completion; however, the L,L-DAP reaction has significant amount of unreacted substrate even after overnight incubation (Figure 5).

PtLYSA was also subjected to differential scanning fluorimetry (DSF) against a pool of potential ligands (Figure 6). Purified *PtLYSA* exhibited a melting temperature (T_m) of 49 °C (± 1) and is stabilized by the potential ligands. The substrate *meso*-DAP increased thermal stability (ΔT_m) of *PtLYSA* by 4.8 °C (± 0.8), indicating a potential interaction between *meso*-DAP and *PtLYSA*. Similarly, *PtLYSA* was stabilized by the product, L-lysine ($\Delta T_m = 5.9$ °C (± 1)). DSF data revealed that the D,D-DAP isomer was able to stabilize *PtLYSA* by 5.8 °C (± 0.6); however, L,L-DAP enhanced the

thermal stability by 1 °C ($T_m = 50\text{ °C} (\pm 0.2)$; Figure 6). These DSF experiments are consistent with the preferred utilization of *meso*-DAP and D,D-DAP as substrates, relative to L,L-DAP. These data are also consistent with the incomplete processing of L,L-DAP observed during *in vitro* assays (~50% turnover to L-lysine after overnight incubation, Figure 5), in contrast to using D,D-DAP and *meso*-DAP to yield D-lysine and L-lysine, respectively. In addition, D-lysine binds to *PtLYSA* and induces a similar thermal stability similar to L-amino acid (Figure 6), suggesting that the catalytic pocket of *PtLYSA* accommodates both enantiomeric forms of lysine.

The use of D,D-DAP as a substrate and the production of D-lysine is remarkable, as reported DAPDC orthologs, thus far, exhibit strong selectivity towards *meso*-DAP as substrate. Structural analyses reveals that amino acids sidechains in DAPDC active sites orient *meso*-DAP for catalysis and restrict D,D-DAP from engaging in a productive orientation with respect to the bound PLP cofactor necessary for decarboxylation reaction (Hu et al., 2008; Gokulan et al., 2003; Ray et al., 2002). In particular, a conserved arginine side chain in DAPDC orthologs appears to impose extensive steric hindrance to bar D,D-DAP; interestingly in diatoms the arginine side chain at this position is replaced by a threonine (T317 in *PtLYSA*).

Crystal Structure of *PtLYSA*

On the basis of reported coordinates of DAPDC enzymes in the RCSB Protein Data Bank, a construct of *PtLYSA* encompassing amino acids 39-476, which lacks the N-terminal signal peptide, was purified and co-crystallized in the presence of D-lysine (Methods). The structure of *PtLYSA* was determined by molecular replacement using *E. coli* DAPDC (PDB: 1KNW) as a search model. The final model refined at 2.78 Å resolution had R_{cryst} and R_{free} values of 0.18 and 0.23, respectively (Table 2).

Crystals contained two molecules of *PtLYSA* in the asymmetric unit (protomers 1 and 2; Figure 7A), suggesting a dimer. Size-exclusion chromatography is consistent with the proposed dimer (elution time corresponding to an apparent molecular weight of ~100 kDa; monomer molecular weight of ~50 kDa) (Supplemental Figure 7), suggesting the protein is a dimer in solution. The two protomers are arranged in a “head-to-tail” configuration related by 2-fold non-crystallographic symmetry. The interface is polar in nature and shows remarkable surface complementarity while burying a total of ~4120 Å² of solvent assessable surface area (Supplemental Figure 8) upon dimerization.

Structure of *Pt*LYSA in relation to DAPDC orthologs

Individual protomers of *Pt*LYSA consist of two domains: an N-terminal 8-fold α/β barrel and a C-terminal β sandwich domain (Figure 7A), with both domains contributing to the dimer interface. The barrel domain lacks the N-terminal β sheet extension of *C. glutamicum* (Son and Kim, 2018), *M. tuberculosis* (Weyand et al., 2009) and *M. jannaschii* (Ray et al., 2002) enzymes; however, the domain is topologically similar to *H. pylori* (Hu et al., 2008), *E. coli* (PDB: 1KNW) and *T. maritima* (PDB: 2YXX) DAPDC, which lack the N-terminal extension. The protomers of *Pt*LYSA are similar to these last three DAPDC orthologs with a root-mean-square deviation (RMSD) between 1.1 and 1.3 Å over 249 aligned C α atoms. Therefore, the structure unambiguously confirmed that *Pt*LYSA belongs to the group IV pyridoxal-5'-phosphate (PLP) dependent decarboxylase family (Kern et al., 1999; Sugio et al., 1995).

Structure-based sequence alignment of the DAPDC orthologs underscore two structurally divergent regions in *Pt*LYSA: 1) the loop segment between β 6 and α 7 (β 6- α 7 loop; amino acids 183-209), and 2) the loop segment between β 7 and α 9 (β 7- α 9 loop henceforth; amino acids 238-245) (Supplemental Figure 9). In *Pt*LYSA, the β 6- α 7 loop has a 7 amino acid insertion, which could only be built completely as a random coil into the electron density of protomer 1 (Figures 7A and 7B). Akin to protomer 2 of *Pt*LYSA, this segment is not included in the structures of DAPDC from *Aquifex aeolicus* (PDB 2P3E) and *Brucella melitensis* (PDB 3VAB). In the structure of *H. pylori* DAPDC (PDB: 2QGH), the analogous segment has been annotated as 'active site loop', which occludes the active site ligands from solvent (Hu et al., 2008). In *H. pylori* this region forms a 3_{10} helix (Figure 7B). In *C. glutamicum* (PDB: 5X7M, Son and Kim, 2018) and *M. tuberculosis* (PDB: 2O0T, Weyand et al., 2009; PDB: 1HKW, Gokulan et al., 2003), this segment forms a two stranded anti-parallel β sheet (Figure 7B). In *M. jannaschii* (PDB: 1TUF, Ray et al., 2002) and *T. maritima* (PDB: 2YXX), this segment adopts a helical conformation; and in *E. coli* (PDB: 1KNW) it forms a similar, but smaller random coil compared to *Pt*LYSA segment (Figure 7B).

The β 7- α 9 loop forms a wall of the active site in all DAPDCs (Figure 8), and the *Pt*LYSA enzymes has a 5 amino acid deletion in this segment (Supplemental Figure 9), which causes an expansion of the *Pt*LYSA active site (Figure 8) compared

to orthologs from *C. glutamicum*, *M. tuberculosis*, *M. jannaschii*, *H. pylori* and *T. maritima*. An analogous deletion has also been observed in the $\beta 7$ - $\alpha 9$ loop of *E. coli* DAPDC (Supplemental Figure 9), in which it adopts a conformation similar to that observed in *PtLYSA* (Figure 8).

Architecture of the substrate binding site in *PtLYSA*

The structural analyses reveal that both protomers of the *PtLYSA* dimer contribute to the active site by reciprocally sharing 8 conserved amino acids: Lys95, His238, Glu314, Tyr430, Cys392, Glu393, Ser394, and Thr317. The Thr317 is unique to *PtLYSA* (Figure 7C, Supplemental Figure 9), as the corresponding residue in all structurally characterized DAPDCs is an arginine, which is part of the EPGR motif (Supplemental Figure 9) (Son and Kim, 2018, Weyand et al., 2009; Hu et al., 2008; Gokulan et al., 2003; Ray et al., 2002). While the sidechains of Lys95, His238, Tyr430 and Thr317 are supplied in *cis* to the active site of the protomers, the side chain of Cys392, Glu393 and Ser394, which form a conserved CESH/SD motif in DAPDC, are shared in *trans* from the other protomer (Figures 7C and 7D). In DAPDC orthologs, the conserved side chain at Lys73 from the Y/FAXKA motif (Supplemental Figure 9) forms a Schiff base with the cofactor, PLP, which exhibits aromatic stacking interaction with the side chain of His238. The side chains of Glu314 (OE) and Tyr430 (OH) contact (H-bonds) the pyridine endocyclic nitrogen (N1) and phosphate (O1P) of the PLP, respectively.

In the active sites of the *PtLYSA* dimer, a sulfate ion occupies the position of the phosphate group of PLP (Figures 7C and 7D). Superimpositions of *PtLYSA* protomers with PLP-bound DAPDC structures reveal that the coordination sphere of the sulfate ion is very similar to that observed in the PLP-PO₄ interaction. However, there is a striking adaptation observed in the polyanion coordination in *PtLYSA*; the sidechain oxygen (OG1) of Thr317 (EPGR motif; Supplemental Figure 9) is engaged in a H-bond interaction with the SO₄ (Figures 7C and 7D). The arginine sidechains in DAPDC orthologs do not participate in such an interaction (Figure 8), instead they contribute to substrate specificity and stabilize the active site by interacting with a tyrosine sidechain OH via a H-bond. The tyrosine is conserved in DAPDC orthologs; however, it is replaced by an arginine, Arg278, in *PtLYSA*.

Additional, but weak, electron density features were observed adjacent to the SO₄ ion in the active sites of *PtLYSA*. The densities were fenced by Lys275, Arg278,

Arg358 and Tyr360, and interpreted and modeled as co-crystallized D-lysine for protomer 1 (Figure 7D). Given the weak density and modest 2.78 Å resolution of the refined coordinates, the assignment of D-lysine remains speculative. It is, however, notable that only Arg358 and Tyr360 are conserved among the DAPDC orthologs (Supplemental Figure 9).

DISCUSSION

Diatoms evolved via serial endosymbiotic events between a photosynthetic symbiont and a non-photosynthetic exobiont that have occurred over the past 1.8 billion years (Armbrust et al., 2004). In addition to these processes, there have been substantial horizontal gene transfer events between diatoms and bacteria (Mock et al., 2017; Raymond et al., 2012; Bowler et al., 2008). The phylogenetic analyses reported within conclusively show the lysine biosynthetic pathway in diatoms is conserved across all species examined by genome or transcriptome sequencing to date. This conserved pathway includes genes likely derived from the symbiont in the primary endosymbiotic event (*DAPL*, *DAPB*, *DAPF*) as well as two genes (*DAPA*, *LYSA*) likely derived from cryptic horizontal gene transfers from Bacteria. *DAPA* appears to have originated from gammaproteobacteria in the *Vibrio* lineage, which is not a common Bacterial HGT partner for diatoms (Bowler, et al. 2009; Bowler et al., 2008). In contrast, the *LYSA* appears to have been transferred from the Verrucamicrobia lineage, which has been predicted to have contributed a substantial number of genes to modern diatom genomes (Bowler et al., 2008). The phylogenetically chimeric pathway is reminiscent of the urea cycle and overall nitrogen assimilation pathways within diatoms (Smith et al., 2019; Allen et al., 2011), both of which are composed of genes from different evolutionary partners of the serial endosymbiotic events. The conservation of this chimeric pathway across all diatoms sequenced to date suggests strong selective pressure for the retention of the pathway and that this event occurred early in the evolution of the diatom lineage.

Subsequent work focused on the terminal step of the pathway, which was hypothesized to be catalyzed by *PtLYSA*. Biochemical assays confirmed the purified protein converts *meso*-DAP to L-lysine. The observation that overexpression of FP-linked fusions of *PtLYSA* in *P. tricornutum* cells results in the co-localization of signal with chloroplast autofluorescence is consistent with the prediction that *PtLYSA* contains a canonical bipartite transit peptide harboring an ASAF cleavage site (Gruber and Kroth, 2017; Apt et al., 2002). This localization also corroborates previous reports that in plants lysine production is carried out in the chloroplast (Mills et al., 1980; Wallsgrove and Mazelis, 1980) and that the three genes upstream of *PtLYSA* (*DAPF*, *DAPL* and *DAPB*) encoding the terminal arm of the *P. tricornutum* lysine pathway also contain predicted N-terminal chloroplast targeting peptides (Table 1). The localization of these enzymes in the chloroplast has potential

biochemical and system-wide consequences. These steps of the pathways will be subject to large changes in redox state (Rosenwasser et al., 2014; Baier and Dietz, 2005). Diurnal swings in chloroplast redox state may have significant implications for *DAPF*, where a dimerization is not only necessary for function but also sensitive to reversible disulfide-bond formation between protomers (Sagong and Kim, 2017). It is possible that redox-responsive disulfide bond disruption deactivates *DAPF* during the dark periods where the local chloroplast environment becomes more reducing, potentially halting the supply of *meso*-DAP. This potential post-translational effect is an additional level of regulation, superimposed upon that of diel gene expression of the entire lysine pathway, which is transcribed at higher levels during the light period in *P. tricornutum* (Smith et al., 2016) and may contribute to regulation of carbon flux and assimilation during the day/night cycle.

The dynamic regulation of the lysine biosynthetic pathway genes at the transcript level suggests an important role for this pathway in diatom metabolism. This notion is supported by the observed changes in the engineered gene expression during the course of this study. Overexpression of a heterologous copy of *PtLYSA* from an episome using a non-diel responsive promoter in a wild-type background induces a change in *P. tricornutum* cells from fusiform to ovoid morphotype, which is associated with cell stress (Lauritano et al., 2015). Controls show that this effect is not due to the expression of fluorescent tag. Similarly, homozygous gene knockouts were not attainable, with only heterozygous exconjugants being attained in media supplemented with exogenous L-lysine when Cas9 editing of *PtLYSA* was attempted. These data demonstrate a functional allele of *PtLYSA* is essential for growth. These heterozygous exconjugants contained alleles coding for a native copy of *PtLYSA* as well as a copy with a 6 amino-acid deletion that is predicted to have minimal effect on the tertiary structure of the homodimeric enzyme. Deletion analyses identified the LEEAA sequence segment (amino acids 72-76) in *PtLYSA* as essential (Supplemental Figure 9). This segment is not conserved among DAPDC orthologs, hence the basis for its essentiality was not immediately apparent. In the crystal structure, this segment forms a helix ($\alpha 2$ in *PtLYSA*, Supplemental Figure 9), which is structurally conserved across the orthologs despite the lack of sequence similarity. The *PtLYSA* structure underscores that this segment plays a secondary but important role in catalysis by buttressing residues in and adjacent to the active site (Supplemental Figure 10). We surmise that analogous deletion in DAPDCs would

result in similar phenotypes in the orthologs. RNAi with constitutive promoters was also unsuccessful, though inducible RNAi successfully generated transgenic lines, which were used to verify that knockdown of *PtLYSA* expression has a detrimental effect on growth (Supplemental Figure 4). Overall, *PtLYSA* appears to be an essential gene in *P. tricornutum* and these results suggest that a diverse gene editing tool kit beyond just CRISPR-Cas guided gene knockouts is necessary to characterize *LYSA* and possibly other essential genes in diatoms.

The activity of *PtLYSA* towards both *meso*-DAP and D,D-DAP (Figure 5) was unexpected, and is consistent with DSF data showing that *PtLYSA* is able to bind both *meso*-DAP and/or D,D-DAP with similar ΔT_m s (Figure 6). It should be noted that while the biochemical assay demonstrates that *PtLYSA* can use L,L-DAP as substrate, and the reaction does not proceed to completion under the conditions employed (Figure 5). Additionally, L,L-DAP did not impact the thermal stability of *PtLYSA* (Figure 6). Metabolomic studies in diatoms are in their infancy, therefore it is not known if diatoms produce D,D-DAP *in vivo*, or encounter it in its environment. Production of D,D-DAP *in vivo* would also be biochemically unprecedented, as D,D-DAP has never been demonstrated to be a physiological substrate for DAPDC enzymes, nor has it been demonstrated to be a potential product of DAPF epimerase activity in any organism (Hor et al., 2013). Therefore, the topic remains an open area of investigation. However, flexibility in substrate utilization may be advantageous under selective pressure, allowing organisms to take advantage of non-canonical substrates or cofactors encountered in the environment.

The structure of *PtLYSA* confirms that the protein forms a “head-to-tail” symmetric homodimer, and buries a large amount of solvent accessible surface area, including the active site (Supplemental Figure 8). The extensive dimer interface of *PtLYSA* along with the *cis-trans* orientation of the active site and its behavior in solution (Supplemental Figure 7) suggests that the protein is an obligate dimer, consistent with orthologs from diverse organisms (Son and Kim, 2018; Peverelli et al., 2016; Griffin, 2012 et al.; Weyand et al., 2009; Hu et al., 2008; Gokulan et al., 2003; Ray et al., 2002). In this aspect, *PtLYSA* is similar to members of the DAPDC superfamily; however, unique aspects of the structure also provide some insight as to the stereochemical origins of the stereopromiscuity. The active site of *PtLYSA* possesses three striking adaptations: 1) an insertion in the active site loop ($\beta 7$ - $\alpha 9$

loop), **2**) a deletion in the $\beta 7$ - $\alpha 9$ loop, and **3**) presence of a unique Thr317 residue within the active site (Supplemental Figure 9). The active site loop insertion appears to afford additional conformational flexibility, which might impact substrate selectivity in *PtLYSA*. Deletion in the $\beta 7$ - $\alpha 9$ loop expands the active site significantly (Supplemental Figure 9), and potentially removes steric hindrances, which could allow multiple stereoisomers to bind; the notion is consistent with the DSF analyses (Figure 6). A deletion in $\beta 7$ - $\alpha 9$ loop has also been observed in *E. coli* DAPDC (Supplemental Figure 9), which was shown to be able to bind D-lysine in the active site (PDB: 1KO0). Notwithstanding, the most remarkable adaptation in the active site of *PtLYSA* is the replacement of the conserved arginine in the EPGR motif with a threonine (Supplemental Figure 9). Similar substitution at this residue are observed in all sequenced diatoms to date, as well as some bacterial (*Leptospira interrogans*), algal (*Ostreococcus lucimarinus*) and cryptophytes (*Guillardia theta*) lineages. The side chain of Thr317 coordinates a SO_4 , which occupies the position of PLP- PO_4 (Figures 7 and 8). The shorter sidechain of Thr317 may relax substrate specificity and structural constraints imposed by the arginine in structurally characterized DAPDC orthologs. Importantly, each of these three active site adaptations are not specific to *P. tricornutum* and are conserved across all putative diatom LYSA proteins.

METHODS

Phylogenetic Analysis

Sequences for the analysis were retrieved from NCBI-nr, PhyloDB, JGI, and MarDB (<https://mmp.sfb.uit.no/databases/mardb/>) using DIAMOND BLAST (Buchfink, 2015). Outgroup (non-diatom) sequences were clustered on 50% identity and 80% coverage using MMseqs2 (Steinegger and Soding, 2017). Datasets were aligned by MAFFT v7.407 (Katoh and Standley 2013) using the L-INS-i refinement and a maximum of 1000 iterations and trimmed by trimAl v1.4 (Capella-Gutiérrez, 2009) allowing 70% sequences to have a gap (-gt 0.3). Maximum likelihood (ML) trees were inferred by IQ-TREE v 1.6.12 (Nguyen, 2015) using the LG+C20+F+G model and the posterior mean site frequency method (Wang, 2018), starting from a LG+F+G guide tree and employing the strategy of rapid bootstrapping followed by a “thorough” ML search with 1,000 ultra-fast bootstrap replicates.

To resolve the topology of LYSA with multiple long branches, the ML tree was compared with a Bayesian topology. The latter was inferred by PhyloBayes MPI v1.7 (Lartillot, 2013) under the CAT-GTR model most robust against long-branch attraction artifacts in three independent chains run for ~40,000 generations when they reached convergence at 0.128 maximum discrepancy. Then, burn-in of 4,000 generations (10 %) was discarded from each chain and a consensus tree was calculated from every-10th-tree subsamples.

Plasmids for Gene Deletion Studies

Vectors for inhibiting the expression of *PtLYSA* via RNAi were designed using previously described plasmids (Allen et al., 2011). The plasmid backbone was linearized via PCR using primers LYS-038/039. 250-bp and 400-bp targeting fragments were amplified from the *PtLYSA* gene using primers LYS-040/041 and LYS-041/042, respectively. Loop-hairpin overlaps targeting *PtLYSA* were assembled using Gibson cloning. The resulting vectors drive loop-hairpin expression using the relatively high expression promoter *pFCPB* coupled with the terminator *tFCPA*. In a subsequent build, this vector was linearized using primers that exclude the *pFCPB* element and then assembled with *pNR* using Gibson cloning.

Plasmids for TALEN were constructed as previously described (Weyman et al., 2015) and introduced via electroporation. Plasmids for *P. tricornutum* CRISPR

were built using Golden Gate-based assembly methods for CRISPR-Cas9 targeting vectors ([dx.doi.org/10.17504/protocols.io.4acgsaw](https://doi.org/10.17504/protocols.io.4acgsaw)).

Plasmids for Localization

PtLYSA-YFP fusion constructs were built with Gateway cloning method (Siaut et al., 2007). Plasmids for episomal overexpression of fusion proteins in *P. tricornutum* were based upon *PtPBR1* and contained an *oriT* to allow for conjugation of the plasmids from bacteria (Karas et al., 2015). The predicted promoter and terminator for *PtLYSA* were amplified from *Phaeodactylum* gDNA and correspond to the genomic coordinate chromosome 20(+):362797-363410 for the promoter and chromosome 20(+):366255-366898 for the terminator. The nitrogen-inducible pNR promoter and native tNR terminator were amplified from *Phaeodactylum* gDNA and correspond to the genomic coordinate chromosome 20(+):362797-363410 for the promoter and chromosome 20(+):366255-366898 for the terminator. The coding sequence for the mTurquoise2 reporter protein (Goedhart et al., 2012) was amplified from plasmid L1_23 (Pollak et al., 2019) using primers LYS-09 and LYS-010 (Supplemental Information) and contained overhangs to produce a C-terminal fusion protein when translated. Plasmids were built using the Gibson assembly method and validated by Sanger sequencing.

Plasmids for Protein Expression, DSF and Crystallography

The coding sequences of full-length, N-terminally $\Delta 24$ and $\Delta 36$ deletion constructs of *PtLYSA* were cloned from *P. tricornutum* genomic DNA using primers described in Table 1 (Supplemental Information) and PrimeStar polymerase (Takara) with primers LYS13-LYS16. These fragments were sub-cloned using Gibson assembly methods into XhoI-linearized PTpBAD-CTHF vector (Brunson et al., 2018). Sequence analysis after cloning (Eurofins) revealed that some $\Delta 24$ and $\Delta 36$ clones had obtained a point mutation at the junction between the terminal F481 and thrombin cleavage linker in the vector backbone, resulting in a F481L mutation. These constructs ($\Delta 24$ -*PtLYSA*-F481L and $\Delta 36$ -*PtLYSA*-F481L) were analyzed alongside their wild-type counterparts for their level of solubility and expression.

DSF and crystallization studies utilized a DNA segment encoding *PtLYSA* (amino acid residues 39-476) was inserted in frame into pNIC28 Bsa4 (pSGC-His) using ligation independent cloning (Gileadi et al., 2008); the gene is fused to a leader

sequence encoding an TEV-cleavable N-terminal His₆ tag. The inserted DNA sequence of the corresponding domain was sequenced (Genewiz) completely to exclude the acquisition of unwanted coding changes during DNA amplification and cloning. The resultant plasmid was transformed into *E. coli* BL21 (DE3) CodonPlus RIL (Novagen). For protein preparation, 2 L of bacterial culture (Super Broth, Teknova) supplemented with 50 µg ml⁻¹ kanamycin, 100 µg ml⁻¹ chloramphenicol and 100 µL L⁻¹ antifoam 204 (Sigma), was grown at 37 °C in LEX 48 airlift bioreactors (Epiphyte3, Canada). After 7 hours of growth (*A*₆₀₀ of 2), the temperature of the cultures was reduced to 20 °C for optimal protein folding, induced with addition of 0.5 mM isopropyl-β-D-thiogalactoside (IPTG; GoldBio) and incubated overnight.

Transformation of plasmid DNA into *P. tricornutum*

The fusion constructs and control plasmids were transferred into *P. tricornutum* cells using the multi-well conjugation method (Diner et al., 2016). Briefly, plasmids were transformed into EPI-300 *E. coli* cells (Lucigen) harboring the pTA-MOB conjugation helper plasmid (Strand et al., 2014). The bacterial strains were then isolated, amplified and mixed with diatom cells to allow for conjugation of plasmids according to published methods (Diner et al., 2016). Diatom exconjugants were selected on 1/2 L1 1% agar medium supplemented with 20 µg mL⁻¹ phleomycin (Gold Biolabs) at a diel cycle of 14:10 at ~50 µE ms⁻¹ light intensity.

Microscopy

All confocal images were captured using a Leica TCS-SP5 confocal microscope (Leica Microsystems). mT2 was detected by excitation with a 458 nm laser and an emission range of 465-510 nm. Chlorophyll autofluorescence was also excited with a 458 nm laser and detected with an emission range of 680-712 nm.

Small-scale expression of PtLYSA

Clones from BL21 *E. coli* (New England Biolabs) transformation plates were picked and cultured overnight in 5 mL LB broth supplemented with 10 µg mL⁻¹ tetracycline at 30 °C and 220 rpm shaking. On the day of testing, these cultures were used to seed 10 mL cultures of Terrific Broth supplemented with 10 µg mL⁻¹ tetracycline at a 1:20 dilution. The cultures were then incubated for 4 hours at 30 °C, at which time they were then moved to 18 °C for an additional 60 minutes to cool. L-arabinose was

added to the cultures to a final concentration of 0.5% and the cells allowed to incubate overnight at 18 °C. Cultures were pelleted the following day and evaluation of protein expression was carried out as described ([dx.doi.org/10.17504/protocols.io.bkhgkt3w](https://doi.org/10.17504/protocols.io.bkhgkt3w)). Eluates and insoluble fractions were analyzed via polyacrylamide electrophoresis and gels stained with Coomassie Brilliant Blue G-250 solution for visualization of proteins.

Purification of P_tLYSA for biochemical analysis

For large-scale purification, 1 L of the $\Delta 36$ -P_tLYSA-F481L BL21 cell line was cultured and induced as described above. Cell pellets were sonicated in 500 mM NaCl, 20 mM Tris-HCl pH 8.0, 10% glycerol and purified using FPLC and a 5 mL HisTrap FF column. Large-scale protein purification from *E. coli* was performed on an ÄKTApurifier instrument (GE Healthcare) with the modules Box-900, UPC-900, R-900 and Frac-900 with all solvents filtered through a nylon membrane 0.2 μ m GDWP (Merck) prior to use. Final eluates were pooled and resuspended in 20 mM HEPES pH 8.0, 300 mM KCl, and 10% glycerol. Concentration was determined by method of Bradford.

Decarboxylation Assay

Enzyme assays for HPLC analysis were conducted in 50 mM HEPES pH 8, 200 mM KCl, and 10% glycerol, at 100 μ L scale. Substrates (DAP, ornithine and arginine) were added at 1 mM concentration and cofactor PLP was added at 50 μ M concentration. Assays were started with the addition of ~50 μ M P_tLYSA protein and allowed to incubate overnight (12-18 hours), after which reactions were quenched upon addition of assay components for Marfey's derivatization.

Marfey's derivatization and RP-HPLC analysis

Marfey's derivatization was carried out by the addition of 20 μ L concentrated Na₂CO₃ (approx. 10% w/v) and 100 μ L of freshly prepared 1% w/v 1-fluoro-2,4-dinitrophenyl-5-L-alanine amide (L-FDAA) in acetone to 50 μ L of the P_tLYSA reactions described above. For the generation of standards and H₂O blanks, 50 μ L of 1 mM aqueous amino acid stock solutions or Milli-Q H₂O was added in place of the P_tLYSA reaction mix. Marfey's derivatization reactions were allowed to incubate for 90 minutes at 37 °C before quenching with 25 μ L 1N HCl and adding 5 μ L Milli-Q

H₂O to bring the total volume to 200 μ L. Reactions were centrifuged (18,000 x g, 10 min) and 10 μ L of the clarified supernatant was injected for analytical reverse-phase HPLC (Agilent Technologies 1200 series, Phenomenex Luna 5u C18(2), 4.6 x 150 mm) at a flow rate of 1 mL min⁻¹ using the following method: 5% B (5 min), 5 – 95% B (20 min), 95 – 5% B (1 min), 5% B (4 min), where A = 0.1% aqueous trifluoroacetic acid, and B = 0.1% trifluoroacetic acid in acetonitrile (Brunson et al., 2018). Reaction products were monitored by UV-detection at 340 nm.

Protein purification of *PtLYSA* for DSF and Crystallization

Cells were harvested by centrifugation at 6,500 x g and suspended in buffer containing 20 mM HEPES (pH 7.5), 500 mM NaCl, 20 mM imidazole, 0.1% IGEPAL, 20% sucrose, 1 mM β -mercaptoethanol (β ME). Cells were disrupted by sonication and cell-debris was removed by centrifugation at 45,000 x g. The supernatants were applied to a chromatography column packed with 5 ml His60 superflow resin (Clontech) that had been equilibrated with buffer A (20 mM HEPES pH 7.5, 20 mM imidazole, 500 mM NaCl, 1 mM β ME). The column was washed with buffer A and the His₆ tagged *PtLYSA* was eluted with buffer B (20 mM HEPES pH 7.5, 350 mM NaCl, 250 mM imidazole, 1 mM β ME). The N-terminal His₆ tags from the protein was removed by overnight digestion at 4 °C with the TEV protease at a 1000:1 (w/w) ratio of *PtLYSA*:TEV. The tag-free protein was then separated from the His₆ tag and TEV protease by a Superdex 200 (16/60) size exclusion chromatography column equilibrated with buffer containing 20 mM HEPES pH 7.5, 150 mM NaCl, and 5 mM DTT. Potential peak fractions containing *PtLYSA* were assessed by SDS-PAGE, pooled concentrated to 30 mg ml⁻¹ using a 30 kDa Amicon Ultra-15 centrifugal filter device (Millipore). For crystallization experiments, the tag-free protein was purified using a Superdex 200 (16/60) gel filtration column in a buffer containing 20 mM Na-Acetate pH 5.5, 350 mM NaCl, and 1 mM β ME. Fractions containing the peak of the protein were pooled and concentrated to 20 mg/ml and buffer exchanged to Na-Acetate (pH 5.5), 150 mM NaCl, and 10 mM DTT using a 30 kDa Amicon Ultra-15 centrifugal filter device (Millipore). Purified *PtLYSA* appeared to be bright yellow, suggesting the presence of co-purified cofactor, PLP from *E. coli* host.

Differential Scanning Fluorimetry

Potential ligands (10 mM) stocks were prepared in H₂O and stored at -20 °C, and diluted to 1 mM working stock in buffer containing 20 mM HEPES, pH 7.5, 150 mM NaCl and 5 mM DTT (DSF buffer). Final DSF reaction mixtures (20 µl final volume) contained 100 µM *Pt*LYSA, 500 mM each ligand, and 5× SYPRO Orange (Invitrogen) in DSF buffer and were distributed in a 384-well PCR plate (Applied Biosystems) with 5 control wells (purified *Pt*LYSA) and 5 wells with *Pt*LYSA-ligand complexes indicated in Figure 6. The fluorescence intensities were measured using an Applied Biosystems 7900HT fast real-time PCR system with excitation at 490 nm and emission at 530 nm. The samples were heated from 25 to 99 °C at a rate of 3 °C min⁻¹. The midpoint of the unfolding transition (*T_m*) was obtained from fitting the melting curve to a Boltzmann equation (Niesen et al., 2007). Δ*T_m* for each specific ligand was calculated as the difference of the *T_m* values measured with or without a ligand (average of 5 measurements).

Protein crystallization and crystal harvesting

Initial crystallization screening of *Pt*LYSA was performed using 800 nl (protein:mother liquor=1:1) sitting drops at a concentration of 12 mg/ml (255 µM) co-crystallized with D-lysine (1 mM) with a Crystal Gryphon (Art Robbins Instruments) and utilizing MCSG sparse matrix crystallization suite (Microlytic). Crystals were eventually obtained in sitting drops by vapor diffusion against well solutions containing 2.0 M Ammonium sulfate and 0.1 M Bis-Tris pH 6.5. Crystals appeared after 12 days at 19 °C and reached maximum dimensions of 15 × 30 × 30 µm³ in another week. For data collection, crystals were cryo-preserved by addition of 30% glycerol to the mother liquor prior to flash-cooling in liquid nitrogen.

Data collection and processing, structure determination, model building, refinement and analysis

Data were collected with an Eiger 9M detector, with a wavelength of 0.98 Å, on the ID-17-1 (AMX) beamline at the National Synchrotron Light Source-II, Brookhaven National Laboratory (Table 2). Data from a single crystal were integrated and scaled using AIMLESS (Winn et al., 2011). Diffraction was consistent with the orthorhombic space group P2₁2₁2₁ (a=85.8, b= 86.3, c=127.2 Å) and extended to a resolution of 2.78 Å, with two molecules in the asymmetric unit (protomer 1/chain A and 2/chain B). Initial phases were determined by molecular replacement with

PHASER (McCoy et al., 2007) using refined coordinates of the *E. coli* DAPDC (PDB: 1KNW, Seq ID 27%) as the search model. After determining the phases an atomic model was built into the density using the automated model building program BUCCANEER (Cowtan, 2006) and manually inspected using COOT (Emsley and Cowtan, 2004). The model was refined with REFMAC5 (CCP4, 1994) and PHENIX (Liebschner et al., 2019). For structural interpretation of *PtLYSA* protomer 1 has been used. Analyses of the structures were performed in COOT and MOLPROBITY (Chen et al., 2010). Crystallographic statistics and RCSB accession codes are provided in Table 2.

Accession Numbers

This study did not generate new genomic data. Accession numbers for genes described in this work include *P. tricornutum* *LYSA*, J21592; *DAPA*, J11151; *DAPB*, J4025; *DAPL*, J22909; *DAPF*, J34852.

Supplemental Data

Supplemental Figure 1. Phylogenetic analysis of the putative diatom lysine biosynthesis pathway components.

Supplemental Figure 2. Diel expression of lysine pathway genes from Table 1 from transcriptomic analysis.

Supplemental Figure 3. Reduction of *PtLYSA* protein content in *P. tricornutum* cells via RNAi results in a lag in growth rate.

Supplemental Figure 4. Sanger sequencing of CRISPR *P. tricornutum* exconjugants.

Supplemental Figure 5. Small-scale solubility testing of *PtLYSA*-CTHF constructs in *E. coli* BL21 cells.

Supplemental Figure 6. *PtLYSA* does not act upon L-ornithine.

Supplemental Figure 7. Structure based primary sequence analyses.

Supplemental Figure 8. Dimer interface of *PtLYSA*.

Supplemental Figure 9. The LEEAA segment of *PtLYSA* maintains integrity of the active site.

ACKNOWLEDGEMENTS

This research is supported by the *Department of Energy*, Office of Biological and Environmental Research (*BER*), grant DE-SC0018344.

AUTHOR CONTRIBUTIONS

V.A.B., J.K.B., A.G., M.A.M., S.M.K.K., and C.L.D. designed research. V.A.B., J.K.B., A.G., M.A.M., E.A.G., Z.F., J.B., S.M.K.K., and C.L.D. performed research. V.A.B., J.K.B., A.G., M.A.M., E.A.G., and C.L.D. contributed new reagents/analytic tools. V.A.B., J.K.B., A.G., M.A.M., Z.F., S.M.K.K., B.S.M., A.E.A., S.C.A., and C.L.D. analyzed data. V.A.B., J.K.B., A.G., M.A.M., S.C.A., and C.L.D. wrote the article.

Figure 1. Lysine biosynthetic pathway in *P. tricornutum* based on predicted gene functions from the genome (Table 1).

Entry of aspartate metabolites into the DAP pathway begins with addition of pyruvate to A-4-S via *DAPA* to produce 4-hydroxy-tetrahydrodipicolinate (HTPA), followed by *DAPB*-mediated reduction to THDPA (2,3,4,5-tetrahydrodipicolinate). DAP aminotransferase (*DAPL*) converts THDPA to L,L-diaminopimelate (L,L-DAP). L,L-DAP is subsequently isomerized to *meso*-DAP by *DAPF* epimerization activity, with lysine produced from *meso*-DAP via decarboxylation by *LYSA*

)LJXUHBK\ORJHQHWLF DQDO\VLV RI WKH SXWDS\W\KZGLDWRP
,Q VFKHPDWLF WUHHV \$ & ERRWVWUDSDV\KSLRUVSR\QWKRZQ
\$ 0/ SK\ORJHQ\ RI GLK\GURGLDAPFRDQDQWM DQGWKDFHHQ DOJD
ZHOO DV UHG DOJDH DUH GLVWDQWO\UDHODDWHGLQVROKGRJ
VWUDPHQRSLOHV DQG GLDWRPV 7KH WRSRORHULDXQJRVWLQ
RI WKH GAPWZLPWK D SRVVLEOXanthronolidesRQVKLS WR
%0/ SK\ORJHQ\ RI GLDPLQRSPLPHSDV\W\K\G\H\FRPRJODURP JUHH
DOJDH H[FOXGLQJ ODQG SODQWV DQG FRPSVR\$K\JHDBQGR
VWUDPHQRSLOH FODGHV FOOpitulaeVerucamicrobDQVHIVRRUDQ
7KLV WRSRORJ\ ZDV FRQVLVWHQW ZLWKS%UDWHV\KRCZQ,QRUHU
EUDQFKHV GW ± GLIIHUHQW WRSRORJHV\$OYRHSQYRDXMGRG
KLVWRU\ RI WKLV HQJ\PH FDQ EH LQIHKGHWKBBW WKHOPDQV
GHILQHGH FURZQ JURXS RI FUISWRSK\WPHQKSLSWRSK\WHHV
DFWLYH VLWH ORRS LQVBiolidophyceaeLGLDWRPLFDGR WKH
&0/ SK\ORJHQGRDPLQRSPLPHODWH DAPLQRKHJGQVWUDEXH\LRQ
DapL IROORZV D SDWWHUQ FRQVLRWWHQRPPRQWK
DQFHVWRU /(&\$ IROORZHG E\ HQGRVQFEHLR\URP HUGLDWQGD
WR VWUDPHQRSLOHV DQG KDSWRSK\WHV
' &RQVHUYHG FKDUDFWHU RI WKH GLDWRP WLSMIFLQRF\$QV\KH
ELWVFRUH SORW VXPPDULJHV WKH FRQVZLWYKHGKUDH\DLG\KHU\LR
WZR LQYDULDQW SKHQ\ODODQLQHV DQGDGLGWLQV\KWRZHK\G
7KH ORJRSORW LV EDVHG RQ DQ DOLJQPHQVHRVHTGHQFVRP D
\$PLQR DFLG QXPEHULQJ EwricofurUGLQXHVQRHWKQ WKH JUH\
PDUNV WKH IODQNLQJ UHJLRQ FRQVHUYHGUBBWRVGE\6\$ K
:HEORJR &URRNV HW DO

Figure 3. *In vivo* localization of C-terminal FP-tagged *PtLYSA* constructs in *P. tricornutum*.

(A) Z-stacked confocal microscopy image of *PtLYSA*-YFP. Yellow, YFP signal; Red; Chlorophyll autofluorescence.

(B) Confocal microscopy of *PtLYSA*-mT2, chlorophyll autofluorescence channel.

(C) Confocal microscopy of *PtLYSA*-mT2, mTurquoise2 channel.

(D) Confocal microscopy of *PtLYSA*-mT2, bright field channel.

(E) Merge of B,C,D

Figure 4. TIDE sequencing analysis results for heterozygous *P. tricornutum* *PtLYSA* mutants produced using two sgRNAs (gLysA-1 and gLysA-2).

(A-C) Heterozygous mutation of an 18-bp deletion paired with a wild-type sequence for three cell lines.

(D) TIDE result for a wild-type sequence of *PtLYSA*.

Figure 5. RP-HPLC ($\lambda = 340$ nm) analyses of L-FDAA (Marfey's) derivatized *PtLYSA* reactions with *meso*-, L,L-, and D,D-DAP, and comparison to similarly derivatized lysine and DAP standards.

Substrates for enzyme reactions were added at 1 mM concentration and similarly 1 mM of each standard was used for subsequent derivatization. Overnight *PtLYSA* reactions of *meso*-DAP and D,D-DAP produced L- and D-lysine, respectively. L,L-DAP reactions also produced L-lysine, but did not proceed to completion.

Figure 6. Differential scanning fluorimetry of *Ptl*LYSA in presence of potential ligands.

(A) Denaturation of *Ptl*LYSA as a function of temperature as observed by increase in fluorescence of the indicator dye SYPRO Orange, which binds nonspecifically to hydrophobic surfaces. At higher temperatures, the intrinsic fluorescence degrades due to the formation of protein aggregates and dye dissociation.

(B) Schematic description of ligands and their calculated ΔT_m values are shown.

Figure 7. Structure of *Pt*LYSA highlighting the active site.

(A) View of the *Pt*LYSA dimer shown in ribbon representation with secondary structure elements indicated by arrows (β strands) and ribbons (helices). The two protomers are colored in light-blue (protomer 1; chain A) and light-pink (protomer 2; chain B). A transparent molecular surface envelops the structure. The sulfate ions that occupies the active sites of the protomers are shown in stick representation and labeled. The ordered active site loop in protomer 1 is colored in green. N and C denote the location of the N and C termini.

(B) Closeup view of the part of the active site in protomer 1 highlighting the bound SO_4 and the active site loop (color coded as in A). Relative locations and structural elements of active site loops in *C. glutamicum* (PDB: 5X7M, in grey), *M. tuberculosis* (PDB: 2O0T, in pink), *E. coli* (PDB: 1KNW, in dark-blue), *T. maritima* (PDB: 2YXX, in gold), *H. pylori* (PDB: 2QGH, in cyan) and *M. jannaschii* (PDB: 1TUF, in magenta) DAPDC orthologs by aligning the respective coordinates on protomer 1. PLP (transparent spheres) was modeled into the protomer 1 based on its position in the *T. maritima* structure to denote the potential location and orientation of the co-factor in the *Pt*LYSA active site.

925 **(C)** Closeup view and interactions of the SO₄ in the active site in *PtLYSA*.
926 Contributing sidechains to the active site form the protomers (color coded as A) are
927 shown in sticks and labeled. Contacts to SO₄ are shown as dashed lines. Modeled
928 PLP is shown in grey lines and transparent spheres.
929

930 **(D)** Closeup view of the active site in *PtLYSA* showing electron density from omit
931 maps (contoured at 1.2 σ ; light-blue mesh) around the SO₄ and putative D-lysine
932 (stick representation, in green). Atomic contacts in *PtLYSA* sidechains with SO₄ are
933 indicated by dashed lines. Potential sidechain contacts to D-lysine in the active site
934 are shown stick representation. All figures depicting structure were generated with
935 PyMol (DeLano et al., 2002).
936
937

Figure 8. Distinctive active site in PtLYSA.

The conserved arginine in DAPDC orthologs is replaced by a threonine, Thr317, (indicated in stick representation and color coded as in Figure 6) in *PtLYSA*. SO_4 in the active site is shown in stick representation. The connecting loop between the strand $\beta 7$ and helix $\alpha 9$ (amino acid 236-244), which forms a wall of the active site, is smaller in comparison to the orthologs with exception of *E. coli* DAPDC (see Figure S8). The smaller sidechain of Thr317 together with the smaller $\beta 7$ - $\alpha 9$ connective loop adopting a distinct conformation (indicated by an arrow) compared to its counterpart create a distinctive and larger *PtLYSA* active site (indicated by a transparent sphere) that can potentially accommodate unique substrate (D,D-DAP) for catalysis and can be exploited as specific druggable packet. Part of the active site loops in the orthologous DAPDC are shown and color coded as in Figure 6 and labeled. Modeled PLP from *T. maritima* (PDB: 2XYY) is shown in grey lines.

A

B

Figure S1. Phylogenetic analysis of the putative diatom lysine biosynthesis pathway components.

(A) ML phylogeny of dihydrodipicolinate reductase *DapB*. With moderate support, plants and green algae are more related to complex algae, including diatoms. The *DapB* homolog in this clade is likely related to delta-proteobacteria or Fibrobacteriales (note the long branch of the latter group). In this schematic, bootstrap support is shown for critical branching points of the outgroups, while >85 branch support is shown in the Bolidophyceae-diatom clade.

(B) ML phylogeny of diaminopimelate epimerase *DapF*. This enzyme is apparently derived from cyanobacteria and in plants and green algae, red algae, stramenopiles and haptophytes constitutes a monophyletic clade, with red algae being moderately supported as the donor of the complex algal homolog. *Paulinella* are rhizarian algae that established a primary endosymbiotic relationship with a cyanobacterium independently on other algal lineages, as apparent from the figure. In this schematic,

1006 bootstrap support is shown for critical branching points of the outgroups, while >85
1007 branch support is shown in the Bolidophyceae-diatom clade.
1008

Figure S2. Diel expression of *P. tricornutum* lysine pathway genes from Table 1 from transcriptomic analysis.

The black bar above represents the dark portion of the day. Adapted from Smith et al., 2016.

Figure S3. Reduction of *PtLYSA* protein content in *P. tricornutum* cells via RNAi results in a lag in growth rate.

(A) Western blot of *PtLYSA* protein in uninduced RNAi conditions (NH₄-repressed) comparing wild-type (WT) versus transgenic cell line (P2-7)

(B) Western blot of *PtLYSA* protein in induced RNAi conditions (NO₃-active) comparing wild-type (WT) versus transgenic cell line (P2-7)

(C) RNAi-induced vs -uninduced growth curve of P2-7 cell line in duplicate. X-axis, time in days; Y-axis, chlorophyll content of batch cultures measured via fluorimetry

Figure S4. Sanger sequencing of CRISPR *P. tricornutum* exconjugants.

Green arrows in schematic at top represent the targeted sites within *PtLYSA* gene and red bar represents expected 18-bp deletion upon editing with two sgRNAs.

Figure S5. Small-scale solubility testing of *PtLYSA*-CTHF constructs in *E. coli* BL21 cells.

Protein expression was induced with 0.5% L-arabinose in Terrific Broth at 30 °C for 16 hours. The arrow indicates the $\Delta 36$ -*PtLYSA*-F481L-CTHF construct employed for biochemical analyses. In, clarified input; E, eluate; FL, full-length protein; EV, empty vector control.

Figure S6. *PtLYSA* does not act upon L-ornithine.

RP-HPLC ($\lambda = 340$ nm) analyses of L-FDAA (Marfey's) derivatized *PtLYSA* reactions with L-ornithine and comparison to similarly derivatized ornithine (time ~21.5 min) and putrescine (time ~23.2 min) standards. Substrates for enzyme reactions were added at 1 mM concentration and similarly 1 mM of each standard was used for subsequent derivatization.

Figure S7. Structure based primary sequence analyses.

Aligned amino acid sequences for PtLYSA and DAPDC from *C. glutamicum* (PDB: 5X7M), *M. tuberculosis* (PDB: 2O0T), *E. coli* (PDB: 1KNW), *T. maritima* (PDB: 2YXX), *H. pylori* (PDB: 2QGH) and *M. jannaschii* (PDB: 1TUF). PtLYSA secondary structure elements (helices as cylinders; strands as arrows) are color coded as in Figure 6 above the sequence. Alignment gaps indicated by (-). Sidechain identity/similarity is denoted by red shading (conserved in all) or red letter (conserved in most). Location of Thr317, the $\beta 7$ - $\alpha 9$ connective and active site loops are outlined. Conserved lysine that forms Schiff's base with PLP highlighted in green shade. Light-blue and light-pink asterisks above the alignment indicate conserved active site residues supplied in *cis* and *trans*, respectively. Green asterisks above denote PtLYSA residues adjacent to putative D-lysine. Three key binding motifs are indicated below the alignment.

Figure S8. Dimer interface of *PtLYSA*.

Structure of *PtLYSA* is shown in surface representation and the protomers are color coded as in Figure 6. The interface was mapped using 1.4 Å probe radius and colored in yellow. Position of the active site loop in protomer 1 is indicated by arrows and labeled.

Figure S9. The LEEAA segment of *PtLYSA* maintains integrity of the active site.

A view of the interactions between the LEEAA (part of helix a6; aa 72-76) and the segment that demarcates the active site in protomer 1 (chain A; color coded as in Figure 6). Stick models of the amino acids that comprise the interface and the active site are shown; the bound SO_4 and water in the active site are shown in stick representation and red sphere, respectively. Part of PLP (transparent spheres and lines) was modeled into the active site based on its position in the *C. glutamicum* structure (PDB: 5X7M) to denote the potential location and orientation of the co-factor in *PtLYSA*. Potential hydrogen bonds and van der Waals interactions are represented as black and red dashed lines, respectively. The interaction of the LEEAA segment at the interface highlights its potential role in maintaining active site integrity.

Pt Gene ID	Function	EC ID	Ancestor	Localization
Phatr3_J11151	<i>DAPA</i> , 4-hydroxytetrahydropicolinate synthase	4.3.3.7	<i>Vibrio</i>	Unclear
Phatr3_J4025	<i>DAPB</i> , 4-hydroxytetrahydropicolinate reductase	1.17.1.8	Plant	Chloroplast
Phatr3_J22909	<i>DAPL</i> , L,L-diaminopimelate aminotransferase	2.6.1.83	Plant	Chloroplast
Phatr3_J34582	<i>DAPF</i> , diaminopimelate epimerase	5.1.1.7	Plant	Chloroplast
Phatr3_J21592	<i>LYSA</i> , diaminopimelate decarboxylase	4.1.1.20	<i>Verrucamicrobia</i>	Chloroplast

Table 1. *P. tricornutum* lysine biosynthetic pathway protein IDs, phylogenetic outgroup, and predicted subcellular localization.

1128

Gene ID	PHATRDRAFT_21592
Co-Crystallized Ligand	D-lysine
Data Collection*	
Source	BNL ID-17-1
Wavelength (Å)	0.98
Number of crystals	1
Space group	P2 ₁ 2 ₁ 2 ₁
Cell dimensions	
a,b,c (Å)	85.8 86.3 127.2
Resolution (Å)	19-2.78 (2.88-2.78)
Completeness (%)	99.4 (98.6)
Total reflections	329040
Unique reflections	24274
Wilson B-factor	29.9
Multiplicity	13.6 (13.4)
R _{merge} (%)	25.7 (85.63)
CC _{1/2} (%)	99.1 (86.1)
CC* (%)	99.8 (96.2)
<l>/s(l)	10.2 (3.0)
Refinement*	
Reflections: work/free	24270 (2349)/1304 (131)
R _{work} /R _{free} (%)	19.8 (26.2)/24.4 (30.1)
Number of TLS groups	14
Number of atoms	
Protein	6479
Ligand	35
Water	49
Average B-factors (Å ²)	
Protein	23.4
Ligand	34.32
Water	19.5
r.m.s.d.	
Bond lengths (Å)	0.003
Bond angles (°)	0.6
Molprobit†	
Favored	98.3% (652 aa)
Allowed	100% (867 aa)
Outliers	none
Clash score	99 th percentile
Molprobit score	98 th percentile
PDB ID Code	7JPJ

1129

1130 *Statistics calculated using PHENIX (Adam et al., 2010); highest resolution shells indicated in
 1131 parentheses

1132 †Calculated with the program MolProbity (Chen et al., 2010)

1133

1134 **Table 2.** Crystallographic data collection and refinement statistics

1135

1136

CITATIONS

- Allen, A. E., C. L. Dupont, M. Oborník, A. Horák, A. Nunes-Nesi, J. P. McCrow, H. Zheng, D. A. Johnson, H. Hu, A. R. Fernie and C. Bowler (2011). Evolution and metabolic significance of the urea cycle in photosynthetic diatoms. Nature 473: 203.
- Apt, K. E., A. R. Grossman and P. G. Kroth-Pancic (1996). Stable nuclear transformation of the diatom *Phaeodactylum tricornutum*. Molecular and General Genetics MGG 252(5): 572-579.
- Apt, K. E., L. Zaslavkaia, J. C. Lippmeier, M. Lang, O. Kilian, R. Wetherbee, A. R. Grossman and P. G. Kroth (2002). *In vivo* characterization of diatom multipartite plastid targeting signals. Journal of Cell Science 115(21): 4061-4069.
- Armbrust, E. V., J. A. Berges, C. Bowler, B. R. Green, D. Martinez, N. H. Putnam, et al. (2004). The Genome of the Diatom *Thalassiosira Pseudonana*: Ecology, Evolution, and Metabolism. Science 306(5693): 79.
- Armbrust, E. V. (2009). The life of diatoms in the world's oceans. Nature 459(7244): 185-192.
- Asada, Y., K. Tanizawa, S. Sawada, T. Suzuki, H. Misono and K. Soda (1981). Stereochemistry of meso-alpha,epsilon-diaminopimelate decarboxylase reaction: the first evidence for pyridoxal 5'-phosphate dependent decarboxylation with inversion of configuration. Biochemistry 20(24): 6881-6886.
- Baier, M. and K.-J. Dietz (2005). Chloroplasts as source and target of cellular redox regulation: a discussion on chloroplast redox signals in the context of plant physiology. Journal of Experimental Botany 56(416): 1449-1462.
- Bhushan, R. and H. Brückner (2004). Marfey's reagent for chiral amino acid analysis: A review. Amino Acids 27(3): 231-247.

1170 Bowler, C., A. E. Allen, J. H. Badger, J. Grimwood, K. Jabbari, A. Kuo, et al. (2008).
 1171 The *Phaeodactylum* genome reveals the evolutionary history of diatom genomes.
 1172 Nature 456: 239.
 1173

1174 Bowler, C., A. Vardi and A. E. Allen (2009). "Oceanographic and Biogeochemical
 1175 Insights from Diatom Genomes." Annual Review of Marine Science 2(1): 333-365.
 1176

1177 Bromke, M. A. (2013). Amino Acid biosynthesis pathways in diatoms. Metabolites
 1178 3(2): 294-311.
 1179

1180 Brunson, J. K., S. M. K. McKinnie, J. R. Chekan, J. P. McCrow, Z. D. Miles, E. M.
 1181 Bertrand, V. A. Bielinski, H. Luhavaya, M. Oborník, G. J. Smith, D. A. Hutchins, A. E.
 1182 Allen and B. S. Moore (2018). Biosynthesis of the neurotoxin domoic acid in a bloom-
 1183 forming diatom. Science 361(6409): 1356-1358.
 1184

1185 Buchfink, B., C. Xie and D. H. Huson (2015). Fast and sensitive protein alignment
 1186 using DIAMOND. Nature Methods 12(1): 59-60.
 1187

1188 Bukhari, A. I. and A. L. Taylor (1971). Genetic Analysis of Diaminopimelic Acid- and
 1189 Lysine-Requiring Mutants of *Escherichia coli*. Journal of Bacteriology 105(3): 844-
 1190 854.
 1191

1192 Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: a tool for
 1193 automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics
 1194 (Oxford, England), 25(15), 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>
 1195

1196 Chang, Y.-F. (1978). Lysine metabolism in the rat brain: the pipecolic acid-forming
 1197 pathway. Journal of Neurochemistry 30(2): 347-354.
 1198

1199 CCP4 (1994). The CCP4 suite: programs for protein crystallography. Acta Crystallogr
 1200 D Biol Crystallogr 50: 760-763.
 1201

1202 Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral,
 1203 G.J., Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-

atom structure validation for macromolecular crystallography. Acta Crystallogr D Biol Crystallogr 66: 12-21.

Chu, L., D. Ewe, C. Rio Bartulos, P. G. Kroth and A. Gruber (2016). Rapid induction of GFP expression by the nitrate reductase promoter in the diatom *Phaeodactylum tricornutum*. PeerJ 4: e2344.

Cowtan, K. (2006). The Buccaneer software for automated model building. 1. Tracing protein chains. Acta Crystallogr D Biol Crystallogr 62: 1002-1011.

Crooks, G. E., Hon, G., Chandonia, J. M., & Brenner, S. E. (2004). WebLogo: a sequence logo generator. Genome Research, 14(6), 1188–1190.
<https://doi.org/10.1101/gr.849004>

Crowther, J. M., Cross, P. J., Oliver, M. R., Leeman, M. M., Bartl, A. J., Weatherhead, A. W., North, R. A., Donovan, K. A., Griffin, M., Suzuki, H., Hudson, A. O., Kasanmascheff, M., & Dobson, R. (2019). Structure-function analyses of two plant *meso*-diaminopimelate decarboxylase isoforms reveal that active-site gating provides stereochemical control. The Journal of Biological Chemistry, 294(21), 8505–8515. <https://doi.org/10.1074/jbc.RA118.006825>

De Lencastre, H., Wu, S.W., Pinho, M.G., Ludovice, A.M., Filipe, S., Gardete, S., Sobral, R., Gill, S., Chung, M., and Tomasz, A. (1999). Antibiotic resistance as a stress response: complete sequencing of a large number of chromosomal loci in *Staphylococcus aureus* strain COL that impact on the expression of resistance to methicillin. Microb Drug Resist 5:163-175.

De Riso, V., Raniello, R., Maumus, F., Rogato, A., Bowler, C., & Falciatore, A. (2009). Gene silencing in the marine diatom *Phaeodactylum tricornutum*. Nucleic Acids Research, 37(14), e96. <https://doi.org/10.1093/nar/gkp448>

DeLano, W.L. (2002). The PyMOL Molecular Graphics System. DeLano Scientific; San Carlos, CA

1238 Dewey, D. L., & Work, E. (1952). Diaminopimelic acid decarboxylase. Nature,
1239 169(4300), 533–534. <https://doi.org/10.1038/169533a0>
1240

1241 Diner, R. E., V. A. Bielinski, C. L. Dupont, A. E. Allen and P. D. Weyman (2016).
1242 Refinement of the Diatom Episome Maintenance Sequence and Improvement of
1243 Conjugation-Based DNA Delivery Methods. Frontiers in bioengineering and
1244 biotechnology 4: 65-65.
1245

1246 Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular
1247 graphics. Acta Crystallogr D Biol Crystallogr 60: 2126-2132.
1248

1249 Gibson, D. G., G. A. Benders, K. C. Axelrod, J. Zaveri, M. A. Algire, M. Moodie, M. G.
1250 Montague, J. C. Venter, H. O. Smith and C. A. Hutchison (2008). One-step assembly
1251 in yeast of 25 overlapping DNA fragments to form a complete synthetic *Mycoplasma*
1252 *genitalium* genome. Proceedings of the National Academy of Sciences
1253 105(51):20404.
1254

1255 Gileadi, O., Burgess-Brown, N.A., Colebrook, S.M., Berridge, G., Savitsky, P., Smee,
1256 C.E., Loppnau, P., Johansson, C., Salah, E., and Pantic, N.H. (2008). High
1257 throughput production of recombinant human proteins for crystallography. Methods
1258 Mol Biol 426; 221-246.
1259

1260 Goedhart, J., D. von Stetten, M. Noirclerc-Savoye, M. Lelimosin, L. Joosen, M. A.
1261 Hink, L. van Weeren, T. W. J. Gadella and A. Royant (2012). Structure-guided
1262 evolution of cyan fluorescent proteins towards a quantum yield of 93%. Nature
1263 Communications 3(1): 751.
1264

1265 Gokulan, K., B. Rupp, M. S. Pavelka, W. R. Jacobs and J. C. Sacchettini (2003).
1266 Crystal Structure of *Mycobacterium tuberculosis* Diaminopimelate Decarboxylase, an
1267 Essential Enzyme in Bacterial Lysine Biosynthesis. Journal of Biological Chemistry
1268 278(20): 18588-18596
1269

1270 Griffin, M. D., J. M. Billakanti, A. Wason, S. Keller, H. D. Mertens, S. C. Atkinson, R.
1271 C. Dobson, M. A. Perugini, J. A. Gerrard and F. G. Pearce (2012). Characterisation

of the first enzymes committed to lysine biosynthesis in *Arabidopsis thaliana*. PLoS One 7(7): e40318.

Gruber, A. and P. G. Kroth (2017). Intracellular metabolic pathway distribution in diatoms and tools for genome-enabled experimental diatom research. Philos Trans R Soc Lond B Biol Sci 372(1728).

Guillou, L., M.-J. Chrétiennot-Dinet, L. K. Medlin, H. Claustre, S. L.-d. Goër and D. Vaultot (1999). Bolidomonas: A new genus with two species belonging to a new algal class, the Bolidophyceae (Heterokonta). Journal of Phycology 35(2): 368-381.

Hor, L., M. G. Peverelli, M. A. Perugini and C. A. Hutton (2013). A new robust kinetic assay for DAP epimerase activity. Biochimie 95(10): 1949-1953.

Hu, T., D. Wu, J. Chen, J. Ding, H. Jiang and X. Shen (2008). The Catalytic Intermediate Stabilized by a “Down” Active Site Loop for Diaminopimelate Decarboxylase from *Helicobacter pylori*: enzymatic characterization with crystal structure analysis. Journal of Biological Chemistry 283(30): 21284-21293

Hudson, A. O., C. Bless, P. Macedo, S. P. Chatterjee, B. K. Singh, C. Gilvarg and T. Leustek (2005). Biosynthesis of lysine in plants: evidence for a variant of the known bacterial pathways. Biochim Biophys Acta 1721(1-3): 27-36.

Hudson, A. O., B. K. Singh, T. Leustek and C. Gilvarg (2006). An LL-diaminopimelate aminotransferase defines a novel variant of the lysine biosynthesis pathway in plants. Plant Physiol 140(1): 292-301.

Hutton, C. A., M. A. Perugini and J. A. Gerrard (2007). Inhibition of lysine biosynthesis: an evolving antibiotic strategy. Molecular BioSystems 3(7): 458-465.

Jander G, and Joshji, V. Aspartate-derived Amino Acid Biosynthesis in *Arabidopsis thaliana*. The Arabidopsis Book. 2009;e0121(10.1199/tab.0121):15.

1305 Karas, B. J., R. E. Diner, S. C. Lefebvre, J. McQuaid, A. P. R. Phillips, C. M.
 1306 Noddings, J. K. Brunson, R. E. Valas, T. J. Deerinck, J. Jablanovic, J. T. F. Gillard, K.
 1307 Beerli, M. H. Ellisman, J. I. Glass, C. A. Hutchison Iii, H. O. Smith, J. C. Venter, A. E.
 1308 Allen, C. L. Dupont and P. D. Weyman (2015). Designer diatom episomes delivered
 1309 by bacterial conjugation. Nature Communications 6: 6925.
 1310
 1311 Katoh, K. and D. M. Standley (2013). MAFFT Multiple Sequence Alignment Software
 1312 Version 7: Improvements in Performance and Usability. Molecular Biology and
 1313 Evolution 30(4): 772-780.
 1314
 1315 Kelland, J. G., M. M. Palcic, M. A. Pickard and J. C. Vederas (1985). Stereochemistry
 1316 of lysine formation by meso-diaminopimelate decarboxylase from wheat germ: use of
 1317 proton-carbon-13 NMR shift correlation to detect stereospecific deuterium labeling.
 1318 Biochemistry 24(13): 3263-3267.
 1319
 1320 Kern, A. D., M. A. Oliveira, P. Coffino and M. L. Hackert (1999). Structure of
 1321 mammalian ornithine decarboxylase at 1.6A resolution: stereochemical implications
 1322 of PLP-dependent amino acid decarboxylases. Structure 7(5): 567-581.
 1323
 1324 Kidron, H., S. Repo, M. S. Johnson and T. A. Salminen (2006). Functional
 1325 Classification of Amino Acid Decarboxylases from the Alanine Racemase Structural
 1326 Family by Phylogenetic Studies. Molecular Biology and Evolution 24(1): 79-89.
 1327
 1328 Kilian, O. and P. G. Kroth (2005). Identification and characterization of a new
 1329 conserved motif within the presequence of proteins targeted into complex diatom
 1330 plastids. Plant J 41(2): 175-183.
 1331
 1332 Kirma, M., W. L. Araújo, A. R. Fernie and G. Galili (2012). The multifaceted role of
 1333 aspartate-family amino acids in plant metabolism. Journal of Experimental Botany
 1334 63(14): 4995-5001.
 1335
 1336 Knorr, S., M. Sinn, D. Galetskiy, R. M. Williams, C. Wang, N. Müller, O. Mayans, D.
 1337 Schleheck and J. S. Hartig (2018). Widespread bacterial lysine degradation

proceeding via glutarate and L-2-hydroxyglutarate. Nature Communications 9(1):
5071.

Kosuge, T. and T. Hoshino (1998). Lysine is synthesized through the α -aminoadipate
pathway in *Thermus thermophilus*. FEMS Microbiology Letters 169(2): 361-367.

Lartillot, N., Rodrigue, N., Stubbs, D., & Richer, J. (2013). PhyloBayes MPI:
phylogenetic reconstruction with infinite mixtures of profiles in a parallel environment.
Systematic Biology 62(4), 611–615. <https://doi.org/10.1093/sysbio/syt022>

Lauritano, C., I. Orefice, G. Procaccini, G. Romano and A. Ianora (2015). Key genes
as stress indicators in the ubiquitous diatom *Skeletonema marinoi*. BMC Genomics
16(1): 411-411.

Liebschner, D., Afonine, P.V., Baker, M.L., Bunkoczi, G., Chen, V.B., Croll, T.I.,
Hintze, B., Hung, L.W., Jain, S., McCoy, A.J., et al. (2019). Macromolecular structure
determination using X-rays, neutrons and electrons: recent developments in Phenix.
Acta Crystallogr D Struct Biol 75, 861-877.

McCoy, A. J., N. E. Adams, A. O. Hudson, C. Gilvarg, T. Leustek and A. T. Maurelli
(2006). L,L-diaminopimelate aminotransferase, a trans-kingdom enzyme shared by
Chlamydia and plants for synthesis of diaminopimelate/lysine. Proceedings of the
National Academy of Sciences 103(47): 17909-17914.

McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and
Read, R.J. (2007). Phaser crystallographic software. J Appl Crystallogr 40: 658-674.

Mills, W. R., P. J. Lea and B. J. Miflin (1980). Photosynthetic Formation of the
Aspartate Family of Amino Acids in Isolated Chloroplasts. Plant Physiology 65(6):
1166-1172.

Mock, T., R. P. Otiilar, J. Strauss, M. McMullan, P. Paajanen, J. Schmutz, et al.
(2017). Evolutionary genomics of the cold-adapted diatom *Fragilariopsis cylindrus*.
Nature 541(7638): 536-540.

1372

1373 Moosburner, M. A., P. Gholami, J. K. McCarthy, M. Tan, V. A. Bielinski and A. E.
 1374 Allen (2020). Multiplexed Knockouts in the Model Diatom *Phaeodactylum* by
 1375 Episomal Delivery of a Selectable Cas9. Frontiers in Microbiology 11: 5.

1376

1377 Nguyen, L.-T., H. A. Schmidt, A. von Haeseler and B. Q. Minh (2014). IQ-TREE: A
 1378 Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood
 1379 Phylogenies. Molecular Biology and Evolution 32(1): 268-274.

1380

1381 Niesen, F.H., Berglund, H., and Vedadi, M. (2007). The use of differential scanning
 1382 fluorimetry to detect ligand interactions that promote protein stability. Nat Protoc 2:
 1383 2212-2221.

1384

1385 Perl, A., O. Shaul and G. Galili (1992). Regulation of lysine synthesis in transgenic
 1386 potato plants expressing a bacterial dihydrodipicolinate synthase in their chloroplasts.
 1387 Plant Molecular Biology 19(5): 815-823.

1388

1389 Peverelli, M. G. and M. A. Perugini (2015). An optimized coupled assay for
 1390 quantifying diaminopimelate decarboxylase activity. Biochimie 115: 78-85.

1391

1392 Peverelli, M. G., T. P. Soares da Costa, N. Kirby and M. A. Perugini (2016).
 1393 Dimerization of Bacterial Diaminopimelate Decarboxylase Is Essential for Catalysis.
 1394 Journal of Biological Chemistry 291(18): 9785-9795.

1395

1396 Pollak, B., A. Cerda, M. Delmans, S. Álamos, T. Moyano, A. West, R. A. Gutiérrez,
 1397 N. J. Patron, F. Federici and J. Haseloff (2019). Loop assembly: a simple and open
 1398 system for recursive fabrication of DNA circuits. New Phytologist 222(1): 628-640.

1399

1400 Poulsen, N. and N. Kröger (2005). A new molecular tool for transgenic diatoms. The
 1401 FEBS Journal 272(13): 3413-3423.

1402

1403 Pratelli, R. and G. Pilot (2014). Regulation of amino acid metabolic enzymes and
 1404 transporters in plants. Journal of Experimental Botany 65(19): 5535-5556.

1405

Price, M. N., G. M. Zane, J. V. Kuehl, R. A. Melnyk, J. D. Wall, A. M. Deutschbauer and A. P. Arkin (2018). Filling gaps in bacterial amino acid biosynthesis pathways with high-throughput genetics. PLOS Genetics 14(1): e1007147.

Ray, S. S., J. B. Bonanno, K. R. Rajashankar, M. G. Pinho, G. He, H. De Lencastre, A. Tomasz and S. K. Burley (2002). Cocystal Structures of Diaminopimelate Decarboxylase: Mechanism, Evolution, and Inhibition of an Antibiotic Resistance Accessory Factor. Structure 10(11): 1499-1508.

Raymond, J. A. and H. J. Kim (2012). Possible role of horizontal gene transfer in the colonization of sea ice by algae. PloS one 7(5): e35968-e35968.

Reyes-Prieto, A. and A. Moustafa (2012). Plastid-localized amino acid biosynthetic pathways of Plantae are predominantly composed of non-cyanobacterial enzymes. Sci Rep 2: 955.

Rosenwasser, S., S. Graff van Creveld, D. Schatz, S. Malitsky, O. Tzfadia, A. Aharoni, Y. Levin, A. Gabashvili, E. Feldmesser and A. Vardi (2014). Mapping the diatom redox-sensitive proteome provides insight into response to nitrogen stress in the marine environment. Proc Natl Acad Sci U S A 111(7): 2740-2745.

Sagong, H. Y. and K. J. Kim (2017). Structural basis for redox sensitivity in *Corynebacterium glutamicum* diaminopimelate epimerase: an enzyme involved in l-lysine biosynthesis. Sci Rep 7: 42318.

Savitsky, P., J. Bray, C. D. O. Cooper, B. D. Marsden, P. Mahajan, N. A. Burgess-Brown and O. Gileadi (2010). High-throughput production of human proteins for crystallization: The SGC experience. Journal of Structural Biology 172(1): 3-13.

Siaut, M., M. Heijde, M. Mangogna, A. Montsant, S. Coesel, A. Allen, A. Manfredonia, A. Falciatore and C. Bowler (2007). Molecular toolbox for studying diatom biology in *Phaeodactylum tricornutum*. Gene 406(1): 23-35.

Smith, S. R., J. T. Gillard, A. B. Kustka, J. P. McCrow, J. H. Badger, H. Zheng, A. M. New, C. L. Dupont, T. Obata, A. R. Fernie and A. E. Allen (2016). Transcriptional Orchestration of the Global Cellular Response of a Model Pennate Diatom to Diel Light Cycling under Iron Limitation. PLoS Genet 12(12): e1006490.

Smith, S. R., C. L. Dupont, J. K. McCarthy, J. T. Broddrick, M. Oborník, A. Horák, Z. Füßy, J. Cihlář, S. Kleessen, H. Zheng, J. P. McCrow, K. K. Hixson, W. L. Araújo, A. Nunes-Nesi, A. Fernie, Z. Nikoloski, B. O. Palsson and A. E. Allen (2019). Evolution and regulation of nitrogen flux through compartmentalized metabolic networks in a marine diatom. Nature Communications 10(1): 4552.

Son, H. F. and K.-J. Kim (2018). Structural basis for substrate specificity of meso-diaminopimelic acid decarboxylase from *Corynebacterium glutamicum*. Biochemical and Biophysical Research Communications 495(2): 1815-1821.

Steinegger, M. and J. Söding (2017). MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. Nature Biotechnology 35(11): 1026-1028.

Strand, T. A., R. Lale, K. F. Degnes, M. Lando and S. Valla (2014). A New and Improved Host-Independent Plasmid System for RK2-Based Conjugal Transfer. PLOS ONE 9(3): e90372.

Sugio, S., G. A. Petsko, J. M. Manning, K. Soda and D. Ringe (1995). Crystal Structure of a D-Amino Acid Aminotransferase: How the Protein Controls Stereoselectivity. Biochemistry 34(30): 9661-9669.

Velasco, A. M., J. I. Leguina and A. Lazcano (2002). Molecular Evolution of the Lysine Biosynthetic Pathways. Journal of Molecular Evolution 55(4): 445-449.

Wallsgrave, R. M. and M. Mazelis (1980). The enzymology of lysine biosynthesis in higher plants: Complete localization of the regulatory enzyme dihydrodipicolinate synthase in the chloroplasts of spinach leaves. FEBS Letters 116(2): 189-192.

1473 Wang, H.-C., B. Q. Minh, E. Susko and A. J. Roger (2017). Modeling Site
1474 Heterogeneity with Posterior Mean Site Frequency Profiles Accelerates Accurate
1475 Phylogenomic Estimation. Systematic Biology 67(2): 216-235.

1476

1477 Weyand, S., Kefala, G., Svergun, D.I., and Weiss, M.S. (2009). The three-
1478 dimensional structure of diaminopimelate decarboxylase from *Mycobacterium*
1479 *tuberculosis* reveals a tetrameric enzyme organisation. J Struct Funct Genomics 10:
1480 209-217.

1481

1482 Weyman, P. D., K. Beerli, S. C. Lefebvre, J. Rivera, J. K. McCarthy, A. L. Heuberger,
1483 G. Peers, A. E. Allen and C. L. Dupont (2015). Inactivation of *Phaeodactylum*
1484 *tricornutum* urease gene using transcription activator-like effector nuclease-based
1485 targeted mutagenesis. Plant Biotechnology Journal 13(4): 460-470.

1486

1487 White, P. J. and B. Kelly (1965). Purification and properties of diaminopimelate
1488 decarboxylase from *Escherichia coli*. The Biochemical Journal 96(1): 75-84.

1489

1490 Winn, M.D., Ballard, C.C., Cowtan, K.D., Dodson, E.J., Emsley, P., Evans, P.R.,
1491 Keegan, R.M., Krissinel, E.B., Leslie, A.G., McCoy, A., *et al.* (2011). Overview of the
1492 CCP4 suite and current developments. Acta Crystallogr D Biol Crystallogr 67: 235-
1493 242.

1494

1495 Xu, H., B. Andi, J. Qian, A. H. West and P. F. Cook (2006). The α -amino adipate
1496 pathway for lysine biosynthesis in fungi. Cell Biochemistry and Biophysics 46(1): 43-
1497 64.

1498

1499 Zhu, X. and G. Galili (2004). Lysine Metabolism Is Concurrently Regulated by
1500 Synthesis and Catabolism in Both Reproductive and Vegetative Tissues. Plant
1501 Physiology 135(1): 129.

Parsed Citations

Allen, A. E., C. L. Dupont, M. Oborník, A. Horák, A. Nunes-Nesi, J. P. McCrow, H. Zheng, D. A. Johnson, H. Hu, A. R. Fernie and C. Bowler (2011). Evolution and metabolic significance of the urea cycle in photosynthetic diatoms. *Nature* 473: 203.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Apt, K. E., A. R. Grossman and P. G. Kroth-Pancic (1996). Stable nuclear transformation of the diatom *Phaeodactylum tricornutum*. *Molecular and General Genetics MGG* 252(5): 572-579.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Apt, K. E., L. Zaslavkaia, J. C. Lippmeier, M. Lang, O. Kilian, R. Wetherbee, A. R. Grossman and P. G. Kroth (2002). In vivo characterization of diatom multipartite plastid targeting signals. *Journal of Cell Science* 115(21): 4061-4069.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Armbrust, E. V., J. A. Berges, C. Bowler, B. R. Green, D. Martinez, N. H. Putnam, et al. (2004). The Genome of the Diatom *Thalassiosira pseudonana*: Ecology, Evolution, and Metabolism. *Science* 306(5693): 79.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Armbrust, E. V. (2009). The life of diatoms in the world's oceans. *Nature* 459(7244): 185-192.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Asada, Y., K. Tanizawa, S. Sawada, T. Suzuki, H. Misono and K. Soda (1981). Stereochemistry of meso-alpha,epsilon-diaminopimelate decarboxylase reaction: the first evidence for pyridoxal 5'-phosphate dependent decarboxylation with inversion of configuration. *Biochemistry* 20(24): 6881-6886.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Baier, M. and K.-J. Dietz (2005). Chloroplasts as source and target of cellular redox regulation: a discussion on chloroplast redox signals in the context of plant physiology. *Journal of Experimental Botany* 56(416): 1449-1462.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bhushan, R. and H. Brückner (2004). Marfey's reagent for chiral amino acid analysis: A review. *Amino Acids* 27(3): 231-247.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bowler, C., A. E. Allen, J. H. Badger, J. Grimwood, K. Jabbari, A. Kuo, et al. (2008). The *Phaeodactylum* genome reveals the evolutionary history of diatom genomes. *Nature* 456: 239.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bowler, C., A. Vardi and A. E. Allen (2009). "Oceanographic and Biogeochemical Insights from Diatom Genomes." *Annual Review of Marine Science* 2(1): 333-365.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bromke, M. A. (2013). Amino Acid biosynthesis pathways in diatoms. *Metabolites* 3(2): 294-311.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Brunson, J. K., S. M. K. McKinnie, J. R. Chekan, J. P. McCrow, Z. D. Miles, E. M. Bertrand, V. A. Bielinski, H. Luhavaya, M. Oborník, G. J. Smith, D. A. Hutchins, A. E. Allen and B. S. Moore (2018). Biosynthesis of the neurotoxin domoic acid in a bloom-forming diatom. *Science* 361(6409): 1356-1358.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Buchfink, B., C. Xie and D. H. Huson (2015). Fast and sensitive protein alignment using DIAMOND. *Nature Methods* 12(1): 59-60.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bukhari, A. I. and A. L. Taylor (1971). Genetic Analysis of Diaminopimelic Acid- and Lysine-Requiring Mutants of *Escherichia coli*. *Journal of Bacteriology* 105(3): 844-854.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* (Oxford, England), 25(15), 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chang, Y.-F. (1978). Lysine metabolism in the rat brain: the pipecolic acid-forming pathway. *Journal of Neurochemistry* 30(2): 347-354.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

CCP4 (1994). The CCP4 suite: programs for protein crystallography. *Acta Crystallogr D Biol Crystallogr* 50: 760-763.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J., Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Crystallogr D Biol Crystallogr* 66: 12-21.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chu, L., D. Ewe, C. Rio Bartulos, P. G. Kroth and A. Gruber (2016). Rapid induction of GFP expression by the nitrate reductase promoter in the diatom *Phaeodactylum tricornutum*. *PeerJ* 4: e2344.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cowtan, K. (2006). The Buccaneer software for automated model building. 1. Tracing protein chains. *Acta Crystallogr D Biol Crystallogr* 62: 1002-1011.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Crooks, G. E., Hon, G., Chandonia, J. M., & Brenner, S. E. (2004). WebLogo: a sequence logo generator. *Genome Research*, 14(6), 1188–1190. <https://doi.org/10.1101/gr.849004>

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Crowther, J. M., Cross, P. J., Oliver, M. R., Leeman, M. M., Bartl, A. J., Weatherhead, A. W., North, R. A., Donovan, K. A., Griffin, M., Suzuki, H., Hudson, A. O., Kasanmascheff, M., & Dobson, R. (2019). Structure-function analyses of two plant meso-diaminopimelate decarboxylase isoforms reveal that active-site gating provides stereochemical control. *The Journal of Biological Chemistry*, 294(21), 8505–8515. <https://doi.org/10.1074/jbc.RA118.006825>

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

De Lencastre, H., Wu, S.W., Pinho, M.G., Ludovice, A.M., Filipe, S., Gardete, S., Sobral, R., Gill, S., Chung, M., and Tomasz, A. (1999). Antibiotic resistance as a stress response: complete sequencing of a large number of chromosomal loci in *Staphylococcus aureus* strain COL that impact on the expression of resistance to methicillin. *Microb Drug Resist* 5:163-175.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

De Riso, V., Raniello, R., Maumus, F., Rogato, A., Bowler, C., & Falcatore, A. (2009). Gene silencing in the marine diatom *Phaeodactylum tricornutum*. *Nucleic Acids Research*, 37(14), e96. <https://doi.org/10.1093/nar/gkp448>

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

DeLano, W.L. (2002). The PyMOL Molecular Graphics System. DeLano Scientific; San Carlos, CA

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dewey, D. L., & Work, E. (1952). Diaminopimelic acid decarboxylase. *Nature*, 169(4300), 533–534. <https://doi.org/10.1038/169533a0>

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Diner, R. E., V. A. Bielinski, C. L. Dupont, A. E. Allen and P. D. Weyman (2016). Refinement of the Diatom Episome Maintenance Sequence and Improvement of Conjugation-Based DNA Delivery Methods. *Frontiers in bioengineering and biotechnology* 4: 65-65.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* 60: 2126-2132.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gibson, D. G., G. A Benders, K. C. Axelrod, J. Zaveri, M. A. Agire, M. Moodie, M. G. Montague, J. C. Venter, H. O. Smith and C. A. Hutchison (2008). One-step assembly in yeast of 25 overlapping DNA fragments to form a complete synthetic *Mycoplasma genitalium* genome. *Proceedings of the National Academy of Sciences* 105(51):20404.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gileadi, O., Burgess-Brown, N.A., Colebrook, S.M., Berridge, G., Savitsky, P., Smeets, C.E., Loppnau, P., Johansson, C., Salah, E., and Pantic, N.H. (2008). High throughput production of recombinant human proteins for crystallography. *Methods Mol Biol* 426: 221-246.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Goedhart, J., D. von Stetten, M. Noirclic-Savoye, M. Lelimosin, L. Joosen, M. A. Hink, L. van Weeren, T. W. J. Gadella and A. Royant (2012). Structure-guided evolution of cyan fluorescent proteins towards a quantum yield of 93%. *Nature Communications* 3(1): 751.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gokulan, K., B. Rupp, M. S. Pavelka, W. R. Jacobs and J. C. Sacchettini (2003). Crystal Structure of *Mycobacterium tuberculosis* Diaminopimelate Decarboxylase, an Essential Enzyme in Bacterial Lysine Biosynthesis. *Journal of Biological Chemistry* 278(20): 18588-18596

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Griffin, M. D., J. M. Billakanti, A. Wason, S. Keller, H. D. Mertens, S. C. Atkinson, R. C. Dobson, M. A. Perugini, J. A. Gerrard and F. G. Pearce (2012). Characterisation of the first enzymes committed to lysine biosynthesis in *Arabidopsis thaliana*. *PLoS One* 7(7): e40318.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gruber, A and P. G. Kroth (2017). Intracellular metabolic pathway distribution in diatoms and tools for genome-enabled experimental diatom research. *Philos Trans R Soc Lond B Biol Sci* 372(1728).

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Guillou, L., M.-J. Chretiennot-Dinet, L. K. Medlin, H. Claustre, S. L.-d. Goër and D. Vaulot (1999). *Bolidomonas*: A new genus with two species belonging to a new algal class, the Bolidophyceae (Heterokonta). *Journal of Phycology* 35(2): 368-381.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hor, L., M. G. Peverelli, M. A. Perugini and C. A. Hutton (2013). A new robust kinetic assay for DAP epimerase activity. *Biochimie* 95(10): 1949-1953.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hu, T., D. Wu, J. Chen, J. Ding, H. Jiang and X. Shen (2008). The Catalytic Intermediate Stabilized by a "Down" Active Site Loop for Diaminopimelate Decarboxylase from *Helicobacter pylori*: enzymatic characterization with crystal structure analysis. *Journal of Biological Chemistry* 283(30): 21284-21293

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hudson, A. O., C. Bless, P. Macedo, S. P. Chatterjee, B. K. Singh, C. Gilvarg and T. Leustek (2005). Biosynthesis of lysine in plants: evidence for a variant of the known bacterial pathways. *Biochim Biophys Acta* 1721(1-3): 27-36.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hudson, A. O., B. K. Singh, T. Leustek and C. Gilvarg (2006). An LL-diaminopimelate aminotransferase defines a novel variant of the lysine biosynthesis pathway in plants. *Plant Physiol* 140(1): 292-301.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hutton, C. A., M. A. Perugini and J. A. Gerrard (2007). Inhibition of lysine biosynthesis: an evolving antibiotic strategy. *Molecular BioSystems* 3(7): 458-465.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jander G, and Joshji, V. Aspartate-derived Amino Acid Biosynthesis in

***Arabidopsis thaliana*. The Arabidopsis Book. 2009;e0121(10.1199/tab.0121):15.**

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Karas, B. J., R. E. Diner, S. C. Lefebvre, J. McQuaid, A. P. R. Phillips, C. M. Noddings, J. K. Brunson, R. E. Valas, T. J. Deerinck, J. Jablanovic, J. T. F. Gillard, K. Beeri, M. H. Ellisman, J. I. Glass, C. A. Hutchison III, H. O. Smith, J. C. Venter, A. E. Allen, C. L. Dupont and P. D. Weyman (2015). Designer diatom episomes delivered by bacterial conjugation. *Nature Communications* 6: 6925.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Katoh, K. and D. M. Standley (2013). MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* 30(4): 772-780.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kelland, J. G., M. M. Palcic, M. A. Pickard and J. C. Vederas (1985). Stereochemistry of lysine formation by meso-diaminopimelate decarboxylase from wheat germ: use of proton-carbon-13 NMR shift correlation to detect stereospecific deuterium labeling. *Biochemistry* 24(13): 3263-3267.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kern, A. D., M. A. Oliveira, P. Coffino and M. L. Hackert (1999). Structure of mammalian ornithine decarboxylase at 1.6Å resolution: stereochemical implications of PLP-dependent amino acid decarboxylases. *Structure* 7(5): 567-581.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kidron, H., S. Repo, M. S. Johnson and T. A. Salminen (2006). Functional Classification of Amino Acid Decarboxylases from the Alanine Racemase Structural Family by Phylogenetic Studies. *Molecular Biology and Evolution* 24(1): 79-89.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kilian, O. and P. G. Kroth (2005). Identification and characterization of a new conserved motif within the presequence of proteins targeted into complex diatom plastids. *Plant J* 41(2): 175-183.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kirma, M., W. L. Araújo, A. R. Fernie and G. Galili (2012). The multifaceted role of aspartate-family amino acids in plant metabolism. *Journal of Experimental Botany* 63(14): 4995-5001.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Knorr, S., M. Sinn, D. Galetskiy, R. M. Williams, C. Wang, N. Müller, O. Mayans, D. Schleheck and J. S. Hartig (2018). Widespread bacterial lysine degradation proceeding via glutarate and L-2-hydroxyglutarate. *Nature Communications* 9(1): 5071.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kosuge, T. and T. Hoshino (1998). Lysine is synthesized through the α -amino adipate pathway in *Thermus thermophilus*. *FEMS Microbiology Letters* 169(2): 361-367.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lartillot, N., Rodrigue, N., Stubbs, D., & Richer, J. (2013). PhyloBayes MPI: phylogenetic reconstruction with infinite mixtures of profiles in a parallel environment. *Systematic Biology* 62(4), 611–615. <https://doi.org/10.1093/sysbio/syt022>

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lauritano, C., I. Orefice, G. Procaccini, G. Romano and A. Ianora (2015). Key genes as stress indicators in the ubiquitous diatom *Skeletonema marinoi*. *BMC Genomics* 16(1): 411-411.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liebschner, D., Afonine, P.V., Baker, M.L., Bunkoczi, G., Chen, V.B., Croll, T.I., Hintze, B., Hung, L.W., Jain, S., McCoy, A.J., et al. (2019). Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* 75, 861-877.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

McCoy, A. J., N. E. Adams, A. O. Hudson, C. Gilvarg, T. Leustek and A. T. Maurelli (2006). L,L-diaminopimelate aminotransferase, a trans-kingdom enzyme shared by *Chlamydia* and plants for synthesis of diaminopimelate/lysine. *Proceedings of the National Academy of Sciences* 103(47): 17909-17914.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read, R.J. (2007). Phaser crystallographic software. *J Appl Crystallogr* 40: 658-674.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mills, W. R., P. J. Lea and B. J. Mifflin (1980). Photosynthetic Formation of the Aspartate Family of Amino Acids in Isolated Chloroplasts. *Plant Physiology* 65(6): 1166-1172.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mock, T., R. P. Otilar, J. Strauss, M. McMullan, P. Paajanen, J. Schmutz, et al. (2017). Evolutionary genomics of the cold-adapted diatom *Fragilariopsis cylindrus*. *Nature* 541(7638): 536-540.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Moosburner, M. A., P. Gholami, J. K. McCarthy, M. Tan, V. A. Bielinski and A. E. Allen (2020). Multiplexed Knockouts in the Model Diatom *Phaeodactylum* by Episomal Delivery of a Selectable Cas9. *Frontiers in Microbiology* 11: 5.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nguyen, L.-T., H. A. Schmidt, A. von Haeseler and B. Q. Minh (2014). IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and Evolution* 32(1): 268-274.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Niesen, F.H., Berglund, H., and Vedadi, M. (2007). The use of differential scanning fluorimetry to detect ligand interactions that promote protein stability. *Nat Protoc* 2: 2212-2221.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Perl, A., O. Shaul and G. Galili (1992). Regulation of lysine synthesis in transgenic potato plants expressing a bacterial dihydrodipicolinate synthase in their chloroplasts. *Plant Molecular Biology* 19(5): 815-823.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Peeverelli, M. G. and M. A. Perugini (2015). An optimized coupled assay for quantifying diaminopimelate decarboxylase activity. *Biochimie* 115: 78-85.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Peeverelli, M. G., T. P. Soares da Costa, N. Kirby and M. A. Perugini (2016). Dimerization of Bacterial Diaminopimelate Decarboxylase Is Essential for Catalysis. *Journal of Biological Chemistry* 291(18): 9785-9795.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pollak, B., A. Cerda, M. Delmans, S. Álamos, T. Moyano, A. West, R. A. Gutiérrez, N. J. Patron, F. Federici and J. Haseloff (2019). Loop assembly: a simple and open system for recursive fabrication of DNA circuits. *New Phytologist* 222(1): 628-640.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Poulsen, N. and N. Kröger (2005). A new molecular tool for transgenic diatoms. *The FEBS Journal* 272(13): 3413-3423.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pratelli, R. and G. Pilot (2014). Regulation of amino acid metabolic enzymes and transporters in plants. *Journal of Experimental Botany* 65(19): 5535-5556.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Price, M. N., G. M. Zane, J. V. Kuehl, R. A. Melnyk, J. D. Wall, A. M. Deutschbauer and A. P. Arkin (2018). Filling gaps in bacterial amino acid biosynthesis pathways with high-throughput genetics. *PLOS Genetics* 14(1): e1007147.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ray, S. S., J. B. Bonanno, K. R. Rajashankar, M. G. Pinho, G. He, H. De Lencastre, A. Tomasz and S. K. Burley (2002). Cocrystal Structures of Diaminopimelate Decarboxylase: Mechanism, Evolution, and Inhibition of an Antibiotic Resistance Accessory Factor. *Structure* 10(11): 1499-1508.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Raymond, J. A. and H. J. Kim (2012). Possible role of horizontal gene transfer in the colonization of sea ice by algae. *PloS one* 7(5): e35968-e35968.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Reyes-Prieto, A. and A. Moustafa (2012). Plastid-localized amino acid biosynthetic pathways of Plantae are predominantly composed of non-cyanobacterial enzymes. *Sci Rep* 2: 955.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rosenwasser, S., S. Graff van Creveld, D. Schatz, S. Malitsky, O. Tzfadia, A. Aharoni, Y. Levin, A. Gabashvili, E. Feldmesser and A. Vardi (2014). Mapping the diatom redox-sensitive proteome provides insight into response to nitrogen stress in the marine environment. *Proc Natl Acad Sci U S A* 111(7): 2740-2745.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sagong, H. Y. and K. J. Kim (2017). Structural basis for redox sensitivity in *Corynebacterium glutamicum* diaminopimelate epimerase: an enzyme involved in L-lysine biosynthesis. *Sci Rep* 7: 42318.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Savitsky, P., J. Bray, C. D. O. Cooper, B. D. Marsden, P. Mahajan, N. A. Burgess-Brown and O. Gileadi (2010). High-throughput production of human proteins for crystallization: The SGC experience. *Journal of Structural Biology* 172(1): 3-13.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Siaut, M., M. Heijde, M. Mangogna, A. Montsant, S. Coesel, A. Allen, A. Manfredonia, A. Falciatore and C. Bowler (2007). Molecular toolbox for studying diatom biology in *Phaeodactylum tricornutum*. *Gene* 406(1): 23-35.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Smith, S. R., J. T. Gillard, A. B. Kustka, J. P. McCrow, J. H. Badger, H. Zheng, A. M. New, C. L. Dupont, T. Obata, A. R. Fernie and A. E. Allen (2016). Transcriptional Orchestration of the Global Cellular Response of a Model Pennate Diatom to Diel Light Cycling under Iron Limitation. *PLoS Genet* 12(12): e1006490.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Smith, S. R., C. L. Dupont, J. K. McCarthy, J. T. Broddrick, M. Oborník, A. Horák, Z. Füßy, J. Cihlář, S. Kleessen, H. Zheng, J. P. McCrow, K. K. Hixson, W. L. Araújo, A. Nunes-Nesi, A. Fernie, Z. Nikoloski, B. O. Palsson and A. E. Allen (2019). Evolution and regulation of nitrogen flux through compartmentalized metabolic networks in a marine diatom. *Nature Communications* 10(1): 4552.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Son, H. F. and K.-J. Kim (2018). Structural basis for substrate specificity of meso-diaminopimelic acid decarboxylase from *Corynebacterium glutamicum*. *Biochemical and Biophysical Research Communications* 495(2): 1815-1821.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Steinegger, M. and J. Söding (2017). MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nature Biotechnology* 35(11): 1026-1028.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Strand, T. A., R. Lale, K. F. Degnes, M. Lando and S. Valla (2014). A New and Improved Host-Independent Plasmid System for RK2-Based Conjugal Transfer. *PLOS ONE* 9(3): e90372.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sugio, S., G. A. Petsko, J. M. Manning, K. Soda and D. Ringe (1995). Crystal Structure of a D-Amino Acid Aminotransferase: How the Protein Controls Stereoselectivity. *Biochemistry* 34(30): 9661-9669.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Velasco, A. M., J. I. Leguina and A. Lazcano (2002). Molecular Evolution of the Lysine Biosynthetic Pathways. *Journal of Molecular Evolution* 55(4): 445-449.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wallsgröve, R. M. and M. Mazelis (1980). The enzymology of lysine biosynthesis in higher plants: Complete localization of the regulatory enzyme dihydrodipicolinate synthase in the chloroplasts of spinach leaves. *FEBS Letters* 116(2): 189-192.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang, H.-C., B. Q. Minh, E. Susko and A. J. Roger (2017). Modeling Site Heterogeneity with Posterior Mean Site Frequency Profiles Accelerates Accurate Phylogenomic Estimation. *Systematic Biology* 67(2): 216-235.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Weyand, S., Kefala, G., Svergun, D.I., and Weiss, M.S. (2009). The three-dimensional structure of diaminopimelate decarboxylase from *Mycobacterium tuberculosis* reveals a tetrameric enzyme organisation. *J Struct Funct Genomics* 10: 209-217.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Weyman, P. D., K. Beeri, S. C. Lefebvre, J. Rivera, J. K. McCarthy, A. L. Heuberger, G. Peers, A. E. Allen and C. L. Dupont (2015). Inactivation of *Phaeodactylum tricornutum* urease gene using transcription activator-like effector nuclease-based targeted mutagenesis. *Plant Biotechnology Journal* 13(4): 460-470.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

White, P. J. and B. Kelly (1965). Purification and properties of diaminopimelate decarboxylase from *Escherichia coli*. *The Biochemical Journal* 96(1): 75-84.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Winn, M.D., Ballard, C.C., Cowtan, K.D., Dodson, E.J., Emsley, P., Evans, P.R., Keegan, R.M., Krissinel, E.B., Leslie, A.G., McCoy, A., et al. (2011). Overview of the CCP4 suite and current developments. *Acta Crystallogr D Biol Crystallogr* 67: 235-242.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xu, H., B. Andi, J. Qian, A. H. West and P. F. Cook (2006). The α -amino adipate pathway for lysine biosynthesis in fungi. *Cell Biochemistry and Biophysics* 46(1): 43-64.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhu, X. and G. Galili (2004). Lysine Metabolism Is Concurrently Regulated by Synthesis and Catabolism in Both Reproductive and Vegetative Tissues. *Plant Physiology* 135(1): 129.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)