Effects of relocation on Immunological and physiological measures in

squirrel monkeys (*Saimiri boliviensis boliviensis*)

Pramod Nehete^{1,2}, Bharti P Nehete¹, Greg K Wilkerson¹, Steve J Schapiro^{1,3}, and Lawrence E Williams¹.

¹Department of Comparative Medicine, the University of Texas MD Anderson Cancer Center, Bastrop, Texas, USA, ²The University of Texas Graduate School of Biomedical Sciences, Houston, Texas, USA, ³Department of Experimental Medicine, University of Copenhagen, Copenhagen, Denmark

Corresponding Authors:

Pramod Nehete,

E-mail: pnehete@mdanderson.org

Department of Comparative Medicine

MD Anderson Cancer Center,

Address: 650 Cool Water Drive

Bastrop, TX 78602

Telephone: 512-332-5200

L. Williams,

lewillia@mdanderson.org

Department of Comparative Medicine,

MD Anderson Cancer Center,

Address: 650 Cool Water Drive

Bastrop, TX 78602

26 Telephone: (512) 321-3991.

27 Running title:

28 Key words: relocation, stress, Immune and physiological measurements

29

30

31 **Abstract**

32 In the present study, we have quantified the effects of transport, relocation and
33 acclimate/adapt to their new surroundings on squirrel monkey. These responses are
34 measured in blood samples obtained from squirrel monkeys, at different time points relative
35 to their relocation from their old home to their new home. A variety of immunological assays
36 are performed on the phenotype and function of peripheral blood mononuclear cells (PBMCs)
37 in a group of squirrel monkeys that were transported by road for approximately 10 hours from
38 one facility to another. Using a panel of human antibodies and a set of standardized human
39 immune assays, we evaluated the phenotype of lymphocyte subsets by flow, mitogen-specific
40 immune responses of PBMCs in vitro, and levels of cytokines at various time points including
41 immediately before transport, immediately upon arrival, and after approximately 150 days of
42 acclimation. We observed significant changes in T cells and subsets, NK and B cells (CD4⁺,
43 CD8⁺, CD4⁺/CD8⁺, CD16⁺, and CD20⁺). Mitogen specific (e.g. PHA, PWM and LPS)
44 proliferation responses, IFN- γ by ELISPOT assay, and cytokines (IL-2, IL-4 and VEGF)
45 significant changes were observed. Changes seen in the serum chemistry measurements
46 mostly complement those seen in the hematology data. The specific goal was to empirically
47 assess the effects of relocation stress in squirrel monkeys in terms of changes in the
48 numbers and functions of various leukocyte subsets in the blood and the amount of time
49 require for acclimating to their new environment. Such data will help to determine when newly
50 arrived animals become available for use in research studies.

51

52 **Introduction**

53 The number of nonhuman primates (NHP) used in U.S. biomedical research reached an all-
54 time high in 2018 year, according to data released in late September by the U.S. Department
55 of Agriculture (USDA) (NIH report released in September,2018. Nonhuman Primate
56 Evaluation and Analysis Part 1: Analysis of Future Demand and Supply, September 21,
57 2018). The rising demand for NHP appears to be driven by researchers studying HIV/AIDS,
58 cancer, the brain, Alzheimer's disease, addiction, Parkinson's, obesity/diabetes, and
59 emerging infectious diseases like Zika and Ebola and to learn better ways to prevent negative
60 pregnancy outcomes, including miscarriage, stillbirth, and premature birth. This research is
61 also helping scientists to uncover information that makes human organ transplants easier and
62 more accessible, literally giving new life to those whose kidneys, hearts, and lungs are failing.
63 Squirrel monkeys, small New World NHP, have served an important role in studying
64 pathogenesis of human disease conditions such as Alzheimer's disease [1-3], malaria [4-6],
65 HIV [7], Creutzfeldt-Jakob disease [8, 9] and Giardia infection [10].
66 Many of these nonhuman primates are raised at one facility and subsequently
67 transported/relocated to another facility for research purposes. Relocating captive nonhuman
68 primates from a familiar home cage or colony room to a novel environment is a potent
69 psychosocial stressor [11-15]. The new and unfamiliar environment presents a sudden,
70 uncontrollable, and unpredictable change. Manipulations of the environments of captive
71 nonhuman primates often have welfare consequences to the animals, including behavioral
72 effects, and for certain manipulations, physiological effects as well. The processes of
73 transporting, relocating, and acclimatizing nonhuman primates across facilities represent
74 manipulations that are likely to have welfare, behavioral, and physiological consequences to
75 the relocated animals [16-19]. Our group and a few others have undertaken a series of

studies that have attempted to quantify 1) the effects of transport and relocation, and 2) the amount of time that is required for NHPs to acclimate to the new environments and management procedures after relocation; whether the relocation was to the next room or halfway around the world [20, 21]. Previously, we have reported on the effects of relocation and transport on immunological measurements in chimpanzees, rhesus and cynomolgus monkeys [22-24]. During the winter of 2008 two colonies of different species of nonhuman primates, approximately 500 squirrel monkeys (*Saimiri sciureus sciureus*, *S. boliviensis boliviensis*, & *S. boliviensis peruviansis*) were transported to the University of Texas MD Anderson Cancer Center, Michale E. Keeling Center for Comparative Medicine and Research at Bastrop, TX (KCCMR) from the University of South Alabama in Mobile, AL.

In the present study, we assessed the effects of relocation stress on dependent variables of relevance in research (i.e., immune responses and hematological and chemistry values) in squirrel monkeys. We measured the physiological indicators of stress associated with relocation and the time course of adaptation to the physical and social environments of the new setting. The focus of this project was to empirically assess the effects of transport and relocation on physiological responses in NHPs, and to quantify acclimation processes. Measuring the effects of transport, relocation, and acclimation should allow investigators to conduct studies that are minimally influenced/confounded by such manipulations.

Materials and Methods:

Animals, Care, Diet and Housing

Subjects were 30 female squirrel monkeys (*Saimiri boliviensis*), aged 3-9 years of age from the breeding colony at University of Texas MD Anderson Cancer Center, Michale E. Keeling Center for Comparative Medicine and Research at Bastrop, TX (KCCMR). Monkeys were

socially housed throughout the study period in social groups in two connecting cages that are 4' wide x 6' tall x 14' long. Animals had *ad libitum* access to New World Primate Diet (Purina #5040) and water. In addition, they were fed either fresh fruit or vegetables daily. Specialty foods, such as seeds, peanuts, raisins, yogurt, cereals, frozen juice cups and peanut butter, were distributed daily to them as enrichment. At no time were the subjects ever food or water deprived. Subjects were also provided with destructible enrichment manipulanda and different travel/perching materials on a rotating basis to promote the occurrence of species-typical behaviors. The monkeys were examined by veterinarians before, during study period and determined to be healthy. Experiments were approved by the Institutional Animal Care and Use Committee of The University of Texas MD Anderson Cancer Center and were carried out according to the principles included in the *Guide for the Care and Use of Laboratory Animals*, the provisions of the Animal Welfare Act, PHS Animal Welfare Policy, and the policies of The University of Texas MD Anderson Cancer Center [25].

Prior to transport the animals were housed in the Primate Research Laboratory at the University of South Alabama, in social groups had been stable for at least one year. The animals were transported in modified Vari-Kennels® that provided perching and bedding appropriate for squirrel monkeys. Each kennel was provided with a hydration system gel pack, fresh vegetable and fruits, and non-human primate food biscuits. The squirrel monkeys were manually caught and placed in the kennels with a social partner between 1600-1700 hours. Entire social groups were moved at the same time. The trip from Mobile, AL to Bastrop, TX was made in a commercial self-contained, USDA approved, trailer that was climate controlled. The trailer left Mobile, AL between 1700-1730 and arrived in Bastrop, TX 0700-0800 the following morning. Beginning at approximately 0800 the kennels were removed from the trailer and placed in front of their new social housing units. Once an inventory was taken and all animals were found to be present and in the correct social

grouping, each kennel was opened, and the animals were released into their new housing. At no point were the animals sedated. The entire colony of 500 animals was moved over a five weeks period following the same routine procedures.

Collection of samples and Peripheral blood mononuclear cells (PBMC)

Ten animals from three of separate shipments, for a total of 30, were sampled for this study. Baseline blood samples were obtained one day prior to the relocation as part of the animals' "pre-shipment" physical exam. The next blood samples "Day 1" were obtained within the first few hours upon arrival at the Keeling Center as part of the animals' first "quarantine" physical exam. The final used for this study was collected at "Day150" post-arrival to monitor the animals' adaptation to their assigned housing conditions at the Keeling Center. Blood samples (3 mL) were collected, 1.5mL in EDTA anti-coagulant tubes and 1.5 mL in coagulation CBC tubes, from the femoral vein at the different time points. The animals were manually restrained for each of the three blood draws, and no sedatives or other chemical restraints were utilized. All blood sample collections occurred in the morning (8-10AM) before the animals were fed. Before the separation of peripheral blood mononuclear cells (PBMC) from the blood samples, plasma was collected and stored immediately at -80°C until analyzed. The PBMC prepared from the blood samples by the standard ficoll-hypaque density-gradient centrifugation were used for various immune assays [23, 26]. All blood samples were processed at the Keeling Center following domestic overnight shipment (for the pre-travel blood collections) or within 2-4 hours of collection at the Keeling Center (for arrival and day 150 post-arrival samples). The PBMCs, freshly prepared from whole blood collected in EDTA tubes, were more than 90% viable as determined by the trypan blue exclusion method. For each immune assay, 10^5 cells /well were used for various immune assay.

Complete blood count and blood chemistry analysis.

149 EDTA whole blood samples were analyzed for a complete blood count on Siemens Advia 120
150 Hematology Analyzer, Tarrytown, NY. The Parameters analyzed included: total WBC, total
151 RBC, hemoglobin, hematocrit, RBC indices, WBC differential counts, and platelet count.

152 Serum Chemistry was analyzed on Chemistry Analyzer (Beckman Coulter AU680®
153 Chemistry Analyzer). The different parameters analyzed were for example, Glucose, Na, K,
154 Cl, CO₂, Cholesterol, Triglycerides, Iron, BUN, Creatinine etc.).

155 **Flow cytometry:**

156 A series of commercially available human monoclonal antibodies that cross react with
157 nonhuman primate mononuclear cells were used in flow cytometry analyses as described
158 previously [10, 26-28]. Briefly, 100µl of whole blood from each sample was added to each
159 12mm×75mm polystyrene tube (Falcon, Lincoln Park, NJ, USA) containing pre-added
160 cocktail of monoclonal antibodies against CD3, CD4 and CD8 CD16 and CD20 (CD3-PE,
161 clone SP-34; CD4-PerCP, clone L200; CD8-FITC, clone SK1, CD16 APC (clone 3G8), and
162 CD20-APC, clone L27 (all from BD Biosciences, San Diego, USA) and incubated for 15 min
163 at room temperature in the dark. Red blood cells were lysed with 1x RBC lysing solution
164 (Becton Dickinson, USA) following the manufacturer's instructions. The samples were
165 washed thoroughly in 1x phosphate-buffer saline (PBS) by centrifugation; then cell sediments
166 were suspended in 1% paraformaldehyde buffer (300ul) and acquired on a on a
167 Fluorescence Activated Cell Sorter (FACS) Calibur flow cytometer (BD Biosciences, San
168 Jose, CA, USA). All samples acquired in this study were compensated using the single-color
169 stained cells. Lymphocytes that were gated on forward scatter versus side scatter dot plot
170 were used to analyze CD3⁺, CD4⁺, and CD8⁺ T cell and CD20⁺ B cell lymphocyte subsets
171 using FlowJo software (Tree Star, Inc., Ashland, OR, USA).
172 For analysis of NK cells, a separate tube with 100ul of blood was stained with separate
173 cocktail of consisting of -CD3 PE (clone SP-34 and CD16 APC (clone 3G8), (all from BD

Pharmingen, San Jose, USA) antibodies, as described above. The gating scheme for T-cell, B and NK markers in peripheral blood from a representative cynomolgus macaques has been identified previously [10]. The absolute number of lymphocytes and monocytes, as obtained from hematologic analysis, was used to convert the percentages identified through FACS analysis into absolute numbers for each of the lymphocyte and monocyte subset populations.

***in vitro* mitogen stimulation of PBMC**

The PBMCs, freshly prepared from whole blood collected in EDTA tubes, were more than 90% viable as determined by the trypan blue exclusion method. For each immune assay, 10^5 cells /well were used for various immune assay. Briefly, aliquots of PBMCs (10^5 /well) were seeded in triplicate wells of 96-well, flat-bottom plates and individually stimulated with the mitogens phytohemagglutinin (PHA), lipopolysaccharide (LPS), and pokeweed mitogen (PWM) (Sigma, St Louis, MO, USA), each at 1 μ g/mL final concentration. The culture medium without added mitogens served as a negative control.

Proliferation Assay

The proliferation of PBMC samples from the monkeys obtained at different time points during the study were determined by the standard [3 H] thymidine incorporation assay, using mitogens PHA, LPS and PWM (each at 1 μ g/mL final concentration). The culture medium served as negative control. Aliquots of the PBMC (10^5 /well) were suspended in RPMI-1640 culture medium supplemented with 10% fetal calf serum and seeded in triplicate wells of U-bottom 96-well plates and incubated with mitogens for 72hr at 37°C in humidified 5% CO₂ atmosphere. During the last 16-18 hr., 1 μ Cr of 3 H thymidine was added. Cells were harvested onto filter strips for estimating 3 H-incorporation and counted using a liquid scintillation counter. The proliferative response in terms of stimulation index (SI) was

calculated as fold-increase in the radioactivity over that of the cells cultured in medium alone.
The responses to antigens were considered positive when the SI values were ≤ 2.0 [29, 30].

ELISPOT Assay for Detecting IFN- γ producing Cells

Freshly-prepared PBMC were stimulated with the different mitogens (PHA, PWM and LPS) to determine the numbers of IFN- γ producing cells by the ELISPOT assay using the methodology reported earlier [10, 31, 32]. Briefly, aliquots of PBMC (10^5 /well) were seeded in duplicate wells of 96-well plates (MABTECH) pre-coated with the primary IFN- γ antibody and stimulated with mitogens PHA, LPS and PWM (each at 1 μ g/mL final concentration). After incubation for 24 hr. at 37°C, the cells were removed, and the wells were thoroughly washed with PBS. Subsequently, 100 μ L of biotinylated secondary antibody to IFN- γ (detection antibody) was added to the wells for 3 hr. at 37°C followed by avidin-peroxidase treatment for another 30 min. Purple colored spots representing individual cells secreting IFN γ were developed using freshly-prepared substrate (0.3 mg/mL of 3-amino-9-ethyl-carbazole) in 0.1 M sodium acetate buffer, containing 0.015% hydrogen peroxide. Plates were washed to stop color development, and spots were counted by an independent agency (Zellnet Consulting, New Jersey, NJ) using the KS-ELISPOT automatic system (Carl Zeiss, Inc. Thornwood, NY) for the quantitative analysis of the number of IFN- γ spot forming cells (SFC) for 10^5 input PBMC. Responses were considered positive when the numbers of spot forming cells (SFC) with the test antigen were at least five and were five above the background control values from cells cultured in the medium alone.

ELISA Assay for Detecting IFN- α producing Cells

Commercial Cytokines kits were used to measure the concentration of IFN- α (PBL Biomedical Laboratories, Piscataway, NJ) in cell culture supernatants following PHA, LPS

and PWM (each at 1 ug/mL final concentration) stimulation of PBMC from squirrel monkeys.

ELISA for IFN- α cytokine were performed according to the manufacturer's instructions.

The minimum IFN- a detectable concentration was 2.9 pg/mL.

NK Assay:

The natural killer activity (NK) was measured as previously described [33]. Briefly, PBMCs from blood were purified by centrifugation on a Ficoll-Hypaque density gradient as described above. Serial two-fold dilutions of the PBMC (effectors) were mixed with ^{51}Cr -labeled target cells K562 in triplicate wells of microtiter plates to attain the E: T ratio of 100:1, 50:1, 25:1 and 12.5:1. After 4-h incubation, 100 μl of supernatant was collected from each well and the amount of ^{51}Cr released was determined using the γ -counter. To account for the maximum release, the cells were incubated with 5% Triton X-100. Spontaneous release was determined from target cells incubated without added effector cells. The % of specific lysis was calculated by the following formula:

$$\% \text{ Specific lysis} = (\text{experimental release} - \text{spontaneous release}) / (\text{maximum release} - \text{spontaneous release}) \times 100.$$

Cytokine multiplex assays

Cytokine were measured in cell-free PBMC supernatant using MILLIPLEX-MAP human cytokine/chemokine magnetic bead panel (EMD Millipore Corporation, Billerica, MA, USA) according to the manufacturer's instructions. There is 91.4%–98.1% homology between the nucleotide sequences of SQM cytokine genes and published sequences of equivalent human and nonhuman primate genes [34, 35]. Briefly, aliquots of PBMC (10^5 /well) were seeded in duplicate wells of 96-well plates and stimulated for 24 hrs. with mitogens PHA, PWM and LPs (each at 1 ug/mL final concentration) supernatant samples were centrifuged (14,000x g for 5 min) and 25 mL of aliquots were used in assay. The 96-well filter plate was blocked with

assay buffer for 10 min at room temperature, washed, and 25 mL of standard or control samples were dispersed into appropriate wells. After adding 25 mL of beads to each well, the plate was incubated on a shaker overnight at 40C. The next day, after washing two times with wash buffer, the plate was incubated with detection antibody for 1 h at room temperature and again incubated with 25 mL of Streptavidin-Phycoerythrin for 30 min at room temperature. After washing two times with wash buffer, 150 mL of sheath fluid was added into each well and multianalyte profiling was performed on the Bio-Plex 200 system (Luminex X MAP technology). Calibration microspheres for classification and reporter readings as well as sheath fluid, assay, and wash buffer were also purchased from Bio-Rad (Hercules, CA, USA). Acquired fluorescence data were analyzed by the Bio-Plex manager 5.0 (Bio-Rad, Hercules, CA, USA). All steps of incubations were performed on a shaker. The minimum detectable concentration was calculated by the Multiplex Analyst immunoassay analysis Software from Millipore. Cytokines were measured using Nonhuman Primate Cytokine kit with IFN- γ , IFN- α , IL-1b, IL-2, IL-4, IL-6, MIP-1b, TNF- α , and VEGF, from Millipore Corporation (Billerica, MA) using the cytokine bead array (CBA) methodology according to the manufacturers' protocols and as described previously [36].

Statistical comparisons

The CBC, chemistry, and immunological data are analyzed using a series of within-subjects One-way Analyses of Variance. The primary comparisons are across the levels of the independent variable; transport and relocation (pre-transport, immediately after transport and after a 150day acclimation referred to as Pre, Day 1, day 150 samples in the results). Two-tailed tests, appropriate correction factors, and planned comparison techniques are utilized to fully explore the data.

Results

To understand the effect of transport and relocation on immune responses of PBMCs of

squirrel monkeys, we performed detailed analyses of cell-mediated immune responses, including assays for 1) Phenotypic analysis by flow cytometry, 2) proliferation, 3) IFN- γ by ELISPOT and IFN- α by ELISA, in response to stimulation with mitogens (e.g., PHA, PWM, and LPS), 4) cytokines in cell supernatant and 5) complete blood count and serum chemistry analysis before, and after relocation.

The lymphocytes and monocytes were first gated based on forward scatter (FCS) versus side scatter (SSC), and then CD3⁺ T cells, CD14⁺ (monocytes), CD3⁺CD16⁺ (NK) cells, CD3⁺CD16⁺ NKT cells, and CD20⁺ B cells were positively identified. The specificity of staining for the various markers was ascertained according to the isotype control antibody staining used for each pair of combination markers, as shown.

Fig 1 (A). Gating scheme for phenotype analyses of the various cell markers in the peripheral blood from a representative squirrel monkey.

Influence of relocation on major lymphocyte subsets in the peripheral blood

We first checked cross activity of a large panel of commercially available antibodies from different commercial companies at the concentration recommended by the supplier using blood from squirrel monkeys as described previously [37]. The reactivity was considered positive if the signal obtained gave a dot plot clearly distinct from the negative control dot plot using an isotype-matched antibody. As positive controls, fresh blood obtained from normal healthy rhesus monkey as donors were stained in parallel. Based on the data showing a high degree of cross reactivity of human monoclonal antibodies to different lymphocyte subsets in squirrel monkeys, we began probing into immunological indicators of stress associated with relocation.

Using the monoclonal antibodies listed in Table 1 we determined the levels of the different lymphocyte subsets and established normal value ranges in the blood of a total of 30 adult

Saimiri monkeys from the breeding unit of MD Anderson Cancer Center at Bastrop. Specifically, we analyzed for the T cells (CD3+), NK cells (CD16+), B cells (CD20+ cells), helper T cells (CD3+, CD4+) and cytotoxic/suppressor T cells (CD3+, CD8+) using human monoclonal antibodies that exhibited cross-reactivity with squirrel monkey PBMC. Details of the specificity, clone names, isotypes and supplier of the commercially available human monoclonal antibodies are shown in Table 1.

Table 1.

Human specific monoclonal antibodies used for squirrel monkey FACS and its reactivity

Antibody	Supplier	Clone	Isotype	Reactivity
CD3	BD	SP34	IgG3, λ	+
CD4	BD	L200	IgG1 κ	+
CD8	BD	RPA-T8	IgG1 κ	+
CD16	BD	3G8	IgG1 κ	+
CD20	BD	L27	IgG1 κ	+

The distribution lymphocyte subsets in blood of squirrel monkeys are shown at day pre shipment, post day 2 and post day150 after relocation in Fig 1B. CD3+ T cell count ($F(1.7,25.9) = 15, p < 0.05$) showed significant changes across time with the Day 1 arrival levels lower than the Pre and day 150 post-arrival samples. CD4+ T cell counts ($F(1.5,22.6) = 105, p < 0.05$), CD8+ T cell counts ($F(1.3,19.3) = 6.54, p < 0.05$), and CD4+CD8+(double positive) T cell counts ($F(1.1,16.6) = 226, p < 0.05$) showed significant changes across time with the day 150 levels significantly higher than both the Pre and Day 1 and Day 150 samples. CD16+ NK cells cell counts ($F(1.7,25.6) = 5.81, p < 0.05$) showed significant changes across time with Day 1 levels significantly lower than the Pre and Day 1 samples. CD20+ B cells counts ($F(1.2,18) = 9.58, p < 0.05$) showed significant changes across time with the Day 1 levels significantly higher than both the Pre and Day 1 samples.

CD3+CD16+ NK T cells cell counts (F (1.7,25.3) =5.66, p<0.05) showed significant changes across time with the Day 1 levels significantly lower than Pre.

Figure 1 (B). Relocation-dependent differences in lymphocytes in squirrel monkeys.

Proliferative responses

Since, we found significant differences in expression of T, and B cells, we investigated functional hallmark of proliferation of PBMCs samples from the squirrel monkeys. We measured proliferation in ³H thymidine incorporation assay (Fig 2A). The proliferative responses to PHA (F (2,36) =47.1, p<0.0001), and PWM (F (2,51) =33.7, p<0.0001) were significantly higher at post day 2 shipment compare to pre and day150. No statistically significant differences were observed for proliferative response stimulation with LPS (Fig 2A).

Figure 2 (A). Proliferative response of PBMCs to mitogens.

ELISPOT assay for detecting mitogen-specific IFN- γ producing cells in squirrel monkeys

Additional functional activity was measured for IFN γ production by PBMCs in response to stimulation with PHA, PWM, and LPS by the cytokine ELISPOT assay. As shown in Fig 2B, squirrel monkey PBMC showed significantly higher numbers of IFN- γ producing cells in response to stimulation with PHA (F(2,51)=136.4,p<0.0001), PWM (F(2,51)=136.4,p<0.0001) and LPS (F(2,51)=113.4,p<0.0001) showed significant changes across time with the Day 150 levels significantly higher than both the Pre and Day 1.

Figure 2 (B). IFN- γ ELISPOT response to mitogens.

ELISA assay for detecting mitogen-specific IFN- α producing cells in squirrel monkeys

Freshly isolated PBMCs were either unstimulated (medium) or stimulated with PHA, PWM and LPS (1 μ g/mL) for 24 hr. and supernatant was collected to measure IFN- α (Fig.3A) by ELISA. In general, Day 150 levels were intermediate between the Pre and Day 1 samples. No significant differences across time were observed for IFN-a ELISA with regard to the PHA, PWM, and LPS assays (Fig 3A).

Figure 3 (A). IFN- α ELISA response to mitogens.

Influence of relocation on natural killer activity. PBMCs from squirrel monkey were analyzed for NK activity using a standard ⁵¹chromium (Cr) release assay. We observed no significant differences were observed in natural killer activity at different effector to Target ratio (E: T100:1, E: T 50:1, and E: T25:1) (Fig 3B).

Figure 3 (B). Relocation-dependent differences in natural killer activity.

Cytokine multiplex assays

Freshly isolated PBMCs were either unstimulated (medium) or stimulated with PHA, PWM and LPS (1 μ g/mL) for 24 hr. and cell supernatant was collected frozen and used for multiplex cytokines assay using cytokines: IL-1 β , IL-2, IL-4, IL-6, MIP-1 β , TNF- α , and VEGF. Only IL-2 (PHA) (F (1.8,25.6) =7.29, p<0.05), IL-4 (PHA)(F (1.8,25.4) =4.3, p<0.05), and VEGF (PHA) (F (1.5,21.6) =5.58, p<0.05) showed significant differences across time with Day 1 arrival levels (Fig 4).

Figure 4. Relocation-dependent differences in Cytokine.

Hematology

The blood samples collected at pre shipment Day 1 and Day150 was subjected hematology and analysis is shown in Fig 5A and 5B. The white blood cell count ($F(1,12) = 12.2, p < 0.05$) and relative count of lymphocytes ($F(1.2,14.4) = 16.9, p < 0.05$) both showed a significant change across time, with the Pre levels significantly higher than both the Day 1 measurements and the post Day 150 samples. The monocyte cell count was significantly different across time ($F(1.5,19.5) = 5.38, p < 0.05$) with a significant difference between the Pre levels and Day 150 levels. The neutrophil count ($F(1.5,20.3) = 50.6, p < 0.05$) and hematocrit ($F(1.8,22.9) = 12.6, p < 0.05$) both showed a significant change across time, with the Pre significantly higher than the post-treatment samples and the Day 150 samples. The eosinophil levels were significantly different across time ($F(1.5,19.1) = 5.83, p < 0.05$) with Day 1 levels significantly higher compared to post Day 150 levels. The hemoglobin levels showed significant changes across time ($F(1.7,21.3) = 9.94, p \leq 0.05$) with Day 1 levels significantly lower than both Pre and Day 1 levels. The red blood cell count ($F(1.6,21.1) = 8.07, p < 0.05$), MCHC ($F(1.9,24.7) = 4.08, p < 0.05$), and RDW ($F(1.7,21.6) = 7.59, p < 0.05$) all were significantly different across time with Pre levels significantly higher than the Day1 levels. MCV, MCH, PLT, MPV, and segmented neutrophil levels showed no significant changes across time.

Serum Chemistry

Similarly, the blood samples collected at pre shipment, Day 1 and Day 150 was subjected to serum chemistry analysis and is shown in Fig 5C, 5D and 5E. Total bilirubin levels ($F(1.3,16.9) = 12.8, p < 0.05$), ALT levels ($F(1.7,23.8) = 26.7, p < 0.05$), CK levels ($F(1,21.6) = 8.9, p < 0.05$), BUN levels ($F(1.4,30.1) = 14.6, p < 0.05$), Glucose levels ($F(1.4,19.7) = 8.22, p < 0.05$), Osmolarity ($F(1.9,27) = 8.09, p < 0.05$), Phosphorus levels ($F(1.4,31.3) = 9.96, p < 0.05$), and Sodium levels ($F(1.7,24.5) = 12.0, p < 0.05$) all showed similar

significant changes across time, with Day 1 levels significantly higher than Pre and Day 1 and Day 150 samples. Triglyceride levels ($F(1.7, 24, 2) = 183$, $p < 0.05$) showed a significant change across time with Day 1 levels significantly lower than both Pre and the Day 1 and 150 samples. Albumin levels ($F(1.6, 23, 1) = 8.58$, $p < 0.05$) and Creatinine levels ($F(1.6, 35, 5) = 15.2$, $p < 0.05$) both showed significant changes across time with Day 1 and Day 150 levels significantly lower than both Pre and Day 1 samples. Globulin levels ($F(1.5, 21, 3) = 14.6$, $p < 0.05$) and Anion Gap ($F(1.8, 39, 4) = 5.48$, $p < 0.05$) showed similar significant changes across time but with post Day 150 levels significantly higher than both Pre and Day 1 arrival samples. Iron levels ($F(1.7, 23, 9) = 5.73$, $p < 0.05$) and cholesterol ($F(1.2, 17, 8) = 1.07$, $p < 0.05$) showed significant changes across time with the Pre samples significantly higher than Day 1 arrival. CO₂ levels ($F(1.4, 31, 1) = 2.73$, $p < 0.05$) showed significant changes across time with post Day 150 levels significantly higher than Pre samples. LDH levels ($F(1.8, 23, 6) = 7.44$, $p < 0.05$) showed significant changes across time with levels Day 1 arrival levels significantly higher than post Day 150 samples. Pre LDH levels were lower than the Day 1 arrival levels but not significantly so. AST levels ($F(1, 14, 3) = 32$, $p < 0.05$) showed significant changes across time with Day 1 arrival samples significantly higher than Pre and post Day 150 samples. The post Day 150 AST sample results were also significantly higher than Pre levels. GGT, ALK, Total protein, calcium, potassium, chloride, and TIBC levels showed no significant changes across time.

Figure 5A-5E. Relocation-dependent differences in Hematology (5A and 5B) and blood Chemistry (5C, D, E).

Effect of Acclimatization:

For most of the hematology measurements all animals had returned to Pre levels at the Day 150 sample. Exceptions include Total Lymphocytes which were higher 150 days post

transport ($F(1,9) = 13.0$, $\eta_p^2 = 0.6$), and RDW which remained lower than Pre ($F(1,9) = 7.6$, $\eta_p^2 = 0.5$).

These hematology effects can mostly be explained by mild dehydration during transport, including changes in the RBC, Hgb, and HCT, even though the animals were provided with water and gel-packs during transport. Transient reductions in the number of WBC's have been seen with stress as part of intense physical exercise. Unpredictable movements of the truck and crate may have tired and stressed the animals leading to the reduction in WBC's seen the day of arrival. However, changes in some of the serum chemistry measurements that are related to liver function may be related. Only two hematology measurements showed any post Day 150 effects, higher WBC and lower RDW in the squirrel monkeys. These results are not immediately explainable but do illustrate the types of post Day 150 changes in baseline values that can occur with relocation. Changes seen in the serum chemistry measurements mostly complement those seen in the hematology data. Mild dehydration during the transport may be the causes for changes in Osmolality and sodium levels in the squirrel monkeys. indirectly related may be changes in albumin, phosphorus, and triglycerides that are usually associated with kidney lesions. Elevated levels of CK and LDH, seen are associated with strenuous exercise. Like the fall in WBC's, this effect is likely due to the animals having to compensate for the unpredictable truck and cage movements. The increase in glucose levels in the squirrel monkeys as well as the decreased levels of iron can be attributed to overall stress and sleep deprivation. Changes in the ALT, AST, and BUN levels in the squirrel monkeys, and total bilirubin in both is usually indicative of liver lesions. Taken together with the platelet changes seen in *squirrel monkeys*, these changes suggest that transport has some transient effect on liver function through a as yet unknown mechanism.

Long term changes seen in BUN, GGT, Chloride, and cholesterol levels in the squirrel monkeys may be an accommodation to the new living arrangements. Although the new cages had a similar volume, they were constructed out of a thermo-neutral material (Tresva®) rather than sheet metal. They also received more cage enrichment and were on a different feeding and cleaning schedule. Any of these factors, or other changes in environment and routine, may be enough to alter the physiology of the animals enough to make it different from the original, pre-shipment, baseline.

Discussion

The data from this study establish clear physiological and immune system effects of transportation and acclimation for squirrel monkeys. Transport involves many factors that can be considered to negatively affect an animal's physiological state, including anesthesia; loading and unloading; separation from familiar social partners and environments; novel noises, smells, and vibrations; and relocation to a new and unfamiliar environment. The negative effects of these events have been assessed in several species [38-41], as changes in serum levels of cortisol and other physiological measures. Others have previously studied the effects of transport, relocation, and/or acclimatization on nonhumans primates [11, 24, 41-45], however these studies were conducted on old world and ape species, and focused on limited sets of behavioral or physiological responses and reported a decrease in lymphocytes in cynomolgus macaques, while Kagira, Ngotho (46) found changes in hematological parameters for recently trapped vervet. There are considerably more data available on the effects of transport and relocation in other, non-primate animal species. These reports found similar effects for mice, rats, dogs, and pigs as a function of transport, relocation, and/or acclimatization as suggested by changes in blood glucose, cholesterol, and blood urea nitrogen [47, 48]; lymphocyte counts [38]; and white blood cell counts, body weights, and natural killer cell activity [15, 40].

456 The data from the present study provide some insight into the time that it takes squirrel to
457 acclimatize to their new surroundings. Some standard clinical chemistry and hematologic
458 values appeared to return to Pre levels by about 6 weeks after arrival, while others did not.
459 Some of the cell-mediated immune responses that were affected by transport and relocation
460 also did not return to Pre levels. The squirrel monkeys were still affected by the transport
461 process weeks after transport and relocation, and probably should not be considered
462 acclimated to their new facility. We reported similar effects in chimpanzees, rhesus and
463 cynomolgus monkey's relocation to Texas [23, 24] . Their conclusion was that the
464 chimpanzees, rhesus and cynomolgus monkeys should not serve as subjects in studies that
465 use the measured parameters as dependent variables, until they have had adequate time to
466 adjust to their new conditions. This required at least 6-8 weeks for the chimpanzees and
467 longer for the monkeys in this study. The current data does not address the time interval
468 between arrival and 150 days. Additional data is needed to fill in this 5-month gap in the
469 monkey data.

470 This study focused on the effects of transportation and relocation as assessed by a variety of
471 parameters that are also likely to be dependent measures in biomedical investigations.
472 Significant changes as a function of transport may have few clinical implications in healthy
473 animals, yet they may have numerous research implications. For example, changes in red
474 blood cell counts in investigations of malaria [49] or changes in CD4+ counts in
475 immunodeficiency virus investigations [50]. The better the effects of transport are
476 understood, the better the refinements to management procedures can be. The development
477 of more refined management techniques will result in enhanced welfare for the animals [24,
478 51-54], enhanced abilities to directly test experimental hypotheses, and may ultimately result
479 in important reductions in the number of primate subjects required to effectively test
480 hypotheses.

Future studies will examine smaller-scale movements of squirrel monkeys within the various housing settings of the KCCMR. We have additional data [24, 55] from reasonably-sized samples of transported and relocated chimpanzees and cynomolgus monkeys that also demonstrate that transport and relocation result in statistically significant changes in a variety of hematological, clinical chemistry, and immunological parameters for these species, although the specific measures affected differ by species.

The data presented here crucial to guide researchers in determining an appropriate acclimation period for their study; however, there are additional factors researchers must consider that can influence how long it takes for acclimation to occur. These factors include the intensity and duration of stress, as well as the species, sex, age, genotype, health status, previous life experience, allometric differences, and even the time of year that acclimation occurs.

Acknowledgement

We thank Virginia Parks and Bethany Brock for providing blood samples for this study. This study is partly supported by the Cattleman for Cancer Research (PN) and the Squirrel Monkey Breeding and Research Resource (LW) at the Michale E Keeling Center for Comparative Medicine and Research at the MD Anderson Cancer Center.

Author Contributions: PN and LW designed, interpreted, and analyzed data. PN and BN performed the experiments. GW and SS provided additional interpretation.

Conflicts of Interest: The authors no conflict of interest.

Funding: P40-OD010938-40 (LW), and Cattleman for Cancer Research grant (PN, LW).

References

1. Elfenbein HA, Rosen RF, Stephens SL, Switzer RC, Smith Y, Pare J, et al. Cerebral beta-amyloid angiopathy in aged squirrel monkeys. *Histology and histopathology*. 2007;22(2):155-67. PubMed PMID: 17149688.
2. Heuer E, Jacobs J, Du R, Wang S, Keifer OP, Cintron AF, et al. Amyloid-Related Imaging Abnormalities in an Aged Squirrel Monkey with Cerebral Amyloid Angiopathy. *Journal of Alzheimer's disease : JAD*. 2017;57(2):519-30. doi: 10.3233/JAD-160981. PubMed PMID: 28269776.
3. Nehete PN, Williams LE, Chitta S, Nehete BP, Patel AG, Ramani MD, et al. Class C CpG Oligodeoxynucleotide Immunomodulatory Response in Aged Squirrel Monkey (*Saimiri Boliviensis* Boliviensis). *Front Aging Neurosci*. 2020;12:36. Epub 2020/03/21. doi: 10.3389/fnagi.2020.00036. PubMed PMID: 32194391; PubMed Central PMCID: PMCPCMC7063459.
4. Collins WE, Sullivan JS, Strobert E, Galland GG, Williams A, Nace D, et al. Studies on the Salvador I strain of *Plasmodium vivax* in non-human primates and anopheline mosquitoes. *Am J Trop Med Hyg*. 2009;80(2):228-35. Epub 2009/02/05. PubMed PMID: 19190218.
5. Riccio EKP, Pratt-Riccio LR, Bianco-Júnior C, Sanchez V, Totino PRR, Carvalho LJM, et al. Molecular and immunological tools for the evaluation of the cellular immune response in the neotropical monkey *Saimiri sciureus*, a non-human primate model for malaria research. *Malaria Journal*. 2015;14(1):166. doi: 10.1186/s12936-015-0688-1.
6. Tougan T, Aoshi T, Coban C, Katakai Y, Kai C, Yasutomi Y, et al. TLR9 adjuvants enhance immunogenicity and protective efficacy of the SE36/AHG malaria vaccine in nonhuman primate models. *Hum Vaccin Immunother*. 2013;9(2):283-90. Epub 2013/01/08. doi: 10.4161/hv.22950. PubMed PMID: 23291928; PubMed Central PMCID: PMCPCMC3859748.
7. LaBonte JA, Babcock GJ, Patel T, Sodroski J. Blockade of HIV-1 infection of New World monkey cells occurs primarily at the stage of virus entry. *J Exp Med*. 2002;196(4):431-45. doi: 10.1084/jem.20020468. PubMed PMID: 12186836.
8. Williams L, Brown P, Ironside J, Gibson S, Will R, Ritchie D, et al. Clinical, neuropathological and immunohistochemical features of sporadic and variant forms of Creutzfeldt-Jakob disease in the squirrel monkey (*Saimiri sciureus*). *J Gen Virol*. 2007;88(Pt 2):688-95. Epub 2007/01/26. doi: 10.1099/vir.0.81957-0. PubMed PMID: 17251588.
9. Ritchie DL, Gibson SV, Abee CR, Kreil TR, Ironside JW, Brown P. Blood transmission studies of prion infectivity in the squirrel monkey (*Saimiri sciureus*): the Baxter study. *Transfusion*. 2016;56(3):712-21. Epub 2015/11/26. doi: 10.1111/trf.13422. PubMed PMID: 26594017.
10. Nehete PN, Wilkerson G, Nehete BP, Chitta S, Ruiz JC, Scholtzova H, et al. Cellular immune responses in peripheral blood lymphocytes of *Giardia* infected squirrel monkey (*Saimiri boliviensis boliviensis*) treated with Fenbendazole. *PLOS ONE*. 2018;13(11):e0198497. doi: 10.1371/journal.pone.0198497.
11. Capitanio JP, Lerche NW. Social separation, housing relocation, and survival in simian AIDS: a retrospective analysis. *Psychosomatic medicine*. 1998;60(3):235-44. doi: 10.1097/00006842-199805000-00001. PubMed PMID: 9625208.
12. Champagne AB, Gunstone RF, Klopfer LE, West LHT, Pines AL. Cognitive Structure and Conceptual Change1985. null p.
13. Crockett CM, Bowers CL, Sackett GP, Bowden DM. Urinary cortisol responses of longtailed macaques to five cage sizes, tethering, sedation, and room change. *Am J Primatol*. 1993;30(1):55-74. Epub 1993/01/01. doi: 10.1002/ajp.1350300105. PubMed PMID: 31941178.
14. Crockett CM, Shimoji M, Bowden DM. Behavior, appetite, and urinary cortisol responses by adult female pigtailed macaques to cage size, cage level, room change, and ketamine sedation. *Am J Primatol*. 2000;52(2):63-80. Epub 2000/10/29. doi: 10.1002/1098-2345(200010)52:2<63::Aid-ajp1>3.0.Co;2-k. PubMed PMID: 11051442.

15. Dalin AM, Magnusson U, Häggendal J, Nyberg L. The effect of transport stress on plasma levels of catecholamines, cortisol, corticosteroid-binding globulin, blood cell count, and lymphocyte proliferation in pigs. *Acta Vet Scand.* 1993;34(1):59-68. Epub 1993/01/01. PubMed PMID: 8342466.
16. Neal Webb SJ, Hau J, Schapiro SJ. Relationships between captive chimpanzee (*Pan troglodytes*) welfare and voluntary participation in behavioural studies. *Applied Animal Behaviour Science.* 2019;214:102-9. doi: <https://doi.org/10.1016/j.applanim.2019.03.002>.
17. Crockett CM, Bowers CL, Shimoji M, Leu M, Bowden DM, Sackett GP. Behavioral responses of longtailed macaques to different cage sizes and common laboratory experiences. *J Comp Psychol.* 1995;109(4):368-83. Epub 1995/12/01. doi: 10.1037/0735-7036.109.4.368. PubMed PMID: 7497695.
18. Worlein JM, Kroeker R, Lee GH, Thom JP, Bellanca RU, Crockett CM. Socialization in pigtailed macaques (*Macaca nemestrina*). *Am J Primatol.* 2017;79(1):1-12. Epub 2016/04/26. doi: 10.1002/ajp.22556. PubMed PMID: 27109591; PubMed Central PMCID: PMC5994344.
19. Golub MS, Hogrefe CE, Germann SL, Capitanio JP, Lozoff B. Behavioral consequences of developmental iron deficiency in infant rhesus monkeys. *Neurotoxicology and Teratology.* 2006;28(1):3-17. doi: <https://doi.org/10.1016/j.ntt.2005.10.005>.
20. Capitanio JP, Kyes RC, Fairbanks LA. Considerations in the selection and conditioning of Old World monkeys for laboratory research: animals from domestic sources. *Ilar j.* 2006;47(4):294-306. Epub 2006/09/12. doi: 10.1093/ilar.47.4.294. PubMed PMID: 16963810.
21. Obernier JA, Baldwin RL. Establishing an appropriate period of acclimatization following transportation of laboratory animals. *Ilar j.* 2006;47(4):364-9. Epub 2006/09/12. doi: 10.1093/ilar.47.4.364. PubMed PMID: 16963816.
22. Nehete PN, Shelton KA, Nehete BP, Chitta S, Williams LE, Schapiro SJ, et al. Effects of transportation, relocation, and acclimation on phenotypes and functional characteristics of peripheral blood lymphocytes in rhesus monkeys (*Macaca mulatta*). *PloS one.* 2017;12(12):e0188694-e. doi: 10.1371/journal.pone.0188694. PubMed PMID: 29261698.
23. Shelton KA, Nehete BP, Chitta S, Williams LE, Schapiro SJ, Simmons J, et al. Effects of Transportation and Relocation on Immunologic Measures in *Cynomolgus* Macaques (*Macaca fascicularis*). *J Am Assoc Lab Anim Sci.* 2019;58(6):774-82. Epub 2019/10/13. doi: 10.30802/aalas-jaalas-19-000007. PubMed PMID: 31604484; PubMed Central PMCID: PMC6926399.
24. Schapiro SJ, Lambeth SP, Jacobsen KR, Williams LE, Nehete BN, Nehete PN. Physiological and Welfare Consequences of Transport, Relocation, and Acclimatization of Chimpanzees (*Pan troglodytes*). *Appl Anim Behav Sci.* 2012;137(3-4):183-93. Epub 2012/07/10. doi: 10.1016/j.applanim.2011.11.004. PubMed PMID: 22773870; PubMed Central PMCID: PMC3388538.
25. Laboratory animal welfare: Public Health Service policy on humane care and use of laboratory animals by awardee institutions; notice. *Federal register.* 1985;50(90):19584-5. Epub 1985/05/09. PubMed PMID: 11655780.
26. Nehete P, Hanley P, Nehete B, Yang G, Ruiz J, Williams L, et al. Phenotypic and Functional Characterization of Lymphocytes from Different Age Groups of Bolivian Squirrel Monkeys (*Saimiri boliviensis boliviensis*). *PloS one.* 2013;8:e79836. doi: 10.1371/journal.pone.0079836.
27. Nehete PN, Shelton KA, Nehete BP, Chitta S, Williams LE, Schapiro SJ, et al. Effects of transportation, relocation, and acclimation on phenotypes and functional characteristics of peripheral blood lymphocytes in rhesus monkeys (*Macaca mulatta*). *PLoS One.* 2017;12(12):e0188694. Epub 2017/12/21. doi: 10.1371/journal.pone.0188694. PubMed PMID: 29261698; PubMed Central PMCID: PMC5736198.
28. Nehete PN, Williams LE, Chitta S, Nehete BP, Patel AG, Ramani MD, et al. Class C CpG Oligodeoxynucleotide Immunomodulatory Response in Aged Squirrel Monkey (*Saimiri Boliviensis Boliviensis*). *Frontiers in Aging Neuroscience.* 2020;12(36). doi: 10.3389/fnagi.2020.00036.

29. Nehete PN, Murthy KK, Satterfield WC, Arlinghaus RB, Sastry KJ. Studies on V3-specific cross-reactive T-cell responses in chimpanzees chronically infected with HIV-1IIB. *AIDS*. 1995;9(6).
30. Oka T, Sastry KJ, Nehete P, Schapiro SJ, Guo JQ, Talpaz M, et al. Evidence for specific immune response against P210 BCR-ABL in long-term remission CML patients treated with interferon. *Leukemia*. 1998;12(2):155-63. Epub 1998/03/31. doi: 10.1038/sj.leu.2400919. PubMed PMID: 9519777.
31. Nehete PN, Chitta S, Hossain MM, Hill L, Bernacky BJ, Baze W, et al. Protection against chronic infection and AIDS by an HIV envelope peptide-cocktail vaccine in a pathogenic SHIV-rhesus model. *Vaccine*. 2001;20(5-6):813-25. Epub 2001/12/12. doi: 10.1016/s0264-410x(01)00408-x. PubMed PMID: 11738745.
32. Nehete PN GR, Nehete B, Sastry KJ. Dendritic cells enhance detection of antigen-specific cellular immune responses by lymphocytes from rhesus macaques immunized with an HIV envelope peptide cocktail vaccine. *Jr Medical Primatology*. 2003;In Press.
33. Nehete PN, Lewis DE, Tang DN, Pollack MS, Sastry KJ. Presence of HLA-C-restricted cytotoxic T-lymphocyte responses in long-term nonprogressors infected with human immunodeficiency virus. *Viral Immunol*. 1998;11(3):119-29. Epub 1999/01/26. doi: 10.1089/vim.1998.11.119. PubMed PMID: 9918403.
34. Heraud JM, Merien F, Mortreux F, Mahieux R, Kazanji M. Immunological changes and cytokine gene expression during primary infection with human T-cell leukaemia virus type 1 in squirrel monkeys (*Saimiri sciureus*). *Virology*. 2007;361(2):402-11. Epub 2007/01/16. doi: 10.1016/j.virol.2006.11.028. PubMed PMID: 17223152.
35. Heraud J-M, Lavergne A, Kazanji M. Molecular cloning, characterization, and quantification of squirrel monkey (*Saimiri sciureus*) Th1 and Th2 cytokines. *Immunogenetics*. 2002;54:20-9. doi: 10.1007/s00251-002-0443-y.
36. Nehete P, Magden ER, Nehete B, Hanley PW, Abee CR. Obesity related alterations in plasma cytokines and metabolic hormones in chimpanzees. *International Journal of Inflammation*. 2014;2014.
37. Nehete PN, Nehete BP, Chitta S, Williams LE, Abee CR. Phenotypic and Functional Characterization of Peripheral Blood Lymphocytes from Various Age- and Sex-Specific Groups of Owl Monkeys (*Aotus nancymae*). *Comp Med*. 2017;67(1):67-78. PubMed PMID: 28222841.
38. Bergeron R, Scott SL, Emond J-P, Mercier F, Cook NJ, Schaefer AL. Physiology and behavior of dogs during air transport. *Can J Vet Res*. 2002;66(3):211-6. PubMed PMID: 12146895.
39. Fazio E, Medica P, Alberghina D, Cavaleri S, Ferlazzo A. Effect of long-distance road transport on thyroid and adrenal function and haematocrit values in Limousin cattle: influence of body weight decrease. *Vet Res Commun*. 2005;29(8):713-9. Epub 2005/12/22. doi: 10.1007/s11259-005-3866-8. PubMed PMID: 16369885.
40. McGlone JJ, Salak JL, Lumpkin EA, Nicholson RI, Gibson M, Norman RL. Shipping stress and social status effects on pig performance, plasma cortisol, natural killer cell activity, and leukocyte numbers. *J Anim Sci*. 1993;71(4):888-96. Epub 1993/04/01. doi: 10.2527/1993.714888x. PubMed PMID: 8478291.
41. Watson SL, McCoy JG, Stavisky RC, Greer TF, Hanbury D. Cortisol response to relocation stress in Garnett's bushbaby (*Otolemur garnettii*). *Contemp Top Lab Anim Sci*. 2005;44(3):22-4. Epub 2005/06/07. PubMed PMID: 15934719.
42. Davenport MD, Lutz CK, Tiefenbacher S, Novak MA, Meyer JS. A rhesus monkey model of self-injury: effects of relocation stress on behavior and neuroendocrine function. *Biol Psychiatry*. 2008;63(10):990-6. Epub 2008/01/01. doi: 10.1016/j.biopsych.2007.10.025. PubMed PMID: 18164279; PubMed Central PMCID: PMC2486411.
43. Fernström AL, Sutian W, Royo F, Fernström AL, Sutian W, Royo F, et al. Stress in cynomolgus monkeys (*Macaca fascicularis*) subjected to long-distance transport and simulated transport housing conditions. *Stress*. 2008;11(6):467-76. doi: 10.1080/10253890801903359.

44. Honess PE, Johnson PJ, Wolfensohn SE. A study of behavioural responses of non-human primates to air transport and re-housing. *Lab Anim.* 2004;38(2):119-32. Epub 2004/04/09. doi: 10.1258/002367704322968795. PubMed PMID: 15070451.
45. Kim CY, Han JS, Suzuki T, Han SS. Indirect indicator of transport stress in hematological values in newly acquired cynomolgus monkeys. *J Med Primatol.* 2005;34(4):188-92. Epub 2005/08/02. doi: 10.1111/j.1600-0684.2005.00116.x. PubMed PMID: 16053496.
46. Kagira JM, Ngotho M, Thuita JK, Maina NW, Hau J. Hematological changes in vervet monkeys (*Chlorocebus aethiops*) during eight months' adaptation to captivity. *Am J Primat.* 2007;69(9):1053-63.
47. van Ruiven R, Meijer G, Wiersma A, Baumans V, van Zutphen L, Ritskes-Hoitinga R. The influence of transportation stress on selected nutritional parameters to establish the necessary minimum period for adaptation in rat feeding studies. *Lab Animal.* 1998;32:446-56.
48. Van Scott MR, Reece SP, Olmstead S, Wardle R, Rosenbaum MD. Effects of acute psychosocial stress in a nonhuman primate model of allergic asthma. *Journal of the American Association for Laboratory Animal Science : JAALAS.* 2013;52(2):157-64. PubMed PMID: 23562098.
49. Barasa M, Gicheru MM, Kagasi EA, Hastings OS. Characterisation of placental malaria in olive baboons (*Papio anubis*) infected with *Plasmodium knowlesi* H strain. *International Journal of Integrative Biology.* 2010;9(2):54-8.
50. The Primate Immune System. A Companion to Anthropological Genetics. p. 309-26.
51. Laule G, Whittaker M. Enhancing nonhuman primate care and welfare through the use of positive reinforcement training. *J Appl Anim Welf Sci.* 2007;10(1):31-8. Epub 2007/05/09. doi: 10.1080/10888700701277311. PubMed PMID: 17484676.
52. Schaffner CM, Smith TE. Familiarity may buffer the adverse effects of relocation on marmosets (*Callithrix kuhlii*): Preliminary evidence. *Zoo Biology.* 2005;24(1):93-100. doi: 10.1002/zoo.20019.
53. Schapiro S, Lambeth S, Thiele E, Rousset O. The effects of behavioral management programs on dependent measures in biomedical research. *American Journal of Primatology.* 2007;69.
54. Veeder CL, Bloomsmith MA, McMillan JL, Perlman JE, Martin AL. Positive reinforcement training to enhance the voluntary movement of group-housed sooty mangabeys (*Cercocebus atys atys*). *J Am Assoc Lab Anim Sci.* 2009;48(2):192-5. Epub 2009/04/23. PubMed PMID: 19383217; PubMed Central PMCID: PMC2679662.
55. Cinque C, De Marco A, Mairesse J, Giuli C, Sanna A, De Marco L, et al. Relocation stress induces short-term fecal cortisol increase in Tonkean macaques (*Macaca tonkeana*). *Primates; journal of primatology.* 2017;58(2):315-21. doi: 10.1007/s10329-016-0590-7. PubMed PMID: 27904967.

FIGURE LEGENDS

Fig 1 (A). Gating scheme for phenotype analyses of the various cell markers in the peripheral blood from a representative squirrel monkey. The lymphocytes were first gated based on forward scatter (FCS) versus side scatter (SSC), and then CD3⁺, CD4⁺, CD8⁺, CD4⁺CD8⁺, and CD20⁺ cells, and NK and NKT cells, were positively identified from the lymphocyte subset. The specificity of staining for the various markers was ascertained according to the isotype control antibody staining used for each pair of combination markers, as shown.

Figure 1 (B). Relocation-dependent differences in lymphocytes in squirrel monkeys.

Aliquots of EDTA whole blood were stained with fluorescence-labeled antibodies to the CD3⁺, CD4⁺, CD8⁺, and CD20⁺ to identify lymphocyte subpopulations Pre- and Day 1 and 150 post-transportation and relocation. Values on the Y-axis are absolute lymphocyte cells. P values were considered statistically significant at $p < 0.05$. *Symbol: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.0001$.*

Figure 2 (A). Proliferative response of PBMCs to mitogens. PBMCs that are isolated from blood samples of the squirrel monkeys were used for determining proliferative response to different mitogens, using the standard [3H] thymidine incorporation assay. The proliferation responses are expressed as Stimulation Index (SI) after blank (i.e., medium only) subtraction. P values were considered statistically significant at $p < 0.05$. *Symbol: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.0001$.*

Figure 2 (B). IFN- γ ELISPOT response to mitogens. In duplicate wells of the 96-well microtiter plates, pre-coated with IFN- γ antibody, were seeded with 10^5 PBMCs from squirrel monkeys stimulated with 1 μ g of each of the mitogens for 36 h at 37°C, and then washed and stained with biotinylated second IFN- γ antibody. The total number of spots forming cells (SFCs) in each of the mitogen-stimulated wells was counted and adjusted to control medium as background. See the Methods section for experiment details. P values were considered statistically significant at $p < 0.05$. *Symbol: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.0001$.*

Figure 3 (A). IFN- α ELISA response to mitogens. In triplicate wells of the 96-well filter plate, PBMC cultures were stimulated with 1 μ g/mL mitogens for 36hr at 37°C and supernatant was collected to measure IFN- α by ELISA. The minimum detectable concentration in pg/mL for IFN α (2.9) was used for considering positive responses. P values were considered statistically significant at $p < 0.05$. *Symbol: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.0001$.*

Figure 3 (B). Relocation-dependent differences in natural killer activity. The NK activity of PBMC isolated from squirrel monkeys' blood was determined by co-culturing with ⁵¹Cr-labeled K562 target cells at different effector to target cell ratios in culture medium. The percentage (%) of specific lysis is

shown at different effector to target ratios. P values $p < 0.05$ were considered statistically significant.

*Symbol: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.0001$.*

Figure 4. Relocation-dependent differences in Cytokine. In duplicate wells of the 96-well filter plate, 25 μ L of cell supernatant was incubated with 25 μ L of cytokine coupled beads overnight at 4°C, followed by washing and staining with biotinylated detection antibody. The plates were read on Biorad 200 with use of Luminex technology, and the results are expressed as pg/mL concentration. The minimum detectable concentrations in pg/mL for IL-2 (0.7), IL-4 (2.7), IL-6 (0.3), IL-10 (6.2), IL-12(P40) (1.2), IFN- γ (2.2), and TNF- α (2.1) were used for considering positive responses. See the Methods section for experimental details. P values were considered statistically significant at $p < 0.05$.

*Symbol: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.0001$.*

Figure 5A-5E. Relocation-dependent differences in Hematology (5A and 5B) and blood Chemistry (5C, D, E). Whole blood by using an automated analyzer Advia (Siemens Healthcare Diagnostics, Tarrytown, NY). Values on the Y-axis are the absolute numbers of lymphocytes and monocytes presented as 10^3 per μ L of whole blood. P values were considered statistically significant at $p < 0.05$. *Symbol: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.0001$.*

Fig 1A

Fig 1B

Fig 2A

Fig 2B

Fig 3A

Fig 3B

Fig 4A

Fig 4B

Fig 5A

Fig 5B

Fig 5C

Fig 5D

Fig 5E