1 **Effect of high-frequency low-intensity pulsed electric field** 2 **on protecting SH-SY5Y cells against hydrogen peroxide** 3 **and β -amyloid-induced cell injury via ERK pathway**

4
5

6 Wei-Ting Chen^{1,2,¶}, Guan-Bo Lin^{1,2,¶}, Yu-Yi Kuo^{1,2,¶}, Chih-Hsiung Hsieh^{1,2},
7 Chueh-Hsuan Lu^{1,2}, Yi-Kun Sun², and Chih-Yu Chao^{1,2,3,*}

8

9 ¹ Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan
10 University, Taipei 10617, Taiwan

11 ² Biomedical & Molecular Imaging Center, National Taiwan University College of Medicine,
12 Taipei 10051, Taiwan

13 ³ Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University,
14 Taipei 10617, Taiwan

15

16 * Corresponding author

17 E-mail: cychao@phys.ntu.edu.tw (CYC)

18 ¶ These authors contributed equally to this work.

19
20
21
22
23
24
25
26
27
28

Abstract

As the most common type of neurodegenerative diseases (NDDs), Alzheimer's disease (AD) is thought to be caused mainly by the excessive aggregation of β -amyloid protein ($A\beta$). However, a growing number of studies have found that reactive oxygen species (ROS) play a key role in the onset and progression of AD. The present study aimed to probe the neuroprotective effect of high-frequency low-intensity pulsed electric field (H-LIPEF) for SH-SY5Y cells against hydrogen peroxide (H_2O_2) and $A\beta$ -induced cytotoxicity. By looking in a systematic way into the frequency- and amplitude-dependent neuroprotective effect of pulsed electric field (PEF), the study finds that H-LIPEF at 200 Hz produces the optimal protective effect for SH-SY5Y cells. The underlying mechanisms were confirmed to be due to the activation of extracellular signal-regulated kinase (ERK) pathway and the downstream prosurvival and antioxidant proteins. Because electric field can be modified to focus on specific area in a non-contact manner, the study proposes that H-LIPEF holds great potential for treating NDDs, whose effect can be further augmented with the administering of drugs or natural compounds at the same time.

48

49 **Introduction**

50 Neurodegenerative diseases (NDDs) feature progressive deterioration of brain function,
51 resulting from the gradual degeneration or death of neurons. Along with increasing human
52 lifespan, deaths caused by NDDs have been on the rise over the past years, impacting the
53 quality of life of patients and their families. The World Health Organization (WHO) predicts
54 that by 2040, NDDs will overtake cancer to be the second leading cause of death, next to
55 cardiovascular diseases.

56 As the most common kind of neurodegenerative disorder among various NDDs,
57 Alzheimer's disease (AD) displays such typical symptoms as the deposition of cortical plaques,
58 formed by aggregation of β -amyloid protein ($A\beta$) or hyperphosphorylation of Tau protein,
59 leading to the formation of neurofibrillary tangles ultimately [1]. Scientists have long suspected
60 that these abnormal structures are major culprit in damaging and killing nerve cells, which
61 causes progressive brain degeneration and deterioration of cognitive function among elderly
62 people. Anti-amyloid therapies have long been pursued by scientists, however, a more nuanced
63 treatment is required because of the continuing clinical failures [2]. In addition, there has been
64 increasing evidence showing that ROS plays a central role in the onset and progression of AD
65 [3]. Therefore, drugs or treatments which could decrease the ROS level and protect neural cells
66 against oxidative damage may be potential strategies to treat AD. Some in vitro studies have

explored the curative effect of antioxidant and antiapoptotic drugs in AD treatment [4, 5]. Clinical trials were also conducted to search for antioxidant supplements or drugs to remedy AD [6-10]. However, there has yet to be a cure for AD or other kinds of NDDs. Existing medications can only avoid deterioration of symptoms or postpone their appearance at best. One blockade in AD drug development is the blood-brain barrier (BBB), which is insurmountable for 98% of small molecule drugs and 100% of large molecule drugs [11]. Therefore, effective AD treatment remains a major challenge for modern medicine.

In our previous study, non-invasive low-intensity pulsed electric field (LIPEF) was employed, in conjunction with fucoidan, in preventing oxidative stress-induced mouse motor neuron death [12]. It found that LIPEF suppressed the hydrogen peroxide (H_2O_2)-enhanced expression of ROCK protein and increased the phosphorylation of Akt in H_2O_2 -treated NSC-34 cell line. The current study employed non-contact LIPEF stimulation in studying human neural cell line SH-SY5Y, a neurodegenerative damage model with extensive application, and probed the neuroprotective effect of LIPEF against oxidative damage caused by H_2O_2 . It also applied AD disease model by $A\beta$ protein administration in neuroprotective study. Interestingly, the study finds that the LIPEF parameters applied in mouse motor neuron are not effective in human neural cell line. The systematic study on frequency- and amplitude-dependent neuroprotective effect of LIPEF was performed, and the result found that high-frequency LIPEF (H-LIPEF) at 200 Hz had the optimal protective effect for SH-SY5Y cells. Since the

alternating current (AC) signal has only one frequency and intensity, it needs time-consuming and painstaking effort for finding the specific AC electric signal to achieve a certain biological effect. Therefore, the approach has only limited applicability, as the specific AC signal is by no means sufficient in targeting complex signal transductions in a biological process or disease treatment. By contrast, the pulsed electric field (PEF) features a composition of multiple sinusoidal subcomponents with diverse frequencies and intensities simultaneously [12]. The study points out that thanks to its multi-frequency components, PEF can facilitate significantly the development of new disease therapeutics in fulfilling different demands.

In the study, we applied the H-LIPEF stimulation to human neural cell line SH-SY5Y, and examined the pro-survival effect of such physical stimulation against oxidative damage caused by H_2O_2 and $A\beta$ -induced cytotoxicity. The results found that exposure of the cells to H-LIPEF at 200 Hz significantly ameliorated the H_2O_2 - and $A\beta$ -induced cytotoxicity in SH-SY5Y cells. Examination of the underlying mechanism also showed that 200 Hz H-LIPEF could activate extracellular signal-regulated kinase (ERK) pathway and downstream neuroprotective proteins. These findings imply that H-LIPEF is a promising electrotherapy, shedding light on a novel approach in treatment of AD or other NDDs.

Materials and methods

Experimental setup for exposure of the cells to non-contact

H-LIPEF

The LIPEF device described previously [13] was used for exposure of electric field with various strengths and frequencies. The cells were placed between two copper flat and parallel electrodes. Consecutive pulses with pulse duration 2 ms, different electric field intensities and different frequencies were applied across the electrodes. Cells treated with continuous LIPEF were kept in a humidified incubator of 5% CO₂ and 95% air at 37°C. The control cells were incubated in a similar LIPEF device without the electric field turned on. After optimization of the pulse parameters, we used the H-LIPEF stimulation to produce the electric field in a non-contact manner, and examined its neuroprotection effect against the oxidative insult caused by H₂O₂ to SH-SY5Y human neural cells.

Cell culture and treatment

The SH-SY5Y cell was purchased from Bioresource Collection and Research Center (Hsinchu, Taiwan) and maintained in MEM/F-12 mixture containing 10% fetal bovine serum (FBS) (HyClone; GE Healthcare Life Sciences) and 1% penicillin-streptomycin, supplemented with 1 mM sodium pyruvate and 0.1 mM non-essential amino acids. Cells were maintained in a humidified incubator composed of 5% CO₂ and 95% air at 37°C. Cells were harvested with 0.05% trypsin–0.5 mM EDTA solution (Gibco; Thermo Fisher Scientific, Inc.) and prepared

for in vitro experiments. To differentiate SH-SY5Y cells, the cells were incubated in low serum medium (0.5% FBS) containing 20 μ M retinoic acid (RA) (Sigma-Aldrich; Merck KGaA) for 5 days. Cells were pretreated with H-LIPEF for 4 h and then challenged with 500 μ M H₂O₂ [14, 15] in the continuous administration of H-LIPEF for another 24 h. For the AD disease model, the cells were pretreated with H-LIPEF for 4 h and then challenged with 25 or 50 μ M A β ₂₅₋₃₅ protein solution [16, 17] in the continuous administration of H-LIPEF for another 4 days. The A β stock solution was prepared by solubilizing the A β ₂₅₋₃₅ peptide (Sigma-Aldrich; Merck KGaA) in sterile deionized water to a concentration of 1 mM and then incubated at 37°C for 3 days to allow self-aggregation before treatment. After the treatment, the cell viability was determined by MTT assay. A β fiberization was determined by Thioflavin T (ThT) (Sigma-Aldrich; Merck KGaA) staining. A β stock solution was diluted with PBS to 25 μ M A β solution containing 10 μ M ThT. The ThT fluorescence was then measured by SpectraMax i3x plate reader (Molecular Devices, LLC.) with excitation filter of 450 nm and emission filter of 490 nm. For the ERK inhibitor experiment, 50 μ M PD98059 (Cell Signaling Technology, Inc.) was treated 1 h before the H-LIPEF treatment.

Cell viability assay

Viability of SH-SY5Y cells after treatments with or without H-LIPEF was assessed by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma-Aldrich; Merck

KGaA) assay. Briefly, the medium was replaced with MTT solution (0.5 mg/mL in SH-SY5Y culture medium) and incubated at 37°C for 4 h. During this period, the ability of mitochondrial dehydrogenases to reduce the MTT to formazan crystal is an indicator of cellular viability. Following MTT reaction, formazan dissolution was performed by using 10% sodium dodecyl sulfate (SDS) and 0.01 M HCl. The optical density of each well was then determined at 570 nm subtracting the background at 690 nm using an ELISA microplate reader. The cell viability was calculated based on the intensity of formazan, and expressed as percentage of the non-treated control.

Flow cytometric detection of apoptotic cells

The mode of cell death was analyzed by flow cytometry with an Annexin V-FITC and propidium iodide (PI) double-staining kit (BD Biosciences). After the treatments, SH-SY5Y cells were collected by trypsinization and resuspended in binding buffer containing Annexin V and PI. Cells were incubated for 15 min at 25°C in the dark before being analyzed by flow cytometry.

ROS level detection

After the treatments, the SH-SY5Y cells were collected and the ROS levels were detected using the fluorescent dye dihydroethidium (DHE) (Sigma-Aldrich; Merck KGaA). Cells were

incubated with 5 μ M DHE dye for 20 min at 37°C in the dark and then the fluorescence intensity was measured by flow cytometry in the PE channel. The ROS levels were expressed as mean fluorescence intensity for comparison.

Mitochondrial membrane potential (MMP) measurement

The detection of MMP was carried out by flow cytometry using 3,3'-dihexyloxacarbocyanine iodide (DiOC₆(3)) (Enzo Life Sciences International Inc.). DiOC₆(3) is a lipophilic cationic fluorescent dye which is selective for the mitochondria of live cells. It allows estimation of the percentage of cells with depleted MMP which reflects the initiation of proapoptotic signal. Cells treated with H₂O₂ and H-LIPEF alone or in combination for 24 h were harvested and suspended in PBS with 20 nM DiOC₆(3). After incubation at 37°C for 15 min in the dark, the fluorescence intensity was immediately measured by flow cytometry.

Western blot analysis

After the treatments, cells were washed with phosphate buffered saline (PBS) and then harvested and lysed in RIPA lysis buffer (EMD Millipore) containing fresh protease and phosphatase inhibitor cocktail (EMD Millipore). The lysates were then centrifuged, and the supernatant was collected and used for the determination of protein concentration. Equal amounts of protein extracts were loaded in the 10% SDS-PAGE wells and transferred to

polyvinylidene fluoride membranes. After drying for 1 h at room temperature (RT), the membranes were probed with ROCK (Abcam, plc.), phosphorylated ERK (p-ERK), total ERK (t-ERK), Nrf2, phosphorylated CREB (p-CREB), total CREB (t-CREB), Bcl-2, Bax (Cell Signaling Technology, Inc.), and GAPDH (GeneTex, Inc.) antibodies overnight at 4°C. Subsequently, the washed membranes were incubated with horseradish peroxidase-coupled secondary antibodies (Jackson ImmunoResearch Laboratories, Inc.) for 1 h at RT. All the antibodies were diluted at the optimal concentration according to the manufacturer's instructions. Finally, protein bands were visualized with an enhanced chemiluminescence substrate (Advansta, Inc.) and detected with the Amersham Imager 600 imaging system (GE Healthcare Life Sciences). The images were analyzed with Image Lab software (Bio-Rad Laboratories, Inc.). For normalization of proteins, GAPDH was used as an internal control.

Statistical analysis

Each data point represents the average from larger or equal to three independent experiments and is expressed as the mean $\pm$ standard deviation. Statistical analyses were performed using OriginPro 2015 software (OriginLab). Differences of statistical significance were analyzed by one-way or two-way analysis of variance (ANOVA), followed by Tukey's post-hoc test. P-value < 0.05 was considered to indicate a statistically significant difference.

Results

In vitro-applied H-LIPEF

The H-LIPEF was applied in a non-contact manner with the apparatus consisting of two thin parallel conductive electrodes mounted on the insulating acrylic frame (Fig 1A). The culture dishes were placed between the parallel electrodes so the cells were not in contact with the electrodes directly. Therefore, the H-LIPEF stimulation device can provide the electric field in a non-invasive manner, avoiding the harmful effects caused by conduction current and Joule effect via contact electrodes. The pulsed electric signal was from a function generator and amplified by a voltage amplifier. In the study, the applied electric signal was characterized by a pulse train waveform (Fig 1B) with various field intensities (1-60 V/cm) and frequencies (2-499 Hz) for optimization of the treatment.

Fig 1. The experimental setup of H-LIPEF. (A) Image of the H-LIPEF stimulation device. The culture dishes were placed in the stimulation device constructed of two parallel electrodes. The blue arrows shown here indicate the schematic representation of non-invasive H-LIPEF exposure on cells. (B) Schematic representation of the pulse train waveform applied on the electrodes.

Effect of H-LIPEF exposure on H₂O₂-induced cytotoxicity in SH-SY5Y cells

The oxidative stress plays a critical role in the etiology and exacerbates the NDDs progression.

We applied H₂O₂ to SH-SY5Y cells as an oxidative stress and looked into the neuroprotective

effect of H-LIPEF technique. As shown in Fig 2A, administration of H₂O₂ on SH-SY5Y cells

for 24 h decreased the cell viability in a concentration-dependent manner. Treatment of

SH-SY5Y cells with 500 μ M H₂O₂ for 24 h reduced the cell viability to 52% compared to the

control cells, which was used in the subsequent experiments. To examine whether H-LIPEF

could produce protection and increase the cell viability against the oxidative insult, cells were

pretreated with H-LIPEF for 4 h and then exposed to H₂O₂ in the continuous administration of

H-LIPEF for another 24 h. In our previous study, the low frequency LIPEF with parameters of

2 Hz, 60V/cm, and 2 ms pulse duration was used in the mouse motor neurons NSC-34 [12], but

the result showed that this parameter combination is ineffective in the SH-SY5Y cells. In this

study, we used H-LIPEF in our experiments, and found that the H-LIPEF stimulation using the

parameters of 200 Hz, 10 V/cm, and 2 ms pulse duration produced significant protection

against the oxidative insult caused by H₂O₂ to SH-SY5Y human neural cell line. To examine

whether the frequency and intensity are both critical parameters in the neuroprotective effect,

the systematic study on frequency- and amplitude-dependent effects was performed. The

intensity of H-LIPEF was firstly fixed at 10 V/cm and the frequency was varied from 2 Hz to

499 Hz. The upper limit was set to 499 Hz since for 2 ms pulse duration, two pulses will overlap for frequency greater or equal to 500 Hz. The results showed that for frequency less than or equal to 100 Hz, their survival rates were not significantly different from the H₂O₂-treated group. At high frequency region, the H-LIPEF conferred protective effect to the H₂O₂-treated SH-SY5Y cells, and the maximum protective effect was at near 200 Hz (Fig 2B). For the intensity-dependent effect, the frequency was fixed at 200 Hz, and the intensity varied from 1~60 V/cm was examined. As shown in Fig 2C, it was obvious that there was a maximum at 10 V/cm, indicating that the best protective effect was at 10 V/cm. Therefore, we conclude that both the frequency and intensity are critical parameters in the neuroprotective effect, and the optimum protective parameters (200 Hz, 10 V/cm, and 2 ms pulse duration) (Fig 2D) were used in the following experiments. To further confirm the mode of cell death induced by H₂O₂, cells were stained with Annexin V-FITC and PI and assessed by flow cytometry. The results showed that H₂O₂-induced apoptotic cells were significantly reduced under H-LIPEF treatment (Fig 3), indicating SH-SY5Y cells were undergoing apoptotic cell death when treating with H₂O₂, and H-LIPEF could ameliorate the toxic effect. SH-SY5Y cell is a human neuroblastoma cell line, which is widely used as a neuronal cell model in studying NDDs. To investigate whether H-LIPEF could also provide neuroprotective effect in differentiated SH-SY5Y cells, SH-SY5Y cells were differentiated with RA for 5 days prior to H₂O₂ or H-LIPEF treatment. As shown in Fig 4A, the differentiated SH-SY5Y cells exhibited neuronal-like morphology with

high levels of neurite outgrowth. And the results showed that H-LIPEF could provide similar neuroprotective effect in the differentiated SH-SY5Y cells (Fig 4B).

Fig 2. Effect of H-LIPEF on H₂O₂-induced cytotoxicity in SH-SY5Y cells. (A)

Concentration-dependent curve of SH-SY5Y cells treated with different concentrations of H₂O₂ and the cell viability was measured by MTT assay at 24 h after the H₂O₂ treatment. (B)

SH-SY5Y cells were pretreated with 10 V/cm PEF at different pulse frequencies and challenged with 500 μ M H₂O₂ for 24 h and then the cell viability was measured by MTT assay.

(C) SH-SY5Y cells were pretreated using 200 Hz PEF with different field intensities and challenged with 500 μ M H₂O₂ for 24 h and then the cell viability was measured by MTT assay.

(D) SH-SY5Y cells were pretreated using 200 Hz H-LIPEF with field intensity of 10 V/cm and challenged with or without 500 μ M H₂O₂. The cell viability was also measured by MTT assay

at 24 h after the H₂O₂ treatment. Data represent the mean $\pm$ standard deviation (n=3).

Concentration-dependent curve of SH-SY5Y cells was statistically analyzed using one-way

ANOVA with Tukey's post hoc test (^{***}P < 0.001). Comparisons of the effect of H₂O₂ and

H-LIPEF on the cell viability were analyzed using two-way ANOVA with Tukey's post hoc test

(^{***}P < 0.001).

Fig 3. Assessment of the mode of cell death via Annexin V-FITC/PI double staining. (A)

SH-SY5Y cells were pretreated using 200 Hz H-LIPEF with field intensity of 10 V/cm and challenged with or without 500 μ M H₂O₂ for 24 h. The apoptotic cells were then assessed by flow cytometric analysis of Annexin V-FITC and PI staining. The cells in the Q2 and Q4 quadrants indicate Annexin V-positive apoptotic cells. (B) Quantification of SH-SY5Y apoptotic cells. Comparisons of the effect of H₂O₂ and H-LIPEF on the apoptotic cells were analyzed using two-way ANOVA with Tukey's post hoc test (**P < 0.001 and **P < 0.01).

Fig 4. Effect of H-LIPEF on H₂O₂-induced cytotoxicity in differentiated SH-SY5Y cells.

(A) Morphology of differentiated SH-SY5Y cells. Scale bar = 50 μ m. (B) Differentiated SH-SY5Y cells were pretreated using 200 Hz H-LIPEF with field intensity of 10 V/cm and challenged with or without 500 μ M H₂O₂ for 24 h. The cell viability was then determined by MTT assay. Comparisons of the effect of H₂O₂ and H-LIPEF on the cell viability were analyzed using two-way ANOVA with Tukey's post hoc test (***P < 0.001).

Effect of H-LIPEF exposure on A β -induced cytotoxicity in SH-SY5Y cells

For AD disease model, SH-SY5Y cells were treated with aggregated A β peptide, and the results showed that 25 or 50 μ M A β treatment significantly reduced the viability of SH-SY5Y cells (Fig 5A). To confirm the A β aggregation state, the ThT dye was used to measure the

β -sheet formation. The result showed that the fluorescence of the prepared A β stock solution was greatly higher than the ThT dye alone control group (Fig 5B), indicating that the A β stock solution was indeed in the β -sheet rich aggregated state. As shown in Fig 5C and 5D, the H-LIPEF treatment greatly improved the viability and restored the cell morphology of A β -treated SH-SY5Y cells, indicating the curative effect of H-LIPEF in AD disease model in vitro. The cell numbers were quantified using the ImageJ software [18] to investigate the effect of H-LIPEF on A β -induced cell number loss. The results showed that H-LIPEF retained the cell numbers of A β -treated SH-SY5Y cells (Fig 5E).

Fig 5. Effect of H-LIPEF on A β -induced cytotoxicity in SH-SY5Y cells. (A) The cell viability of SH-SY5Y cells treated with 25 or 50 μ M A β for 4 days. (B) The A β aggregation state was assessed by the ThT fluorescence. (C) The SH-SY5Y cells were pretreated with H-LIPEF for 4 h and then exposed to 50 μ M A β in the continuous administration of H-LIPEF for another 4 days, and the cell viability was measured by MTT assay. (D) Representative light microscopy images of SH-SY5Y cells after A β and/or H-LIPEF treatment. The integrity of the cells was destructed by A β , and the H-LIPEF treatment caused the protective effect and retained the cell morphology. Scale bar = 100 μ m. (E) Quantification of the cell numbers in the light microscopy images by ImageJ software. Data represent the mean $\pm$ standard deviation (n=3). Cell viability of SH-SY5Y cells treated with different concentration of A β and the ThT

fluorescence experiment were statistically analyzed using one-way ANOVA with Tukey's post hoc test ($^{***}P < 0.001$). Comparisons of the effect of A β and H-LIPEF on the cell viability and the cell numbers were analyzed using two-way ANOVA with Tukey's post hoc test ($^{***}P < 0.001$ and $^{**}P < 0.01$).

H-LIPEF attenuates H₂O₂-induced ROS generation

Since the onset of AD is associated with ROS accumulation, we further examined the effect of H-LIPEF on the ROS level regulation by flow cytometry (Fig 6A). Quantification results (Fig 6B) showed that the ROS level was increased to 175% of the control level after exposure to 500 μ M H₂O₂ for 24 h. Compared to the group of H₂O₂ alone, the H-LIPEF treatment greatly ameliorated the H₂O₂-induced ROS level elevation, indicating that H-LIPEF could have the ability to decrease the ROS level.

Fig 6. Effect of H-LIPEF on H₂O₂-induced ROS generation in SH-SY5Y cells. (A) ROS level was measured by flow cytometry with DHE fluorescent dye. (B) Quantification of the ROS levels after H-LIPEF, H₂O₂, or H-LIPEF+H₂O₂ treatment. Data represent the mean $\pm$ standard deviation (n=3). Comparisons of the effect of H₂O₂ and H-LIPEF on the ROS generation were analyzed using two-way ANOVA with Tukey's post hoc test ($^{***}P < 0.001$).

H-LIPEF attenuates MMP loss in H₂O₂-treated cells

Substantial evidence suggests that mitochondria play important roles in NDDs through the accumulation of net cytosolic ROS production [19]. In healthy cells, the levels of MMP are kept relatively stable, and the depletion of MMP suggests the loss of mitochondrial membrane integrity. Apoptotic signals are usually initiated by such mitochondrial dysfunctions. To examine whether the neuroprotective effect of H-LIPEF is associated with mitochondrial functions, the MMP is further analyzed by the flow cytometry with DiOC₆(3) fluorescent dye. As shown in Fig 7A, it was found that H₂O₂ caused mitochondrial dysfunction, decreasing the MMP and therefore decreased the intensity of the DiOC₆(3) fluorescent signal while H-LIPEF alone did not change the MMP. From the quantification results (Fig 7B), we found that the cells with decreased MMP drastically increased after the H₂O₂ treatment, and H-LIPEF pretreatment noticeably suppressed the H₂O₂-induced dissipation of MMP, so the cells with decreased MMP reduced significantly. The results indicate that H-LIPEF could prevent the mitochondria from damaging by the oxidative stress and thus caused the blockage of the apoptotic signaling transduction.

Fig 7. Effect of H-LIPEF on H₂O₂-induced MMP reduction in SH-SY5Y cells. (A) MMP was analyzed by flow cytometry with DiOC₆(3) fluorescent dye. (B) Quantification of the cells with decreased MMP after H-LIPEF, H₂O₂, or H-LIPEF+H₂O₂ treatment. Data represent the

mean $\pm$ standard deviation (n=3). Comparisons of the effect of H₂O₂ and H-LIPEF on the MMP were analyzed using two-way ANOVA with Tukey's post hoc test (^{***}P < 0.001 and ^{**}P < 0.01).

Effect of H-LIPEF treatment on ROCK protein expression in SH-SY5Y cells

In our previous study, the neuroprotective effect of LIPEF was found to be associated with the inhibition of the H₂O₂-induced activation of ROCK pathway in the motor neuron cell line NSC-34 [12]. In this work, we examined the ROCK protein expressions in order to investigate whether the protective effect of high frequency LIPEF on human neuroblastoma SH-SY5Y cells was related to the ROCK pathway. As shown in Fig 8, the expression of ROCK protein is slightly higher after H₂O₂ treatment but almost unaffected after the treatment of both H-LIPEF and H₂O₂. This phenomenon points out that although the H-LIPEF treatment can protect SH-SY5Y cells, the main mechanism is not associated with the ROCK pathway. Therefore, other prosurvival protein pathways should be considered in the following experiments.

Fig 8. Effect of H-LIPEF on expressions of ROCK protein in SH-SY5Y cells. (A) Western blot analysis of ROCK protein expressions. (B) Quantification of ROCK expression level after H-LIPEF, H₂O₂, or H-LIPEF+H₂O₂ treatment. The expression levels were normalized to

GAPDH and each relative expression level was compared with control. Data represent the mean $\pm$ standard deviation (n=3).

Effect of H-LIPEF exposure on ERK-Nrf2/CREB signalling pathways in H₂O₂-treated SH-SY5Y cells

We then studied the signalling pathways that could participate in the neuroprotective mechanism of H-LIPEF stimulation. ERK is one of the major cascades of mitogen-activated protein kinases (MAPK) family and has been known to be widely associated with cell survival [20]. Upon phosphorylation, the active ERK proteins enter into the nucleus where they can phosphorylate certain transcription factors, thus regulating the expression of specific genes that contribute to cell survival [21]. Therefore, in our study, we examined the phosphorylation of ERK and the downstream transcription factors including cAMP response element binding protein (CREB) and nuclear factor erythroid 2-related factor 2 (Nrf2) by western blot analysis.

In comparison to the untreated control cells, as shown in Fig 9A, the normalized level of p-ERK (p-ERK/t-ERK) was found to slightly increase in the cells exposed to H-LIPEF alone for 24 h. In contrast, the normalized level of p-ERK (p-ERK/t-ERK) in the cells was greatly decreased when exposed to 500 μ M H₂O₂ for 24 h without the treatment of H-LIPEF. In addition, the result showed that the H-LIPEF treatment induced a significant recovery of the p-ERK level in the cells exposed to 500 μ M H₂O₂. This result indicates that the

neuroprotection provided by H-LIPEF could be related to the recovery of the H₂O₂-induced decline of p-ERK. In the following, we further examined the expression levels of Nrf2 and p-CREB proteins using western blot analysis. Nrf2 is a key transcription factor that regulates the expression of antioxidant proteins participating in the protection of oxidative damage triggered by injury and inflammation. As shown in Fig 9B, the level of Nrf2 was also increased in H-LIPEF treated SH-SY5Y cells compared to the untreated cells. In contrast, the level of Nrf2 in the cells exposed to 500 μM H₂O₂ for 24 h but without the treatment of H-LIPEF was decreased. Meanwhile, the H-LIPEF treatment was found to recover the expression of Nrf2 in the cells exposed to 500 μM H₂O₂ obviously. This result showed that the neuroprotective effect was due in part to the H-LIPEF induced Nrf2 expression and the antioxidant effect. Next, CREB is also an important transcription factor triggering the expression of prosurvival proteins. As shown in Fig 9C, in comparison to the untreated control cells, the level of p-CREB was dramatically increased when cells were under the H-LIPEF treatment alone. On the other hand, the level of p-CREB in cells was decreased when exposed to 500 μM H₂O₂ for 24 h without the treatment of H-LIPEF. Moreover, the result found that the H-LIPEF treatment obviously recovered the expression of p-CREB in the cells exposed to 500 μM H₂O₂. Furthermore, it is widely known that CREB phosphorylation regulates Bcl-2 family proteins including anti-apoptotic Bcl-2 and pro-apoptotic Bax proteins, and the ratio of Bcl-2/Bax determines whether neurons undergo survival or death after an apoptotic stimulus. Therefore, the

expressions of Bcl-2 and Bax protein levels were also examined in the study. As shown in Fig 9D, for SH-SY5Y cells under the H-LIPEF treatment alone, the expression levels of Bcl-2 and Bax were increased and decreased respectively, leading to a higher elevation of the Bcl-2/Bax ratio compared to the untreated control cells. Meanwhile, in comparison to the untreated control cells, the level of Bcl-2/Bax ratio of SH-SY5Y cells was found to decrease when exposed to 500 μ M H₂O₂, indicating that the cells had undergone apoptosis. Moreover, it was observed that the H-LIPEF treatment notably restored the expression of Bcl-2/Bax ratio in the cells exposed to 500 μ M H₂O₂. To sum up, the results showed that the H-LIPEF treatment induced the anti-apoptotic effect by regulating the Bcl-2 family proteins via modulating the ERK/CREB pathways and thus promoted the cell survival against oxidative stress.

Fig 9. Effect of H-LIPEF on ERK/Nrf2 and ERK/CREB signalling pathways and related protein expressions. Western blot analysis of p-ERK (A), Nrf2 (B), p-CREB (C), Bcl-2 and Bax (D) protein expressions. The expression levels of p-ERK and p-CREB were normalized to t-ERK and t-CREB, respectively. The expression level of Nrf2 was normalized to GAPDH and each relative expression level was compared with control. Data represent the mean $\pm$ standard deviation (n=3). Comparisons of the effect of H₂O₂ and H-LIPEF on different protein levels were analyzed using two-way ANOVA with Tukey's post hoc test (*** P < 0.001, ** P < 0.01 and * P < 0.05).

Effect of PD98059 on the H-LIPEF activated ERK pathway and protective effect

To confirm the neuroprotective effect of H-LIPEF is related to the ERK pathway, we inhibited the MAPK kinases MEK1 and MEK2, the upstream kinases of ERK, by the inhibitor PD98059 [22] in the study. In Fig 10A, we found that H-LIPEF could significantly increase the cell viability of SH-SY5Y cells to about 80% of the control level under H₂O₂ stress. In the presence of MEK inhibitor, the p-ERK level was indeed reversed and the neuroprotective effect of H-LIPEF treatment against H₂O₂-induced cell apoptosis was abrogated (Fig 10A and 10B). The results indicate that the ERK pathway was involved to mediate the neuroprotective effect of H-LIPEF treatment against H₂O₂ toxicity. On the whole, the data point out that H-LIPEF treatment may produce neuroprotective effect via activation of the ERK signalling pathway in the H₂O₂-treated SH-SY5Y human neural cells.

Fig 10. Effect of PD98059 on the neuroprotective effect of H-LIPEF and p-ERK protein expression induced by H-LIPEF stimulation. (A) H-LIPEF increased the cell viability of SH-SY5Y cells under H₂O₂ oxidative stress. The neuroprotective effect of H-LIPEF treatment was abrogated by addition of the MEK inhibitor PD98059. (B) The western blot analysis showed that the inhibitor PD98059 reversed the activated level of p-ERK induced by H-LIPEF

treatment. The expression levels were normalized to t-ERK. We used the abbreviation “PD” to represent the MEK inhibitor PD98059 in the figure. Data represent the mean $\pm$ standard deviation (n=3). Comparisons of the effect of PD on the neuroprotective effect of H-LIPEF and p-ERK protein levels were analyzed using two-way ANOVA with Tukey’s post hoc test ($^{***}P < 0.001$ and $^{**}P < 0.01$).

Discussion

The focus of this study was to probe the neuroprotective effect of H-LIPEF on SH-SY5Y cells. The results demonstrated that application of high-frequency (~200 Hz) and low-intensity (~10 V/cm) pulsed electric field produced the most significant protection for SH-SY5Y human neural cells against H₂O₂- and A β -induced apoptosis. The western blot results exhibited that H-LIPEF triggered the ERK signalling pathway and mediated the activation of CREB and Nrf2 transcription factors.

As the most common form of age-related NDDs, AD features excessive aggregation of A β and intracellular microtubule-associated protein tau, known as neurofibrillary tangles. With many questions remaining unanswered, some researchers believe that the amyloid cascade hypothesis offers the best explanation for the cause of AD [23], maintaining that A β is the causative factor of Alzheimer’s pathology and that the neurofibrillary tangles, cell loss, vascular damage, and dementia all result directly from such deposition. Although the

466 mechanism of A β -induced cytotoxicity has yet to be elucidated completely, several studies
 467 have proven that the oxidative stress plays a vital role in A β -induced apoptosis [24-26].
 468 Therefore, researchers have been striving to find various antioxidants that can protect against
 469 oxidative stress and alleviate the symptoms of NDDs [27-29]. However, the therapeutic effect
 470 of most antioxidant compounds is limited, due to their incapability to cross the BBB. And the
 471 antioxidant therapies have continually failed in treatment of AD. Therefore, effective AD
 472 treatment strategy is urgently needed for modern medicine. Among different ROS species,
 473 H₂O₂ is one of the molecules most related to the progression of AD. In the study, H₂O₂ was
 474 applied as the oxidative stress which caused cell death at certain dosage. The results found that
 475 H₂O₂ treatment increased the ROS level and H-LIPEF treatment significantly ameliorated the
 476 H₂O₂-induced ROS level elevation, underscoring the antioxidant ability of H-LIPEF. The study
 477 employed MTT and MMP assays to assess the cell viability and mitochondria integrity, finding
 478 in both results that H-LIPEF provided the neuroprotective effect and increased the viability and
 479 MMP in the H₂O₂-treated neural cells. Furthermore, A β was used to investigate the
 480 neuroprotective effect of H-LIPEF in the AD disease model. The MTT and light microscopy
 481 results also showed that H-LIPEF greatly improved the viability and retained the cell integrity
 482 in the A β -treated cells. Since the electric field can penetrate the human body without the
 483 restriction of BBB, H-LIPEF promises a potent alternative therapeutic medicine in NDDs
 484 treatment.

485 The electric stimulation has a long history of medical applications, including wound
486 healing [30], muscle relaxation [31], and increase of blood circulation [32], often involving
487 passage of electric current through the injury parts via direct contact between the body and
488 implanted electrodes. However, the invasive or contact manner significantly dampens the
489 feasibility of the treatment or even causes harms to cells or body, such as electrolysis, burns
490 and muscle spasms [33]. Therefore, it is desirable for electrotherapy to be carried out in a
491 non-invasive or non-contact manner. Our previous study demonstrated a non-contact electric
492 stimulation device [13], which operates as a parallel-plate capacitor generating electric field
493 between two parallel electrodes (Fig 1A). Therefore, the electric field with such capacitive
494 coupling mode can pass through biological samples to achieve specific biological effect
495 without current actually flowing through cells or media via contact with electrodes. It is
496 noteworthy that the strength of electric field passing through culture dishes or media is the
497 same as the voltage for electrodes. Besides, the electric field waveform employed was a PEF,
498 which is characterized by its multi-frequency component nature. Pulse train signal is composed
499 of multiple sinusoidal subcomponents with different frequencies and intensities [12]. When
500 PEF enters a biological sample, it encounters different proteins, cells and tissues with different
501 dielectric constants and is therefore dispersed to multiple frequencies electrically.
502 Multi-frequency sinusoidal waves may interact with molecules, proteins, or organelles
503 simultaneously to induce specific bioeffect in the PEF exposure. In our previous studies, the

504 non-contact LIPEF exhibited promising potential in activating anticancer agent in pancreatic
505 cancer treatment [34] and also in the neuroprotection of mouse motor neurons [12]. In the
506 current study, LIPEF with different parameters was applied to the human neural cells for
507 examination of its neuroprotective effect. Unexpectedly, the LIPEF parameter applied in mouse
508 motor neurons NSC-34 turned out to be ineffective for human SH-SY5Y cells. In fact,
509 scientists have found that mouse-based model is not applicable to humans [35], which provides
510 the reason why some drugs or treatments that successfully treat mice do not work for people.
511 Therefore, it is critical to find treatments for animal models or cell types closely related to
512 humans. The study found that for human SH-SY5Y cells, the neuroprotective effect of LIPEF
513 was more pronounced in the high-frequency region, especially near 200 Hz, quite different
514 from the parameter at 2 Hz for NSC-34 mouse cell line. For higher-frequency LIPEF, it
515 consists more high-frequency components in its composition. Therefore, the neuroprotective
516 effect of H-LIPEF for human SH-SY5Y neural cells may be attributable to the employment of
517 high-frequency components participating in the treatment. The results also showed that the
518 therapeutic window of the H-LIPEF intensity for SH-SY5Y cells was near 10 V/cm, which was
519 smaller than the most effective parameter (60 V/cm) for NSC-34 cells. Therefore, it is evident
520 that the pulse frequency and the field intensity are critical parameters for therapeutic
521 applications. The parameters (frequency, intensity, and pulse duration) of LIPEF need to be
522 fine-tuned and optimized for different types of cells or diseases. Furthermore, the study

proposes combination of multiple pulses with different parameters to treat the targeted region simultaneously.

As for the neuroprotective mechanism, it was found previously that the LIPEF treatment obviously suppressed the H₂O₂-enhanced expression of ROCK protein in NSC-34 cells. However, for SH-SY5Y cells, the H-LIPEF simulation did not have significant impact on ROCK protein expression. It is thus known that the neuroprotective mechanism is not through the ROCK pathway. In the western blot results, the study proved that such neuroprotective effect induced by H-LIPEF was through activation of ERK pathway. The ERK pathway plays an important role in regulating of proliferation and differentiation as well as survival of various cell types [36]. It transmits external cues, including growth factors into signalling cascade via the phosphorylation of ERK. Several studies have suggested that the stimulation of ERK pathway may be a potential way for neuroprotection [20, 21]. Here, the study showed that the H-LIPEF stimulation induced a significant recovery of the p-ERK level in the cells under H₂O₂ oxidative stress. The result indicates that the neuroprotection effect produced by H-LIPEF could be related to the recovery of the H₂O₂-induced reduction of p-ERK. Moreover, it should be noted that the normalized p-ERK level (p-ERK/t-ERK) stimulated by H-LIPEF alone was only slightly higher than the untreated control cells without significant difference (Fig 9A). The study found that the slight enhanced level of p-ERK did not cause cell proliferation as assessed by MTT assay (Fig 2D). Therefore, the result confirmed that the neuroprotective effect of

H-LIPEF stimulation was not caused by the increased cell numbers, but by specific protein pathways induced by H-LIPEF on SH-SY5Y cells. Furthermore, since the ERK pathway is initiated by the binding of epidermal growth factor (EGF) to the epidermal growth factor receptor (EGFR), we speculate that H-LIPEF could have an interaction with EGFR and cause the conformation change of EGFR, thus triggering the ERK pathway. Previous study had shown that magnetic field could alter the conformation of TRPV1 ion channel, making it more favorable for capsaicin binding [37]. The results implicate that the PEF or magnetic field with different parameters could activate different signalling pathways. In the study, H-LIPEF with 200 Hz frequency and 10 V/cm field intensity was found to significantly recover the p-ERK protein level and produce the neuroprotective effect. The phosphorylation of ERK protein could result in the activation of its kinase activity and lead to phosphorylation of many downstream targets including certain transcription factors containing CREB and Nrf2 required for neuroprotection.

CREB is an important transcription factor activated by phosphorylation from ERK or other kinases. It triggers expression of neurotrophins and antiapoptotic proteins such as brain-derived neurotrophic factor (BDNF) and B-cell lymphoma 2 (Bcl-2) [38]. Besides, CREB downregulation is related to the pathology of AD [39], and thus increasing the CREB expression has been considered as a potential therapeutic strategy for AD. In this study, the result found that the H-LIPEF treatment notably recovered the expression of p-CREB in the

cells exposed to 500 μM H_2O_2 . On one hand, it is known that CREB phosphorylation regulates Bcl-2 family proteins including anti-apoptotic Bcl-2 and pro-apoptotic Bax proteins. The antiapoptotic protein, Bcl-2, has been proposed to be a pivotal component for neuroprotection [40]. It suppresses apoptosis by heterodimerizing with Bax protein and neutralizes the effects of Bax. When Bcl-2 is present in excess, cells are protected against apoptosis. It should be also pointed out that when Bax is excessive and the homodimers of Bax dominate, cells are susceptible to programmed cell death. Therefore, it is the ratio of Bcl-2 to Bax which determines the fate of a cell [41]. In the results, we found that the H-LIPEF treatment obviously restored the expression of Bcl-2/Bax ratio in the cells exposed to 500 μM H_2O_2 , indicating the prosurvival effect of H-LIPEF in SH-SY5Y cells.

Another possible mechanism for neuroprotection of the H-LIPEF stimulation may be associated with the Nrf2 protein. Nrf2 is also an important transcription factor activated by ERK pathway. It regulates the expression of various antioxidant proteins and maintains their redox states in mammalian cells [42]. The results showed that the H-LIPEF treatment obviously recovered the expression of Nrf2 in the cells exposed to 500 μM H_2O_2 , indicating that the neuroprotective effect was due in part to the H-LIPEF induced Nrf2 expression and the antioxidant effect.

In the study, our results confirmed the neuroprotective effect of H-LIPEF stimulation on SH-SY5Y cells. Because electric field can be modified to focus on a targeted region, it affirms

the applicability of H-LIPEF treatment for different human organs. Moreover, it found that H-LIPEF technique can significantly reduce treatment risk and time, thanks to its non-contact and multi-frequency nature. In addition, the study also demonstrated the ability of H-LIPEF in rescuing the A β -induced cytotoxicity. Since the accumulation of misfolded proteins in brain is also a typical symptom of some NDDs, the study could shed light on future exploration for optimal LIPEF conditions in the treatment of other NDDs.

In summary, the study demonstrates for the first time that the physical stimulation by H-LIPEF can protect SH-SY5Y human neural cells from H₂O₂- and A β -induced cytotoxicity. The underlying molecular mechanism for the neuroprotective effect is partially attributed to the activation of ERK pathway and the function of some downstream proteins, such as CREB, Nrf2, and Bcl-2. Further studies are necessary to ascertain the optimal LIPEF conditions for other NDDs and the involvement of other proteins in the neuroprotective effect. The combinations of H-LIPEF with NDD drugs or natural compounds are also advisable for future studies.

Funding

This work was supported by grants from Ministry of Science and Technology (MOST 109-2112-M-002-004, MOST 108-2112-M-002-016, and MOST 105-2112-M-002-006-MY3 to CYC) and Ministry of Education (MOE 106R880708 to CYC) of the Republic of China. The

fundere had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests

The authors have declared that no competing interests exist.

Acknowledgments

The authors would like to acknowledge the service provided by the Research Core Facilities 3 Laboratory of the Department of Medical Research at National Taiwan University hospital for use of flow cytometry system.

Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Wei-Ting Chen, Guan-Bo Lin, Yu-Yi Kuo, Chih-Yu Chao.

Formal analysis: Wei-Ting Chen, Guan-Bo Lin, Yu-Yi Kuo, Chih-Hsiung Hsieh, Chueh-Hsuan Lu, Yi-Kun Sun, Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Wei-Ting Chen, Guan-Bo Lin, Yu-Yi Kuo, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

618 **Supervision:** Chih-Yu Chao.

619 **Validation:** Wei-Ting Chen, Guan-Bo Lin, Yu-Yi Kuo.

620 **Writing – original draft:** Wei-Ting Chen, Guan-Bo Lin, Chih-Yu Chao.

621 **Writing – review & editing:** Chih-Yu Chao.

622

623

624 **References:**

625

626 1. Ross CA, Poirier MA. Protein aggregation and neurodegenerative disease. Nat Med. 2004;
627 10:S10-S17.

628 2. Mullane K, Williams M. Alzheimer's disease beyond amyloid: Can the repetitive failures
629 of amyloid-targeted therapeutics inform future approaches to dementia drug discovery?
630 Biochem Pharmacol. 2020; 177:113945.

631 3. Kim GH, Kim JE, Rhie SJ, Yoon S. The role of oxidative stress in neurodegenerative
632 diseases. Exp Neurol. 2015; 24:325-340.

633 4. Hemming ML, Selkoe DJ. Amyloid beta-protein is degraded by cellular
634 angiotensin-converting enzyme (ACE) and elevated by an ACE inhibitor. J Biol Chem. 2005;
635 280:37644-37650.

636 5. Alghazwi M, Smid S, Musgrave I, Zhang W. In vitro studies of the neuroprotective
637 activities of astaxanthin and fucoxanthin against amyloid beta (A β 1-42) toxicity and
638 aggregation. Neurochem Int. 2019; 124:215-224.

639 6. Goozee K, Shah T, Sohrabi HR, Rainey-Smith S, Brown B, Verdile G, et al. Examining
640 the potential clinical value of curcumin in the prevention and diagnosis of Alzheimer's disease.
641 Br J Nutr. 2016; 115:449-465.

642 7. Bachurin SO, Bovina EV, Ustyugov AA. Drugs in clinical trials for Alzheimer's disease:
643 the major trends. Med Res Rev. 2017; 37:1186-1225.

644 8. Kryscio RJ, Abner EL, Caban-Holt A, Lovell M, Goodman P, Darke AK, et al.
645 Association of antioxidant supplement use and dementia in the prevention of Alzheimer's
646 disease by vitamin E and selenium trial (PREADVISE). JAMA Neurol. 2017; 74:567-573.

647 9. Farina N, Llewellyn D, Isaac MGEKN, Tabet N. Vitamin E for Alzheimer's dementia and
648 mild cognitive impairment. Cochrane Database Syst Rev. 2017; 4:CD002854.

649 10. Polidori MC, Nelles G. Antioxidant clinical trials in mild cognitive impairment and

- 650 Alzheimer's disease - challenges and perspectives. *Curr Pharm Design*. 2014; 20:3083-3092.
- 651 11. Yang Y, Zhang X, Ye D, Laforest R, Williamson J, Liu Y, et al. Cavitation dose painting
652 for focused ultrasound-induced blood-brain barrier disruption. *Sci Rep*. 2019; 9:2840.
- 653 12. Hsieh CH, Lu CH, Kuo YY, Lin GB, Chao CY. The protective effect of non-invasive low
654 intensity pulsed electric field and fucoidan in preventing oxidative stress-induced motor neuron
655 death via ROCK/Akt pathway. *PLoS One*. 2019; 14:e0214100.
- 656 13. Hsieh CH, Lu CH, Chen WT, Ma BL, Chao CY. Application of non-invasive low strength
657 pulsed electric field to EGCG treatment synergistically enhanced the inhibition effect on
658 PANC-1 cells. *PLOS ONE*. 2017; 12:e0188885.
- 659 14. Wang CM, Yang CQ, Cheng BH, Chen J, Bai B. Orexin-A protects SH-SY5Y cells
660 against H₂O₂-induced oxidative damage via the PI3K/MEK1/2/ERK1/2 signaling pathway. *Int*
661 *J Immunopathol Pharmacol*. 2018; 32:2058738418785739.
- 662 15. Zhao J, Liu Y, Wei X, Yuan C, Yuan X, Xiao X. A novel WD-40 repeat protein WDR26
663 suppresses H₂O₂-induced cell death in neural cells. *Neurosci Lett*. 2009; 460:66-71.
- 664 16. Gray NE, Morre J, Kelley J, Maier CS, Stevens JF, Quinn JF, et al. Caffeoylquinic acids in
665 *Centella asiatica* protect against amyloid- β toxicity. *J Alzheimers Dis*. 2014; 40:359-373.
- 666 17. Zhang JJ, Zhang RF, Meng XK. Protective effect of pyrroloquinoline quinone against
667 A β -induced neurotoxicity in human neuroblastoma SH-SY5Y cells. *Neurosci Lett*. 2009;
668 464:165-169.
- 669 18. Venter C, Niesler C. Rapid quantification of cellular proliferation and migration using
670 ImageJ. *Biotechniques*. 2019; 66:99-102.
- 671 19. Manczak M, Anekonda TS, Henson E, Park BS, Quinn J, Reddy PH. Mitochondria are a
672 direct site of A β accumulation in Alzheimer's disease neurons: implications for free radical
673 generation and oxidative damage in disease progression. *Hum Mol Genet*. 2006;
674 15:1437-1449.
- 675 20. Mohamed SK, Ahmed AAE, Elmorsy EM, Nofal S. ERK activation by zeranol has
676 neuroprotective effect in cerebral ischemia reperfusion. *Life Sci*. 2019; 227:137-144.
- 677 21. Almeida RD, Manadas BJ, Melo CV, Gomes JR, Mendes CS, Grãos MM, et al.
678 Neuroprotection by BDNF against glutamate-induced apoptotic cell death is mediated by ERK
679 and PI3-kinase pathways. *Cell Death Differ*. 2005; 12:1329-1343.
- 680 22. Caunt CJ, Sale MJ, Smith PD, Cook SJ. MEK1 and MEK2 inhibitors and cancer therapy:
681 the long and winding road. *Nat Rev Cancer*. 2015; 15:577-592.
- 682 23. Karran E, Mercken M, Strooper BD. The amyloid cascade hypothesis for Alzheimer's
683 disease: an appraisal for the development of therapeutics. *Nat Rev Drug Discov*. 2011;
684 10:698-712.
- 685 24. Cheignon C, Tomas M, Bonnefont-Rousselot D, Faller P, Hureau C, Collin F. Oxidative
686 stress and the amyloid beta peptide in Alzheimer's disease. *Redox Biol*. 2018; 14:450-464.
- 687 25. Butterfield DA, Swomley AM, Sultana R. Amyloid β -peptide (1-42)-induced oxidative

688 stress in Alzheimer disease: importance in disease pathogenesis and progression. *Antioxid*
689 *Redox Signal*. 2013; 19:823-835.

690 26. Arimon M, Takeda S, Post KL, Svirsky S, Hyman BT, Berezovska O. Oxidative stress and
691 lipid peroxidation are upstream of amyloid pathology. *Neurobiol Dis*. 2015; 84:109-119.

692 27. Muthaiyah B, Essa MM, Chauhan V, Chauhan A. Protective effects of walnut extract
693 against amyloid beta peptide-induced cell death and oxidative stress in PC12 cells. *Neurochem*
694 *Res*. 2011; 36:2096-2103.

695 28. Wang C, Li J, Liu Q, Yang R, Zhang JH, Cao YP, et al. Hydrogen-rich saline reduces
696 oxidative stress and inflammation by inhibit of JNK and NF- κ B activation in a rat model of
697 amyloid-beta-induced Alzheimer's disease. *Neurosci Lett*. 2011; 491:127-132.

698 29. Chan KH, Lam KSL, Cheng OY, Kwan JSC, Ho PWL, Cheng KKY, et al. Adiponectin is
699 protective against oxidative stress induced cytotoxicity in amyloid-beta neurotoxicity. *PLoS*
700 *One*. 2012; 7:e52354-e52354.

701 30. Gentzkow GD, Miller KH. Electrical stimulation for dermal wound healing. *Clin Podiatr*
702 *Med Surg*. 1991; 8:827-841.

703 31. Windsor RE, Lester JP, Herring SA. Electrical stimulation in clinical practice. *Physician*
704 *Sportsmed*. 1993; 21:85-93.

705 32. Park RJ, Son H, Kim K, Kim S, Oh T. The effect of microcurrent electrical stimulation on
706 the foot blood circulation and pain of diabetic neuropathy. *J Phys Ther Sci*. 2011; 23:515-518.

707 33. Antov Y, Barbul A, Mantsur H, Korenstein R. Electroendocytosis: exposure of cells to
708 pulsed low electric fields enhances adsorption and uptake of macromolecules. *Biophys J*. 2005;
709 88:2206-2223.

710 34. Lu CH, Lin SH, Hsieh CH, Chen WT, Chao CY. Enhanced anticancer effects of low-dose
711 curcumin with non-invasive pulsed electric field on PANC-1 cells. *OncoTargets Ther*. 2018;
712 11:4723-4732.

713 35. Uhl EW, Warner NJ. Mouse models as predictors of human responses: evolutionary
714 medicine. *Curr Pathobiol Rep*. 2015; 3:219-223.

715 36. Sun Y, Liu WZ, Liu T, Feng X, Yang N, Zhou HF. Signaling pathway of MAPK/ERK in
716 cell proliferation, differentiation, migration, senescence and apoptosis. *J Recept Signal*
717 *Transduct*. 2015; 35:600-604.

718 37. Chen WT, Lin GB, Lin SH, Lu CH, Hsieh CH, Ma BL, et al. Static magnetic field
719 enhances the anticancer efficacy of capsaicin on HepG2 cells via capsaicin receptor TRPV1.
720 *PLoS One*. 2018; 13:e0191078.

721 38. Finkbeiner S. CREB couples neurotrophin signals to survival messages. *Neuron*. 2000;
722 25:11-14.

723 39. Rosa E, Fahnstock M. CREB expression mediates amyloid β -induced basal BDNF
724 downregulation. *Neurobiol Aging*. 2015; 36:2406-2413.

725 40. Howard S, Bottino C, Brooke S, Cheng E, Giffard RG, Sapolsky R. Neuroprotective

726 effects of bcl-2 overexpression in hippocampal cultures: interactions with pathways of
 727 oxidative damage. J Neurochem. 2002; 83:914-923.
 728 41. Pepper C, Hoy T, Bentley DP. Bcl-2/Bax ratios in chronic lymphocytic leukaemia and
 729 their correlation with in vitro apoptosis and clinical resistance. Br J Cancer. 1997; 76:935-938.
 730 42. Kubben N, Zhang W, Wang L, Voss TC, Yang J, Qu J, et al. Repression of the antioxidant
 731 NRF2 pathway in premature aging. Cell. 2016; 165:1361-1374.

Figure 1

A

bioRxiv preprint doi: <https://doi.org/10.1101/2019.12.30.891499>; this version posted October 7, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

parallel electrode

acrylic frame

parallel electrode

B

Figure 2

Figure 3

bioRxiv preprint doi: <https://doi.org/10.1101/2019.12.30.891499>; this version posted October 7, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Figure 4

bioRxiv preprint doi: <https://doi.org/10.1101/2019.12.30.891499>; this version posted October 7, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

A

B

Figure 5

bioRxiv preprint doi: <https://doi.org/10.1101/2019.12.30.891499>; this version posted October 7, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Figure 6

bioRxiv preprint doi: <https://doi.org/10.1101/2019.12.30.891499>; this version posted October 7, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Figure 7

Figure 8

A

bioRxiv preprint doi: <https://doi.org/10.1101/2019.12.30.891499>; this version posted October 7, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

B

Figure 9

A

B

C

D

Figure 10

bioRxiv preprint doi: <https://doi.org/10.1101/2019.12.30.891499>; this version posted October 7, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.