In-stream habitat availability for river dolphins in response to flow: use of ecological integrity to manage river flows

Shambhu Paudel^{ab*}, John L. Koprowski^{c,d}, & Usha Thakuri^b

^{a,c} School of Natural Resources and the Environment, University of Arizona, 1064 East Lowell Street, Tucson, Arizona 85721, USA

^b Tribhuvan University, Institute of Forestry, Post Box:43, Pokhara, Nepal

^d Haub School of Environment and Natural Resources, University of Wyoming, Bim Kendall House, 804 E Fremont St, Laramie, WY 82072

*Corresponding author

Email addresses:

^aspaudel@email.arizona.edu ^c5quirrel@ag.arizona.edu

Abstract

Population decline and extinction risk of riverine cetaceans are primarily associated with flow alteration. Previous studies predominantly highlighted maintenance of adequate flow for low water seasons when habitats contract and the risk of local extinction escalates. Although riverine cetaceans are sensitive to reduction in river flow, no studies quantify the relationships between flow and ecology of riverine cetaceans to mitigate the potential adverse impacts of flow alteration. We quantify the relationships between flow and the ecology of river cetaceans with reference to Ganges River dolphins (GRD; *Platanista gangetica gangetica*) usable area availability (AWS) for the low water season at wider flows (50-575 m³/s) at finer spatial and temporal scales. This study reveals that distribution of usable area to GRD highly regulated by the adequate flow level as the interaction of appropriate velocity and depth at particular geomorphic that likely offer energetically efficient modes of locomotion to increase surfacing behavior efficiency, suggesting the hydro-physical environment as a major determinant of river cetaceans distribution and abundance. Flow and AWS relationships indicate that the flow during dry season negatively contributed to AWS, whereas that of pre-monsoon maximized the AWS, suggesting that modifying flow regimes does alter in-stream habitats at varying spatial scales and may influence life-history strategies. Substantial fragmentation in suitable pool availability and loss of longitudinal connectivity exhibited by dry season flow suggested a higher risk of adverse biological effects during the dry season, which may reduce population viability by reducing survivorship and reproduction failure. Owing to riverine cetaceans' dependence on the attribute of freshwater flow, they can be expected to be more affected by flow regulations as interactive effects. Considering the seasonal effects and changes in the availability of usable areas by flow alteration, adopting effective habitat retention plans by water-based development projects appears critical to avoid further ecological risks in aquatic species conservation. Identifying priority riverscapes for river cetaceans and prioritizing investment opportunities is an essential first step towards effective riverine cetacean conservation.

Introduction

As the human population size increases, freshwater species and ecosystems are increasingly threatened by many development activities, including habitat alteration, river diversions, fragmentation and flow regulation, expansion of agricultural and urban landscapes, and climate change [1]. All these anthropogenic consequences determine the quality and quantity of freshwater, which affects ecological processes and dynamics that determine freshwater ecosystem productivity and integrity. As the withdrawals and diversions are the main threats to freshwater availability and quality [2, 3, 4], freshwater biodiversity is primarily threatened by habitat fragmentation and flow regulation [5]. Worldwide, agriculture accounts for about 70% of all freshwater usage, compared to 20% and 10% by industry and domestic use, respectively [6]. Accelerated water extraction or diversion combined with changes in climate, human population, and changing land use puts immense pressure on river basin biodiversity and ecosystem services through habitat loss and fragmentation [7].

Globally, a sharp decline in freshwater species, especially megafauna with large body size (adult body weight ≥ 30 kg) and complex habitat requirements, is commonly attributed to water diversion and extraction. From 1970 to 2012, populations of vertebrate freshwater declined by 88% due to habitat degradation induced by flow regulation [8, 9]. Several river basins in Southern Asia (Yangtze River, Ganges River, Indus River, Brahmaputra River) and Northern South America (Amazon, Orinoco, and Tocantins-Araguaia) have been impaired by large water extraction projects (i.e., hydropower and agriculture irrigation projects) with negative impacts on aquatic species conservation [1, 10, 11]. Such mega-projects threaten to freshwater species with extinction, isolation, and reduction in abundance and range, especially in the case of apex predators and highly migratory riverine cetaceans [12, 13, 10, 11]. As river cetaceans are sensitive to river flow, reduced in-stream habitat quality due to the lack of adequate river flow further escalated endangerment and extinction risk by increasing entanglement and adverse biological effects through fisheries-freshwater cetaceans interactions [14]. Risks to riverine cetaceans are more severe during the low water season when their habitats are fragmented and when dispersal ability is reduced [15]. Extinction of Yangtze River dolphins and a sharp decline in the populations of several riverine cetaceans [Ganges River dolphin (*Platanista gangetica gangetica*), Indus River dolphin (*Platanista gangetica minor*), Irrawaddy dolphin (*Orcaella*

brevirostris), Amazonian river dolphin (*Inia geoffrensis*), Bolivian river dolphin (*Inia boliviensis*), Tucuxi (*Sotalia fluviatilis*), Araguaian boto (*Inia araguaiaensis*)] are the example of consequences of inadequate flow in the Anthropocene [16, 17, 18, 13, 10, 11].

As habitat loss and fragmentation had significant adverse biological impacts on river cetaceans, previous studies recognized the sensitivity (presence/absence) of riverine cetaceans to flow and their hydraulic properties (i.e., velocity, depth, and cross-sectional area) [19, 20]. Even though river flow directly regulates in-stream habitat structure that has a substantial impact on the organization and structure of biological communities [21], relationships between flow and ecological requirements of river cetaceans (particularly flows that determine the suitably usable area to riverine cetaceans) have never been discussed and quantified. At the same time, appropriation of freshwater flows urged globally to sustain societal benefits and freshwater biodiversity by balancing the needs of humans, freshwater species and ecosystems by setting limits for freshwater withdrawals and diversions [22]. Thus, there is an immediate need to understand the relationships between flow and riverine cetaceans' hydro-physical requirements at a finer scale to understand the effects of flow alternation on in-stream habitat availability while balancing riverine cetaceans' conservation and societal demands in the Anthropocene.

Previous studies on aquatic ecosystem species conservation heavily consider ecological requirements of fish communities, thereby generalizing the flows over the entire ecosystem [23, 24]. Such an estimated flow for fish communities may not be ecologically adequate and relevant to the species that are at the top trophic level with large body size, i.e., freshwater cetaceans. Furthermore, studies on habitat selection by freshwater cetaceans conservatively link presence and abundance data to coarse-scale (~1-3 km) and satellite image driven hydro-physical variables and visually characterize habitat structure [South Asian river dolphins [19, 20, 25], Irrawaddy water dolphin [19, 26], Amazonian river dolphin [15, 27], Bolivian river dolphin [28], Yangtze finless porpoise [29], Tucuxi [15] with hydro-physical habitat preference, however do not assess quality and quantity of the hydro-physical parameters at discrete points occupied by river cetaceans. Even though species are sensitive to certain flow velocities and depths [30], previous studies on freshwater cetaceans did not account for these parameters when examining the cetaceans' habitat preferences as these parameters play an important role in determining the availability of suitable in-stream habitat [20]. Lack of empirical relationships between river flow

and riverine cetaceans' ecological responses may further elevate extinction risks by limiting our ability to establish thresholds on environmental flows that sustain remaining riverine endangered cetaceans.

We quantify the relationships between flow and river cetacean ecology to determine the availability of suitable in-stream physical habitat as a response to flow. These results on an apex predator may contribute to maintaining the aquatic integrity by offering delineation of suitable habitats and enhancing longitudinal connectivity that facilitates species dispersal across the diverse hydro-physical habitats. We tested the effect of the individual hydro-physical parameters (flow, depth, velocity, cross-sectional area, width, and wetted perimeter) on the availability of the in-stream usable habitat based on the preference of a riverine apex predator, Ganges River dolphins (GRD), for a wide range of hydro-physical parameters measured at fine spatial and temporal scale. We assumed that GRD prefers a particular combination (velocity and depth at a specific flow) of the abiotic environment that determines the quality and availability of in-stream physical habitat. We, thus, developed the Habitat Suitability Curve (HSC) using a combination of the depth and velocity preferred by GRD, which allows us to quantify the availability of suitable in-stream habitat in the form of area-weighted suitability (AWS, m^2/m). Our study quantifies the hydro-physical habitat selection of GRD in relation to properties of wider level river flow, which could be essential to determine adequate river flows that are particularly significant for the conservation of in-stream habitats and critical to endangered freshwater cetaceans' life activities during the low water season. Thus, the results of this study are essential to balance aquatic species' conservation while minimizing the ecological risks induced through water extraction or diversion by development projects.

Materials and methods

Study area

This study was conducted in the Karnali River system of Nepal, where mega hydropower projects are under construction, and several are proposed and completed (Fig 1). Details about the current water-based projects and ecological significance are described in [10]. This river segment, particularly downstream, is highly significant to endangered aquatic species like GRD

(*Platanista gangetica gangetica*), Gharial (*Gavialis gangeticus*), Smooth Indian otter (*Lutra perspicillata*), and diverse fish species. Current water extraction or diversion projects typically adopt a traditional biodiversity impact assessment but do not assess more detailed habitat relationships that include flow [31]. Additionally, water management plans adopted by such mega-projects are based on the proportion of total discharge at a particular season or time, undermining the aquatic species' ecological requirements across space and time. Thus, balancing aquatic species conservation and development is the foremost priority of the concerned stakeholders.

Fig 1. A study map showing the location of development structures and segments took for hydro-physical measurements.

Available hydro-physical measurements

We conducted hydro-physical measurements across a 55 km continuous segment of the Karnali River of Nepal to the border with India during low water season when available habitat is reduced and river dolphin-anthropogenic activity interactions are intensified [32]. As the risk of changes in physical habitat may not truly represent the change in flow during the high flood season (or during daily fluctuation), we measured hydro-physical parameters two months from the termination of the monsoon (September). Field measurements of hydro-physical variables were conducted in December of 2017 and March and May of 2018 representing different flows (medium, low and high respectively) within the low water season. Therefore, our hydro-physical measurements represent a stable channel environment in response to flow. Flow was classified into different categories based on 39 years (1977-2015) of discharge data available for the study site from the Department of Hydrology, Government of Nepal. We stratified the study site into three segments (upper, middle, and lower, Fig 1) based on anthropogenic activities and hydro-physical characteristics of the river segment. The Mega hydropower project (Upper Karnali) is located upstream of the study site, whereas the Rani Jamuna Irrigation project water extraction point is in the upper segment of the study site. Within each segment, the study reach (the linear segment where cross-sections are established) was established in such a way that the length of each reach was at least higher than the mean width (so the number varies among segments) of the respective segment. Further, we also tried to maintain a relatively similar flow at the top and

bottom of the reach to create hydro-physically homogenous reaches. Within each reach, random cross-sections were established to capture the hydraulic properties based on flow variation and length of the reach.

We used SonTek RiverSurveyor S5 systems to measure open channel hydro-physical parameters (total discharge- m^3/s ; velocity- m/s ; depth- m , wetted perimeter- m ; total width- m , cross-sectional area (CSA)- m^2) at each cross section, which used multiple acoustic frequencies along with precise bandwidth control for the most robust and continuous shallow-to-deep discharge measurements ever. The details of the operation of S5 are given in [10]. An intelligent algorithm with high ping rates of S5 looks at water depth, velocity, and turbulence. It then acoustically adapts to those conditions to ensure extremely robust data collection by compensating for vessel motion due to surface conditions. All discharge computations are collected within the S5. Total discharge at each cross-section, along with its attributes (velocity and depth) were measured at a cell size between 0.02 to 0.5 m, with potential measurement errors of 0.25 % and 1 % in velocity and depth, respectively. The ADP S5 hydraulic data were imported into Excel databases (Microsoft v. 2010) to format for System for Environmental Flow Analysis (SEFA, version 1.5; Aquatic Habitat Analysts Inc.) software, where we estimated AWS for each cross-section and reach using HSC (described in a later section). The habitat type (HT), i.e., pool, run, and riffle, was classified based on the Froude number (Fr), where Froude is an index of hydraulic turbulence (the ratio of velocity by the acceleration of gravity). Points with Froude numbers exceeding 0.41 were considered riffles, points with Froude numbers less than 0.18 were considered pools, and intermediate values were classified as run habitats.

We used basic descriptive statistics (mean, range, standard deviation (SD)) to characterize the hydro-physical parameters. Variation of these parameters across the season and habitat type was examined using F-test at a 5% level of significance. All these analyses were done in R-Studio version 1.3.95.

GRD occupied habitat assessment

We measured the hydro-physical characteristics of the space used by GRD only in the main Karnali River, which is occupied by GRD most of the year. The tributary, Mohana segment, excluded from the study as this tributary is only used by GRD during the monsoon season to avoid excessively high flow rates in the main stream. Also, across most of the year, this tributary

exhibited seasonal characteristic (almost dry in the dry season). During each survey period, we used SonTek RiverSurveyor S5 for the measurement of hydro-physical parameters of GRD occupied. We used the occupied habitat's hydro-physical parameters to derive the Habitat Suitability Curve (HSC); the procedure of HSC development is described in [10]. We mainly derived a suitable curve as a function of depth and velocity, which allows quantification of the quality of aquatic habitat in the form of area-weighted suitability (AWS, m^2/m) at wider flow levels. This research was conducted under a research permit issued to the principal author of this paper by the Department of National Parks and Wildlife Conservation (DNPWC), Government of Nepal. All the observation procedures comply with regulations developed by the DNPWC.

Area weighted suitability model

Because of complex relationships between response (AWS) and explanatory variables, the risk of violation of statistical assumptions (risk of non-homogeneity of the variance of a response variable, skewness in the distribution of response) is very high. To address this, we used generalized additive models for location, scale, and shape (GAMLSS) to build relationships between response (AWS) and explanatory variables (flow, velocity, depth, season (S), habitat type (HT), width, wetted perimeter, cross-sectional area (CSA)) which permits us to address those statistical issues within the model [33]. First, we selected the distribution of the response variable that can adequately describe the nature of the response variable by building linear models of explanatory variables using available potential distribution type (gamma, Box-Cox Cole-Green orig., normal, inverse gamma, lognormal) within the GAMLSS package. The model with proper distribution type was selected using generalized Akaike information criterion (GAIC). We found gamma distribution to be the best distribution family that can describe our response variable adequately. We used the gamma distribution for all our subsequent analysis.

We select significant model terms – linear, smoothing, and interactive – separately to develop the initial model that consists of all potential linear, smoothing, and interactive terms. A significant linear term was selected by developing a full linear model of seven explanatory variables. Then we used the *dropterm* function to get the final significant linear terms, which adopts stepwise

model selection steps based on GAIC. To obtain the significant smoothing term, the smoothing parameters (except categorical variables) were selected by developing the initial null model, and then each smoother was added, one at a time, to the null model to get significant smoothing terms. Further, we determine whether two-way interactions are needed in the model using *stepGAIC* function in between intercept (lower model) and the most complicated model with all two-way interactions (upper). After having a full initial model with significant terms (liner, interactive, and smoothers), we again used the *dropterm* function to get the final model with the lowest GAIC.

We use *drop1* function available in GAMLSS package to check for the approximate significance of the contribution of the smoothers (including the linear and interactive terms) at a 5% level of significance. To check the adequacy of the fitted GAMLSS model, we used a worm plot (*wp* function), which is a de-trended QQ-plot of the residuals [34]. For an adequate fitted model, we would expect the dots to be close to the middle horizontal line and 95 % of them to lie between the upper and lower dotted curves, which act as 95 % pointwise confidence intervals, with no systematic departure. All modeling procedures were completed using *gamlss* package in R-Studio version 1.3.95.

Results

Hydro-physical parameters

Hydro-physical parameters differ over season, except in depth (Table 1). All parameters were high in May and lowest in March. Across space or habitat type, hydro-physical parameters differ, except in flow (Table 2). Higher depth and cross-sectional area, along with lowest velocity were recorded in deep pool habitat, followed by run and riffle habitats.

Table 1. Hydro-physical parameters across seasons. Variation on parameters across season tested using F-test and P value of significance reported.

	December (N=70)	March (N=60)	May(N=47)	Total (N=177)	P.value
Flow(m³/s)					$F_{2,174}=22.05$, $P<0.001$
Mean (SD)	245.94 (81.52)	203.14(64.56)	326.33 (139.48)	252.78 (106.73)	
Range	82.96–420.34	81.52–475.61	50.60–665.49	50.60–665.49	
Depth(m)					$F_{2,174}=1.91$, $P=0.15$
Mean (SD)	1.95 (0.79)	1.92 (0.74)	1.70 (0.47)	1.87 (0.71)	
Range	0.93–4.71	0.88–5.02	1.05–3.12	0.88–5.02	
Velocity(m/s)					$F_{2,174}=4.89$, $P=0.009$
Mean (SD)	0.96 (0.33)	0.83 (0.29)	1.03 (0.35)	0.94 (0.33)	
Range	0.454–2.120	0.335–2.111	0.374–1.822	0.335–2.120	
Width(m)					$F_{2,174}=10.23$, $P<0.001$
Mean (SD)	128.17 (46.54)	126.18 (56.18)	180 (102.37)	141.26 (72.06)	
Range	47.50–271.62	56.56–390.50	55.01–490.15	47.50–490.15	
CSA(m²)					$F_{2,174}=3.78$, $P=0.025$
Mean (SD)	252.56 (136.35)	233.31 (102.97)	306.67 (181.26)	260.40 (142.40)	
Range	68.45–760.94	74.79–593.67	80.72–774.76	68.45–774.76	
WP(m)					$F_{2,174}=10.14$, <0.001
Mean (SD)	129.29 (46.52)	127.23 (55.96)	180.87 (102.63)	142.29 (72.06)	
Range	48.25–272.68	57.61–391.19	57.13–491.93	48.25–491.93	

262

263

264

265

266

267

268

269

Table 2. Hydro-physical parameters across habitat type. Variation on parameters across season tested using F-test and P value of significance reported.

	Pool(N=65)	Riffle(N=20)	Run(N=92)	Total(N=177)	P.value (F-test)
Flow(m³/s)					F _{2,174} =0.36, 0.695
Mean (SD)	261.78 (94.97)	245.99 (97.43)	247.90 (116.68)	252.78 (106.73)	
Range	50.60–481.55	121.34– 475.61	95.68– 665.49	50.60–665.49	
Depth(m)					F _{2,174} =58.16, P< 0.001
Mean (SD)	2.46 (0.80)	1.39 (0.24)	1.57 (0.35)	1.87 (0.71)	
Range	1.17–5.02	0.93–1.92	0.88–2.87	0.88–5.02	
Velocity(m/s)					F _{2,174} =203, P< 0.001
Mean (SD)	0.66 (0.14)	1.59 (0.28)	0.99 (0.18)	0.94 (0.33)	
Range	0.33–1	1.12–2.12	0.53–1.41	0.33–2.12	
Width(m)					F _{2,174} =3.59, P=0.03
Mean (SD)	147.23 (64.87)	101.22 (34.05)	145.74 (80.27)	141.26 (72.06)	
Range	55.01–441.05	47.50–171.49	56.56–490.15	47.50–490.15	
CSA(m²)					F _{2,174} =28.55, P< 0.001
Mean (SD)	346.27 (139.31)	139.07 (45.61)	226.11(124.27)	260.40 (142.40)	
Range	80.72–760.94	68.45–246.03	74.79–774.76	68.45–774.76	
WP(m)					F _{2,174} =3.61, P=0.029
Mean (SD)	148.73 (64.65)	102.16(34.29)	146.46 (80.35)	142.29 (72.06)	
Range	57.13–442.75	48.25–172.08	57.61–491.93	48.25–491.93	

272

273 *GAMLSS model of area-weighted suitability*

274 The model with the lowest GAIC value consists of the linear (flow, velocity, habitat type,
275 season), smoother (depth, cross-sectional area) and interactive terms (depth and velocity,
276 velocity and season, habitat type, and season) [Model 1, Table 3]. As indicated by the summary
277 statistics of the quantile residuals of the best-fitted model (the mean of the model is nearly zero,
278 variance almost one, coefficient of skewness is almost zero, and coefficient of kurtosis is almost
279 3), the residuals are approximately normally distributed, as for an adequate model [33].
280 Furthermore, the Filliben correlation coefficient (or the normal probability plot correlation
281 coefficient) is almost 1. 95% of the points lie between the two elliptic curves of the Worm plot

(Fig 2). All these statistical and visual graphs suggest that the fitted distribution (or the fitted terms) of the model is adequate to explain the response variable.

Table 3. GAMLSS models with df (degree of freedom) and GAIC (generalized Akaike Information Criteria) for the response variable area-weighted suitability (AWS, m²/m) and explanatory variables – flow (m³/s), velocity (m/s), depth(m), habitat type (HT- deep pool, riffle, run), cross-sectional area (m²) and season (December, March and May). The best fitted model is highlighted in bold, which consist of linear, smoother (as indicated by pb), and interactive terms of the explanatory variables.

Model ID	Model	df	GAIC
1	AWS ~ Flow + Velocity + HT + Season + pb(Depth)+pb(CSA) + Depth:Velocity + Velocity:Season + HT:Season	27.73	1258.52
2	AWS ~ Flow + Depth + Velocity + CSA + HT + Season + pb(Depth) +pb(CSA) + Depth:Velocity + Flow:Season + Velocity:Season + HT:Season	29.66	1262.49
3	AWS ~ Flow + Depth + Velocity + CSA + HT + Season + pb(Depth) + Depth:Velocity + Flow:Season + Velocity:Season + HT:Season	27.06	1276.33
4	AWS ~ Flow + Depth + Velocity + CSA + HT + Season + Depth:Velocity +Velocity:Season + Depth:HT + Flow:CSA + HT:Season	25	1286.28
5	AWS ~ Flow + pb(Depth) + pb(Velocity) + CSA + HT + Season + Depth:Velocity + Flow:Season + Velocity:Season	22.97	1290.27
6	AWS ~ Flow + pb(Depth) + Velocity + CSA + HT + Season+Depth:Velocity + Flow:Season	20.7	1292.98
7	AWS ~ Flow + Depth + Velocity + CSA + HT + Season + Depth:Velocity + Velocity:Season + Depth:HT + Flow:CSA	21	1300
8	AWS ~ Flow + Depth + Velocity + CSA + HT + Season + Depth:Velocity +Flow:Season + Velocity:Season + Depth:HT + Flow:CSA	23	1303.54
9	AWS ~ Flow + Depth + Velocity + CSA + HT + Season + Depth:Velocity +Flow:Season + Velocity:Season + Depth:HT	22	1309.94
10	AWS ~ Flow + Depth + Velocity + CSA + HT + Season + Depth:Velocity + Flow:Season	18	1313.73
11	AWS~Flow + Depth + Velocity + CSA + HT + Season + Depth:Velocity +Flow:Season + Velocity:Season	20	1315.13
12	AWS ~ Flow + Depth + Velocity + CSA + HT + Season + Depth:Velocity	16	1324.36
13	AWS ~ Velocity + CSA + HT + Season	13	1339.85
14	AWS~Flow+Depth+Velocity+CSA+HT+Season	15	1343.82

Fig 2. A worm plot depicts that points of the worm plot are inside the 95 % pointwise confidence intervals (curves) and close to the middle horizontal line, indicating adequacy of the fitted model.

Highly significant (the level of significance $P < 0.001$) linear predictors of the AWS were flow, velocity, season (March), depth, cross-sectional area, and of interactive terms were velocity and depth, and velocity and season (March). Flow and velocity exhibited linear negative relationships with AWS. In contrast, the depth and cross-sectional area revealed non-linear relationships, indicating both positive and negative contributions to AWS (Fig 3). Highest AWS was recorded in both December and March, and pools and riffle habitats exhibited higher AWS. The highest contribution to the variation of AWS was the interactions of the velocity and season (March) [$\beta=32.15$, 95 % CI range: 5.89- 175.43], velocity and depth ($\beta: 4.52$, 95 % CI: 2.26-9.03), followed by linear terms – cross-sectional area ($\beta=1.01$, 95 % CI: 1-1.01), flow ($\beta=0.992$, 95 % CI: 0.990-0.994), depth ($\beta=0.36$, 95 % CI: 0.21-0.60) and velocity ($\beta=0.03$, 95 % CI: 0.009-0.159) respectively. The cross-sectional area $< 250 \text{ m}^2$ showed a negative contribution to the AWS, and above this point showed a positive linear contribution to the AWS (Fig 3). By season, peak dry season contributed negatively to the AWS. In contrast, both December and May had a positive contribution to the AWS in which the contribution of May was higher than December. Contribution of the deep pool and riffle habitats were higher to AWS than run (Fig 3). We model the contribution of wider flows ($50\text{-}665 \text{ m}^3/\text{s}$) to AWS, in which flows between 300 and $550 \text{ m}^3/\text{s}$ showed maximum positive contribution to AWS with the peak value of AWS at $450 \text{ m}^3/\text{s}$ (Fig 4). Relationships between flow and AWS reveal that a small and large value of flow has the same value of AWS. Velocity $> 1\text{m/s}$ showed a negative contribution to the AWS in which 0.6 m/s had the highest positive contribution to AWS. Depth of $2\text{--}4 \text{ m}$ maximized contribution to AWS, in which 2.5 m had the highest contribution to AWS (Fig 5).

Fig 3. A plot of the final fitted model shows that the predictor (AWS) declines linearly as flow and velocity increase, and AWS exhibited nonlinear relationships with depth and cross-sectional area. The contribution of the season to AWS was higher for both December and May, whereas March contributed negatively to AWS. Deep pool and riffle contributed positively, but run habitat holds a negative contribution to AWS.

Fig 4. Centile curves for AWS (m^2/m) against the flow (m^3/s) for different percentiles (5-95 %) shows that maximum AWS is obtained with the flow around $450 \text{ m}^3/\text{s}$. Vertical line represents an average of 90 % exceedance flow of low (March), medium (December), and high (May) flow periods, suggesting highest AWS was available during May and AWS was substantially reduced during March and December due to reduced flow. It indicates that the flow of March (peak dry season) needs to be protected as it is under critical flow to protect river dolphins, which contributes negatively to AWS.

Fig 5. A plot of centile curves for AWS (m^2/m) fitted against depth and velocity separately indicates AWS is maximized when depth is between 2 and 3.5 m and velocity 0.3–0.8 m/s. 95 centile value of AWS $>150 \text{ m}^2/\text{m}$ when depth and velocity were 2.7 m and 0.6 m/s, respectively.

Interactive effects of velocity and depth on area-weighted suitability

The velocity $<0.25 \text{ m/s}$ and depth with 2.5 m yielded the highest AWS (Fig 6). Depth between 2 and 3 m and higher cross-sectional area ($>600 \text{ m}^2$) predicted the highest AWS. Velocity between 0.5 and 1 m/s with a higher cross-sectional area ($>600 \text{ m}^2$) yielded the highest AWS.

Fig 6. A surface plot of the AWS (m^2/m) showing the interactive effects of (a) velocity and depth, (b) cross-sectional area and velocity, (c) cross-sectional area, and depth. The plot suggests that depth between 2 and 3 m combined with velocity $<0.5 \text{ m/s}$ maximized the AWS. Similarly, velocity and depth with a similar range along with a cross-sectional area $>600 \text{ m}^2$ optimized the AWS.

Discussion

~~Understanding the fine-scale habitat selection of large and mobile aquatic predators –~~

Lack of information on the fine-scale habitat selection of large and mobile aquatic predators – river cetaceans – limits our ability to understand the broad scale impacts of human-caused changes to the dynamic riverine ecosystem. As river cetaceans exhibit strong dependence on hydro-physical attributes, including geomorphic and seasonal effects, they are essential to informing the decision-making process of flow regulation management, including the identification and protection of river cetaceans' priority habitats. Our results can be used to incorporate knowledge about in-stream habitat selection and the potential effects of declining

freshwater flow on the ecology of the large top predator of the riverine ecosystem into the design of habitat retention plan and water development projects that may alter the natural flow regime at the cost of aquatic species conservation. Incorporation of the ecology of a key mobile aquatic predator may help to conserve other lower trophic or similar aquatic species that share the river's physical habitat template and ecosystem [35].

Previous studies on riverine cetaceans' habitat selection or preference commonly reported the aggregation of riverine cetaceans in certain hydro-physical habitats [32, 36], especially deep pools. The majority of the studies attributed such habitat selection to the higher biological productivity and prey abundance [37, 38]. However, the ecological significance of such habitats to river dolphins has never been described explicitly. Here, we characterize the hydro-physical parameters of the deep pools that may be significant to the ecology of riverine cetaceans. We demonstrate that deep pools offer higher cross-sectional areas combined with the deepest depth and considerably reduced water velocity, and in turn, yield the highest usable area to river dolphins. The ecological significance of deep pools that offer the highest usable area is linked to the diving physiology of dolphins as the selection of deeper usable areas appears to be a mechanism that enables diving river cetaceans with limited oxygen stores to extend the duration of a dive [39]. Further, reduced velocity of water in deep pools in comparison to other habitats (run, riffle) may support river dolphins to reduce their swimming speed as the elevated swimming speeds during dive or swimming decreases the duration of a dive due to rapid depletion of limited oxygen reserves [40]. As a result, frequent diving in deep pools by riverine cetaceans is ecologically and physiologically important by reducing the metabolic cost of dives. Such energetically efficient modes of locomotion in deep pools provide an advantage during periods of diving (submerge) and will presumably increase foraging efficiency as the animals perform progressively longer dives.

Global population decline and isolation risks of riverine cetaceans is frequently attributed to declining flows due to dams or barrages [16, 32, 18, 41, 42, 11]. As a result, the need for adequate flow to conserve the remaining endangered riverine cetaceans is heightened globally [19, 20, 43, 42, 10]. However, to our knowledge, information on quantitative relationships between flow and river cetaceans is scant, particularly for the low water season when habitat availability reduces, and longitudinal connectivity worsens. The linear negative relationships

between flow and AWS combined with physical effects of flow variability in the cross-sectional area of the channel exemplified that immediate effects of flow fluctuation (either increase or decrease) are the loss of suitably usable area (i.e. habitat loss). All flows are not contributing in the same fashion to AWS, offering the opportunity to balance water extraction and conservation. Availability of AWS as a function of river flow shows that the flow of May (average 90 % exceedance flow= 439.4 m³/s) offers maximum AWS compared to the flows of December (average 90 % exceedance flow=335 m³/s) and March (average 90 % exceedance flow=221 m³/s) [Fig 4]. Flows in March represent critically low flow (negative contribution to AWS), which substantially reduces AWS. Flow from December to April seems more critical for river dolphin conservation. This flow variation across season suggests that flow is the major determinant of river cetaceans' hydro-physical habitat, which substantially alters the habitat at varying spatial and temporal scales. Further, the variation in flow is likely to control the migration strategies of aquatic species [44]. This is demonstrated by GRD in our study site. As the monsoon starts with average 90% exceedance flow of 672 m³/s in June, unusable habitats in the main stream with flow >600 m³/s (Fig 4) trigger GRD to migrate to a tributary (Mohana River; starts from the first week of June; personal observation), which offers seasonal refuge and suggests that a seasonal high flow has a major influence on shaping the life-history patterns of riverine cetaceans. Similar high flow-regulated seasonal migration patterns are exhibited by river dolphins in an Amazonian floodplain [45]. As the riverine cetaceans' birthing peak is during low water season [46], such high flow with the potential of physical disturbance immediately after the birth may pose a risk of recruitment failure for young calves [44].

The requirement of adequate flow is commonly cited to protect riverine cetaceans and avoid the risk of population isolations that may cause recruitment failure and intensify local extinction [19, 20, 11, 10]. Our results may assist to determine the adequate flow across seasons that might have potential ecological significance to maintain the longitudinal connectivity essential to sustain the viability of river dolphins and other aquatic taxa that share the ecosystem. As water extraction and dam construction limits aquatic species dispersal ability to move freely through the stream network, extinction risk and endangerment of river cetaceans whose existence relies on flow will particularly be high during low water season. Our results could help river managers to predict how far in-stream usable areas can be altered from their natural state before water extraction or withdrawal occurs. As the species' abundance and distribution vary with stream depth and

velocity [47], the microhabitat preferences of river cetaceans as a function of velocity and depth were poorly known. Our model shows that the distribution of suitable habitats (i.e. pools) of GRD is largely determined by the interactive effects of velocity and depth at a particular landform. The velocity (<0.5 m/s) combined with a depth between 2 and 3 m yielded a higher useable area to GRD in the study site (Fig 6). Further, a greater cross-sectional area (>600 m²) with a similar velocity and depth ranges offer maximum suitable areas to GRD. Such complex habitat selection behavior (depth, velocity, and cross-sectional area) exhibited by Ganges River dolphins in a dynamic aquatic environment indicates GRD are habitat specialists with narrow habitat breadth and environmental plasticity. Given the often-cited patchy distribution pattern of GRD [49], such interactive effects may play a significant role in the distribution and abundance of river cetaceans. Understanding riverine cetaceans' specific in-stream habitat preference in relation to flow may offer an effective riverine conservation management plan for a heavily regulated river system [44].

Loss in habitat quality and the limited ability of species to migrate to new habitats is a major challenge for aquatic species conservation [50]. Our results indicate that vulnerability of river cetaceans is increased by fragmentation and loss of longitudinal connectivity when flow reduction is intensified either naturally or artificially. Higher risk of deep pool fragmentation and loss of longitudinal connectivity across suitable pools are commonly suggested conservation issues for riverine cetaceans conservation, particularly in the dry season [17, 11, 10]. Effective flow regulation plans are critical to riverine cetacean conservation and management to avoid further acceleration of extinction risk [19, 20, 11, 10]. A pseudo-2D map of the segments (surface map of depth as a function of flow) for the low flow period (March) clearly indicates the substantial fragmentation in deep pools' (>2) availability along with potential dolphin passage barriers (depth <0.5 m) as a response to reduce flow (Fig 7). Barriers to passage may limit GRD dispersal ability to suitable habitats, which may exacerbate the rate of endangerment and local extinction of the remaining populations of endangered GRD [51]. We suggest that conservation agencies must be adopted effective assessment procedures to determine the true impact of flow extraction or diversion projects to intensify conservation of aquatic biota in these vanishing riverine ecosystems.

Fig 7. A pseudo-2D view of the depth (m) surface showing in contours and shading for segments (a) upper, (b) medium, (c) lower using interpolation technique. Light yellow indices depth <0.5 m and dark blue indicate deep pools >2 m, respectively. Sparsely located dark blue and substantially present light yellow over the continuum of the segments indicate acute habitat fragmentation and loss of connectivity across suitable pools during low water (peak dry season-March) when flow reduced.

The biological consequences of habitat loss and fragmentation in declining river cetaceans are evident. Births and mating activities of riverine cetaceans generally peak during low water season when all animals are concentrated along the main stream [46], likely reducing pairing success, which may hinder reproduction processes in isolated habitats and be a contributing factor to the low occupancy and decline of species, especially *Platanista gangetica* species. Further, spatial overlap during the low water season between river cetaceans and fisheries may intensify, leading to entanglement and reduction of survivorship [15, 14, 10]. Further diversion or withdrawal of flow from the mainstream contributes to habitat loss and undoubtedly leads to local extirpation. Combined effects of habitat loss with anthropogenic disturbances (fishing pressure and flow regulation by development projects) may lead to rapid population decline and local extinction of river cetaceans [51]. Rapid loss (~50 %) of *Platanista gangetica* species is the clear evidence of such effects [17, 10]. Even though conserving the natural flow is unrealistic given societal demands [52], the seasonal effects and changes on the availability of usable areas to river cetaceans must be a central guiding element to conserve riverine biodiversity. Identifying priority riverscapes for river cetaceans and prioritizing investment opportunities could be an essential first step towards effective conservation of riverine biodiversity [53].

Our research addresses a critical limitation in the ability to predict and quantify the riverine cetaceans' response to flow regulation. We demonstrated quantitative relationships between flow and cetaceans ecology in the form of usable area availability in response to flow. These findings should permit development of a proactive riverine cetacean management plan that incorporates habitat manipulation and protection, including the creation of migration/movement corridors. Owing to the strong dependence on the attribute of freshwater flow, riverine cetaceans are expected to be more affected by flow regulation as the interactive effects of flow and depth at particular landforms play a significant role to determine the usable area. Freshwater withdrawal and extraction by large-scale hydropower or irrigation projects will undoubtedly cause major ecological changes in the dynamic riverine ecosystem, which decreases viability of riverine

cetaceans populations. River cetaceans are more sensitive to the attributes of freshwater flow, and the decline of freshwater flow may further put endangered cetaceans under risk of local extirpation, as exemplified by a sharp decline in abundance and range of GRD in a study site as flow regulation progress [32]. Knowledge of key hydro-physical habitat components may support development of a riverine habitat retention management plan in response to top predator or mega-species life history requirements [10].

Acknowledgments

Thanks to the numerous agencies -- Rufford Foundation (UK), WWF-EFN Program (USA), Institute of Forestry (Tribhuvan University, Nepal), Commonwealth Scientific and Industrial Research Organization (CSIRO-Australia), WWF-Nepal, University of Arizona – School of Natural Resources and the Environment – for field funding support. We also thank Rajesh Sigdel, RC Gautam and various local stakeholders for assisting data collection process.

508

References

- 509 [1] Dudgeon D. Multiple threats imperil freshwater biodiversity in the Anthropocene. *Current*
510 *Biology*. 2019 Oct 7;29(19):R960-7.
- 511 [2] Destouni G, Jaramillo F, Prieto C. Hydroclimatic shifts driven by human water use for food
512 and energy production. *Nature Climate Change*. 2013 Mar;3(3):213-7.
- 513 [3] Jaramillo F, Destouni G. Local flow regulation and irrigation raise global human water
514 consumption and footprint. *Science*. 2015 Dec 4;350(6265):1248-51.
- 515 [4] Destouni G, Jarsjö J. Zones of untreatable water pollution call for better appreciation of
516 mitigation limits and opportunities. *Wiley Interdisciplinary Reviews: Water*. 2018
517 Nov;5(6):e1312.
- 518 [5] Winemiller KO, McIntyre PB, Castello L, Fluet-Chouinard E, Giarrizzo T, Nam S, Baird IG,
519 Darwall W, Lujan NK, Harrison I, Stiassny ML. Balancing hydropower and biodiversity
520 in the Amazon, Congo, and Mekong. *Science*. 2016 Jan 8;351(6269):128-9.
- 521 [6] FAO. 2016. AQUASTAT database. <http://www.fao.org/nr/water/aquastat/data/>.
- 522 [7] Elmhagen B, Destouni G, Angerbjörn A, Borgström S, Boyd E, Cousins SA, Dalén L, Ehrlén
523 J, Ermold M, Hambäck PA, Hedlund J. Interacting effects of change in climate, human
524 population, land use, and water use on biodiversity and ecosystem services. *Ecology and*
525 *Society*. 2015 Mar 1;20(1).
- 526 [8] Carrizo SF, Jähnig SC, Bremerich V, Freyhof J, Harrison I, He F, Langhans SD, Tockner K,
527 Zarfl C, Darwall W. Freshwater megafauna: Flagships for freshwater biodiversity under
528 threat. *Bioscience*. 2017 Oct 1;67(10):919-27.
- 529 [9] He F, Zarfl C, Bremerich V, David JN, Hogan Z, Kalinkat G, Tockner K, Jähnig SC. The
530 global decline of freshwater megafauna. *Global Change Biology*. 2019 Nov;25(11):3883-
531 92.
- 532 [10] Paudel S, Koprowski JL. Factors affecting the persistence of endangered Ganges River
533 dolphins (*Platanista gangetica gangetica*). *Ecology and Evolution*. 2020 Mar;10(6):3138-
534 48.
- 535 [11] Paschoalini M, Almeida RM, Trujillo F, Melo-Santos G, Marmontel M, Pavanato HJ,
536 Guerra FM, Ristau N, Zerbini AN. On the brink of isolation: Population estimates of the
537 Araguaian river dolphin in a human-impacted region in Brazil. *Plos one*. 2020 Apr
538 22;15(4):e0231224.
- 539 [12] Braulik GT, Noureen U, Arshad M, Reeves RR. Review of status, threats, and conservation
540 management options for the endangered Indus River blind dolphin. *Biological*
541 *Conservation*. 2015 Dec 1;192:30-41.
- 542 [13] Krützen M, Beasley I, Ackermann CY, Lieckfeldt D, Ludwig A, Ryan GE, Bejder L, Parra
543 GJ, Wolfensberger R, Spencer PB. Demographic collapse and low genetic diversity of the
544 Irrawaddy dolphin population inhabiting the Mekong River. *Plos one*. 2018 Jan
545 3;13(1):e0189200.

- [14] Kelkar N, Krishnaswamy J, Choudhary S, Sutaria D. Coexistence of fisheries with river dolphin conservation. *Conservation Biology*. 2010 Aug;24(4):1130-40.
- [15] Martin AR, da Silva VM, Salmon DL. Riverine habitat preferences of botos (*Inia geoffrensis*) and tucuxis (*Sotalia fluviatilis*) in the central Amazon. *Marine Mammal Science*. 2004 Apr;20(2):189-200.
- [16] Turvey ST, Pitman RL, Taylor BL, Barlow J, Akamatsu T, Barrett LA, Zhao X, Reeves RR, Stewart BS, Wang K, Wei Z. First human-caused extinction of a cetacean species?. *Biology letters*. 2007 Oct 22;3(5):537-40.
- [17] Braulik GT, Noureen U, Arshad M, Reeves RR. Review of status, threats, and conservation management options for the endangered Indus River blind dolphin. *Biological Conservation*. 2015 Dec 1;192:30-41.
- [18] Pavanato HJ, Melo-Santos G, Lima DS, Portocarrero-Aya M, Paschoalini M, Mosquera F, Trujillo F, Meneses R, Marmontel M, Maretti C. Risks of dam construction for South American river dolphins: a case study of the Tapajós River. *Endangered species research*. 2016 Sep 21;31:47-60.
- [19] Smith BD, Braulik G, Strindberg S, Mansur R, Diyan MA, Ahmed B. Habitat selection of freshwater-dependent cetaceans and the potential effects of declining freshwater flows and sea-level rise in waterways of the Sundarbans mangrove forest, Bangladesh. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 2009 Mar;19(2):209-25.
- [20] Braulik GT, Reichert AP, Ehsan T, Khan S, Northridge SP, Alexander JS, Garstang R. Habitat use by a freshwater dolphin in the low-water season. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 2012 Jun;22(4):533-46.
- [21] Maddock I. The importance of physical habitat assessment for evaluating river health. *Freshwater biology*. 1999 Mar;41(2):373-91.
- [22] Aldaya MM, Martínez-Santos P, Llamas MR. Incorporating the water footprint and virtual water into policy: Reflections from the Mancha Occidental Region, Spain. *Water Resources Management*. 2010 Mar 1;24(5):941-58.
- [23] Rosenfeld JS. Developing flow–ecology relationships: Implications of nonlinear biological responses for water management. *Freshwater Biology*. 2017 Aug;62(8):1305-24.
- [24] Bruckerhoff LA, Leasure DR, Magoulick DD. Flow–ecology relationships are spatially structured and differ among flow regimes. *Journal of applied ecology*. 2019 Feb;56(2):398-412.
- [25] Choudhary S, Dey S, Dey S, Sagar V, Nair T, Kelkar N. River dolphin distribution in regulated river systems: implications for dry-season flow regimes in the Gangetic basin. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 2012 Jan;22(1):11-25.
- [26] Peter C, Poh AN, Ngeian J, Tuen AA, Minton G. Identifying habitat characteristics and critical areas for Irrawaddy dolphin, *Orcaella brevirostris*: Implications for conservation. In *Naturalists, explorers and field scientists in South-East Asia and Australasia 2016* (pp. 225-238). Springer, Cham.

- 585 [27] Araújo CC, da Silva VM. Spatial distribution of river dolphins, *Inia geoffrensis* (Iniidae), in
586 the Araguaia River (central Brazil). *Mammalia*. 2014 Nov 1;78(4):481-6.
- 587 [28] Guizada L, Aliaga-Rossel E. Abundance of the Bolivian River Dolphin (*Inia boliviensis*) in
588 Mamore River, Upper Madeira Basin. *Aquatic Mammals*. 2016 Jul 1;42(3)
- 589 [29] Chen M, Yu D, Lian Y, Liu Z. Population abundance and habitat preference of the Yangtze
590 finless porpoise in the highest density section of the Yangtze River. *Aquatic Conservation:*
591 *Marine and Freshwater Ecosystems*. 2020 Jun;30(6):1088-97.
- 592 [30] Thomas JA, Bovee KD. Application and testing of a procedure to evaluate transferability of
593 habitat suitability criteria. *Regulated Rivers: Research & Management*. 1993 Aug;8(3):285-
594 94.
- 595 [31] Rani Jamara Kulariya Irrigation Project, 2017. *Environmental Assessment*. Volume 2. Nepal:
596 Department of Irrigation.
- 597 [32] Paudel S, Pal P, Cove MV, Jnawali SR, Abel G, Koprowski JL, Ranabhat R. The
598 Endangered Ganges River dolphin *Platanista gangetica gangetica* in Nepal: abundance,
599 habitat and conservation threats. *Endangered Species Research*. 2015 Nov 25;29(1):59-68.
- 600 [33] Stasinopoulos MD, Rigby RA, Heller GZ, Voudouris V, De Bastiani F. Flexible regression
601 and smoothing: using GAMLSS in R. CRC Press; 2017 Apr 21.
- 602 [34] Buuren SV, Fredriks M. Worm plot: a simple diagnostic device for modelling growth
603 reference curves. *Statistics in medicine*. 2001 Apr 30;20(8):1259-77.
- 604 [35] Turvey ST, Risley CL, Barrett LA, Yujiang H, Ding W. River dolphins can act as
605 population trend indicators in degraded freshwater systems. *PLoS One*. 2012 May
606 29;7(5):e37902.
- 607 [36] Gomez-Salazar C, Trujillo F, Portocarrero-Aya M, Whitehead H. Population, density
608 estimates, and conservation of river dolphins (*Inia* and *Sotalia*) in the Amazon and
609 Orinoco river basins. *Marine Mammal Science*. 2012 Jan;28(1):124-53.
- 610 [37] Smith BD, Haque AA, Hossain MS, Khan A. River dolphins in Bangladesh: conservation
611 and the effects of water development. *Environmental management*. 1998 May
612 1;22(3):323-35.
- 613 [38] Biswas SP, Boruah S. Ecology of the river dolphin (*Platanista gangetica*) in the upper
614 Brahmaputra. *Hydrobiologia*. 2000 Jul 1;430(1-3):97-111.
- 615 [39] Skrovan RC, Williams TM, Berry PS, Moore PW, Davis RW. The diving physiology of
616 bottlenose dolphins (*Tursiops truncatus*). II. Biomechanics and changes in buoyancy at
617 depth. *Journal of Experimental Biology*. 1999 Oct 15;202(20):2749-61.
- 618 [40] Williams TM, Haun JE, Friedl WA. The diving physiology of bottlenose dolphins (*Tursiops*
619 *truncatus*). I. Balancing the demands of exercise for energy conservation at depth. *Journal*
620 *of Experimental Biology*. 1999 Oct 15;202(20):2739-48.

- [41] Choudhury NB, Mazumder MK, Chakravarty H, Choudhury AS, Boro F, Choudhury IB. The endangered Ganges river dolphin heads towards local extinction in the Barak river system of Assam, India: A plea for conservation. *Mammalian Biology*. 2019 Jan 1;95(1):102-11.
- [42] Trujillo-González F, Mosquera-Guerra F, Franco N. River dolphins: Species that indicate the state of health of the aquatic ecosystems of the Amazon and Orinoco regions. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales*. 2019 Jun;43(167):199-211.
- [43] Dey, M., Krishnaswamy, J., Morisaka, T., & Kelkar, N. (2019). Interacting effects of vessel noise and shallow river depth elevate metabolic stress in Ganges river dolphins. *Scientific reports*, 9(1), 1-13.
- [44] Bunn SE, Arthington AH. Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environmental management*. 2002 Oct 1;30(4):492-507.
- [45] Mintzer VJ, Lorenzen K, Frazer TK, da Silva VM, Martin AR. Seasonal movements of river dolphins (*Inia geoffrensis*) in a protected Amazonian floodplain. *Marine Mammal Science*. 2016 Apr;32(2):664-81.
- [46] Martin AR, Da Silva VM. Reproductive parameters of the Amazon river dolphin or boto, *Inia geoffrensis* (Cetacea: Iniidae); an evolutionary outlier bucks no trends. *Biological Journal of the Linnean Society*. 2018 Mar 2;123(3):666-76.
- [47] Jowett IG, Richardson J, McDowall RM. Relative effects of in-stream habitat and land use on fish distribution and abundance in tributaries of the Grey River, New Zealand. *New Zealand journal of marine and freshwater research*. 1996 Dec 1;30(4):463-75.
- [49] Smith BD. 1990 Status and conservation of the Ganges River dolphin *Platanista gangetica* in the Karnali River, Nepal. *Biological Conservation*. 1993 Jan 1;66(3):159-69.
- [50] Schwartz MW. Modelling effects of habitat fragmentation on the ability of trees to respond to climatic warming. *Biodiversity & Conservation*. 1993 Feb 1;2(1):51-61.
- [51] Battin J. When good animals love bad habitats: ecological traps and the conservation of animal populations. *Conservation Biology*. 2004 Dec;18(6):1482-91.
- [52] Walker KF, Thoms MC. Environmental effects of flow regulation on the lower River Murray, Australia. *Regulated Rivers: Research & Management*. 1993 May;8(1-2):103-19.
- [53] Sowa SP, Annis G, Morey ME, Diamond DD. A gap analysis and comprehensive conservation strategy for riverine ecosystems of Missouri. *Ecological monographs*. 2007 Aug;77(3):301-34.

Figure 1

Figure 3

Figure 4

Figure 5

Figure 6

Figure 2

Figure 7