Immunomagnetic Sequential Ultrafiltration (iSUF) Platform for Enrichment and Purification of Extracellular Vesicles from Biofluids

Jingjing Zhang,¹ Luong TH Nguyen,¹ Richard Hickey,¹ Nicole Walters,¹ Xinyu Wang,¹ Kwang Joo Kwak,¹ L. James Lee,¹ Andre F. Palmer,¹ Eduardo Reátegui^{1,2*}

¹ William G. Lowrie Department of Chemical and Biomolecular Engineering, The Ohio State University, Columbus OH 43210

² Comprehensive Cancer Center, The Ohio State University, Columbus, OH 43210

*corresponding author: reategui.8@osu.edu

Abstract

Extracellular vesicles (EVs) derived from tumor cells have the potential to provide a much-needed source of non-invasive molecular biomarkers for liquid biopsies. However, current methods for EV isolation have limited specificity towards tumor-derived EVs that limit their clinical use. Here, we present an approach called immunomagnetic sequential ultrafiltration (iSUF) that consists of sequential stages of purification and enrichment of EVs (nonspecifically and specifically) in approximately 2 h. In iSUF, EVs present in different volumes of biofluids (0.5 mL to 100 mL) can be significantly enriched (up to 1000 times), with up to 99 % removal of contaminating proteins (e.g., albumin). The EV recovery rate by iSUF for cell culture media (CCM), serum, and urine corresponded to $98.0\% \pm 3.6\%$, $96.0\% \pm 2.0\%$ and $94.0\% \pm 1.9\%$, respectively ($p > 0.05$). The final step of iSUF enables the separation of tumor-specific EVs by incorporating immunomagnetic beads specific to a target subpopulation of EVs. Serum from a small cohort of clinical samples from metastatic breast cancer (BC) patients and healthy donors were processed by the iSUF platform and the isolated EVs from patients showed significantly higher expression levels of BC biomarkers (i.e., HER2, CD24, and miR21).

Introduction

Extracellular vesicles (EVs) are increasingly recognized as relevant diagnostic and therapeutic entities present in different biofluids¹. EVs are lipid particles with sizes that vary from 30 nm to a few microns². EVs are endogenously shed from the surface of cells through distinct mechanisms, leading to different types of vesicles³. Multivesicular bodies that contain smaller vesicles can fuse with the plasma membrane to release their internal vesicles (i.e., exosomes)⁴. Larger lipid vesicles can directly bud from the plasma membrane as microvesicles⁵. EVs carry various biological cargo, including proteins, RNA, and DNA fragments, giving EVs unique roles in regulating cell-cell communication⁶. Moreover, it has been shown that tumor EVs (tEVs) can tune cellular microenvironments at distant sites to promote angiogenesis, invasiveness, immunosuppression, and metastasis⁷⁻¹⁰.

Different proof of concept studies have used tEVs to develop liquid biopsy assays to diagnose and monitor cancer at different stages^{11,12}. EVs are more abundant than other circulating biomarkers (e.g., circulating tumor cells), and they are structurally more robust¹³. However, tEVs present in biofluids are surrounded by massive amounts of normal EVs (nEVs; secreted by healthy cells), and other biomolecules (e.g., albumin, lipoproteins, globulins)¹⁴, thus novel purification methods are required to isolate tEVs¹⁵. A recent survey on the methods used for isolation and characterization of EVs from research laboratories around the world reveals that more than 80% of researchers use ultracentrifugation (UC) for the isolation of EVs and western blotting for protein characterization¹⁶. Although UC and density gradient methods can be used to process different biofluids, they are labor-intensive, produce protein aggregate contaminants, and

are nonspecific towards EV type (derived from a tumor or normal cells)¹⁷⁻¹⁹. Other EV isolation methods, including polymeric or salt precipitation kits²⁰, size exclusion chromatography (SEC) columns (e.g., qEVs)²¹, and nano/microdevices have limitations²². Precipitation kits have low EV recovery rates, lack specificity, and have low purity²³. qEVs can separate EVs into different size fractions with high purity and low protein contamination, but have low EV recovery rates and are nonspecific for EV subpopulation²⁴. Recently, immunoaffinity methods that were developed for cell separation have been adapted for specific EV isolation²⁵. Microfluidic and plasmonic devices have been functionalized with antibodies to target different EV populations^{26,27}. However, the majority of these approaches target tetraspanins and annexins, which are ubiquitous proteins present in all EVs²⁸. Other attempts used epithelial cell adhesion molecule (EpCAM); however, this antigen is also expressed on normal epithelial EVs²⁹. Recently, we demonstrated the use of nanostructured polymeric brushes conjugated with epidermal growth factor receptor (EGFR) and integrated into a microfluidic channel to enhance specificity towards tumor-derived EVs isolated from glioblastoma (GBM) patients³⁰. Although this approach can achieve a remarkable 94% specificity towards tEVs, the limited amount of biofluid processed (1 to 1.5 mL of serum or plasma) and the retention of albumin, significantly limits its use for proteomics³¹.

Compromises have to be made when using a particular technology/methodology for the isolation of EVs³². Currently, there is a trade-off between sample volume and specificity in EV isolation technologies that limits quantitative molecular analysis of EV contents, ultimately impacting the utility of EVs in cancer diagnostics³³. Here, we present a novel approach termed immunomagnetic sequential ultrafiltration (iSUF) that

overcomes current limitations for EV enrichment and purification. iSUF combines three stages of ultrafiltration and immunoaffinity separation: a tangential flow filtration (TFF) step, a standard centrifugal enrichment step, and a magnetic-bead antibody-based EV capture step. Using iSUF, we demonstrate that small or large volumes of biofluid can be processed (~ 500 μ L or > 100 mL) while concomitantly removing up to 99 % of contaminating proteins (e.g., albumin, lipoproteins, globulins). We have demonstrated the use of iSUF for the enrichment of EVs present in three different types of biofluids: cell culture media (CCM), serum, and urine for which the sample processing time was under approximately 2 h. Another feature of iSUF is that it can enrich EVs up to 1000 times with an EV recovery rate higher than 94 %, which overcomes the limitations of other commercially available methods. To further validate the clinical utility of iSUF, we have processed serum samples from 10 metastatic breast cancer (BC) patients and demonstrated the presence of HER2, CD24 and miR21 biomarkers at significantly higher levels compared to healthy controls ($p < 0.05$).

Materials and Methods

Ethics

Healthy donors (HDs) were enrolled via an approved Institutional Review Board at The Ohio State University (IRB# 2018H0268 and 2019C0189). All HDs participants provided written informed consent. We also confirm that all experiments were carried out following relevant guidelines and regulations.

Materials

Hollow fiber cartridges (molecular weight cut off, MWCO: 500 kDa, material: polysulfone) were supplied by Repligen (Rancho Dominguez, CA). Amicon® ultra-15

centrifugal filter units (MWCO: 3, 10, and 30 kDa) were purchased from MilliporeSigma (Burlington, MA). Streptavidin-coated magnetic particles (3.0-3.9 μm) were obtained from Spherotech (Lake Forest, IL). For capturing EVs, Cetuximab was purchased from ImClone LLC (Branchburg, NJ), EpCAM, CD63, HER2 were obtained from R&D Systems (Minneapolis, MN). Antibodies were biotinylated using an EZ-Link™ micro Sulfo-NHS-biotinylation kit (ThermoFisher Scientific, Waltham, MA). For surface marker EV detection, CD24 was obtained from Novus Biological LLC (Centennial, CO), and HER2 was obtained from Cell Signaling Technology (Danvers, MA).

Cell culture and supernatant collection

U-251 glioblastoma (GBM), MCF-7 breast, and A375 melanoma cancer cell lines were supplied by American Type Culture Collection (ATCC, Manassas, VA). Cell lines were cultured in their recommended culture medium³⁴ containing 10% FBS and 1% penicillin-streptomycin at 37°C in a 5% CO₂ incubator. For isolation of EVs from CCM, U251, MCF7, and A375 were grown in T75 flasks to 90% cell confluence, followed by washing the cells twice with PBS. Culture medium with 10 % EV-depleted FBS was added to cells for 24 h. CCM was centrifuged at 1, 000 x g for 5 min at room temperature (RT) to discard cell debris before further processing. EV-depleted FBS was prepared by using the permeate of FBS filtered by tangential flow filtration (TFF) (MWCO: 300 kDa).

Healthy donor serum collection

10 mL of whole blood from healthy donors was collected into BD SST serum tubes (Thermo Fisher Scientific, Waltham, MA). Tubes were rocked 10 times and then gently placed upright to coagulate for 60 min. Then, the tubes were centrifuged at RT at 1,100 × g for 10 min. The serum was subsequently aspirated carefully and stored in 1 mL

aliquots at -80 °C. Additionally, we purchased six vials of healthy donor serum samples from ZenBio (Research Triangle Park, NC) collected according to FDA guidelines.

Healthy donor urine collection

Urine was also collected from the same healthy donors above, by either a first-morning or second-morning standard collection protocol³⁵. The urine volume collected was 10 to 100 mL. Urine was collected in sterilized 50-mL centrifuge tubes containing 4.2 mL protease inhibitor - a mixture of 1.67 mL 100 mM sodium azide (NaN₃), 2.5 mL phenylmethylsulfonyl fluoride (PMSF), and 50 µl Leupeptin (Millipore Sigma)³⁶. After collection, urine samples were frozen at -80 °C until processing time.

Cancer patient samples

1 mL of serum was collected from 10 metastatic BC patients. Samples were stored at -80 °C until use. All patient samples were collected from the biospecimens biobank through the Total Cancer Care (TCC) Program at the James Comprehensive Cancer Center at The Ohio State University.

Processing biofluids using the iSUF platform

The schematic workflow of the iSUF platform is shown in **Fig. 1**. In stage 1, TFF was used to enrich and diafiltrate EVs from the biofluid. Thus, biofluid from the sample feed reservoir was removed as filtrate/permeate from the TFF filter. Diafiltration is a fractionation process that removes smaller molecules (filtrate/permeate) through the filter and leaves larger molecules in the reservoir by adding a diafiltration solution into the reservoir at the same rate as the filtrate is generated. Briefly, a TFF pump circulates the biofluid through a hollow fiber filter cartridge at a controlled flow rate. Sample fractionation

depends on the hollow fiber membrane pore size (MWCO), which should be large enough to permeate proteins and free nucleic acids while small enough to retain EVs. During the enrichment step, freely permeable molecules are partially removed. To remove the remaining contaminants, a diafiltration step with PBS is necessary. The diafiltration processing time is proportional to the biofluid volume in the system³⁷, so diafiltration started with a total biofluid volume of 7 mL, which was the sum of the dead volume of the self-build TFF system (2 mL in the product container) and 5 mL remaining in the tubing that can be subject to further processing later. The liquid remaining in the tubing was necessary to protect the EVs from drying out and to enable constant volume diafiltration. Hence, CCM and urine were pre-enriched to a total volume of 7 mL. For the processing of serum, 0.5 mL of sample was diluted to a total volume of 7 mL in PBS and then processed with diafiltration. The input flow rate was kept at 35 mL/min using a peristaltic pump (Cole-Parmer, Vernon Hills, IL). The sample volume after stage 1 was approximately 2 mL (dead volume of the TFF system). At stage 2, ultra-centrifugal units with 10 kDa MWCO were used to further enrich the samples to 100 μ L at 3,000 $\times$ g for 20 min. For specific isolation of subpopulations of EVs, stage 3 of iSUF, Streptavidin-coated magnetic beads were functionalized with biotinylated capture antibodies (e.g., EpCAM, HER2, EGFR) overnight at 4 °C to target tEVs in spiked samples as well as patient samples. 100 μ L of the processed sample (after stage 2) was incubated with the antibody-coated beads for 1 h at RT. The efficiency of the iSUF platform for isolating tEVs was evaluated using flow cytometry and fluorescence microscopy. EVs were also characterized for their size, concentration, morphology, and molecular content (e.g., protein, RNA).

Processing biofluids using ultracentrifugation (UC)

CCM, serum, and urine samples were filtered using a syringe filter (pore size: 1.0 μm) and transferred to ultracentrifuge tubes (Beckman Coulter, Brea, CA) gently using a syringe and blunt needle (Becton, Dickinson and Company, Franklin Lakes, NJ). Ultracentrifuge tubes were sealed with a cordless tube topper (Beckman Coulter) after balancing, then were placed in a Type 55.2 Ti rotor (Beckman Coulter) and centrifuged in the Optima L-80 XP ultracentrifuge (Beckman Coulter) for 90 min at 4 °C at 100,000 $\times g$. The supernatants were discarded carefully after UC, and pellets were re-suspended in 100 μL of PBS.

Processing biofluids using commercially available EV isolation kits

Using the Total Exosome Isolation Reagent (TEIR, Invitrogen, Carlsbad, CA), EVs were isolated from 0.5 mL serum according to the manufacturer's instructions. Briefly, 0.5 mL of serum was mixed with a proprietary reagent provided in a kit and incubated for 30 min at 4 °C. After mixing, the sample was centrifuged at 10,000 $\times g$ for 10 min at RT. EVs pellets were resuspended in 100 μL PBS. For size-exclusion chromatography, 0.5 mL serum was loaded into a qEV column (Izon Science, Medford, MA), and flushed with PBS, fractions 7-12 were collected and pooled in 3 mL according to the manufacturer's instructions.

EV quantification

The size and concentration of EVs were measured using a tunable resistive pulse sensing (TRPS) method (qNano, Izon Science). Different nanopore stretchable membranes (NP150, NP300, NP600, NP800, and NP1000) were selected to cover the

wide size range of EVs³⁸. Samples were first filtered through 1- μ m filters before processed in the qNano instrument. EVs with a size range of 70 nm to 1000 nm were characterized.

Immunofluorescence staining

After EVs were captured on the functionalized magnetic beads using a cocktail of a recombinant chimeric EGFR monoclonal antibody (Cetuximab, Erbitux®, ImClone LLC, Branchburg, NJ), a goat EpCAM/TROP-1 polyclonal antibody (#AF960, R&D Systems, Minneapolis, MN) and a goat ErbB2/Her2 polyclonal antibody (#AF1129, R&D Systems), they were blocked with 3% (w/v) BSA and 0.05% (v/v) Tween® 20 in PBS for 1 h at RT, and then stained with either a rabbit HER2/ErbB2 monoclonal antibody – PE conjugate (#98710S, Cell Signaling Technology, Danvers, MA) or a mouse CD24 monoclonal antibody - Alexa Fluor™ 594 conjugate (#NB10077903AF594, Novus Biologicals™) for 1 h at RT.

Molecular beacon design and quantification

Molecular beacon (MB) (listed 5' – 3') targeting miR-21 used in this study was T+CA A+CA /iCy3/ +TCA +GT+C T+GA TAA GCT AAC TTA TCA GAC TGA /3BHQ_2/. Locked nucleic acid (LNA) nucleotides (positive sign (+) bases) were incorporated into oligonucleotide strands to improve the thermal stability and nuclease resistance of MBs for incubation at 37 ° C. The designed MBs were custom synthesized and purified by Sigma-Aldrich. An aqueous solution of MBs in PBS was vigorously mixed with a lipid formulation of dioleoyl-3-trimethylammonium propane (DOTAP), cholesterol, phosphatidylcholine (POPC), and 1, 2-distearoyl-sn-glycero-3-phosphoethanolamine-poly(ethylene glycol) (DSPE- PEG) in 200 proof ethanol, and then sonicated for 5 min using an ultrasonic bath. The MB/lipid mixture was subsequently injected into PBS,

vortexed, and sonicated for 5 min. Finally, it was dialyzed with a 20 kDa MWCO dialysis bag to remove free MBs. After EVs were captured on magnetic beads, they were incubated with the prepared MBs for 2 h at 37 °C before imaging.

Flow cytometry and fluorescence microscopy.

tEVs from U251 GBM cells were stained with a lipophilic fluorescent dye, SP-DiOC18(3) (ThermoFisher Scientific) for 20 min, the excess dye was washed out. 100 μ L fluorescent EVs ($\sim 10^{11}$ particles/mL) were then spiked into 500 μ L of serum ($\sim 10^{12}$ particles/mL) and 100 mL of urine ($\sim 10^9$ particles/mL) from healthy donors. Each of the samples was processed by iSUF and recovered in 100 μ L of PBS (stage 1 to 3). Non-spiked fluorescent tEVs were also captured on functionalized beads as a positive control. After washing with PBS, the captured EVs were analyzed by imaging flow cytometry (Amnis, ImageStream^X Mark II Imaging Flow Cytometer, Luminexcorp, Austin, TX); also images were taken using a fluorescence microscope (Nikon Eclipse Ti Inverted Microscope System) with a 100x oil immersion lens. For comparison, samples were also processed by UC and resuspended in 100 μ L PBS. For total RNA quantification, captured EVs were lysed, and RNA was extracted and quantified using the same procedures mentioned above.

Definition of terms used in this study.

$$TFF \text{ Recovery Rate (\%)} = \frac{C_{TFF} \times V_{TFF}}{C_{E0} \times V_{E0}} \times 100\% \quad \text{Equation (1)}$$

Where V_{E0} and C_{E0} are the initial volume and concentration of the sample; V_{TFF} (in 2 mL) and C_{TFF} (measured by qNano pore size [NP150, NP300, NP600, NP800, and NP1000]) are the final volume and concentration of the TFF product.

$$\text{Remaining Protein (\%)} = \frac{C_{Pt} \times V_t}{C_{P0} \times V_0} \times 100\% \quad \text{Equation (2)}$$

Where V_0 and C_{P0} are the initial volume of the sample and concentration of free proteins in the sample, respectively; V_t and C_{Pt} are the volume of the TFF retentate and concentration of free proteins in the retentate at a specified time t , respectively. Concentration of free proteins was measured using the bicinchoninic acid (BCA) assay. The free protein removal efficiency can be calculated by subtracting the percentage of remaining protein in the final TFF product from 100%.

$$\text{Enrichment Factor} = \frac{C_{Ef}}{C_{E0}} \quad \text{Equation (3)}$$

Where C_{E0} is the EV concentration in the initial sample; C_{Ef} is the EV concentration in the final iSUF product. Concentration of samples were measured by qNano pore size [NP150, NP300, NP600, NP800, and NP1000]).

$$\text{Recovery Rate (\%)} = \frac{C_{Ef} \times V_{Ef}}{C_{E0} \times V_{E0}} \times 100\% \quad \text{Equation (4)}$$

Where V_{E0} is the initial volume of the product; V_{Ef} is the volume of the final iSUF product.

$$\text{Purity} = \frac{C_{Ef}}{C_{Pf}} \times 100\% \quad \text{Equation (5)}$$

Where C_{Ef} is the final concentration of the iSUF product (in 100 μL); C_{Pf} is the remained free protein concentration in the product.

$$\text{Coefficient of variation} = \frac{\sigma}{\mu} \times 100\% \quad \text{Equation (6)}$$

The coefficient of variation (CV) is a measure of relative variability. It is the ratio of the standard deviation (σ) to the mean (μ)

Statistical Analysis.

Data are expressed as the mean $\pm$ STD. A significant test between different mean values was evaluated using one-way ANOVA in JMP Pro 16 software provided by The

Ohio State University. Differences between samples were considered statistically significant for $p < 0.05$.

Results

Optimization of the iSUF platform

To overcome current limitations of the enrichment and purification of EVs and on-demand EV subpopulation characterization, we developed the iSUF platform (**Fig. 1 A**) which includes three stages: (1) tangential flow filtration (TFF) for the enrichment and purification of EVs, (2) centrifugation for further enrichment of EVs, and (3) immunomagnetic affinity selection for desired EV subpopulation isolation. We used the iSUF platform to process various volumes (0.5-100 mL) of different biofluids (CCM, serum, and urine).

To design stage 1, many parameters of TFF processing required optimization, including the selection of membrane pore size (MWCO), sample processing temperature, sample flow rate, pressure, and sample protein concentration. We tested membrane filters with two MWCO sizes (300 and 500 kDa) to determine the optimal MWCO that maximizes the removal of free proteins and nucleic acids while reducing processing time. Our experiments showed that 500 kDa filter membranes were able to remove up to 99 % of free proteins with over 99% EV recovery rate (Equation 1, **Supplementary Fig. 1**). When a 300 kDa membrane filter was used, only 80 % of free proteins were removed. Moreover, a 500 kDa membrane filter was chosen since it processed samples 2~3 times faster than a 300 kDa membrane filter (**Supplementary Table 1**). The TFF stage 1 of iSUF was run at 4°C to minimize EV degradation³⁹. We further tried to optimize sample processing time, which was highly dependent on the flow rate. The flow rate was linearly

associated with the shear rate generated by the filter based on the manufacturer's protocol (Repligen), which exerted a shear force on the EVs. Then, we used a flow rate of 35 mL/min to maintain a shear rate below 5000/s⁴⁰. High flow rates increased the system pressure, mainly when the protein concentration of the sample was high (> 15 mg/mL). We kept the pressure of the system below 10 psig to avoid leakage and maximize the lifespan of the 500 kDa filter. We used dilutions of fetal bovine serum (FBS) to test the effect of protein concentration on system pressure at 35 mL/min. Our results showed that protein concentration must be equal or lower than 15 mg/mL to maintain the pressure of the system below 10 psig to protect the filter (**Supplementary Fig. 2**).

At stage 2 of iSUF, EVs were loaded into centrifugal filter units and were spun down at 3,000 x g. We compared the recovery rates of EVs and processing time for different filter pore sizes. A 3 kDa filter unit obtained over 99% recovery rate and took 60 min to spin down, while a 10 kDa unit obtained a 95% recovery rate in 20 min, and a 30 kDa filter obtained only a 70% recovery rate and took 15 min to spin down (**Supplementary Fig. 3**). We selected the 10 kDa filter unit to maintain a high recovery rate while reducing sample processing time. To enrich tEVs (stage 3 of iSUF), we tested incubation of 3 µm magnetic beads with different concentrations of biotinylated antibodies (10, 20, 100 µg/mL) for 1 h and 2 h at RT, and overnight at 4 °C. Overnight incubation with 100 µg/mL of antibody exceeded the bead-antibody binding efficiency, while 20 µg/mL of antibody was able to yield a high bead-antibody binding efficiency (>90%). Then, we examined the volume ratio of beads to EVs at 5 µL /100 µL, 20 µL /100 µL, and 80 µL /100 µL during the bead/tEVs incubation step. The 20/100 µL ratio achieved the highest bead-tEVs capture efficiency (>90%).

iSUF platform for processing biofluids

A flow rate of 35 mL/min for stage 1 of iSUF was applied since the protein concentrations of the different biofluids were below 15 mg/mL (**Supplementary Table 2**). We tested the ability of our platform to purify and enrich EVs from three different biofluids (i.e., CCM, serum, and urine; **Fig. 1B**). 50 mL of CCM, and 100 mL of urine were enriched to 7 mL. Subsequently, PBS diafiltration buffer was used to remove the remaining protein contaminants from CCM and urine in 80 min and 100 min, respectively. The percentage of remaining protein is defined as the mass of free proteins in the TFF retentate at a specified time divided by their initial mass in the sample (Equation 2). For serum, the initial high concentration of proteins (> 80 mg/mL) required an initial dilution of 0.5 mL of the sample in 7 mL of PBS. Subsequently, PBS diafiltration buffer was used to remove the remaining protein contaminants from serum in 120 min. We first tested iSUF with a 10% BSA solution for which an SDS-PAGE gel showed extensive removal of albumin (**Supplementary Fig. 4**). Moreover, analysis of the purified samples by an SDS-PAGE gel indicated that iSUF removed BSA from CCM, human serum albumin (HSA) and globulins from serum, and Tamm-Horsfall glycoprotein (THF) and HSA from urine. (**Supplementary Fig. 5, 6, 7**).

Concentration, size distribution and microscopy characterization of EVs

For 50 mL of CCM, 0.5 mL of serum, and 100 mL of urine, the enrichment factors were 489 ± 18 , 4.8 ± 0.1 , and 942 ± 19 , respectively ($n = 5$; **Fig. 2A**) (Equation 3). Accordingly, the EV recovery rate (the ratio of the total number of EVs post-processing by iSUF to the total number of EVs pre-processing) was $98\% \pm 3.6\%$, $96\% \pm 2.0\%$ and $94\% \pm 1.9\%$ for CCM, serum, and urine, respectively (Equation 4). Considering that EVs

are heterogeneous in size, we also tested the EF across a wide size range of EVs (70 nm - 1 μ m) in CCM with similar results ($n = 5$; $p > 0.05$; **Fig. 2B**). Moreover, the EV concentrations obtained by iSUF was greater than the concentrations obtained by UC. This difference was consistent across the 70 nm to 1 μ m size range ($n = 5$; $p < 0.05$; **Fig. 2C**), with iSUF enriching EVs at two to three orders of magnitude higher than UC (approximately 10^{11} EVs concentrated by iSUF and 10^9 for UC). Similar results were obtained with comparisons for the enrichment of EVs present in serum and urine. With iSUF, EVs from serum and urine were enriched almost at the same level (10^{12} EVs/mL; **Fig. 2D**). We confirmed the presence of EVs by using atomic force and electron microscopy on the different biofluids processed. The majority of isolated EVs exhibited a round morphology with heterogeneous size distribution. Cryo transmission electron microscopy images (TEM) images of isolated EVs showed the presence of a double-layered lipid membrane, a representative characteristic of EVs. (**Fig. 2E & 2F**).

To further test our iSUF platform, we performed comparative studies with different commercially available EV isolation methods: qEV, TEI, and UC. 0.5 mL of serum from healthy donors were processed with different EV isolation platforms. EVs demonstrated similar smaller and larger size distribution for all platforms, but iSUF obtained a higher EV concentration than the other methods within the 70 nm - 1 μ m size range (**Fig. 3A**). We also compared the mean size of subpopulations of EVs based on their physical characteristics using different methods³⁸. **Fig. 3B** shows the mean size of small EVs (sEVs) and medium/large EVs (m/l EVs). For both size ranges, EVs processed by iSUF were significantly smaller than UC ($n = 5$; $p < 0.05$). For l/m EVs, EVs purified by UC were larger than all other techniques ($n = 5$; $p < 0.05$). Then, we compared the total

concentration and purity of isolated EVs using different methods. iSUF enriched EVs significantly more efficiently than all the tested methods ($n = 5$; $p < 0.05$; **Fig. 3C**). The concentration of EVs enriched by iSUF was 51 ± 18 , 7.0 ± 2.0 , and 56 ± 21 times higher than qEV, TEI, and UC. The purity of isolated EVs was normalized and evaluated in terms of the ratio between EV concentration and remaining contaminating protein concentration present in samples after purification using the different methods (Equation 5). The purity of the isolated EVs by iSUF was 100-10000 times higher than the different tested methods (**Fig. 3C**).

Repeatability experiments for the EV recovery rate and protein removal efficiency from samples processed by iSUF were also conducted (CCM, $n = 10$; serum, $n = 10$; Urine, $n = 10$). Both sEVs and m/l EVs in all biofluids demonstrated a mean value above 94% for the EV recovery rate. The recovery rate of sEVs in CCM, serum, and urine exhibited 4.2%, 3.7%, and 2.4% coefficient variations, respectively, while m/l EVs showed 3.7%, 3.3%, and 2.5% coefficient variations (Equation 6). Protein removal efficiencies were also tested, CCM, serum, and urine presented up to 99% removal efficiency with coefficient variations of 0.9%, 4.0%, and 2.1% (**Supplementary Fig. 8**).

Molecular content quantification and characterization of EVs

We quantified the total amount of protein and RNA present in EVs isolated by iSUF for different biofluids. For CCM, the quantity of RNA obtained was 11 ± 8.2 ng/mL, thus giving a 9-fold higher concentration of RNA when compared to UC (**Fig. 4A**). For protein analysis, the protein concentration was 6 ± 1.8 μ g/mL, almost 8-fold more protein obtained than when the same biofluid was processed by UC (**Fig. 4B**). Total RNA and protein quantification were carried out for serum and urine from 5 healthy donors. 0.5 mL

of serum and 100 mL of urine processed by iSUF produced 54 ± 40 ng/mL and 0.3 ± 0.1 ng/mL of RNA, respectively (**Fig. 4C**). The protein concentration was 1250 ± 480 μ g/mL for serum and 3.1 ± 3.0 μ g/mL for urine (**Fig. 4D**). The differences in protein concentration of EVs in original CCM, processed by UC, and processed by iSUF were also demonstrated using CD63 and CD9 western blot analysis. Also, the expression level of CD9 and CD63 in original urine and serum samples, and after iSUF processing demonstrated superiority enrichment and purification performance of the iSUF (**Fig. 4E**). Full-length western blot images are displayed in (**Supplementary Fig. 9**). Moreover, we compared the RNA content obtained from EVs enriched from 0.5 mL of serum using different commercially available methods (**Supplementary Table 3**). The total RNA content obtained using iSUF was 54 ± 41 ng, which was significantly higher than other methods that only obtained 10 ± 9.8 ng (**Fig. 4F**).

Next, we processed different volumes of serum and urine with iSUF to identify the equivalent volumes of biofluid that will produce comparable concentrations of EVs, total RNA, and total protein. We started with 0.5 mL of serum and different volumes of urine (i.e., 100, 75, 65, 10 mL). For different volumes of urine and 0.5 mL of serum, the enriched concentrations of EVs were comparable ($\sim 10^{12}$ particles/mL, **Supplementary Fig. 10**). The total RNA content obtained from EVs from serum and urine also exhibited comparable values of 60 ± 35 ng and 75 ± 40 ng, respectively. However, analysis of protein content of the isolated EVs from both biofluids showed that the protein content of EVs isolated from urine was 10-fold lower than the protein content of EVs in serum, which was 0.4 ± 0.1 mg and 40 ± 10 mg, respectively.

Immunomagnetic affinity selection of the iSUF platform

tEVs are surrounded by many normal EVs (nEVs) that require removal to perform accurate molecular analysis⁴¹. One way to isolate tEVs is to exploit the presence of specific surface markers⁴². Our iSUF platform enables the separation of subpopulations of EVs by capturing them on magnetic beads through immunoaffinity. To demonstrate our approach, we used a model system that consisted of spiking fluorescently labeled tEVs from a cancer cell line (i.e., U251 EVs) in 0.5 mL of serum and 100 mL of urine from healthy donors (n = 3). 3- μ m magnetic beads were functionalized with EGFR as the capture antibody. All samples were processed through all stages of iSUF, and the captured subpopulation of EVs was characterized and quantified using TIRF and imaging flow cytometry (**Fig. 5A**). A comparable fluorescent signal was obtained between the positive control (tEVs directly from U251 cell supernatant) and spiked samples processed in serum and urine, which verified the high recovery efficiency of tEVs (88 ± 1 for serum, and 93 ± 2 for urine), and their purity. The biofluids spiked with tEVs were also processed by UC, which yielded significantly lower fluorescent signal (3.1 ± 0.1 times lower, **Fig. 5A**). Furthermore, to determine tEV RNA isolation efficiency by iSUF, lysed U251-EVs (positive control) and bead-isolated U251-EVs were extracted. iSUF showed over 90% RNA isolation efficiency for the EGFR⁺ tEVs (**Fig. 5B**). Moreover, the RNA content of bead-extracted tEVs from serum using the iSUF platform was 7-fold higher than UC, which was mainly attributed to the high yield of total EVs recovered in iSUF (**Fig. 5C**). The high RNA content of tEVs from serum isolated using iSUF also confirmed the ability of iSUF to enrich and retrieve tumor targets from mixed populations of EVs.

Detection of proteins and miRNA in EVs from clinical samples

The emergence of targeted therapies require a precise characterization of the molecular subtypes of a patient's tumor⁴³. For BC patients (e.g., luminal A/B, triple-negative), the molecular subtype is typically determined by testing a tissue biopsy for the presence or absence of three essential proteins: estrogen receptor (ER), progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER2)⁴⁴. However, a tissue-based tumor profile is subject to sampling bias, provides only a snapshot of tumor heterogeneity, and cannot be obtained repeatedly⁴⁵. These limitations restrict early cancer detection and significantly contribute to the overdiagnosis and overtreatment of BC patients⁴⁶. The analysis of EVs present in biofluids of cancer patients provides a minimally invasive method to quantify different biomarkers that would enable precise diagnosis or response to treatment⁴⁷. To test the potential use of the iSUF platform in screening biomarkers for BC, we processed 0.5 mL of serum from 10 metastatic BC patients and quantified the expression levels of HER2, CD24, and miR21 on patient isolated EVs⁴⁸⁻⁵⁰. After stage 3 of iSUF, magnetic beads were isolated using a magnet and incubated with detection antibodies and MBs. Then, the magnetic beads were processed with imaging flow cytometry and TIRF microscopy. Similar volumes of serum from healthy donors were processed and analyzed. The age of the healthy donors was between 21 and 68 years old. Although the age range was wider than from our BC patient cohort, a comparison of levels of expression for HER2, CD24, and miR21 between healthy donors samples did not show statistically significant differences ($p > 0.05$, **Supplementary Fig. 11**). We obtained representative images and histograms that quantify the number of beads and their corresponding fluorescent intensities (**Fig. 6A & 6B**) for BC patient and healthy donor samples. Both protein and miRNA show significantly

higher expression in BC patient samples. ($p < 0.05$; **Fig. 6B**), which suggested that iSUF can differentiate tumor biomarkers (**Supplementary Table 4**). Moreover, we have characterized the CD9 and CD63 levels of protein expression in serum samples from both cohort of samples. Although in this case, similar levels of expression were obtained ($p > 0.05$, **Fig. 6A & 6B**).

Discussion

Recently, EVs have been explored for diagnostic and therapeutic applications, including liquid biopsy assays for cancer diagnostics, and nanocarriers for drugs and nucleic acids^{51,52}. The large number of tEVs compared to other rare biomarkers (i.e., circulating tumor cells, CTCs) makes them more statistical reliable⁵³. Innovative methods developed for the enrichment and purification of EVs should remove all contaminants (e.g., free proteins), have a high yield, work amongst different biofluids, and maintain the integrity of the EVs. We have engineered a new platform that includes a TFF enrichment and purification stage, a centrifugal-unit enrichment stage, and a magnetic separation stage for specific isolation of subpopulations of EVs (e.g., tEVs). In stages 1 and 2, iSUF performs enrichment and purification of all EVs (e.g., tEVs and nEVs). In stage 3, tEVs are isolated based on their tumor-specific surface markers. 300 kDa and 500 kDa TFF filters were initially selected for this study because the molecular weights of the major free proteins in CCM (e.g., BSA, 65 kDa), serum (e.g., HSA, 65 kDa) and urine (e.g., THP, 98 kDa) were below 300 kDa⁵⁴⁻⁵⁶. After testing, a 500 kDa TFF filter was finally chosen for subsequent experiments because of its minimal EV loss and faster processing time (~ 2 h).

Different biofluids required different TFF diafiltration buffer volumes and different scale of TFF hollow fiber filters to remove most contaminants. CCM and urine contain relatively low concentrations of protein contaminants^{57,58}, so a small total diafiltration buffer volume was required to remove the majority of free proteins. Unlike CCM and urine, the serum has a significant amount of free proteins⁵⁹. Therefore, a large total diafiltration buffer was necessary to remove the free protein in serum. iSUF was able to achieve up to 99 % of free protein removal efficiency. Free proteins removed by iSUF were shown as decreased bands on the SDS-PAGE gel. Other bands were increasing on the SDS-PAGE gel because those correspond to different proteins present on EVs, which were enriched by the iSUF⁶⁰⁻⁶⁴. Also, there are Very-low-density lipoproteins (VLDL) that overlap in size with EVs that also might be enriched^{2,65,66}.

Although TFF can concentrate and purify samples, the final product volume is mostly dependent on the dead volume of the specific TFF system. Therefore, further enrichment is necessary to concentrate the samples (e.g., spin down to ~ 100 µL) for applications such as the diagnosis of rare biomarkers⁶⁷. Centrifugal units with different MWCOs (3, 10, and 30 kDa) were evaluated in terms of processing time and yield in our study. A larger MWCO size required shorter centrifugation time, but more EVs were lost because prolonged high-speed centrifugation elongated EVs into an oval shape which made them squeeze through the filter membrane^{68,69}. The centrifugal unit with the 10 kDa MWCO was optimal with the shortest processing time and highest yield.

We also compared EVs purified from serum and urine samples of healthy donors. EVs from urine and serum can serve as prognosis biomarkers for clinical analysis^{70,71}. Serum is the most commonly used biofluid in the clinical setting with the highest EV

concentration⁷². Compared to serum, urine collection is minimally invasive and can be obtained in larger volumes⁷³, but it usually suffers from much lower EV concentration⁷⁴. Interestingly, using our iSUF platform, the EV RNA concentration in the final products of urine samples (originating from ~ 100 mL of collected sample volume) were comparable to those of serum samples (originating from 0.5 mL of collected sample volume). This suggests that clinical diagnosis by urinary EVs is possible. However, there is still a concern towards urinary EV collection because of large variabilities in the urine volume and its EV concentration. More efforts are needed to come up with a gold standard protocol for urine collection and processing. For protein in urinary EVs, we found that the amount of protein was lower than the amount of protein obtained from serum EVs, which might be explained by the degradation of EV membrane proteins by urine proteases⁷⁵.

In this study, we compared the performance of iSUF with three different EV isolation techniques (TEI, qEV, and UC) to determine EV concentration, purity, and quantity of RNA recovered. As other authors have discussed^{76,77}, we found that UC, a traditional method for EV isolation, has raised concerns about the integrity, yield, and purity of EVs after purification. Interestingly, we found UC-isolated EVs were larger than other platforms, one possible explanation is the presence of extensive levels of protein aggregates that cause bias. For our UC method, washing steps using multi-rounds ultracentrifugation were not applied, since it was shown that purity of the samples did not increase significantly when adding washing steps, and a significant loss of enriched EVs could occur^{54,78-82}. Moreover, although qEV obtained a relatively pure product, its recovery rate was low; TEI obtained a relatively higher EV number, while retaining

massive amounts of free protein (**Fig. 3C**). Therefore, low EV concentration and purity will impact the accuracy of molecular analysis of EVs⁸³.

We used the MISEV 2018 guidelines to define EVs: small EVs (sEVs) and medium and large EVs (m/l EVs)³⁸. Following these definitions, we found that the concentration and average size of EVs in patient samples and healthy controls varied (**Supplementary Table 4**). Different factors may explain these differences. Subtypes and stages of BC in the patient cohort, may contribute to the difference in the metrics. Moreover, we obtained our patient samples from a biobank, we had minimal control of the storage conditions, thus it was possible that some degradation of the samples could occur. It is known that factors such as material of the container, temperature of storage, and cycles of freezing and thawing may also influence the concentration and mean size of EVs⁸⁴⁻⁸⁷. Considering all these aspects, the size distributions and concentrations of EVs that we obtained for the different samples are within the reported values in the literature. Previously concentrations of EVs in serum from cancer patients are between $10^9 - 10^{11}$ particles/mL

88-91.

Differentiating EV particles and other non-EV particles (e.g., lipoproteins) when measuring size distribution and concentration is challenging for all current methods, including TRPS and nanoparticle tracking analysis NTA^{65,92-96}. This is because the size range of different lipoproteins (e.g., 7 – 13 nm (HDL); 21 – 27 nm (LDL); and 30 – 90 nm (VLDL))⁹⁷ may overlap with the reported size ranges for sEVs and m/l EVs³⁸. Thus, current methods for EV size characterization and concentration have certain limitations^{87,98-100}. For iSUF, we used the TRPS method to measure size distribution and concentration of EVs. Different nanopore stretchable membranes (NP150, NP300,

NP600, NP800, and NP1000) were selected, covering a broad size range of EVs such as 70 nm to 1000 nm¹⁰¹. The 70 nm cut-off were chosen since EVs in samples processed by iSUF, exhibited a range above 70 nm. Choosing a small stretchable membrane such as NP100 and NP80 that have limits of detection of 50 nm and 40 nm, respectively, did not change the size ranges of EVs in our samples after iSUF (**Supplementary Fig. 12**). Thus, iSUF intrinsically filters out the majority of lipoproteins and EVs that overlap in size. This effect is beneficial since we are trying to characterize EVs with minimal contribution of lipoproteins. We consider the 70 nm cut-off used for the TRPS measurements as appropriate. Moreover, above this value the different characterization metrics are considered accurate. The particle count measurements can verify this for each one of the methods used in our study (**Supplementary Fig. 13**). A linear relationship indicates an accurate measurement³⁰. In **Fig 3**, we found that the lower limit of detection varies for the different methods, indicating an intrinsic loss of EVs during processing samples with different methods. However, iSUF shows better recovery for sEVs and m/l EVs. The wide upper range showed by UC could be indicative that we measure aggregates of EVs¹⁰²⁻¹⁰⁵. Our measurements are in agreement with MISEV 2018 since sEVs, and m/l EVs are in the 100 ~ 150 nm and larger than 200 nm, respectively^{38,62,106-110}.

We are interested in the enrichment and isolation of tEVs to characterize tumor-related proteins and RNAs. Like other immunoaffinity methods¹¹¹⁻¹¹³, iSUF captured and isolated EVs using specific tumor surface proteins. iSUF differentiated metastatic BC patients from healthy donors by detecting significantly higher expression levels of proteins and RNA biomarkers present in EVs (e.g., HER2, CD24, and miR21). Based on previous reports, HER2 and miR21 are cancer-associated protein and microRNA species, and are

known to be overexpressed in metastatic BCs^{114,115}. Compared to miR21 and HER2, CD24 is relatively less investigated in BC but was previously identified as being released from BC stem cells¹¹⁶. Furthermore, a recent study indicated that serum CD24 is elevated among BC patients¹¹⁷. Moreover, it is important to note that one of these biomarkers may not be a reliable predictor of BC alone. However, the combination of several biomarkers can serve as a tool for BC risk assessment.

In conclusion, iSUF was proposed for rapid, efficient, and specific isolation of EVs from different biofluids. The EV recovery rate mean value was above 94%, with 90% specific isolation of tEVs and negligible concentrations of free proteins and nucleic acids. Although iSUF does not process a sample in the shortest time (80 min ~ 120 min), its versatility working with different biofluids, sample volumes, and high purity after sample processing constitutes unmatched advantages over current methods used in the field. Overall, we found that the iSUF platform isolated and enriched EVs from a scaled-up sample volume with high purity and yield in a sterile and quick manner simultaneously, and isolated tEVs with high specificity, while other current methods could not guarantee all of those conditions at the same time. Furthermore, we recognize that the iSUF platform has potentially broad clinical applications beyond liquid biopsies for cancer diagnosis or monitoring.

Conflicts of interest

J.Z., L.T.H.N., R. H., K.K., A.F.P., J.L.L., and E.R. have a provisional patent application relevant to this study.

Acknowledgments

We acknowledge all the patients and healthy volunteers who participated in this study. This work was supported by the U.S. National Institutes of Health (NIH) grants UG3TR002884 (ER). and R01HL126945, R01HL138116, and R01EB021926 (AFP). Additional support for E.R. was provided by the William G. Lowrie Department of Chemical and Biomolecular Engineering and the Comprehensive Cancer Center at The Ohio State University.

Contributions

E.R and A.F.P. conceived the idea. E.R., L.T.H.N, and J.Z design the study. J.Z. performed most of the experiments and data analysis, prepared all tables and figures, and drafted the original manuscript with inputs from all authors. L.T.H.N established protocols for protein/RNA analysis and immunomagnetic beads and performed some of the experiments. R. H. performed equipment installation and calibration. N.W. performed sample collection and supported with editing the manuscript. X.W. developed the quantification code for the third stage of iSUF. K.K. designed molecular beacons for targeting miR21. A.F.P. supervised the TFF experiments and edited the manuscript. L.J.L. supervised the design of molecular beacons and edited the manuscript. E.R. supervised the whole study and edited the manuscript. All authors provided critical feedback and helped to shape the research, analysis, and manuscript.

References

- 1 Raposo, G. & Stoorvogel, W. J. J. o. C. B. Extracellular vesicles: exosomes, microvesicles, and friends. **200**, 373-383 (2013).
- 2 Mathieu, M., Martin-Jaular, L., Lavieu, G. & Théry, C. Specificities of secretion and uptake of exosomes and other extracellular vesicles for cell-to-cell communication. *Nature Cell Biology* **21**, 9-17, doi:10.1038/s41556-018-0250-9 (2019).
- 3 Yáñez-Mó, M. *et al.* Biological properties of extracellular vesicles and their physiological functions. *Journal of Extracellular Vesicles* **4**, 27066, doi:10.3402/jev.v4.27066 (2015).
- 4 Théry, C., Zitvogel, L. & Amigorena, S. Exosomes: composition, biogenesis and function. *Nat Rev Immunol* **2**, 569-579, doi:10.1038/nri855 (2002).
- 5 Cocucci, E., Racchetti, G. & Meldolesi, J. Shedding microvesicles: artefacts no more. *Trends Cell Biol* **19**, 43-51, doi:10.1016/j.tcb.2008.11.003 (2009).
- 6 Hessvik, N. P. & Llorente, A. Current knowledge on exosome biogenesis and release. *Cellular and Molecular Life Sciences* **75**, 193-208, doi:10.1007/s00018-017-2595-9 (2018).
- 7 Filipazzi, P., Bürdek, M., Villa, A., Rivoltini, L. & Huber, V. Recent advances on the role of tumor exosomes in immunosuppression and disease progression. *Semin Cancer Biol* **22**, 342-349, doi:10.1016/j.semcancer.2012.02.005 (2012).
- 8 Kahlert, C. & Kalluri, R. Exosomes in tumor microenvironment influence cancer progression and metastasis. *J Mol Med (Berl)* **91**, 431-437 (2013).
- 9 Li, K., Chen, Y., Li, A., Tan, C. & Liu, X. Exosomes play roles in sequential processes of tumor metastasis. *Int J Cancer* **144**, 1486-1495, doi:10.1002/ijc.31774 (2019).
- 10 Wang, J. *et al.* Multiple myeloma exosomes establish a favourable bone marrow microenvironment with enhanced angiogenesis and immunosuppression. *J Pathol* **239**, 162-173, doi:10.1002/path.4712 (2016).
- 11 Brock, G., Castellanos-Rizaldos, E., Hu, L., Coticchia, C. & Skog, J. J. T. C. R. Liquid biopsy for cancer screening, patient stratification and monitoring. **2015** **4**, 280-290 (2015).
- 12 Li, A., Zhang, T., Zheng, M., Liu, Y. & Chen, Z. Exosomal proteins as potential markers of tumor diagnosis. *Journal of Hematology & Oncology* **10**, 175, doi:10.1186/s13045-017-0542-8 (2017).
- 13 Zhang, W. *et al.* Liquid Biopsy for Cancer: Circulating Tumor Cells, Circulating Free DNA or Exosomes? *Cell Physiol Biochem* **41**, 755-768, doi:10.1159/000458736 (2017).
- 14 Théry, C., Amigorena, S., Raposo, G. & Clayton, A. Isolation and characterization of exosomes from cell culture supernatants and biological fluids. *Curr Protoc Cell Biol* **Chapter 3**, Unit 3 22, doi:10.1002/0471143030.cb0322s30 (2006).
- 15 Li, P., Kaslan, M., Lee, S. H., Yao, J. & Gao, Z. Progress in Exosome Isolation Techniques. *Theranostics* **7**, 789-804, doi:10.7150/thno.18133 (2017).
- 16 Momen-Heravi, F. in *Extracellular Vesicles* 25-32 (Springer, 2017).
- 17 Mol, E. A., Goumans, M. J., Doevendans, P. A., Sluijter, J. P. G. & Vader, P. Higher functionality of extracellular vesicles isolated using size-exclusion chromatography

- compared to ultracentrifugation. *Nanomedicine* **13**, 2061-2065, doi:10.1016/j.nano.2017.03.011 (2017).
- 18 Yuana, Y., Levels, J., Grootemaat, A., Sturk, A. & Nieuwland, R. Co-isolation of extracellular vesicles and high-density lipoproteins using density gradient ultracentrifugation. *Journal of extracellular vesicles* **3**, 10.3402/jev.v3.23262, doi:10.3402/jev.v3.23262 (2014).
- 19 Koh, Y. Q., Almughlliq, F. B., Vaswani, K., Peiris, H. N. & Mitchell, M. D. Exosome enrichment by ultracentrifugation and size exclusion chromatography. *Front Biosci (Landmark Ed)* **23**, 865-874, doi:10.2741/4621 (2018).
- 20 Furi, I., Momen-Heravi, F. & Szabo, G. Extracellular vesicle isolation: present and future. *Annals of translational medicine* **5**, 263-263, doi:10.21037/atm.2017.03.95 (2017).
- 21 Gámez-Valero, A. *et al.* Size-Exclusion Chromatography-based isolation minimally alters Extracellular Vesicles' characteristics compared to precipitating agents. *Sci Rep* **6**, 33641 (2016).
- 22 Kang, Y. T. *et al.* High-purity capture and release of circulating exosomes using an exosome-specific dual-patterned immunofiltration (ExoDIF) device. *Nanoscale* **9**, 13495-13505, doi:10.1039/c7nr04557c (2017).
- 23 Rekker, K. *et al.* Comparison of serum exosome isolation methods for microRNA profiling. *Clinical Biochemistry* **47**, 135-138, doi:https://doi.org/10.1016/j.clinbiochem.2013.10.020 (2014).
- 24 Stranska, R. *et al.* Comparison of membrane affinity-based method with size-exclusion chromatography for isolation of exosome-like vesicles from human plasma. *Journal of Translational Medicine* **16**, 1, doi:10.1186/s12967-017-1374-6 (2018).
- 25 Yang, F., Liao, X., Tian, Y. & Li, G. Exosome separation using microfluidic systems: size-based, immunoaffinity-based and dynamic methodologies. *Biotechnol J* **12**, doi:10.1002/biot.201600699 (2017).
- 26 Liga, A., Vliegthart, A. D., Oosthuyzen, W., Dear, J. W. & Kersaudy-Kerhoas, M. Exosome isolation: a microfluidic road-map. *Lab Chip* **15**, 2388-2394, doi:10.1039/c5lc00240k (2015).
- 27 Im, H. *et al.* Label-free detection and molecular profiling of exosomes with a nano-plasmonic sensor. *Nature Biotechnology* **32**, 490-495, doi:10.1038/nbt.2886 (2014).
- 28 Lin, J. *et al.* Exosomes: novel biomarkers for clinical diagnosis. *TheScientificWorldJournal* **2015**, 657086-657086, doi:10.1155/2015/657086 (2015).
- 29 Jiang, L. *et al.* EpCAM-dependent extracellular vesicles from intestinal epithelial cells maintain intestinal tract immune balance. *Nature Communications* **7**, 13045, doi:10.1038/ncomms13045 (2016).
- 30 Reategui, E. *et al.* Engineered nanointerfaces for microfluidic isolation and molecular profiling of tumor-specific extracellular vesicles. *Nat Commun* **9**, 175, doi:10.1038/s41467-017-02261-1 (2018).
- 31 Abramowicz, A., Widlak, P. & Pietrowska, M. Proteomic analysis of exosomal cargo: the challenge of high purity vesicle isolation. *Mol Biosyst* **12**, 1407-1419, doi:10.1039/c6mb00082g (2016).

- 32 Ziaei, P., Berkman, C. E. & Norton, M. G. Review: Isolation and Detection of Tumor-Derived Extracellular Vesicles. *ACS Applied Nano Materials* **1**, 2004-2020, doi:10.1021/acsanm.8b00267 (2018).
- 33 Gurunathan, S., Kang, M.-H., Jeyaraj, M., Qasim, M. & Kim, J.-H. Review of the Isolation, Characterization, Biological Function, and Multifarious Therapeutic Approaches of Exosomes. *Cells* **8**, 307, doi:10.3390/cells8040307 (2019).
- 34 ATCC. Cell Culture Guides. *ATCC online* (2016).
- 35 Zhou, H. *et al.* Collection, storage, preservation, and normalization of human urinary exosomes for biomarker discovery. *Kidney Int* **69**, 1471-1476 (2006).
- 36 Pisitkun, T., Shen, R. F. & Knepper, M. A. Identification and proteomic profiling of exosomes in human urine. *Proc Natl Acad Sci U S A* **101**, 13368-13373 (2004).
- 37 Schwartz, L. S., K. Introduction to Tangential Flow Filtration for Laboratory and Process Development Applications. *PALL Corp.* (2014).
- 38 Théry, C. *et al.* Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *Journal of Extracellular Vesicles* **7**, 1535750, doi:10.1080/20013078.2018.1535750 (2018).
- 39 Lee, M., Ban, J.-J., Im, W. & Kim, M. Influence of storage condition on exosome recovery. *Biotechnology and Bioprocess Engineering* **21**, 299-304, doi:10.1007/s12257-015-0781-x (2016).
- 40 Laboratories, S. MicroKros and MidiKros Hollow Fiber membranes for Tangential Laboratory Separations.
- 41 Soung, Y. H., Ford, S., Zhang, V. & Chung, J. Exosomes in Cancer Diagnostics. *Cancers (Basel)* **9** (2017).
- 42 Xu, R., Greening, D. W., Zhu, H. J., Takahashi, N. & Simpson, R. J. Extracellular vesicle isolation and characterization: toward clinical application. *J Clin Invest* **126**, 1152-1162 (2016).
- 43 Guo, L. & Guo, N. Exosomes: Potent regulators of tumor malignancy and potential bio-tools in clinical application. *Critical Reviews in Oncology/Hematology* **95**, 346-358, doi:https://doi.org/10.1016/j.critrevonc.2015.04.002 (2015).
- 44 Onitilo, A. A., Engel, J. M., Greenlee, R. T. & Mukesh, B. N. Breast cancer subtypes based on ER/PR and Her2 expression: comparison of clinicopathologic features and survival. *Clin Med Res* **7**, 4-13 (2009).
- 45 Halvaei, S. *et al.* Exosomes in Cancer Liquid Biopsy: A Focus on Breast Cancer. *Mol Ther Nucleic Acids* **10**, 131-141 (2018).
- 46 Shimomura, A. *et al.* Novel combination of serum microRNA for detecting breast cancer in the early stage. *Cancer Sci* **107**, 326-334 (2016).
- 47 Lässer, C. Exosomes in diagnostic and therapeutic applications: biomarker, vaccine and RNA interference delivery vehicle. *Expert Opin Biol Ther* **15**, 103-117, doi:10.1517/14712598.2015.977250 (2015).
- 48 Krishnamurti, U. & Silverman, J. F. HER2 in Breast Cancer: A Review and Update. *Advances in Anatomic Pathology* **21** (2014).
- 49 Li, S., Yang, X., Yang, J., Zhen, J. & Zhang, D. Serum microRNA-21 as a potential diagnostic biomarker for breast cancer: a systematic review and meta-analysis. *Clin Exp Med* **16**, 29-35, doi:10.1007/s10238-014-0332-3 (2016).

- 50 Sims, A. H., Howell, A., Howell, S. J. & Clarke, R. B. Origins of breast cancer subtypes and therapeutic implications. *Nature Clinical Practice Oncology* **4**, 516-525, doi:10.1038/ncponc0908 (2007).
- 51 Kalluri, R. The biology and function of exosomes in cancer. *The Journal of Clinical Investigation* **126**, 1208-1215, doi:10.1172/JCI81135 (2016).
- 52 De Toro, J., Herschlik, L., Waldner, C. & Mongini, C. Emerging roles of exosomes in normal and pathological conditions: new insights for diagnosis and therapeutic applications. *Front Immunol* **6**, 203, doi:10.3389/fimmu.2015.00203 (2015).
- 53 Sheridan, C. Exosome cancer diagnostic reaches market. *Nature Biotechnology* **34**, 359-360, doi:10.1038/nbt0416-359 (2016).
- 54 Lobb, R. J. *et al.* Optimized exosome isolation protocol for cell culture supernatant and human plasma. *Journal of extracellular vesicles* **4**, 27031-27031, doi:10.3402/jev.v4.27031 (2015).
- 55 Grigor'eva, A. E. *et al.* Contamination of exosome preparations, isolated from biological fluids. *Biochemistry (Moscow), Supplement Series B: Biomedical Chemistry* **11**, 265-271, doi:10.1134/S1990750817030040 (2017).
- 56 Cheruvanky, A. *et al.* Rapid isolation of urinary exosomal biomarkers using a nanomembrane ultrafiltration concentrator. *Am J Physiol Renal Physiol* **292**, F1657-1661, doi:10.1152/ajprenal.00434.2006 (2007).
- 57 Langdon, S. P. Cancer Cell Culture. (2003).
- 58 Yang, C. Y. *et al.* Diagnostic Accuracy of Urine Protein/Creatinine Ratio Is Influenced by Urine Concentration. *PLoS One* **10**, e0137460, doi:10.1371/journal.pone.0137460 (2015).
- 59 Busher., J. T. Serum Albumin and Globulin. *Clinical Methods: The History, Physical, and Laboratory Examinations* (1990).
- 60 Yoshioka, Y. *et al.* Comparative marker analysis of extracellular vesicles in different human cancer types. *Journal of extracellular vesicles* **2**, 10.3402/jev.v3402i3400.20424, doi:10.3402/jev.v2i0.20424 (2013).
- 61 Kowal, J. *et al.* Proteomic comparison defines novel markers to characterize heterogeneous populations of extracellular vesicle subtypes. *Proceedings of the National Academy of Sciences* **113**, E968, doi:10.1073/pnas.1521230113 (2016).
- 62 Willms, E. *et al.* Cells release subpopulations of exosomes with distinct molecular and biological properties. *Scientific Reports* **6**, 22519, doi:10.1038/srep22519 (2016).
- 63 Andreu, Z. & Yáñez-Mó, M. Tetraspanins in extracellular vesicle formation and function. *Frontiers in immunology* **5**, 442-442, doi:10.3389/fimmu.2014.00442 (2014).
- 64 Rosa-Fernandes, L., Rocha, V. B., Carregari, V. C., Urbani, A. & Palmisano, G. A Perspective on Extracellular Vesicles Proteomics. *Frontiers in Chemistry* **5**, 102 (2017).
- 65 Brennan, K. *et al.* A comparison of methods for the isolation and separation of extracellular vesicles from protein and lipid particles in human serum. *Scientific Reports* **10**, 1039, doi:10.1038/s41598-020-57497-7 (2020).
- 66 Menard, J. A., Cerezo-Magaña, M. & Belting, M. Functional role of extracellular vesicles and lipoproteins in the tumour microenvironment. *Philosophical*

- Transactions of the Royal Society B: Biological Sciences* **373**, 20160480, doi:10.1098/rstb.2016.0480 (2018).
- 67 Alix-Panabières, C. & Pantel, K. Clinical prospects of liquid biopsies. *Nature Biomedical Engineering* **1**, 0065, doi:10.1038/s41551-017-0065 (2017).
- 68 Chernyshev, V. S. et al. Size and shape characterization of hydrated and desiccated exosomes. *Anal Bioanal Chem* **407**, 3285-3301, doi:10.1007/s00216-015-8535-3 (2015).
- 69 Skliar, M. & Chernyshev, V. S. Imaging of Extracellular Vesicles by Atomic Force Microscopy. *J Vis Exp*, doi:10.3791/59254 (2019).
- 70 Smith, E. R., Zurakowski, D., Saad, A., Scott, R. M. & Moses, M. A. Urinary biomarkers predict brain tumor presence and response to therapy. *Clin Cancer Res* **14**, 2378-2386, doi:10.1158/1078-0432.Ccr-07-1253 (2008).
- 71 Su, Y. Y. et al. Upregulated expression of serum exosomal miR-375 and miR-1307 enhance the diagnostic power of CA125 for ovarian cancer. *J Ovarian Res* **12**, 6, doi:10.1186/s13048-018-0477-x (2019).
- 72 Macías, M. et al. Comparison of six commercial serum exosome isolation methods suitable for clinical laboratories. Effect in cytokine analysis. *Clin Chem Lab Med* **57**, 1539-1545, doi:10.1515/ccm-2018-1297 (2019).
- 73 Dear, J. W., Street, J. M. & Bailey, M. A. Urinary exosomes: a reservoir for biomarker discovery and potential mediators of intrarenal signalling. *Proteomics* **13**, 1572-1580, doi:10.1002/pmic.201200285 (2013).
- 74 Gu, C. Y. et al. Clinical significance of urine prostatic exosomal protein in the diagnosis of prostate cancer. *Am J Cancer Res* **9**, 1074-1078 (2019).
- 75 Musante, L. et al. Proteases and protease inhibitors of urinary extracellular vesicles in diabetic nephropathy. *J Diabetes Res* **2015**, 289734, doi:10.1155/2015/289734 (2015).
- 76 Buschmann, D. et al. Evaluation of serum extracellular vesicle isolation methods for profiling miRNAs by next-generation sequencing. **7**, 1481321 (2018).
- 77 Helwa, I. et al. A Comparative Study of Serum Exosome Isolation Using Differential Ultracentrifugation and Three Commercial Reagents. *PLOS ONE* **12**, e0170628, doi:10.1371/journal.pone.0170628 (2017).
- 78 Patel, G. K. et al. Comparative analysis of exosome isolation methods using culture supernatant for optimum yield, purity and downstream applications. *Scientific Reports* **9**, 5335, doi:10.1038/s41598-019-41800-2 (2019).
- 79 Kırbaş, O. K. et al. Optimized Isolation of Extracellular Vesicles From Various Organic Sources Using Aqueous Two-Phase System. *Scientific Reports* **9**, 19159, doi:10.1038/s41598-019-55477-0 (2019).
- 80 Davis, C. N. et al. The importance of extracellular vesicle purification for downstream analysis: A comparison of differential centrifugation and size exclusion chromatography for helminth pathogens. *PLOS Neglected Tropical Diseases* **13**, e0007191, doi:10.1371/journal.pntd.0007191 (2019).
- 81 Liu, C. et al. Field-Free Isolation of Exosomes from Extracellular Vesicles by Microfluidic Viscoelastic Flows. *ACS Nano* **11**, 6968-6976, doi:10.1021/acsnano.7b02277 (2017).

- 82 Tang, Y.-T. *et al.* Comparison of isolation methods of exosomes and exosomal RNA from cell culture medium and serum. *International journal of molecular medicine* **40**, 834-844, doi:10.3892/ijmm.2017.3080 (2017).
- 83 Syn, N. L., Wang, L., Chow, E. K., Lim, C. T. & Goh, B. C. Exosomes in Cancer Nanomedicine and Immunotherapy: Prospects and Challenges. *Trends Biotechnol* **35**, 665-676, doi:10.1016/j.tibtech.2017.03.004 (2017).
- 84 Jeyaram, A. & Jay, S. M. Preservation and Storage Stability of Extracellular Vesicles for Therapeutic Applications. *The AAPS journal* **20**, 1-1, doi:10.1208/s12248-017-0160-y (2017).
- 85 Cheng, Y., Zeng, Q., Han, Q. & Xia, W. Effect of pH, temperature and freezing-thawing on quantity changes and cellular uptake of exosomes. *Protein & Cell* **10**, 295-299, doi:10.1007/s13238-018-0529-4 (2019).
- 86 Maroto, R. *et al.* Effects of storage temperature on airway exosome integrity for diagnostic and functional analyses. *Journal of Extracellular Vesicles* **6**, 1359478, doi:10.1080/20013078.2017.1359478 (2017).
- 87 Ramirez, M. I. *et al.* Technical challenges of working with extracellular vesicles. *Nanoscale* **10**, 881-906, doi:10.1039/C7NR08360B (2018).
- 88 Gonzalez-Villasana, V. *et al.* Presence of Circulating miR-145, miR-155, and miR-382 in Exosomes Isolated from Serum of Breast Cancer Patients and Healthy Donors. *Disease Markers* **2019**, 6852917, doi:10.1155/2019/6852917 (2019).
- 89 Moloney, B. M. *et al.* Investigating the Potential and Pitfalls of EV-Encapsulated MicroRNAs as Circulating Biomarkers of Breast Cancer. *Cells* **9**, doi:10.3390/cells9010141 (2020).
- 90 Jung, H. H., Kim, J.-Y., Lim, J. E. & Im, Y.-H. Cytokine profiling in serum-derived exosomes isolated by different methods. *Scientific Reports* **10**, 14069, doi:10.1038/s41598-020-70584-z (2020).
- 91 Vardaki, I. *et al.* Periostin is identified as a putative metastatic marker in breast cancer-derived exosomes. **7** (2016).
- 92 Johnsen, K. B., Gudbergsson, J. M., Andresen, T. L. & Simonsen, J. B. What is the blood concentration of extracellular vesicles? Implications for the use of extracellular vesicles as blood-borne biomarkers of cancer. *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer* **1871**, 109-116, doi:https://doi.org/10.1016/j.bbcan.2018.11.006 (2019).
- 93 Buzás, E. I., Gardiner, C., Lee, C. & Smith, Z. J. Single particle analysis: Methods for detection of platelet extracellular vesicles in suspension (excluding flow cytometry). *Platelets* **28**, 249-255, doi:10.1080/09537104.2016.1260704 (2017).
- 94 Khatun, Z., Bhat, A., Sharma, S. & Sharma, A. Elucidating diversity of exosomes: biophysical and molecular characterization methods. *Nanomedicine* **11**, 2359-2377, doi:10.2217/nnm-2016-0192 (2016).
- 95 Aguilera-Rojas, M., Badewien-Rentzsch, B., Plendl, J., Kohn, B. & Einspanier, R. Exploration of serum- and cell culture-derived exosomes from dogs. *BMC Veterinary Research* **14**, 179, doi:10.1186/s12917-018-1509-x (2018).
- 96 Bachurski, D. *et al.* Extracellular vesicle measurements with nanoparticle tracking analysis – An accuracy and repeatability comparison between NanoSight NS300 and ZetaView. *Journal of Extracellular Vesicles* **8**, 1596016, doi:10.1080/20013078.2019.1596016 (2019).

- 97 German, J. B., Smilowitz, J. T. & Zivkovic, A. M. Lipoproteins: When size really matters. *Current opinion in colloid & interface science* **11**, 171-183, doi:10.1016/j.cocis.2005.11.006 (2006).
- 98 Maas, S. L. N. *et al.* Possibilities and limitations of current technologies for quantification of biological extracellular vesicles and synthetic mimics. *Journal of Controlled Release* **200**, 87-96, doi:<https://doi.org/10.1016/j.jconrel.2014.12.041> (2015).
- 99 Hartjes, T. A., Mytnyk, S., Jenster, G. W., van Steijn, V. & van Royen, M. E. Extracellular Vesicle Quantification and Characterization: Common Methods and Emerging Approaches. *Bioengineering* **6**, doi:10.3390/bioengineering6010007 (2019).
- 100 Koritzinsky, E. H., Street, J. M., Star, R. A. & Yuen, P. S. T. Quantification of Exosomes. *Journal of cellular physiology* **232**, 1587-1590, doi:10.1002/jcp.25387 (2017).
- 101 SCIENCE, I. Measure Nanoparticles with Unparalleled Precision. <https://izon.com/nanoparticlemeasurement/>.
- 102 Yi, Y. W. *et al.* Advances in Analysis of Biodistribution of Exosomes by Molecular Imaging. *International Journal of Molecular Sciences* **21**, doi:10.3390/ijms21020665 (2020).
- 103 Linares, R., Tan, S., Gounou, C., Arraud, N. & Brisson, A. R. High-speed centrifugation induces aggregation of extracellular vesicles. *Journal of Extracellular Vesicles* **4**, 29509, doi:10.3402/jev.v4.29509 (2015).
- 104 Yamashita, T., Takahashi, Y., Nishikawa, M. & Takakura, Y. Effect of exosome isolation methods on physicochemical properties of exosomes and clearance of exosomes from the blood circulation. *European Journal of Pharmaceutics and Biopharmaceutics* **98**, 1-8, doi:<https://doi.org/10.1016/j.ejpb.2015.10.017> (2016).
- 105 Bobrie, A., Colombo, M., Krumeich, S., Raposo, G. & Théry, C. Diverse subpopulations of vesicles secreted by different intracellular mechanisms are present in exosome preparations obtained by differential ultracentrifugation. *Journal of Extracellular Vesicles* **1**, 18397, doi:10.3402/jev.v1i0.18397 (2012).
- 106 Lane, R. *et al.* Cell-derived extracellular vesicles can be used as a biomarker reservoir for glioblastoma tumor subtyping. *Communications Biology* **2**, 315, doi:10.1038/s42003-019-0560-x (2019).
- 107 Dobra, G. *et al.* Small Extracellular Vesicles Isolated from Serum May Serve as Signal-Enhancers for the Monitoring of CNS Tumors. *International Journal of Molecular Sciences* **21**, doi:10.3390/ijms21155359 (2020).
- 108 Simon, T., Kumaran, A., Veselu, D.-F. & Giamas, G. Three Method-Combination Protocol for Improving Purity of Extracellular Vesicles. *International Journal of Molecular Sciences* **21**, doi:10.3390/ijms21093071 (2020).
- 109 Nair, S. & Salomon, C. Extracellular vesicles as critical mediators of maternal-fetal communication during pregnancy and their potential role in maternal metabolism. *Placenta*, doi:<https://doi.org/10.1016/j.placenta.2020.06.011> (2020).
- 110 Chiam, K. *et al.* Serum outperforms plasma in small extracellular vesicle microRNA biomarker studies of adenocarcinoma of the esophagus. *World journal of gastroenterology* **26**, 2570-2583, doi:10.3748/wjg.v26.i20.2570 (2020).

- 111 Yoo, C. E. *et al.* A direct extraction method for microRNAs from exosomes captured by immunoaffinity beads. *Analytical Biochemistry* **431**, 96-98, doi:<https://doi.org/10.1016/j.ab.2012.09.008> (2012).
- 112 Sharma, P. *et al.* Immunoaffinity-based isolation of melanoma cell-derived exosomes from plasma of patients with melanoma. **7**, 1435138 (2018).
- 113 Simpson, R. J., Lim, J. W. E., Moritz, R. L. & Mathivanan, S. Exosomes: proteomic insights and diagnostic potential. *Expert Review of Proteomics* **6**, 267-283, doi:10.1586/epr.09.17 (2009).
- 114 Schwarzenbach, H. Clinical Relevance of Circulating, Cell-Free and Exosomal microRNAs in Plasma and Serum of Breast Cancer Patients. *Oncol Res Treat* **40**, 423-429, doi:10.1159/000478019 (2017).
- 115 Hannafon, B. N. *et al.* Plasma exosome microRNAs are indicative of breast cancer. *Breast Cancer Research* **18**, 90, doi:10.1186/s13058-016-0753-x (2016).
- 116 Ricardo, S. *et al.* Breast cancer stem cell markers CD44, CD24 and ALDH1: expression distribution within intrinsic molecular subtype. *J Clin Pathol* **64**, 937-946, doi:10.1136/jcp.2011.090456 (2011).
- 117 Rupp, A. K. *et al.* Loss of EpCAM expression in breast cancer derived serum exosomes: role of proteolytic cleavage. *Gynecol Oncol* **122**, 437-446, doi:10.1016/j.ygyno.2011.04.035 (2011).

Figure Captions

Figure 1. The Immunomagnetic sequential ultrafiltration (iSUF) platform. A) Schematic representation of iSUF stages 1, 2, 3, including tangential flow filtration (TFF) purification, centrifugal unit enrichment, and tEV immunoaffinity isolation, respectively. In stage 1, biofluids were processed using a 500 kDa TFF filter, EVs were retained in the retentate and enriched in 7 mL while free proteins and nucleic acids permeated through the TFF filter. Then the PBS valve was opened to start TFF diafiltration until removing up to 99 % of free proteins. Finally, EVs were recovered in 2 mL of PBS after flushing the TFF system with air. In stage 2, EVs were centrifuged using a 10 kDa centrifugal unit at 3,000 x g and enriched in 100 μ L. In stage 3, tEVs were captured using antibodies

immobilized to streptavidin-coated magnetic beads (i.e., EGFR) and subsequently pulled out with a magnet. B) iSUF stage 1 purification performance for different biofluids. Cell supernatant (CCM), urine, and serum took 80, 100, and 120 min to obtain up to 99 % efficiency of free protein and nucleic acid removal.

Figure 2. iSUF enrichment performance and characterization of EVs separated from CCM, serum and urine using iSUF. A) Enrichment factor (EF) for EVs present in CCM, serum, and urine after iSUF ($n = 5$ for each biofluid; $p > 0.05$) (mean $\pm$ STD). Enrichment factors (EFs) were calculated as the ratio of EV concentration in biofluids present before and after iSUF processing. B) EFs for different size ranges of EVs for CCM ($n = 3$; $p >$

0.05) (mean $\pm$ STD). C) qNano measurements of the size distribution and concentration of EVs in the original CCM, after iSUF processing, and after UC processing. The EV concentration after iSUF was significantly higher than in the original CCM and after UC processing ($n = 3$; $p < 0.05$) (mean $\pm$ STD). D) qNano measurements of the size distribution and concentration of EVs in serum and urine after iSUF processing. E) AFM and SEM images of EVs from U251 CCM after iSUF processing and UC processing. Images were also obtained for EVs in serum and urine after iSUF processing. F) Transmission-EM images of EVs present in CCM after iSUF processing.

Figure 3. Comparison of size distribution and purity of EVs isolated with different platforms. A) Size distribution of EVs isolated from serum using different platforms. iSUF demonstrated the highest EV concentration within the 70 nm – 1 μ m size range. B) Left. Mean size boxplot for small EVs (sEVs). sEVs isolated using qEV and TEI demonstrated no differences with iSUF or UC, UC-sEVs showed larger mean size than iSUF-sEVs ($n = 5$; $p < 0.05$) (mean $\pm$ STD). Right. Mean size boxplot for medium/large EVs (m/l EVs). UC-m/l EVs were larger than iSUF, qEV and TEI m/l EVs ($n = 5$; $p < 0.05$) (mean $\pm$ STD). C) Concentration and purity of isolated EVs using different platforms. Black boxplots were the absolute concentration of EVs isolated from 0.5 mL of serum (left-axis); red dots corresponding to the right y-axis were purities defined as the EV concentration divided by the remaining free protein concentration. iSUF isolated EVs from serum most efficiently

with high purity ($n = 5$; $p < 0.05$) (mean $\pm$ STD), qEV yielded pure EVs, but at a lower concentration, TEI isolated more EVs but at relatively low purity, and UC recovered EVs with the lowest concentration and purity.

Figure 4. Characterization of protein and RNA content of EVs isolated from different biofluids. A) Quantity of total RNA extracted from CCM-EVs using UC and iSUF (n = 6; p < 0.05). B) Comparison of proteins extracted from CCM-EVs using UC and iSUF (n = 6; p < 0.05). C) Quantification of total RNA extracted from serum and urine EVs isolated by iSUF. D) Quantification of EV proteins isolated from serum and urine EVs isolated by iSUF. E) Western blot analysis of CD63 and CD9 expression in original CCM, CCM-EVs processed using iSUF and UC, original serum and urine, serum and urine-EVs processed

using iSUF. The expression level of CD63 and CD9 was calculated using an equal mass (40 µg) of protein lysates from samples. F). Total RNA of EVs isolated using different EV isolation methods. Black boxplots were the absolute amount of RNA isolated from 0.5 mL of serum. iSUF isolated the most RNA contents from serum EVs ($n = 5$; $p < 0.05$) (mean $\pm$ STD).

Figure 5. Characterization of stained tEVs isolated from U251-CCM and spiked in different biofluids. A) Upper. Fluorescence images of tEVs on magnetic beads using TIRF microscopy. Lower. Comparison of fluorescence intensities quantified using imaging flow cytometry. Stained U251-EVs isolated by iSUF was a positive control. tEVs isolated from serum and urine samples using iSUF demonstrated no significant difference ($n = 3$; $p > 0.05$), while UC isolated tEVs differed with iSUF and the positive control ($n = 3$; $p < 0.05$) (mean $\pm$ STD). B) Quantification of total RNA isolated from U251-EVs before and after bead extraction showed no differences ($n = 3$; $p > 0.05$). C) Total RNA extracted from serum samples using iSUF was significantly higher than using UC ($n = 3$; $p < 0.05$) (mean $\pm$ STD).

Figure 6. Characterization of tEVs isolated breast cancer (BC) patient serum using iSUF. A) Left. Characteristic fluorescence images of HER2, CD24, miR21, CD63, and CD9 expression on isolated EVs immobilized on magnetic beads from BC patients and

healthy donors. Right. Fluorescence intensity histograms of HER2, CD24, miR21, CD63, and CD9 of isolated EVs for BC patients (n = 10) and healthy donors (n = 14). B) Total fluorescence intensity quantification of HER2, CD24, miR21, CD63, and CD9 expression level on isolated EVs from BC patients and healthy donors. Patients demonstrated higher expression levels of HER2, CD24, and miR21 than healthy donors ($p < 0.05$). Patients and HDs demonstrated similar expression level of CD63 and CD9 ($p > 0.05$). P: patient; H: healthy donor.

Immunomagnetic Sequential Ultrafiltration (iSUF) Platform for Enrichment and Purification of Extracellular Vesicles from Biofluids

Jingjing Zhang,¹ Luong TH Nguyen,¹ Richard Hickey,¹ Nicole Walters,¹ Xinyu Wang,¹ Kwang Joo Kwak,¹ L. James Lee,¹ Andre F. Palmer,¹ Eduardo Reátegui^{1,2*}

¹ William G. Lowrie Department of Chemical and Biomolecular Engineering, The Ohio State University, Columbus OH 43210

² Comprehensive Cancer Center, The Ohio State University, Columbus, OH 43210

*corresponding author: reategui.8@osu.edu

Supplementary Information

Size distribution and concentration of EVs

A tunable resistive pulse sensing (TRPS) method (Izon Sciences, Boston, MA) was employed to quantify the size and concentration of EVs. 45 μ L of biofluid was pipetted into nanopore membranes (NP150, NP300, NP600, NP800, and NP1000), and then pressure (10 mbar) and voltage (0.38V, 0.32V, 0.26V, 0.18V and 0.12V) were applied. Every single EV causes a resistive pulse that can be used to calculate EV size and concentration. Polystyrene nanoparticles of different known sizes and concentrations were used for calibration.

Atomic force microscopy (AFM)

A clean mica substrate was vapor-phase coated with 3-aminopropyltriethoxysilane (APTES, Millipore Sigma, Burlington, MA) in a vacuum chamber and then dried overnight at 65 °C. Subsequently, 10 μ L of purified EVs were incubated on the surface for 30 min at RT. Unbound EVs were extensively rinsed with PBS and then with DI water. The samples were air-dried again before imaging using an AFM (Asylum Research MFP-3D-BIO AFM, Oxford Instruments, Abingdon, United Kingdom).

Scanning electron microscopy (SEM)

Clean coverslips were soaked in 0.25 mg/mL Zetag solution (BASF, Southfield, MI, USA) for 30 min, followed by overnight air drying at RT. Purified EVs were attached to the coated coverslip for 30 min at RT by physisorption. EVs were fixed in 2% glutaraldehyde (MilliporeSigma, Burlington, MA) and 0.1 M sodium cacodylate solution (Electron Microscopy Sciences, Hatfield, PA, USA) for 3h. After washing with 0.1 M sodium cacodylate solution, EVs were incubated in 1% osmium tetroxide (Electron Microscopy Sciences) and 0.1 M sodium cacodylate for 2h. The sample was

subsequently rinsed with 0.1 M sodium cacodylate solution before dehydration in increasing concentrations of ethanol (50, 70, 85, 95, and 100%, ThermoFisher Scientific) for 30 min each. Next, the samples were transferred to a CO₂ critical point dryer (tousimis, Rockville, MD, USA). Finally, the samples were coated with ~ 2 nm of gold using a sputtering machine (Leica EM ACE 600, Buffalo Grove, IL) and imaged using SEM (Apreo ii, FEI, Thermo Fisher Scientific).

Transmission electron microscopy (TEM)

3 µL EVs purified from CCM using iSUF were applied to a glow discharged lacey carbon-coated copper grid (400 mesh, Pacific Grid-Tech, San Francisco, CA) and flash-frozen in liquid ethane using an automated vitrification device (FEI Vitrobot Mark IV, FEI, Hillsboro, OR). The sample was then visualized in a Glacios™ Cryo-TEM (ThermoFisher Scientific).

Protein extraction and quantification

EV samples were lysed in RIPA buffer (Abcam) with the addition of Thermo Scientific™ Halt™ Protease and Phosphatase Inhibitor Cocktails for 15 min on ice. EV samples (with/without lysis) were then pipetted to a 96-well plate, and their protein concentrations were quantified using a Pierce™ Rapid Gold BCA Protein Assay kit (ThermoFisher Scientific). EV protein concentration was determined by subtracting the amount of free protein (without lysis) from the total (with lysis) in the purified EV sample.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE)

Proteins in the final product were denatured and reduced in the presence of NuPAGE Reducing Agent in NuPAGE LDS Sample Buffer at 95 °C for 10 min. The proteins were then separated in a mini gel tank (ThermoFisher Scientific) using NuPAGE 4-12% Bis-Tris Protein Gel in

NuPAGE MOPS SDS Running Buffer for 50 min at 200 V. After separation, the proteins were stained with Coomassie Brilliant Blue G-250 Dye.

Western blotting

After separation by SDS-PAGE, proteins were transferred onto a polyvinylidene fluoride (PVDF) membrane (ThermoFisher Scientific) and then blocked with 3% bovine serum albumin (BSA) and 0.05% Tween in PBS for 1 hr at RT. Primary antibodies against tetraspanin surface markers such as CD63, CD81, and CD9 were incubated with the EVs overnight at 4 °C (Santa Cruz Biotechnology, Inc Dallas, TX, USA). The next day, the PVDF membrane was incubated with an HRP conjugated secondary antibody for 1h at RT. Finally, the sample was incubated with SuperSignal™ West Femto Maximum Sensitivity Substrate for 5 min at RT before imaging using a C-digit blot scanner (LI-COR, Lincoln, NE, USA).

RNA quantification

Total RNA was extracted with QIAzol Lysis Reagent and then purified using a miRNeasy Mini kit according to the manufacturer's protocol (Qiagen, Germantown, MD, USA). After purification, the RNA concentration was quantified using a Qubit microRNA Assay Kit at excitation/emission wavelengths of 500/525 nm.

Supplementary Figure 1. EV recovery rate after TFF processing. Cell culture medium (n = 5), serum (n = 5), and urine (n = 5) were purified using TFF and enriched into a volume of 2 mL.

Supplementary Figure 2. Pressure in the TFF stage of iSUF when processing bovine serum albumin (BSA) solutions at different concentrations using a fixed flow rate of 35 mL/min. 15 mg/mL was the maximum protein concentration in the TFF system to maintain the system pressure below 10 psig.

Supplementary Figure 3. EV recovery rate and processing time of centrifugal units (iSUF, stage 2) with different MWCOs. 3 kDa, 10 kDa, and 30 kDa MWCO centrifugal units obtained over 99%, 95%, and 70% recovery rate, respectively. They took 60 min (3 kDa MWCO), 20 min (10 kDa MWCO), and 15 min (30 kDa MWCO) to spin down the sample to a final volume of 100 μ L.

Supplementary Figure 4. SDS-PAGE of 10% BSA solution before and after iSUF processing.

BSA was extensively removed after iSUF processing.

Supplementary Figure 5. SDS-PAGE of CCM before and after iSUF processing. BSA was extensively removed after iSUF processing.

Supplementary Figure 6. SDS-PAGE of serum samples before and after iSUF processing. HSA was extensively removed after iSUF processing.

Supplementary Figure 7. SDS-PAGE of urine samples before and after iSUF processing. HSA was extensively removed after iSUF processing.

Supplementary Figure 8. A). Repeatability evaluation of EV recovery rate for samples processed by iSUF.

Cell culture medium (n = 10), serum (n = 10), and urine (n = 10) were purified by iSUF and enriched into a final volume of 100 μ L. **B).** Repeatability evaluation of protein removal efficiency for samples processed by iSUF.

Supplementary Figure 9. Full-length western blots for cropped images in Fig. 4E.

Supplementary Figure 10. Comparison of EV concentration, protein, and RNA content in iSUF-urine and iSUF-serum samples from seven healthy donors. The EV concentration and RNA content showed comparable values between urine and serum ($n = 10$; $p > 0.05$), while protein content in urine was 10 times lower than in serum.

Supplementary Figure 11. Levels of expression of BC biomarkers in young and old healthy donor serum samples. For older HDs, the mean age was 58.7 ± 5.6 years (SD); for younger HDs, the mean age was 24.8 ± 2.1 years (SD). There was not a statistically significant difference in fluorescence signal for both cohorts ($p > 0.05$). **A)** Characteristic fluorescence images of HER2, CD24, miR21. **B)** Total fluorescence intensity quantification of HER2, CD24, miR21.

Supplementary Figure 12. Size distribution of EVs in a CCM sample processed by iSUF and measured by a TRPS method (qNano) using an NP100 and NP150 stretchable membrane.

Supplementary Figure 13. Particle count plot of EVs isolated by iSUF, TEI, qEV, and UC. The graph indicates an overall constant particle detection.

Supplementary Table 1. Comparison of 300 and 500 kDa TFF filter performance on CCM, urine, and serum.

TFF filter MWCO	CCM (50 mL)		Urine (100 mL)		Serum (0.5 mL)	
	Protein removal efficiency	Processing time	Protein removal efficiency	Processing time	Protein removal efficiency	Processing time
300 kDa	>78%	150 min	>80%	200 min	>75%	300 min
500 kDa	up to 99%	80 min	up to 99%	100 min	up to 99%	120 min

Supplementary Table 2. Protein concentration of the TFF enrichment step.

	U251 Supernatant	Urine	Serum
Initial protein concentration (mg/mL)	1.20 ± 0.13 (in 50 mL)	1.40 ± 0.50 (in 10~100 mL)	10.0 ± 2.0 (dilute 0.5 mL of serum to total volume of 7mL in PBS)
Protein concentration after TFF enrichment step (mg/mL)	2.30 ± 0.38	1.90 ± 0.31	Not applied

Supplementary Table 3. EV concentration, purity, and RNA content comparison among different EV purification platforms.

	Concentration (particles/mL)	Free protein (μ g/mL)	Purity (particles/ μ g)	RNA (ng)
qEV	8.19E10 $\pm$ 2.62E10	15.39 $\pm$ 4.06	5.94E09 $\pm$ 3.08E09	8.29 $\pm$ 7.58
UC	7.46E10 $\pm$ 2.25E10	11841.0 $\pm$ 2375.6	6.77E06 $\pm$ 3.30E06	5.52 $\pm$ 5.02
TEI	5.70E11 $\pm$ 2.79E11	67.2 $\pm$ 29.8	1.32E10 $\pm$ 1.43E10	15.23 $\pm$ 8.24
iSUF	4.15E12 $\pm$ 4.66E11	6.32 $\pm$ 4.28	1.32E12 $\pm$ 1.26E12	53.72 $\pm$ 41.02

Supplementary Table 4.

Breast cancer patient information

	P1	P2	P3	P4	P5
Concentration (sEVs/mL)	1.06E11	9.67E11	1.77E10	4.81E10	2.54E11
Mean size (nm)	124	130	127	139	120
Concentration (m/IEVs/mL)	9.87E9	1.01E11	9.37E9	1.21E10	9.08E10
Mean size (nm)	321	354	320	317	340
Age	66	55	81	60	78
Sex	F	F	F	F	F
Histology	Ductal carcinoma in site	Infiltrating duct carcinoma, NOS	Lobular carcinoma, NOS	Lobular carcinoma, NOS	Infiltrating duct carcinoma, NOS
	P6	P7	P8	P9	P10
Concentration (sEVs/mL)	5.78E10	1.27E11	3.19E10	9.13E10	3.02E9
Mean size (nm)	135	135	140	120	128
Concentration (m/IEVs/mL)	4.30E9	9.99E9	2.54E9	1.15E10	3.21E9
Mean size (nm)	331	334	348	350	317

Age	63	68	32	62	50
Sex	F	F	F	F	F
Histology	Carcinoma	Carcinoma	Infiltrating duct carcinoma, NOS	Adenocarci noma	Adenocarci noma

Healthy donor information

	H1	H2	H3	H4	H5
Concentration (sEVs/mL)	3.64E12	1.57E12	7.93E11	4.69E11	1.27E12
Mean size (nm)	128	134	145	119	110
Concentration (m/IEVs/mL)	8.87E10	1.21E10	8.93E9	2.41E10	8.82E10
Mean size (nm)	354	327	378	327	301
Age	25	24	21	24	27
Sex	F	F	F	F	F
	H6	H7	H8	H9	H10
Concentration (sEVs/mL)	2.49E12	3.85E11	4.67E12	2.37E11	9.81E10
Mean size (nm)	128	134	119	122	107

Concentration (m/IEVs/mL)	2.87E11	1.54E11	1.01E11	1.59E10	1.06E10
Mean size (nm)	324	354	320	317	305
Age	28	24	25	53	53
Sex	F	F	F	F	F
	H11	H12	H13	H14	
Concentration (sEVs/mL)	2.69E10	3.21E10	4.19E10	1.07E10	
Mean size (nm)	124	120	134	127	
Concentration (m/IEVs/mL)	8.54E9	3.17E9	2.88E9	9.46E8	
Mean size (nm)	328	317	322	300	
Age	58	59	61	68	
Sex	F	F	F	F	