Extreme Genomic Makeover: Evolutionary History of Maternally-transmitted Clam Symbionts

Short title:

Evolution of maternally-transmitted symbiont genomes

M Perez¹, C Breusing², B Angers¹, YJ Won³, and CR Young⁴

¹ Department of Biological Sciences, Université de Montréal, Montreal, Canada

² Graduate School of Oceanography, University of Rhode Island, Narragansett, USA

³ Division of EcoScience, Ewha Womans University, Seoul, South Korea

⁴ National Oceanography Centre, Southampton, UK

Classification:

Evolution, Genetics, Microbiology, Selection, Recombination, Symbiosis, Genomics, Vesicomidae

Abstract

Given their recent switch to a vertically-transmitted intracellular lifestyle, the chemosynthetic bacteria associated with deep-sea vesicomid clams are an excellent model system to study the processes underlying reductive genome evolution. In this study, we provide the first estimates of the relative contributions of drift, recombination and selection in shaping the ongoing reductive genome evolution in these symbionts. To do so, we compared the genomes of endosymbionts associated with 11 vesicomid clam species to that of closely related free-living bacteria and their respective hosts' mitochondria. Our investigation confirmed that neutral evolutionary processes were the dominant driver of reductive genome evolution in this group and highlighted the important role of horizontal gene transfer in mitigating genome erosion. Finally, a genome-wide screen for episodic positive selection across the symbiont phylogeny revealed the pervasive role of selective processes in maintaining symbiont functional integrity.

Introduction

The evolution of biological complexity includes many examples of symbiotic associations. For example, the early evolution of the eukaryotic cell involved multiple endosymbiotic events leading to mitochondria and plastids^{1,2}. More recent examples include associations of metazoans with intracellular bacteria³⁻⁶, including the well-studied associations of insects and *Buchnera* proteobacterial symbionts⁷. These associations have profound consequences for both host and symbiont, ranging from alterations of sex-ratio in insect hosts to providing nutrients that are otherwise unavailable in the host's habitat. Some intracellular symbionts are transmitted from parent to offspring of hosts through the germline (i.e. vertical transmission), while others are acquired from the environment every generation⁶. The mode of transmission strongly affects the evolution of the microbial partner in these symbioses, as the genomes of vertically transmitted symbionts all seem to follow the same process of reductive genome evolution (RGE) regardless of their phylogenetic origin, host, or habitat. Compared to their free-living counterparts, the genomes of host-restricted symbionts are smaller, contain fewer genes, and are enriched in AT^{8,9}. A prime example is the genomes of cellular organelles such as mitochondria and plastids which are extremely streamlined compared to their bacterial cousins¹⁰. Symbiont genome evolution is thought to follow two main stages¹¹. Following host restriction, symbionts undergo rapid genome erosion as they lose non-essential genes through pseudogenization and deletions^{12,13}. Then, symbionts enter a "stabilizing phase". At this point, their genomes are streamlined, redundant genes and functions are lost¹⁴, and the effective rate of deletion diminishes¹⁵. This process might be largely neutral due to the reduced effective population size of host-restricted taxa.

The pea aphid/*Buchnera* symbiosis and several other insect/bacteria models support the neutral hypothesis. Captured symbionts experience successive bottleneck events during their transmission that reduce their effective population size and increase genetic clonality. As a consequence, genetic drift increases relative to selection in these taxa¹⁶⁻¹⁸. Under these circumstances, elevated mutation load (i.e. the Muller's ratchet¹⁹) and genetic erosion might lead to the functional death of the symbiont lineage^{17,20-22} unless compensating mechanisms such as gene transfer to the host nucleus or compensatory mutations alleviate the genetic load. Likewise, deep-sea taxa exhibit evidence of nearly neutral processes affecting evolutionary rates due to reduced population sizes in vertically transmitted symbionts²³. Other metazoan/microbial symbioses highlight the importance of selection in shaping reductive genome evolution. For instance, symbiont traits that are beneficial for the host are likely to experience increased selective pressures, while selection may be relaxed on genes that are functionally redundant⁸. Red Queen dynamics are expected to occur in obligate symbioses to maintain the host-symbiont specificity and the functioning of cyto-nuclear interactions through speciation²⁰. Unfortunately, the role of positive selection has often been ignored in studies of symbiont genome evolution and broad screens for positive selection have almost never been performed.

The intracellular sulfur-oxidizing bacteria associated with deep-sea vesicomyid clams (Bivalvia: Vesicomyidae: Pliocardiinae) represent an ideal model to address the neutral and selective processes driving reductive genome evolution. The symbionts are found within the epithelial cells of their host's gills and provide them with chemosynthetically derived food. They are vertically transmitted to the next generation through the eggs^{24,25} and generally show co-speciation with their hosts

87 ^{26,27}. It is assumed that symbiont capture in these animals was a single event that,
88 based on fossil and molecular information, happened before their radiation about 45
89 Mya ²⁸, an acquisition that is much more recent than that of other well-studied models
90 such as the aphid/*Buchnera* (~ 200 Mya ²⁹) and nematode/*Wolbachia* (~100Mya ³⁰)
91 symbioses. Today, the hosts represent the most diverse group of deep-sea bivalves ³¹,
92 with 173 described species present in a variety of reducing habitats worldwide from
93 hydrocarbon seeps on continental margins to hydrothermal vents on mid-ocean ridges
94 ^{32–34}. A comparative study of the first two sequenced vesicomyid symbiont genomes
95 ^{35,36} indicated that they possessed intermediate genome sizes and level of AT
96 enrichment compared to other host-restricted symbionts ¹¹. The symbionts of deep-sea
97 vesicomyid clams group into two divergent clades: Clade I (associated with hosts of
98 the *gigas* group), and Clade II (associated with all other lineages of vesicomyid hosts)
99 ³⁷. The genomic characteristics of Clade I symbionts indicate that this group is in an
100 advanced state of reductive genome evolution compared to Clade II. However, in
101 contrast to the well-studied pea aphid/*Buchnera* association, which has been in a state
102 of stasis for 50 Myrs ³⁸, the evolutionary processes responsible for remodeling the
103 genomes of vertically transmitted symbionts appear to be still operating in the
104 vesicomyid clam symbiosis. Conspicuous bottlenecks during transmission ²⁵ and loss
105 of DNA repair genes in several lineages ³⁷ suggest that neutral processes and
106 mutational pressures are driving RGE in vesicomyid symbionts, although this
107 hypothesis has not been formally tested.

108
109 In this study, we aim to assess the relative contribution of neutral and selective
110 processes to genome evolution in the maternally transmitted symbionts of deep-sea
111 vesicomyid clams. Specifically, we test the hypotheses that genetic drift is the main
112 driver of RGE in these symbionts and that diversifying selection has shaped their
113 genome to maintain host-symbiont epistasis throughout the evolutionary history of the
114 symbiosis. To do so, we applied comparative methods to the symbiont genomes of 11
115 vesicomyid deep-sea clam taxa representative of the diversity of Clade I and Clade II,
116 the mitochondrial genomes of their respective hosts, and two of their close free-living
117 relatives: the environmentally acquired gill symbiont of the hydrothermal vent mussel
118 *Bathymodiolus thermophilus* and the free-living bacteria of the SUP05 group, which
119 are marine chemoautotrophic Gammaproteobacteria found in hypoxic waters ^{39,40}.

120
121

122 Results

123 *Host mitochondrial and symbiont phylogenies*

124 Host mitochondrial genomes from the lineages examined in this study possess
125 identical gene orders and contents as previously published mitochondrial genomes
126 ^{41,42}. The phylogeny constructed with mitochondrial genome data (Figure 1A) is
127 congruent with the known host phylogenetic relationships based on multilocus
128 sequence data and the *COI* phylogeny ³¹. Structural variation is, however, present. We
129 observe the previously described noncoding structural variation, hypothesized to be
130 the control region, between the *tRNA^{Trp}* or *tRNA^{His-2}* and *ND6* loci ⁴¹⁻⁴³ but we were
131 unable to resolve this region with the current sequence data. We also found the *COX2*
132 gene varies in length among taxa (range: 1005-1452bp). All protein-coding genes in
133 the mitochondrial genomes were screened for selection using the adaptive branch-site
134 random effects likelihood method. Interestingly, the *COX2* gene exhibited evidence
135 for episodic diversifying selection on multiple branches of the phylogeny.

136
137 Genome size and GC content for the 11 symbiont assemblies in our study varied from
138 1.02Mb to 1.59 Mb and 31% to 37% GC, respectively (Table 1). The number CDS
139 ranged from 939 in *Ca. V. okutanii* to 2210 in *Ca. R. phaseoliformis*. Following initial
140 nomenclature, the symbiont lineages are referred to by the previously erected genera
141 for this group, *Candidatus Vesicomysocius* for Clade I, and *Candidatus Ruthia* for
142 Clade II symbionts, followed by host species names ^{35,36,44}. This classification at the
143 genus level is coherent with both the phylogenetic definition based on 16S identity
144 (inter-genus identity < 95% ⁴⁵) and functional definition based on criteria of genetic
145 isolation ⁴⁶ (see Symbiont genome structure and recombination)

146
147 Examination of the mitochondrial and symbiont phylogenies (Figure 1) shows good
148 concordance for all lineages except one. The symbiont lineages of *Ca. V. diagonalis*
149 and *Ca. V. extenta* are nearly identical whereas their respective host mitochondrial
150 lineages are divergent. The donor lineage in this recent symbiont replacement appears
151 to be *A. diagonalis*. It is noteworthy that these clams were both collected from sites in
152 Monterey Canyon. Pairwise comparison of mitochondrial and symbiont genome-wide
153 synonymous divergence indicates faster evolutionary rates in the mitochondria
154 compared to the symbionts in almost every holobiont pair (Figure 2). Within the
155 symbionts, we detect signatures of elevated substitution rates on the branch leading to
156 Clade I: the symbiont pairs across the Clade I- Clade II bipartition have significantly
157 higher divergence than the others even when controlled for host divergence ($1 < dS_{mito}$
158 < 2).

159

160 *Symbiont genome structure and recombination*

161 Free living bacteria associated with *B. thermophilus* and *Ca. T. autotrophicus* shared
162 about 1 Mbp of their genomes with the clam symbionts. Permutation analysis of
163 locally collinear blocks (i.e. long fragments of aligned genomes) with GRIMM
164 (<http://grimm.ucsd.edu/cgi-bin/grimm.cgi>) showed that at least 18 inversion events
165 occurred between the genome of the *B. thermophilus* symbiont and that of the *Ca. R.*
166 *magnifica* reference. Fewer rearrangements (3 inversions) were observed between
167 SUP05 and *Ca. R. magnifica*.

168

Genome structure among the clam symbionts was also variable (Figure 1B). The previously reported *Ca. V. okutanii* genome³⁶ possesses one inversion compared to that of *Ca. R. magnifica*³⁵ but that of *Ca. V. okutanii*'s closest relative, *Ca. V. soyoae*, does not. The genomes of *Ca. R. pacifica* and *Ca. R. rectimargo* share a single inversion distinct to that of *Ca. V. okutanii*. Two other inversions were found in the *Ca. V. gigas* genome. Finally, read-mapping to the consensus assemblies for *Ca. R. phaseoliformis* and *Ca. R. southwardae* suggested the presence of intra-host structural variation in these symbionts.

Applying Bayesian concordance analysis to all core protein-coding genes, we detect a large amount of recombination among symbiont lineages, though recombination is not randomly distributed. We observe no recombination between members of Clade I and II, but recombination is occurring within these genera (Figure 1B). Strikingly, much less topological concordance was found in Clade II – more than 40 different topologies were necessary to fully represent the diversity of conflicting phylogenetic signals – compared to that of Clade I whose phylogeny was fully represented by 5 different trees. Within Clade I, conflict originates from the uncertainty of the position of *Ca. V. gigas*. Only 50% of the genes support its position in the phylogenetic tree issued from the concatenated core genome alignment (Figure 1B). Other well supported positions for this species are at the base of the clade (supported by 27% of genes) and closer to the group composed of *Ca. V. soyoae* and *Ca. V. okutanii* (supported by 20% of genes). Within Clade II, only the grouping of the sister species *Ca. R. rectimargo* and *Ca. R. pacifica* is supported by the topologies of all genes while the positions of other species have low support.

Gene conservation across symbionts and free-living bacteria

Genes of free-living and horizontally-transmitted bacteria missing in vesicomylid symbionts

The genomes of the free-living bacteria contained many large (> 5kb) contiguous sections that were not found in the symbionts. These genomic islands were mostly composed of unannotated genes and mobile elements (transposases, integrases, prophage genes) (Table S1). We found more selfish genetic elements in the genome of the *Bathymodiolus* symbiont than in that of SUP05. The genomic islands found in the two genomes also encoded several gene clusters of particular functional interest described below.

Unsurprisingly for a bacterium living in a metal-rich hydrothermal environment, the *B. thermophilus* symbiont genome possesses genes for resistance against heavy-metal toxicity such as a multi-copper oxidase (*mmcO*), a copper ion exporting ATPase (*copB*), cobalt-zinc-cadmium resistance proteins (*czcD* and *czcCBA*), and a chromate transport protein (*chrA*). The genomic islands of the mussel symbiont also carried full operons for three different defense systems; a type I restriction and modification system (*hsdRMS*), a CRISPR-Cas type II system (*cas9*, *cas1*, *cas2*, *cas4*), and a type II toxin-antitoxin system (*vapCB*). Finally, this genome possesses a 23kb hydrogenase operon that has 83% and 82% identity to that of the symbionts of *Bathymodiolus septemdierum*⁴⁷ and *B. puteoserpentis*⁴⁸, respectively. The representative SUP05 genome contained a 21kb motility locus, comprising a type IV pilus biogenesis operon (*pilA*, *pilB*, *pilC*, *pilT*, *pilQ*, *pilY1*), and a toxin-antitoxin locus (*higAB*), that was not found in the other genomes. Furthermore, this genome possessed two

additional smaller genomic islands (6kb and 15 kb) encoding a nitric oxide reductase (*norCBQD*), and a periplasmic nitrate reductases (*napAB*), respectively, which clustered with sulfur covalently binding protein genes (*soxYZ*).

Gene content in vesicomymid symbionts

The symbionts of Clade I and Clade II possessed essentially a subset of the genes found in the free-living lineages. Indeed, sequence-based comparisons of free-living lineages to the symbionts revealed that many genes present only within the symbiont lineages are hypothetical genes with unknown function resulting from the degeneration of ancestral genes, as indicated by premature stop codons, frameshifts, and loss of neighboring genes (Table S1). These pseudogenes were more prevalent in the genomes of Clade I than Clade II symbionts. In many instances, homologous regions within the Clade I symbiont genomes were instead characterized by large deletions. In general, patterns of gene decay were more variable within Clade II than Clade I. Genes were overall more conserved within *Ca. R. southwardae*, *Ca. R. phaseoliformis* and *Ca. R. pliocardia* than in other lineages. Among the *Ca. Ruthia* symbionts, gene degeneration was most pronounced in *Ca. R. magnifica*, which possessed a conservation pattern closer to that of the Clade I lineages (Figure S1B).

Genome-wide pattern of relaxed selection

Codon usage bias was reduced in the symbiont lineages compared to their free-living relatives. Furthermore, symbionts in Clade I showed reduced bias and variance compared to Clade II (Figure 3A). The CDC values of core protein-coding genes were significantly correlated between lineage pairs both at the clade and species level (Pearson's test p-value <0.001; Figure 3B, Table S2), suggesting that the reduction in codon usage bias in the vertically transmitted symbionts result from a genome-wide reduction of the efficacy of purifying selection.

RELAX analysis revealed intensified selection in the vesicomymid symbionts compared to free-living bacteria for less than 5% of the core orthologous genes, while relaxed selection was detected in more than half of the core gene set (Figure 3C, Table S3). The magnitude of relaxation ($k < 1$) was in the range of that observed in insect endosymbionts⁴⁹ but was not correlated to codon bias. Genes exhibiting intensified and relaxed selection represented a multitude of metabolic functions, but genes under relaxed selection were enriched in the protein metabolism, nucleoside and nucleotides, and DNA metabolism categories while genes under intensifying selection were more likely to be associated with respiration, cell wall and capsule, and sulfur metabolism. However, we did not find increased relaxation in the symbionts of Clade I compared to Clade II. Indeed, fewer genes exhibited significant change in selection pressure (intensified or relaxed) between these groups than between symbionts and free-living bacteria, and about the same proportion of genes under relaxed and intensified selection was found in both clades.

Genome-wide screen for positive selection

The symbiont genes that passed the inclusion criteria to be screened for selection (see methods) included 652 loci. The application of the adaptive BS-REL method yielded 223 genes with significant evidence for episodic diversifying selection along branches in the phylogeny. Selection is distributed throughout the evolutionary history of the group (Figure S1A, and Table S4) with most selection occurring on the branches discriminating free-living bacteria, Clade I, and Clade II (branches a, b, and c in

Figure 4), as well as within the *B. thermophilus* symbiont and SUP05 lineage (43 and 37 genes, respectively). Eighty-five percent of the loci that exhibited unequivocal evidence of selection was assigned to SEED categories (Figure 4, Table S5). Within each clade and along each of the main branches, these selected loci were not equally represented amongst cellular functions of the core genome (hypergeometric tests p-values < 0.001). Genes in overrepresented functional categories are presented in Table 2. The complete list of selected genes is available in Table S4.

Selection within free-living bacteria

Amongst the free-living lineages, a larger than expected number of genes associated with protein metabolism, respiration, and sulfur metabolism were under selection (Fisher tests p-value < 0.05). These included genes involved in ribosome assembly, t-RNA biogenesis, protein folding, oxidative phosphorylation, sulfur oxidation, and dissimilatory sulfate reduction. On the bipartition between the free-living and symbiont groups, additional genes associated with protein metabolism were positively selected, including ribosomal protein genes, and the t-RNA ligase genes.

Selection within symbionts

Many genes coding for chaperones, ribosomal proteins, and t-RNA ligases were under selection within the symbiont phylogeny. In addition, we found evidence for selection in metabolic genes that are central to the chemosynthetic role of these symbionts. Several genes involved in sulphur metabolism (i.e. *dsrA*, *dsrP*, *soxB*, *cobB-cbiA/dsrN*) and electron donating/accepting reactions were under selection. Two genes involved in ammonia assimilation (*gltB*, and *glnD*) also exhibited evidence of selection within both symbiont clades. Within Clade II and along the branch partitioning this group, there was an over-representation of selected genes involved in *de novo* purine and pyrimidine biosynthesis, carbon fixation, and DNA recombination and repair. Selection within Clade I favored additional genes broadly associated with DNA metabolism. Notably, 60 genes showed evidence for positive selection in multiple branches of the phylogeny, including 44 genes within the symbiont phylogeny. These genes were mostly associated with protein metabolism.

Discussion

Reductive genome evolution is still ongoing in the clam symbionts and is driven by neutral processes

Comparative analyses of the first two reference genomes of vesicomid clam endosymbionts revealed variation in genome structure, genome characteristics, and genome composition between distantly-related symbiont species¹¹ suggesting that RGE might still be ongoing in this group. Our results confirm these early findings and reveal additional genomic variation among the deeply diverging lineages. These findings expand the ranges of genome size, genome content and GC% considerably.

As in other models of recently acquired bacteria^{22,50}, gene content differed greatly between vesicomid symbiont genomes indicating that the different lineages are independently losing genes. The presence of structural variation and putative pseudogenes (Figure S2) within the vesicomid symbiont genomes suggest that these symbionts have not yet reached a stable streamlined state as those of the *Buchnera* or *Paulinella* symbionts^{15,38}. Comparing the clam symbionts to their free-living relatives revealed reduced GC%, a reduction in codon usage bias, pseudogenization, and

evidence for reduced purifying selection in the vast majority of genes. Taken together, these observations support the nearly neutral theory of RGE, driven by a reduction of effective population size in these taxa.

Finally, in agreement with the findings of Stewart et al.^{27,51}, Decker et al.⁵², and Ozawa et al.⁵³, we detected no recombination between Clade I and II symbionts even though some of the host taxa co-occur^{54–56}. These findings imply that there is enough molecular and ecological divergence between the two clades for clonal interference and/or strong host-symbiont epistatic interactions to constrain symbiont exchange^{20,52}. Thus, our results support the nomenclature initially put forward by Newton *et al.*³⁵ and Kuwahara *et al.*³⁶ classifying the symbionts from Clade I and II into two distinct bacterial genera, *Ca. Vesicomysocius* and *Ca. Ruthia*. For clarity, we will keep referring to these two genera as Clade I and Clade II in the rest of the discussion.

Reductive genome evolution is exacerbated in non-recombining symbionts

Clade I symbionts are in a more advanced state of RGE than the others. Indeed, compared to Clade II, their genomes are smaller and lower in GC%, possess fewer genes and pseudogenes, and exhibit less codon usage bias. The genomes of Clade I symbionts are also more homogeneous. Patterns of gene conservation suggest that much of the loss in this group happened after its speciation but before its radiation, a period of roughly 20Mys^{26,31}. Together with increased substitution rate on its diverging branch these results show that the ancestral Clade I lineage experienced an episodic acceleration of reductive genome evolution. It is likely that the increased level of genome reduction in Clade I results from a reduction of homologous recombination in the ancestor of the group exacerbating Muller's ratchet⁵⁷. Drift-driven loss of recombination machinery may have strongly reduced the rate of genetic exchange among the symbionts in this genus. Indeed, essential genes of the RecF and RecBCD pathways for homologous recombination appear to be lost in all of the Clade I symbionts³⁷ and while horizontal transfer of genetic material is widespread among symbionts within Clade II it is almost absent in Clade I.

Strong linkage disequilibrium forces whole genomes to sweep in populations that lack genetic exchange capabilities. Hence, the loss of homologous recombination genes should favor symbiont replacement in cases where the divergence between “native” and foreign symbionts is low (i.e. when the foreign symbionts are not too easily outcompeted by those that have co-evolved with the host). In fact, we find multiple examples of symbiont replacement among symbionts of Clade I. For instance, individual clams of the species *P. extenta* have acquired the symbionts of the sympatric species *A. diagonalis*. Likewise, Breusing et al.⁵⁶ found a population of *A. gigas* carrying the symbionts of the host species *P. soyoae*. Symbiont replacement occurs in several vertically transmitted symbioses^{58–61} and is speculated to constitute a mechanism for escaping the evolutionary rabbit hole caused by Muller's ratchet^{20,58,62}. The present data support this notion, and future population genomic studies could determine the prevalence of symbiont replacement and relative rates of recombination in these taxa on more recent time scales.

Despite the lack of recombining machinery in Clade I, one lineage in this genus, *Ca. V. gigas*, showed evidence for recombination. It is puzzling how recombination might be occurring in this species. Breusing et al.⁵⁶ recently found evidence of unidirectional introgression from *P. soyoae* into *A. gigas*. This mechanism might

enable *A. gigas* symbionts to come into contact with other symbionts. Perhaps the recombination in this species is enabled via host-encoded proteins⁶³. Transfer of symbiont genes to the host nuclear genome is possible and should be investigated in future studies. Indeed, evidence for such transfer was recently found by Ip *et al.*⁴⁴ who identified *Bathymodiolus* symbiont gene homologs in the genome of the *A. marissinica*.

Putative ecological and evolutionary consequences of RGE

The Muller's ratchet has been hypothesized to lead to a progressive loss of fitness in host restricted symbionts²⁰. Sympatric populations of symbionts from Clade I and II represent an excellent model to test this hypothesis because of their contrasting reductive stages. For instance, comparisons of the sulfide physiology of the host species *P. soyoae* and *C. pacifica*, which occupy different micro-niches in the same habitat, reveal that *P. soyoae* individuals have lower sulfide oxidation capacities than those of *C. pacifica*⁵⁵. This could be the consequence of a less efficient sulfide metabolism in *Ca. V. soyoae* resulting from a more advanced state reductive genome evolution in this species compared to *Ca. R. pacifica*. If RGE in the symbionts can restrict their host's ecological range, contrasting degrees of RGE may put constraints on the potential for genetic exchange across different holobiont species and even promote speciation²⁰. Future observational and experimental studies could help define the evolutionary constraints imposed by both host and symbiont physiology and clarify the role of reductive genome evolution in niche partitioning and speciation.

Selective processes in the evolutionary history of the symbionts

Contrasting patterns of gene conservation between the symbionts and their free-living relatives are caused by a shift in selective regime in the host-associated bacteria. Genes enabling bacteria to face the challenges of a free-living environment, such as detoxification, anti-viral defense and inter-species competition, were not conserved in the vesicomyid clam symbionts. Furthermore, different patterns of pseudogenization in Clade I and Clade II likely translate to different physiological adaptations at the level of the holobiont. For example, Breusing *et al.* [in review] found that the two vesicomyid symbiont clades show enzymatic differences related to sulfide oxidation and nitrate reduction and have contrasting dependencies on nickel and vitamin B12 in accordance with adaptations to different ecological niches. In addition, episodes of diversifying selection on genes associated with respiration, ammonia assimilation, and chemosynthesis might reflect the constraints imposed by the diverse selective pressures of host physiology throughout their radiation and niche expansion.

Selective constraints are expected to affect genes involved in host-symbiont interactions. Interspecific communication between eukaryotes and microbes generally involve molecules with distinct motifs produced by the symbiont (e.g., Nod factors, lipopolysaccharides, or peptidoglycans) that are sensed by special receptor in the host^{64,65}. These molecular pathways must experience reciprocal adaptations to persist through speciation and niche expansion. Diversifying selection acting on genes involved in the mediation of host-symbiont interactions such as lipopolysaccharides and peptidoglycans was observed in divergent clades of *Wolbachia*⁶⁶ and many facultative endosymbionts⁶⁷. In a recent study, Chong *et al.*⁶⁸ performed a genome-wide screen for selection in the *Buchnera* symbionts from the aphid subfamily Aphidinae. Of the 371 protein-coding genes tested, the authors detected 29 positively

selected genes representing a variety of metabolic functions including two outer membrane porins (OmpF and OmpA), which are assumed to be important for host interaction.

Surprisingly, in the clam symbionts, we did not detect selection on proteins associated with host-symbiont interactions but found instead a pervasive pattern of diversifying selection that affected many loci related to housekeeping functions such as DNA and RNA metabolism, transcription and translation. Many ribosomal proteins and chaperones showed evidence for episodic positive selection repeatedly throughout the symbiont phylogeny. These results could indicate that the accumulation of slightly deleterious mutations in the symbiont genomes initiates a selective pressure for compensatory mutations^{69,70}. Evidence for such mutations exist in several organelles and symbiont models⁷⁰⁻⁷³. For instance, in insect endosymbionts, positively selected loci of the chaperonin GroEL are suspected to permit better protein binding and allow proper protein folding despite mutations affecting their conformation⁷¹. Alternatively, these signatures of selection might be in response to other generalized selection pressures such as differences in host habitat (e.g., depth). However, the host mitochondria do not overall seem to be similarly affected making this alternative less likely. Regardless, the pervasive nature of episodic diversifying selection at the level of amino acids in the symbiont genomes suggests that increased drift due to effective size reduction is not the sole driver of molecular evolution in these taxa.

Conclusion

The vertically transmitted symbionts of deep-sea vesicomyid clams are an ideal model to study the processes of reductive genome evolution, as they constitute a highly diverse group of host-restricted bacteria with varying degrees of genomic reduction. We show that both neutral and selective processes have played a role in the evolutionary history of these symbionts and that factors affecting their clonality have strongly influenced the rate of genome evolution. While the vesicomyid clams have yet to be successfully bred in aquaria, significant progress has been made towards their cultivation⁷⁴. Examination of the symbionts at the population-level, both within and across individual hosts, will help to decipher the contributions of host physiology, genetic drift, symbiont fitness, cytonuclear incompatibilities, and horizontal gene transfer to their evolution. Additionally, experimental studies on host-symbiont interactions and holobiont metabolism will shed further light onto the role of these symbionts in the ecological partitioning of their hosts.

Acknowledgements

The Monterey Bay Aquarium Research Institute kindly provided samples from the Vrijenhoek collection for this study. We thank the ships' crews and submersible pilots involved in the collections for this study, without whose efforts this work would not have been possible. N. Pratt and A. Baylay contributed to sequencing at the National Oceanography Centre Genomics Facility. This work was supported by NERC National Capability funding. MP acknowledges the support of the National Science and Engineering Research Council (NSERC) of Canada's Alexander Graham Bell graduate scholarship and Michael Smith Foreign Study Supplements. CB's contribution was supported through a postdoctoral fellowship of the German Research Foundation (BR 5488/1-1) and a grant from the United States National Science Foundation awarded to CB's mentor Roxanne Beinart at the University of Rhode

461 Island (OCE-1736932). BA acknowledges support from NSERC research grant
462 #238600. The genome of *Bathymodiolus thermophilus* was sequenced as part of a
463 project titled ‘Understanding the deep-sea biosphere on seafloor hydrothermal vents
464 in the Indian Ridge (No. 20170411)’ funded to YJW by the Ministry of Oceans and
465 Fisheries, Korea.
466

467 **Materials and Methods**

468 ***Sample collection and sequencing***

469 Host taxa examined in this study were chosen from the deepest diverging lineages
470 within the Vesicomidae that are distributed globally in the northern hemisphere
471 (Figure S3) and are representative of the known host diversity³¹. Specimens of nine
472 clam species were collected between 1996 and 2004 over eight research expeditions
473 (Table 3, Figure S3). Depths of sampling locations ranged from 650–3550m. Samples
474 were dissected aboard ship and then frozen at -80C or were frozen whole at -80C.
475 DNA was extracted from symbiont bearing gill tissue using the DNeasy Blood &
476 Tissue extraction kit (Qiagen, Hilden, Germany) following the manufacturer's
477 protocol. Host species identification was initially confirmed by sequencing the host
478 mitochondrial *COI* gene using vesicomid-specific primers²⁸.

479
480 Mixed host and symbiont DNA samples were sequenced in-house on a MiSeq
481 instrument. Genomic DNA libraries were prepared using the KAPA Hyperplus
482 Library Preparation kit (KAPA Biosystems, Wilmington, MA, US) according to kit
483 instructions. Read quality of genomic data was assessed using FastQC³⁸.

485 ***Mitochondrial and symbiont genome reconstruction and annotation***

486 Initial symbiont and mitochondrial assemblies were constructed from the same
487 metagenomic libraries (Table 3) using Velvet⁷⁶, manually optimizing for *k*-mer size
488 distribution and read depth. Some assemblies were also constructed using the read
489 mapping and assembly functions in Geneious version 10.1.3⁷⁷.

490
491 Scaffolding and circularization of the symbiont genomes were performed by mapping,
492 extracting and reassembling reads mapping to the extremities of contigs using
493 Bowtie2⁷⁸, Samtools⁷⁹ and SPAdes⁸⁰, respectively. Mitochondrial genomes were
494 assembled de novo with MITObim⁸¹ using as seed a set of initial contigs constructed
495 using the read mapping and assembly functions in Geneious version 10.1.3⁷⁷.
496 Mitochondrial genome annotations were produced by the GeSeq application⁸² using
497 ARWEN v1.2.3 for tRNA prediction, and manually curated with the aid of previously
498 annotated mitochondrial genomes^{41,42} in Geneious. Mitogenome assembly statistics
499 are presented in Table S6. The symbiont genomes were annotated in RAST⁸³.

500 ***Structural variation and phylogenomic analyses***

501 Host mitochondrial and symbiont genomes were aligned with Progressive Mauve⁸⁴.
502 Progressive Mauve and GRIMM (<http://grimm.ucsd.edu/cgi-bin/grimm.cgi>⁸⁵) were
503 used to identify large-scale structural differences among genomes. Locally collinear
504 blocks (LCBs) longer than 100bp and found in all genomes were extracted with
505 Mauve's stripSubsetLCBs program, aligned with Mafft⁸⁶ and concatenated into host
506 mitochondrial and bacterial core genomes. Phylogenetic trees were produced from
507 these core genomes using the GTR model and 100 bootstraps in PhyML-3.1⁸⁷.

508
509 We compared host and symbiont evolutionary rates by estimating the divergence at
510 synonymous sites for each host pair. Using the Biopython toolkit⁸⁸, we extracted the
511 nucleic and amino acid sequences of 13 conserved mitochondrial and 718 bacterial
512 core protein-coding genes (see below). Amino acid sequences were then aligned with
513 Muscle⁸⁹ and reverse translated into codon alignments using the "build" function

from the Biopython codonalign package. The mitochondrial and bacterial codon-based alignments were then each concatenated into two genome-wide alignments with complete (no gaps, no N) lengths of 10417 bp and 662118 bp, respectively. We assessed substitution saturation by plotting transitions and transversions against adjusted genetic distance. Pairwise synonymous (dS) substitution rates were computed using the Maximum-Likelihood method⁹⁰ implemented in the Biopython codonalign package. The source code was slightly modified to accommodate for ambiguous bases in the mitochondrial genomes.

Identification of bacterial core genes

Because of low structural differences among genomes, orthologous genes could be inferred based on homology and position⁹¹. A list of positional homologs with a minimum identity of 30% and a minimum coverage of 60% was exported from the Mauve alignments. Additional maps with a stricter identity criterion (60% identity, 80% coverage) were produced from the alignments of multiple subsets of symbiont genomes. The consensus of these orthologous maps yielded 749 core genes (Figure S2, Table S1) including 718 core protein-coding genes ranging from 138 bp to 4554 bp (average 975bp) (Table S3).

Bayesian concordance analyses

We used Bucky v.1.2⁹² to estimate the proportion of core protein-coding genes supporting each topology. Putative recombination breakpoints within the 718 core protein-coding genes previously found were identified with GARD and the KH test^{93,94}. Using a false positive discovery rate threshold of 5%, recombination was found in 66 genes which were thus split into multiple contiguous non-recombining gene segments at the inferred breakpoints prior to phylogenetic inference. Bucky takes as input the posterior distribution of topologies for each gene (or gene segment). These distributions were each obtained from 800 trees generated in MrBayes v.3.2.7a⁹⁵ using a Gamma + I rate variation across sites. These trees represented a well-mixed sample of the tree space after convergence of four independent Markov Chain Monte Carlo (MCMC) chains which were each run for 2,000,000 generations after an initial 100,000 generations burn-in period. Trees were sampled every 10,000 generations to avoid autocorrelation. Parameter optimization for the MCMCs was performed by assessing convergence and mixing of both continuous parameters of the model and tree topologies using the R package RWTY v.1.0.2⁹⁶. In Bucky, two independent MCMC runs were carried out using the prior assumption that all genes shared the same topology (alpha=0). MCMC runs performed 1,000,000 updates after an initial 10% burn-in period. One cold and three heated chains (swapping frequency=10) were used to improve mixing and convergence of all of the MCMC runs.

Relaxed and positive selection detection

Relaxation of the strength of selection was detected in the symbiont genomes by two independent methods. First we use the Codon Deviation Coefficient (CDC)⁹⁷ to quantify codon usage bias on all protein-coding genes (Table S2) because this index does not require a priori knowledge of gene expression and is not biased by GC content. Second, we used RELAX⁴⁹ on individual core genes. RELAX detects

change in the strength of selection between two groups by observing change in the distribution of ω (dN/dS ratio) classes in a branch-site random effects likelihood (BS-REL) framework between a set of test and reference branches. We compared *Ca. Vesicomysocius*, *Ca. Ruthia*, and both clades together to the group composed of the free-living lineages.

To reduce false positives in phylogenetic selection tests⁹⁸, genes with significant evidence of recombination (see Bayesian concordance analyses) were excluded from these analyses. Episodic diversifying selection on individual lineages was identified on the remaining non-recombining 652 protein-coding genes using the adaptive Branch-site Random Effects Likelihood method (aBSRel⁹⁹). The Holm-Bonferroni correction for multiple testing was applied and threshold for detection was set to 10% false positive discovery rate. We used the hypergeometric test function `dmvhyper` from `extraDistr` v.1.8.11¹⁰⁰ to test whether the genes under relaxed or positive selection represented a random subsample of all core genes according to SEED categories⁸³. The Fisher test¹⁰¹ was applied to find SEED categories that were over-represented in the genes under relaxed or positive selection.

578 ***Data availability***

579 Symbiont genomes and Sequence Read Archives (SRAs) are available at the National
580 Center for Biotechnology Information (NCBI) under the BioProject PRJNA641445.
581 The mitochondrial genomes were deposited in GenBank under the references
582 MT947381-MT947391.

583
584 Genome alignment files and Rmarkdown scripts of downstream analyses are available
585 at https://github.com/maepz/VesicSymb_Evolution
586

587 **Authors contributions**

588 CRY designed the study; CRY and YJW contributed to data collection; MP, CB, and
589 CRY performed analysis, BA contributed to data interpretation. And all authors co-
590 wrote the manuscript.

591

592 **Competing interests**

593 The authors declare no competing interests.

References

1. Roger, A. J., Muñoz-Gómez, S. A. & Kamikawa, R. The Origin and Diversification of Mitochondria. *Current Biology* **27**, R1177–R1192 (2017).
2. McFadden, G. I. Origin and Evolution of Plastids and Photosynthesis in Eukaryotes. *Cold Spring Harb Perspect Biol* **6**, a016105 (2014).
3. Cavanaugh, C. M. Microbial Symbiosis: Patterns of Diversity in the Marine Environment. *American Zoologist* **34**, 79–89 (1994).
4. Clavijo, J. M., Donath, A., Serôdio, J. & Christa, G. Polymorphic adaptations in metazoans to establish and maintain photosymbioses. *Biological Reviews* **93**, 2006–2020 (2018).
5. Garate, L., Sureda, J., Agell, G. & Uriz, M. J. Endosymbiotic calcifying bacteria across sponge species and oceans. *Scientific Reports* **7**, 43674 (2017).
6. Bright, M. & Bulgheresi, S. A complex journey: transmission of microbial symbionts. *Nat Rev Micro* **8**, 218–230 (2010).
7. Baumann, P. *et al.* Genetics, physiology, and evolutionary relationships of the genus Buchnera: intracellular symbionts of aphids. *Annu. Rev. Microbiol.* **49**, 55–94 (1995).
8. Wernegreen, J. J. Endosymbiont evolution: predictions from theory and surprises from genomes. *Annals of the New York Academy of Sciences* **1360**, 16–35 (2015).
9. Fisher, R. M., Henry, L. M., Cornwallis, C. K., Kiers, E. T. & West, S. A. The evolution of host-symbiont dependence. *Nature Communications* **8**, 15973 (2017).
10. Husnik, F. & Keeling, P. J. The fate of obligate endosymbionts: reduction, integration, or extinction. *Current Opinion in Genetics & Development* **58–59**, 1–8 (2019).
11. Kuwahara, H. *et al.* Reductive genome evolution in chemoautotrophic intracellular symbionts of deep-sea Calyptogena clams. *Extremophiles* **12**, 365–374 (2008).
12. Itoh, T., Martin, W. & Nei, M. Acceleration of genomic evolution caused by enhanced mutation rate in endocellular symbionts. *PNAS* **99**, 12944–12948 (2002).
13. Moran, N. A., McLaughlin, H. J. & Sorek, R. The Dynamics and Time Scale of Ongoing Genomic Erosion in Symbiotic Bacteria. *Science* **323**, 379–382 (2009).
14. Mendonça, A. G., Alves, R. J. & Pereira-Leal, J. B. Loss of Genetic Redundancy in Reductive Genome Evolution. *PLOS Computational Biology* **7**, e1001082 (2011).
15. Lhee, D. *et al.* Evolutionary dynamics of the chromatophore genome in three photosynthetic Paulinella species. *Scientific Reports* **9**, 2560 (2019).
16. Wernegreen, J. J. & Moran, N. A. Evidence for genetic drift in endosymbionts (Buchnera): analyses of protein-coding genes. *Mol Biol Evol* **16**, 83–97 (1999).
17. Moran, N. A. Accelerated evolution and Muller’s ratchet in endosymbiotic bacteria. *PNAS* **93**, 2873–2878 (1996).
18. Kuo, C.-H., Moran, N. A. & Ochman, H. The consequences of genetic drift for bacterial genome complexity. *Genome Res.* **19**, 1450–1454 (2009).
19. Muller, H. J. The relation of recombination to mutational advance. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **1**, 2–9 (1964).
20. Bennett, G. M. & Moran, N. A. Heritable symbiosis: The advantages and perils of an evolutionary rabbit hole. *PNAS* **112**, 10169–10176 (2015).
21. Andersson, S. G. E. & Kurland, C. G. Reductive evolution of resident genomes. *Trends in Microbiology* **6**, 263–268 (1998).

- 643 22. Andersson, J. O. & Andersson, S. G. Genome degradation is an ongoing process
644 in Rickettsia. *Mol Biol Evol* **16**, 1178–1191 (1999).
- 645 23. Peek, A. S., Vrijenhoek, R. C. & Gaut, B. S. Accelerated evolutionary rate in
646 sulfur-oxidizing endosymbiotic bacteria associated with the mode of symbiont
647 transmission. *Mol Biol Evol* **15**, 1514–1523 (1998).
- 648 24. Cary, S. C. & Giovannoni, S. J. Transovarial inheritance of endosymbiotic
649 bacteria in clams inhabiting deep-sea hydrothermal vents and cold seeps. *PNAS*
650 **90**, 5695–5699 (1993).
- 651 25. Ikuta, T. *et al.* Surfing the vegetal pole in a small population: extracellular vertical
652 transmission of an ‘intracellular’ deep-sea clam symbiont. *Royal Society Open*
653 *Science* **3**, 160130 (2016).
- 654 26. Peek, A. S., Feldman, R. A., Lutz, R. A. & Vrijenhoek, R. C. Cospeciation of
655 chemoautotrophic bacteria and deep sea clams. *PNAS* **95**, 9962–9966 (1998).
- 656 27. Stewart, F. J., Young, C. R. & Cavanaugh, C. M. Evidence for homologous
657 recombination in intracellular chemosynthetic clam symbionts. *Mol. Biol. Evol.*
658 **26**, 1391–1404 (2009).
- 659 28. Peek, A. S., Gustafson, R. G., Lutz, R. A. & Vrijenhoek, R. C. Evolutionary
660 relationships of deep-sea hydrothermal vent and cold-water seep clams (Bivalvia:
661 Vesicomidae): results from the mitochondrial cytochrome oxidase subunit I.
662 *Marine Biology* **130**, 151–161 (1997).
- 663 29. Moran, N. A., Munson Mark A., Baumann Paul & Ishikawa Hajime. A molecular
664 clock in endosymbiotic bacteria is calibrated using the insect hosts. *Proceedings*
665 *of the Royal Society of London. Series B: Biological Sciences* **253**, 167–171
666 (1993).
- 667 30. Ferri, E. *et al.* New Insights into the Evolution of Wolbachia Infections in Filarial
668 Nematodes Inferred from a Large Range of Screened Species. *PLOS ONE* **6**,
669 e20843 (2011).
- 670 31. Johnson, S. B., Krylova, E. M., Audzijonyte, A., Sahling, H. & Vrijenhoek, R. C.
671 Phylogeny and origins of chemosynthetic vesicomid clams. *Systematics and*
672 *Biodiversity* **15**, 346–360 (2017).
- 673 32. Krylova, E. M. & Sahling, H. Vesicomidae (Bivalvia): Current Taxonomy and
674 Distribution. *PLoS ONE* **5**, e9957 (2010).
- 675 33. Audzijonyte, A., Krylova, E. M., Sahling, H. & Vrijenhoek, R. C. Molecular
676 taxonomy reveals broad trans-oceanic distributions and high species diversity of
677 deep-sea clams (Bivalvia: Vesicomidae: Pliocardiinae) in chemosynthetic
678 environments. *Systematics and Biodiversity* **10**, 403–415 (2012).
- 679 34. MolluscaBase. MolluscaBase. Vesicomidae Dall & Simpson, 1901.
680 <http://www.marinespecies.org/aphia.php?p=taxdetails&id=23140#distributions>
681 (2019).
- 682 35. Newton, I. L. G. *et al.* The Calyptogena magnifica Chemoautotrophic Symbiont
683 Genome. *Science* **315**, 998–1000 (2007).
- 684 36. Kuwahara, H. *et al.* Reduced Genome of the Thioautotrophic Intracellular
685 Symbiont in a Deep-Sea Clam, Calyptogena okutanii. *Current Biology* **17**, 881–
686 886 (2007).
- 687 37. Kuwahara, H. *et al.* Loss of genes for DNA recombination and repair in the
688 reductive genome evolution of thioautotrophic symbionts of Calyptogena clams.
689 *BMC Evolutionary Biology* **11**, 285 (2011).
- 690 38. Tamas, I. *et al.* 50 Million Years of Genomic Stasis in Endosymbiotic Bacteria.
691 *Science* **296**, 2376–2379 (2002).

- 692 39. Roeselers, G. *et al.* Complete genome sequence of Candidatus Ruthia magnifica.
693 *Stand Genomic Sci* **3**, 163–173 (2010).
- 694 40. Anantharaman, K., Breier, J. A., Sheik, C. S. & Dick, G. J. Evidence for hydrogen
695 oxidation and metabolic plasticity in widespread deep-sea sulfur-oxidizing
696 bacteria. *PNAS* **110**, 330–335 (2013).
- 697 41. Liu, H., Cai, S., Zhang, H. & Vrijenhoek, R. C. Complete mitochondrial genome
698 of hydrothermal vent clam Calyptogena magnifica. *Mitochondrial DNA Part A* **27**,
699 4333–4335 (2016).
- 700 42. Ozawa, G. *et al.* Updated mitochondrial phylogeny of Pteriomorph and
701 Heterodont Bivalvia, including deep-sea chemosymbiotic Bathymodiolus mussels,
702 vesicomyid clams and the thyasirid clam Conchocele cf. bisecta. *Mar Genomics*
703 **31**, 43–52 (2017).
- 704 43. Yang, M., Gong, L., Sui, J. & Li, X. The complete mitochondrial genome of
705 Calyptogena marissinica (Heterodonta: Veneroida: Vesicomyidae): Insight into
706 the deep-sea adaptive evolution of vesicomyids. *PLoS One* **14**, (2019).
- 707 44. Ip, J. C.-H. *et al.* Host-Endosymbiont Genome Integration in a Deep-Sea
708 Chemosymbiotic Clam. *Molecular Biology and Evolution* (2020)
709 doi:10.1093/molbev/msaa241.
- 710 45. Stackebrandt, E. & Goebel, B. M. Taxonomic Note: A Place for DNA-DNA
711 Reassociation and 16S rRNA Sequence Analysis in the Present Species Definition
712 in Bacteriology. *International Journal of Systematic and Evolutionary*
713 *Microbiology*, **44**, 846–849 (1994).
- 714 46. Cohan, F. M. Sexual Isolation and Speciation in Bacteria. *Genetica* **116**, 359–370
715 (2002).
- 716 47. Fujiwara, Y. *et al.* Phylogenetic characterization of endosymbionts in three
717 hydrothermal vent mussels: influence on host distributions. *Marine Ecology*
718 *Progress Series* **208**, 147–155 (2000).
- 719 48. Petersen, J. M. *et al.* Hydrogen is an energy source for hydrothermal vent
720 symbioses. *Nature* **476**, 176–180 (2011).
- 721 49. Wertheim, J. O., Murrell, B., Smith, M. D., Kosakovsky Pond, S. L. & Scheffler,
722 K. RELAX: Detecting Relaxed Selection in a Phylogenetic Framework. *Mol Biol*
723 *Evol* **32**, 820–832 (2015).
- 724 50. Burke, G. R. & Moran, N. A. Massive Genomic Decay in Serratia symbiotica, a
725 Recently Evolved Symbiont of Aphids. *Genome Biol Evol* **3**, 195–208 (2011).
- 726 51. Stewart, F. J., Young, C. R. & Cavanaugh, C. M. Lateral Symbiont Acquisition in
727 a Maternally Transmitted Chemosynthetic Clam Endosymbiosis. *Mol Biol Evol*
728 **25**, 673–687 (2008).
- 729 52. Decker, C., Olu, K., Arnaud-Haond, S. & Duperron, S. Physical Proximity May
730 Promote Lateral Acquisition of Bacterial Symbionts in Vesicomyid Clams. *PLoS*
731 *One* **8**, (2013).
- 732 53. Ozawa, G. *et al.* Ancient occasional host switching of maternally transmitted
733 bacterial symbionts of chemosynthetic vesicomyid clams. *Genome biology and*
734 *evolution* **9**, 2226–2236 (2017).
- 735 54. Kojima, S. The distribution and the phylogenies of the species of genus
736 Calyptogena and those of vestimentiferans around Japan. *JAMSTEC Journal of*
737 *Deep-Sea Research* **11**, 243–248 (1995).
- 738 55. Goffredi, S. K. & Barry, J. P. Species-specific variation in sulfide physiology
739 between closely related Vesicomyid clams. *Marine Ecology Progress Series* **225**,
740 227–238 (2002).

- 741 56. Breusing, C., Johnson, S. B., Vrijenhoek, R. C. & Young, C. R. Host
742 hybridization as a potential mechanism of lateral symbiont transfer in deep-sea
743 vesicomyid clams. *Molecular Ecology* **28**, 4697–4708 (2019).
- 744 57. Naito, M. & Pawlowska, T. E. Defying Muller’s Ratchet: Ancient Heritable
745 Endobacteria Escape Extinction through Retention of Recombination and
746 Genome Plasticity. *mBio* **7**, e02057-15 (2016).
- 747 58. Pérez-Brocal, V. *et al.* A small microbial genome: the end of a long symbiotic
748 relationship? *Science* **314**, 312–313 (2006).
- 749 59. Koga, R. & Moran, N. A. Swapping symbionts in spittlebugs: evolutionary
750 replacement of a reduced genome symbiont. *ISME J* **8**, 1237–1246 (2014).
- 751 60. Sudakaran, S., Kost, C. & Kaltenpoth, M. Symbiont Acquisition and Replacement
752 as a Source of Ecological Innovation. *Trends in Microbiology* **25**, 375–390 (2017).
- 753 61. Chong, R. A. & Moran, N. A. Evolutionary loss and replacement of Buchnera, the
754 obligate endosymbiont of aphids. *The ISME Journal* **12**, 898 (2018).
- 755 62. Latorre, A. & Manzano-Marin, A. Dissecting genome reduction and trait loss in
756 insect endosymbionts. *Annals of the New York Academy of Sciences* **1389**, 52–75
757 (2017).
- 758 63. Saki, M. & Prakash, A. DNA damage related crosstalk between the nucleus and
759 mitochondria. *Free Radical Biology and Medicine* **107**, 216–227 (2017).
- 760 64. Chu, H. & Mazmanian, S. K. Innate immune recognition of the microbiota
761 promotes host-microbial symbiosis. *Nature immunology* **14**, 668–675 (2013).
- 762 65. Oldroyd, G. E. Speak, friend, and enter: signalling systems that promote
763 beneficial symbiotic associations in plants. *Nature Reviews Microbiology* **11**,
764 252–263 (2013).
- 765 66. Brownlie, J. C., Adamski, M., Slatko, B. & McGraw, E. A. Diversifying selection
766 and host adaptation in two endosymbiont genomes. *BMC Evolutionary Biology* **7**,
767 68 (2007).
- 768 67. Dale, C. & Moran, N. A. Molecular Interactions between Bacterial Symbionts and
769 Their Hosts. *Cell* **126**, 453–465 (2006).
- 770 68. Chong, R. A., Park, H. & Moran, N. A. Genome Evolution of the Obligate
771 Endosymbiont Buchnera aphidicola. *Mol Biol Evol* (2019)
772 doi:10.1093/molbev/msz082.
- 773 69. Wagner, G. P. & Gabriel, W. Quantitative Variation in Finite Parthenogenetic
774 Populations: What Stops Muller’s Ratchet in the Absence of Recombination?
775 *Evolution* **44**, 715–731 (1990).
- 776 70. Rand, D. M., Haney, R. A. & Fry, A. J. Cytonuclear coevolution: the genomics of
777 cooperation. *Trends in Ecology & Evolution* **19**, 645–653 (2004).
- 778 71. Fares, M. A., Moya, A. & Barrio, E. Adaptive evolution in GroEL from distantly
779 related endosymbiotic bacteria of insects. *Journal of Evolutionary Biology* **18**,
780 651–660 (2005).
- 781 72. Howe, D. K. & Denver, D. R. Muller’s Ratchet and compensatory mutation in
782 *Caenorhabditis briggsae* mitochondrial genome evolution. *BMC Evolutionary*
783 *Biology* **8**, 62 (2008).
- 784 73. Castillo, D. M. & Pawlowska, T. E. Molecular Evolution in Bacterial
785 Endosymbionts of Fungi. *Mol Biol Evol* **27**, 622–636 (2010).
- 786 74. Ohishi, K. *et al.* Long-term Cultivation of the Deep-Sea Clam *Calyptogena*
787 *okutanii*: Changes in the Abundance of Chemoautotrophic Symbiont, Elemental
788 Sulfur, and Mucus. *The Biological Bulletin* **230**, 257–267 (2016).
- 789 75. Andrews, S. *et al.* *FastQC*. (2012).

790 76. Zerbino, D. R. & Birney, E. Velvet: algorithms for de novo short read assembly
791 using de Bruijn graphs. *Genome Res.* **18**, 821–829 (2008).
792 77. Kears, M. *et al.* Geneious Basic: An integrated and extendable desktop software
793 platform for the organization and analysis of sequence data. *Bioinformatics* **28**,
794 1647–1649 (2012).
795 78. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*
796 *Meth* **9**, 357–359 (2012).
797 79. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
798 **25**, 2078–2079 (2009).
799 80. Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its
800 Applications to Single-Cell Sequencing. *Journal of Computational Biology* **19**,
801 455–477 (2012).
802 81. Hahn, C., Bachmann, L. & Chevreux, B. Reconstructing mitochondrial genomes
803 directly from genomic next-generation sequencing reads—a baiting and iterative
804 mapping approach. *Nucleic Acids Res* **41**, e129–e129 (2013).
805 82. Tillich, M. *et al.* GeSeq – versatile and accurate annotation of organelle genomes.
806 *Nucleic Acids Res* **45**, W6–W11 (2017).
807 83. Overbeek, R. *et al.* The SEED and the Rapid Annotation of microbial genomes
808 using Subsystems Technology (RAST). *Nucleic Acids Res* **42**, D206–D214 (2014).
809 84. Darling, A. E., Mau, B. & Perna, N. T. progressiveMauve: Multiple Genome
810 Alignment with Gene Gain, Loss and Rearrangement. *PLoS ONE* **5**, e11147
811 (2010).
812 85. Tesler, G. GRIMM: genome rearrangements web server. *Bioinformatics* **18**, 492–
813 493 (2002).
814 86. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software
815 Version 7: Improvements in Performance and Usability. *Mol Biol Evol* **30**, 772–
816 780 (2013).
817 87. Guindon, S. *et al.* New Algorithms and Methods to Estimate Maximum-
818 Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. *Syst Biol* **59**,
819 307–321 (2010).
820 88. Cock, P. J. A. *et al.* Biopython: freely available Python tools for computational
821 molecular biology and bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).
822 89. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
823 throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).
824 90. Goldman, N. & Yang, Z. A codon-based model of nucleotide substitution for
825 protein-coding DNA sequences. *Mol Biol Evol* **11**, 725–736 (1994).
826 91. Lemoine, F., Lespinet, O. & Labedan, B. Assessing the evolutionary rate of
827 positional orthologous genes in prokaryotes using synteny data. *BMC Evol Biol* **7**,
828 237 (2007).
829 92. Larget, B. R., Kotha, S. K., Dewey, C. N. & Ané, C. BUCKY: Gene tree/species
830 tree reconciliation with Bayesian concordance analysis. *Bioinformatics* **26**, 2910–
831 2911 (2010).
832 93. Kishino, H. & Hasegawa, M. Evaluation of the maximum likelihood estimate of
833 the evolutionary tree topologies from DNA sequence data, and the branching
834 order in hominoidea. *J Mol Evol* **29**, 170–179 (1989).
835 94. Kosakovsky Pond, S. L., Posada, D., Gravenor, M. B., Woelk, C. H. & Frost, S. D.
836 W. GARD: a genetic algorithm for recombination detection. *Bioinformatics* **22**,
837 3096–3098 (2006).

- 838 95. Ronquist, F. *et al.* MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and
839 Model Choice Across a Large Model Space. *Systematic Biology* **61**, 539–542
840 (2012).
- 841 96. Warren, D. L., Geneva, A. J. & Lanfear, R. RWTY (R We There Yet): An R
842 package for examining convergence of Bayesian phylogenetic analyses.
843 *Molecular Biology and Evolution* msw279 (2017) doi:10.1093/molbev/msw279.
- 844 97. Zhang, Z. *et al.* Codon Deviation Coefficient: a novel measure for estimating
845 codon usage bias and its statistical significance. *BMC Bioinformatics* **13**, 43
846 (2012).
- 847 98. Pond, S. L. K., Frost, S. D. W. & Muse, S. V. HyPhy: hypothesis testing using
848 phylogenies. *Bioinformatics* **21**, 676–679 (2005).
- 849 99. Smith, M. D. *et al.* Less Is More: An Adaptive Branch-Site Random Effects
850 Model for Efficient Detection of Episodic Diversifying Selection. *Mol Biol Evol*
851 **32**, 1342–1353 (2015).
- 852 100. Wołodźko, T. *extraDistr: Additional Univariate and Multivariate*
853 *Distributions*. (2019).
- 854 101. Fisher, S. R. A. Confidence Limits for a Cross-Product Ratio. *Australian*
855 *Journal of Statistics* **4**, 41–41 (1962).
- 856

857 **Tables**

858 Table 1 Annotation statistics for symbiont and free-living genomes in this study.

859

Sample name	# of contigs (N50 Mbp)	Mean coverage	Genome size (Mbp)	GC %	# of CDS	# of tRNA	# of rRNA	% non- annotated CDS	NCBI Accession number	Reference
<i>Ca. V. okutanii</i>	1	9	1.02	32	939	35	3	7	AP009247	Kuwahara et al. (2007)
<i>Ca. V. soyae</i> (kilmeri)	1	110	1.02	32	983	36	3	11	CP060686	this paper
<i>Ca. V. extenta</i>	1	137	1.02	31	995	36	3	9	CP060685	this paper
<i>Ca. V. diagonalis</i>	1	110	1.02	31	1005	36	3	10	CP060680	this paper
<i>Ca. V. gigas</i>	1	153	1.04	31	979	36	3	10	CP060682	this paper
<i>Ca. R. magnifica</i>	1	14	1.16	34	976	36	3	7	CP000488	Newton et al. (2007)
<i>Ca. R. pliocardia</i>	1	113	1.23	37	1642	36	3	31	CP060688	this paper
<i>Ca. R. southwardae</i>	39 (0.63)	159	1.59	37	2035	36	3	28	JACRUS00	this paper
<i>Ca. R. phaseoliiformis</i>	8 (0.37)	118	1.53	37	2210	36	3	39	JACRUR00	this paper
<i>Ca. R. rectimargo</i>	1	91	1.23	37	1476	37	3	29	CP060684	this paper
<i>Ca. R. pacifica</i>	1	140	1.18	37	1456	35	3	30	CP060683	this paper
<i>Ca. B. thermophilus</i>	1	126	2.83	39	2067	36	3	43	CP024634	
<i>Ca. T. autotrophicus</i> (SUP05)	1	106	1.51	39	1506	35	3	32	CP010552	Shah and Morris (2015)

860 Table 2 Overrepresented functional categories for genes exhibiting significant
861 evidence for episodic diversifying selection

	Overrepresented function	gene	reference locus_tag
Within free-living	t-RNA biogenesis	<i>pheS</i>	Rmag_0643
		<i>tyrS</i>	Rmag_0132
		<i>valS</i>	Rmag_0464
		<i>fnt</i>	Rmag_0785
	ribosome assembly	<i>rplB</i>	Rmag_0168
		<i>rplO</i>	Rmag_0184
		<i>rpsM</i>	Rmag_0187
		<i>rpsS</i>	Rmag_0169
	protein folding	<i>dnaJ</i>	Rmag_0352
		<i>dnaK</i>	Rmag_0353
		<i>htpG</i>	Rmag_0493
		<i>clpB</i>	Rmag_0787
	oxidative phosphorylation	<i>ccmE</i>	Rmag_0659
		<i>ccmF</i>	Rmag_0272
		<i>CYTB/petB</i>	Rmag_0010
		<i>nhd</i>	Rmag_0224
	sulfur oxidation	<i>soxB</i>	Rmag_0156
		<i>soxY</i>	Rmag_0807
	dissimilatory sulfate reduction	<i>aprM</i>	Rmag_0086
		<i>aprAB</i>	Rmag_0088, Rmag_0087
		<i>dsrAB</i>	Rmag_0870, Rmag_0869
bipartition FL-SYMB	ribosomal proteins	<i>rplJ</i>	Rmag_0813
		<i>rpsA</i>	Rmag_0592
		<i>rpsC</i>	Rmag_0171
		<i>rpsH</i>	Rmag_0179
	t-RNA ligases	<i>ileS</i>	Rmag_0340
		<i>cysS</i>	Rmag_0097
		<i>thrS</i>	Rmag_0648
		<i>glyS</i>	Rmag_0721
within symbionts	chaperones, ribosomal proteins	<i>rplC</i>	Rmag_0165
		<i>rplD</i>	Rmag_0166
		<i>rplF</i>	Rmag_0180
		<i>rplL</i>	Rmag_0812
		<i>rpsC</i>	Rmag_0171
		<i>rpsP</i>	Rmag_0990
		<i>glyS</i>	Rmag_0721
		<i>argS</i>	Rmag_0079
	and t-RNA ligases	<i>aspS</i>	Rmag_0396
		<i>gltX</i>	Rmag_0051
		<i>ileS</i>	Rmag_0340
		<i>metG</i>	Rmag_0570
		<i>trpS</i>	Rmag_0338
		<i>dsrA</i>	Rmag_0870
		<i>dsrP</i>	Rmag_0859
		<i>soxB</i>	Rmag_0156
	sulphur metabolism		

		<i>cobB-cbiA</i>	Rmag_0858
	electron	<i>nuoFG</i>	Rmag_0242, Rmag_0243
	donating/accepting	<i>rnfABC</i>	Rmag_0139, Rmag_0140, Rmag_0141
	reactions	<i>rnfE</i>	Rmag_0788
	ammonia	<i>gltB</i>	Rmag_0333, Rmag_1018
	assimilation	<i>glnD</i>	Rmag_0475
within Clade II	de novo purine and	<i>purO</i>	Rmag_0969
	pyrimidine	<i>purL</i>	Rmag_0837
	biosynthesis	<i>purA</i>	Rmag_0531
		<i>purC</i>	Rmag_0392
		<i>carB</i>	Rmag_0875
		<i>pyrDII</i>	Rmag_0963
	carbon fixation	<i>shmt/glyA</i>	Rmag_0632
		<i>rbcL</i>	Rmag_0701
		<i>cbbOQ</i>	Rmag_0699, Rmag_0700
	DNA	<i>ihfB</i>	Rmag_0591
	recombination and	<i>yebC</i>	Rmag_0394
	repair	<i>pcrA/uvrD</i>	Rmag_0080, Rmag_0320
		<i>recJ</i>	Rmag_0649
		<i>uvrA</i>	Rmag_0263
within Clade I	DNA metabolism	<i>parC</i>	Rmag_0302
		<i>dnaX</i>	Rmag_0466
		<i>ihfB</i>	Rmag_0591
		<i>exoI</i>	Rmag_0946

862

863

864 Table 3 Sampling information and genome accession numbers for taxa in this study

865

Species	Accession	Locality ^a	Dive #	Lat	Long	Depth (m)	Year
Clade I							
<i>Ca. Vesicomysocius okutanii</i>	AP009247.1	Sagami Bay		35.2	139.5	1157	2004
<i>Phreagena okutanii</i> (mtDNA)	AP014555	Sagami Bay		35.0150	139.222	852	2007
<i>Ca. Vesicomysocius soyae</i> (kilmeri)	CP060686	Monterey Canyon (s)	V2059	36.7762	-122.084	985	2001
<i>Phreagena soyae</i> (mtDNA)	MT947390						
<i>Ca. Vesicomysocius extenta</i>	CP060685	Monterey Canyon (s)	T406	36.6088	-122.437	2889	2002
<i>Phreagena extenta</i> (mtDNA)	MT947388						
<i>Ca. Vesicomysocius diagonalis</i>	CP060680	Monterey Canyon (s)	T488	36.2254	-122.885	3455	2002
<i>Archivesica diagonalis</i> (mtDNA)	MT947381						
<i>Ca. Vesicomysocius gigas</i>	CP060682	Guaymas Basin (v)	T548	27.3400	-111.270	1754	2003
<i>Archivesica gigas</i> (mtDNA)	MT947383						
Clade II							
<i>Ca. Ruthia magnifica</i>	CP000488.1	East Pacific Rise (v)		9.8505	-104.300	2500	2004
<i>"Calypptogena" magnifica</i> (mtDNA)	KR862368	East Pacific Rise (v)		20.8305	-109.103	2601	2003
<i>Ca. Ruthia pliocardia</i>	CP060688	Blake Spur (s)	A3710	32.4948	-76.185	2155	2001
<i>Pliocardia</i> sp. Blake Ridge (mtDNA)	MT947391						
<i>Ca. Ruthia southwardae</i>	JACRUS00	Logatchev, MAR (v)	A3133	14.7532	-44.980	3038	1997
<i>Abyssogena southwardae</i> (mtDNA)	MT947385						
<i>Ca. Ruthia phaseoliformis</i>	JACRUR00	Aleutian Trench (s)	TVG	54.3050	-157.213	3550	1996
<i>Abyssogena phaseoliformis</i> (mtDNA)	MT947384						
<i>Abyssogena phaseoliformis</i> (mtDNA)	AP014557	Japan trench		39.1052	143.893	5347	2009
<i>Ca. Ruthia rectimargo</i>	CP060684	Monterey Canyon (s)	V2338	36.6816	-122.120	1540	2003
<i>Calypptogena rectimargo</i> (mtDNA)	MT947387						
<i>Ca. Ruthia pacifica</i>	CP060683	Monterey Canyon (s)	V2555	36.7739	-122.049	650	2004
<i>Calypptogena pacifica</i> (mtDNA)	MT947386						
<i>Abyssogena mariana</i> (mtDNA)	LC126311	Mariana trench		11.6569	143.049	5633	2013
"free-living"							
<i>Ca. B. thermophilus</i>	CP024634	East Pacific Rise (v)		9.82	-104.30	2518	2000
<i>Ca. T autotrophicus</i>	CP010552	Effingham Inlet		49.0369	-125.208	60	2013

866 Table S6 Assembly and annotation statistics for mitochondrial genomes in this study.

Sample name	# of contigs (N50 Mbp)	Mean coverage	Genome size (Mbp)	GC %	# of CDS	# of tRNA	# of rRNA	NCBI Accession number	Reference
<i>Phreagena okutani</i>	1	n.a.	16336	34	13	23	2	AP014555	Ozawa et al (2017)
<i>Phreagena soyoae</i>	1	25	19254	34	13	23	2	MT947390	this paper
<i>Phreagena extenta</i>	1	6	18098	33	13	22	2	MT947388	this paper
<i>Archivesica diagonalis</i>	1	8	20322	33	13	22	2	MT947381	this paper
<i>Archivesica gigas</i>	1	7	15625	35	13	21	2	MT947383	this paper
<i>“Calypptogena” magnifica*</i>	1	n.a.	19738	32	13	22	2	KR862368	Liu et al (2016)
<i>Pliocardia</i> sp.	1	20	18885	28	13	22	2	MT947391	this paper
<i>Abyssogena southwardae</i>	1	15	19082	29	13	24	2	MT947385	this paper
<i>Abyssogena phaseoliformis</i>	1	10	17997	31	13	23	2	MT947384	this paper
<i>Abyssogena phaseoliformis*</i>	1	n.a.	19424	30	13	24	2	AP014557	Ozawa et al (2017)
<i>Calypptogena rectimargo</i>	1	22	19326	32	13	25	2	MT947387	this paper
<i>Calypptogena pacifica</i>	1	18	19897	31	13	23	2	MT947386	this paper
<i>Abyssogena mariana</i>	1	n.a.	15927	30	13	23	2	LC126311	Ozawa et al (2017)

*Complete mitogenome

867 **Figures and captions**

868
869
870 Figure 1 Host and symbiont phylogenomic estimates. A) Neighbor-joining phylogeny
871 based on genetic distance (GTR model) between genome-wide alignments of
872 mitochondrial genomes (15272 bp). Numbers in black are bootstrap values. B)
873 Neighbor-joining phylogeny based on genetic distance (GTR model) between
874 genome-wide alignments of symbiont (Clade I; *Ca. Vesicomysocius* in blue, Clade
875 II; *Ca. Ruthia* in green) and free-living (in red) genomes (761866 bp). Chromosome
876 schemes showing genome inversions and assembly fragmentation are displayed at the
877 end of the branches. Refer to text for a description of the genome structures. Numbers
878 in red are the genome-wide mean covariance factors; they represent the percentage of
879 protein-coding genes supporting each split of the phylogeny.

882
883 Figure 2 Relationship between symbiont and mitochondrial divergence. For each
884 holobiont pair, host and symbiont divergences are expressed as pairwise synonymous
885 substitutions rates (dS) in their respective genomes.

886

887

888 Figure 3 Codon bias in symbionts and free-living. A) Codon Deviation Coefficient
 889 (CDC) spectra for each genome (all protein-coding genes). Within the free-living,
 890 yellow: *Ca. B. thermophilus*; red: *Ca. T. autotrophicus*. B) Correlation between the
 891 average CDC of free-living, *Ca. Ruthia* (green) and *Ca. Vesicomysocius* (blue) core
 892 genes. Linear regressions are shown. CDC values vary from 0 (no bias) to 1
 893 (maximum bias). C) Selection parameter (k) spectra of genes for which a significant
 894 change in selection was detected by RELAX. Note that k is on a log scale.

895

896

Figure 4 SEED category distribution of core genes under episodic diversifying selection within phylogenetic clades (A, B, C, D), and on partitioning branches (C, D, E, F). A) Distribution of all non-recombining core genes (dark grey, 652 loci) and loci under selection within the free-living, *Ca. Ruthia*, and *Ca. Vesicomysocius* clades (light grey, 168 loci). The number of loci selected within each clade is represented in the inset. B) Genes under selection within the free-living. C) Genes under selection within *Ca. Ruthia*. D) Genes under selection within the *Ca. Vesicomysocius*. E) Distribution of all non-recombining core genes (dark grey, 652 loci) and loci under selection on all partitioning branches (light grey, 80 loci). The number of loci selected on each branch is represented in the inset. F) Genes under selection on branch a. G) Genes under selection on branch b. H) Genes under selection on branch c. Note that genes may be represented in multiple functional categories and multiple clades or branches. SEED categories significantly overrepresented (in red) and underrepresented (in blue) in the groups compared to the core genome are highlighted. Refer to text for further breakdown of these categories. NA: no functional annotation.

A) Core gene sequences

B) Gene conservation pattern

Figure S1 Distance-based neighbour-joining trees established from A) a concatenated alignment of 652 non-recombining core protein-coding genes sequences (618342 bp, HKY nucleotide substitution model). In red are the number of genes under episodic diversifying selection in each branch; B) the presence/absence of positionally orthologous genes (Jaccard distance on 6110 genes). Numbers above branches are bootstrap support values.

Figure S2 Venn diagram representing the 6110 unique and shared putative genes amongst the free-living, *Ca. Vesicomysocius* and *Ca. Ruthia*. The outer circles represent the pan-genome while the inner circles represent the core-genome of the groups. Free-living: *Ca. B. thermophilus* and *Ca. T. autotrophicus*; *Ca. Ruthia*: *Ca. R. magnifica*, *Ca. R. phaseoliformis*, *Ca. R. pacifica*, *Ca. R. rectiomargo*, *Ca. R. pliocardia*, and *Ca. R. southwardae*; *Ca. Vesicomysocius*: *Ca. V. okutanii*, *Ca. V. soyoae*, *Ca. V. diagonalis-extenta*, and *Ca. V. gigas*. Groups in which more than 50% of the genes are unannotated are identified in red. The complete orthology is available in Table S1.

Figure S3 Sampling locations. Inset depicts samples collected from varying depths in Monterey Bay.