1 Potential linkage between *Toxoplasma gondii* Infection and physical education scores 2 of College Students

3 Zhijin Sheng¹, Yu Jin², Yinan Du³, Xinlei Yan^{4*}, Yong Yao^{5*}

4 1. Department of Physical Education, College of Humanistic Medicine, Anhui Medical University,
5 Hefei, 230032, Anhui, P. R China.

6 2. Anhui Provincial Institute of Parasitic Diseases, Hefei, 230000, Anhui, P. R China.

7 3. Department of Human Parasitology, Anhui Medical University, Hefei, 230032, Anhui, P. R China.

8 4. Food Science and Engineering College of Inner Mongolia Agricultural University, Hohhot
9 010018, China.

10 5. College of Life Sciences, Anhui Medical University, Hefei, 230032, Anhui, P. R China.

11 *Corresponding author: Xinlei Yan (Email: yanxinlei1987620@foxmail.com) or Yong Yao (Email:
12 yaoyongahmu@hotmail.com)

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

ABSTRACT:

Objective *Toxoplasma gondii* is a worldwide protozoan parasite that could infect virtually all warm-blooded animals, including humans. Our study aimed to investigate the prevalence of *T. gondii* infection in college students at Anhui province, China. Moreover, growing studies demonstrated the association between *T. gondii* infection and host behavioral changes. We also studied the linkage between *T. gondii* and scores of college students. **Methods** 2704 serum samples of medical school students attending physical education lessons were collected from September 2017 to September 2019 and evaluated for *T. gondii* IgG antibodies using an enzyme-linked immunosorbent assay (ELISA). We also analysed PE scores of *T. gondii* infected students and *T. gondii* uninfected students. **Results** The overall seroprevalence of *T. gondii* was 11.5%. The main risk factors related to *T. gondii* infections were cat in the household and gardening or agriculture activity. Furthermore, in basketball group and football group, scores of *T. gondii* seropositive students were significantly higher than that of seronegative students, while in other sports there is no difference between scores of *T. gondii* infected students and *T. gondii* uninfected students. **Conclusion** This is the first report of *T. gondii* seroprevalence in college students in Anhui province, China.

Key Words: *Toxoplasma gondii*, College students, Seroprevalence, Physical education

Introduction

Toxoplasma gondii is a kind of intracellular protozoan parasite, which chronically infected about one-third of the world's population. Humans can be infected by *T. gondii* via three major routes including consumption of unwell-cooked meat or raw meat containing *T. gondii* tissue cysts, ingesting foods or water contaminated with oocysts shed by cats, and congenital transmission to fetus during women pregnancy [1].

In immunocompetent persons, *T. gondii* infection is usually asymptomatic or induces only mild clinical symptoms; however, it may lead to devastating conditions, such as lymphadenitis, meningoencephalitis, or ocular toxoplasmosis in immunosuppressive patients (Transplant recipients, cancer patients, and HIV/AIDS patients). If pregnant women are infected with *T. gondii*, this parasite could cross placental barrier to influence fetal development, resulting in intracranial calcification, mental retardation, chronic chorioretinitis, hydrocephalus, and even fetal death [2].

Though *T. gondii* infection shows no apparent clinical manifestations in immunocompetent people, growing evidence showed that *T. gondii* is associated with increased risk-taking behaviours, in both humans and experimental animals. In humans, latent chronic infection with *T. gondii* has been previously linked with suicidal self-directed violence [3], trait aggression and impulsivity [4], and bipolar disorder [5]. Stefanie K. Johnson *et al.* [6] demonstrated the linkage between *T. gondii* infection and complex human behaviours, including those relevant to business, entrepreneurship and economic productivity. In animals, it has been reported that *Toxoplasma* infection not only reduces a mouse innate aversion to predator odors [7, 8], but also, surprisingly, results in the development of an fatal attraction in rats [9]. The behavioural alterations in *T. gondii* infected rodents may be associated with brain inflammation, because *T. gondii* chronically infects host in the form of tissue cysts in muscle and central nerve system (CNS) [8].

In China's high education system, physical education (PE) lesson is one optional course, and college students could select sport of their interest, such as football, basketball, and table tennis. Sports need not only physical quality, but also psychological quality. The seroprevalence of *T. gondii* in college students varies from 22.3% in Brazil [10] to 4.8% in USA [11]. Na Yang *et al.* [12] showed that the seroprevalence of *T. gondii* among the newly enrolled undergraduates students in China was 1.63%. However, the impact of *T. gondii* latent infection on study scores of young college students has not been investigated. In present study, we investigated the prevalence of *T. gondii* infection among college students attending PE lessons and analysed the correlation between *T. gondii* infection and scores of participants.

Materials and Methods

Participants

A total of 2704 whole blood samples of undergraduates of Anhui Medical University (AHMU)

originated from Anhui province of China (between east longitudes of 114°54' to 119°37' and north latitudes of 29°41' to 34°38') were collected from September 2017 to September 2019 survey the presence of *T. gondii* specific antibodies. The age of the involved students ranged from 20 to 22 years. At AHMU, physical education is optional courses, college students could select which kind of sport they like taking by themselves. In present study, scores of PE of students from six groups (Football, basketball, tennis, table tennis, badminton, and volleyball) were extracted from their annual examination records. Teachers for PE do not know the infection status of students.

Ethics Statement

This study was approved by the institutional review board at the Anhui Medical University (#2017USHAEC-026) with written informed consent from all participants. All subjects gave written informed consent in accordance with the Declaration of Helsinki.

Serological Testing

The sera of all participants were tested for the specific IgG to *T. gondii* using commercial ELISA Kit for IgG Antibody to *Toxoplasma* (Haitai Biomed, Zhuhai, China), according to manufacturer's instruction. Briefly, the serum sample was diluted with a ratio of 1:100, followed by adding to test well in the antigen-coated plate and incubation at 37 °C for 30 min. After intensive washing for three times with washing solution, 50 µL peroxidase-conjugated anti-human IgG was added to the wells. After 30 min incubation at 37 °C, each well was washed with washing solution for three times. Then, 50 µL "A" solution and 50 µL "B" solution were added to test wells, and the plate was incubated at 37 °C. Ten min later, reaction was stopped by adding 50 µL stopping solution. Microplates were read at an optical density (OD) of 450 nm in the MK3 microplate reader (Thermo Fisher Scientific, Waltham, MA, USA) and ratios (OD 450 value of serum sample/OD 450 value of negative control) were calculated after correction for the OD 450 value of the blank well. The serum samples were considered positive when the ratio was ≥ 2.1 .

Questionnaire

The questionnaire contained information of basic demographic data, including age, gender, and residence before university. Possible risk factors, including drinking unboiled water, raw or not well-cooked meat (including lamb, beef, pork, fish) and raw vegetable consumption, cat contacts, gardening or agricultural activities and living in urban areas or countryside.

Statistical Analysis

For the statistical analysis, the SPSS 20.0 software package (IBM, Armonk, NY, United States) was used. Statistical analyses of *T. gondii* prevalence in different variables were performed by χ^2 -test. PE scores of students of seropositive students and seronegative students were analysed using *t*-test. *P*-values less than 0.05 were considered statistically significant.

Results

Seroprevalence of *T. gondii* Infection

First, we used ELISA to detect the seroprevalence of *T. gondii* infection among college students attending PE lessons. As shown in Table 1, the overall seroprevalence of *T. gondii* among participants at AHMU was 11.5% (95% CI [10.30-12.70]). There was significant difference in the seroprevalence of *T. gondii* between male students (13.14%, 95% CI [11.51-14.77]) and female students (8.90%, 95% CI [7.17-10.63]) ($\chi^2=10.916$, $p=0.001$). In the univariate analysis, we found two variables associated with anti-*T. gondii* IgG positivity, including cat in the household ($p=0.000$) and gardening or agricultural activity ($p=0.047$) (Table 2).

T. gondii seroprevalence among the students for football, tennis, basketball, table tennis, badminton, and volleyball were 15.64%, 6.53%, 16.63%, 5.46%, 8.62% and 13.56%, respectively. No difference exists between seroprevalence of male and female students in football, tennis, table tennis, badminton, and volleyball classes, while for students in basketball classes, seropositive rate of *T. gondii* in male students (18.75%, 95% CI [14.91-24.59]) was significantly higher than that of female students (9.40%, 95% CI [4.03-14.77]) ($\chi^2=5.050$, $p=0.025$).

We next analysed whether there was difference in scores of *T. gondii* infected students and *T. gondii* uninfected students. As we can see in Figure 2, in basketball (85.32 ± 3.23 vs 83.08 ± 7.19 , $p=0.000$) and football (87.00 ± 7.43 vs 84.80 ± 6.84 , $p=0.010$), scores of *T. gondii* seropositive students were significantly higher than that of seronegative students, while in other sports there is no difference between scores of *T. gondii* infected students and *T. gondii* uninfected students. The similar pattern was observed that scores of seropositive male students were higher than that of seronegative male students, in football group and basketball group (Figure 2 A and B). For female students in basketball group (Figure 2A), seropositive students presented higher scores (85.23 ± 3.13), compared to seronegative students (80.08 ± 8.90).

Discussion

The influence of *T. gondii* chronic infection on host psycho-behaviours has been a hot point in parasitological study. Our results in present study showed that seroprevalence of *T. gondii* infection of undergraduates attending PE lessons at AHMU was 11.5%. In addition, through analysing PE scores of students from different sport groups, we found that compared with seronegative students, scores of seropositive students in basketball group and football group were apparently higher.

Consistent with previous study showing that the overall anti-*T. gondii* IgG prevalence in China was 12.3% [13], our present study indicated that seropositive rate of *T. gondii* of college student was 11.5%. However, it has been reported that seropositive rate of *T. gondii* in newly enrolled

undergraduates from Shenyang Agricultural University (SYAU, China) is 1.63% [12], much lower than our results. This might be due to the different methods used in our study (MAT vs ELISA). A study conducted in Henan province (China) using ELISA indicated that the overall seroprevalence of *T. gondii* in primary school children was 9.51% [14]. Furthermore, students in SYAU are enrolled from all provinces around China, whereas students participating in our study are mainly from Anhui province. The primary geographical location of involved student population may contribute to difference in seroprevalence of *T. gondii*. In Korea, 22.5% of children living in the rural areas of Pyin Oo Lwin and Naung Cho, Myanmar were positive for *T. gondii* IgG [15]; In Brazilian, the IFAT method showed a seroprevalence of 22.3% in college students [10]; in Jordan, *T. gondii* IgG antibodies were detected in 66.5% of undergraduate female university students [16]. In addition, like veterinary students having a high prevalence of antibodies to *T. gondii* (5.6%) at Virginia Tech [11], medical students would have a greater chance at exposure to the parasite than an average population of undergraduate students due to increased contact with clinical samples and experimental animals. Though we found that contact with cat (cat in the household) and exposure to soil (gardening or agriculture) were two main factors for college student infection with *T. gondii* in our study, the detailed risk factors for medical students with high seropositive rate of *T. gondii* infection is worth further investigation.

Interestingly, our study found that *T. gondii* chronic infected rates of college students at contact sport groups (basketball and football) were slightly higher than that of non-contact sport groups (badminton, tennis and table tennis). *T. gondii* infection is associated with increased risk-taking behaviours, potentially due to hormonal or neurological changes resulting from parasitic tissue cysts in host CNS [17]. These physiological changes may directly or indirectly enhance infected students to select more challenging sports. We also observed that *T. gondii* seropositive students had higher scores than seronegative students, in basketball group and football group. This result could be partly explained by previous study showing that nations with higher infection rate of *T. gondii* had a lower fraction of respondents citing ‘fear of failure’ in inhibiting new business ventures. In mice and rats, latent *Toxoplasma* infection converted the aversion to feline odors into attraction [18]. A recent report indicates that *T. gondii* lowers general anxiety in infected mice, increases explorative behaviours, and alters predator aversion without selectivity toward felids, which had a positive correlation with the cyst load in host brain [8]. Given that we have no chance to count tissue cysts in human brains, novel methods need to be developed for detection of neuropathological levels of *T. gondii* chronic infected humans.

In conclusion, our study suggests the relatively high seroprevalence of *T. gondii* infection in college students at Anhui province, China. Though we found that PE scores (basketball and football) of seropositive students were higher, the relationship between student PE scores and *T. gondii* infection should be determined by further well-designed investigations.

Conflict of interest statement

All authors declare that no conflict of interest exists in this study.

Acknowledgements

Our study is financially supported by Research on the humanities and social sciences of Anhui Province (Grant #: SK2017A0163), Anhui high education quality elevation project (Grant #: 2017jyxm0143), Research Foundation for Universities at Anhui (Grant #: KJ2019A0264) and National Natural Science Foundation of China (Grant #: 31701162), and Program of Inner Mongolia Natural Science Foundation of China (No. 2018BS03015).

References

1. Wang D, Liu Y, Jiang T, Zhang G, Yuan G, He J, et al. Seroprevalence and genotypes of *Toxoplasma gondii* isolated from pigs intended for human consumption in Liaoning province, northeastern China. *Parasit Vectors*. 2016;9:248. Epub 2016/05/01. doi: 10.1186/s13071-016-1525-2. PubMed PMID: 27129860; PubMed Central PMCID: PMC4851807.
2. Rorman E, Zamir CS, Rilkis I, Ben-David H. Congenital toxoplasmosis--prenatal aspects of *Toxoplasma gondii* infection. *Reprod Toxicol*. 2006;21(4):458-72. Epub 2005/11/29. doi: 10.1016/j.reprotox.2005.10.006. PubMed PMID: 16311017.
3. Hinze-Selch D, Daubener W, Erdag S, Wilms S. The diagnosis of a personality disorder increases the likelihood for seropositivity to *Toxoplasma gondii* in psychiatric patients. *Folia Parasitol (Praha)*. 2010;57(2):129-35. Epub 2010/07/09. doi: 10.14411/fp.2010.016. PubMed PMID: 20608475.
4. Cook TB, Brenner LA, Cloninger CR, Langenberg P, Igbide A, Giegling I, et al. "Latent" infection with *Toxoplasma gondii*: association with trait aggression and impulsivity in healthy adults. *J Psychiatr Res*. 2015;60:87-94. Epub 2014/10/13. doi: 10.1016/j.jpsychires.2014.09.019. PubMed PMID: 25306262.
5. Pearce BD, Kruszon-Moran D, Jones JL. The relationship between *Toxoplasma gondii*

222 infection and mood disorders in the third National Health and Nutrition Survey. *Biol Psychiatry*.
 223 2012;72(4):290-5. Epub 2012/02/14. doi: 10.1016/j.biopsych.2012.01.003. PubMed PMID:
 224 22325983; PubMed Central PMCID: PMC4750371.

225 6. Johnson SK, Fitz MA, Lerner DA, Calhoun DM, Beldon MA, Chan ET, et al. Risky
 226 business: linking *Toxoplasma gondii* infection and entrepreneurship behaviours across
 227 individuals and countries. *Proc Biol Sci*. 2018;285(1883). Epub 2018/07/28. doi:
 228 10.1098/rspb.2018.0822. PubMed PMID: 30051870; PubMed Central PMCID:
 229 PMCPMC6083268.

230 7. Ingram WM, Goodrich LM, Robey EA, Eisen MB. Mice infected with low-virulence strains
 231 of *Toxoplasma gondii* lose their innate aversion to cat urine, even after extensive parasite
 232 clearance. *PLoS One*. 2013;8(9):e75246. Epub 2013/09/24. doi:
 233 10.1371/journal.pone.0075246. PubMed PMID: 24058668; PubMed Central PMCID:
 234 PMCPMC3776761.

235 8. Boillat M, Hammoudi PM, Dogga SK, Pages S, Goubran M, Rodriguez I, et al.
 236 Neuroinflammation-Associated Aspecific Manipulation of Mouse Predator Fear by *Toxoplasma*
 237 *gondii*. *Cell Rep*. 2020;30(2):320-34 e6. Epub 2020/01/16. doi: 10.1016/j.celrep.2019.12.019.
 238 PubMed PMID: 31940479; PubMed Central PMCID: PMCPMC6963786.

239 9. Berdoy M, Webster JP, Macdonald DW. Fatal attraction in rats infected with *Toxoplasma*
 240 *gondii*. *Proc Biol Sci*. 2000;267(1452):1591-4. Epub 2000/09/28. doi: 10.1098/rspb.2000.1182.
 241 PubMed PMID: 11007336; PubMed Central PMCID: PMCPMC1690701.

242 10. Rodrigues JP, Frei F, Navarro IT, Silva LP, Marcelino MY, de Andrade-Junior HF, et al.
 243 Seroepidemiological analysis of toxoplasmosis in college students. *J Venom Anim Toxins Incl*

- 244 Trop Dis. 2015;21(1):1. Epub 2015/01/15. doi: 10.1186/1678-9199-21-1. PubMed PMID:
245 25587248; PubMed Central PMCID: PMC4293106.
- 246 11. Rosypal AC, Houk AE, Zajac AM, Lindsay DS. Prevalence of IgG Antibodies to
247 Toxoplasma gondii in Veterinary and Undergraduate Students at Virginia Tech, Blacksburg,
248 Virginia. Zoonoses Public Health. 2015;62(7):553-6. Epub 2015/03/11. doi: 10.1111/zph.12184.
249 PubMed PMID: 25753511.
- 250 12. Yang N, Wang D, Xing M, Li C, Li J, Wu A, et al. Seroepidemiology and Risk Factors of
251 Toxoplasma gondii Infection among the Newly Enrolled Undergraduates and Postgraduate
252 Students in China. Front Microbiol. 2017;8:2092. Epub 2017/11/11. doi:
253 10.3389/fmicb.2017.02092. PubMed PMID: 29123513; PubMed Central PMCID:
254 PMCPMC5662622.
- 255 13. Xiao Y, Yin J, Jiang N, Xiang M, Hao L, Lu H, et al. Seroepidemiology of human
256 Toxoplasma gondii infection in China. BMC Infect Dis. 2010;10:4. Epub 2010/01/09. doi:
257 10.1186/1471-2334-10-4. PubMed PMID: 20055991; PubMed Central PMCID:
258 PMCPMC2818656.
- 259 14. Wang S, Yao Z, Li H, Li P, Wang D, Zhang H, et al. Seroprevalence and risk factors of
260 Toxoplasma gondii infection in primary school children in Henan province, central China.
261 Parasite. 2020;27:23. Epub 2020/04/08. doi: 10.1051/parasite/2020018. PubMed PMID:
262 32254019; PubMed Central PMCID: PMC7137538.
- 263 15. Thai TL, Jun H, Park SH, Le HG, Lee J, Ahn SK, et al. Seroprevalence of Toxoplasma
264 gondii among School Children in Pyin Oo Lwin and Naung Cho, Upper Myanmar. Korean J
265 Parasitol. 2019;57(3):303-8. Epub 2019/07/10. doi: 10.3347/kjp.2019.57.3.303. PubMed PMID:

31284355; PubMed Central PMCID: PMC6616164.

16. Obaidat MM, Al-Sheyab NA, Bani Salman AE, Lafi SQ. Seroepidemiology and risk factors of *Toxoplasma gondii* infection in undergraduate university female students in Jordan. *Epidemiol Infect.* 2015;143(9):1898-903. Epub 2014/12/30. doi: 10.1017/S0950268814003604. PubMed PMID: 25543692.

17. Prandovszky E, Gaskell E, Martin H, Dubey JP, Webster JP, McConkey GA. The neurotropic parasite *Toxoplasma gondii* increases dopamine metabolism. *PLoS One.* 2011;6(9):e23866. Epub 2011/10/01. doi: 10.1371/journal.pone.0023866. PubMed PMID: 21957440; PubMed Central PMCID: PMC3177840.

18. Vyas A, Kim SK, Giacomini N, Boothroyd JC, Sapolsky RM. Behavioral changes induced by *Toxoplasma* infection of rodents are highly specific to aversion of cat odors. *Proc Natl Acad Sci U S A.* 2007;104(15):6442-7. Epub 2007/04/04. doi: 10.1073/pnas.0608310104. PubMed PMID: 17404235; PubMed Central PMCID: PMC1851063.

Figure legends

Figure 1. Geographic picture of Anhui province, China.

Table 1. Seroprevalence of anti-*T. gondii* IgG antibody in 2704 college students at different PE groups in Anhui province, China.

Table 2. Univariate analysis of the factors associated with *T. gondii* seroprevalence of 2704 college students in Anhui province, China.

Figure 2. PE scores of 2704 college students in Anhui province, China.

Figure 1

Figure 2

	Male			Female			Total		
	No. positive	No. negative	Positive rate (%, 95% CI)	No. positive	No. negative	Positive rate (%, 95% CI)	No. positive	No. negative	Positive rate (%, 95% CI)
Basketball	75	325	18.75 (14.91-24.59)	11	106	9.40 (4.03-14.77)*	86	431	16.63 (13.41-19.86)
Football	64	303	17.44 (13.54-21.34)	15	123	10.87 (5.61-16.13)	79	426	15.64 (12.46-18.82)
Volleyball	34	212	13.82 (9.48-18.16)	28	183	13.27 (8.66-17.89)	62	395	13.56 (10.42-16.72)
Badminton	17	179	8.67 (4.70-12.65)	18	192	8.57 (4.75-12.39)	35	371	8.62 (5.88-11.36)
Tennis	17	219	7.20 (3.88-10.53)	9	153	5.56 (1.99-9.12)	26	372	6.53 (4.09-8.97)
Table tennis	11	203	5.14 (2.16-8.12)	12	195	5.80 (2.59-9.00)	23	398	5.46 (3.28-7.64)
Total	218	1441	13.14 (11.51-14.77)	93	952	8.90 (7.17-10.63) *	311	2393	11.50 (10.30-12.70)

*The female vs the male, $p < 0.05$.

Table 1

Variable	No. positive	No. negative	Positive rate (%; 95% CI)	<i>P</i> -value
Geographical location				
North of the Yangtze River	132	1096	10.75 (9.01-12.48)	0.276
South of the Yangtze River	179	1297	12.12 (10.46-13.79)	
Living in rural areas or city				
City	156	1236	11.21 (9.55-12.87)	0.630
Rural areas	155	1157	11.81 (10.07-13.56)	
Cat in the household				
Yes	150	856	14.91(12.71-17.12)	0.000
No	161	1537	9.48 (8.09-10.88)	
Drink unboiled water				
Yes	158	1201	11.63 (9.92-13.33)	0.857
No	153	1192	11.37 (9.68-13.07)	
Undercooked beef meat/lamb consumption				
Yes	19	153	11.04 (6.31-15.77)	0.847
No	292	2240	11.53 (10.29-12.78)	
Undercooked pork meat consumption				
Yes	9	80	10.11 (3.73-16.50)	0.865
No	302	2313	11.55 (10.32-12.77)	
Undercooked vegetables consumption				
Yes	53	412	11.40 (8.50-14.30)	0.939
No	258	1981	11.52 (10.20-12.85)	
Gardening or agriculture				
Yes	175	1203	12.70 (10.94-14.46)	0.047
No	136	1190	10.26 (8.62-11.85)	

Table 2