Research Article

Sediment-associated microbial community profiling: sample pre-processing through sequential membrane filtration for 16s rDNA amplicon sequencing

Running Head: Pre-processing for Sediment Microbial Community Profiling

Authors and Affiliations

Joeselle M. Serrana^{1,2,3} and Kozo Watanabe^{1,2,3*}

¹Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 3, Matsuyama, Ehime 790-8577, Japan

²Graduate School of Science and Engineering, Ehime University, Bunkyo-cho 3, Matsuyama, Ehime 790-8577, Japan

³Biological Control Research Unit, Center for Natural Sciences and Environmental Research, De La Salle University, 2401 Taft Avenue, Manila 1004, Philippines

***corresponding author**

Prof. Kozo Watanabe, PhD

E-mail Address: watanabe.kozo.mj@ehime-u.ac.jp

Phone & Fax Number: +81 (0) 89 927 9847

ABSTRACT

Sequential membrane filtration as a pre-processing step for the isolation of microorganisms could provide good quality and integrity DNA that can be preserved and kept at ambient temperatures before community profiling through culture-independent molecular techniques, e.g., 16s rDNA amplicon sequencing. Here, we assessed the impact of pre-processing sediment samples by sequential membrane filtration (from 10, 5 to 0.22 μm pore size membrane filters) for 16s rDNA-based community profiling of sediment-associated microorganisms. Specifically, we examined if there would be method-driven differences between non- and pre-processed sediment samples regarding the quality and quantity of extracted DNA, PCR amplicon, resulting high-throughput sequencing reads, microbial diversity, and community composition. We found no significant difference in the quality and quantity of extracted DNA and PCR amplicons between the two methods. Although we found a significant difference in raw and quality-filtered reads, read abundance after bioinformatics processing (i.e., denoising and the chimeric-read filtering steps) were not significantly different. These results suggest that read abundance after these read processing steps were not influenced by sediment processing or lack thereof. Although the non- and pre-processed sediment samples had more unique than shared amplicon sequence variants (ASVs), we report that their shared ASVs accounted for 74% of both methods' absolute read abundance. More so at the genus level, the final collection filter identified most of the genera (95% of the reads) captured from the non-processed samples, with a total of 51 false-negative (2%) and 59 false-positive genera (3%). Accordingly, the diversity estimates and community composition were not significantly different between the non- and pre-processed samples. We demonstrate that while there were differences in shared and unique taxa, both methods revealed comparable microbial diversity and community composition. We also suggest the inclusion of sequential filters (i.e., pre- and mid-filters) in the community profiling, given the additional taxa not detected from the non-processed and the final collection filter. Our observations highlight the feasibility of pre-processing sediment samples for community analysis and the need to further assess sampling strategies to help conceptualize appropriate study designs for sediment-associated microbial community profiling.

Keywords: sediment-associated, microbial communities, river sediments, sequential membrane filtration, 16s rDNA amplicon sequencing.

INTRODUCTION

Microorganisms have long been recognized as useful bioindicators for biomonitoring and ecological assessment of freshwater ecosystems (Payne, 2013; Amleida et al., 2014; Pawlowski et al., 2016). Recent studies took advantage of high-throughput sequencing (HTS) to characterize freshwater sediment-associated microorganisms for impact assessment of anthropogenic activities and environmental factors on diversity and composition and their functions (e.g., Stern et al., 2017, Liao et al., 2019). In particular, 16s rDNA amplicon sequencing is a relatively faster and cheaper approach providing substantially higher taxonomic resolution (Singer et al., 2016), with the capability of detecting unculturable, rare, and novel microorganisms (Browne et al., 2016) in comparison to the conventional strategies, e.g., culture-dependent methods (Fransoza et al., 2015) for microbial community profiling.

Most studies would directly extract microbial DNA from sediment samples, amplify a target hypervariable region of the 16s rDNA gene through polymerase chain reaction (PCR), process for amplicon library construction, and sequence on a high-throughput platform (e.g., Illumina-based technologies). One major challenge with such an approach would be the isolation and capture of good quality and quantity DNA from sediment samples (Harnpicharnchai et al., 2007; Solomon et al., 2016), which mostly contain impurities that inhibit amplification through PCR (Albers et al., 2013). Various commercial extraction kits are available for the rapid processing of environmental samples tailored to yield abundant and high-quality DNA minimizing the effects of enzyme inhibitors, e.g., humic acid, polysaccharides, metals, etc. that must be removed before amplification with the help of proprietary chemicals (Kosch and Summer, 2013; Ni et al., 2016; Lear et al., 2018). However, most of these kits commonly rely on DNA-binding steps via silica spin columns for DNA purification and concentration. This procedure possibly results in DNA loss due to the competitive column-binding of organic matter (Lloyd, MacGregor, and Teske, 2010) that has been reported to selectively retain high molecular-weight DNA fragments (Rohland et al., 2018). Furthermore, collected sediments and other organic matter usually result in a large sample volume that requires proper processing so that DNA representing the whole community can be extracted, similar to environmental DNA samples (Aylagas et al., 2016).

Pre-processing sediment samples by multi-level or sequential membrane filtration have been reported to efficiently isolate high-quality DNA while reducing inhibitory enzyme

compounds (Solomon et al., 2016; Kachiprath et al., 2017; Mathai et al., 2019; Sakami, 2019). Sequential filtration has been used to concentrate microbial biomass and assess communities based on size fractions using filter membranes with different pore sizes (Padilla et al., 2015; Bae, Lyons, and Onstad, 2019). A pre-filter of larger pore size (1.0 to 30 μm) and a collection filter of smaller size (0.22 μm) are commonly used in-line series of filters (Stewart et al., 2012; Liu et al., 2017) to efficiently capture viruses, bacteria, and parasites based on size exclusion (Hill et al., 2007). DNA is then extracted from the final collection filter to separate targeted microorganisms from the comparatively larger eukaryotic cells (e.g., Smith et al., 2017) or to remove large particle-associated microbes from the free-living fraction (e.g., Teeling et al., 2012; Smith et al., 2013; Orsi et al., 2015; Padilla et al., 2015; Schultz et al., 2020).

Previous studies have characterized and compared the microbial community structure of various collection strategies against *in situ* or on-site filtration of particle or sediment collected samples, mainly from marine environments (e.g., Puigcorb et al., 2020; Torres-Beltrn et al., 2019). On-site filtration keeps the sampled microbial communities *in situ* conditions while reducing the time between collection and storage (Puigcorb et al., 2020). The microorganisms from environmental samples should be inactivated right after collection without significant damage to their DNA (Song et al., 2016). Managing this time is critical to prevent bacterial overgrowth or taxonomically biased DNA damage and degradation (Hugerth and Andersson, 2017).

Integrating filtration as a pre-processing step for the isolation of microorganisms could provide good quality and integrity DNA from sediment samples that can be preserved sufficiently well and kept at ambient temperatures before DNA extraction and library construction for HTS-analyses. Most of the studies on applying pre-processing sediment samples by sequential membrane filtration focused on the quality assessment and efficiency of the extracted metagenomic DNA. Solomon et al. (2016) demonstrated that community DNA with minimal shearing was obtained from pre-processing marine sediment samples and performed PCR amplification of the 16S rDNA gene to confirm that the filtration method isolated high-quality DNA. A similar protocol was employed to process arctic sediment samples to characterize the bacterial community structure by 16S rDNA amplicon sequencing (Kachiprath et al., 2017). However, there is no comprehensive information on the potential biases of sequential membrane filtration on the retained microbial taxa compared to its non-processed counterpart, specifically whether sample pre-processing via sequential filtration compare to non-processed

community profiles for quantitative measurements of freshwater microbial diversity and community structure.

Here, we examined if there would be method-driven differences between non- and pre-processed sediment samples (represented by the collection filter) by sequential membrane filtration for microbial community profiling through 16S rDNA amplicon sequencing. Specifically, we evaluated the impact of pre-processing on the quality and quantity of extracted DNA, PCR amplicon, resulting HTS-reads, microbial diversity, and community composition with the non-processed sediment as the basis of comparison. Given the assumption that membrane filters of different size fractions (i.e., samples filtered from membranes of different pore sizes) retain different microbial biomass, we also assessed the difference in relative abundances, composition, and diversity of microbial taxa retained between each filter fractions. We provided the first comparison of the two approaches using 16S rDNA amplicon sequencing for sediment-associated microbial community profiling. Understanding the influence of pre-processing sediment samples for community analysis would be vital for conceptualizing appropriate study designs for sediment-associated microbial community profiling through molecular methods.

MATERIALS AND METHODS

Sediment Collection and Sample Pre-processing

The Trinity River is a large gravel-bed river impounded by the Trinity Dam (164 m a.b.l. and 3020 million m³ storage) and the smaller Lewiston Dam (28 m a.b.l. and 18 million m³ storage) in northern California, USA. It is under current dam operating guidelines with a mean annual flood of approximately 180 m³/s (Gaeuman et al., 2014). Sediment samples from three sites (i.e., sites A and C are from up-welling zones; site B from a down-welling zone) were collected approximately 10 cm below the submerged surface of selected gravel bars in the Trinity River assessed in the study of Serrana et al. (2020). The samples were stored in 50 ml sterile falcon tubes and immediately fixed with 99.5% molecular grade ethanol upon collection. The collected sediment samples were mainly composed of coarse sediments ranging from 1 to 5 mm in diameter, containing smaller sand grains and fine particulate mass. Pre-processing of sediment samples was done two to four hours after collection.

The experimental procedure of the sediment-associated microbial community profiling employed in this study is illustrated in **Figure 1**. Subsamples of ~600 mg each were aliquoted for sequential membrane filtration. The subsamples were resuspended in separate 50 ml solutions containing 0.22 μm filtered river water with Tween 20 (at a concentration of 1 ml l⁻¹ v/v), agitated and mixed via a magnetic stirrer for 30 min. The resuspended subsamples were then filtered through a pre-filter with a 10 μm pore size (Nuclepore™ hydrophilic membrane filter paper; Whatman, Tokyo, Japan), followed by a mid-filter of 5 μm pore size (Mixed cellulose ester membrane filter; Merck Millipore, USA) and finally through a 0.22 μm collection filter (Cellulose mixed ester membrane filter; Merck Millipore, USA). The pre-processed samples were then kept in 2 ml microcentrifuge tubes, immediately fixed with 99.5% molecular grade ethanol. For non-processed sediments, triplicate subsamples of 200 mg were taken from the collected samples preserved in 50 ml Falcon tubes with 99.5% molecular grade ethanol.

DNA Extraction, PCR Amplification, and Sequencing

Before DNA extraction, the membrane filters were taken out from the collection tubes and dried at room temperature until most of the preserving ethanol evaporated. The membrane filter tubes (ethanol with finer particulate mass) and the subsampled non-processed sediments were then subjected to high speed (12,000 rpm) centrifugation for 30 min to resuspend the remaining fine particles and sediments to the bottom of each tube. The supernatant was removed carefully, and the tubes were dried at room temperature to evaporate the remaining ethanol. The dried membrane filters were cut into smaller pieces using sterile scissors and placed back into their original tubes. The samples were then suspended in a buffer consisting of 10 mM EDTA, 50 mM Tris-HCl, 50 mM Na₂HPO₄·7H₂O at pH 8.0 to remove PCR inhibitors (Zhou et al. 1996; Poulain et al., 2015). Genomic DNA was extracted from both the non-processed and filtered subsamples following the protocol of Zhou et al. (1996) (as employed in Solomon et al., 2016). The DNA extracted from non-processed sediment subsamples were combined accordingly before amplification. The quality and quantity of total DNA extracted was initially assessed with a NanoDrop spectrophotometer (NanoDrop 2000, Thermo Scientific).

Amplicon library preparation was carried out through a one-step PCR amplification using modified fusion primers of the V4 hypervariable region of the 16S SSU rRNA gene (i.e., 515F and 806R; Caporaso et al., 2012), with 12-base error-correcting Golay codes on

both forward and reverse primers. The PCR was performed with high-fidelity Phusion polymerase (Thermo Fisher Scientific Inc.) in a T100 Thermal Cycler (Bio-Rad Laboratories, USA). The 25 µl PCR reaction mixture consisted of 5 µl of 5X Phusion GC Buffer, 1.25 µl each of the forward and reverse primers (10 µM), two µl dNTPs (2.5 mM), 0.75 µl DMSO, 0.25 µl Phusion Polymerase (1 U) and one µl of template DNA. The PCR condition followed was initial denaturation at 98°C for 3 min, 25 cycles of denaturation at 98°C for 15 s, annealing at 50°C for 30 s, and extension at 72°C for 30 s, followed by a final extension period at 72°C for 7 min.

Post-amplification, library-quality control was performed by checking the library size distribution via the High-Sensitivity DNA chip (Agilent BioAnalyzer). The libraries were purified and size selected using SPRI beads (AmpureXP, Beckman Coulter Genomics). Amplicon size was ~400-bp. Triplicate quantitative PCR reactions at appropriate dilutions were performed to quantify the amplicon libraries with the KAPPA Illumina Library qPCR Quantification kit (Kappa Biosystems, Wilmington, MA, USA). Negative control was used to monitor contamination from DNA extraction and PCR to post-amplification library quantity and quality verification; however, no quantifiable amplicon was detected for further analysis. The purified amplicon libraries were then normalized, and equimolar amounts were pooled. The 4 nM pooled library was sequenced at the Advanced Research Support Center (ADRES) of Ehime University using the Illumina MiSeq platform with paired-end reads of 300-bp per read.

Read Processing and Taxonomic Assignment

The raw sequence reads generated on the Illumina MiSeq platform were demultiplexed via the command-line tool Cutadapt v.2.1 (Martin, 2011). The 3,805,575 demultiplexed sequences were quality screened, processed, and inferred amplicon sequence variants (ASVs) with the denoising pipeline of the DADA2 v.1.12 package (Callahan et al., 2016) in R v.3.6.2 (R Core Team, 2019). Based on the read error profiles, the reverse reads have poor read quality. Low read abundance with acceptable overlaps between the reads can be accounted for after quality filtering; therefore, only the forward reads were used in the subsequent analysis. Primer contaminants were excluded, and the reads were filtered based on quality and identified sequence variants likely to be derived from sequencing error. ASVs were inferred from the sequence data, subsequently removing chimeric sequences and singletons. The DADA2 pipeline was implemented to use sequence error models to correct amplicon errors in ASVs. Reads with a maximum

expected error greater than 5 were discarded as a quality filtering measure and truncated at a read length of 100-bp. The remaining ASV sequences were aligned to the SILVA database (Pruesse et al., 2007) through the SILVA ACT: Alignment, Classification, and Tree Service online server (www.arb-silva.de/aligner) (Pruesse, Peplies, and Glöckner, 2012). For this analysis, the small subunit (SSU) category was selected, and a minimum similarity identity of 0.95 was set with ten neighbors per query sequence. Sequences below 70% identity were rejected and discarded. The least common ancestor (LCA) method was used for the taxonomic assignment. Chloroplasts, mitochondria, and unclassified ASVs were removed, resulting in a total of 2,875 taxonomically assigned ASVs.

The raw sequence data were deposited into the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under the accession number PRJNA559761. The ASV matrix, the taxonomy, and sample table generated in this study have been deposited in the Figshare data repository (10.6084/m9.figshare.13088834).

Statistical Analysis and Data Visualization

Statistical analyses were performed in R v.3.6.2 (R Core Team, 2019). The significant differences in the quality and quantity of extracted DNA and PCR amplicon libraries, and the HTS-reads for each read processing steps between sites (i.e., A, B, C), and filters [i.e., non-processed (NP), pre-filter (10 µm filter, "10"), mid-filter (5 µm, "5"), and the collection filter (0.22 µm, "0.22")] were tested via two-way analysis of variance (ANOVA), and pairwise comparisons via multiple T-tests in the presence of significant main effects using the `stat_compare_mean()` in the `ggpubr` package (Kassambara, 2018). The correlation between the extracted DNA and PCR amplicon library concentration and purity, and between HTS-read count per processing step (i.e., raw reads, quality filtering, denoising, chimera removal, taxonomic assignment, and ASV count) were tested with Pearson correlation analyses on log-transformed data. A correlogram with significant tests was calculated and visualized with the `Hmisc` and `corrplot` packages (Harrell and Harrell, 2019).

Before subsequent statistical analyses, the ASV table was normalized at median sequencing depth. The shared and unique taxonomic assignment and ASVs between the groups were visualized with Venn diagrams and UpSetR plots (Lex et al., 2014). The boxplots were illustrated via `ggplot2` (Wickham, Chang, and Wickham, 2016). The spatial differences between the microbial communities were visualized using non-metric

dimensional scaling (NMDS) based on Bray-Curtis distances with the `plot_ordination()` function from the `phyloseq` package (McMurdie and Holmes, 2013), and in a hierarchical clustering dendrogram based on the average-linkage algorithm using the `hclust()` function. PERMANOVA (permutational multivariate analysis of variance) (vegan; Oksanen et al., 2013) was performed to identify significant differences in community composition between filters based on the NMDS ordination.

Alpha diversity metrics (i.e., Chao1 richness, Shannon diversity, Pielou's J evenness, Berger-Parker's dominance, and rarity index) were calculated and visualized based on the ASV dataset to identify the changes in community structure between the non-processed and filtered samples using the `plot_alpha_diversities()` function (microbiomeutilities; Sudarshan, Shetty and Lahti, 2018). Significant differences between the alpha diversity of sites and filters were also tested via ANOVA and pairwise comparisons via multiple t-tests in the presence of significant main effects. Linear discriminant analysis (LDA) effect size (LEfSe) (Segata et al., 2011) was performed using the Python LEfSe package (parameters: $p < 0.05$, $q < 0.05$, $LDA > 2.0$) to identify which microbial taxa significantly explained differences in community composition between the filter groups (i.e., NP, 10, 5, 0.22). The LEfSe algorithm was used to determine indicator taxa considering both the abundance and occurrence of a particular taxon.

RESULTS

DNA Yield, PCR Amplicon, and HTS-read Abundance

The initial concentration and the ratio of absorbance (at 260/280 and 260/230) to assess the purity of extracted DNA were measured via spectrophotometry (**Table 1**; **Supplementary Figure 1A** and **1B**). The DNA yield between sites (A, B, and C) and filters (NP, 10, 5, and 0.22) was higher for sites A and B, and NP and 0.22 filters, but a significant difference between the observed values were only reported for the sites. A ratio of ~1.8 is generally accepted as pure DNA for the 260/280 ratio. Although sites B and C, and filters 10 and 5 reported a relatively high 260/280 ratio, ANOVA showed no significant difference in DNA purity between sites and between filters. The 260/230 ratio was also relatively low for all samples given the accepted range of 2.0-2.2 for pure nucleic acid indicative of the presence of contaminants, e.g., EDTA, carbohydrates, and phenol. It was notable that the mean PCR amplicon library concentration of NP was relatively lower than those of the filtered samples, given that it has higher extracted DNA concentration. However, the PCR amplicon library concentrations quantified via qPCR

were not significantly different between sites and between filters. The correlation between extracted DNA and PCR amplicon library concentrations was not significant (Pearson correlation: $r = -0.024$, $p = 0.94$) (**Supplementary Figure S2**).

Based on the site and filter grouping, sites A and C and filters NP, 10, and 0.22 had higher read abundances (from raw reads to reads with taxonomic assignment) and ASV counts than site B and filter 5, respectively (**Supplementary Figure S3** and **Supplementary Table 1**). ANOVA showed no significant difference in read and ASV counts between the sites, while the raw, filtered (ANOVA; $p < 0.05$), denoised, and non-chimeric reads (ANOVA; $p < 0.10$) were significantly different between the filters. Although the amplicon libraries were normalized to equimolar concentrations before HTS, the NP samples had significantly higher absolute raw read abundance than the filtered samples (t-test: $p < 0.05$). After quality filtering, NP was only significantly different against the 5 filters (t-test: $p = 0.047$). Furthermore, the correlations between the read abundances from raw reads to each processing step were all significantly ($p < 0.05$) positive with strong (Pearson's $r > 0.60$) to very strong (Pearson's $r > 0.80$) correlations (**Supplementary Figure S2**).

ASV Richness, Taxonomic Diversity, and Community Composition

From the 2,875 ASVs, 2,871 were identified as bacteria, while 4 ASVs were assigned as archaea (i.e., Nitrosopumilales and Woesearchaeales). We identified a total of 324 microbial genera from 232 families under 161 orders, 85 classes, and 39 phyla, including unclassified taxa (e.g., Unclassified Bacteria). **Figure 2A** presents the relative abundance of the sediment-associated microbial phyla grouped per filter. Phyla with high relative sequence abundances include the Proteobacteria, Bacteroidota, and Acidobacteria (**Figure 2B**). Rhodobacteriaceae and Vicinamibacteriaceae predominantly represented non-processed sediments. Whereas Chitinophagaceae, Microscillaceae, and *Flavobacterium* dominate the 10, 5, and 0.22 filters, respectively (**Supplementary Figure S4**).

To explore the difference between the non-processed and collection filter samples, the shared and unique ASVs and taxa (e.g., Phylum, Class, Order, Family, and Genus) assigned per filter were visualized via Venn diagrams (**Figure 3A** and **Supplementary Figure S5**) and UpSetR plots (**Figure 3B** and **Supplementary Figure S6**). Notably, the 10 filters always showed the highest ASV count throughout the sites (**Table 1**). When grouped by filter type, the 10 filters had the highest unique ASV count with 978, followed

by 0.22, NP, and 5 with 594, 492, and 121 unique ASVs, respectively. The NP and 0.22 collection filters shared 63 ± 89 (Mean $\pm$ SD) or a total of 239 ASVs (74% of reads shared) having 257 ± 143 (total of 493; 16% of reads) and 215 ± 81 (total of 595; 10% of reads) unique ASVs, respectively. When aggregated at the genus level, the two methods shared 35 ± 34 or a total of 108 genera (95% of reads) with 54 ± 40 (total of 51; 2% of reads) and 39 ± 1 (total of 59; 3% of reads) unique genera, respectively. Also, the 10 and 5 filters shared 449 ASVs, and no ASV was shared between all four filters.

Alpha diversity based on Chao1 richness, Shannon diversity, Pielou's evenness, Berger-Parker's dominance, and the rarity index is presented in **Supplementary Figure S7**. ANOVA showed no significant difference between the sites and between filters in richness, diversity, evenness, dominance, and rarity estimates. Both the NMDS ordinations of the genus and ASV datasets indicated that the samples cluster based on the filters as visualized in the ordination space (**Supplementary Figure S8**). Notably, filters 10 and 5, and NP and 0.22 clustered closely together. The hierarchical clustering of samples based on the ASV dataset also showed the separation of NP and 0.22 against the 10 and 5 filters (**Figure 2C**). However, PERMANOVA showed no significant difference in the community composition of both the genus ($R^2 = 0.21$, $p = 0.245$) and ASV ($R^2 = 0.22$, $p = 0.062$) datasets.

Indicator Taxa Analysis

LEfSe was performed to identify taxa significantly explained differences in the community compositions between the filter groups. Thirty-five significantly discriminative features out of 51 were selected before internal Wilcoxon, and 25 had an LDA score > 2 . A cladogram showing the 25 microbial taxa's phylogenetic distribution significantly associated with each filter group is presented in **Figure 4A**. The corresponding LDA values for each taxon are shown in **Figure 4B**.

LEfSe analysis showed that the taxa from four families (i.e., Crocinitomicaceae, Env. OPS 17, Pseudomonadaceae, Rhizobiales *Incertae Sedis*), and two genera (i.e., *Polymorphobacter*, *Pseudomonas*) were significantly abundant in NP compared to other filter groups. For the sequential membrane filters, phylum Elusimicrobiota, four classes [e.g., Subgroup 22 (Acidobacteriota), JG30-KF-CM66 (Chloroflexi)], four orders (e.g., Chitinophagales, Sphingobacteriales), family Acetobacteraceae, and three genera [i.e., DEV114 (Pedosphaeraceae), *Ferruginibacter*, *Phenylobacterium*] were significantly more abundant for the 10 μ m filter, while three orders (i.e., Gemmatales, Haliangiales,

Pirellulales), family Haliangiaceae), and two genera (i.e., *Haliangium*, *Fimbriiglobus*) were significantly more abundant for the 0.22 μ m filter. No taxa were found to be significantly abundant for the 5 μ m filter.

DISCUSSION

In this study, we assessed whether riverine sediment-associated microorganisms would differ between non-processed and pre-processed samples by sequential membrane filtration. We provided the first comparison of the two approaches using 16s rDNA amplicon sequencing for microbial community profiling. We report that although the non- and pre-processed samples (represented by the final collection filter, 0.22) had more unique than shared ASVs, the latter accounted for a total of 239 ASVs that includes 74% of the reads between the two methods. More so at the genus level, the non- and pre-processed samples had a relatively high percentage of total shared genus count (108 genera, 50%) that accounts for 95% of the reads' absolute abundance. This showed that the final collection filter (0.22) captured most of the abundant genus identified from the non-processed samples. Notably, the collection filter detected a total of 59 more unique genera (3% of the reads). These false-positive detections suggested that the pre-processed samples can detect taxa not captured from the non-processed approach.

A range of mechanisms potentially drove the false-positive detections. First, this could be due to the effectiveness of the multiple filtration process to reduce inhibitory compounds. Sequential-filter isolation techniques have been employed to improve the yield of environmental DNA by reducing the concentration of inhibitory compounds, e.g., humic acid, polysaccharides, metals, etc. (Solomon et al., 2016; Kachiprath et al., 2017; Hunter et al., 2019). Specifically, sediment samples contain high humic substances, which are the primary compounds co-extracted with DNA that inhibits enzymes (e.g., *Taq* polymerase) in PCR reactions (Matheson et al., 2010). The reduction of these inhibition compounds could have led to the generation of false-positive taxa in relation to the non-processed samples. However, we observed no significant difference in the quality of extracted DNA to support reduced inhibitory compounds' influence on the false-positive detections. Other reasons, such as sequencing depth (the total number of usable reads from the sequencing machine), have been reported to influence the rate of false-positive detections in metabarcoding studies (Ficetola, Taberlet, and Coissac, 2016). Insufficient sequence depth could result in the non-detection of rare taxa. For example, singletons (single sequence detection, or an OTU/ASV only present in one sample) are usually

considered erroneous sequences or artifacts and are usually removed for subsequent analysis. Increasing the sequencing depth might result in an increase in these reads' abundances in the sample. Also, method-specific or unique taxa could result from having abundant taxa with polymorphisms (Laroche et al., 2017). On the other hand, setting a more stringent parameter for quality filtering could reduce the rate of detecting false positives (Ficetola et al., 2015; Serrana et al., 2018). Given that we employed a relatively lax read quality filtering parameter in this study, the false positive detection could result from low-quality passing reads.

On the other hand, the false-negative taxa (51 genera; 2% of the reads) absent from the collection filter could be microbial groups that passed through the 0.22 μm pore-sized filter. As previously reported by Maejima et al. (2018), isolated bacteria from lake water samples belonging to the Proteobacteria, Bacteroidetes, Firmicutes, and Actinobacteria passed through a 0.22 μm pore size filter. The filtered fractions from < 0.2 μm filtered samples that were usually considered "sterile" were found to still contain miniature cells, ultramicrobacteria (i.e., bacteria whose cell size are smaller than 0.1 μm^3) and slender filamentous bacteria (e.g., Oligoflexia, Proteobacteria) overlooking a broad diversity of filterable agents (Wang et al., 2007; Nakai 2020). However, we observed that the false-negative taxa had very low read abundance, which could be due to smaller cell size leading to low DNA yield. This suggests that the microbial groups that possibly passed through the 0.22 μm pore-sized collection filter were mostly low abundant taxa. Nonetheless, we observed a low read abundance of these false-positive and negative detections. As demonstrated from the diversity and community composition analyses employed in this study, these method-specific taxa would unlikely affect these results.

On the other hand, the pre- and mid-filters had a relatively high count of 449 shared and 978 and 121 unique ASVs, respectively. The non-processed samples only had 1 ASV shared with the pre- and mid-filter, similar to the collection filter. The clear separation between NP and 0.22 against the 10 and 5 filters was also observed in the NMDS ordination and the hierarchical clustering. At the genus-level, the pre- and mid-filters had 57 and 6 unique genera. These values added with the genera shared between the two filters makes a total of 106 captured solely from the pre- and mid-inline filtration. The very low ASV and low genera shared between non-processed and collection filter against the pre- and mid- filters suggested that a huge part of the sediment microbial community is underrepresented or lost from the community profile. A previous study comparing the prokaryotic and eukaryotic diversity and community composition between pre- and

collection filters from lake water samples suggested the possible “pre-filter” bias in the community structure from the collected biomass (Lanzen et al., 2013). They reported contrasting read abundance even though most operational taxonomic units (OTUs) were shared between filters. Sequential filtration of sediments might be a stochastic process where taxa are presumably retained according to cell size rather than their abundance, with the rare taxa retained along the previous filtration step (Pinto et al., 2020). We presented a stronger pre- and mid-filter community composition bias, given that very few ASVs and taxa were shared between the in-line filters against the non- and pre-processed samples. Since we observed that certain sediment-associated microbial taxa were not captured from the non-processed samples, and if only the collection filter is considered to represent the pre-processed samples' microbial community profile, we suggest the inclusion of pre-filters in microbial communities' profiling.

Statistical analyses revealed that groups based on filter were not significantly different in the richness, diversity, and evenness estimates of alpha diversity. Although shared taxa between the two methods were relatively low, community structures based on Bray-Curtis distance were also not significantly different between the two methods. Bray-Curtis dissimilarity is sensitive to differences in abundance between taxa, where abundant taxa are weighted more than the rare ones (Ricotta and Podani, 2017). Although the overall microbial community composition was not significantly different between the two methods, the significantly abundant indicator taxa detected between the filter types were different, primarily due to the variations in the detection of low abundance taxa. Based on LEfSe, representatives from the Alphaproteobacteria (i.e., Rhizobiales *Incertae Sedis*, and *Polymorphobacter*), *Pseudomonas* (Pseudomonadaceae) and the Crocinitomicaceae and the uncultured eubacterium env. OPS 17 were significantly more abundant in the non-processed sediment samples. The taxa affiliated with the Alphaproteobacteria have shown a consistent preference for a particle-attached lifestyle (Mestre et al., 2018). The pre-filter (10 µm filter) had the most significantly more abundant taxa with representatives from Acetobacteraceae (Alphaproteobacteria), Acidobacteriota, Bacteroidota, Chloroflexi, and Elusimicrobiota. Candidate microbial divisions and Chloroflexi have been reported to be primarily recovered when particle samples were subjected to filtration in situ (Torres-Beltrán et al., 2019). The collection filter had significantly more abundant *Fimbrigiobus* (Gemmatales), Pirellulales, and *Haliangium* (Haliangiales) sequences. The first two taxa are classified as members of the Planctomycetes, while the latter belongs to the Myxococcota. A study evaluating the

influence of standard filtration practices on marine particles also reported that proportional abundances in the pre-filter fraction of Myxococcales (Deltaproteobacteria) and Planctomycetes increased with filter volume (Padilla et al., 2015). Furthermore, in-situ filtration (0.4 μ m filter) increased the capture of Planctomycetes by fivefold compared to on-ship in-line filtration (Torres-Beltrán et al., 2019).

The isolation and capture of good quality and quantity DNA from sediment samples are very challenging (Harnpicharnchai et al., 2007; Solomon et al., 2016), and the preservation medium and the time between collection and storage is critical for particle or sediment-associated microorganisms to prevent biased overgrowth and DNA damage before HTS sample processing (Song et al., 2016). We observed that extracted DNA concentration varied between sites and filters and was relatively high for the NP filters. However, no significant difference was observed for the DNA yield between the two methods. PCR amplicon concentration and quality were also not significantly different between the non-processed and processed samples. Hence, we report that the quantity and quality of extracted DNA and its PCR amplicon libraries were not significantly different between the non-processed and processed samples. We should note that we used the same DNA extraction method for both non-processed and processed samples, employing the method of Zhou et al. (1996), which includes the removal of PCR inhibitors, i.e., humic compounds. The chosen DNA extraction method could present different impacts on the characterization of the overall microbial community composition (Ushio, 2019). Previous studies have investigated the influence of filter types and pore sizes on DNA yield from aquatic ecosystems (i.e., on environmental DNA, e.g., Robson et al., 2016; Li et al., 2018). Filters of different pore sizes did not affect the amount of total DNA recovered and detected species from environmental DNA (Li et al., 2018).

The PCR amplicon libraries were normalized before sequencing to assure an even read distribution for all samples. However, the raw HTS-reads and quality-filtered reads were significantly different between methods, with the non-processed significantly having the highest raw read abundance. Interestingly, after the denoising and the chimeric-read filtering steps, the retained reads from the non-processed sample declined and were not significantly different between methods. This suggested that the retained read abundance after the bioinformatics step was not significantly influenced by sediment processing or lack thereof. Previous studies have reported that higher GC content and larger insert size decreased the abundance of reads retained after quality filtering (Huptas et al., 2016). Moreover, fragment length may also impact the base qualities of

Illumina reads (Tan et al., 2019). The decline in read abundance of NP (from being significantly different from the others to insignificant difference) after quality filtering suggests the possibility of the extracted DNA having either high GC content or large fragments which reduced the base qualities of the reads.

Our time from collection to processing and ethanol preservation of the filtered samples were from two to four hours. Previous studies reported that larger processing time between sample collection and filter storage might allow the growth of opportunistic prokaryotic groups introducing bias by microbial population turnover within the sample (Puigcorb et al., 2020). Here the sediments processed for sequential membrane filtration were from samples that have already been preserved in ethanol; hence, this bias was not tested in our experimental design. We recommend further assessment of sediment pre-processing by comparing different filter types and combinations, preservation medium, sample volume, and the influence of various processing time for further method evaluation. This will fully present the capability and viability of on-site sequential membrane filtration as a processing step against the direct collection and preservation of riverine sediment samples.

CONCLUSION

In the present study, we found no significant difference in the quantity and quality of extracted DNA and PCR amplicon between non- and pre-processed sediment samples. Raw and quality-filtered reads were significantly different between methods, but read abundance after bioinformatics processing were not significantly different. These results suggest that read abundance after the bioinformatics steps were not significantly influenced by sediment processing or lack thereof. We report that although the non- and pre-processed sediment samples had more unique than shared ASVs, both methods shared a total of 239 ASVs that accounts for 74% of the reads. More so at the genus level, the final collection filter also detected most of the genus identified from the non-processed samples, with 51 false-negatives (2% of the reads) and 59 false-positive genera (3% of the reads). All of the alpha diversity indices estimated, and the microbial community composition was not significantly different between the non- and pre-processed samples. These results demonstrate that while differences in shared and unique ASVs and microbial taxa were detected, both methods revealed comparable microbial diversity and community composition. We also suggest the inclusion of sequential filters (i.e., pre- and mid-filters) in the community profiling, given the additional

taxa not detected from the non-processed and the final collection filter. We presented the feasibility of pre-processing sediment samples for community analysis and the need for further assessment sampling strategies to help conceptualize appropriate study designs for sediment-associated microbial community profiling.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

JMS performed field sampling and sample processing. JMS and KW conceptualized the study, analyzed the data, and wrote the manuscript.

Funding

This work was supported by the Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research (Grant No. 17H01666, 19K21996, and 19H02276).

Acknowledgments

We are grateful to Dr. Bin Li of the Molecular Ecology and Health Laboratory (MEcoH), Ehime University, members of the Disaster Prevention Research Institute (DPRI), Kyoto University, and Dr. David Gaeuman of the Trinity River Restoration Program (TRRP) for their assistance during the field survey. We thank Dr. Naohito Tokunaga of the Division of Analytical Bio-Medicine for his assistance in performing high-throughput sequencing on the Illumina MiSeq platform of the Advanced Research Support Center (ADRES), Ehime University.

References

- Albers, C. N., Jensen, A., Bælum, J., and Jacobsen, C. S. (2013). Inhibition of DNA polymerases used in Q-PCR by structurally different soil-derived humic substances. *Geomicrobiology Journal*, 30(8), 675-681.
- Almeida, S. F., Elias, C., Ferreira, J., Tornés, E., Puccinelli, C., Delmas, F., ... and Mancini, L. (2014). Water quality assessment of rivers using diatom metrics across Mediterranean Europe: a methods intercalibration exercise. *Science of the Total Environment*, 476, 768-776.
- Aylagas, E., Mendibil, I., Borja, Á., and Rodríguez-Ezpeleta, N. (2016). Marine sediment sample pre-processing for macroinvertebrates metabarcoding: mechanical enrichment and homogenization. *Frontiers in Marine Science*, 3, 203.

Pre-processing for Sediment Microbial Community Profiling

- 545 Bae, S., Lyons, C., and Onstad, N. (2019). A culture-dependent and metagenomic approach of household
546 drinking water from the source to point of use in a developing country. *Water research X*, 2, 100026.
- 547 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., ... and Bai, Y.
548 (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.
549 *Nature biotechnology*, 1.
- 550 Browne, H. P., Forster, S. C., Anonye, B. O., Kumar, N., Neville, B. A., Stares, M. D., ... and Lawley, T. D.
551 (2016). Culturing of 'unculturable' human microbiota reveals novel taxa and extensive sporulation.
552 *Nature*, 533(7604), 543-546.
- 553 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P. (2016).
554 DADA2: high-resolution sample inference from Illumina amplicon data. *Nature methods*, 13(7), 581.
- 555 Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., ... and Gormley, N.
556 (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq
557 platforms. *The ISME journal*, 6(8), 1621.
- 558 Ficitola, G. F., Pansu, J., Bonin, A., Coissac, E., Giguët-Covex, C., De Barba, M., ... & Rayé, G. (2015).
559 Replication levels, false presences and the estimation of the presence/absence from eDNA
560 metabarcoding data. *Molecular ecology resources*, 15(3), 543-556.
- 561 Ficitola, G. F., Taberlet, P., & Coissac, E. (2016). How to limit false positives in environmental DNA and
562 metabarcoding?. *Molecular ecology resources*, 16(3), 604-607.
- 563 Franzosa, E. A., Hsu, T., Sirota-Madi, A., Shafquat, A., Abu-Ali, G., Morgan, X. C., & Huttenhower, C.
564 (2015). Sequencing and beyond: integrating molecular'omics' for microbial community
565 profiling. *Nature Reviews Microbiology*, 13(6), 360-372.
- 566 Gaeuman, D. (2014). High-flow gravel injection for constructing designed in-channel features. *River*
567 *Research and Applications*, 30(6), 685-706.
- 568 Harnpicharnchai, P., Thongaram, T., Sriprang, R., Champreda, V., Tanapongpipat, S., and Eurwilaichitr,
569 L. (2007). An efficient purification and fractionation of genomic DNA from soil by modified trouting
570 method. *Letters in applied microbiology*, 45(4), 387-391.
- 571 Harrell Jr, F. E., and Harrell Jr, M. F. E. (2019). Package 'Hmisc'. CRAN2018, 2019, 235-6.
- 572 Hill, V. R., Kahler, A. M., Jothikumar, N., Johnson, T. B., Hahn, D., and Cromeans, T. L. (2007). Multistate
573 evaluation of an ultrafiltration-based procedure for simultaneous recovery of enteric microbes in 100-
574 liter tap water samples. *Applied and Environmental Microbiology*, 73(13), 4218-4225.
- 575 Hunter, M. E., Ferrante, J. A., Meigs-Friend, G., & Ulmer, A. (2019). Improving eDNA yield and inhibitor
576 reduction through increased water volumes and multi-filter isolation techniques. *Scientific Reports*,
577 9(1), 1-9.
- 578 Hugerth, L. W., & Andersson, A. F. (2017). Analysing microbial community composition through amplicon
579 sequencing: from sampling to hypothesis testing. *Frontiers in Microbiology*, 8, 1561.
- 580 Huptas, C., Scherer, S., & Wenning, M. (2016). Optimized Illumina PCR-free library preparation for
581 bacterial whole genome sequencing and analysis of factors influencing de novo assembly. *BMC*
582 *research notes*, 9(1), 269.
- 583 Kachiprath, B., Puthumana, J., Gopi, J., Solomon, S., Krishnan, K. P., & Philip, R. (2018). Amplicon
584 sequencing based profiling of bacterial diversity from Krossfjorden, Arctic. *Data in brief*, 21, 2522-
585 2525.
- 586 Kassambara, A. (2018). ggpubr: "ggplot2" based publication ready plots. R package version 0.1, 7.

Kosch, T. A., and Summers, K. (2013). Techniques for minimizing the effects of PCR inhibitors in the chytridiomycosis assay. *Molecular ecology resources*, 13(2), 230-236.

Lanzen, A., Simachew, A., Gessesse, A., Chmolowska, D., Jonassen, I., & Øvreås, L. (2013). Surprising prokaryotic and eukaryotic diversity, community structure and biogeography of Ethiopian soda lakes. *PLoS One*, 8(8), e72577.

Laroche, O., Wood, S. A., Tremblay, L. A., Lear, G., Ellis, J. I., & Pochon, X. (2017). Metabarcoding monitoring analysis: the pros and cons of using co-extracted environmental DNA and RNA data to assess offshore oil production impacts on benthic communities. *PeerJ*, 5, e3347.

Lear, G., Dickie, I., Banks, J., Boyer, S., Buckley, H. L., Buckley, T. R., ... and Kamke, J. (2018). Methods for the extraction, storage, amplification and sequencing of DNA from environmental samples. *New Zealand Journal of Ecology*, 42(1), 10-50A.

Lex, A., Gehlenborg, N., Strobel, H., Vuilleumot, R., and Pfister, H. (2014). UpSet: visualization of intersecting sets. *IEEE transactions on visualization and computer graphics*, 20(12), 1983-1992.

Li, J., Lawson Handley, L. J., Read, D. S., & Hänfling, B. (2018). The effect of filtration method on the efficiency of environmental DNA capture and quantification via metabarcoding. *Molecular Ecology Resources*, 18(5), 1102-1114.

Liu, L., Liu, M., Wilkinson, D. M., Chen, H., Yu, X., & Yang, J. (2017). DNA metabarcoding reveals that 200-µm-size-fractionated filtering is unable to discriminate between planktonic microbial and large eukaryotes. *Molecular ecology resources*, 17(5), 991-1002.

Liao, H., Yu, K., Duan, Y., Ning, Z., Li, B., He, L., and Liu, C. (2019). Profiling microbial communities in a watershed undergoing intensive anthropogenic activities. *Science of The Total Environment*, 647, 1137-1147.

Lloyd, K. G., MacGregor, B. J., and Teske, A. (2010). Quantitative PCR methods for RNA and DNA in marine sediments: maximizing yield while overcoming inhibition. *FEMS microbiology ecology*, 72(1), 143-151.

Maejima, Y., Kushimoto, K., Muraguchi, Y., Fukuda, K., Miura, T., Yamazoe, A., ... & Shintani, M. (2018). Proteobacteria and Bacteroidetes are major phyla of filterable bacteria passing through 0.22 µm pore size membrane filter, in Lake Sanaru, Hamamatsu, Japan. *Bioscience, Biotechnology, and Biochemistry*, 82(7), 1260-1263.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet journal*, 17(1), 10-12.

Mathai, P. P., Dunn, H. M., Magnone, P., Zhang, Q., Ishii, S., Chun, C. L., & Sadowsky, M. J. (2019). Association between submerged aquatic vegetation and elevated levels of *Escherichia coli* and potential bacterial pathogens in freshwater lakes. *Science of the Total Environment*, 657, 319-324.

Matheson, C. D., Gurney, C., Esau, N., & Lehto, R. (2010). Assessing PCR inhibition from humic substances. *The Open Enzyme Inhibition Journal*, 3(1), 38-45.

McMurdie, P. J., and Holmes, S. (2013). phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PloS one*, 8(4), e61217.

Mestre, M., Ruiz-González, C., Logares, R., Duarte, C. M., Gasol, J. M., & Sala, M. M. (2018). Sinking particles promote vertical connectivity in the ocean microbiome. *Proceedings of the National Academy of Sciences*, 115(29), E6799-E6807.

Pre-processing for Sediment Microbial Community Profiling

- Nakai, R. (2020). Size Matters: Ultra-small and Filterable Microorganisms in the Environment. *Microbes and Environments*, 35(2), ME20025.
- Ni, C., Horton, D. J., Rui, J., Henson, M. W., Jiang, Y., Huang, X., and Learman, D. R. (2016). High concentrations of bioavailable heavy metals impact freshwater sediment microbial communities. *Annals of Microbiology*, 66(3), 1003-1012.
- Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R. B., ... and Oksanen, M. J. (2013). Package 'vegan'. *Community ecology package*, version, 2(9), 1-295.
- Orsi, W. D., Smith, J. M., Wilcox, H. M., Swalwell, J. E., Carini, P., Worden, A. Z., & Santoro, A. E. (2015). Ecophysiology of uncultivated marine euryarchaea is linked to particulate organic matter. *The ISME journal*, 9(8), 1747-1763.
- Padilla, C. C., Ganesh, S., Gantt, S., Huhman, A., Parris, D. J., Sarode, N., and Stewart, F. J. (2015). Standard filtration practices may significantly distort planktonic microbial diversity estimates. *Frontiers in Microbiology*, 6, 547.
- Pawlowski, J., Lejzerowicz, F., Apotheloz-Perret-Gentil, L., Visco, J., and Esling, P. (2016). Protist metabarcoding and environmental biomonitoring: time for change. *European Journal of Protistology*, 55, 12-25.
- Payne, R. J. (2013). Seven reasons why protists make useful bioindicators. *Acta Protozoologica*, 52(3).
- Pinto, F., Zolfo, M., Beghini, F., Armanini, F., Asnicar, F., Silverj, A., ... & Segata, N. (2020). A step-by-step sequence-based analysis of virome enrichment protocol for freshwater and sediment samples. *bioRxiv*. doi: <https://doi.org/10.1101/2020.09.17.302836>.
- Poulain, A. J., Aris-Brosou, S., Blais, J. M., Brazeau, M., Keller, W. B., and Paterson, A. M. (2015). Microbial DNA records historical delivery of anthropogenic mercury. *The ISME journal*, 9(12), 2541.
- Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., and Glöckner, F. O. (2007). SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic acids research*, 35(21), 7188-7196.
- Pruesse, E., Peplies, J., and Glöckner, F. O. (2012). SINA: accurate high-throughput multiple sequence alignment of ribosomal RNA genes. *Bioinformatics*, 28(14), 1823-1829.
- Puigcorb, V., Ruiz-Gonzlez, C., Masqu, P., & Gasol, J. M. (2020). Sampling Device-Dependence of Prokaryotic Community Structure on Marine Particles: Higher Diversity Recovered by in situ Pumps Than by Oceanographic Bottles. *Frontiers in Microbiology*, 11, 1645.
- R Core Team (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Ricotta, C., & Podani, J. (2017). On some properties of the Bray-Curtis dissimilarity and their ecological meaning. *Ecological Complexity*, 31, 201-205.
- Robson, H. L., Noble, T. H., Saunders, R. J., Robson, S. K., Burrows, D. W., & Jerry, D. R. (2016). Fine-tuning for the tropics: application of eDNA technology for invasive fish detection in tropical freshwater ecosystems. *Molecular ecology resources*, 16(4), 922-932.
- Rohland, N., Glocke, I., Aximu-Petri, A., and Meyer, M. (2018). Extraction of highly degraded DNA from ancient bones, teeth and sediments for high-throughput sequencing. *Nature protocols*, 13(11), 2447-2461.
- Sakami, T. (2019). Collection of Microbial DNA from Marine Sediments. In *Marine Metagenomics* (pp. 17-20). Springer, Singapore.

- 670 Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., and Huttenhower, C. (2011).
671 Metagenomic biomarker discovery and explanation. *Genome biology*, 12(6), 1-18.
- 672 Serrana, J. M., Yaegashi, S., Kondoh, S., Li, B., Robinson, C. T., & Watanabe, K. (2018). Ecological
673 influence of sediment bypass tunnels on macroinvertebrates in dam-fragmented rivers by DNA
674 metabarcoding. *Scientific reports*, 8(1), 1-10.
- 675 Serrana, J. M., Li, B., Sumi, T., Takemon, Y., and Watanabe, K. (2020). Profiling the microbial community
676 structure and functional diversity of a dam-regulated river undergoing gravel bar restoration. *bioRxiv*.
677 doi: <https://doi.org/10.1101/2020.01.25.919381>.
- 678 Schultz, D., Zühlke, D., Bernhardt, J., Francis, T. B., Albrecht, D., Hirschfeld, C., ... & Riedel, K. (2020). An
679 optimized metaproteomics protocol for a holistic taxonomic and functional characterization of
680 microbial communities from marine particles. *Environmental Microbiology Reports*.
- 681 Singer, E., Bushnell, B., Coleman-Derr, D., Bowman, B., Bowers, R. M., Levy, A., ... and Hallam, S. J.
682 (2016). High-resolution phylogenetic microbial community profiling. *The ISME journal*, 10(8), 2020-
683 2032.
- 684 Smith, M. W., Zeigler Allen, L., Allen, A. E., Herfort, L., & Simon, H. M. (2013). Contrasting genomic
685 properties of free-living and particle-attached microbial assemblages within a coastal ecosystem.
686 *Frontiers in microbiology*, 4, 120.
- 687 Smith, M. W., Herfort, L., Fortunato, C. S., Crump, B. C., & Simon, H. M. (2017). Microbial players and
688 processes involved in phytoplankton bloom utilization in the water column of a fast-flowing, river-
689 dominated estuary. *MicrobiologyOpen*, 6(4), e00467.
- 690 Solomon, S., Kachiprath, B., Jayanath, G., Sajeevan, T. P., Singh, I. B., and Philip, R. (2016). High-quality
691 metagenomic DNA from marine sediment samples for genomic studies through a pre-processing
692 approach. *3 Biotech*, 6(2), 160.
- 693 Song, S. J., Amir, A., Metcalf, J. L., Amato, K. R., Xu, Z. Z., Humphrey, G., & Knight, R. (2016). Preservation
694 methods differ in fecal microbiome stability, affecting suitability for field studies. *MSystems*, 1(3).
- 695 Stern, N., Ginder-Vogel, M., Stegen, J. C., Arntzen, E., Kennedy, D. W., Larget, B. R., and Roden, E. E.
696 (2017). Colonization habitat controls biomass, composition, and metabolic activity of attached
697 microbial communities in the Columbia River hyporheic corridor. *Applied and environmental*
698 *microbiology*, 83(16).
- 699 Stewart, F. J., Dalsgaard, T., Young, C. R., Thamdrup, B., Revsbech, N. P., Ulloa, O., ... and DeLong, E.
700 F. (2012). Experimental incubations elicit profound changes in community transcription in OMZ
701 bacterioplankton. *PLoS One*, 7(5), e37118.
- 702 Sudarshan A. Shetty, and Leo Lahti. (2018). *microbiomeutilities*: An R package for utilities to guide in-depth
703 marker gene amplicon data analysis (Version 0.99.00). Zenodo. DOI:10.5281/zenodo.1471685.
- 704 Tan, G., Opitz, L., Schlapbach, R., & Rehauer, H. (2019). Long fragments achieve lower base quality in
705 Illumina paired-end sequencing. *Scientific reports*, 9(1), 1-7.
- 706 Teeling, H., Fuchs, B. M., Becher, D., Klockow, C., Gardebrecht, A., Bennke, C. M., ... & Weber, M. (2012).
707 Substrate-controlled succession of marine bacterioplankton populations induced by a phytoplankton
708 bloom. *Science*, 336(6081), 608-611.
- 709 Torres-Beltrán, M., Mueller, A., Scofield, M., Pachiadaki, M. G., Taylor, C., Tyshchenko, K., ... & Hyun, J.
710 H. (2019). Sampling and processing methods impact microbial community structure and potential

711 activity in a seasonally anoxic fjord: Saanich Inlet, British Columbia. *Frontiers in Marine Science*, 6,
712 132.

713 Ushio, M. (2019). Use of a filter cartridge combined with intra-cartridge bead-beating improves detection
714 of microbial DNA from water samples. *Methods in Ecology and Evolution*, 10(8), 1142-1156.

715 Wang, Y., Hammes, F., Boon, N., & Egli, T. (2007). Quantification of the filterability of freshwater bacteria
716 through 0.45, 0.22, and 0.1 µm pore size filters and shape-dependent enrichment of filterable
717 bacterial communities. *Environmental Science & Technology*, 41(20), 7080-7086.

718 Wickham, H., Chang, W., and Wickham, M. H. (2016). Package 'ggplot2'. Create Elegant Data
719 Visualisations Using the Grammar of Graphics. Version, 2(1), 1-189.

720 Zhou, J., Bruns, M. A., and Tiedje, J. M. (1996). DNA recovery from soils of diverse composition. *Appl.*
721 *Environ. Microbiol.*, 62(2), 316-322.

722 **Supplementary Material**

723 The Supplementary Material for this article is submitted as an attachment: MS-Sediment-
724 Filtering_SuppMat.pdf.

725 **Data Availability Statement**

726 The raw sequence data were deposited into the National Center for Biotechnology
727 Information (NCBI) Sequence Read Archive (SRA) under the accession number
728 PRJNA559761. The ASV matrix, the taxonomy and sample table generated in this study
729 have been deposited in the Figshare data repository (10.6084/m9.figshare.13088834).

730 FIGURES

731

732 **Figure 1.** Schematic overview of the experimental procedure of the sediment-associated microbial community profiling employed in this
 733 study. Collection of sediment samples **(A)**. Sequential membrane filtration from 10, 5 to 0.22 µm pore size filters as pre-processing step
 734 **(B)**. DNA extraction following the protocol of Zhou et al. (1996) (as employed in Solomon et al., 2016) with some modifications **(C)**. One-
 735 step PCR amplification of the 16s rRNA V4 hypervariable region **(D)**. Sequencing through the Illumina MiSeq Platform **(E)**. Bioinformatics
 736 and statistical data analysis were done in R (R Core Team, 2019) **(F)**.

Pre-processing for Sediment Microbial Community Profiling

Figure 2. Relative abundance of microorganisms identified by 16s rDNA amplicon sequencing (A). Compositions are illustrated at the phylum level. The chord diagram indicating the log-transformed abundance of the top three Phylum detected for each filters (B). Hierarchical clustering dendrogram of the similarity in community composition across the sampling sites (C).

Figure 3. Venn Diagrams (A) and UpSetR plots (B) showing shared and unique ASVs and genus between the non-processed (NP) and pre-processed samples (represented by the collection filter, 0.22 μ m), and between all groups (NP, 10, 5, and 0.22 μ m) sediment samples. The bars in the upset plot show the overlap between the indicated sample below.

Pre-processing for Sediment Microbial Community Profiling

Figure 4. Linear Discriminant Analysis (LDA) Effect Size (LEfSe) plot of indicator taxa identified from non-processed (NP), and sequential filtered (10, 5, and 0.22 μ m) sediment samples. Cladogram representing the hierarchical structure of the indicator taxa identified between the non-processed and filtered samples (filter) **(A)**. Each filled circle represents one indicator taxa. Blue, indicator taxa statistically overrepresented in "NP"; red indicator taxa statistically overrepresented in "0.22"; green, indicator taxa statistically overrepresented in "10". Identified indicator taxa grouped by filter and ranked by effect size **(B)**. The threshold for LDA score was >2.0.

758 **TABLES**

759 **Table 1.** Quality and quantity of extracted DNA, PCR amplicon, and HTS-read and amplicon sequence variant (ASV) count per sediment
 760 sample.

Site	Code	Filter Type	Extracted DNA ^a			Amplicon Library			Read Processing				Taxonomic Assignment	
			ng/ul	A ₂₆₀ /A ₂₈₀	A ₂₆₀ /A ₂₃₀	qPCR (nM) ^b	Size (bp)	Band	Raw Reads	Quality Filtered	Denoiised	Non-Chimeric	Reads w/ Tax. ID	ASVs w/ Tax ID
A	ANP	NP	307.1	1.4	0.46	4.88	410	1	1,166,991	560,891	456,284	446,586	424,397	461
	A02	0.22	479.7	1.37	0.61	14.16	409	2	305,045	171,106	154,579	147,490	143,019	436
	A05	5	110.1	1.43	5	45.97	408	1	73,962	40,433	32,181	31,458	30,146	183
	A10	10	619.2	1.45	0.37	56.53	413	1	260,341	148,596	112,796	105,220	96,643	633
B	BNP	NP	625.5	1.45	0.3	56.79	413	1	446,984	135,890	87,857	86,948	85,109	75
	B02	0.22	397.9	1.61	0.75	27.33	414	1	79,771	29,699	22,339	22,153	20,745	104
	B05	5	44.3	2.53	0.06	8.34	415	1	146,808	101,535	86,288	82,169	77,590	465
	B10	10	35.8	3.08	0.04	17.38	415	1	182,665	135,387	125,620	112,804	105,066	1,071
C	CNP	NP	107.1	1.67	0.13	1.44	396	1	790,386	381,687	285,719	275,483	113,715	426
	C02	0.22	2.3	1.86	0.09	76.9	412	2	112,123	61,001	54,375	50,476	46,631	295
	C05	5	5.1	1.03	0.17	43.23	411	3	23,323	14,327	12,515	10,757	9,582	141
	C10	10	7.6	5.1	0.03	4.4	388	2	217,176	138,449	97,116	94,199	89,877	460
Total									3,805,575	1,919,001	1,527,669	1,465,743	1,242,520	2,875

^aInitial quantification and quality assessment of extracted DNA via NanoDrop Spectrophotometer. ^bAmplicon library quantification via Kappa Illumina Library Quantification Kit.

^cDNA Assay for fragment size quantification and quality via Agilent 2100 BioAnalyzer High Sensitivity DNA Kit. "NP" stands for non-processed sediment samples; "10" for the pre-filter (10 µm), "5" for the mid-filter (5 µm), and "0.22" for the collection filter (0.22 µm).

761