1 Nutrient Export and Periphyton Biomass in a Stream-Lake Basin from 2 the Patagonian Andean Region

3

4

5 Alejandro Sosnovsky¹, Mailén Elizabeth Lallement¹, Magalí Rechencq¹, María Valeria
6 Fernández¹, Eduardo Enrique Zattara¹ & Claudia Silvina Feijoó²

7

8 ¹Grupo de Evaluación y Manejo de Recursos Ícticos, INIBIOMA, CONICET-
9 Universidad Nacional del Comahue, Quintral 1250, R8400FRF San Carlos de
10 Bariloche, Argentina

11

12 ²Programa Biogeoquímica de Ecosistemas Dulceacuícolas, INEDES-CONICET,
13 Departamento de Ciencias Básicas, Universidad Nacional de Luján, Ruta Nacional N° 5
14 y Av. Constitución B6700ZBA Luján, Argentina

15

16 Corresponding author: Alejandro Sosnovsky

17 e-mail: alejandro.sosnovsky@comahue-conicet.gob.ar

18 Telephone number: +54 294 423374 (452), Fax number: +54-2944-422111

19

Nutrient Export and Periphyton Biomass in a Stream-Lake Basin from the Patagonian Andean Region

Abstract: We characterized how land use influenced dissolved nutrients and periphytic algal biomass in an Andean basin from Northwest Patagonia. Nutrient export, especially dissolved inorganic nitrogen increased with human population density. However, no correlation between nutrient concentration and algal biomass was found, which could instead be limited by light availability. Our results suggest that local N-limited ecosystems are liable to eutrophication by increased demographic pressure and that alternative wastewater treatment strategies are necessary for sustainable growth.

Keywords: Ecosystem Functioning; Water Quality; Eutrophication; Baseflow period; Spatial studies, On-site wastewater treatment systems

Physicochemical characteristics of a stream reflect the interactions between water and the basin it flows through. For example, a stream's N:P ratio depends on water flow paths of each individual watershed (Green *et al.*, 2007). Eutrophication, the enrichment of surface waters with nutrients, is a key human-induced stressor in streams and often a consequence of intensified land use (Dodds & Smith, 2016). Stream eutrophication results in higher export and concentration of phosphorus (P) and nitrogen (N), usually followed by increased algal biomass and Cyanobacteria blooms (Álvarez *et al.*, 2017). Since algal intracellular nutrient concentration and stream nutrient concentration are directly linked, ambient N:P ratio indicate which of these elements could act as a limiting factor to primary productivity of the ecosystem. A review of field studies found that the optimal C:N:P molar ratio of freshwater benthic algae is 158:18:1 (Kahlert, 1998), suggesting that P is limiting when environmental concentration of N:P exceed 18:1, and N is limiting when ratio is < 18:1. Thus, N:P ratio is a simple metric to evaluate the impact of land use changes on nutrient limitation in stream ecosystems.

The ecological characteristics of Northwest Patagonia region are driven by the Andean range. This region has a temperate-cold climate with markedly seasonal precipitation regimes (Paruelo *et al.*, 1998). Its lakes and forests are generally N-limited ecosystems (Díaz *et al.*, 2007; Diehl *et al.*, 2008), while its Andisols-type soils are characterized by high capacity to retain P (Satti *et al.*, 2003). The region harbours a

53 wide network of stream ecosystems subject to a yearly cycle formed by three
54 contrasting hydrological periods: i) stormflow, when chemistry is controlled by lateral
55 flow through the landscape, ii) meltflow, when chemistry is largely influenced by
56 melting snow in the uplands and iii) baseflow, when chemistry is controlled by
57 groundwater inflow (Ahearn *et al.*, 2004; Sosnovsky *et al.*, 2020). Although a
58 considerable fraction of the region has been under an increasing influence from human
59 activities since urban settlement started by the early 1900's, it is startling that studies
60 linking this influence to eutrophication-driving nutrients like N and P are still scarce
61 (Baffico, 2001; Miserendino *et al.*, 2011; Sosnovsky *et al.*, 2019). We hypothesize that
62 more intensive land use will translate into higher nutrient export and concentration of
63 nutrients in fluvial ecosystems, resulting in increased periphyton biomass on an
64 otherwise nutrient-limited ecosystem. Thus, we aimed to determine during the baseflow
65 period if i) dissolved inorganic nutrient export (i.e., N and P) varied with human
66 population density and ii) changes in nutrient concentration were reflected by changes
67 in periphytic algal biomass.

68 Research was conducted in a suburban area within the city of San Carlos de
69 Bariloche, Northwest Patagonia, Argentina. This city has approximately 110 000
70 inhabitants distributed across several drainage basins. One of them, Gutiérrez stream's
71 drainage basin ($41^{\circ}09'36.18''S$ $71^{\circ}24'37.19''W$), is large (162 km^2) and includes a large
72 and deep lake (Gutiérrez Lake, 17 km^2 , 111 m maximum depth) (Fig. 1). Settlements in
73 this basin are not connected to the city's main sewer system, using instead economic on-
74 site wastewater treatment systems (OWTS) to treat and dispose domestic wastewater.
75 Thus, we can assume that OWTS densities are directly related to local population
76 density within Gutiérrez's basin. Wastewater undergoes primary treatment in a septic
77 tank via sedimentation and anaerobic digestion in OWTS. Secondary treatment of the
78 septic tank effluents occurs within a rubble bed. This bed has a large aerobic zone,
79 favouring nitrification processes over denitrification ones. The lake of this basin has 7
80 inflow perennial streams; its seasonal average concentrations of soluble reactive
81 phosphorus (SRP) and dissolved inorganic nitrogen (DIN) are 3.4 mg m^{-3} and 4.6 mg m^{-3} ,
82 respectively (Díaz *et al.*, 2007). Gutiérrez stream, the outflow of this lake, flows 9 km
83 through a wide gently sloped valley and has a single perennial affluent, the Cascada
84 stream. Gutiérrez's riparian zone is extensively colonized by the exotic crack willow,
85 *Salix fragilis*, and runs through a series of populated areas. In addition, a trout farm with
86 a total production of $\sim 10\text{ tons year}^{-1}$ pours its sewage into Gutiérrez stream.

Fig. 1. a) Gutiérrez stream basin with the sampled sub-basins highlighted. Sampling sites at tributaries and mainstem of Gutiérrez stream are shown: 1) Torrontegui stream, 2) Melgarejo stream, 3) Pescadero stream, 4) Loncochinoco stream, 5) Cascada stream. b) 6-10) Gutiérrez stream. Population zones are shown in grey. c) Export of nutrients and concentration of Chlorophyll a (Chl a). The vertical dotted line represents the lake, to its left are its tributary streams, to its right are the Gutiérrez stream and its tributary. The Y axes corresponding to dissolved inorganic nitrogen (NO₃ + NO₂ + NH₄) and Chl a have two segments for a better visualization of the data.

87

88

89 We surveyed water quality along the Gutiérrez stream basin during year 2016's
90 baseflow period. Demographic density (inhabitants· km⁻²) was calculated from 2010's
91 National Census data (<https://www.indec.gob.ar>). Water samples were taken at four of
92 the headwater lake's tributary streams (1-4), at the Cascada stream (5) and at five points
93 along the main stream (6-10; Table 1, Fig. 1), measured *in-situ* discharge (Q), electrical
94 conductivity (CE), temperature (T) and turbidity, and estimated canopy cover from
95 riparian vegetation using a concave gridded mirror. We filtered 0.5 liters of water

96 through a MGF Munktel filter to determine the concentration of DIN and SRP. We
 97 calculated DIN as the sum of ammonium (NH₄), nitrates (NO₃) and nitrites (NO₂). We
 98 analysed NH₄ by the phenol method, NO₃ + NO₂ with a FUTURA Autoanalyzer
 99 (Alliance Instruments, Frepillon, France) through a reaction with sulfanilamide with a
 100 previous Cu-Cd-reduction for NO₃ and SRP by the ascorbic acid method (A.P.H.A.,
 101 2005). The export nutrients were calculated by the following formula:

$$Nutrient\ export = \frac{[N] Q t}{A}$$

102
 103 Where [N] is nutrient concentration, Q is discharge, t is time and A is drainage basin
 104 area; nutrient export was standardized to kg of nutrient per km² and year (kg km⁻² yr⁻¹).

105 We also determined chlorophyll *a* (Chl *a*) concentration, corrected for
 106 pheopigments (Lorenzen, 1967), as a proxy for periphytic algal biomass. For periphyton
 107 sampling, two representative boulders were chosen and scraped, using a piece of
 108 cylindrical plastic pipe measuring 16 cm² in cross-section and a brush to standardize the
 109 sampled area and remove the algae. Relationships between i) population density and
 110 nutrient export, and ii) nutrient concentration and Chl *a* concentration were modelled
 111 using standard linear regression, log-transforming variables that failed tests for normal
 112 distribution.

Number/Site	Drainage basin (km ²)	Discharge (m ³ s ⁻¹)	EC (μS cm ⁻¹)	Temperature (°C)	Turbidity (NTU)	Nutrient concentration (mg m ⁻³)				N:P (molar)	Canopy (%)
						Nh ₄	NO ₂ + NO ₃	DIN	SRP		
1) Str. Torrontegui	17.0	0.158	92	6.7	0.32	17.1	0.9	18.0	3.7	8.1	7
2) Str. Melgarejo	2.6	0.016	198	8.1	1.47	nd	1.1	1.1	6.5	0.2	0
3) Str. Pescadero	11.3	0.006	74	9.6	0.41	nd	1.7	1.7	4.6	0.2	66
4) Str. Loncochinoco	9.4	0.099	146	9.3	1.14	4.4	488.2	492.6	12.0	21.0	0
5) Str. Cascada	13.0	0.003	87	7.6	0.07	nd	27.9	27.9	8.3	1.7	15
6) Gutiérrez rise	133.1	1.051	70	14.8	0.17	nd	0.5	0.5	3.7	0.1	3
7) Gutiérrez	144.2	1.132	76	12.6	0.24	1.0	14.8	15.8	7.4	1.3	0
8) Gutiérrez	158.3	0.231	88	11.5	0.83	30.2	78.3	108.5	12.0	7.6	100
9) Gutiérrez	160.9	0.938	80	11.7	0.32	12.0	326.0	338.1	11.0	16.6	44
10) Gutiérrez mouth	161.6	0.937	80	11.7	0.35	5.3	452.3	457.6	5.5	42.4	0

Table 1 Morphometric and limnological characteristics of the streams (Str.). Electrical Conductivity (EC); Ammonium (Nh₄); Nitrates + Nitrites (NO₃ + NO₂); Soluble reactive phosphorus (SRP); Dissolved inorganic nitrogen (DIN), Phosphorus:Nitrogen ratio (N:P). Nd: not detectable

Tributary streams to Gutiérrez Lake and stream show a highly heterogeneous distribution of variable values (Fig. 1a, Table 1, supplementary table ST1). These streams had considerably lower temperature and discharge values relative to those found along stream Gutiérrez. In contrast, their nutrient concentrations were higher than those measured at the stream's headwater (site 6) (Table 1). Torrontegui (site 1) and Melgarejo (site 2) streams, draining sparsely populated basins, showed the lowest concentration of $\text{NO}_3 + \text{NO}_2$ and Chl *a* (Fig. 1a, Table 1). Despite the low nutrient export at Gutiérrez stream headwater, we detected an increasing trend in DIN export and N:P ratio towards the stream mouth (Fig. 1c, Table 1). This trend was mainly driven by increased $\text{NO}_3 + \text{NO}_2$ and not NH_4 ; ammonium export was highest at the two sites immediately downstream of the trout farm. The increase in DIN in stream G was not accompanied by increases in either SRP exports or Chl *a* concentration (Fig. 1c, Table 1). Instead, Chl *a* concentration was low in sites shaded by a dense riparian cover (sites 8 and 9, Figure 1c, Table 1).

Integrating our results across the whole study basin, we observed that the stream ecosystem responded to an increase in land use intensity, given that DIN and SRP exports were higher at sites draining more densely populated areas (Fig. 2a-b). This relationship was more noticeable for DIN: the linear regression between DIN and population density was significantly higher ($P = 0.002$, compared to $P = 0.018$ for SRP) and explained a larger amount of variance ($R^2 = 0.72$, compared to $R^2 = 0.52$ for SRP). Similarly, the increase in population density was reflected in an increase in the N:P ratio, which surpasses 18: 1 once population reaches ~ 35 inhabitants per km^2 (Fig. 2c). This potential effect of land use on nutrient load was not followed by a response from periphytic algal biomass (DIN-Chl *a* $R^2 = 0.02$, $n = 10$, $F = 0.17$, $P = 0.69$; SRP-Chl *a* $R^2 = 0.01$, $n = 10$, $F = 0.05$, $P = 0.82$).

Fig. 2. Linear regressions Human density - Nutrients export. a) Dissolved inorganic nitrogen export (DIN_{exp}) b) Soluble reactive phosphorus export (SRP_{exp}) c) Nitrogen:Phosphorus ratio in a molar basis. Dotted line indicates N:P = 18:1.

140

141

142 Our results reflect a positive relationship between population density and exports of DIN
 143 and SRP. On the one hand, the steepest response shown by DIN values indicates that
 144 nitrogen was more readily incorporated into the stream ecosystem. This may be partially
 145 explained by leeching of nitrates from OWTS towards groundwater (Beal *et al.*, 2005; De
 146 & Toor, 2017). In contrast to this pervasive and diffuse source of nitrates to the stream,
 147 ammonium increase is better explained by single point sources, likely excretion from fishes
 148 at the trout farm. On the other hand, OWTS and Andisol soils seem to be better at retaining
 149 phosphorus derived from human activities. Thus, the specific interactions between human
 150 settlement, waste treatment strategies and natural environmental characteristics in this area
 151 skew nutrient export ratios towards nitrogen enrichment. In mountain areas as the study

region, this enrichment becomes most relevant during the baseflow, when groundwater flow path makes the largest contribution to total stream discharge (Ahearn *et al.*, 2004).

The range of ambient N:P ratio has been used to define the transition between N and P limitation in benthic algae (Schanz & Juon, 1983). If the N:P ratio is greater than 20 then P can be assumed to be in limiting supply. If N:P is less than 10 then N can be assumed to be in limiting supply. The distinction of the limiting nutrient when ambient N:P ratios are between 10 and 20 is not precise. The N:P ratio recorded on our study spanned a wide range, from <1 to 42. Thus, the observed lack of response from periphytic algal biomass to the increase in nutrient concentration could be because the limiting nutrient differed between sites, being phosphorous, nitrogen or a colimitation by both them. However, N:P ratio as an indicator of nutrient limitation should be viewed with caution because nutrient requirements for different algal species can vary widely (Allan & Castillo, 2007). In addition, stream ecosystems found within a forest matrix can be limited by light availability (Hill *et al.*, 2001); indeed, we found that sites with low Chl *a* concentration despite high nutrient loads were usually shadier. Thus, we can postulate that in stream ecosystems as the one we studied, the response of primary producers to increased nutrient availability is modulated by the nature and structure of the riparian vegetation. Recently, a significant fraction of the invasive *S. fragilis* population has been removed presenting a great opportunity to test this hypothesis; and more generally, to measure the impact of riparian vegetation on stream productivity.

This work presents some points which deserve further study. NH₄ concentration in Torrontegui stream was unusually high and merits deeper investigation. Moreover, measurements were limited to the baseflow period, which might have different dynamics relating population density and nutrient exports than the stormflow and meltflow periods of the local hydrologic cycle. Furthermore, our understanding about how periphytic algae respond to nutrient availability would benefit from studying species and chemical composition of these primary producers (Carrillo *et al.*, 2018). Finally, the study of the environmental C:N:P relationships will give us a better comprehension of the auto-heterotrophic dynamics of Andean fluvial ecosystems of Patagonia.

Our research yields insights into stream eutrophication, as it links land uses with physicochemical characteristics of a N-limited drainage basin. The headwater lake acted as

an efficient nutrient trap, while residential land use caused increased nutrient load to the basin. Whether this nutrient increase results in algal growth is modulated by the degree of canopy shading. We can conclude that in the Northwest Patagonian region, OWTS as those currently used within Gutiérrez stream basin are likely contributing to eutrophication of streams during the baseflow period. Thus, it would be necessary to consider and evaluate the use of alternative OWTS, with higher denitrification capacity, in order to reduce release of nitrates into groundwater. Furthermore, inhibition of algal response to nitrogen increases by canopy shading could allow build-up of nutrient concentrations to go unnoticed, resulting in potentially harmful blooms if the canopy is removed due to changes in land use. Thus, special care should be taken in preserving the riparian zone of these streams, especially since nitrate concentration of the groundwater might be reduced by denitrification in this zone (Hill, 1996). Given that the Gutiérrez basin is part of a touristic region that can seasonally double its population density, devising mitigation and remediation measurements for human impact is imperative to achieve environmental sustainability.

Acknowledgements

We acknowledge technician Pablo Alvear for field assistance. Sosnovsky, A., Rechencq, M., Fernández, M.V. and Zattara, E.E. are CONICET researchers. This work was funded by Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET resol. N° 3646/13).

References

- A.P.H.A. 2005. *Standard Methods for the Examination of Water and Wastewater*. American Public Health Association. Washington DC.
- AHEARN, D. S., R. W. SHEIBLEY, R. A. DAHLGREN & K. E. KELLER. 2004. Temporal dynamics of stream water chemistry in the last free-flowing river draining the western Sierra Nevada, California. *Journal of Hydrology*, 295: 47-63. DOI:10.1016/j.jhydrol.2004.02.016

213 ÁLVAREZ, X., E. VALERO, R. M. B. SANTOS, S. G. P. VARANDAS, L. F. SANCHES
214 FERNANDES & F. A. L. PACHECO. 2017. Anthropogenic nutrients and
215 eutrophication in multiple land use watersheds: Best management practices and
216 policies for the protection of water resources. *Land Use Policy*, 69: 1-11.
217 DOI.org/10.1016/j.landusepol.2017.08.028

218 ALLAN, J. D. & M. M. CASTILLO. 2007. *Stream Ecology. Structure and Function of*
219 *Running Waters*. Springer. Dordrecht, The Netherlands.

220 BAFFICO, G. 2001. Variations in the periphytic community structure and dynamics of
221 Lake Nahuel Huapi (Patagonia, Argentina). *Hydrobiologia*, 455 (1-3): 79-85.
222 DOI:10.1023/A:1011991402794

223 BEAL, C. D., E. A. GARDNER & N. W. MENZIES. 2005. Process, performance, and
224 pollution potential: A review of septic tank-soil absorption systems. *Australian*
225 *Journal of Soil Research*, 43: 781-802. DOI.org/10.1071/SR05018

226 CARRILLO, U., V. DÍAZ VILLANUEVA & B. E. MODENUTTI. 2018. Sustained effects
227 of volcanic ash on biofilm stoichiometry, enzyme activity and community
228 composition in North-Patagonia streams. *Science of the Total Environment*, 621:
229 235-244. DOI.org/10.1016/j.scitotenv.2017.11.270

230 DE, M. & G. S. TOOR. 2017. Nitrogen transformation in the mounded drainfields of drip
231 dispersal and gravel trench septic systems. *Ecological Engineering*, 102: 352-360.
232 DOI.org/10.1016/j.ecoleng.2017.02.039

233 DÍAZ, M. M., F. L. PEDROZO, C. S. REYNOLDS & P. F. TEMPORETTI. 2007.
234 Chemical composition and the nitrogen-regulated trophic state of Patagonian lakes.
235 *Limnologia*, 37: 17-27. DOI:10.1016/j.limno.2006.08.006

236 DIEHL, P., M. J. MAZZARINO & S. FONTENLA. 2008. Plant limiting nutrients in
237 Andean-Patagonian woody species: effects of interannual rainfall variation, soil
238 fertility and mycorrhizal infection. *Forest Ecology and Management*, 225 (7): 2973-
239 2980. DOI.org/10.1016/j.foreco.2008.02.003

240 DODDS, W. K. & V. H. SMITH. 2016. Nitrogen, phosphorus, and eutrophication in
241 streams. *Inland Waters*, 6: 155-164. DOI: 10.5268/IW-6.2.909

242 GREEN, M. B., J. L. NIEBER, G. JOHNSON, J. MAGNER & B. SCHAEFER. 2007.
 243 Flow path influence on an N:P ratio in two headwater streams: a paired watershed
 244 study. *Journal of Geophysical Research*, 112. DOI:10.1029/2007JG000403

245 HILL, A. R. 1996. Nitrate removal in stream riparian zones. *Journal of Environmental*
 246 *Quality*, 25 (4): 743-755. DOI.org/10.2134/jeq1996.00472425002500040014x

247 HILL, W. R., P. J. MULHOLLAND & E. R. MARZOLF. 2001. Stream ecosystem
 248 response to forest leaf emergence in spring. *Ecology*, 82 (8): 2306-2319.

249 KAHLERT, M. 1998. C:N:P ratios of freshwater benthic algae. *Archiv für Hydrobiologie*,
 250 *Special Issues, Advances in Limnology*, 51: 105-114.

251 LORENZEN, C. J. 1967. Determination of chlorophyll and pheopigments:
 252 spectrophotometric equations. *Limnology and Oceanography*, 12: 343-346.

253 MISERENDINO, M. L., R. CASAUX, M. ARCHANGELSKY, Y. C. DI PRINZIO, C.
 254 BRAND & M. A. KUTSCHKER. 2011. Assessing land-use effects on water
 255 quality, in-stream habitat, riparian ecosystems and biodiversity in Patagonian
 256 northwest streams. *Science of the Total Environment*, 409: 612-624.
 257 DOI:10.1016/j.scitotenv.2010.10.034

258 PARUELO, J. M., A. BELTRAN, E. JOBBÁGY, O. SALA & R. GOLLUSCIO. 1998. The
 259 climate of Patagonia: general patterns and controls on biotic processes. *Ecología*
 260 *Austral*, 8 (2): 85-101.

261 SATTI, P., M. J. MAZZARINO & M. GOBBI. 2003. Soil N dynamics in relation to leaf
 262 litter quality and soil fertility in north-western Patagonian forests. *Journal of*
 263 *Ecology*, 91 (2): 173-181. DOI.org/10.1046/j.1365-2745.2003.00756.x

264 SCHANZ, F. & H. JUON. 1983. Two different methods of evaluating nutrient limitantions
 265 of periphyton bioassays, using water from the River Rhine and eight of its
 266 tributaries. *Hydrobiologia*, 102: 187-195. DOI.org/10.1007/BF00006346

267 SOSNOVSKY, A., M. E. LALLEMENT, M. RECHENCQ, E. E. ZATTARA, M. V.
 268 FERNÁNDEZ, R. DAGA, M. J. SUAREZ, S. LEIVA & R. J. C. CANTET. 2019.
 269 Topography and Land Use Modulate Hydrological and Nutrient Export Dynamics
 270 in two Andean Streams. *BioRxiv*. DOI: <https://doi.org/10.1101/742411>

SOSNOVSKY, A., M. RECHENCQ, V. FERNÁNDEZ, M. J. SUAREZ & R. J. C.
CANTET. 2020. Hydrological and Physico-chemical dynamics in two Andean
streams. *Limnetica*, 39 (1): 17-33. DOI: 10.23818/limn.39.02

Number/Site	Population (inhabitants km ⁻²)	Nutrient export (kg km ⁻² y ⁻¹)				Chlorophyll (mg m ⁻²)
		Nh ₄	NO ₂ + NO ₃	DIN	SRP	
1) Str. Torrontegui	0	5.0	0.3	5.3	1.08	7
2) Str. Melgarejo	0	nd	0.2	0.2	1.27	7
3) Str. Pescadero	4	nd	nd	nd	0.08	117
4) Str. Loncochinoco	45	1.5	162.2	163.7	3.97	52
5) Str. Cascada	13	nd	0.2	0.2	0.06	77
6) Gutiérrez rise	7	nd	0.1	0.1	0.92	7
7) Gutiérrez 7	19	0.3	3.7	4.0	1.83	66
8) Gutiérrez 8	18	1.4	3.6	5.0	0.55	12
9) Gutiérrez 9	23	2.2	59.9	62.1	2.03	4
10) Gutiérrez mouth	28	1.0	83.1	84.1	1.02	29

Table 1 annexed Data supporting Figures 1 and 2. Population density and limnological characteristics of the study sites (Str.) Streams. Ammonium (Nh₄); Nitrates + Nitrites (NO₃ + NO₂); Soluble reactive phosphorus (SRP). Nd: not detectable.