PhyKIT: a UNIX shell toolkit for processing and analyzing phylogenomic data

Jacob L. Steenwyk^{1,*}, Thomas J. Buida III², Abigail L. Labella¹, Yuanning Li¹, Xing-Xing Shen³, & Antonis Rokas^{1,*}

¹ Vanderbilt University, Department of Biological Sciences, VU Station B #35-1634, Nashville, TN 37235, United States of America

² 9 City Place #312, Nashville, TN 37209, United States of America

³ Ministry of Agriculture Key Lab of Molecular Biology of Crop Pathogens and Insects, Institute of Insect Sciences, Zhejiang University, Hangzhou 310058, China

*Correspondence should be addressed to: jacob.steenwyk@vanderbilt.edu or antonis.rokas@vanderbilt.edu

ORCiDs

J.L. Steenwyk: 0000-0002-8436-595X

T.J. Buida III: 0000-0001-9367-6189

Abigail L. Labella: 0000-0003-0068-6703

Y. Li: 0000-0002-2206-5804

X.-X. Shen: 0000-0001-5765-1419

A. Rokas: 0000-0002-7248-6551

Running title: PhyKIT: a toolkit for examining phylogenomic datasets

Keywords: molecular phylogenetics, phylogenomics, multiple sequence alignment, phylogenetic signal, network, gene-gene covariation, evolutionary rate, polytomy, tree distance

Abstract

Diverse disciplines in biology process and analyze multiple sequence alignments (MSAs) and phylogenetic trees to evaluate their information content, infer evolutionary events and processes, and predict gene function. However, automated processing of MSAs and trees remains a challenge due to the lack of a unified toolkit. To fill this gap, we introduce PhyKIT, a toolkit for the UNIX shell environment with 30 functions that process MSAs and trees, including but not limited to estimation of mutation rate, evaluation of sequence composition biases, calculation of the degree of violation of a molecular clock, and collapsing bipartitions (internal branches) with low support. To demonstrate the utility of PhyKIT, we detail three use cases: (1) summarizing information content in MSAs and phylogenetic trees for diagnosing potential biases in sequence or tree data; (2) evaluating gene-gene covariation of evolutionary rates to identify functional relationships, including novel ones, among genes; and (3) identify lack of resolution events or polytomies in phylogenetic trees, which are suggestive of rapid radiation events or lack of data. We anticipate PhyKIT will be useful for processing, examining, and deriving biological meaning from increasingly large phylogenomic datasets. PhyKIT is freely available on GitHub (<https://github.com/JLSteenwyk/PhyKIT>) and documentation including user tutorials are available online (<https://jlsteenwyk.com/PhyKIT>).

Introduction

Multiple sequence alignments (MSAs) and phylogenetic trees are widely used in numerous disciplines, including bioinformatics, evolutionary biology, molecular biology, and structural biology. As a result, the development of user-friendly software that enables biologists to process and analyze MSAs and phylogenetic trees is an active area of research (Kapli *et al.* 2020).

In recent years, numerous methods have proven useful for diagnosing potential biases and inferring biological events in genome-scale phylogenetic (or phylogenomic) datasets. For example, methods that evaluate sequence composition biases in MSAs (Phillips and Penny 2003), signatures of clock-like evolution in phylogenetic trees (Liu *et al.* 2017), phylogenetic treeness (Lanyon 1988; Phillips and Penny 2003), taxa whose long branches may cause variation in their placement on phylogenetic trees (Struck 2014), and others have assisted in summarizing the information content in phylogenomic datasets and improved phylogenetic inference (Felsenstein 1978; Philippe *et al.* 2011; Salichos and Rokas 2013; Doyle *et al.* 2015; Liu *et al.* 2017; Smith *et al.* 2018; Walker *et al.* 2019).

Other methodological innovations include identifying significant gene-gene covariation of evolutionary rate, which has been shown to accurately and sensitively identify genes that have shared functions, are co-expressed, and/or are part of the same multimeric complexes (Sato *et al.* 2005; Clark *et al.* 2012). Furthermore, gene-gene covariation serves as a powerful evolution-based genetic screen for predicting gene function (Brunette *et al.* 2019). Lastly, a recently developed method has enabled the identification of unresolved internal branches or polytomies in species trees (Sayyari and Mirarab 2018; One Thousand Plant Transcriptomes Initiative 2019);

such branches can stem from rapid radiation events or from lack of data (Rokas and Carroll 2006).

Despite the wealth of information in MSAs and phylogenetic trees, there is a dearth of tools, especially ones that allow to conduct these analyses in a unified framework. For example, to utilize the functions mentioned in the previous paragraphs, a combination of web-server applications, ‘hard-coded’ scripts available through numerous repositories and supplementary material, standalone software, and/or extensive programming in languages including R, Python, or C is currently required (Cock *et al.* 2009; Junier and Zdobnov 2010; Revell 2012; Talevich *et al.* 2012; Struck 2014; Kück and Longo 2014; Wolfe and Clark 2015; One Thousand Plant Transcriptomes Initiative 2019). As a result, integrating these functions into bioinformatic pipelines is challenging, reducing their accessibility to the scientific community.

To facilitate the integration of these methods into bioinformatic pipelines, we introduce PhyKIT, a UNIX shell toolkit with 30 functions (Table 1) with broad utility for analyzing and processing MSAs and phylogenetic trees. Current functions implemented in PhyKIT include measuring topological similarity of phylogenetic trees, creating codon-based MSAs, concatenating sets of MSAs into phylogenomic datasets, editing and/or viewing alignments and phylogenetic trees, and identifying putatively spurious homologs in MSAs. We highlight three uses of PhyKIT: (1) calculating diverse statistics that summarize the information content and potential biases (e.g., sequence- or phylogeny-based biases) in MSAs and phylogenetic trees; (2) creating a gene-gene covariation network of evolutionary rates; and (3) inferring the presence of polytomies from

phylogenomic data. The diverse functions implemented in PhyKIT will likely be of interest to bioinformaticians, molecular biologists, evolutionary biologists, and others.

Materials and Methods

PhyKIT is a command line tool for the UNIX shell environment written in the Python programming language (<https://www.python.org/>). PhyKIT requires few dependencies (Biopython (Cock *et al.* 2009) and SciPy (Virtanen *et al.* 2020)) making it user-friendly to install and integrate into existing bioinformatic pipelines. Furthermore, the online documentation of PhyKIT comes complete with tutorials that detail how to use various functions. Lastly, PhyKIT is modularly designed to allow straightforward integration of additional functions in future versions.

PhyKIT has 30 different functions that help process and analyze MSAs and phylogenetic trees (Table 1). The 30 functions can be grouped into broad categories that assist in conducting analyses of MSAs and phylogenies or in processing/editing them. For example, “analysis” functions help examine information content biases, gene-gene covariation, and polytomies in phylogenomic datasets; “processing/editing” functions help prune tips from phylogenies, collapse poorly supported bipartitions in phylogenetic trees, concatenate sets of MSAs into a single data matrix, or create codon-based alignments from protein alignments and their corresponding nucleotide sequences.

Detailed information about each one of PhyKIT’s functions and tutorials for using the software can be found in the online documentation (<https://jlsteenwyk.com/PhyKIT>). Here, we focus on

three specific groups of functions implemented in PhyKIT that enable researchers to summarize information content in phylogenomic datasets, create gene-gene evolutionary rate covariation networks, and identifying polytomies in phylogenomic data.

Evaluating information content and biases in phylogenomic datasets

MSAs and phylogenetic trees are frequently examined to evaluate their information content and potential biases in characteristics such as sequence composition or branch lengths (Phillips and Penny 2003; Philippe *et al.* 2011; Struck 2014; Doyle *et al.* 2015; Shen *et al.* 2016a; Liu *et al.* 2017; Smith *et al.* 2018). PhyKIT implements numerous functions for doing so. Here, we demonstrate the application of 14 functions:

(1) *Alignment length*. The length of a multiple sequence alignment, which is associated with robust bipartition support and tree accuracy (Shen *et al.* 2016a; Walker *et al.* 2019);

(2) *Alignment length with no gaps*. The length of a multiple sequence alignment after excluding sites with gaps, which is associated with robust bipartition support and tree accuracy (Shen *et al.* 2016a);

(3) *Degree of violation of a molecular clock (DVMC)*. A metric used to determine the clock-like evolution of a gene using the standard deviation of branch lengths for a single gene tree (Liu *et al.* 2017). DVMC is calculated using the following formula:

$$DVMC = \sqrt{\frac{1}{N-1} \sum_{j=1}^N (i_j - \bar{i})^2}$$

where N represents the number of tips in a phylogenetic tree, i_j being the distance between the root of the tree and species j , and $\bar{i}$ represents the average root to tip distance. DVMC can be used to identify genes with clock-like evolution for divergence time estimation (Liu *et al.* 2017);

(4) *Internal branch lengths*. Summary statistics of internal branch lengths in a phylogenetic tree are reported including mean, median, 25th percentile, 75th percentile, minimum, maximum, standard deviation, and variance values. Examination of internal branch lengths is useful in evaluating phylogenetic tree shape;

(5) *Long branch score*. A metric that examines the degree of taxon-specific long branch attraction (Struck 2014; Weigert *et al.* 2014). Long branch scores of individual taxa are calculated using the following formula:

$$LB_i = \left(\frac{\overline{PD}_i}{\overline{PD}_{all}} - 1 \right) \times 100$$

where $\overline{PD}_i$ represents the average pairwise patristic distance of taxon i to all other taxa, $\overline{PD}_{all}$ represents the average patristic distance across all taxa, and LB_i represents the long branch score of taxon i . Long branch scores can be used to evaluate heterogeneity in tip-to-root distances and identify taxa that may be susceptible to long branch attraction;

(6) *Pairwise identity*. Pairwise identity is a crude approximation of the evolutionary rate of a gene and is calculated by determining the average number of sites in an MSA that are the same character state between all pairwise combinations of taxa. This can be used to group genes based on their evolutionary rates (e.g., faster-evolving genes vs. slower-evolving ones) (Chen *et al.* 2017);

(7) *Patristic distances*. Patristic distances refer to all distances between all pairwise combinations of tips in a phylogenetic tree (Fourment and Gibbs 2006), which can be used to evaluate the rate of evolution in gene trees or taxon sampling density in species trees;

(8) *Parsimony-informative sites*. Parsimony-informative sites are those sites in an MSA that have a least two character states (excluding gaps) that occur at least twice (Kumar *et al.* 2016); the

number of parsimony-informative sites is associated with robust bipartition support and tree accuracy (Shen *et al.* 2016a; Steenwyk *et al.* 2020);

(9) *Variable sites*. Variable sites are those sites in an MSA that contain at least two different character states (excluding gaps) (Kumar *et al.* 2016); the number of variable sites is associated with robust bipartition support and tree accuracy (Shen *et al.* 2016a);

(10) *Relative composition variability*. Relative composition variability is the average variability in the sequence composition among taxa in an MSA. Relative composition variability is calculated using the following formula:

$$Relative\ composition\ variability = \sum_{i=1}^c \sum_{j=1}^n \frac{|c_{ij} - \bar{c}_i|}{s \times n}$$

where c is the number of different character states per sequence type, n is the number of taxa in an MSA, c_{ij} is the number of occurrences of the i th character state for the j th taxon, $\bar{c}_i$ is the average number of the i th c character state across n taxa, and s refers to the total number of sites (characters) in an MSA. Relative composition variability can be used to evaluate potential sequence composition biases in MSAs, which in turn violate assumptions of site composition homogeneity in standard models of sequence evolution (Phillips and Penny 2003);

(11) *Saturation*. Saturation refers to when an MSA contains many sites that have experienced multiple substitutions in individual taxa. Saturation is estimated from the slope of the regression line between patristic distances and pairwise identities. Saturated MSAs have reduced phylogenetic information and can result in issues of long branch attraction (Lake 1991; Philippe *et al.* 2011);

(12) *Total tree length*. Total tree length refers to the sum of internal and terminal branch lengths and is calculated using the following formula:

$$total\ tree\ length = \sum_{i=1}^a l_i + \sum_{j=1}^b l_j$$

175

176 Where l_i is the branch length of the i th branch of a internal branches and l_j is the branch length of
177 the j th branch of b terminal branches. Total tree length measures the inferred total amount or rate
178 of evolutionary change in a phylogenetic tree;

179 (13) *Treeness*. Treeness (also referred to as stemminess) is a measure of the inferred relative
180 amount or rate of evolutionary change that has taken place on internal branches of a phylogenetic
181 tree (Lanyon 1988; Phillips and Penny 2003) and is calculated using the following formula:

$$treeness = \sum_{u=1}^b \frac{l_u}{l_t}$$

182 where l_u is the branch length of the u th branch of b internal branches, and l_t refers to the total
183 branch length of the phylogenetic tree. Treeness can be used to evaluate how much of the total
184 tree length is observed among internal branches;

185 (14) *Treeness divided by relative composition variability*. This function combines two metrics to
186 measure both composition bias and other biases that may negatively influence phylogenetic
187 inference. High treeness divided by relative composition variability values have been shown to
188 be less susceptible to sequence composition biases and are associated with robust bipartition
189 support and tree accuracy (Phillips and Penny 2003; Shen *et al.* 2016a).

190

191 **Calculating gene-gene evolutionary rate covariation or coevolution**

192 Genes that share similar rates of evolution through speciation events (or coevolve) tend to have
193 similar functions, expression levels, or are parts of the same multimeric complexes (Sato *et al.*
194 2005; Clark *et al.* 2012). Thus, identifying significant coevolution between genes (i.e.,

identifying genes that are significantly correlated in their evolutionary rates across speciation events) can be a powerful evolution-based screen to determine gene function (Brunette *et al.* 2019).

To measure gene-gene evolutionary rate covariation, PhyKIT implements the mirror tree method (Pazos and Valencia 2001; Sato *et al.* 2005), which examines whether two trees have correlated branch lengths. Specifically, PhyKIT calculates the Pearson correlation coefficient between branch lengths in two phylogenetic trees that share the same tips and topology. To account for differences in taxon representation between the two trees, PhyKIT first automatically determines which taxa are shared and prunes one or both such that the same set of taxa is present in both trees. PhyKIT requires that the two input trees have the same topology, which is typically the species tree topology inferred from whole genome or proteome data. Thus, the user will typically first estimate a gene's branch lengths by constraining the topology to match that of the species tree. When running this function, users should be aware that many biological factors, such as horizontal transfer (Doolittle and Baptiste 2007), incomplete lineage sorting (Degnan and Salter 2005), and introgression / hybridization (Sang and Zhong 2000), can lead to gene histories that deviate from the species tree. In these cases, constraining a gene's history to match that of a species may lead to errors in the covariation analysis.

Due to factors including time since speciation and mutation rate, correlations between uncorrected branch lengths result in a high frequency of false positive correlations (Sato *et al.* 2005; Clark *et al.* 2012; Chikina *et al.* 2016). To ameliorate the influence of these factors, PhyKIT first transforms branch lengths into relative rates. To do so, branch lengths are corrected

by dividing the branch length in the gene tree by the corresponding branch length in the species tree. Previous work revealed that one or a few outlier branch length values can be responsible for false positive correlations and should be removed prior to analysis (Clark *et al.* 2012). Thus, PhyKIT removes outlier data points defined as having corrected branch lengths greater than five (i.e., removing gene tree branch lengths that are five or more times greater than their corresponding species tree branch lengths). Lastly, values are converted into relative rates using a Z-transformation. The resulting relative rates are used when calculating Pearson correlation coefficients.

Identifying polytomies in phylogenomic data

Rapid radiations or diversification events have occurred throughout the tree of life including among mammals, birds, plants, and fungi (Jarvis *et al.* 2014; Liu *et al.* 2017; One Thousand Plant Transcriptomes Initiative 2019; Li *et al.* 2020). Polytomies correspond to internal branches whose length is 0 (or statistically indistinguishable from 0) and can be driven either by biological (e.g., rapid radiations) or analytical (e.g., low amount of data) factors. Thus, polytomies are useful for inferring rapid radiation or diversification events and exploring incongruence in phylogenies (Sayyari and Mirarab 2018; One Thousand Plant Transcriptomes Initiative 2019; Li *et al.* 2020).

To identify polytomies, a modified approach to a previous strategy was implemented (Sayyari and Mirarab 2018). More specifically, the support for three alternative topologies is calculated among all gene trees from a phylogenomic dataset. For example, in species tree $((A,B),C), D)$; if examining the presence of a polytomy at the ancestral bipartition of tips A, B , and C , PhyKIT

will determine the number of gene trees that support $((A,B),C)$, $((A,C),B)$, and $((B,C),A)$; using the rooted gene trees provided by the user. Equal support for the three topologies (i.e., the presence of a polytomy) among a set of gene trees is assessed using a Chi-squared test. Failing to reject the null hypothesis is indicative of a polytomy (Sayyari and Mirarab 2018). Note that this approach is distinct from the approach of Sayyari and Mirarab to identify polytomies because PhyKIT uses a gene-based signal rather than a quartet-based signal. The difference between the two methods is that each gene contributes equally to the inference of a polytomy when a gene-based signal is used, whereas genes with greater taxon representation (which contain a greater number of quartets) will contribute a greater signal during polytomy identification when a quartet-based signal is used.

Results and Discussion

We outline three example uses of PhyKIT: 1) summarizing information content and identifying potential biases in animal, plant, yeast, and filamentous fungal phylogenomic datasets (Shen *et al.* 2016b; Steenwyk *et al.* 2019; Laumer *et al.* 2019; One Thousand Plant Transcriptomes Initiative 2019), 2) constructing a network of significant gene-gene covariation, which reveals genes of shared functions from empirical data spanning ~550 million years of evolution among fungi (Shen *et al.* 2020), and 3) illustrating how to identify polytomies using simulated and empirical data (Steenwyk *et al.* 2019).

Summarizing information content and biases in phylogenomic data

Examining information content in phylogenomic datasets can help diagnose potential biases that stem from low signal-to-noise ratios, multiple substitutions, non-clocklike evolution, and other

biological or analytical factors. To demonstrate the utility of PhyKIT to summarize the information content in phylogenomic datasets, we calculated 14 different metrics known to help diagnose potential biases in phylogenomic datasets or be associated with accurate and well supported phylogenetic inferences (Felsenstein 1978; Phillips and Penny 2003; Philippe *et al.* 2011; Struck 2014; Doyle *et al.* 2015; Shen *et al.* 2016a; Liu *et al.* 2017; Smith *et al.* 2018) using four empirical phylogenomic datasets from animals (201 tips; 2,891 genes) (Laumer *et al.* 2019), budding yeast (332 taxa; 2,408 genes) (Shen *et al.* 2018), filamentous fungi (93 taxa; 1,668 genes) (Steenwyk *et al.* 2019), and plants (1,124 taxa; 403 genes) (One Thousand Plant Transcriptomes Initiative 2019) (Figure 1, Table 1).

Examination of the distributions of the values of the 14 different metrics revealed inter- and intra-dataset heterogeneity (Figure 1). For example, inter-dataset heterogeneity was observed among animal and plant datasets, which had the lowest and highest average pairwise identity across alignments, respectively; intra-dataset heterogeneity was observed in the uniform distribution of pairwise identities in the budding yeast datasets. Similarly, inter-dataset heterogeneity was observed in estimates of saturation where the budding yeast and filamentous fungal MSAs were less saturated by multiple substitutions than the plant and animal datasets; intra-data heterogeneity was also observed in all four datasets. Varying degrees of inter- and intra-dataset heterogeneity was observed for other information content statistics, which may be due biological (e.g., mutation rate) or analytical factors (e.g., taxon sampling, distinct alignment, trimming, and tree inference strategies).

In summary, PhyKIT is useful for examining the information content of phylogenomic datasets. For example, the generation of different phylogenomic data submatrices by selecting subsets of genes or taxa with certain properties (e.g., retention of genes with the highest numbers of parsimony-informative sites or following removal of taxa with high long branch scores) can facilitate the exploration of the robustness of species tree inference or estimating time since divergence (Salichos and Rokas 2013; Liu *et al.* 2017; Shen *et al.* 2018, 2020; Steenwyk *et al.* 2019; Walker *et al.* 2019; Li *et al.* 2020).

A network of gene-gene covariation reveals neighborhoods of genes with shared function

Genes with similar evolutionary histories often have shared functions, are co-expressed, or are parts of the same multimeric complexes (Sato *et al.* 2005; Clark *et al.* 2012). Using PhyKIT, we examined gene-gene covariation using 815 genes spanning 1,107 genomes and ~563 million years of evolution among fungi (Shen *et al.* 2020). By examining 331,705 pairwise combinations of genes, we found 298 strong signatures of gene-gene covariation (defined as $r > 0.825$). The two genes with the strongest signatures of covariation were *SEC7* and *TAO3* ($r = 0.87$), suggesting that their protein products have similar or shared functions. Supporting this hypothesis, Sec7p contributes to cell-surface growth in the model yeast *Saccharomyces cerevisiae* (Novick and Schekman 1979) and genes with the Sec7 domain are transcriptionally coregulated with yeast-hyphal switches in the human pathogen *C. albicans* (Song *et al.* 2008). Similarly, Tao3p in both *S. cerevisiae* and *C. albicans* is part of a RAM signaling network, which controls hyphal morphogenesis, polarized growth, and cell-cycle related processes including cell separation, cell proliferation, and phase transitions (Bogomolnaya *et al.* 2006; Song *et al.* 2008).

Complex relationships of gene-gene covariation can be visualized as a network (Figure 2). Examination of network neighborhoods identified groups of genes that have shared functions and are parts of the same multimeric complexes. For example, the proteins encoded by *NDC80* and *NUF2* are part of the same kinetochore-associated complex termed the NDC80 complex—which is required for efficient mitosis (Sundin *et al.* 2011)—and significantly covary with one another ($r = 0.84$). Similarly, multiple genes that encode proteins involved in DNA replication and repair (i.e., *POL2*, *MSH6*, *RAD26*, *CDC9*, and *EXO1*) were part of the same network neighborhood, consistent with previous work suggesting an intimate interplay between DNA replication and multiple DNA repair pathways (Tsubouchi and Ogawa 2000; Lujan *et al.* 2012; Boiteux and Jinks-Robertson 2013). Similarly, network neighborhoods of genes involved in ribosome biogenesis, Golgi apparatus-related transport, and control of DNA replication were identified (Figure 2).

Taken together, these results indicate PhyKIT is a useful tool for evaluating gene-gene covariation and predicting genes' functions (Sato *et al.* 2005; Clark *et al.* 2012; Brunette *et al.* 2019). Thus, we anticipate PhyKIT will be helpful for evaluating gene-gene covariation and conducting evolution-based screens for gene functions across the tree of life.

Identifying polytomies in phylogenomic datasets

Rapid radiations or diversification events have occurred throughout the tree of life (Jarvis *et al.* 2014; Liu *et al.* 2017; One Thousand Plant Transcriptomes Initiative 2019; Li *et al.* 2020). One approach to identifying rapid radiations is by testing for the existence of polytomies in species

trees (Sayyari and Mirarab 2018; One Thousand Plant Transcriptomes Initiative 2019; Li *et al.* 2020). Polytomies can also arise when the amount of data at hand is insufficient for resolution (Walsh *et al.* 1999). To demonstrate the utility of PhyKIT to identify polytomies, we tested it using a simulated set of phylogenies that had a branch whose length was extremely small (Figure 3A). We found that PhyKIT was able to conservatively identify the simulated polytomy. Our results demonstrate PhyKIT can accurately identify a polytomy and provide further support that equal support among alternative topologies can be used as a means to identify a rapid radiation.

We next examined if there is evidence of polytomies in the evolutionary history of filamentous fungi from the genera *Aspergillus* and *Penicillium*. We examined three branches. The first two branches—one dating back ~110 million years ago (Figure 3B), and another dating back ~25 million years ago (Figure 3C)—were not polytomies. In contrast, examination of a ~60 million-year-old branch involving *Lanata-divaricata*, *Citrina*, and *Exilicaulis* (Figure 3D), which are major lineages (or sections) in the genus *Penicillium*, was consistent with a polytomy. Given the large number of gene trees used in our analysis (n=1,668), these results are consistent with a rapid radiation or diversification event in the history of *Penicillium* species.

In summary, these results suggest that PhyKIT is useful in identifying polytomies in simulated and empirical datasets. PhyKIT can also be useful for exploring incongruence in phylogenies by calculating gene support frequencies for alternative topologies. Calculations of gene-based support for various topologies can be used in diverse applications, including identifying putative introgression / hybridization events and conducting phylogenetically-based genome-wide association (PhyloGWAS) studies (Pease *et al.* 2016; Steenwyk *et al.* 2019).

355

356 **Conclusion**

357 We have developed PhyKIT, a comprehensive toolkit for processing and analyzing MSAs and
358 trees in phylogenomic datasets. PhyKIT is freely available on GitHub

359 (<https://github.com/JLSteenwyk/PhyKIT>) with extensive documentation and user tutorials

360 (<https://jlsteenwyk.com/PhyKIT>). PhyKIT is a fast and flexible toolkit for the UNIX shell

361 environment, which allows it to be easily integrated into bioinformatic pipelines. We anticipate

362 PhyKIT will be of interest to biologists from diverse disciplines and with varying degrees of

363 experience in analyzing MSAs and phylogenies. In particular, PhyKIT will likely be helpful in

364 addressing one of the greatest challenges in biology, building, understanding, and deriving

365 meaning from the tree of life.

366

367 **Data Availability**

368 All data used will become available in figshare (doi: 10.6084/m9.figshare.13118600) upon
369 publication.

370

371 **Acknowledgements**

372 We thank the Rokas lab for helpful discussion and feedback. J.L.S. and A.R. were funded by the

373 Howard Hughes Medical Institute through the James H. Gilliam Fellowships for Advanced

374 Study program. X.X.S. was supported by National Natural Science Foundation of China (No.

375 32071665) and the Fundamental Research Funds for the Central Universities (No.

376 2020QNA6019).

377

References

- Bogomolnaya L. M., R. Pathak, J. Guo, and M. Polymenis, 2006 Roles of the RAM signaling network in cell cycle progression in *Saccharomyces cerevisiae*. *Curr. Genet.* 49: 384–92.
<https://doi.org/10.1007/s00294-006-0069-y>
- Boiteux S., and S. Jinks-Robertson, 2013 DNA Repair Mechanisms and the Bypass of DNA Damage in *Saccharomyces cerevisiae*. *Genetics* 193: 1025–1064.
<https://doi.org/10.1534/genetics.112.145219>
- Brunette G. J., M. A. Jamalruddin, R. A. Baldock, N. L. Clark, and K. A. Bernstein, 2019 Evolution-based screening enables genome-wide prioritization and discovery of DNA repair genes. *Proc. Natl. Acad. Sci.* 116: 19593–19599. <https://doi.org/10.1073/pnas.1906559116>
- Chen M.-Y., D. Liang, and P. Zhang, 2017 Phylogenomic Resolution of the Phylogeny of Laurasiatherian Mammals: Exploring Phylogenetic Signals within Coding and Noncoding Sequences. *Genome Biol. Evol.* 9: 1998–2012. <https://doi.org/10.1093/gbe/evx147>
- Chikina M., J. D. Robinson, and N. L. Clark, 2016 Hundreds of Genes Experienced Convergent Shifts in Selective Pressure in Marine Mammals. *Mol. Biol. Evol.* 33: 2182–2192.
<https://doi.org/10.1093/molbev/msw112>
- Clark N. L., E. Alani, and C. F. Aquadro, 2012 Evolutionary rate covariation reveals shared functionality and coexpression of genes. *Genome Res.* 22: 714–720.
<https://doi.org/10.1101/gr.132647.111>
- Cock P. J. A., T. Antao, J. T. Chang, B. A. Chapman, C. J. Cox, *et al.*, 2009 Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics* 25: 1422–1423. <https://doi.org/10.1093/bioinformatics/btp163>
- Degnan J. H., and L. A. Salter, 2005 Gene tree distributions under the coalescent process.

401 Evolution (N. Y). 59: 24–37. <https://doi.org/10.1111/j.0014-3820.2005.tb00891.x>

402 Doolittle W. F., and E. Bapteste, 2007 Pattern pluralism and the Tree of Life hypothesis. Proc.

403 Natl. Acad. Sci. 104: 2043–2049. <https://doi.org/10.1073/pnas.0610699104>

404 Doyle V. P., R. E. Young, G. J. P. Naylor, and J. M. Brown, 2015 Can We Identify Genes with

405 Increased Phylogenetic Reliability? Syst. Biol. 64: 824–837.

406 <https://doi.org/10.1093/sysbio/syv041>

407 Felsenstein J., 1978 Cases in which Parsimony or Compatibility Methods will be Positively

408 Misleading. Syst. Biol. 27: 401–410. <https://doi.org/10.1093/sysbio/27.4.401>

409 Fourment M., and M. J. Gibbs, 2006 PATRISTIC: a program for calculating patristic distances

410 and graphically comparing the components of genetic change. BMC Evol. Biol. 6: 1.

411 <https://doi.org/10.1186/1471-2148-6-1>

412 Hunter J. E., and S. H. Cohen, 2007 Package: igraph. Educ. Psychol. Meas.

413 <https://doi.org/10.1177/001316446902900315>

414 Jarvis E. D., S. Mirarab, A. J. Aberer, B. Li, P. Houde, *et al.*, 2014 Whole-genome analyses

415 resolve early branches in the tree of life of modern birds. Science (80-.). 346: 1320–1331.

416 <https://doi.org/10.1126/science.1253451>

417 Junier T., and E. M. Zdobnov, 2010 The Newick utilities: high-throughput phylogenetic tree

418 processing in the UNIX shell. Bioinformatics 26: 1669–1670.

419 <https://doi.org/10.1093/bioinformatics/btq243>

420 Kapli P., Z. Yang, and M. J. Telford, 2020 Phylogenetic tree building in the genomic age. Nat.

421 Rev. Genet. <https://doi.org/10.1038/s41576-020-0233-0>

422 Kück P., and G. C. Longo, 2014 FASconCAT-G: extensive functions for multiple sequence

423 alignment preparations concerning phylogenetic studies. Front. Zool. 11: 81.

<https://doi.org/10.1186/s12983-014-0081-x>

Kumar S., G. Stecher, and K. Tamura, 2016 MEGA7: Molecular Evolutionary Genetics Analysis

Version 7.0 for Bigger Datasets. Mol. Biol. Evol. <https://doi.org/10.1093/molbev/msw054>

Lake J. A., 1991 The order of sequence alignment can bias the selection of tree topology. Mol.

Biol. Evol. <https://doi.org/10.1093/oxfordjournals.molbev.a040654>

Lanyon S. M., 1988 The Stochastic Mode of Molecular Evolution: What Consequences for

Systematic Investigations? Auk 105: 565–573. <https://doi.org/10.1093/auk/105.3.565>

Laumer C. E., R. Fernández, S. Lemer, D. Combosch, K. M. Kocot, *et al.*, 2019 Revisiting

metazoan phylogeny with genomic sampling of all phyla. Proc. R. Soc. B Biol. Sci. 286:

20190831. <https://doi.org/10.1098/rspb.2019.0831>

Li Y., J. L. Steenwyk, Y. Chang, Y. Wang, T. Y. James, *et al.*, 2020 A genome-scale phylogeny

of Fungi; insights into early evolution, radiations, and the relationship between taxonomy

and phylogeny. bioRxiv 2020.08.23.262857. <https://doi.org/10.1101/2020.08.23.262857>

Liu L., J. Zhang, F. E. Rheindt, F. Lei, Y. Qu, *et al.*, 2017 Genomic evidence reveals a radiation

of placental mammals uninterrupted by the KPg boundary. Proc. Natl. Acad. Sci. 114:

E7282–E7290. <https://doi.org/10.1073/pnas.1616744114>

Lujan S. A., J. S. Williams, Z. F. Pursell, A. A. Abdulovic-Cui, A. B. Clark, *et al.*, 2012

Mismatch Repair Balances Leading and Lagging Strand DNA Replication Fidelity, (C. E.

Pearson, Ed.). PLoS Genet. 8: e1003016. <https://doi.org/10.1371/journal.pgen.1003016>

Novick P., and R. Schekman, 1979 Secretion and cell-surface growth are blocked in a

temperature-sensitive mutant of *Saccharomyces cerevisiae*. Proc. Natl. Acad. Sci. U. S. A.

76: 1858–62. <https://doi.org/10.1073/pnas.76.4.1858>

One Thousand Plant Transcriptomes Initiative, 2019 One thousand plant transcriptomes and the

phylogenomics of green plants. *Nature* 574: 679–685. <https://doi.org/10.1038/s41586-019-1693-2>

Pazos F., and A. Valencia, 2001 Similarity of phylogenetic trees as indicator of protein–protein interaction. *Protein Eng. Des. Sel.* 14: 609–614. <https://doi.org/10.1093/protein/14.9.609>

Pease J. B., D. C. Haak, M. W. Hahn, and L. C. Moyle, 2016 Phylogenomics Reveals Three Sources of Adaptive Variation during a Rapid Radiation, (D. Penny, Ed.). *PLOS Biol.* 14: e1002379. <https://doi.org/10.1371/journal.pbio.1002379>

Philippe H., H. Brinkmann, D. V. Lavrov, D. T. J. Littlewood, M. Manuel, *et al.*, 2011 Resolving Difficult Phylogenetic Questions: Why More Sequences Are Not Enough, (D. Penny, Ed.). *PLoS Biol.* 9: e1000602. <https://doi.org/10.1371/journal.pbio.1000602>

Phillips M. J., and D. Penny, 2003 The root of the mammalian tree inferred from whole mitochondrial genomes. *Mol. Phylogenet. Evol.* 28: 171–185. [https://doi.org/10.1016/S1055-7903\(03\)00057-5](https://doi.org/10.1016/S1055-7903(03)00057-5)

Revell L. J., 2012 phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* 3: 217–223. <https://doi.org/10.1111/j.2041-210X.2011.00169.x>

Robinson D. F., and L. R. Foulds, 1981 Comparison of phylogenetic trees. *Math. Biosci.* 53: 131–147. [https://doi.org/10.1016/0025-5564\(81\)90043-2](https://doi.org/10.1016/0025-5564(81)90043-2)

Rokas A., and S. B. Carroll, 2006 Bushes in the Tree of Life. *PLoS Biol.* 4: e352. <https://doi.org/10.1371/journal.pbio.0040352>

Salichos L., and A. Rokas, 2013 Inferring ancient divergences requires genes with strong phylogenetic signals. *Nature* 497: 327–331. <https://doi.org/10.1038/nature12130>

Sang T., and Y. Zhong, 2000 Testing Hybridization Hypotheses Based on Incongruent Gene

Trees, (R. Olmstead, Ed.). Syst. Biol. 49: 422–434.
<https://doi.org/10.1080/10635159950127321>

Sato T., Y. Yamanishi, M. Kanehisa, and H. Toh, 2005 The inference of protein-protein interactions by co-evolutionary analysis is improved by excluding the information about the phylogenetic relationships. Bioinformatics 21: 3482–3489.
<https://doi.org/10.1093/bioinformatics/bti564>

Sayyari E., and S. Mirarab, 2018 Testing for Polytomies in Phylogenetic Species Trees Using Quartet Frequencies. Genes (Basel). 9. <https://doi.org/10.3390/genes9030132>

Shen X.-X., L. Salichos, and A. Rokas, 2016a A Genome-Scale Investigation of How Sequence, Function, and Tree-Based Gene Properties Influence Phylogenetic Inference. Genome Biol. Evol. 8: 2565–2580. <https://doi.org/10.1093/gbe/evw179>

Shen X.-X., X. Zhou, J. Kominek, C. P. Kurtzman, C. T. Hittinger, *et al.*, 2016b Reconstructing the Backbone of the Saccharomycotina Yeast Phylogeny Using Genome-Scale Data. G3 Genes|Genomes|Genetics 6: 3927–3939. <https://doi.org/10.1534/g3.116.034744>

Shen X.-X., D. A. Opulente, J. Kominek, X. Zhou, J. L. Steenwyk, *et al.*, 2018 Tempo and Mode of Genome Evolution in the Budding Yeast Subphylum. Cell 175: 1533-1545.e20.
<https://doi.org/10.1016/j.cell.2018.10.023>

Shen X.-X., J. L. Steenwyk, A. L. Labella, D. A. Opulente, X. Zhou, *et al.*, 2020 Genome-scale phylogeny and contrasting modes of genome evolution in the fungal phylum Ascomycota. bioRxiv. <https://doi.org/10.1101/2020.05.11.088658>

Smith S. A., J. W. Brown, and J. F. Walker, 2018 So many genes, so little time: A practical approach to divergence-time estimation in the genomic era, (H. Escriva, Ed.). PLoS One 13: e0197433. <https://doi.org/10.1371/journal.pone.0197433>

493 Song Y., S. A. Cheon, K. E. Lee, S.-Y. Lee, B.-K. Lee, *et al.*, 2008 Role of the RAM network in
 494 cell polarity and hyphal morphogenesis in *Candida albicans*. *Mol. Biol. Cell* 19: 5456–77.
 495 <https://doi.org/10.1091/mbc.e08-03-0272>

496 Steenwyk J. L., X.-X. Shen, A. L. Lind, G. H. Goldman, and A. Rokas, 2019 A Robust
 497 Phylogenomic Time Tree for Biotechnologically and Medically Important Fungi in the
 498 Genera *Aspergillus* and *Penicillium*, (J. P. Boyle, Ed.). *MBio* 10.
 499 <https://doi.org/10.1128/mBio.00925-19>

500 Steenwyk J. L., T. J. Buida, Y. Li, X.-X. Shen, and A. Rokas, 2020 ClipKIT: a multiple sequence
 501 alignment-trimming algorithm for accurate phylogenomic inference. *bioRxiv*
 502 2020.06.08.140384. <https://doi.org/10.1101/2020.06.08.140384>

503 Struck T. H., 2014 TreSpEx—Detection of Misleading Signal in Phylogenetic Reconstructions
 504 Based on Tree Information. *Evol. Bioinforma.* 10: EBO.S14239.
 505 <https://doi.org/10.4137/EBO.S14239>

506 Sundin L. J. R., G. J. Guimaraes, and J. G. Deluca, 2011 The NDC80 complex proteins Nuf2 and
 507 Hec1 make distinct contributions to kinetochore-microtubule attachment in mitosis. *Mol.*
 508 *Biol. Cell* 22: 759–68. <https://doi.org/10.1091/mbc.E10-08-0671>

509 Talevich E., B. M. Invergo, P. J. Cock, and B. A. Chapman, 2012 Bio.Phylo: A unified toolkit
 510 for processing, analyzing and visualizing phylogenetic trees in Biopython. *BMC*
 511 *Bioinformatics* 13: 209. <https://doi.org/10.1186/1471-2105-13-209>

512 Tsubouchi H., and H. Ogawa, 2000 Exo1 Roles for Repair of DNA Double-Strand Breaks and
 513 Meiotic Crossing Over in *Saccharomyces cerevisiae*, (T. D. Fox, Ed.). *Mol. Biol. Cell* 11:
 514 2221–2233. <https://doi.org/10.1091/mbc.11.7.2221>

515 Virtanen P., R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, *et al.*, 2020 SciPy 1.0:

fundamental algorithms for scientific computing in Python. Nat. Methods.
<https://doi.org/10.1038/s41592-019-0686-2>

Walker J. F., N. Walker-Hale, O. M. Vargas, D. A. Larson, and G. W. Stull, 2019 Characterizing
gene tree conflict in plastome-inferred phylogenies. PeerJ 7: e7747.
<https://doi.org/10.7717/peerj.7747>

Walsh H. E., M. G. Kidd, T. Moum, and V. L. Friesen, 1999 Polytomies and the Power of
Phylogenetic Inference. Evolution (N. Y). 53: 932. <https://doi.org/10.2307/2640732>

Weigert A., C. Helm, M. Meyer, B. Nickel, D. Arendt, *et al.*, 2014 Illuminating the Base of the
Annelid Tree Using Transcriptomics. Mol. Biol. Evol. 31: 1391–1401.
<https://doi.org/10.1093/molbev/msu080>

Wolfe N. W., and N. L. Clark, 2015 ERC analysis: web-based inference of gene function via
evolutionary rate covariation: Fig. 1. Bioinformatics btv454.
<https://doi.org/10.1093/bioinformatics/btv454>

Figure 1. Summary of information content in four empirical phylogenomic datasets.

metrics implemented in PhyKIT help summarize the information content and identify potential biases in phylogenomic datasets. Each graph displays a violin plot with a black point representing the mean. Error bars indicate one standard error above and below the mean; however, these are difficult to see in nearly all graphs because they were often near the mean. Abbreviations are as follows: Aln. len.: alignment length; Aln. len. no gaps: alignment length excluding sites with gaps; DVMC: degree of violation of a molecular clock; Internal branch len.: average internal branch length; Patristic distances: average patristic distance in a gene tree;

541 Percent PI Sites: percentage of parsimony-informative sites in an MSA; Percent var. sites:
542 percentage of variable sites in an MSA; RCV: relative composition variability.

Figure 2. Gene-gene covariation network inferred from ~550 million years of evolution

across 1,107 fungi. A network of significant gene-gene coevolution identifies network

neighborhoods representative of associated functional categories. For example, the *NDC80* and

NUF2 genes (toward the top right of the network) were identified to be significantly coevolving

with one another ($r = 0.84$, $p < 0.01$, Pearson's correlation test); they both encode proteins that

are part of the same multimeric kinetochore-associated complex (green). Similarly, genes that are

DNA replication factors (orange), contribute to DNA replication and repair processes (yellow),

551 participate in Golgi apparatus-related transport (brown), or ribosome biogenesis (pink) were
552 found to be neighbors in the network. Network visualization was done with the igraph package,
553 v1.2.4.2 (Hunter and Cohen 2007), in R, v3.6.2 (<https://www.r-project.org/>).

554

Figure 3. Identifying polytomies from phylogenomic data. (Ai) A cladogram of a simulated species phylogeny with tip names A-G. The red branch has a very short branch length of 2×10^{-5} substitutions per site. (Aii) Phylogram of the same phylogeny shows that all other branches are much longer (≥ 1.0 substitutions per site). (Aiii) After reconstructing the evolutionary history from 1,000 alignments simulated from the phylogeny in Aii, the hypothesis of a polytomy was tested using gene support frequencies for three alternative rooted topologies defined by the clades of green, orange, and purple taxa. Failure to reject the null hypothesis of equal support among genes for each topology is indicative of a polytomy ($\chi^2 = 5.97$, p-value = 0.05, Chi-

564 squared test). (B-D) The same approach was then used to examine if there is evidence for a
 565 polytomy at three different branches in a phylogeny of filamentous fungi. (D) Support for a
 566 polytomy ($\chi^2 = 0.514$, p-value = 0.77, Chi-squared test) was observed for the relationships
 567 between three different sections of *Penicillium* fungi. These results demonstrate the utility of
 568 gene-support frequencies for evaluating polytomies and examining incongruence in
 569 phylogenomic datasets.
 570

571 **Table 1. Summary of 30 functions implemented in PhyKIT**

<u>Description</u>	<u>Name</u>	<u>Function Alias(es)</u>	<u>Type of function</u>	<u>Input data</u>	<u>Citation</u>
Calculate MSA length	alignment_length	aln_len; al	analytic	alignment	NA
Calculate MSA length after removing sites with gaps	alignment_length_no_gaps	aln_len_no_gaps; alng	analytic	alignment	NA
Combine numerous MSAs into one concatenated matrix	create_concatenation_matrix	create_concat; cc	processing/editing	alignment	NA
Calculate guanine/cytosine content in an MSA	gc_content	gc	analytic	alignment	NA
Calculate summary statistics* for pairwise identity among sequences in an MSA	pairwise_identity	pairwise_id; pi	analytic	alignment	(Chen <i>et al.</i> 2017)
Calculate the number and percentage of parsimony-informative sites in an MSA	parsimony_informative_sites	pis	analytic	alignment	NA
Calculate relative composition variability in an MSA	relative_composition_variability	rel_comp_var; rev	analytic	alignment	(Phillips and Penny 2003)
Rename FASTA entries in an MSA file	rename_fasta_entries	rename_fasta	processing/editing	alignment	NA
Thread DNA sequences over a protein MSA	thread_dna	pal2nal; p2n	processing/editing	alignment	NA
Calculate the number and percentage of variable sites in an MSA	variables_sites	vs	analytic	alignment	NA
Calculate summary statistics* for bipartition support values from a set of phylogenetic trees	bipartition_support_stats	bss	analytic	phylogeny	NA
Multiply all branch lengths of a phylogenetic tree by a specific number	branch_length_multiplier	blm	processing/editing	phylogeny	NA

Collapse bipartitions with bipartition support lower than a user-specified value in a phylogenetic tree	collapse_branches	collapse; cb	processing/editing	phylogeny	NA
Calculates the correlation of evolutionary rates between two genes trees with the same topology	covarying_evolutionary_rates	cover	analytic	phylogeny	(Clark <i>et al.</i> 2012)
Calculate the degree a gene tree violates a molecular clock-like rate of evolution	degree_of_violation_of_a_molecular_clock	dvmc	analytic	phylogeny	(Liu <i>et al.</i> 2017)
Calculate summary statistics* for internal branch lengths in a phylogenetic tree	internal_branch_stats	ibs	analytic	phylogeny	NA
Create numeric labels for bipartitions in a phylogenetic tree	internode_labeler	il	processing/editing	phylogeny	NA
Calculate summary statistics* for long branch scores for taxa in a phylogenetic tree	long_branch_score	lb_score; lbs	analytic	phylogeny	(Weigert <i>et al.</i> 2014)
Calculate the total length of a phylogenetic tree	total_tree_length	tree_len	analytic	phylogeny	NA
Calculate summary statistics* for all patristic distances (or all tip-to-tip distances) in a phylogenetic tree	patristic_distances	pd	analytic	phylogeny	(Fourment and Gibbs 2006)
Use gene support frequencies to test for the existence of a polytomy	polytomy_test	polyt_test; polyt; ptt	analytic	phylogeny	(Sayyari and Mirarab 2018)
Print a phylogenetic tree using ASCII text in the UNIX shell environment	print_tree	print; pt	processing/editing	phylogeny	NA

Prune tips from a phylogenetic tree	prune_tree	prune	processing/editing	phylogeny	NA
Rename tip names in a phylogenetic tree	rename_tree_tips	rename_tree; rename_tips	processing/editing	phylogeny	NA
Calculate raw and normalized Robinson-Foulds distance scores between two phylogenetic trees	robinson_foulds_distance	rf_distance; rf_dist; rf	analytic	phylogeny	(Robinson and Foulds 1981)
Identify putatively spurious sequences by identifying terminal branches with an outlier branch length	spurious_sequence	spurious_seq; ss	analytic	phylogeny	(Jarvis <i>et al.</i> 2014)
Print all tip names in a phylogenetic tree	tip_labels	tree_labels; labels; tl	processing/editing	phylogeny	NA
Calculate treeness or stemminess of a phylogenetic tree	treeness	tness	analytic	phylogeny	(Lanyon 1988)
Calculate saturation, a measure of multiple substitutions across many sites in an MSA	saturation	sat	analytic	alignment and phylogeny	(Philippe <i>et al.</i> 2011)
Calculate treeness divided by relative composition variability, a measure of composition bias susceptibility	treeness_over_rcv	toverr	analytic	alignment and phylogeny	(Phillips and Penny 2003)

572 *Summary statistics reported are mean, median, 25th percentile, 75th percentile, minimum, maximum, standard deviation, and variance

573 values. All functions that calculate summary statistics also have verbose options that print out the raw data used to calculate summary

574 statistics.