

Non-lesioned subcortical brain volumes are associated with post-stroke sensorimotor behavior across 28 cohorts worldwide: An ENIGMA Stroke Recovery study

Sook-Lei Liew^{1,2*}, PhD, OTR/L, Artemis Zavaliangos-Petropulu^{3,4}, BA, Nicolas Schweighofer⁵, PhD, Neda Jahanshad⁴, PhD, Catherine E. Lang⁶, PhD, Keith R. Lohse⁷, PhD, Nerisa Banaj⁸, PhD, Giuseppe Barisano^{3,9}, MD, Lee A. Baugh¹⁰⁻¹³, PhD, Anup K. Bhattacharya¹⁴, MD, Bavrina Bigjahan¹⁵, MSc, Michael R. Borich¹⁶, PhD, Lara A. Boyd¹⁷, PhD, Amy Brodtmann^{18,19}, PhD, MD, Cathrin M. Buetefisch^{16,20,21}, PhD, MD, Winston D. Byblow²², PhD, Jessica M. Cassidy²³, PhD, Valentina Ciullo⁸, PhD, Adriana B. Conforto^{24,25}, PhD, Richard C. Craddock²⁶, PhD, Adrienne N. Dula, PhD,²⁶ Natalia Egorova^{18,27}, PhD, Wuwei Feng, MD²⁸, Kelene A. Fercho^{29,30}, PhD, Chris M. Gregory²⁸, PhD, PT, Colleen A. Hanlon^{31,32}, PhD, Kathryn S. Hayward^{18,33,34}, PhD, Jess A. Holguin¹, OTD, Brenton Hordacre³⁵, PhD, Darryl H. Hwang^{15,36}, PhD, Steven A. Kautz^{28,37}, PhD, Mohamed Salah Khelif¹⁸, PhD, Bokkyu Kim³⁸, PhD, Hosung Kim², PhD, Amy Kuceyeski³⁹, PhD, Jingchun Liu, MD,⁴⁰ David Lin⁴¹, MD, Martin Lotze⁴², MD, Bradley J. MacIntosh^{43,44}, PhD, John L. Margetis¹, OTD, Feroze B. Mohamed⁴⁵, PhD, Jan Egil Nordvik⁴⁶, PhD, Matthew A. Petoe^{47,48}, PhD, Fabrizio Piras⁸, PhD, Sharmila Raju⁴⁹, MSc, Ander Ramos-Murguialday^{51,51}, PhD, Kate P. Reville⁵², PhD, Pamela Roberts^{1,53,54}, PhD, OTR/L, SCFES, FAOTA, CPHQ, FNAP, FACRM, Andrew D. Robertson^{55,56}, PhD, Heidi M. Schambra⁴⁹, MD, Na Jin Seo^{28,37,57}, PhD, Mark S. Shiroishi^{4,15}, MD, MSc, Surjo R. Soekadar⁵⁸, MD, Gianfranco Spalletta^{8,59}, PhD, MD, Cathy M. Stinear⁶⁰, PhD, Anisha Suri⁶¹, MSc, Wai Kwong Tang⁶², MD, Gregory T. Thielman⁶³, EdD, Vincent N. Thijs^{18,64}, PhD, MD, Daniela Vecchio⁸, PhD, Junping Wang, MD, Nick S. Ward⁶⁵, MD, Lars T. Westlye^{66,67}, PhD, Carolee J. Winstein^{5,68}, PhD, George F. Wittenberg^{69,70}, PhD, MD, Kristin A. Wong⁷¹, MD, Chunshui Yu^{40,72}, MD, Steven L. Wolf⁷³⁻⁷⁵, PhD, Steven C. Cramer^{54,68}, MD, and Paul M. Thompson⁴, PhD

Affiliations

1. Chan Division of Occupational Science and Occupational Therapy, University of Southern California, Los Angeles, CA, USA.

2. Mark and Mary Stevens Neuroimaging and Informatics Institute, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA.
3. Neuroscience Graduate Program, University of Southern California, Los Angeles, CA, USA.
4. Imaging Genetics Center, Mark and Mary Stevens Neuroimaging and Informatics Institute, Keck School of Medicine, University of Southern California, Marina del Rey, CA, USA.
5. Biokinesiology and Physical Therapy, Ostrow School of Medicine, University of Southern California, Los Angeles, CA, USA.
6. Departments of Physical Therapy, Occupational Therapy, and Neurology, Washington University School of Medicine, St. Louis, MO, USA.
7. Department of Health and Kinesiology, University of Utah, Salt Lake City, UT, USA.
8. Laboratory of Neuropsychiatry, IRCCS Santa Lucia Foundation, Rome, Italy.
9. Laboratory of Neuro Imaging, Mark and Mary Stevens Neuroimaging and Informatics Institute, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA.
10. Basic Biomedical Sciences, Sanford School of Medicine, University of South Dakota, Vermillion, SD, USA.
11. Sioux Falls VA Health Care System, Sioux Falls, SD, USA.
12. Center for Brain and Behavior Research, Vermillion, SD, USA.
13. Sanford Research, Sioux Falls, SD, USA.
14. Mallinckrodt Institute of Radiology, Washington University School of Medicine in St. Louis, St. Louis, MO, USA.
15. Department of Radiology, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA.
16. Department of Rehabilitation Medicine, Emory University, Atlanta, GA, USA.
17. Department of Physical Therapy & the Djavad Mowafaghian Centre for Brain Health, Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada.
18. Florey Institute of Neuroscience and Mental Health, University of Melbourne, Heidelberg, VIC, Australia.
19. Eastern Cognitive Disorders Clinic, Monash University, Melbourne, VIC, Australia.

20. Department of Neurology, School of Medicine, Emory University, Atlanta, GA, USA.
21. Department of Radiology, Emory University, Atlanta, GA, USA.
22. Department of Exercise Sciences and Centre for Brain Research, University of Auckland, Auckland, New Zealand.
23. Allied Health Science, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA.
24. Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, São Paulo, SP, Brazil.
25. Hospital Israelita Albert Einstein, São Paulo, SP, Brazil.
26. Department of Diagnostic Medicine, Dell Medical School, University of Texas at Austin, Austin, Texas, USA.
27. Melbourne School of Psychological Sciences, University of Melbourne, Melbourne, VIC, Australia.
28. Department of Health Sciences & Research, Medical University of South Carolina, Charleston, SC, USA.
29. Civil Aerospace Medical Institute, US Federal Aviation Administration, Oklahoma City, OK, USA.
30. Division of Basic Biomedical Sciences, Sanford School of Medicine, University of South Dakota, Vermillion, SD, USA.
31. Cancer Biology, Wake Forest School of Medicine, Winston Salem, NC, USA.
32. College of Health Professions, Medical University of South Carolina, Charleston, SC, USA.
33. Department of Physiotherapy, University of Melbourne, Heidelberg, VIC, Australia.
34. NHMRC CRE in Stroke Rehabilitation and Brain Recovery, University of Melbourne, Heidelberg, VIC, Australia.
35. Innovation, IMplementation and Clinical Translation (IIMPACT) in Health, Allied Health and Human Performance, University of South Australia, Adelaide, SA, Australia.
36. Department of Biomedical Engineering, Viterbi School of Engineering, University of Southern California, Los Angeles, CA, USA.
37. Ralph H. Johnson VA Medical Center, Charleston, SC, USA.
38. Department of Physical Therapy Education, College of Health Professions, SUNY Upstate

- Medical University, Syracuse, NY, USA.
39. Department of Radiology, Weill Cornell Medicine, New York, NY, USA.
 40. Department of Radiology, Tianjin Medical University General Hospital, Tianjin, China.
 41. Center for Neurotechnology and Neurorecovery, Department of Neurology, Massachusetts General Hospital, Boston, MA, USA.
 42. Department of Diagnostic Radiology, University Medicine Greifswald, Greifswald, Germany.
 43. Hurvitz Brain Sciences, Sunnybrook Research Institute, Toronto, ON, Canada.
 44. Department of Medical Biophysics, University of Toronto, Toronto, ON, Canada.
 45. Jefferson Integrated Magnetic Resonance Imaging Center, Department of Radiology, Thomas Jefferson University, Philadelphia, PA, USA.
 46. CatoSenteret Rehabilitation Center, Son, Norway.
 47. Bionics Institute, Melbourne, VIC, Australia.
 48. Department of Medicine and Centre for Brain Research, University of Auckland, Auckland, New Zealand.
 49. Department of Neurology, Langone School of Medicine, New York University, New York, NY, USA.
 50. TECNALIA, Basque Research and Technology Alliance (BRTA), Neurotechnology Laboratory, San Sebastian Donostia, Spain.
 51. Institute of Medical Psychology and Behavioral Neurobiology, University of Tübingen, Tübingen, Germany.
 52. Facility for Education and Research in Neuroscience, Emory University, Atlanta, GA, USA.
 53. Department of Physical Medicine and Rehabilitation, Cedars-Sinai, Los Angeles, CA, USA.
 54. California Rehabilitation Institute, Los Angeles, CA, USA.
 55. Canadian Partnership for Stroke Recovery, Sunnybrook Research Institute, University of Toronto, Toronto, ON, Canada.
 56. Department of Kinesiology, University of Waterloo, Waterloo, ON, Canada.
 57. Department of Rehabilitation Sciences, Medical University of South Carolina, Charleston, SC, USA.

58. Clinical Neurotechnology Laboratory, Dept. of Psychiatry and Psychotherapy, Charité - University Medicine Berlin, Berlin, Germany.
59. Division of Neuropsychiatry, Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, TX, USA.
60. Department of Medicine, University of Auckland, Auckland, New Zealand.
61. Department of Electrical and Computer Engineering, Swanson School of Engineering, University of Pittsburgh, Pittsburgh, PA, USA.
62. Department of Psychiatry, Faculty of Medicine, the Chinese University of Hong Kong, Hong Kong, China.
63. Department of Physical Therapy and Neuroscience, University of the Sciences, Philadelphia, PA, USA.
64. Department of Neurology, Austin Health, Heidelberg, VIC, Australia.
65. UCL Queen Square Institute of Neurology, London, UK.
66. Department of Psychology, University of Oslo, Oslo, Norway.
67. NORMENT, Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway.
68. Department of Neurology, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA.
69. Department of Neurology, University of Pittsburgh, Pittsburgh, PA, USA.
70. Neurology, Department of Veterans Affairs Pittsburgh Healthcare System, Pittsburgh, PA, USA.
71. Physical Medicine & Rehabilitation, Dell Medical School, University of Texas at Austin, Austin, TX, USA.
72. Tianjin Key Laboratory of Functional Imaging, Tianjin Medical University General Hospital, Tianjin, China.
73. Division of Physical Therapy Education, Department of Rehabilitation Medicine, Department of Medicine, and Department of Cell Biology, Emory University School of Medicine, Atlanta, GA, USA.
74. Nell Hodgson Woodruff School of Nursing, Emory University, Atlanta, GA, USA.
75. Center for Visual and Neurocognitive Rehabilitation, Atlanta VA Health Care System, Decatur, GA, USA.

***Corresponding Author Contact Information:**

Sook-Lei Liew, PhD, OTR/L

Assistant Professor and Director, Neural Plasticity and Neurorehabilitation Laboratory

University of Southern California

Address: 2025 Zonal Ave, Los Angeles, CA 90033

Telephone: 323-865-1755

Email: sliew@usc.edu

ABSTRACT

Up to two-thirds of stroke survivors experience persistent sensorimotor impairments. Recovery relies on the integrity of spared brain areas to compensate for damaged tissue. Subcortical regions play critical roles in the control and regulation of sensorimotor circuits. Identifying relationships between sensorimotor behavior and non-lesioned subcortical volumes will reveal new neural targets for improving outcomes.

We pooled high-resolution T1-weighted MRI brain scans and behavioral data in 828 individuals with unilateral stroke from 28 cohorts worldwide (age: median 63, interquartile range 19 years; 516 males, 312 females). Cross-sectional analyses using linear mixed-effects models related post-stroke sensorimotor behavior to non-lesioned subcortical volumes. We analyzed subacute (≤ 90 days) and chronic (≥ 180 days) stroke; sub-analyses in chronic stroke were performed on class of sensorimotor deficit (impairment, activity limitations) and side of lesioned hemisphere, with exploratory analyses in early stroke (≤ 21 days) and across time (Bonferroni-corrected, $p < 0.004$).

Worse sensorimotor behavior was associated with a smaller ipsilesional thalamic volume in both subacute ($n=274$, $d=0.46$) and early stroke ($n=179$; $d=0.68$). In chronic stroke ($n=404$), worse sensorimotor behavior was associated with smaller ipsilesional putamen ($d=0.52$) and nucleus accumbens ($d=0.39$) volumes, and a larger ipsilesional lateral ventricle volume ($d=-0.42$), representing atrophy. In chronic stroke, worse sensorimotor impairment specifically (measured by the Fugl-Meyer Assessment; $n=256$) was associated with a smaller ipsilesional putamen ($d=0.72$), and larger lateral ventricle ($d=-0.41$), while several measures of activity limitations ($n=116$) showed no significant relationships. Side of lesion (left=214, right=190) had no impact. The full cohort ($n=828$) revealed associations of sensorimotor behavior with the ipsilesional nucleus accumbens ($d=0.23$), putamen ($d=0.33$), thalamus ($d=0.33$), and lateral ventricle ($d=-0.23$).

This analysis identified significant relationships between sensorimotor behavior and key subcortical regions at different times post-stroke. While further longitudinal studies are needed, these findings may represent brain imaging markers of resilience and reserve and provide putative neuroanatomical targets for improving sensorimotor outcomes post-stroke.

Keywords: stroke, rehabilitation, sensorimotor behavior, MRI, subcortical volumes

INTRODUCTION

Despite intensive research efforts and decades of clinical trials, stroke remains a leading cause of adult long-term disability worldwide (Virani *et al.*, 2020), and interventions to improve sensorimotor outcomes have yielded variable results. Although baseline impairment, corticospinal tract integrity, and lesion overlap with cortical regions are reliable prognostic indicators (Boyd *et al.*, 2017), effective therapies to improve sensorimotor outcomes are sparse. Even when accounting for these predictors, many interventional studies still show large inter-individual variability in response, suggesting that personalized neuroanatomical targets should be considered during rehabilitation.

While the role of cortical regions in post-stroke sensorimotor behavior has been widely examined, less attention has been paid to subcortical structures, such as the thalamus and basal ganglia. These structures not only play a critical role in the maintenance and regulation of sensorimotor circuits and motor learning, but they also subserve cognition, metabolic regulation, and reward—all of which have been implicated as contributors to post-stroke outcomes, including sensorimotor functioning and recovery (Fries *et al.*, 1993; Binkofski *et al.*, 1996; Shelton and Reding, 2001; Kuceyeski *et al.*, 2016). Each structure in the cortico-striatal-thalamic circuit has a distinct role in sensorimotor control and possibly outcomes. For instance, the thalamus is integral to the regulation of metabolism, sleep and wakefulness, cognitive processing, and integrating sensorimotor information (Jones, 2012), and thalamic metabolism has been shown to be disordered in the early weeks after stroke (Binkofski *et al.*, 1996; Carmichael *et al.*, 2004). Similarly, the basal ganglia (e.g., caudate, putamen, globus pallidus, and nucleus accumbens) are heavily involved in motor control, learning, and reward, with distinct roles for each nuclei (Alexander *et al.*, 1991; Lanciego *et al.*, 2012). Direct damage to the thalamus and basal ganglia is associated with poor sensorimotor behavior and recovery (Fries *et al.*, 1993; Boyd and Winstein, 2004), but the role of each spared subcortical nuclei is unclear.

To date, these subcortical structures have been studied only in modestly-sized samples, with varying results, and with measurements across multiple regions often aggregated as one (e.g., combined analysis of the thalamus and basal ganglia). However, each nucleus has a characteristic distribution of neurotransmitters, and identifying specific non-lesioned subcortical nuclei could provide more precise neurobiological targets for therapeutics to potentiate recovery.

Inter-individual variability and the heterogeneity of brain changes after stroke pose challenges to the identification of neural targets in spared tissue. Addressing this issue requires large, diverse, and appropriately-powered sample sizes with high-resolution brain MRIs. Although acute stroke research has successfully utilized pooled approaches with individual patient data to examine acute treatment outcomes (Goyal *et al.*, 2016; Campbell *et al.*, 2019), stroke rehabilitation research has been slower to adopt this type of approach due to the complexity of combining elaborate rehabilitation research protocols, differences in the site and size of infarcts, diversity of the patient populations recruited, and variety of the stroke neuroimaging and behavioral measures collected. To address these challenges, we formed the international ENIGMA Stroke Recovery Working Group to harmonize and combine diverse individual patient data, including high-resolution structural brain MRIs and behavioral outcome measures, across multiple research centers (Liew *et al.*, 2020). This combined analysis pools individual patient data across research sites using a harmonized analytical pipeline and includes both published and unpublished data. Compared to traditional single-site analyses or retrospective meta-analyses, this approach allows for greater statistical rigor, testing of more sophisticated hypotheses (e.g., subgroup analyses), and less bias due to the inclusion of both published and unpublished data across diverse cohorts (Berlin *et al.*, 2002; Ioannidis, 2017). Furthermore, pooled analyses with multi-site data increase heterogeneity, which improves generalizability of findings, reduces research inefficiency by leveraging previously collected data to examine novel questions, and advances the field faster than is achievable by prospective studies (Glasziou *et al.*, 2014).

The current study pools data from 828 individuals across 28 cohorts worldwide from the ENIGMA Stroke Recovery Working Group to examine relationships between sensorimotor behavioral measures and volumes of the ipsilesional and contralesional thalamus, putamen, caudate, pallidum, and nucleus accumbens. Enlargement of the lateral ventricles was also examined as an indirect measure of atrophy and vascular integrity (Hijdra and Verbeeten Jr, 1991; Apostolova *et al.*, 2012). Given the neurobiological events unique to early and subacute stroke compared to chronic stroke, data were analyzed separately for individuals in the subacute (≤ 90 days) and chronic (≥ 180 days) stages (Bernhardt *et al.*, 2017). As an exploratory measure, we also analyzed relationships early after stroke (≤ 21 days), before post-stroke secondary structural atrophy is thought to be observed (Egorova *et al.*, 2019), to estimate whether subacute associations are driven by early post-stroke changes or likely existed prior to the stroke.

We hypothesized the thalamus relates to sensorimotor behavior in early and subacute phases after stroke, given its multiple roles in supporting cellular repair (Azari *et al.*, 1996; Binkofski *et al.*, 1996). We further expected to find smaller subcortical volumes, reflecting atrophy of structures associated with sensorimotor control, and larger ventricles, reflecting general atrophy, to be related to chronic sensorimotor behavior (Gauthier *et al.*, 2012). Furthermore, as sensorimotor behavior encompasses multiple classes of the International Classification of Functioning, Disability, and Health (ICF), separate subgroup analyses in chronic stroke were conducted to examine if there are specific neural correlates of loss of body structures and function (i.e., *sensorimotor impairment*) versus loss of activity in daily tasks (i.e., *activity limitations*) (McDougall *et al.*, 2010). Here, we anticipated that subcortical nuclei that play a significant role in direct sensorimotor control, such as the putamen, more strongly relate to impairment, while regions associated with reward and motivation, such as the nucleus accumbens, may be more strongly related to activity limitations. Finally, in chronic stroke, we also examined the impact of the side of the lesion. Based on evidence of hemispheric specialization for motor behavior after stroke (Sainburg *et al.*, 2016), we hypothesized that the side of the lesion would modify the relationship between non-lesioned subcortical tissue volume and sensorimotor behavior.

MATERIALS AND METHODS

Study design

The current cross-sectional pooled analysis used data from the ENIGMA Stroke Recovery Working Group, which was frozen for this analysis on May 22, 2020. A detailed overview of ENIGMA Stroke Recovery procedures and methods are reported elsewhere (Liew *et al.*, 2020). The data were collected across 28 different research studies (i.e., cohorts) at 16 different research institutes in 10 countries, in accordance with the Declaration of Helsinki and in compliance with local ethics review boards at each institute (see *Supplementary Table 1* for details).

ENIGMA Stroke Recovery Dataset

Participants with at least one sensorimotor behavioral outcome measure (see *Behavioral Data Analysis, Supplementary Table 1*) and a segmented high-resolution (e.g., 1-mm isotropic) T1-

weighted (T1w) structural MRI of the brain (see *MRI Data Analysis*) were included, yielding an initial dataset of 1,285 individuals. Only participants with unilateral ischemic stroke or intracerebral hemorrhage were included, and individuals identified as having bilateral lesions or lesions in the brainstem or cerebellum were excluded from this analysis. For any longitudinal observations, only the first time-point was used; the resulting dataset was therefore cross-sectional. Each brain region was manually inspected for quality and overlap with the lesion (see *MRI Data Analysis*). Any individuals missing covariates of age (n=50) or sex (n=89) were also excluded, yielding a final N=828 (age: median 63, interquartile range (IQR) 19 years; 516 males, 312 females). As the relationships between brain volume and sensorimotor behavior were expected to change with time after stroke, the data were divided into subacute stroke (≤ 90 days post-stroke) and chronic stroke (≥ 180 days post-stroke). Exploratory analyses looking only at early stroke (≤ 21 days post-stroke) and across all times after stroke are included in *Supplementary Materials*.

MRI Data Analysis

To extract subcortical volumes, brain imaging software package FreeSurfer (version 5.3) was used to segment subcortical regions of interest (ROIs) from the T1w MRIs (Fischl *et al.*, 2002). Twelve ROIs were extracted: the left and right thalamus, caudate, putamen, pallidum, nucleus accumbens, and lateral ventricles. For all analyses, these were characterized as ipsilesional and contralesional based on the lesioned hemisphere. Total intracranial volume (ICV) was also quantified using FreeSurfer outputs. ENIGMA scripts developed in-house were used to extract the volume of each ROI for each individual and to generate quality control (QC) triplanar images of each segmented ROI as done previously (Hibar *et al.*, 2015; Thompson *et al.*, 2020) (<http://enigma.ini.usc.edu/protocols/>). Given the variability of post-stroke neuroanatomy following a lesion, trained research team members (A.Z.-P., A.S.) performed visual QC for each ROI in each subject. Any regions intersecting the lesion were marked “lesioned” and any regions not properly segmented by FreeSurfer were marked “failed”; regions falling in either category were excluded from further analysis (for the full QC protocol, see Appendix 1 in (Liew *et al.*, 2020)). Sample sizes for each analysis and brain region are reported.

Behavioral Data Analysis

Across cohorts, behavioral data were collected within approximately 72 hours of the MRI. To maximize the utility of the full dataset, a *primary sensorimotor behavior score* was defined for each study cohort using the measure reported in that cohort that was most commonly represented in the dataset overall (see *Supplementary Materials*). From this measure, a fraction of the maximum possible score was calculated, such that 0 represented the worst sensorimotor performance (severe deficits) and 1 represented the best sensorimotor performance (no deficits). The most common measure across cohorts was the Fugl-Meyer Motor Assessment of Upper Extremities (FMA-UE) (Fugl-Meyer *et al.*, 1975).

In chronic stroke, we also identified behavioral measures that specifically captured impairment and activity limitation. Impairment was measured by the FMA-UE, whereas activity limitation was measured by the Action Research Arm Test (ARAT) (Lyle, 1981) and Wolf Motor Function Test (WMFT) (Wolf *et al.*, 2001); data with any of these measures were used. These data were not examined in early stroke due to the limited sample sizes with these measures.

Statistical Analysis

To examine the relationships between sensorimotor behavior and non-lesioned subcortical volumes, we performed linear mixed-effects regressions. A separate regression model was run for the volume of each subcortical ROI (outcome) using sensorimotor behavior (e.g., primary sensorimotor behavior score, sensorimotor impairment, or activity limitations) as the primary predictor of interest. After ruling out collinearity (variance inflation factor ≤ 2.5), normalized age, ICV, and sex were included as fixed effects. Research cohort was included as a random effect. In chronic stroke, the effect of side was also examined; an interaction term between sensorimotor behavior and side of lesioned hemisphere was added to the model predicting subcortical volume. This was not examined in subacute stroke due to the smaller sample size. A likelihood ratio test (LRT) was performed to compare models with and without random effects and showed that the random effects were always significant. The regression assumptions of linearity, normality of the residuals, and homogeneity of the residual variance were checked via visual inspection of residuals versus fits plots as well as qq-plots for both individual observations and research cohorts. Potential influential values for both observations and cohorts were assessed using Cook's distance with recommended thresholds (Nieuwenhuis *et al.*, 2012). As we detected influential observations in almost all analyses, we re-ran the analyses using robust mixed-effect

regression, which reduces the weight of influential observations in the models without excluding data (Greco *et al.*, 2019). Results did not differ between original and robust regression models. The results of the robust regression models can be found in *Supplementary Materials*.

For all analyses, beta coefficients are presented for the factor of interest (e.g., sensorimotor behavior, sensorimotor impairment, or activity limitations), along with the sample size (n), standard error (SE), degrees of freedom (df), standardized effect size (*d*), t-value, and uncorrected p-value. Statistical significance was adjusted for multiple comparisons across the 12 ROIs using a Bonferroni correction ($p < 0.004$). Any significant fixed covariates are also reported.

All statistical analyses were conducted in R (version 3.6.3; R Core Team, 2020 (Team, 2020)). The follow R libraries were used: the *lme* function from *nlme* was used for the linear mixed-effects regressions (Pinheiro J, 2020), the *rlmer* function from *robustlmm* was used for the robust linear mixed-effects regressions (Koller, 2016), and the *rstatix* library was used for the Wilcoxon rank sum test (Kassambara, 2019). In addition, *influence.ME* was used to detect influential values (Nieuwenhuis *et al.*, 2012) and *dplyr* (Wickham *et al.*, 2020) and *tidyverse* (Wickham *et al.*, 2019) libraries were used for data organization.

Data Availability

The deidentified summary data and code that support the findings of this study are available upon reasonable request from the corresponding author. The data are not all publicly available in a repository as they may contain information that could compromise the privacy of research participants. There are also data sharing restrictions imposed by some of the (i) ethical review boards of the participating sites, and consent documents; (ii) national and trans-national data sharing laws; and (iii) institutional processes, some of which require a signed DTA for limited and predefined data use. However, we welcome sharing data with researchers, requiring that they become members of the ENIGMA Stroke Recovery working group and submit an analysis plan for a secondary project for group review. Once this analysis plan is approved, access to the relevant data will be provided contingent on data availability, local PI approval and compliance with all supervening regulatory boards.

RESULTS

Data from 828 individuals from 28 cohorts worldwide were included (see Table 1 for an overview of cohort characteristics).

In subacute stroke (≤ 90 days; $n=274$), worse post-stroke sensorimotor behavior was significantly associated with smaller volumes of the ipsilesional thalamus ($n=274$, $d=0.46$, $p=0.002$; Table 2; Figure 1). A secondary analysis in individuals within the first 21 days post-stroke ($n=179$, $d=0.68$, $p<0.001$) demonstrated the same result (see *Supplementary Materials*).

In chronic stroke (≥ 180 days; $n=404$), worse sensorimotor behavior was related to smaller volumes of the ipsilesional putamen ($d=0.52$, $p<0.001$) and ipsilesional nucleus accumbens ($d=0.39$, $p=0.002$), and a larger volume of the ipsilesional lateral ventricle ($d=-0.42$, $p<0.001$; Table 2; Figure 1).

In chronic stroke, we examined brain-behavior relationships using a measure of impairment (the FMA-UE scale; $n=256$) and two measures of activity limitation (WMFT, ARAT; $n=116$). Worse sensorimotor impairment was associated with smaller ipsilesional putamen ($d=0.72$, $p=0.001$) and larger ipsilesional lateral ventricle volumes ($d=-0.41$, $p=0.002$; Table 3; Figure 1). We found no significant relationships between subcortical nuclei and measures of activity limitations (Table 3).

In chronic stroke, we further analyzed the differences between individuals with left hemisphere stroke (LHS, $n=214$) versus right hemisphere stroke (RHS, $n=190$) by including lesioned hemisphere as an interaction term in the model. There were no significant effects of the side of the lesioned hemisphere on the relationship between sensorimotor behavior and subcortical volumes, and no main effects of the lesioned hemisphere (see *Supplementary Materials*). Inclusion of the lesioned hemisphere into the model did not change the main effects of sensorimotor behavior. There were no differences in sensorimotor behavior between left and right hemisphere stroke groups (see *Supplementary Materials*).

Finally, an exploratory analysis of the entire cohort ($N=828$) demonstrated significant relationships between worse sensorimotor behavior and smaller volumes of the ipsilesional thalamus ($d=0.33$, $p=0.001$), putamen ($d=0.33$, $p<0.001$), and nucleus accumbens ($d=0.23$, $p=0.004$), and a larger lateral ventricle volume ($d=-0.23$, $p=0.001$). See *Supplementary Materials* for additional details.

DISCUSSION

We report the first international, multi-site pooled analysis with individual patient data using high-resolution structural brain imaging in stroke rehabilitation research and the largest study to date relating subcortical brain measures to post-stroke sensorimotor behavior. We identified novel, significant relationships between post-stroke sensorimotor behavior and the volumes of spared deep gray matter structures including the ipsilesional thalamus, putamen, and nucleus accumbens, as well as atrophy as indexed by enlargement of the ipsilesional lateral ventricle. Notably, all significant relationships were found only in the ipsilesional hemisphere. These findings suggest that, post-stroke, subcortical brain alterations related to sensorimotor behavior occur most prominently in the hemisphere directly affected by the stroke. This was observed despite the fact that, after stroke, atrophy and reorganization has been observed bilaterally (Brodtmann *et al.*, 2020). The identification of sensorimotor relationships with specific ipsilesional subcortical nuclei may provide novel neuromodulatory or pharmacological targets to improve stroke outcomes.

Our results support the hypothesis that non-lesioned deep gray structures serve distinct roles in subacute versus chronic stroke, which is not surprising given the cascade of neurobiological and neuroinflammatory processes that occur early after stroke (Murphy and Corbett, 2009; Ward, 2017). Within 90 days after stroke, only the ipsilesional thalamus showed detectable associations with post-stroke sensorimotor behavior, in line with recent research suggesting a role for the thalamus in subacute recovery (Brodtmann *et al.*, 2020). A smaller thalamic volume could reflect cell loss and thalamic dysfunction, thereby limiting resources critical for early recovery (Fries *et al.*, 1993; Brodtmann *et al.*, 2020). Importantly, this relationship persists, and is stronger, when looking at only the first 21 days post-stroke. As non-lesioned brain volumes within six weeks after stroke are assumed to be similar to those before the stroke (Egorova *et al.*, 2019), this finding suggests that larger thalamic volumes prior to stroke could provide a neuroprotective effect. Thalamic atrophy was recently associated with loss of extrinsic and intrinsic connectivity between the thalamus and the rest of the brain, suggesting that thalamic measures may serve as an index of global brain function (Mahajan *et al.*, 2020). Future research using longitudinal

datasets with greater spatial specificity could relate changes in specific thalamic nuclei to sensorimotor recovery to identify targets for neuroprotective or early stroke therapies.

In chronic stroke, smaller volumes of the ipsilesional putamen and nucleus accumbens were consistently associated with worse sensorimotor behavior. Brain atrophy, as indexed by a larger ipsilesional ventricle volume, was also negatively associated with sensorimotor behavioral measures. This is the first large-scale validation of these specific subcortical volume measures as correlates of sensorimotor behavioral outcomes in chronic stroke. This finding augments existing stroke literature, which has typically examined changes across combined subcortical regions, without differentiating roles of the individual basal ganglia nuclei and thalamus. Direct damage to the putamen has been related to post-stroke gait impairment (Alexander *et al.*, 2009), upper limb impairment (Lee *et al.*, 2015), and spasticity (Cheung *et al.*, 2016), deficits which overlap with the behavioral measures used here. Secondary atrophy of the putamen has been reported after cortical stroke and associated with infarct volume (Baudat *et al.*, 2020) and post-stroke cognitive deficits (Lopes *et al.*, 2012). The relationship between chronic sensorimotor behavioral deficits and the volume of the non-lesioned ipsilesional putamen after stroke, however, has not been reported. The putamen plays a key role in corticostriatal circuits, receiving sensorimotor cortical signals and relaying them to the thalamus. Here we report that secondary atrophy in the putamen relates to sensorimotor behavior generally and impairment specifically, as evidenced by the association with the FMA-UE in chronic stroke. Although further research is needed, given its diverse roles in cognitive and sensorimotor function, the volume of the putamen, when spared by stroke, may represent a global measure of general corticostriatal circuit integrity in chronic stroke survivors.

While the ipsilesional nucleus accumbens was significantly related to chronic sensorimotor behavior in general, it was not related to sensorimotor impairment (FMA-UE) alone, nor was it associated with activity limitations (e.g., ARAT, WMFT), as hypothesized. However, the analyses on impairment and activity limitations had less statistical power to detect relationships. As the nucleus accumbens is a key component of the ventral striatum and implicated in dopaminergic modulation of reward-based behaviors (Robbins and Everitt, 1992), this region may impact more complex aspects of motor performance, such as motivation and participation, compared to impairment or activity. A number of studies show decreases in ventral striatal processes such as reward sensitivity, motivation, and apathy after stroke (Rochat *et al.*, 2013),

and post-stroke hypoactivity in the nucleus accumbens has been identified during reward-based decision-making tasks (Widmer *et al.*, 2019). This effect was observed despite no direct lesions in the nucleus accumbens, suggesting that secondary dysfunction of this network can impact behavior after stroke. It is likely that the nucleus accumbens impacts sensorimotor behavior by influencing reward and motivation (Sawada *et al.*, 2015), which could influence use of the affected limb in daily tasks. Although pharmacological methods to modulate the dopaminergic system and promote motor recovery following stroke have been widely studied, there are large individual differences in outcomes (Gower and Tiberi, 2018). Future research may investigate whether individual differences in the volume and connectivity of the nucleus accumbens predict who may benefit from dopaminergic treatment.

The third important finding in chronic stroke is the association between an enlarged ipsilesional lateral ventricle and poor sensorimotor behavior. This relationship was only significant at the chronic stage and was exclusive to the ipsilesional lateral ventricle, which may be due to hydrocephalus ex vacuo. Ventricular enlargement post-stroke may also be influenced by small vessel disease (i.e., leukoaraiosis), although this is typically observed bilaterally (Hijdra and Verbeeten Jr, 1991). Enlargement of the bilateral lateral ventricles has also been associated with generalized brain atrophy that occurs during aging and with impaired cognitive function (Lenzi *et al.*, 1994; Förstl *et al.*, 1995). The contrast between ipsilesional and contralesional ventricles may provide unique insight into the specific impact of the stroke versus general aging on chronic stroke sensorimotor outcomes.

Our results also suggest that there are distinct brain-behavior relationships for different ICF dimensions of sensorimotor behavior. Chronic motor impairment, as measured by the FMA-UE, was associated with a smaller ipsilesional putamen and larger ipsilesional ventricle, which may provide an indication of corticostriatal circuit functions as well as overall brain health essential for sensorimotor control. In contrast, there were no subcortical associations with activity limitations in the current study. This could be related to the smaller sample size (n=116 versus n=256 for sensorimotor impairment). Alternatively, activity limitations may be more strongly related to the integrity or function of distributed regions across whole brain networks rather than subcortical structures (Dong *et al.*, 2006; Van Meer *et al.*, 2010), given that functional performance can be influenced by psychosocial factors to a greater degree than impairment measures.

Finally, findings did not indicate a significant effect of lesioned hemisphere on the relationship between chronic sensorimotor behavior and spared subcortical volumes. These results are surprising, given that the large majority of patients were likely left hemisphere dominant for motor control (Kim *et al.*, 1993), and previous research has identified post-stroke hemispheric specializations and roles in sensorimotor control (Schaefer *et al.*, 2007; Mani *et al.*, 2013; Sainburg *et al.*, 2016). However, previous research has primarily focused on cortical regions and functional activity, rather than subcortical structures. Side of stroke injury may not directly impact sensorimotor relationships with spared subcortical volumes.

Limitations and Future Directions

A key limitation of pooling multi-site individual patient data is inconsistent variables across cohorts, limiting subgroup analyses and reducing the number of included covariates. Models only included the covariates age, sex, and intracranial volume; however, many additional demographic variables, such as duration and type of rehabilitation received, handedness, race, educational level, and comorbidities, may influence these relationships. In addition, larger sample sizes for specific behavioral measures would provide greater support for the current findings. Related, small samples ($n < 50$) at earlier time points of stroke (i.e., ≤ 7 days, defined as acute (Bernhardt *et al.*, 2017)) with sensorimotor behavioral outcomes limited our ability to specifically examine acute brain-behavior relationships or subacute relationships with impairment versus activity limitations in the current analysis. Therefore, the ENIGMA Stroke Recovery Working Group recommends following consensus guidelines for greater harmonization of prospectively-collected data to facilitate more precise pooled analyses across all times after stroke in the future (Kwakkel *et al.*, 2017).

Lesion overlap with subcortical regions, and poor segmentation of subcortical regions due to lesion-induced distortions, resulted in a variable sample size for each ROI, potentially limiting the power to detect relationships in regions with smaller samples. Furthermore, excluding individuals with lesioned or incorrectly segmented ROIs may disproportionately exclude individuals with larger lesions, who may be more severely affected. This could have biased the sample towards more mild-to-moderately impaired patients. Future studies using lesion masks for each observation could address these issues and also provide additional information about lesion location and volume and as well as direct lesion overlap with each subcortical region.

Finally, many of these subcortical regions are also critical for and related to post-stroke cognition, mood, sleep, learning and other traits of interest. While this analysis was limited to sensorimotor behavioral measures to maximize available data for analysis, these findings may not be unique to sensorimotor behavior. Future studies should assess the relationship between these subcortical volumes and additional stroke outcome measures.

Conclusion

This international collaborative analysis revealed significant relationships between post-stroke sensorimotor behavior and volumetric measures of the non-lesioned ipsilesional thalamus, putamen, nucleus accumbens, and lateral ventricle at different times after stroke – brain metrics that may reflect overall brain health and network integrity and could lead to the identification of novel neural targets for pharmacological or behavioral modulation in stroke rehabilitation.

FIGURES




Figure 1. Relationship between post-stroke sensorimotor behavior and non-lesioned subcortical volumes. Non-lesioned subcortical regions (1D, bottom right) that relate to sensorimotor behavior from linear mixed-effects models of people with subacute (1A, top left) and chronic (1B, bottom left) stroke. Non-lesioned subcortical volume relationships with chronic sensorimotor impairment is shown in 1C (top right). There were no significant volume relationships with chronic activity limitations. Colors represent the beta estimate (β) for sensorimotor behavior from each model, with warmer colors representing more positive beta estimates and cooler colors representing more negative beta estimates.

TABLES

Cohort ID	<i>n</i>	Females / Males	Median Age (IQR, min-max)	Median Sensorimotor Score (IQR, min-max)
1	39	10 / 29	61 (17, 31-80)	0.65 (0.23, 0.0-0.9)
2	12	06 / 06	70 (12, 39-85)	0.50 (0.41, 0.2-0.7)
3	14	06 / 08	60 (15, 33-85)	0.25 (0.22, 0.1-0.6)
4	19	06 / 13	44 (15, 30-68)	0.14 (0.17, 0.0-0.5)
7	42	14 / 28	56 (14, 18-80)	0.82 (0.35, 0.4-1.0)
8	8	02 / 06	62 (10, 39-75)	0.55 (0.35, 0.0-1.0)
9	93	29 / 64	70 (16, 24-88)	1.00 (0.07, 0.0-1.0)
10	24	05 / 19	59 (13, 42-74)	1.00 (0.02, 0.7-1.0)
11	29	10 / 19	57 (11, 44-71)	1.00 (0.05, 0.1-1.0)
12	57	31 / 26	71 (17, 31-97)	0.65 (0.71, 0.0-1.0)
13	44	22 / 22	72 (18, 33-91)	0.12 (0.32, 0.0-1.0)
15	14	06 / 08	57 (11, 45-74)	0.72 (0.25, 0.4-0.8)
17	16	05 / 11	59 (04, 45-68)	0.55 (0.23, 0.2-0.7)
18	11	05 / 06	59 (07, 46-73)	0.65 (0.22, 0.5-0.9)
19	13	03 / 10	62 (21, 33-74)	0.84 (0.08, 0.8-0.9)
20	22	08 / 14	70 (13, 49-79)	0.91 (0.14, 0.3-1.0)
22	17	04 / 13	59 (30, 25-72)	0.63 (0.50, 0.0-0.8)
23	13	07 / 06	58 (08, 31-90)	0.42 (0.17, 0.3-0.8)
24	21	11 / 10	63 (13, 32-78)	0.95 (0.00, 0.6-1.0)
25	26	10 / 16	65 (18, 37-88)	0.97 (0.20, 0.0-1.0)
26	24	14 / 10	49 (20, 25-71)	0.64 (0.14, 0.3-0.8)
28	26	07 / 19	62 (11, 23-75)	0.75 (0.25, 0.3-1.0)
31	35	09 / 26	58 (12, 21-86)	0.52 (0.31, 0.2-0.9)
32	7	03 / 04	62 (16, 38-72)	0.95 (0.44, 0.2-1.0)
34	15	06 / 09	58 (11, 32-80)	0.82 (0.20, 0.6-1.0)
35	15	06 / 09	64 (18, 31-83)	0.64 (0.52, 0.2-0.9)
38	81	34 / 47	66 (19, 30-89)	0.85 (0.60, 0.0-1.0)
41	91	33 / 58	70 (15, 32-89)	1.00 (0.02, 0.8-1.0)
TOTAL	828	312 / 516	63 (19, 18-97)	0.82 (0.48, 0-1)

Table 1. Summary of research cohort characteristics. Age and sensorimotor behavioral score data are shown as median (interquartile range (IQR), minimum-maximum values)

SUBACUTE AND CHRONIC STROKE

SUBACUTE STROKE (≤ 90 days)								
Brain Region	n	beta (CI)	SE	df	t-value	p-value	d	Significant covariates
<i>Ipsilesional</i>								
Caudate	194	-0.01 (-0.51-0.48)	0.25	180	-0.06	0.954	-0.01	ICV
Lateral ventricle	274	0.18 (-0.14-0.51)	0.16	259	1.13	0.258	0.14	Age, ICV
Nucleus accumbens	245	0.24 (-0.14-0.62)	0.19	231	1.26	0.210	0.17	Age
Pallidum	223	0.21 (-0.26-0.67)	0.24	209	0.87	0.387	0.12	ICV
Putamen	201	0.39 (-0.09-0.88)	0.25	187	1.61	0.109	0.24	Age, ICV
Thalamus	210	0.69 (0.27-1.11)	0.21	197	3.21	0.002	0.46	Age, ICV
<i>Contralesional</i>								
Caudate	219	0.22 (-0.20-0.64)	0.21	205	1.04	0.298	0.15	ICV
Lateral ventricle	274	0.15 (-0.18-0.49)	0.17	259	0.92	0.361	0.11	Age, ICV
Nucleus accumbens	253	0.15 (-0.23-0.52)	0.19	239	0.77	0.443	0.10	Age, ICV
Pallidum	250	0.50 (0.07-0.92)	0.22	236	2.30	0.022	0.30	ICV
Putamen	229	0.37 (-0.05-0.79)	0.21	215	1.75	0.081	0.24	Age, ICV
Thalamus	217	0.09 (-0.33-0.50)	0.21	204	0.41	0.679	0.06	Age, ICV
CHRONIC STROKE (≥ 180 days)								
Brain Region	n	beta (CI)	SE	df	t-value	p-value	d	Significant covariates
<i>Ipsilesional</i>								
Caudate	193	0.27 (-0.28-0.82)	0.28	169	0.98	0.330	0.15	ICV
Lateral ventricle	404	-0.70 (-1.04--0.36)	0.17	378	-4.04	<0.001	-0.42	Age, ICV
Nucleus accumbens	289	0.72 (0.27-1.18)	0.23	264	3.15	0.002	0.39	Age
Pallidum	225	0.30 (-0.23-0.84)	0.27	200	1.11	0.267	0.16	ICV
Putamen	207	1.01 (0.45-1.57)	0.28	183	3.54	<0.001	0.52	Age
Thalamus	169	0.08 (-0.60-0.75)	0.34	146	0.22	0.827	0.04	Age
<i>Contralesional</i>								
Caudate	345	0.08 (-0.31-0.48)	0.20	320	0.41	0.679	0.05	ICV
Lateral ventricle	404	-0.39 (-0.70--0.07)	0.16	378	-2.42	0.016	-0.25	Age, ICV
Nucleus accumbens	344	0.21 (-0.22-0.65)	0.22	319	0.96	0.339	0.11	Age
Pallidum	359	0.20 (-0.20-0.60)	0.20	334	0.97	0.332	0.11	Sex, ICV
Putamen	355	0.21 (-0.18-0.60)	0.20	330	1.06	0.291	0.12	Age, ICV
Thalamus	329	-0.24 (-0.60-0.12)	0.18	304	-1.29	0.196	-0.15	Age, ICV

Table 2. Relationships between non-lesioned subcortical volumes and sensorimotor behavior in subacute and chronic stroke. Results from linear mixed-effects models of individuals with subacute stroke (top) and chronic stroke (bottom). Results in bold indicate significance with a Bonferroni correction for multiple comparisons ($p < 0.004$). The beta coefficient for sensorimotor behavior (beta) with 95% confidence interval (CI), along with the sample size (n), standard error (SE), degrees of freedom (df), standardized effect size (d), t-value, and uncorrected p-value are reported, in addition to significant fixed covariates including age, sex, and intracranial volume (ICV).

CHRONIC SENSORIMOTOR IMPAIRMENT AND ACTIVITY LIMITATIONS

SENSORIMOTOR IMPAIRMENT IN CHRONIC STROKE								
Brain Region	n	beta (CI)	SE	df	t-value	p-value	d	Significant covariates
<i>Ipsilesional</i>								
Caudate	94	0.92 (-0.06-1.89)	0.49	77	1.87	0.065	0.43	ICV
Lateral ventricle	256	-0.74 (-1.20--0.27)	0.24	237	-3.13	0.002	-0.41	Age, ICV
Nucleus accumbens	171	0.58 (0.01-1.15)	0.29	153	2.02	0.045	0.33	Age
Pallidum	120	0.76 (0.01-1.51)	0.38	102	2.02	0.046	0.40	-
Putamen	104	1.50 (0.61-2.39)	0.45	87	3.34	0.001	0.72	-
Thalamus	84	0.33 (-0.72-1.38)	0.53	68	0.62	0.537	0.15	-
<i>Contralesional</i>								
Caudate	222	0.06 (-0.44-0.57)	0.26	204	0.25	0.806	0.03	ICV
Lateral ventricle	256	-0.51 (-0.88--0.14)	0.19	237	-2.70	0.007	-0.35	Age, ICV
Nucleus accumbens	222	0.21 (-0.31-0.73)	0.26	204	0.80	0.425	0.11	Age
Pallidum	231	0.20 (-0.33-0.73)	0.27	213	0.74	0.459	0.10	Sex
Putamen	229	0.10 (-0.38-0.58)	0.24	211	0.41	0.681	0.06	Age, ICV
Thalamus	211	-0.40 (-0.88-0.07)	0.24	193	-1.67	0.096	-0.24	Age, ICV
ACTIVITY LIMITATIONS IN CHRONIC STROKE								
Brain Region	n	beta (CI)	SE	df	t-value	p-value	d	Significant covariates
<i>Ipsilesional</i>								
Caudate	52	-0.63 (-1.80-0.53)	0.58	44	-1.09	0.280	-0.33	-
Lateral ventricle	116	-0.71 (-1.46-0.04)	0.38	108	-1.88	0.062	-0.36	Age, ICV
Nucleus accumbens	86	0.77 (-0.31-1.85)	0.54	78	1.42	0.159	0.32	-
Pallidum	64	0.71 (-0.25-1.67)	0.48	56	1.47	0.146	0.39	-
Putamen	65	0.71 (-0.62-2.04)	0.67	57	1.06	0.292	0.28	-
Thalamus	56	0.94 (-0.36-2.25)	0.65	48	1.45	0.153	0.42	-
<i>Contralesional</i>								
Caudate	96	-0.07 (-0.98-0.84)	0.46	88	-0.15	0.885	-0.03	-
Lateral ventricle	116	-0.72 (-1.44-0.01)	0.37	108	-1.95	0.054	-0.38	Age, ICV
Nucleus accumbens	107	-0.34 (-1.17-0.49)	0.42	99	-0.81	0.420	-0.16	Age
Pallidum	103	-0.15 (-0.98-0.68)	0.42	95	-0.35	0.728	-0.07	Sex
Putamen	100	0.06 (-0.91-1.03)	0.49	92	0.12	0.903	0.03	Age
Thalamus	92	0.28 (-0.51-1.06)	0.39	84	0.71	0.482	0.15	Age, ICV

Table 3. Relationships between non-lesioned subcortical volumes and two measures of sensorimotor behavior (impairment, activity limitations). Results from linear mixed-effects models in individuals with chronic stroke of sensorimotor impairment (top) compared to activity limitations (bottom). Results in bold indicate significance with a Bonferroni correction for multiple comparisons ($p < 0.004$). The beta coefficient for sensorimotor impairment/activity limitations (beta) with 95% confidence interval (CI), along with the sample size (n), standard error (SE), degrees of freedom (df), standardized effect size (d), t-value, and uncorrected p-value are reported, in addition to significant fixed covariates including age, sex, and intracranial volume (ICV).

ACKNOWLEDGEMENTS

We thank all of the members of the ENIGMA Stroke Recovery working group, all of the participants, as well as Sophia Thomopolous and Bethany Lo for their assistance.

FUNDING

S.-L.L. is supported by NIH K01 HD091283; NIH R01 NS115845.

N.S. is supported by NIH R56 NS100528.

N.J. is supported by NIH R01 AG059874; NIH R01 MH117601.

A.B. is supported by National Health and Medical Research Council (NHMRC) of Australia GNT1020526; GNT1094974; Heart Foundation Future Leader Fellowship 100784.

C.M.B is supported by NIH R21 HD067906; NIH R01 NS090677.

W.D.B. is supported by Health Research Council of New Zealand (09/164R).

J.M.C is supported by NIH R00 HD091375.

A.B.C. is supported by NIH R01 NS076348; IIEP-2250-14.

N.E. is supported by Australian Research Council DE180100893.

W.F. is supported by NIH P20 GM109040

C.A.H. is supported by NIH P20 GM109040.

K.S.H. is supported by National Health and Medical Research Council (NHMRC) of Australia #1088449; NIH R01 NS115845.

B.H. is supported by National Health and Medical Research Council (NHMRC) fellowship (1125054).

S.A.K. is supported by NIH 1IK6RX003075; NIH P20 GM109040.

B.K is supported by NIH R01 HD065438; NIH R56 NS100528.

H.K. is supported by a BrightFocus Faculty Award.

B.J.M. is supported by Canadian Partnership for Stroke Recovery; Canadian Institutes of Health Research; Natural Sciences and Engineering Research Council; Brain & Behavior Research Foundation.

A.R.-M. is supported by Basque Government Elkartek MODULA; Bundesministerium für Bildung und Forschung BMBF AMORSA (FKZ 16SV7754); and the Fortüne-Program of the University of Tübingen (2452-0-0).

F.P. is supported by the Italian Ministry of Health, Grants RC 2016, 2017, 2018, 2019.

K.P.R. is supported by NIH R21 HD067906; NIH R01 NS090677.

H.M.S. is supported by NIH R01 NS110696; NIH R01 LM013316; NIH K02 NS104207.

N.J.S. is supported by NIH U54 GM104941; NIH P20 GM109040.

S.R.S. is supported by the European Research Council (ERC, Grant number 759370).

G.S. is supported by Italian Ministry of Health grant RC 15-16-17-18-19-20/A.

C.M.S. is supported by the Health Research Council of New Zealand.

L.T.W. is supported by the South-Eastern Norway Regional Health Authority (2014097, 2015044, 2015073); the Norwegian ExtraFoundation for Health and Rehabilitation (2015/FO5146); the Research Council of Norway (249795, 262372); and the European Research Council under the European Union's Horizon 2020 Research and Innovation program (ERC StG, Grant 802998).

G.F.W. is supported by the Department of Veterans Affairs RR&D Program.

S.L.W. is supported by NIH R01 NS115845.

P.M.T. is supported by NIH U54 EB020403.

COMPETING INTERESTS

N.J. and P.M.T. are MPI of a research grant from Biogen, Inc for work unrelated to the contents of this manuscript. S.C.C. has served as a consultant for Constant Therapeutics, MicroTransponder, Neuroolutions, SanBio, Stemedica, Fujifilm Toyama Chemical Co., NeuExcell, Medtronic, and TRCare. M.A.P. has received Research Funding & Travel Grant (Bionic Vision Technologies Pty Ltd) unrelated to the contents of this manuscript.

REFERENCES

- Alexander GE, Crutcher MD, DeLong MR. Basal ganglia-thalamocortical circuits: parallel substrates for motor, oculomotor, “prefrontal” and “limbic” functions. *Progress in brain research*: Elsevier; 1991. p. 119-46.
- Alexander LD, Black SE, Patterson KK, Gao F, Danells CJ, McIlroy WE. Association between gait asymmetry and brain lesion location in stroke patients. *Stroke* 2009; 40(2): 537-44.
- Apostolova LG, Green AE, Babakchanian S, Hwang KS, Chou Y-Y, Toga AW, *et al.* Hippocampal atrophy and ventricular enlargement in normal aging, mild cognitive impairment and Alzheimer's disease. *Alzheimer disease and associated disorders* 2012; 26(1): 17.
- Azari NP, Binkofski F, Pettigrew KD, Freund HJ, Seitz RJ. Enhanced regional cerebral metabolic interactions in thalamic circuitry predicts motor recovery in hemiparetic stroke. *Human brain mapping* 1996; 4(4): 240-53.
- Baudat C, Maréchal B, Corredor-Jerez R, Kober T, Meuli R, Hagmann P, *et al.* Automated MRI-based volumetry of basal ganglia and thalamus at the chronic phase of cortical stroke. *Neuroradiology* 2020: 1-10.
- Berlin JA, Santanna J, Schmid CH, Szczech LA, Feldman HI. Individual patient \square versus group \square level data meta \square regressions for the investigation of treatment effect modifiers: ecological bias rears its ugly head. *Statistics in medicine* 2002; 21(3): 371-87.
- Bernhardt J, Hayward KS, Kwakkel G, Ward NS, Wolf SL, Borschmann K, *et al.* Agreed definitions and a shared vision for new standards in stroke recovery research: the stroke recovery and rehabilitation roundtable taskforce. *International Journal of Stroke* 2017; 12(5): 444-50.
- Binkofski F, Seitz R, Arnold S, Classen J, Benecke R, Freund HJ. Thalamic metabolism and corticospinal tract integrity determine motor recovery in stroke. *Annals of neurology* 1996; 39(4): 460-70.

Boyd LA, Hayward KS, Ward NS, Stinear CM, Rosso C, Fisher RJ, *et al.* Biomarkers of stroke recovery: Consensus-based core recommendations from the Stroke Recovery and Rehabilitation Roundtable. *Neurorehabilitation and neural repair* 2017; 31(10-11): 864-76.

Boyd LA, Winstein CJ. Providing explicit information disrupts implicit motor learning after basal ganglia stroke. *Learning & memory* 2004; 11(4): 388-96.

Brodtmann A, Khlif MS, Egorova N, Veldsman M, Bird LJ, Werden E. Dynamic Regional Brain Atrophy Rates in the First Year After Ischemic Stroke. *Stroke* 2020; 51(9): e183-e92.

Campbell BC, Ma H, Ringleb PA, Parsons MW, Churilov L, Bendzus M, *et al.* Extending thrombolysis to 4·5–9 h and wake-up stroke using perfusion imaging: a systematic review and meta-analysis of individual patient data. *The Lancet* 2019; 394(10193): 139-47.

Carmichael ST, Tatsukawa K, Katsman D, Tsuyuguchi N, Kornblum HI. Evolution of diaschisis in a focal stroke model. *Stroke* 2004; 35(3): 758-63.

Cheung DK, Climans SA, Black SE, Gao F, Szilagyi GM, Mochizuki G. Lesion characteristics of individuals with upper limb spasticity after stroke. *Neurorehabilitation and neural repair* 2016; 30(1): 63-70.

Dong Y, Dobkin BH, Cen SY, Wu AD, Winstein CJ. Motor cortex activation during treatment may predict therapeutic gains in paretic hand function after stroke. *Stroke* 2006; 37(6): 1552-5.

Egorova N, Liem F, Hachinski V, Brodtmann A. Predicted brain age after stroke. *Frontiers in Aging Neuroscience* 2019; 11: 348.

Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, *et al.* Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron* 2002; 33(3): 341-55.

Förstl H, Zerfaß R, Geiger-Kabisch C, Sattel H, Besthorn C, Hentschel F. Brain atrophy in normal ageing and Alzheimer's disease. *The British journal of psychiatry* 1995; 167(6): 739-46.

Fries W, Danek A, Scheidtmann K, Hamburger C. Motor recovery following capsular stroke: role of descending pathways from multiple motor areas. *Brain* 1993; 116(2): 369-82.

Fugl-Meyer AR, Jaasko L, Leyman I, Olsson S, Steglind S. The post-stroke hemiplegic patient. 1. a method for evaluation of physical performance. *Scandinavian journal of rehabilitation medicine* 1975; 7(1): 13-31.

Gauthier LV, Taub E, Mark VW, Barghi A, Uswatte G. Atrophy of spared gray matter tissue predicts poorer motor recovery and rehabilitation response in chronic stroke. *Stroke* 2012; 43(2): 453-7.

Glasziou P, Altman DG, Bossuyt P, Boutron I, Clarke M, Julious S, *et al.* Reducing waste from incomplete or unusable reports of biomedical research. *The Lancet* 2014; 383(9913): 267-76.

Gower A, Tiberi M. The intersection of central dopamine system and stroke: potential avenues aiming at enhancement of motor recovery. *Frontiers in synaptic neuroscience* 2018; 10: 18.

Goyal M, Menon BK, van Zwam WH, Dippel DW, Mitchell PJ, Demchuk AM, *et al.* Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *The Lancet* 2016; 387(10029): 1723-31.

Greco L, Luta G, Krzywinski M, Altman N. Analyzing outliers: robust methods to the rescue. *Nature methods* 2019; 16(4): 275-7.

Hibar DP, Stein JL, Renteria ME, Arias-Vasquez A, Desrivières S, Jahanshad N, *et al.* Common genetic variants influence human subcortical brain structures. *Nature* 2015; 520(7546): 224.

Hijdra A, Verbeeten Jr B. Leukoaraiosis and ventricular enlargement in patients with ischemic stroke. *Stroke* 1991; 22(4): 447-50.

Ioannidis J. Next-generation systematic reviews: prospective meta-analysis, individual-level data, networks and umbrella reviews. *BMJ Publishing Group Ltd and British Association of Sport and Exercise Medicine*; 2017.

Jones EG. *The thalamus*: Springer Science & Business Media; 2012.

Kassambara A. *rstatix: Pipe-Friendly Framework for Basic Statistical Tests*. 2019.

Kim S, Ashe J, Hendrich K, Ellermann J, Merkle H, Ugurbil K, *et al.* Functional magnetic resonance imaging of motor cortex: hemispheric asymmetry and handedness. *Science* 1993; 261: 615-7.

Koller M. robustlmm: an R package for robust estimation of linear mixed-effects models. *Journal of statistical software* 2016; 75(6): 1-24.

Kuceyeski A, Navi BB, Kamel H, Raj A, Relkin N, Togliola J, *et al.* Structural connectome disruption at baseline predicts 6-months post-stroke outcome. *Human brain mapping* 2016; 37(7): 2587-601.

Kwakkel G, Lannin NA, Borschmann K, English C, Ali M, Churilov L, *et al.* Standardized measurement of sensorimotor recovery in stroke trials: consensus-based core recommendations from the stroke recovery and rehabilitation roundtable. *Neurorehabilitation and neural repair* 2017; 31(9): 784-92.

Lanciego JL, Luquin N, Obeso JA. Functional neuroanatomy of the basal ganglia. *Cold Spring Harbor perspectives in medicine* 2012; 2(12): a009621.

Lee KB, Kim JS, Hong BY, Kim YD, Hwang BY, Lim SH. The motor recovery related with brain lesion in patients with intracranial hemorrhage. *Behavioural Neurology* 2015; 2015.

Lenzi G, Padovani A, Di Piero V, Bragoni M, De Biasi C, Altieri M, *et al.* White matter changes and ventricular enlargement on magnetic resonance imaging: Comparison between normal elderly subjects and patients with cerebrovascular disorders. *Journal of Stroke and Cerebrovascular Diseases* 1994; 4(4): 245-9.

Liew S-L, Zavaliangos-Petropulu A, Jahanshad N, Lang CE, Hayward KS, Lohse K, *et al.* The ENIGMA Stroke Recovery Working Group: Big data neuroimaging to study brain-behavior relationships after stroke. *Human brain mapping* 2020.

Lopes MA, Firbank MJ, Widdrington M, Blamire AM, Kalaria RN, T O'Brien J. Post-stroke dementia: the contribution of thalamus and basal ganglia changes. *International psychogeriatrics* 2012; 24(4): 568.

Lyle RC. A performance test for assessment of upper limb function in physical rehabilitation treatment and research. *International journal of rehabilitation research* 1981; 4(4): 483-92.

Mahajan KR, Nakamura K, Cohen JA, Trapp BD, Ontaneda D. Intrinsic and extrinsic mechanisms of thalamic pathology in multiple sclerosis. *Annals of Neurology* 2020.

Mani S, Mutha PK, Przybyla A, Haaland KY, Good DC, Sainburg RL. Contralesional motor deficits after unilateral stroke reflect hemisphere-specific control mechanisms. *Brain* 2013; 136: 1288-303.

McDougall J, Wright V, Rosenbaum P. The ICF model of functioning and disability: incorporating quality of life and human development. *Developmental neurorehabilitation* 2010; 13(3): 204-11.

Murphy TH, Corbett D. Plasticity during stroke recovery: from synapse to behaviour. *Nature reviews neuroscience* 2009; 10(12): 861.

Nieuwenhuis R, Te Grotenhuis H, Pelzer B. Influence. ME: tools for detecting influential data in mixed effects models. 2012.

Pinheiro J BD, DebRoy S, Sarkar D, R Core Team. *_nlme: Linear and Nonlinear Mixed Effects Models_*. R package version 2020; 3: 1-149.

Robbins TW, Everitt BJ. Functions of dopamine in the dorsal and ventral striatum. *Seminars in Neuroscience*; 1992: Elsevier; 1992. p. 119-27.

Rochat L, Van der Linden M, Renaud O, Epiney J-B, Michel P, Sztajzel R, *et al*. Poor reward sensitivity and apathy after stroke: implication of basal ganglia. *Neurology* 2013; 81(19): 1674-80.

Sainburg RL, Maenza C, Winstein C, Good D. Motor lateralization provides a foundation for predicting and treating non-paretic arm motor deficits in stroke. *Progress in Motor Control*: Springer; 2016. p. 257-72.

Sawada M, Kato K, Kunieda T, Mikuni N, Miyamoto S, Onoe H, *et al*. Function of the nucleus accumbens in motor control during recovery after spinal cord injury. *Science* 2015; 350(6256): 98-101.

Schaefer SY, Haaland KY, Sainburg RL. Ipsilesional motor deficits following stroke reflect hemispheric specializations for movement control. *Brain* 2007; 130(8): 2146-58.

Shelton FtdN, Reding MJ. Effect of lesion location on upper limb motor recovery after stroke. *Stroke* 2001; 32(1): 107-12.

Team RC. R: A language and environment for statistical computing. . R Foundation for Statistical Computing, Vienna, Austria 2020.

Thompson PM, Jahanshad N, Ching CRK, Salminen LE, Thomopoulos SI, Bright J, *et al.* ENIGMA and global neuroscience: A decade of large-scale studies of the brain in health and disease across more than 40 countries. *Translational Psychiatry* 2020; 10(1): 100.

Van Meer MP, Van Der Marel K, Wang K, Otte WM, El Bouazati S, Roeling TA, *et al.* Recovery of sensorimotor function after experimental stroke correlates with restoration of resting-state interhemispheric functional connectivity. *Journal of Neuroscience* 2010; 30(11): 3964-72.

Virani SS, Alonso A, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP, *et al.* Heart disease and stroke statistics—2020 update: a report from the American Heart Association. *Circulation* 2020: E139-E596.

Ward NS. Restoring brain function after stroke—bridging the gap between animals and humans. *Nature Reviews Neurology* 2017; 13(4): 244.

Wickham H, Averick M, Bryan J, Chang W, McGowan LDA, François R, *et al.* Welcome to the Tidyverse. *Journal of Open Source Software* 2019; 4(43): 1686.

Wickham H, Francois R, Henry L, Müller K. dplyr: A grammar of data manipulation. R package version 102 2020.

Widmer M, Lutz K, Luft AR. Reduced striatal activation in response to rewarding motor performance feedback after stroke. *NeuroImage: Clinical* 2019; 24: 102036.

Wolf SL, Catlin PA, Ellis M, Archer AL, Morgan B, Piacentino A. Assessing Wolf motor function test as outcome measure for research in patients after stroke. *Stroke* 2001; 32(7): 1635-9.

SUPPLEMENTARY MATERIALS

COHORT ID	SITE	COUNTRY	N	PRIMARY SENSORIMOTOR MEASURE
1	UNIVERSITY OF SOUTHERN CALIFORNIA	USA	39	FMA-UE
2	UNIVERSITY OF SOUTHERN CALIFORNIA	USA	12	FMA-UE
3	UNIVERSITY OF SOUTHERN CALIFORNIA	USA	14	FMA-UE
4	UNIVERSITY OF TUEBINGEN	GERMANY	19	FMA-UE
7	UNIVERSITY COLLEGE LONDON	UK	42	Action Research Arm Test
8	UNIVERSITY OF SOUTHERN CALIFORNIA	USA	6	Manual Muscle Test
9	UNIVERSITY OF OSLO	NORWAY	93	NIHSS Motor Score
10	TIANJIN MEDICAL UNIVERSITY	CHINA	24	FMA-UE
11	TIANJIN MEDICAL UNIVERSITY	CHINA	29	FMA-UE
12	UNIVERSITY OF AUCKLAND	NEW ZEALAND	57	FMA-UE
13	UNIVERSITY OF AUCKLAND	NEW ZEALAND	44	FMA-UE
15	MEDICAL UNIVERSITY OF SOUTH CAROLINA	USA	14	FMA-UE
17	MEDICAL UNIVERSITY OF SOUTH CAROLINA	USA	16	FMA-UE
18	MEDICAL UNIVERSITY OF SOUTH CAROLINA	USA	10	FMA-UE
19	UNIVERSITY OF GRIEFSWALD	GERMANY	13	Motricity Index
20	UNIVERSITY OF GRIEFSWALD	GERMANY	21	Motricity Index
22	UNIVERSITY OF GRIEFSWALD	GERMANY	17	Bogenhausen Dysphagia Score
23	UNIVERSITY OF THE SCIENCES	USA	13	FMA-UE
24	EMORY UNIVERSITY	USA	21	Manual Muscle Test
25	UNIVERSITY OF TORONTO	CANADA	26	Grip Strength
26	SAO PAULO UNIVERSITY	BRAZIL	24	FMA-UE
28	UNIVERSITY OF CALIFORNIA, IRVINE	USA	26	FMA-UE
31	UNIVERSITY OF CALIFORNIA, IRVINE	USA	35	FMA-UE
32	UNIVERSITY OF CALIFORNIA, IRVINE	USA	7	FMA-UE
34	UNIVERSITY OF SOUTHERN CALIFORNIA	USA	15	FMA-UE
35	HOSPITAL ISRAELITA ALBERT EINSTEIN	BRAZIL	15	FMA-UE
38	IRCCS SANTA LUCIA FOUNDATION	ITALY	79	Barthel Index
41	FLOREY INSTITUTE OF NEUROSCIENCE AND MENTAL HEALTH	AUSTRALIA	91	NIHSS Total

Supplementary Table 1. Additional cohort details. Research sites (institutions, countries) and primary sensorimotor assessment used are listed for each cohort. FMA-UE = Fugl-Meyer Assessment of Upper Extremities.

EARLY STROKE (≤ 21 days)

Brain Region	n	beta (CI)	SE	df	t-value	p-value	<i>d</i>	Significant covariates
<i>Ipsilesional</i>								
Caudate	135	-0.09 (-0.67-0.48)	0.29	125	-0.32	0.749	-0.06	ICV
Lateral ventricle	182	0.25 (-0.11-0.61)	0.18	172	1.37	0.173	0.21	Age, ICV
Nucleus accumbens	165	0.19 (-0.23-0.60)	0.21	155	0.90	0.369	0.14	Age
Pallidum	157	0.12 (-0.39-0.63)	0.26	147	0.46	0.644	0.08	ICV
Putamen	143	0.25 (-0.28-0.79)	0.27	133	0.93	0.354	0.16	Age, ICV
Thalamus	137	0.79 (0.38-1.20)	0.21	128	3.82	<0.001	0.68	Age, ICV
<i>Contralesional</i>								
Caudate	147	0.17 (-0.29-0.64)	0.24	137	0.74	0.461	0.13	ICV
Lateral ventricle	182	0.19 (-0.20-0.57)	0.19	172	0.96	0.337	0.15	Age, ICV
Nucleus accumbens	170	0.30 (-0.09-0.69)	0.20	160	1.53	0.127	0.24	Age
Pallidum	171	0.65 (0.19-1.11)	0.23	161	2.79	0.006	0.44	ICV
Putamen	158	0.26 (-0.21-0.72)	0.24	148	1.10	0.274	0.18	Age, ICV
Thalamus	150	0.20 (-0.28-0.67)	0.24	141	0.82	0.411	0.14	Age, ICV

Supplementary Table 2. Relationships between non-lesioned subcortical volumes and sensorimotor behavior in early stroke (≤ 21 days; $n=179$). Results from linear mixed-effects models of individuals with early stroke. Uncorrected p-values shown. Results in bold indicate significance with a Bonferroni correction for multiple comparisons ($p<0.004$). The beta coefficient for sensorimotor behavior (beta) with 95% confidence interval (CI), along with the sample size (n), standard error (SE), degrees of freedom (df), standardized effect size (*d*), t-value, and uncorrected p-value are reported, in addition to significant fixed covariates including age, sex, and intracranial volume (ICV).

Brain Region	n	df	Interaction between Sensorimotor Behavior & Lesioned Hemisphere			Main Effect Lesioned Hemisphere			Main Effect Sensorimotor Behavior				Significant Covariates
			beta	SE	p-value	beta	SE	p-value	beta	SE	p-value	d	
<i>Ipsilesional</i>													
Caudate	193	167	0.14	0.52	0.789	-0.17	0.44	0.693	0.26	0.28	0.365	0.14	ICV
Lateral ventricle	404	376	0.21	0.29	0.487	-0.13	0.23	0.572	-0.70	0.17	<0.001	-0.42	Age, ICV
Nucleus accumbens	289	262	0.21	0.39	0.593	-0.13	0.31	0.684	0.73	0.23	0.002	0.39	Age
Pallidum	225	198	-0.53	0.48	0.272	0.52	0.39	0.191	0.35	0.27	0.207	0.18	ICV
Putamen	207	181	-0.03	0.53	0.953	-0.28	0.44	0.525	0.90	0.29	0.002	0.47	ICV
Thalamus	169	144	0.17	0.63	0.792	-0.95	0.51	0.065	0.05	0.32	0.887	0.02	Age, ICV
<i>Contralesional</i>													
Caudate	345	318	0.12	0.33	0.731	-0.14	0.26	0.583	0.06	0.21	0.760	0.03	ICV
Lateral ventricle	404	376	0.27	0.28	0.343	0.06	0.21	0.789	-0.35	0.16	0.030	-0.22	Age, ICV
Nucleus accumbens	344	317	0.38	0.35	0.282	-0.32	0.27	0.236	0.18	0.22	0.423	0.09	Age
Pallidum	359	332	0.07	0.32	0.819	-0.43	0.25	0.083	0.09	0.20	0.672	0.05	Sex, ICV
Putamen	355	328	-0.08	0.31	0.796	0.14	0.24	0.553	0.23	0.20	0.243	0.13	Age, ICV
Thalamus	329	302	0.25	0.30	0.405	0.47	0.23	0.045	-0.17	0.18	0.353	-0.11	Age, ICV

Supplementary Table 3. Relationships between non-lesioned subcortical volumes, lesioned hemisphere, and sensorimotor behavior in chronic stroke. Results from linear mixed-effects models including an interaction term between lesioned hemisphere and sensorimotor behavior in people with chronic stroke. There were no significant interactions or main effects of lesioned hemisphere. The significant main effects for sensorimotor behavior remained, similar to those shown in Table 2. Results in bold indicate significance with a Bonferroni correction for multiple comparisons ($p < 0.004$). The beta coefficients (beta), standard error (SE), and uncorrected p-value for the interaction between lesioned hemisphere and sensorimotor behavior, as well as main effects for lesioned hemisphere and sensorimotor behavior, are reported, along with the sample size (n), degrees

of freedom (df), standardized effectsize (d), and significant fixed covariates including age, sex, and intracranial volume (ICV).

Difference in sensorimotor behavior between chronic left and right hemisphere stroke groups

We examined whether there was a difference in sensorimotor behavior between chronic left hemisphere stroke (LHS; n=214) and right hemisphere stroke (RHS; n=190). The distribution of sensorimotor behavior scores violated the Wilkes-Shapiro test of normality for both groups (LHS: $W=0.89$, $p<0.001$, RHS: $W=0.89$, $p<0.001$). We therefore used a nonparametric Wilcoxon rank sum test to compare independent group samples. The median sensorimotor behavior score in LHS was 0.80 (IQR=0.39) and in RHS was 0.74 (IQR=0.49). The Wilcoxon test showed no significant effect of lesioned hemisphere ($p=0.29$, effect size $r=0.053$).

ALL STROKE

Brain Region	n	beta (CI)	SE	df	t-value	p-value	d	Significant covariates
<i>Ipsilesional</i>								
Caudate	482	0.15 (-0.18-0.48)	0.17	451	0.89	0.375	0.08	Sex, ICV
Lateral ventricle	828	-0.39 (-0.63--0.16)	0.12	796	-3.26	0.001	-0.23	Age, ICV
Nucleus accumbens	655	0.41 (0.13-0.68)	0.14	624	2.91	0.004	0.23	Age
Pallidum	546	0.29 (-0.04-0.61)	0.16	515	1.75	0.081	0.15	ICV
Putamen	490	0.64 (0.28-1.00)	0.18	459	3.53	<0.001	0.33	Age, ICV
Thalamus	462	0.58 (0.25-0.91)	0.17	433	3.47	0.001	0.33	Age, ICV
<i>Contralesional</i>								
Caudate	689	0.01 (-0.26-0.28)	0.14	658	0.08	0.939	0.01	ICV
Lateral ventricle	828	-0.26 (-0.49--0.03)	0.11	796	-2.27	0.024	-0.16	Age, ICV
Nucleus accumbens	727	0.10 (-0.15-0.36)	0.13	696	0.78	0.436	0.06	Age, ICV
Pallidum	743	0.28 (0.02-0.54)	0.13	712	2.08	0.038	0.16	Age, ICV
Putamen	704	0.19 (-0.07-0.45)	0.13	673	1.45	0.147	0.11	Age, ICV
Thalamus	663	-0.02 (-0.28-0.24)	0.13	633	-0.13	0.898	-0.01	Age, ICV

Supplementary Table 4. Relationships between non-lesioned subcortical volumes and sensorimotor behavior post-stroke across all times after stroke (N=828). Results from linear mixed-effects models of individuals across all times after stroke. Uncorrected p-values shown. Results in bold indicate significance with a Bonferroni correction for multiple comparisons ($p<0.004$). The beta coefficient for sensorimotor behavior (beta) with 95% confidence interval (CI), along with the sample size (n), standard error (SE), degrees of freedom (df), standardized effect size (d), t-value, and uncorrected p-value are reported, in addition to significant fixed covariates including age, sex, and intracranial volume (ICV).

ROBUST REGRESSIONS FOR SUBACUTE AND CHRONIC STROKE

SUBACUTE STROKE (≤ 90 days)						
Brain Region	n	beta (CI)	SE	t-value	p-value	Significant covariates
<i>Ipsilesional</i>						
Caudate	194	0.07 (-0.37 – 0.51)	0.22	0.33	0.742	ICV
Lateral ventricle	274	0.15 (-0.12 – 0.42)	0.14	1.11	0.267	Age, ICV
Nucleus accumbens	245	0.18 (-0.21 – 0.57)	0.20	0.90	0.366	Age
Pallidum	223	0.17 (-0.22 – 0.57)	0.20	0.87	0.387	ICV
Putamen	201	0.40 (-0.06 – 0.86)	0.23	1.72	0.086	Age, ICV
Thalamus	210	0.69 (0.26 – 1.11)	0.22	3.17	0.002	Age, ICV
<i>Contralesional</i>						
Caudate	219	0.23 (-0.20 – 0.66)	0.22	1.04	0.297	ICV
Lateral ventricle	274	0.06 (-0.24 – 0.36)	0.15	0.37	0.714	Age, ICV
Nucleus accumbens	253	0.19 (-0.17 – 0.55)	0.18	1.05	0.294	Age, ICV
Pallidum	250	0.59 (0.17 – 1.01)	0.22	2.73	0.006	ICV
Putamen	229	0.31 (-0.08 – 0.70)	0.20	1.57	0.116	Age, ICV
Thalamus	217	0.09 (-0.25 – 0.44)	0.18	0.52	0.603	Age, ICV
CHRONIC STROKE (≥ 180 days)						
Brain Region	n	beta (CI)	SE	t-value	p-value	Significant covariates
<i>Ipsilesional</i>						
Caudate	193	0.31 (-0.21 – 0.83)	0.26	1.18	0.240	ICV
Lateral ventricle	404	-0.71 (-1.02 – -0.41)	0.16	-4.58	<0.001	Age, ICV
Nucleus accumbens	289	0.64 (0.20 – 1.07)	0.22	2.88	0.004	Age
Pallidum	225	0.50 (0.04 – 0.97)	0.24	2.13	0.033	ICV
Putamen	207	1.02 (0.48 – 1.55)	0.27	3.72	<0.001	Age, ICV
Thalamus	169	0.11 (-0.53 – 0.75)	0.33	0.33	0.740	Age
<i>Contralesional</i>						
Caudate	345	0.14 (-0.27 – 0.55)	0.21	0.67	0.501	ICV
Lateral ventricle	404	-0.36 (-0.63 – -0.09)	0.14	-2.62	0.009	Age, ICV
Nucleus accumbens	344	0.22 (-0.19 – 0.63)	0.21	1.07	0.285	Age
Pallidum	359	0.32 (-0.01 – 0.64)	0.17	1.91	0.056	Age, ICV
Putamen	355	0.22 (-0.15 – 0.60)	0.19	1.16	0.244	Age, ICV
Thalamus	329	-0.20 (-0.57 – 0.17)	0.19	-1.06	0.288	Age, ICV

Supplementary Table 5. Robust regressions to examine relationships between non-lesioned subcortical volumes and sensorimotor behavior in subacute and chronic stroke. Results from robust linear mixed-effects models of individuals with subacute stroke (top) and chronic stroke (bottom). Results in bold indicate significance with a Bonferroni correction for multiple comparisons ($p < 0.004$). The beta coefficient for sensorimotor behavior (beta) with 95% confidence interval (CI), along with the sample size (n), standard error (SE), t-value, and uncorrected p-value are reported, in addition to significant fixed covariates including age, sex, and intracranial volume (ICV).

ROBUST REGRESSIONS FOR CHRONIC SENSORIMOTOR IMPAIRMENT AND ACTIVITY LIMITATIONS

CHRONIC SENSORIMOTOR IMPAIRMENT						
Brain Region	n	beta (CI)	SE	t-value	p-value	Significant covariates
<i>Ipsilesional</i>						
Caudate	194	0.98 (0.08 – 1.88)	0.46	2.14	0.032	ICV
Lateral ventricle	274	-0.70 (-1.13 – -0.27)	0.22	-3.2	0.001	Age, ICV
Nucleus accumbens	245	0.63 (0.08 – 1.18)	0.28	2.24	0.025	Age
Pallidum	223	0.79 (0.07 – 1.51)	0.37	2.16	0.031	-
Putamen	201	1.54 (0.60 – 2.47)	0.48	3.23	0.001	-
Thalamus	210	0.56 (-0.33 – 1.44)	0.45	1.23	0.22	-
<i>Contralateral</i>						
Caudate	219	0.02 (-0.52 – 0.56)	0.27	0.08	0.939	ICV
Lateral ventricle	274	-0.46 (-0.80 – -0.13)	0.17	-2.72	0.006	Age, ICV
Nucleus accumbens	253	0.22 (-0.32 – 0.76)	0.28	0.80	0.423	Age
Pallidum	250	0.31 (-0.13 – 0.75)	0.22	1.39	0.166	ICV
Putamen	229	0.02 (-0.46 – 0.51)	0.25	0.09	0.932	Age, ICV
Thalamus	217	-0.34 (-0.84 – 0.15)	0.25	-1.36	0.173	Age, ICV
ACTIVITY LIMITATIONS						
Brain Region	n	beta (CI)	SE	t-value	p-value	Significant covariates
<i>Ipsilesional</i>						
Caudate	193	-0.48 (-1.64 – 0.69)	0.60	-0.80	0.425	-
Lateral ventricle	404	-0.70 (-1.32 – -0.09)	0.31	-2.24	0.025	Age, ICV
Nucleus accumbens	289	0.68 (-0.30 – 1.67)	0.50	1.36	0.172	-
Pallidum	225	0.88 (-0.04 – 1.80)	0.47	1.88	0.060	-
Putamen	207	0.87 (-0.32 – 2.05)	0.60	1.43	0.152	-
Thalamus	169	1.19 (-0.08 – 2.46)	0.65	1.84	0.066	-
<i>Contralateral</i>						
Caudate	345	0.14 (-0.74 – 1.02)	0.45	0.32	0.750	ICV
Lateral ventricle	404	-0.72 (-1.31 – -0.13)	0.30	-2.38	0.017	Age, ICV
Nucleus accumbens	344	-0.33 (-1.13 – 0.46)	0.41	-0.82	0.413	Age
Pallidum	359	-0.07 (-0.93 – 0.79)	0.44	-0.16	0.874	Sex
Putamen	355	0.33 (-0.54 – 1.20)	0.44	0.75	0.454	Age
Thalamus	329	0.19 (-0.57 – 0.95)	0.39	0.49	0.626	Age, Sex

Supplementary Table 6. Robust regressions to examine relationships between non-lesioned subcortical volumes and two measures of sensorimotor behavior (impairment, activity limitations). Results from robust linear mixed-effects models in individuals with chronic stroke showing sensorimotor impairment (top) compared to activity limitations (bottom). Results in bold indicate significance with a Bonferroni correction for multiple comparisons ($p < 0.004$). The beta coefficient for sensorimotor behavior (beta) with 95% confidence interval (CI), along with the sample size (n), standard error (SE), t-value, and uncorrected p-value are reported, in addition to significant fixed covariates including age, sex, and intracranial volume (ICV).