Non-canonical odor coding ensures unbreakable mosquito attraction to humans

Meg A. Younger^{1,2,7*}, Margaret Herre^{1,2,7*}, Alison R. Ehrlich^{1,4}, Zhongyan Gong^{1,5}, Zachary N. Gilbert¹, Saher Rahiel¹, Benjamin J. Matthews^{1,3,6}, Leslie B. Vosshall^{1,2,3*}

SUMMARY

Female *Aedes aegypti* mosquitoes show strong innate attraction to humans. This chemosensory behavior is critical to species survival because females require a blood-meal to reproduce. Humans, the preferred host of *Ae. aegypti*, produce a complex blend of odor cues along with carbon dioxide (CO₂) that attracts females ready to bite. Mosquitoes detect these cues with heteromeric ligand-gated ion channels encoded by three different chemosensory receptor gene families. A common theme in other species is that olfactory neurons express a single receptor that defines their chemical specificity and that they extend axons that converge upon dedicated glomeruli in the first sensory processing center in the brain. Such an organization permits the brain to segregate olfactory information and monitor activity of individual glomeruli to interpret what smell has been encountered. We have discovered that *Ae. aegypti* uses an entirely different organizational principle for its olfactory system. Using genetic strains that label subpopulations of olfactory neurons, we found that many neurons co-express multiple members of at least two of the chemosensory receptor families. This unexpected co-expression is functional, as assessed by *in vivo* calcium imaging showing that a given glomerulus is activated by multiple ligands detected by different receptor families. This has direct functional consequences for mosquito behavior. Mutant mosquitoes that cannot sense CO₂ can be behaviorally activated by a volatile amine that stimulates the CO₂ glomerulus. This non-canonical olfactory system organization featuring overlapping receptor expression may explain the female mosquito's robust and 'unbreakable' attraction to humans.

Key words: *Aedes aegypti*; mosquito; odor coding; olfaction; GCaMP imaging; behavior

INTRODUCTION

Increased global travel, a growing world population, and rising temperatures are all combining to increase the emergence and transmission of novel disease-causing pathogens. This includes vector-borne diseases, in which pathogens are spread by "vector" organisms such as mosquitoes, ticks, sandflies, and fleas. Diseases spread by these arthropods collectively account for more than 700,000 deaths every year (WHO, 2020). Female *Aedes aegypti* mosquitoes spread arboviruses including dengue, Zika, yellow fever, and chikungunya. Only female mosquitoes bite, and they do so because they require a blood-meal for reproduction (Allan et al., 1987). *Ae. aegypti* prefer to bite human hosts, which contributes to their effectiveness as a disease vector (Brown et al., 2014; Gouck, 1972; McBride et al., 2014). To

identify human hosts, mosquitoes rely heavily on chemosensory cues, including carbon dioxide (CO₂) emitted from breath, and human body odor, which is a mixture of hundreds of different individual odorants including lactic acid and volatile amines such as ammonia (Acree et al., 1968; Bernier et al., 2000; Davis, 1984; Gallagher et al., 2008; Geier et al., 1999; Smallegange et al., 2005; Smith et al., 1970).

Insects detect such chemosensory cues using membrane receptors encoded by three large gene families, Olfactory Receptors (ORs), Ionotropic Receptors (IRs), and Gustatory Receptors (GRs). All three gene families encode ionotropic ligand-gated ion channels, in contrast to the metabotropic G

¹Laboratory of Neurogenetics and Behavior, The Rockefeller University, New York, NY 10065 USA, ²Kavli Neural Systems Institute, New York, NY 10065 USA, ³Howard Hughes Medical Institute, New York, NY 10065 USA ⁴Current address: New York University School of Medicine, 550 First Ave, New York, NY 10016 ⁵Current address: Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, CA 94720 USA ⁶Current address: University of British Columbia, Department of Zoology, Vancouver, BC V6S 0K3 Canada ⁷These authors contributed equally *Correspondence: meg.younger@gmail.com (M.A.Y.), margo.herre@gmail.com (M.H.), leslie.vosshall@rockefeller.edu (L.B.V.)

protein-coupled odorant receptors utilized by vertebrates and *C. elegans* nematodes (Ihara et al., 2013). ORs are odorant-gated ion channels (Butterwick et al., 2018; Sato et al., 2008; Wicher et al., 2008) that are formed by a heteromultimeric complex of the conserved co-receptor Orco and a ligand-sensitive OR (Benton et al., 2006; Larsson et al., 2004; Neuhaus et al., 2005; Sato et al., 2008). IRs are variant ionotropic glutamate receptors that are formed by one or more of three conserved co-receptors, Ir25a, Ir8a, and Ir76b, and ligand-sensitive subunits that determine the selectivity of the receptor complex (Abuin et al., 2011; Benton et al., 2009; Silbering et al., 2011). Although GRs are primarily taste receptors (Clyne et al., 2000; Montell, 2009; Scott et al., 2001), several GRs form a complex that detects carbon dioxide (CO₂) (Jones et al., 2007; Kwon et al., 2007). CO₂ is an important volatile human host cue that activates and attracts mosquitoes (Gillies, 1980). In *Ae. aegypti*, *Gustatory Receptor 3* (*Gr3*) encodes an essential subunit of the CO₂ receptor, and *Gr3* mutant mosquitoes lose all sensitivity to CO₂ (McMeniman et al., 2014).

Because mosquitoes specialize on humans and require blood to reproduce, the drive to find humans is strong and innate. Indeed, even mosquitoes genetically engineered to eliminate genes critical for peripheral detection of host sensory cues are able to find and bite people. Animals lacking the olfactory receptor co-receptor (*Orco*), the obligate co-receptor required for the function of the entire family of ORs, show strong attraction to humans (DeGennaro et al., 2013). Deleting the *Ir8a* co-receptor reduces but does not eliminate attraction to humans (Raji et al., 2019). Similarly, while mosquitoes lacking the obligate CO₂ receptor subunit, *Gr3*, do not respond to CO₂ and show impaired behavioral responses in laboratory assays, they are highly effective in finding humans in a more naturalistic semi-field setting (McMeniman et al., 2014). It is not known how the mosquito olfactory system is seemingly infallible in its ability to detect humans.

The cloning of the first odorant receptors by Buck and Axel in 1991 (Buck and Axel, 1991) led to the subsequent discovery that each olfactory sensory neuron expresses a unique odorant receptor that specifies its functional properties. In the well-studied olfactory systems of *Drosophila melanogaster* flies (Clyne et al., 1999; Gao and Chess, 1999; Vosshall et al., 1999) and *Mus musculus* mice (Bashkurova and Lomvardas, 2019; Chess et al., 1994), olfactory sensory neurons are thought to express a single olfactory receptor. In both species, neurons expressing a given receptor project axons to dedicated olfactory glomeruli in the first sensory processing center in the brain, the antennal lobe in insects (Couto et al., 2005; Fishilevich and Vosshall, 2005; Vosshall et al., 2000) and the olfactory bulb in vertebrates (Mombaerts et al., 1996; Ressler et al., 1994; Vassar et al., 1994). This “one-receptor – to – one – neuron – to – one – glomerulus” organization is believed to be a widespread motif in both invertebrate and vertebrate olfactory systems, and the convergence onto discrete glomeruli is hypothesized to permit the brain to parse which subpopulation of olfactory

neurons is activated by a given odor (Rubin and Katz, 1999; Uchida et al., 2000; Wang et al., 2003).

Consistent with this organization, the number of expressed chemosensory receptors in the OR and IR gene families roughly correlates to the number of olfactory glomeruli. This holds true in *Drosophila melanogaster* flies (~60 receptors/~55 glomeruli) (Benton et al., 2009; Laissue et al., 1999; Robertson et al., 2003), the honey bee *Apis mellifera* (~180 receptors/~160 glomeruli) (Flanagan and Mercer, 1989; Robertson et al., 2010), and the tobacco hornworm *Manduca sexta* (~60 receptors/~70 glomeruli) (Grosse-Wilde et al., 2011). Based on these studies, it is widely thought that merely counting the number of antennal lobe glomeruli in a new species would be reasonably predictive of the number of chemosensory receptors found in its genome. In *Ae. aegypti*, however, there is a striking mismatch between the number of expressed chemosensory receptors and the number of antennal lobe glomeruli, with at least twice as many receptors as available glomeruli (Bohbot et al., 2007; Ignell et al., 2005; Matthews et al., 2018; Shankar and McMeniman, 2020; Zhao et al., 2020). How is the mosquito olfactory system organized to accommodate so many receptors and does this departure from rules established in other species explain their exquisite ability to locate human hosts?

In this study, we developed a CRISPR-Cas9-based genetic knock-in strategy in *Ae. aegypti* to generate a suite of genetically-modified mosquito strains that label molecularly distinct populations of olfactory sensory neurons. We used these strains to understand how human odor is encoded in the mosquito olfactory system. Combining anatomical analysis of these lines with extensive RNA *in situ* expression analysis, we found that the olfactory system of *Ae. aegypti* does not have the expected “one-receptor-to-one-neuron-to-one-glomerulus” organization seen in other organisms. We frequently observed co-expression of multiple chemosensory receptors from at least two of the three receptor gene superfamilies within individual olfactory sensory neurons. Antennal lobe glomeruli were often innervated by olfactory sensory neurons expressing different receptors. We hypothesized that this unconventional organization underlies the robust, seemingly unbreakable, properties of the *Ae. aegypti* olfactory system in detecting human odor. To test this, we examined a minimal mixture that drives host-seeking behavior, the combination of CO₂ and lactic acid. We found that the neurons that sense CO₂ express not only the CO₂ receptor gene *Gr3*, but also members of the IR family of chemosensory receptors that detect acids and amines. Using *in vivo* functional calcium imaging we found that these CO₂ neurons also sense volatile amines, consistent with their co-expression of *Gr3* and IRs. Moreover, triethyl amine can substitute for CO₂ to drive attraction to lactic acid in mutants that cannot sense CO₂. This sensory organization, in which multiple receptors that respond to very different types of chemosensory stimuli are co-expressed, suggests a redundancy in the code for human odor. We speculate that this design supports the robust human host-seeking seen in this olfactory specialist.

RESULTS

Mismatch in chemosensory receptor and olfactory glomerulus number suggests a novel olfactory organization

In the mosquito, olfactory cues are sensed by olfactory sensory neurons in the antenna and the maxillary palp, whose axons project to the ipsilateral antennal lobe of the brain (Distler and Boeckh, 1997; Ignell et al., 2005) (Figure 1A-B, S1A-C). The antennal lobe, the insect equivalent of the vertebrate olfactory bulb, is organized into discrete olfactory glomeruli in which axons from peripheral olfactory sensory neurons terminate and synapse with local interneurons and projection neurons that relay olfactory information to the higher brain (Stocker, 1994). Previous studies used morphological criteria to define 50 (Ignell et al., 2005), 60 (Zhao et al., 2020), or 81 (Shankar and McMeniman, 2020) discrete olfactory glomeruli in the female *Ae. aegypti* antennal lobe. In this study, we define approximately 65 olfactory glomeruli (64.9 ± 0.9 , mean $\pm$ SEM), obtained by counting antennal lobe glomeruli in the left hemisphere of 12 female *Ae. aegypti* brains stained to reveal synaptic neuropil (Figure 1B,I,K, S2-5). The glomerulus count ranged from 60-72 glomeruli per antennal lobe, indicating a high level of variability in the organization of the antennal lobe (see Methods). We generated 3-D reconstructions of complete antennal lobes and saw considerable variability in the size and shape of the glomeruli (Figure S1). We were able to consistently identify certain landmark glomeruli, most notably the three glomeruli that are innervated by the maxillary palp (Ignell et al., 2005; Shankar and McMeniman, 2020) (Figure S1D-K).

The canonical “one-receptor-to-one-neuron-to-one glomerulus” organization posits that the number of chemosensory receptors should roughly match the number of glomeruli in the antennal lobe (Figure 1D). While there is not yet a clear consensus on the number of olfactory glomeruli in *Ae. aegypti*, it ranges from 50 to 81. How does this relate to the number of chemosensory receptors expressed? In the updated *Ae. aegypti* genome (Matthews et al., 2018), there are 117 OR, 135 IR, and 72 GR genes for a total of 324 structural genes that could function in the olfactory system (Figure 1E). We have reanalyzed published RNA expression data (Matthews et al., 2016) using multiple thresholds to estimate the number of receptors expressed in the antenna and maxillary palp. Even at the conservative threshold of 5 transcripts per million (TPM), the mosquito olfactory system expresses 102 chemosensory receptors, and moving the threshold to 2, 1, or 0.5 TPM increases the number of receptors plausibly expressed to 134, 156, and 178, respectively (Figure 1F,G, Data File 1). Thus, there are many more chemosensory receptors expressed in the olfactory system than available antennal lobe glomeruli, suggesting that the organization of the *Ae. aegypti* olfactory system must differ from the canonical scheme. We speculate that the mismatch can be resolved by expressing multiple receptors per neuron or having multiple different types of neurons co-converge on a single glomerulus or both (Figure 1H).

To begin to distinguish between these two organizational principles, we generated a collection of CRISPR-Cas9 gene-targeted strains that label subpopulations of olfactory neurons using the Q-system, a binary expression system similar to Gal4/UAS (Brand and Perrimon, 1993) that uses cell type-specific expression of the QF2 transcription factor to induce expression of an effector from the QF2 binding QUAS enhancer (Potter et al., 2010). We introduced an in-frame insertion that replaced the stop codon of each of the co-receptors *Orco*, *Ir25a*, *Ir8a*, and *Ir76b*, as well as the CO₂ receptor subunit *Gr3* with the transcription factor QF2 (Figure 1I, S2-S6. See Data File 1 for a full description of all genotypes by figure) (Matthews et al., 2019; Potter et al., 2010; Riabinina et al., 2016). These five new gene-sparing knock-in strains were designed to cause minimal disruption to the locus to increase the likelihood that they would faithfully report expression of the endogenous gene. We crossed these QF2 driver lines individually to a QUAS-CD8:GFP reporter to label neuronal membranes and visualized axonal projection patterns in the antennal lobe.

Orco, *Ir25a*, *Ir8a*, and *Ir76b* co-receptor driver lines were expressed in olfactory sensory neurons with distinct projection patterns in the antennal lobe (Figure 1I-K). Unexpectedly, neurons that expressed *Ir25a* projected to almost all of the glomeruli in the antennal lobe ($89.9 \pm 1.4\%$, mean $\pm$ SEM, $n = 3$) (Figure 1I-K, S3), and expression overlapped extensively with glomeruli labeled by *Orco* (Figure 1I-K, S2, S3). While these co-receptor driver lines labeled glomeruli in the same regions from brain to brain, the interindividual expression patterns were not identical, consistent with the variability in glomerular anatomy that we have observed (see Methods). Neurons that detect CO₂ are located in the maxillary palp (Grant et al., 1995; Lu et al., 2007; Omer and Gillies, 1971) and we saw that *Gr3*-expressing neurons projected to a large glomerulus in the posterior antennal lobe, Glomerulus 1 (Figure 1I-K, see Methods for full discussion of glomerulus nomenclature) which is also innervated by *Ir25a*-expressing neurons. We also noted the presence of a second small glomerulus that was often innervated by *Gr3*-expressing neurons in the antenna (Figure S6B). These initial findings point to the overlap of OR-, IR-, and GR-expressing neurons in the antennal lobe of *Ae. aegypti*, which are believed to remain segregated in *Drosophila melanogaster*.

Co-expression and co-convergence of *Orco* and *Ir25a* in the mosquito olfactory system

The high degree of overlap between glomeruli labeled by *Orco*- and *Ir25a*-expressing olfactory sensory neurons suggests that there is either widespread *Orco* and *Ir25a* co-expression within individual sensory neurons or that *Orco* and *Ir25a* are expressed in different neurons whose axons co-converge onto individual antennal lobe glomeruli or both (Figure 1H). To distinguish between these possibilities, we adapted the Split-QF2 system (Riabinina et al., 2019) for use in the mosquito. This system “splits” the transcription factor QF2 into two components, the DNA binding domain (QF2-DBD) and the activation domain (QF2-AD) each tagged with a synthetic leucine zipper (Figure 2A,B). When both the QF2-DBD and QF2-AD are co-expressed in the same cell,

Figure 1: Mismatch in chemosensory receptor and olfactory glomerulus number suggests a novel olfactory organization (A) *Ae. aegypti* female with sensory structures highlighted in yellow boxes. (B) Approximate number of antennal lobe glomeruli per brain hemisphere innervated by the indicated sensory structure, derived from quantification of the left antennal lobe in 12 brains presented in (I-J) and Figure S2-S5. See also Figure S1. (C) Cartoons of insect chemosensory gene families. (D) Cartoon of canonical olfactory system organization. (E-G) Stacked bar plots of the number of chemosensory genes in the *Ae. aegypti* genome (E), and the number expressed above the indicated TPM thresholds in the antenna (F) and maxillary palp (G). (H) Two models of olfactory system organization that can account for the observation that there are far more chemosensory receptors than olfactory glomeruli in *Ae. aegypti*. (I) Maximum-intensity projections of confocal Z-stacks of antennal lobes in the left-brain hemisphere of the indicated genotype with immunofluorescent labeling of GFP (green) and the nc82 monoclonal antibody, which recognizes Brp (magenta). Brp is used throughout this paper as a synaptic marker (Wagh et al., 2006). Scale Bar: 50 μm. Orientation: d=dorsal, m=medial. (J) 2-D representation of the boundary of each glomerulus that is GFP positive and GFP negative. See also Figure S2-S6. (K) Cartoon schematic of the glomeruli receiving projections from olfactory sensory neurons expressing the indicated chemosensory receptor.

the two domains associate via the leucine zipper, reconstitute a functional QF2 protein, initiate transcription at the QUAS enhancer, and drive expression of a reporter gene (Figure 2C).

Using the same stop-codon replacement approach that we used to generate the QF2-lines, we inserted the QF2-AD into the *Ir25a* locus (here referred to as *IR25a-QF2-AD*) and the QF2-DBD into the *Orco* locus (here referred to as *Orco-QF2-DBD*). When either *IR25a-QF2-AD* or *Orco-QF2-DBD* was used to drive expression of *dTomato*, we did not see fluorescence in the antenna, maxillary palp, or the antennal lobe (Figure 2D-F, S7). Therefore, neither QF2-DBD nor QF2-AD alone can activate expression from the QUAS enhancer. However, when *Orco-QF2-DBD* and *IR25a-QF2-AD* were crossed into the same animal, we saw expression of *dTomato* in antennal and maxillary palp neurons, as well as axonal projections in the antennal lobe (Figure 2D-F, S7, S8). Nearly half of the glomeruli in the antennal lobe were labelled with dTomato (Figure 2G-I, S7, S8). This points to widespread *Orco* and *Ir25a* co-expression within *Ae. aegypti* olfactory sensory neurons.

While examining these antennal lobes we consistently noticed that a cluster of anterior dorsal glomeruli was unlabeled when the *Orco-QF2-DBD*, *IR25a-QF2-AD* combination was used to drive expression in animals. However, these glomeruli were labelled when either *Orco-QF2* or *IR25a-QF2* were used to drive expression (Figure 2J,K). This observation strongly suggests that separate populations of *Orco*-positive and *Ir25a*-positive neurons co-converge onto these glomeruli. Therefore, the mosquito olfactory system shows evidence of both co-expression and co-convergence, motifs that are not typically seen in the conventional model organisms *Drosophila melanogaster* flies or *Mus musculus* mice (Figure 2L). It is also possible that a given glomerulus shows both co-convergence of molecularly distinct neurons while also receiving projections from neurons co-expressing multiple chemosensory co-receptors.

Extensive co-expression of chemosensory co-receptors in the antenna

The observation that nearly half of antennal lobe glomeruli receive projections from neurons co-expressing *Ir25a* and *Orco* in gene-targeted strains suggested that there is extensive co-expression of the IR and OR chemoreceptor families. To confirm the existence of such co-expressing neurons, we carried out whole mount antennal immunostaining with an antibody to the endogenous *Orco* protein and GFP expressed from each sensory neuron driver. We observed extensive co-expression of *Orco* and *Ir25a*, with substantially fewer cells co-expressing either *Orco* and *Ir8a* or *Orco* and *Ir76b* (Figure 3A-D), even after accounting for fewer total *Ir76b* and *Ir8a* cells (Figure 3B-D, Data File 1).

To confirm and extend these results, we performed RNA *in situ* hybridization in wild-type antennae with probes designed to target endogenous *Orco*, *Ir76b*, and *Ir25a* transcripts (Figure 3E). These experiments replicated patterns of co-expression observed in immunostained antennae

(Figure 3A-D), with almost half of *Orco* cells co-expressing *Ir25a*, and few *Orco* cells co-expressing *Ir76b* (Figure 3F-H, Data File 1) indicating that widespread co-expression is not an artifact of the genetic reporter lines.

By examining the expression pattern of these three transcripts simultaneously, we were also able to elucidate the relationship between *Ir25a* and *Ir76b* expression. In *Drosophila melanogaster*, *Ir76b* is co-expressed with *Ir25a* in olfactory sensory neurons in the antenna and in gustatory neurons in the labellum (Abuin et al., 2011; Lee et al., 2018). In *Ae. aegypti*, we found that 100% of *Ir76b*-expressing neurons in the antenna also express *Ir25a* (Figure 3F,H). We also observed that in addition to widespread co-receptor co-expression, some mosquito olfactory neurons express just one co-receptor (Figure 3I). Our data demonstrate not only extensive co-expression between *Orco* and IR co-receptors, but also complexity in the rules that govern receptor expression in antennal olfactory sensory neurons.

Coordinated co-expression of chemosensory receptors in the maxillary palp

We have documented extensive chemosensory co-receptor co-expression, but to form functional odorant-sensitive IR or OR complexes, olfactory sensory neurons must express both co-receptors and ligand-sensitive receptors (Abuin et al., 2011; Benton et al., 2009; Larsson et al., 2004; Neuhaus et al., 2005). To simultaneously monitor the extent of co-expression of both co-receptors and ligand-sensitive receptors, we carried out multiplexed whole mount RNA *in situ* hybridization (Choi et al., 2018) in the maxillary palp, the olfactory organ that detects the potent host cue CO₂ as well as other host odors (Grant et al., 1995; Lu et al., 2007; McMeniman et al., 2014; Omer and Gillies, 1971) (Figure 4A,B). The maxillary palp expresses many fewer chemosensory receptors genes than the antenna, with 18 receptors detected at the 1 TPM threshold in the maxillary palp compared to 138 in the antenna at the same threshold, simplifying the task of selecting genes for expression analysis (Figure 1F,G). We performed RNA *in situ* hybridization with probes for 10 of the 18 chemosensory co-receptors and ligand-sensitive receptors that were present in maxillary palp RNA-seq at a threshold of TPM>1. This technique visualized gene expression with sufficient sensitivity that even *Or71* and *Ir75g* (present at 1.93 and 1.67 TPM, respectively) were readily detected (Figure 4C-I, S9).

We found no overlap in expression of *Orco* and *Gr3* in the maxillary palp, but *Ir25a* was expressed in all *Orco* and all *Gr3* cells (Figure 4D). Previous work in *Anopheles gambiae* suggested that *Orco*-expressing neurons in the maxillary palp can be evenly divided into two non-overlapping groups: an *Or8* population and an *Or49* population (Lu et al., 2007). We show definitively that *Or8* and *Or49* are in segregated populations of *Orco*-expressing neurons in *Ae. aegypti* (Figure 4E), and when combined with the results of the previous experiment (Figure 4D), that these cells are also all *Ir25a*-positive. Additional RNA *in situ* hybridization experiments revealed that *Or8*- and *Or49*-expressing cells also often express *Ir76b*, with a bias towards expression in *Or8*-

Figure 2. Genetic evidence for widespread *Orco* and *Ir25a* co-expression

(A) Schematic of the Split-QF2 system (left) and diagrams of *Orco* and *Ir25a* gene loci with exons (grey boxes), introns (grey lines) and CRISPR-Cas9 gRNA site (arrowhead) used to insert *T2A-QF2-AD* (light blue) and *T2A-QF2-DBD* (medium blue). AD and DBD gene maps are not to scale. (B-C) Schematic of the Split-QF2 system (B) and outcome of gene expression in olfactory sensory neurons of the indicated genotypes (C). (D-E) Maximum-intensity projections of confocal Z-stacks of antennae (D) and maxillary palp (E) of the indicated genotypes showing intrinsic dTomato fluorescence, with transmitted light overlay. See also Figure S7A,B. (F-G) Maximum-intensity projections of confocal Z-stacks of antennal lobes from the left brain hemisphere of the indicated genotype with immunofluorescent labeling of dTomato (green) and Brp (synaptic marker, magenta). See also Figure S8. (H-I) 2-D representation of the boundary of each glomerulus in (G) that is GFP positive and GFP negative (H) and quantification (I). $n=3$, mean $\pm$ SEM. See also Figure S7C. (J) Single confocal sections of the anterior dorsal region of the antennal lobe in the left-brain hemisphere of the indicated genotype with immunofluorescent labeling of dTomato (green) and Brp (synaptic marker, magenta). (K) Schematic of the 5 glomeruli shown in (J) with GFP positive glomeruli (green). (L) Two models of *Ae. aegypti* olfactory system organization supported by data in this figure. Scale bars: 50 μ m, except (J): 25 μ m. Orientation: proximal left (D), distal left (E); (F,G,H,J): d=dorsal, m=medial.

Figure 3. Extensive co-expression of chemosensory co-receptors in the antenna

(A) Orco and GFP immunostaining in antennae of the indicated genotypes with cartoon schematic indicating cell identity. (B-D) Quantification of antennal cells in the indicated genotypes co-expressing Orco protein and GFP presented as violin plots showing raw cell counts (B)(mean with range), stacked bar plots (C)(mean $\pm$ SEM), and Euler diagrams with area scaled to mean cells/region (D). $n=6-8$ antennal segments, 34-68 cells/region. (E) RNA *in situ* hybridization in wild-type antennae with the indicated probes. (F-H) Quantification of wild-type antennal cells expressing the indicated genes as violin plots showing raw cell counts (F)(mean with range), stacked bar plots (G)(mean $\pm$ SEM), and Euler diagrams with area scaled to mean cells/region (H). $n=4$ antennal segments, 45-63 cells/region. (I) Cartoon schematic of olfactory neuron populations identified in this figure. Scale bars: 10 μ m.

expressing cells (Figure 4F, S9, Data File 1). Taken together these data show that olfactory sensory neurons express co-receptors *Orco*, *Ir25a*, *Ir76b* and either of the ligand-sensitive subunits *Or49* or *Or8*.

When we analyzed IR ligand-sensitive subunit expression, we found that *Ir100a* and *Ir93a* are also expressed in a subset of *Or49*-expressing neurons (Figure 4G,H, S9, Data File 1), suggesting that these cells can form functional OR and IR complexes with their respective co-receptors in the same neuron. Further, we identified co-expression of *Or71* and *Or49*, demonstrating that more than one ligand-sensitive OR can also be expressed in an olfactory sensory neuron in *Ae. aegypti* (Figure S9O). Lastly, we found that 65% of *Gr3*-expressing cells also express *Ir75g*. Taken together with the finding that *Gr3*-expressing cells express *Ir25a* (Figure 4D, 4I, S9N, Data File 1) this raises the possibility that a functional IR complex can be found in CO₂-sensitive neurons.

Volatile amine sensing by CO₂ neurons that lack the *Gr3* CO₂ receptor

We set out to investigate the functional relationships of these extensive receptor co-expression patterns by focusing on the maxillary palp, which is both simpler in organization than the antenna and also uniquely important because it is the sensory appendage that detects CO₂.

To delineate the organization of maxillary palp projections in the brain, we used our QF2 and Split-QF2 driver lines to examine sensory innervation of antennal lobe glomeruli (Figure 5A-H). Glomerulus 2 and Glomerulus 3 received input from both *Orco*-expressing neurons and *Ir25a*- and *Ir76b*-expressing neurons (Figure 5A-G), consistent with data presented in Figure 4. Co-expression of *Orco* and *Ir25a* in neurons that project to these two glomeruli was confirmed using the Split-QF2 system (Figure 2). In *Orco*-QF2-DBD, *Ir25a*-QF2-AD animals, Glomerulus 2 and Glomerulus 3 were labeled, but Glomerulus 1 was not (Figure 5H). We also discovered that Glomerulus 1, which is the largest glomerulus in the antennal lobe (Shankar and McMeniman, 2020), received input from *Gr3*-expressing sensory afferents. Glomerulus 1 was also innervated by *Ir25a*-expressing sensory neurons (Figure 5A-H), consistent with the co-expression of *Gr3* and *Ir25a* in olfactory neurons (Figure 4D). This finding raised the possibility that Glomerulus 1 may respond to additional odorant ligands independent of CO₂-sensitive *Gr3*.

Given the extensive receptor co-expression observed in the maxillary palp (Figure 4,5I), we next explored the possibility that functional expression of multiple receptor types would allow maxillary palp CO₂-sensitive neurons to respond to additional odorants. We set out to examine odorants detected by maxillary palp neurons by developing an *in vivo* calcium imaging preparation with the genetically-encoded calcium sensor GCaMP6s (Chen et al., 2013), allowing us to monitor stimulus-evoked activity in the antennal lobe of a mosquito as it was exposed to different odorants or to CO₂ (Figure 5J). We imaged the axon terminals of maxillary palp sensory neurons in the antennal lobe with two-photon microscopy

using a custom-built olfactometer to deliver odorants and CO₂ within a continuous air stream. We expected that Glomerulus 1 would respond to CO₂ because it receives projections from *Gr3*-expressing neurons. Indeed, in preliminary experiments when we imaged GCaMP6s responses in axonal termini of *Gr3*-expressing neurons in Glomerulus 1, we observed low fluorescence at baseline and concentration-dependent responses to CO₂, with no apparent desensitization across trials (data not shown).

To image activity simultaneously in CO₂-sensitive Glomerulus 1 and the non-CO₂-sensitive Glomerulus 3, we used the *Ir25a*>GCaMP6s strain, which expresses GCaMP6s in all maxillary palp olfactory neurons (Figure 4). We saw consistent activation of Glomerulus 3 by 1-octen-3-ol and 3-octanone (Figure 5K,L), odorants that have been shown in *in vitro* heterologous expression experiments to activate Or8-Orco from both *Ae. aegypti* and *Anopheles gambiae* (Bohbot and Dickens, 2009; Lu et al., 2007). Therefore Glomerulus 3 is likely to be innervated by *Or8*-expressing neurons (Figure 4E). From a panel of candidate odorants, we found that Glomerulus 3 also responded to two volatile amines, hexyl amine and triethyl amine (Figure 5K,L). Volatile amines, including polyamines, have been proposed to be IR ligands in *Drosophila melanogaster* (Geier et al., 1999; Hussain et al., 2016; Min et al., 2013; Silbering et al., 2011). These findings are consistent with the hypothesis that IRs and ORs are functionally co-expressed in maxillary palp neurons and that this co-expression enables these cells to respond to ligands that activate both classes of receptors.

We next turned to the CO₂-sensitive Glomerulus 1 (Figure 5M) and imaged stimulus-evoked activity in *Gr3* heterozygotes, which have no defects in CO₂ sensation, and in *Gr3* homozygous mutants, which lose all physiological and behavioral responses to CO₂ (McMeniman et al., 2014). We used *Gr3* heterozygotes as the control genotype in these experiments because this mutation was introduced in a different wild-type strain than our QF2/QUAS reagents and this genetic configuration better controls for genetic background (see Methods). As expected, we saw robust activation of Glomerulus 1 by CO₂ in *Gr3* heterozygotes, and a complete loss of response to CO₂ in *Gr3* homozygous mutant animals (Figure 5N-P). CO₂-sensing neurons in the maxillary palp respond to multiple odorants in *Aedes*, *Culex*, and *Anopheles* mosquitoes (Lu et al., 2007; Tauxe et al., 2013; Turner et al., 2011) and it has been proposed that *Gr3* is a broadly-tuned receptor that responds to many odorants. We examined the response to a recently identified CO₂-neuron activator, acetone (Ghaninia et al., 2019). This odorant activated Glomerulus 1 in the *Gr3* heterozygote but the response to acetone was abolished in the homozygous *Gr3* mutant (Figure 5Q,R). This suggests that the CO₂ receptor can interact with non-CO₂ ligands. We examined the response to volatile amines, and saw consistent responses to hexyl amine and triethyl amine (Figure 5Q,R). The response to both of these volatile amines was unaffected in *Gr3* homozygous mutant animals (Figure 5Q,R). We speculate that hexyl amine and triethyl amine are activating the CO₂-sensitive neuron by stimulating IRs expressed in this neuron,

Figure 4. Coordinated co-expression of chemosensory receptors in the maxillary palp

(A) Schematic of human chemosensory cues: CO₂ detection requires Gr3. **(B)** Cartoon indicating maxillary palp (green). **(C)** Maxillary palp expression of *Orco* revealed by whole-mount RNA *in situ* hybridization. Orientation: fourth segment, distal right. **(D-I)** Whole-mount maxillary palp RNA *in situ* hybridization with the indicated probes, cartoon schematic indicating cell identity, and quantification of co-expression shown as Euler diagrams, area scaled to mean. n=5 maxillary palps, 26-65 cells/dorso-lateral maxillary palp. See also Figure S9. Scale bars: 50 μm (C), 25 μm (D-I).

Figure 5: Volatile amine sensing by CO₂ neurons that lack the Gr3 CO₂ receptor

(A-B) Cartoon (A) and left 3-D antennal lobe reconstruction (B) showing 3 glomeruli that are innervated by the maxillary palp. Image in (B) is reprinted in Figure S1G. (C-H) Single confocal sections through the center of Glomerulus 1 (top) or Glomerulus 2 and Glomerulus 3 (bottom) in left antennal lobes of the indicated genotypes. Sections are taken from Z-stacks presented in Figure 1C (C-G) and Figure 2G (H). (I) Schematic of sensory neuron gene expression and glomerular convergence based on (C-H). (J) Side view schematic of live-imaging preparation of Glomerulus 3. (K) GCaMP6s fluorescence traces from Glomerulus 3 presented with the indicated stimuli. Black bar indicates stimulus presentation, mean ± SEM, n=9-10 sweeps from 3 animals/stimulus. (L) Summary of data in (K) indicating the maximum ΔF/F (arbitrary units). Lines denote median and quartiles, *p<0.05 one-way ANOVA with Bonferroni correction compared to water control. (M) Side view schematic of live-imaging preparation of Glomerulus 1. (N) Two-photon image of Glomerulus 1 in the left antennal lobe of *Ir25a>dTomato-T2A-GCaMP6s* animals with dTomato (left), GCaMP6s fluorescence at baseline (middle), and after CO₂ presentation (right) with region of interest (yellow), in the indicated *Gr3* genotypes. (O) GCaMP6s traces from Glomerulus 1 from animals of the indicated genotype presented with CO₂ at the time indicated by the black bar, mean ± SEM, n=9-10 sweeps from 3 animals. (P) Summary of data in (O) indicating the maximum ΔF/F (arbitrary units). Lines denote median and quartiles, *p<0.05 one-way ANOVA with Bonferroni correction. (Q) GCaMP6s fluorescence traces from Glomerulus 1 presented with the indicated stimuli. Black bar indicates stimulus presentation, mean ± SEM, n=5-10 sweeps from 3 animals/stimulus. (R) Summary of data in (Q) indicating the maximum ΔF/F (arbitrary units). Lines denote median and quartiles. *p<0.05 one-way ANOVA with Bonferroni correction. (S) Summary of Glomerulus 1 responses with and without *Gr3* function. CO₂ concentration in all panels is 2%; see Methods for details of odorant concentrations. Scale bars: 25 μm.

which we have shown include at least *Ir25a* and *Ir75g* (Figure 4I, 5S).

Volatile amines substitute for CO₂ in activating mosquito attraction to humans

Female mosquitoes integrate multiple sensory cues to drive strong attraction to humans. Human odor is a blend of hundreds of volatile odorants, including lactic acid and volatile amines (Acree et al., 1968; Bernier et al., 2000; Dormont et al., 2013). While body heat, CO₂ emitted in the breath, and individual odorants produced by human skin individually elicit weak to no mosquito attraction, the combination of multiple sensory cues leads to robust attraction in laboratory behavioral assays (Corfas and Vosshall, 2015; Dekker et al., 2005; McMeniman et al., 2014). Among these cues, CO₂ is the most potent synergistic stimulus, sensitizing mosquitoes to gate attraction to both heat and odorants (Dekker et al., 2005; McMeniman et al., 2014). The discovery that triethyl amine and hexyl amine activate Glomerulus 1 in *Gr3* homozygous mutants that are unable to detect CO₂ led us to hypothesize that volatile amines could substitute for CO₂ in sensitizing *Gr3* mutant mosquitoes to human body odor (Figure 6A).

To test this, we used a behavioral assay (Basrur et al., 2020) to measure mosquito attraction to various stimuli (Figure 6B). In control experiments, we baited the assay with either CO₂ or lactic acid, a human host volatile, or both CO₂ and lactic acid combined, and monitored how many mosquitoes accumulated in the trap. As previously shown, either CO₂ or lactic acid alone only weakly attracted mosquitoes, but the combination of lactic acid with CO₂ caused potent dose-dependent attraction in two different wild-type strains of *Ae. aegypti* (Figure 6C, Data File 1) (Acree et al., 1968; Bernier et al., 2000; Davis, 1984; Geier et al., 1999; Majeed et al., 2014; McMeniman et al., 2014; Smith et al., 1970). Female mosquito attraction to humans is suppressed for 72-96 hours after ingestion of a blood-meal (Davis, 1984; Duvall et al., 2019; Klowden, 1981). We saw the same state-dependent suppression of attraction to CO₂ and lactic acid by blood-fed mosquitoes, confirming that these stimuli serve as a proxy for the attraction to human host odor (Figure 6C).

We next tested the hypothesis that activating Glomerulus 1 with volatile amines in CO₂-insensitive *Gr3* mutants could sensitize these animals to respond to lactic acid. In control experiments, we observed that neither wild-type nor *Gr3* mutants were attracted to CO₂ alone, and that only wild-type animals and not *Gr3* mutants showed synergistic attraction to CO₂ and lactic acid (Figure 6D). As previously observed, both wild-type and *Gr3* mutants showed strong attraction to a live human host in the presence of CO₂ (McMeniman et al., 2014). Live human hosts emit many sensory cues, effectively compensating for the inability of *Gr3* mutant mosquitoes to detect CO₂ (McMeniman et al., 2014).

Both CO₂ and triethyl amine activate Glomerulus 1 in wild-type animals but in *Gr3* mutants only triethyl amine and not CO₂ activates this glomerulus. We asked if triethyl amine alone or in combination with CO₂ was attractive to wild type

or *Gr3* mutant mosquitoes. Neither wild type nor *Gr3* mutant mosquitoes showed any attraction to CO₂ alone, triethyl amine alone, or the combination of both (Figure 6E). This was expected because CO₂ alone—and by extension the activation of Glomerulus 1—is insufficient to mediate attraction in this assay. We next asked if triethyl amine can substitute for CO₂ in synergizing with lactic acid to attract *Gr3* mutant mosquitoes. We tested a range of concentrations of triethyl amine with a single concentration of lactic acid and quantified the number of attracted mosquitoes. Lactic acid alone and both low and high concentrations of triethyl amine were not attractive to *Gr3* mutants. However, moderate concentrations of triethyl amine stimulated strong attraction when supplied together with lactic acid (Figure 6F). This sensitization to lactic acid triggered by triethyl amine was not further enhanced by the addition of CO₂, a stimulus that *Gr3* mutants do not detect (Figure 6G). We speculate that CO₂-insensitive *Gr3* mutants rely on a compensatory pathway that uses *Ir25a* and unknown ligand-sensitive IRs to activate Glomerulus 1 with volatile amines.

To ask if this compensatory pathway is active in wild-type mosquitoes, we repeated these experiments and were surprised to discover that the concentrations of triethyl amine that stimulated *Gr3* mutant attraction to lactic acid had no effect on wild-type animals (Figure 6H,I). These results suggest that when mosquitoes lack receptors for potent host cues such as CO₂, they rely more heavily on redundant cues detected by the same neurons via the expression of a diverse array of chemosensory receptors to ensure that host-seeking behavior is not compromised (Figure 6J).

DISCUSSION

A compensatory pathway promotes attraction behavior in the CO₂ receptor mutant

It has only recently become possible to generate targeted mutations in chemosensory genes in *Ae. aegypti*, and a striking finding from studies that generated such mutants is how difficult it is to disrupt human host-seeking. Mutations in the co-receptor *Orco* (DeGennaro et al., 2013) and the co-receptor *Ir8a* (Raji et al., 2019) both eliminate entire classes of ligand-specific receptors. In spite of this, *Orco* mutants show normal attraction to humans and mutations in *Ir8a* reduce but do not abolish mosquito attraction to humans. *Gr3* mutants are completely unable to sense CO₂ yet they can also find human hosts, even when released into a large semi-field environment (McMeniman et al., 2014). These findings raise the question: Why is it so difficult to break the mosquito olfactory system and prevent mosquitoes from finding humans? We propose that this is due to redundancy in receptor expression at the periphery. A recent paper (Zhao et al., 2020) examined *Ae. aegypti* antennal lobe responses of *Orco*-expressing neurons to physiological concentrations of odor collected from several individual humans, a variety of non-human animals, and floral odors. They identified a glomerulus that responded to odor blends from all human individuals, animals, as well as honey and milkweed odor. It is tempting to speculate that this

Figure 6: Volatile amines substitute for CO₂ in activating mosquito attraction to humans

(A) Humans emit a complex blend of volatile odorants, including CO₂, lactic acid, and volatile amines. (B-C) Schematic of behavioral assay (B) and percent wild-type mosquitoes of the indicated blood-feeding status attracted to the indicated stimuli in the assay (C). Lactic acid was supplied at 10 μ L. Mean with range, n=11-15 trials, 19-21 mosquitoes/trial, *p<0.05, two-way ANOVA followed by Tukey's HSD. (D-I) Attraction of mosquitoes of the indicated genotype to the indicated stimuli. Lactic acid was supplied at 1 μ L in D,F-I. Triethyl amine was presented at the indicated concentration in F, H and at 50 μ L in E,G,I. Mean with range, n=7-16 trials, 15-22 mosquitoes/trial. *p<0.05, ns=not significant (*p>0.05), two-way ANOVA followed by Tukey's HSD. (J) Model of the compensatory pathway where volatile amines substitute for CO₂ to enable attraction to humans by CO₂-insensitive Gr3 mutants.

glomerulus is innervated by olfactory sensory neurons that co-express multiple receptors. We have probed IR and GR redundancy in the CO₂ system and found that *Gr3* mutant mosquitoes that do not detect CO₂ can still detect triethyl amine, and have demonstrated that triethyl amine can act as a behavioral substitute for CO₂ in driving attraction to lactic acid in *Gr3* mutants. This activation was strongly dependent on the dose of triethyl amine. Although human skin releases a large number of odorants, including volatile amines, we do not know how the concentrations used in our laboratory behavioral assays relate to those produced by human skin.

Surprisingly, this substitution of triethyl amine for CO₂ does not drive attraction in wild-type animals. One possible mechanism to explain this finding might be that in the absence of *Gr3*, the sensitivity of the circuit shifts to enhance the behavioral effect of compounds that activate the IRs co-expressed in the CO₂-sensitive neuron. If this were the case, it remains to be seen how the CO₂-sensing circuitry is altered, whether it is the sensory neuron or downstream neurons involved in host-seeking, and if changes occur at the level of the receptor, synapse, or the intrinsic firing properties of the sensory neurons themselves (Marder and Goaillard, 2006). CO₂-sensing neurons in the maxillary palp rapidly reset their intrinsic firing to adjust to changes in background levels of CO₂ (Majeed et al., 2014) and perhaps the absence of background CO₂ sensation in the *Gr3* mutant enhances the signal to noise ratio of other odorants.

Coordinated co-expression between IR, OR, and GR ligand-sensitive receptors

We identified co-expression of co-receptors and ligand-sensitive receptors belonging to distinct chemosensory families in single neurons in the maxillary palp. This co-expression poses a gene regulatory problem for an olfactory neuron. For ORs and IRs to form functional chemoreceptors, at least one co-receptor and one ligand-sensitive receptor must be expressed in a cell. We have demonstrated that two IRs, *Ir93a* and *Ir100a*, are expressed nearly exclusively with *Or49* and the co-receptors *Orco* and *Ir25a*, whereas *Ir75g* is co-expressed with the coreceptor *Ir25a* and the CO₂ receptor subunit *Gr3*. Thus, the transcriptional landscape in *Ae. aegypti* maxillary palp olfactory neurons is not only permissive to co-expression, but ensures certain receptors are only expressed with others.

How might this complex code of chemosensory receptor co-expression be regulated? In vertebrates, an elaborate epigenetic silencing mechanism ensures that each olfactory neuron expresses only a single allele of a single odorant receptor (Bashkurova and Lomvardas, 2019). In contrast, *Drosophila* is thought to use a more conventional transcription factor code in which the specification of a neuron and the expression of its chemosensory receptor is tightly regulated (Jafari and Alenius, 2015; Li et al., 2016; Ray et al., 2008). Single-cell sequencing data generated from developing *Drosophila melanogaster* olfactory neurons demonstrates a complex regulatory landscape wherein dependent on cell type, a set of transcription factors govern receptor

expression, axon targeting, or both (Li et al., 2020). Two recent studies (McLaughlin et al., 2020; Task et al., 2020) also document extensive co-expression of receptors in *Drosophila melanogaster*, calling into question the rules that regulate olfactory organization in this insect. It is yet to be determined if mosquito orthologues of these transcription factors have been co-opted to regulate the co-expression we observe or if this novel olfactory organization demands a distinct transcriptional mechanism.

The maxillary palp has been implicated in responses to CO₂ (Acree et al., 1968; Gillies, 1980; Grant et al., 1995), thermosensation (Roth, 1951), mechanosensation (Bohbot et al., 2014), detection of attractive monomolecular odorants such as 1-octen-3-ol (Syed and Leal, 2007; Takken and Kline, 1989; Vythilingam et al., 1992), as well as blends of odorants extracted from human hosts (Tauxe et al., 2013). Maxillary palp transcriptomes offer insights into how these tiny sensory organs detect a broad range of stimuli (Bohbot et al., 2014; Matthews et al., 2018), pointing to a transcriptional landscape that is difficult to reconcile with the prior understanding of neuronal identity in the maxillary palp. It was thought that all basiconic sensilla, which detect volatile odorants, house the sensory dendrites of three neurons that form identical repeating units: one large CO₂-sensitive neuron that expresses *Gr3*, and two smaller neurons that express either an *Orco-Or8* receptor or an *Orco-Or49* receptor (Lu et al., 2007; Mclver, 1972). How do these repeating units of just three neurons enable the detection of so many stimuli and accommodate the expression of so many chemosensory genes? We demonstrate through multiplexed RNA *in situ* hybridization experiments that the receptor composition of these neurons is far more complex, and they can be subdivided into many more than three cell types. We did not evaluate the relationship of every combination of chemosensory genes expressed in the maxillary palp and hypothesize that comprehensive gene expression profiling including the application of single cell RNA-sequencing will reveal even more cell types. The extensive co-expression of IRs, ORs, and GRs documented in this work likely enables maxillary palp neurons to detect volatile odorants belonging to multiple chemical classes and may explain how CO₂-sensitive neurons also respond to human skin odor (Dormont et al., 2013; Tauxe et al., 2013), and why mosquitoes unable to detect CO₂ can still locate a human host (McMeniman et al., 2014).

Receptor co-expression as a mechanism for redundancy or blend integration

We hypothesize that receptor co-expression is used broadly to detect redundant cues that are present in human odor, a blend that can vary from individual to individual and contains hundreds of different chemicals (Bernier et al., 1999; Bernier et al., 2000). We have not yet explored the role of OR-IR co-expression or the expression of multiple IRs or ORs in single olfactory neurons. It is possible that receptor co-expression is always used to form a highly redundant detection system for different cues that represent the same ecological target: human hosts. This motif has the benefit of limiting the number of neurons needed to detect varied odorants with the same meaning, however, in exchange it may sacrifice the

ability to distinguish between cues detected by receptors expressed in the same sensory neurons. In *Drosophila melanogaster* there are rare cases of OR-OR or OR-IR co-expression, which is the exception rather than the rule. *Or35a* is co-expressed with *Ir76b* (Silbering et al., 2011), and while these neurons respond to many odorants (Silbering et al., 2011; Yao et al., 2005), the role of co-expression remains unknown. In *Drosophila melanogaster*, *Or49a* and *Or85f* are co-expressed in a specific olfactory sensory neuron population where they play redundant roles in predator avoidance (Ebrahim et al., 2015). Intriguingly, recent data from *Drosophila melanogaster* also show *Orco* and *Ir25a* co-expression in the antenna and maxillary palp (McLaughlin et al., 2020; Task et al., 2020), suggesting that broad co-receptor co-expression may be a conserved motif in insects and upending previous data that suggested strict segregation of OR and IR sensory neuron populations in flies.

While extensive co-expression of different classes of olfactory receptors within olfactory neurons documented in this study is a novel organizational principle, there are counterexamples of receptor co-expression within a single class of chemosensory receptor. Karner et al. have documented polycistronic expression of multiple odorant receptors in *Anopheles gambiae* sensory neurons (Karner et al., 2015). The nematode *C. elegans* copes with a very large number of chemosensory receptor genes and a very small number of sensory neurons by extensive receptor co-expression (Troemel et al., 1995; Vidal et al., 2018).

What is the functional significance of co-expression of different classes of olfactory receptors, the ORs, IRs, and GRs, in *Aedes aegypti*? It is possible that co-expression is exploited to integrate the detection of multiple odors in the sensory neuron itself. In *Drosophila melanogaster*, IRs that form functional complexes can depolarize resting membrane potential *in vitro* (Ai et al., 2013). If the same phenomenon is occurring in *Ae. aegypti* neurons that express both IR and OR complexes, it is possible that an IR complex could render a neuron more sensitive to OR ligands. Certain odors repress OR activity relative to baseline in *Drosophila melanogaster* (Hallem and Carlson, 2006; Yao et al., 2005) and it is possible that multiple receptors housed in a single neuron could be a source of blend-specific repression. Individual odorant molecules found in human odor exist in many blends and this might be a way to limit the attraction to monomolecular components of human odor that are found in aversive blends or are produced by other animals. Recent data in *Drosophila* show that, in maxillary palp sensilla innervated by *Orco* neurons, responses to several odorants change in *Ir25a* mutant flies (Task et al., 2020). The direction of the response is sensilla-specific. In some neurons, responses are higher in *Ir25a* mutants, in other neurons, responses are lower, suggesting that *Ir25a* may be implicated in gain control in *Orco* neurons (Task et al., 2020). These ideas remain to be tested in the mosquito, but it is tempting to speculate that this could be used as a method to detect ratios of odors present in a blend, a key determinant in how attractive many monomolecular compounds are to mosquitoes (Bernier et al., 2007).

We demonstrate that *Ae. aegypti* mosquitoes employ both receptor co-expression in primary sensory neurons as well as co-convergence of neurons that express distinct classes of chemosensory receptors onto the same glomerulus. Consistent with these data, we also identified olfactory neurons in the antenna that express ORs without IRs, IRs without ORs, and neurons that express both ORs and IRs. What is the functional difference between integrating odorant information in the primary sensory neuron versus at the first synapse in the antennal lobe? Co-convergence of olfactory sensory neurons onto the antennal lobe could allow for the early integration of olfactory cues while still retaining discrete input channels that could be selectively modulated during changes in behavioral state, such as the suppression of host-seeking after a blood meal. The role of co-convergence in this system remains to be tested.

In summary, our study reveals unexpected complexity in the gene expression and functional organization of the mosquito olfactory system. The potential redundancy afforded by a system in which many neurons express receptors that respond to non-overlapping sets of odorant ligands may greatly increase the sensitivity and robustness of the mosquito olfactory system. Future attempts to refine the design of repellents to ward off mosquitoes or attractant traps to lure them will have to reckon with the complexity of this system.

DATA AND RESOURCE AVAILABILITY

Supplementary Figures S1-S9 accompany the paper. All raw data in the paper are provided in Data File 1. Additional raw data and custom scripts are available at https://github.com/VosshallLab/Younger_Herre_Vosshall2020. Plasmids are available from Addgene (accession #162520-162526).

ACKNOWLEDGMENTS

We thank Emily Dennis, Laura Duvall, Itzel Ishida, Philip Kidd, Erica Korb, Carolyn McBride, Christopher Potter, Darya Task, Zhilei Zhao, and members of the Vosshall Lab for comments on the manuscript; Gloria Gordon and Libby Mejia for expert mosquito rearing; Maria Elena (Ellen) de Obaldia for early assistance with the Quatroport olfactometer; Javier Marquina-Solis for antennal lobe tracing assistance; Christina Pyrgaki, Carlos Rico, Katarzyna (Kate) Cialowicz, and Alison North at the Rockefeller Bio-Imaging Resource Center for assistance with confocal imaging; Daniel Gross, James Petrillo, and Peer Strogies at the Rockefeller Precision Instrumental Technologies (PIT) resource center for advice and fabrication of the olfactometer and other imaging components used in the GCaMP imaging experiments; Olena Riabinina and Christopher Potter for providing advice and unpublished Q-system reagents; Carolyn McBride, Matthew DeGennaro and members of the *Aedes* Toolkit Group for advice and discussion; Caroline Jiang for advice on statistical analysis; Nipun Basrur, Priya Rajasethupathy, Andrea Terceros, Harry Choi and Molecular Instruments for advice on RNA *in situ* hybridization experiments; Rob A Harrell II at the Insect Transgenesis Facility at the University of Maryland for embryo injections; Cory Root, Raphael Cohn, Ari Zolin, Gaby Maimon, and Vanessa Ruta for advice on calcium imaging and data analysis; Alan Grant and Greg Suh for advice on CO₂ delivery; Frances Weis-Garcia and the members of the MSKCC Antibody and Bioresource Core Facility for preparation of the nc82/Brp monoclonal antibody.

FUNDING

This work was supported in part by grant # UL1 TR000043 from the National Center for Advancing Translational Sciences (NCATS, National Institutes of Health (NIH) Clinical and Translational Science Award (CTSA) program. Funding for this study was provided by Jane Coffin Childs Postdoctoral Fellowships (M.A.Y. and B.J.M.), The Grass Foundation (M.A.Y.), a Leon Levy Neuroscience Fellowship (M.A.Y.), and by a pilot grant and postdoctoral fellowship from the Kavli Neural Systems Institute (M.A.Y.), a predoctoral fellowship from the Kavli Neural Systems Institute (M.H.), NIH NIDCD grant F30DC017658 (M.H.). B.J.M. received support from National Sciences and Engineering Research Council (NSERC), award RGPIN-2020-05423. M.H. is supported by a Medical Scientist Training Program grant from the National Institute of General Medical Sciences of the NIH under award number T32GM007739 to the Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program. Antibody purification carried out at the MSKCC Antibody and Bioresource Core Facility was supported by a Cancer Center Core Grant 5 P30 CA008748-54. L.B.V. is an investigator of the Howard Hughes Medical Institute.

AUTHOR CONTRIBUTIONS

M.A.Y. carried out all antennal lobe immunostaining and conducted and analyzed all calcium imaging experiments. M.H. carried out all peripheral tissue immunostaining and RNA *in situ* hybridization experiments. M.A.Y. and M.H. together conducted and analyzed all behavior data, except those in Figure 6C, which were carried out by A.R.E. B.J.M. provided gene and transcript analysis in Figure 1. Z.G. cloned and isolated Split-QF2 lines with M.H. Z.N.G. cloned and isolated QF2 stop codon replacement lines with B.J.M. and M.A.Y. S.R. worked with M.H. to generate RNA *in situ* hybridization data. M.A.Y., M.H., and L.B.V. together conceived the study, designed the figures, and wrote the paper with input from all authors.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Abuin, L., Bargeton, B., Ulbrich, M.H., Isacoff, E.Y., Kellenberger, S., and Benton, R. (2011). Functional architecture of olfactory ionotropic glutamate receptors. *Neuron* 69, 44-60.

Acree, F., Jr., Turner, R.B., Gouck, H.K., Beroza, M., and Smith, N. (1968). L-Lactic acid: a mosquito attractant isolated from humans. *Science* 161, 1346-1347.

Ai, M., Blais, S., Park, J.Y., Min, S., Neubert, T.A., and Suh, G.S. (2013). Ionotropic glutamate receptors IR64a and IR8a form a functional odorant receptor complex in vivo in *Drosophila*. *J Neurosci* 33, 10741-10749.

Allan, S.A., Day, J.F., and Edman, J.D. (1987). Visual ecology of biting flies. *Annu Rev Entomol* 32, 297-316.

Bashkurova, E., and Lomvardas, S. (2019). Olfactory receptor genes make the case for inter-chromosomal interactions. *Curr Opin Genet Dev* 55, 106-113.

Basrur, N.S., De Obaldia, M.E., Morita, T., Herre, M., von Heynitz, R.K., Tsitohay, Y.N., and Vosshall, L.B. (2020). *fruitless* mutant male mosquitoes gain attraction to human odor. *bioRxiv* 10.1101/2020.09.04.282434.

Benton, R., Sachse, S., Michnick, S.W., and Vosshall, L.B. (2006). Atypical membrane topology and heteromeric function of *Drosophila* odorant receptors in vivo. *PLoS Biol* 4, e20.

Benton, R., Vannice, K.S., Gomez-Diaz, C., and Vosshall, L.B. (2009). Variant ionotropic glutamate receptors as chemosensory receptors in *Drosophila*. *Cell* 136, 149-162.

Bernier, U.R., Booth, M.M., and Yost, R.A. (1999). Analysis of human skin emanations by gas chromatography/mass spectrometry. 1. Thermal desorption of attractants for the yellow fever mosquito (*Aedes aegypti*) from handled glass beads. *Anal Chem* 71, 1-7.

Bernier, U.R., Kline, D.L., Allan, S.A., and Barnard, D.R. (2007). Laboratory comparison of *Aedes aegypti* attraction to human odors and to synthetic human odor compounds and blends. *J Am Mosq Control Assoc* 23, 288-293.

Bernier, U.R., Kline, D.L., Barnard, D.R., Schreck, C.E., and Yost, R.A. (2000). Analysis of human skin emanations by gas chromatography/mass spectrometry. 2. Identification of volatile compounds that are candidate attractants for the yellow fever mosquito (*Aedes aegypti*). *Anal Chem* 72, 747-756.

Bohbot, J., Pitts, R.J., Kwon, H.W., Rutzler, M., Robertson, H.M., and Zwiebel, L.J. (2007). Molecular characterization of the *Aedes aegypti* odorant receptor gene family. *Insect Mol Biol* 16, 525-537.

Bohbot, J.D., and Dickens, J.C. (2009). Characterization of an enantioselective odorant receptor in the yellow fever mosquito *Aedes aegypti*. *PLoS One* 4, e7032.

Bohbot, J.D., Sparks, J.T., and Dickens, J.C. (2014). The maxillary palp of *Aedes aegypti*, a model of multisensory integration. *Insect Biochem Mol Biol* 48, 29-39.

Brand, A.H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* 118, 401-415.

Brown, J.E., Evans, B.R., Zheng, W., Obas, V., Barrera-Martinez, L., Egizi, A., Zhao, H., Caccone, A., and Powell, J.R. (2014). Human impacts have shaped historical and recent evolution in *Aedes aegypti*, the dengue and yellow fever mosquito. *Evolution* 68, 514-525.

Buck, L., and Axel, R. (1991). A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* 65, 175-187.

Butterwick, J.A., Del Marmol, J., Kim, K.H., Kahlson, M.A., Rogow, J.A., Walz, T., and Ruta, V. (2018). Cryo-EM structure of the insect olfactory receptor Orco. *Nature* 560, 447-452.

Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohian, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., et al. (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* 499, 295-300.

Chess, A., Simon, I., Cedar, H., and Axel, R. (1994). Allelic inactivation regulates olfactory receptor gene expression. *Cell* 78, 823-834.

- Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., and Pierce, N.A. (2018). Third-generation *in situ* hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development* **145**, 10.1242/dev.165753.
- Clyne, P.J., Warr, C.G., and Carlson, J.R. (2000). Candidate taste receptors in *Drosophila*. *Science* **287**, 1830-1834.
- Clyne, P.J., Warr, C.G., Freeman, M.R., Lessing, D., Kim, J., and Carlson, J.R. (1999). A novel family of divergent seven-transmembrane proteins: candidate odorant receptors in *Drosophila*. *Neuron* **22**, 327-338.
- Cohn, R., Morante, I., and Ruta, V. (2015). Coordinated and compartmentalized neuromodulation shapes sensory processing in *Drosophila*. *Cell* **163**, 1742-1755.
- Corfas, R.A., and Vosshall, L.B. (2015). The cation channel TRPA1 tunes mosquito thermotaxis to host temperatures. *Elife* **4**, 10.7554/eLife.11750.
- Couto, A., Alenius, M., and Dickson, B.J. (2005). Molecular, anatomical, and functional organization of the *Drosophila* olfactory system. *Curr Biol* **15**, 1535-1547.
- Davis, E.E. (1984). Regulation of sensitivity in the peripheral chemoreceptor systems for host-seeking behavior by a haemolymph-borne factor in *Aedes aegypti*. *J Insect Physiol* **30**, 179-183.
- DeGennaro, M., McBride, C.S., Seeholzer, L., Nakagawa, T., Dennis, E.J., Goldman, C., Jasinskiene, N., James, A.A., and Vosshall, L.B. (2013). *orco* mutant mosquitoes lose strong preference for humans and are not repelled by volatile DEET. *Nature* **498**, 487-491.
- Dekker, T., Geier, M., and Carde, R.T. (2005). Carbon dioxide instantly sensitizes female yellow fever mosquitoes to human skin odours. *J Exp Biol* **208**, 2963-2972.
- Distler, P., and Boeckh, J. (1997). Central projections of the maxillary and antennal nerves in the mosquito *Aedes aegypti*. *J Exp Biol* **200**, 1873-1879.
- Dormont, L., Bessiere, J.M., McKey, D., and Cohuet, A. (2013). New methods for field collection of human skin volatiles and perspectives for their application in the chemical ecology of human-pathogen-vector interactions. *J Exp Biol* **216**, 2783-2788.
- Duvall, L.B., Ramos-Espiritu, L., Barsoum, K.E., Glickman, J.F., and Vosshall, L.B. (2019). Small-molecule agonists of *Ae. aegypti* neuropeptide Y receptor block mosquito biting. *Cell* **176**, 687-701.
- Ebrahim, S.A., Dweck, H.K., Stokl, J., Hofferberth, J.E., Trona, F., Weniger, K., Rybak, J., Seki, Y., Stensmyr, M.C., Sachse, S., et al. (2015). *Drosophila* avoids parasitoids by sensing their semiochemicals via a dedicated olfactory circuit. *PLoS Biol* **13**, e1002318.
- Fişek, M., and Wilson, R.I. (2014). Stereotyped connectivity and computations in higher-order olfactory neurons. *Nat Neurosci* **17**, 280-288.
- Fishilevich, E., and Vosshall, L.B. (2005). Genetic and functional subdivision of the *Drosophila* antennal lobe. *Curr Biol* **15**, 1548-1553.
- Flanagan, D., and Mercer, A.R. (1989). An atlas and 3-D reconstruction of the antennal lobes in the worker honey bee, *Apis mellifera* L. (Hymenoptera : Apidae). *Int J Insect Morphol & Embryol* **18**, 145-159.
- Gallagher, M., Wysocki, C.J., Leyden, J.J., Spielman, A.I., Sun, X., and Preti, G. (2008). Analyses of volatile organic compounds from human skin. *Br J Dermatol* **159**, 780-791.
- Gao, Q., and Chess, A. (1999). Identification of candidate *Drosophila* olfactory receptors from genomic DNA sequence. *Genomics* **60**, 31-39.
- Geier, M., Bosch, O.J., and Boeckh, J. (1999). Ammonia as an attractive component of host odour for the yellow fever mosquito, *Aedes aegypti*. *Chem Senses* **24**, 647-653.
- Ghaninia, M., Majeed, S., Dekker, T., Hill, S.R., and Ignell, R. (2019). Hold your breath - Differential behavioral and sensory acuity of mosquitoes to acetone and carbon dioxide. *PLoS One* **14**, e0226815.
- Gillies, M.T. (1980). The role of carbon dioxide in host-finding in mosquitoes (Diptera:Culicidae): a review. *Bull Entomol Res* **70**, 525-532.
- Gouck, H.K. (1972). Host preferences of various strains of *Aedes aegypti* and *Aedes simpsoni* as determined by an olfactometer. *Bull World Health Org* **47**, 680-683.
- Grabe, V., Strutz, A., Baschwitz, A., Hansson, B.S., and Sachse, S. (2015). Digital in vivo 3D atlas of the antennal lobe of *Drosophila melanogaster*. *J Comp Neurol* **523**, 530-544.
- Grant, A.J., Wigton, B.E., Aghajanian, J.G., and O'Connell, R.J. (1995). Electrophysiological responses of receptor neurons in mosquito maxillary palp sensilla to carbon dioxide. *J Comp Physiol [A]* **177**, 389-396.
- Grosse-Wilde, E., Kuebler, L.S., Bucks, S., Vogel, H., Wicher, D., and Hansson, B.S. (2011). Antennal transcriptome of *Manduca sexta*. *Proc Natl Acad Sci U S A* **108**, 7449-7454.
- Hallem, E.A., and Carlson, J.R. (2006). Coding of odors by a receptor repertoire. *Cell* **125**, 143-160.
- Hussain, A., Zhang, M., Üçpınar, H.K., Svensson, T., Quillery, E., Gompel, N., Ignell, R., and Grunwald Kadow, I.C. (2016). Ionotropic chemosensory receptors mediate the taste and smell of polyamines. *PLoS Biol* **14**, e1002454.
- Ignell, R., Dekker, T., Ghaninia, M., and Hansson, B.S. (2005). Neuronal architecture of the mosquito deutocerebrum. *J Comp Neurol* **493**, 207-240.
- Ihara, S., Yoshikawa, K., and Touhara, K. (2013). Chemosensory signals and their receptors in the olfactory neural system. *Neuroscience* **254**, 45-60.

- Ito, K., Shinomiya, K., Ito, M., Armstrong, J.D., Boyan, G., Hartenstein, V., Harzsch, S., Heisenberg, M., Homberg, U., Jenett, A., *et al.* (2014). A systematic nomenclature for the insect brain. *Neuron* 81, 755-765.
- Jafari, S., and Alenius, M. (2015). Cis-regulatory mechanisms for robust olfactory sensory neuron class-restricted odorant receptor gene expression in *Drosophila*. *PLoS Genet* 11, e1005051.
- Jones, W.D., Cayirlioglu, P., Kadow, I.G., and Vosshall, L.B. (2007). Two chemosensory receptors together mediate carbon dioxide detection in *Drosophila*. *Nature* 445, 86-90.
- Jové, V., Gong, Z., Hol, F.J.H., Zhao, Z., Sorrells, T.R., Carroll, T.S., Prakash, M., McBride, C.S., and Vosshall, L.B. (2020). Sensory discrimination of blood and floral nectar by *Aedes aegypti* mosquitoes. *Neuron* 10.1016/j.neuron.2020.09.019.
- Karner, T., Kellner, I., Schultze, A., Breer, H., and Krieger, J. (2015). Co-expression of six tightly clustered odorant receptor genes in the antenna of the malaria mosquito *Anopheles gambiae*. *Front Ecol Evol* 3, 10.3389/fevo.2015.00026.
- Kistler, K.E., Vosshall, L.B., and Matthews, B.J. (2015). Genome engineering with CRISPR-Cas9 in the mosquito *Aedes aegypti*. *Cell Rep* 11, 51-60.
- Klowden, M. (1981). Initiation and termination of host-seeking inhibition in *Aedes aegypti* during oöcyte maturation. *J Insect Physiol* 27, 799-803.
- Kwon, H.W., Lu, T., Rutzler, M., and Zwiebel, L.J. (2006). Olfactory responses in a gustatory organ of the malaria vector mosquito *Anopheles gambiae*. *Proc Natl Acad Sci U S A* 103, 13526-13531.
- Kwon, J.Y., Dahanukar, A., Weiss, L.A., and Carlson, J.R. (2007). The molecular basis of CO₂ reception in *Drosophila*. *Proc Natl Acad Sci U S A* 104, 3574-3578.
- Laissue, P.P., Reiter, C., Hiesinger, P.R., Halter, S., Fischbach, K.F., and Stocker, R.F. (1999). Three-dimensional reconstruction of the antennal lobe in *Drosophila melanogaster*. *J Comp Neurol* 405, 543-552.
- Larsson, M.C., Domingos, A.I., Jones, W.D., Chiappe, M.E., Amrein, H., and Vosshall, L.B. (2004). *Or83b* encodes a broadly expressed odorant receptor essential for *Drosophila* olfaction. *Neuron* 43, 703-714.
- Lee, Y., Poudel, S., Kim, Y., Thakur, D., and Montell, C. (2018). Calcium taste avoidance in *Drosophila*. *Neuron* 97, 67-74 e64.
- Li, H., Li, T., Horns, F., Li, J., Xie, Q., Xu, C., Wu, B., Kebschull, J.M., McLaughlin, C.N., Kolluru, S.S., *et al.* (2020). Single-cell transcriptomes reveal diverse regulatory strategies for olfactory receptor expression and axon targeting. *Curr Biol* 30, 1189-1198.e1185.
- Li, Q., Barish, S., Okuwa, S., Maciejewski, A., Brandt, A.T., Reinhold, D., Jones, C.D., and Volkan, P.C. (2016). A functionally conserved gene regulatory network module governing olfactory neuron diversity. *PLoS Genet* 12, e1005780.
- Lu, T., Qiu, Y.T., Wang, G., Kwon, J.Y., Rutzler, M., Kwon, H.W., Pitts, R.J., van Loon, J.J., Takken, W., Carlson, J.R., *et al.* (2007). Odor coding in the maxillary palp of the malaria vector mosquito *Anopheles gambiae*. *Curr Biol* 17, 1533-1544.
- Majeed, S., Hill, S.R., and Ignell, R. (2014). Impact of elevated CO₂ background levels on the host-seeking behaviour of *Aedes aegypti*. *J Exp Biol* 217, 598-604.
- Manning, L., and Doe, C.Q. (2017). Immunofluorescent antibody staining of intact *Drosophila* larvae. *Nat Protoc* 12, 1-14.
- Marder, E., and Goaillard, J.M. (2006). Variability, compensation and homeostasis in neuron and network function. *Nat Rev Neurosci* 7, 563-574.
- Matthews, B.J., Dudchenko, O., Kingan, S.B., Koren, S., Antoshechkin, I., Crawford, J.E., Glassford, W.J., Herre, M., Redmond, S.N., Rose, N.H., *et al.* (2018). Improved reference genome of *Aedes aegypti* informs arbovirus vector control. *Nature* 563, 501-507.
- Matthews, B.J., McBride, C.S., DeGennaro, M., Despo, O., and Vosshall, L.B. (2016). The neurotranscriptome of the *Aedes aegypti* mosquito. *BMC Genomics* 17, 32.
- Matthews, B.J., Younger, M.A., and Vosshall, L.B. (2019). The ion channel *ppk301* controls freshwater egg-laying in the mosquito *Aedes aegypti*. *Elife* 8, e43963.
- McBride, C.S., Baier, F., Omondi, A.B., Spitzer, S.A., Lutomiah, J., Sang, R., Ignell, R., and Vosshall, L.B. (2014). Evolution of mosquito preference for humans linked to an odorant receptor. *Nature* 515, 222-227.
- McIver, S.B. (1972). Fine structure of pegs on the palps of female culicine mosquitoes. *Can J Zool* 50, 571-576.
- McLaughlin, C.N., Brbić, M., Xie, Q., Li, T., Horns, F., Kolluru, S.S., Kebschull, J.M., Vacek, D., Xie, A., Li, J., *et al.* (2020). Single-cell transcriptomes of developing and adult olfactory receptor neurons in *Drosophila*. *bioRxiv* 10.1101/2020.10.08.332130.
- McMeniman, C.J., Corfas, R.A., Matthews, B.J., Ritchie, S.A., and Vosshall, L.B. (2014). Multimodal integration of carbon dioxide and other sensory cues drives mosquito attraction to humans. *Cell* 156, 1060-1071.
- Min, S., Ai, M., Shin, S.A., and Suh, G.S. (2013). Dedicated olfactory neurons mediating attraction behavior to ammonia and amines in *Drosophila*. *Proc Natl Acad Sci U S A* 110, E1321-1329.
- Mombaerts, P., Wang, F., Dulac, C., Chao, S.K., Nemes, A., Mendelsohn, M., Edmondson, J., and Axel, R. (1996). Visualizing an olfactory sensory map. *Cell* 87, 675-686.
- Montell, C. (2009). A taste of the *Drosophila* gustatory receptors. *Curr Opin Neurobiol* 19, 345-353.
- Neuhaus, E.M., Gisselmann, G., Zhang, W., Dooley, R., Storkuhl, K., and Hatt, H. (2005). Odorant receptor heterodimerization in the olfactory system of *Drosophila melanogaster*. *Nat Neurosci* 8, 15-17.

- Omer, S.M., and Gillies, M.T. (1971). Loss of response to carbon dioxide in palpectomized female mosquitoes. *Ent Exp & Appl* 14, 251-252.
- Potter, C.J., Tasic, B., Russler, E.V., Liang, L., and Luo, L. (2010). The Q system: a repressible binary system for transgene expression, lineage tracing, and mosaic analysis. *Cell* 141, 536-548.
- Prieto-Godino, L.L., Rytz, R., Cruchet, S., Bargeton, B., Abuin, L., Silbering, A.F., Ruta, V., Dal Peraro, M., and Benton, R. (2017). Evolution of acid-sensing olfactory circuits in *Drosophilids*. *Neuron* 93, 661-676.e666.
- Raji, J.I., Melo, N., Castillo, J.S., Gonzalez, S., Saldana, V., Stensmyr, M.C., and DeGennaro, M. (2019). *Aedes aegypti* mosquitoes detect acidic volatiles found in human odor using the *IR8a* pathway. *Curr Biol* 29, 1253-1262 e1257.
- Ray, A., van der Goes van Naters, W., and Carlson, J.R. (2008). A regulatory code for neuron-specific odor receptor expression. *PLoS Biol* 6, e125.
- Ressler, K.J., Sullivan, S.L., and Buck, L.B. (1994). Information coding in the olfactory system: evidence for a stereotyped and highly organized epitope map in the olfactory bulb. *Cell* 79, 1245-1255.
- Riabinina, O., Task, D., Marr, E., Lin, C.C., Alford, R., O'Brochta, D.A., and Potter, C.J. (2016). Organization of olfactory centres in the malaria mosquito *Anopheles gambiae*. *Nat Commun* 7, 13010.
- Riabinina, O., Vernon, S.W., Dickson, B.J., and Baines, R.A. (2019). Split-QF system for fine-tuned transgene expression in *Drosophila*. *Genetics* 212, 53-63.
- Robertson, H.M., Gadau, J., and Wanner, K.W. (2010). The insect chemoreceptor superfamily of the parasitoid jewel wasp *Nasonia vitripennis*. *Insect Mol Biol* 19 Suppl 1, 121-136.
- Robertson, H.M., Warr, C.G., and Carlson, J.R. (2003). Molecular evolution of the insect chemoreceptor gene superfamily in *Drosophila melanogaster*. *Proc Natl Acad Sci U S A* 100 Suppl 2, 14537-14542.
- Roth, L.M. (1951). Loci of sensory end-organs used by mosquitoes (*Aedes aegypti* (L.) and *Anopheles quadrimaculatus* say) in receiving host stimuli. *Ann Entomol Soc Am* 44, 59-74.
- Rubin, B.D., and Katz, L.C. (1999). Optical imaging of odorant representations in the mammalian olfactory bulb. *Neuron* 23, 499-511.
- Sato, K., Pellegrino, M., Nakagawa, T., Nakagawa, T., Vosshall, L.B., and Touhara, K. (2008). Insect olfactory receptors are heteromeric ligand-gated ion channels. *Nature* 452, 1002-1006.
- Schaefer, M.L., Finger, T.E., and Restrepo, D. (2001). Variability of position of the P2 glomerulus within a map of the mouse olfactory bulb. *J Comp Neurol* 436, 351-362.
- Scott, K., Brady, R., Jr., Cravchik, A., Morozov, P., Rzhetsky, A., Zuker, C., and Axel, R. (2001). A chemosensory gene family encoding candidate gustatory and olfactory receptors in *Drosophila*. *Cell* 104, 661-673.
- Shankar, S., and McMeniman, C.J. (2020). An updated antennal lobe atlas for the yellow fever mosquito *Aedes aegypti*. *PLoS Negl Trop Dis* 14, e0008729.
- Silbering, A.F., Rytz, R., Grosjean, Y., Abuin, L., Ramdya, P., Jefferis, G.S., and Benton, R. (2011). Complementary function and integrated wiring of the evolutionarily distinct *Drosophila* olfactory subsystems. *J Neurosci* 31, 13357-13375.
- Smallegange, R.C., Qiu, Y.T., van Loon, J.J., and Takken, W. (2005). Synergism between ammonia, lactic acid and carboxylic acids as kairomones in the host-seeking behaviour of the malaria mosquito *Anopheles gambiae sensu stricto* (Diptera: Culicidae). *Chem Senses* 30, 145-152.
- Smith, C.N., Smith, N., Gouck, H.K., Weidhaas, D.E., Gilbert, I.H., Mayer, M.S., Smittle, B.J., and Hofbauer, A. (1970). L-lactic acid as a factor in the attraction of *Aedes aegypti* (Diptera: Culicidae) to human hosts. *Ann Entomol Soc Am* 63, 760-770.
- Stocker, R.F. (1994). The organization of the chemosensory system in *Drosophila melanogaster*: a review. *Cell Tissue Res* 275, 3-26.
- Stocker, R.F., Lienhard, M.C., Borst, A., and Fischbach, K.F. (1990). Neuronal architecture of the antennal lobe in *Drosophila melanogaster*. *Cell Tissue Res* 262, 9-34.
- Strotmann, J., Conzelmann, S., Beck, A., Feinstein, P., Breer, H., and Mombaerts, P. (2000). Local permutations in the glomerular array of the mouse olfactory bulb. *J Neurosci* 20, 6927-6938.
- Syed, Z., and Leal, W.S. (2007). Maxillary palps are broad spectrum odorant detectors in *Culex quinquefasciatus*. *Chem Senses* 32, 727-738.
- Takken, W., and Kline, D.L. (1989). Carbon dioxide and 1-octen-3-ol as mosquito attractants. *J Am Mosq Control Assoc* 5, 311-316.
- Tanaka, N.K., Endo, K., and Ito, K. (2012). Organization of antennal lobe-associated neurons in adult *Drosophila melanogaster* brain. *J Comp Neurol* 520, 4067-4130.
- Task, D., Lin, C.C., Afify, A., Li, H., Vulpe, A., Menuz, K., and Potter, C.J. (2020). Widespread polymodal odorant receptor expression in *Drosophila* olfactory neurons. *bioRxiv in press*.
- Tauxe, G.M., MacWilliam, D., Boyle, S.M., Guda, T., and Ray, A. (2013). Targeting a dual detector of skin and CO₂ to modify mosquito host seeking. *Cell* 155, 1365-1379.
- Troemel, E.R., Chou, J.H., Dwyer, N.D., Colbert, H.A., and Bargmann, C.I. (1995). Divergent seven transmembrane receptors are candidate chemosensory receptors in *C. elegans*. *Cell* 83, 207-218.
- Turner, S.L., Li, N., Guda, T., Githure, J., Carde, R.T., and Ray, A. (2011). Ultra-prolonged activation of CO₂-sensing neurons disorients mosquitoes. *Nature* 474, 87-91.

Uchida, N., Takahashi, Y.K., Tanifuji, M., and Mori, K. (2000). Odor maps in the mammalian olfactory bulb: domain organization and odorant structural features. *Nat Neurosci* 3, 1035-1043.

Vassar, R., Chao, S.K., Sitcheran, R., Nunez, J.M., Vosshall, L.B., and Axel, R. (1994). Topographic organization of sensory projections to the olfactory bulb. *Cell* 79, 981-991.

Vidal, B., U. Aghayeva, H. Sun, C. Wang, L. Glenwinkel, E. A. Bayer, and Hobert, O. (2018). An atlas of *Caenorhabditis elegans* chemoreceptor expression. *PLoS Biol* 16, e2004218.

Vosshall, L.B., Amrein, H., Morozov, P.S., Rzhetsky, A., and Axel, R. (1999). A spatial map of olfactory receptor expression in the *Drosophila* antenna. *Cell* 96, 725-736.

Vosshall, L.B., Wong, A.M., and Axel, R. (2000). An olfactory sensory map in the fly brain. *Cell* 102, 147-159.

Vythilingam, I., Chiang, G.L., and Chan, S.T. (1992). Evaluation of carbon dioxide and 1-octen-3-ol as mosquito attractants. *Southeast Asian J Trop Med Public Health* 23, 328-331.

Wagh, D.A., Rasse, T.M., Asan, E., Hofbauer, A., Schwenkert, I., Dürbeck, H., Buchner, S., Dabauvalle, M.C., Schmidt, M., Qin, G., et al. (2006). Bruchpilot, a protein with homology to ELKS/CAST, is required for structural integrity and function of synaptic active zones in *Drosophila*. *Neuron* 49, 833-844.

Wang, J.W., Wong, A.M., Flores, J., Vosshall, L.B., and Axel, R. (2003). Two-photon calcium imaging reveals an odor-evoked map of activity in the fly brain. *Cell* 112, 271-282.

WHO (2020). Vector-borne diseases. World Health Organization, Geneva <http://www.who.int/mediacentre/factsheets/fs387/en/>.

Wicher, D., Schafer, R., Bauernfeind, R., Stensmyr, M.C., Heller, R., Heinemann, S.H., and Hansson, B.S. (2008). *Drosophila* odorant receptors are both ligand-gated and cyclic-nucleotide-activated cation channels. *Nature* 452, 1007-1011.

Yao, C.A., Ignell, R., and Carlson, J.R. (2005). Chemosensory coding by neurons in the coeloconic sensilla of the *Drosophila* antenna. *J Neurosci* 25, 8359-8367.

Zhao, Z., Zung, J.L., Kriete, A.L., Iqbal, A., Younger, M.A., Matthews, B.J., Merhof, D., Thiberge, S., Strauch, M., and McBride, C.S. (2020). Chemical signatures of human odour generate a unique neural code in the brain of *Aedes aegypti* mosquitoes. *bioRxiv* 10.1101/2020.11.01.363861.

Zou, D.J., Chesler, A., and Firestein, S. (2009). How the olfactory bulb got its glomeruli: a just so story? *Nat Rev Neurosci* 10, 611-618.

MATERIALS AND METHODS

Human and animal ethics statement:

Blood-feeding procedures and behavioral experiments with live hosts were approved and monitored by The Rockefeller University Institutional Animal Care and Use Committee (IACUC protocol 17018) and Institutional Review Board (IRB protocol LV-0652), respectively. Human volunteers gave their written informed consent to participate.

Mosquito rearing and maintenance

Aedes aegypti wild-type laboratory strains (Liverpool and Orlando), CRISPR-Cas9 knock-in, and piggyBAC QUAS transgenic strains were maintained and reared at 25 – 28°C, 70-80% relative humidity with a photoperiod of 14 hr light: 10 hr dark as previously described (DeGennaro et al., 2013). Adult mosquitoes were provided constant access to 10% sucrose. For routine strain maintenance, animals were primarily blood-fed on live mice and occasionally on live human volunteers. Newly generated strains were blood-fed on human volunteers until they were established. All experiments were conducted on female mosquitoes. Detailed genotypes used in each figure, and the use of Liverpool or Orlando as the wild-type strain in a given figure can be found in [Data File 1](#).

Generation of chemosensory receptor QF2 and Split-QF2 knock-in strains

T2A-QF2 gene-sparing stop codon replacement lines were generated using the strategy outlined in Matthews et al. (Matthews et al., 2019). sgRNAs were placed as close to the stop codon as possible and donor constructs were designed to remove the stop codon and replace it with an in-frame cassette containing the T2A ribosomal skipping sequence and the QF2 transcription factor or Split-QF2 domains, comprising the QF2 activation domain QF2-AD, or the QF2 DNA-binding domain QF2-DBD. This strategy spares the function of the gene at the locus being targeted, expresses QF2 or Split-QF2 domains in the cells specified by enhancers at the locus. Insertions were marked by the 3xP3 enhancer expressing a fluorescent protein.

To identify effective sgRNAs, 5 candidate sgRNAs per gene were first injected into separate pools of 500 Liverpool embryos and CRISPR-Cas9-mediated cut rate was evaluated as previously described (Kistler et al., 2015). Either a single sgRNA or 2 sgRNAs with the highest cut rates were then chosen to be injected with donor plasmids to target chemosensory gene loci using homology-directed repair. sgRNAs targeted the respective gene near the stop codon, target sequence with protospacer adjacent motif (PAM) underlined:

Ir25a: GTTTGTGTGCGTGTCCGTA TGG

Ir76b: GTATTACACTTATCTAAATA TGG

Ir8a: GTCACGCTTGTGTACAGGGCGG, TGG

GAACAATTTGAACAAGGTCTG TGG

Gr3: GTTAGTGATGCATAATATGA CGG

Orco: GTCACCTACTTCATGGTGT TGG

sgRNA DNA template was prepared by annealing oligonucleotides as described (Kistler et al., 2015). *In vitro* transcription was performed using HiScribe Quick T7 kit (NEB E2050S) following the manufacturer's directions. Following transcription and DNase treatment for 15 min at 37°C, sgRNA was purified using RNase-free SPRI beads (Ampure RNAclean, Beckman-Coulter A63987), and eluted in Ultrapure water (Invitrogen, 10977-015). Donor plasmids were constructed by Gibson assembly using the following fragments for QF2 lines:

1) *pUC19* digested with XbaI and BamHI

2) Left and right homology arms: *Gr3* (left: 1.9 kb, right: 1.6 kb), *Ir25a* (left: 1.8 kb, right: 1.6 kb), *Ir76b* (left: 1.2 kb, right: 2.2 kb), *Ir8a* (left: 1.7 kb, right: 1.7 kb), *Orco* (left: 1.2 kb, right: 1.3 kb) generated by polymerase chain reaction (PCR) using Liverpool genomic DNA as a template

3) A 2.6 kb fragment containing T2A-QF2-SV40, 3xP3-dsRed, PCR-amplified from a previously assembled vector (*ppk10779-T2A-QF2-SV40*, 3xP3-dsRed, Addgene accession #130667)

For Split-QF2 lines, donor plasmids were constructed by generating fragments using PCR from the indicated template with indicated primers in [Data File 1](#) and assembled using NEBuilder HiFi DNA Assembly (NEB E5520S):

Ir25a-T2A-QFAD::Zip+-SV40-3xP3-eYFP-SV40 was composed of:

1. Plasmid backbone with *Ir25* homology arms from *Ir25a-T2A-QF2* plasmid (6 kb)
 2. *T2A-QFAD::Zip+-SV40* sequence from (Riabinina et al., 2019), fragment synthesized by Genewiz, sequence in Data File 1 (1.5 kb)
 3. *3xP3-EYFP-SV40* from pDSAY (Addgene, #62291) (1.2 kb)
- Orco-T2A-Zip-::QFDBD-SV40-3xP3-dsRED-SV40* was composed of:

1. Plasmid backbone with *Orco* homology arms and *3xP3-dsRED-SV40* from *Orco-T2A-QF2* plasmid (6.3 kb)
2. *T2A-Zip-::QFDBD-SV40* synthesized by Genewiz, sequence in Data File 1 (1.5 kb)

For all QF2 and Split-QF2 constructs, the stop codon of the endogenous gene was removed and the PAM sequences corresponding to the sgRNAs used for injection were modified by PCR mutagenesis during Gibson assembly by introducing synonymous codon substitutions to protect the sequence from Cas9 cleavage while retaining the amino acid identity. Plasmids were isolated using an endotoxin-free plasmid midiprep kit (Macherey-Nagel) for QF2 lines and NucleoBond Xtra Midi Endotoxin-Free plasmid kit (Clontech 740420.50) for Split-QF2 lines and eluted in ultrapure water prior to injection. Donor plasmids are available at Addgene (accession numbers #162520-162526).

Approximately 2000 wild-type Liverpool strain *Ae. aegypti* embryos were injected with a mix containing recombinant Cas9 protein (PNA Bio, CP01) at 300 ng/μL, sgRNAs at 40 ng/μL and donor DNA plasmid (300 ng/μL for QF2 lines, 600 ng/μL for Split-QF2 lines) at the Insect Transformation Facility at the University of Maryland Institute for Bioscience & Biotechnology Research. Embryos were hatched and surviving G0 males and females were crossed to wild-type Liverpool mosquitoes and their G1 offspring were screened for fluorescence indicating positive stable germ line transformants.

For QF2 lines, the fidelity of insertion was verified by PCR and Sanger sequencing. One representative line for each chemosensory receptor QF2 knock-in was selected for further study. QF2-driven expression patterns were examined by crossing to *QUAS-CD8:GFP-3xP3-ECFP* and/or *QUAS-dTomato-T2A-GCaMP6s-3xP3-ECFP*. All lines were outcrossed to wild-type Liverpool mosquitoes for at least 3 generations prior to being used in experiments.

For Split-QF2 lines, a single family with the correct insertion was confirmed by PCR and Sanger sequencing for *Ir25a-QF2-AD* and *Orco-QF2-DBD*. To propagate these lines, a male founder was chosen to cross to wild-type Liverpool females. Animals were then back-crossed to Liverpool for at least 2 additional generations. To evaluate if the Split-QF2 system was functional in *Ae. aegypti*, *Ir25a-QF2-AD* was crossed to *QUAS-dTomato-T2A-GCaMP6s*. The resulting *Ir25a-QF2-AD*, *QUAS-dTomato-T2A-GCaMP6s* animals were then crossed to *Orco-QF2-AD*. Expression of the dTomato reporter was observed in larval antennae and subsequently confirmed in adult antennae and brains.

QUAS transgenic strains

QUAS-CD8:GFP-3xP3-ECFP and *QUAS-dTomato-T2A-GCaMP6s-3xP3-ECFP* transgenic strains were described previously (Matthews et al., 2019). Two independent insertions of the *QUAS-dTomato-T2A-GCaMP6s-3xP3-ECFP* reporter line (Jové et al., 2020; Matthews et al., 2019) were used in this study. These are located on different chromosomes and were used according to the crossing scheme needed for a given experiment. See Data File 1 for details.

Transcript abundance estimates of *Ae. aegypti* OR, IR, and GR genes

Expression values for adult sugar-fed, non-blood-fed female sensory tissues were retrieved from the *Ae. aegypti* L5 genome GitHub repository (<https://github.com/VosshallLab/AGWG-AaegL5>) at this link: <https://github.com/VosshallLab/AGWG-AaegL5/raw/master/AGWG%20AaegL5%20Chemoreceptor%20TPM.xlsx>. These

expression values reflect libraries from a previous transcriptome study (Matthews et al., 2016) that had been aligned to the *Ae. aegypti* genome (AaegL5) and chemoreceptor geneset annotation reported in units of Transcripts Per Million (TPM) (Matthews et al., 2018). The number of genes from each of three gene families (ORs, IRs, and GRs) with expression values above the indicated threshold were plotted in Figure 1F,G and are available in Data File 1.

Whole brain fixation and immunostaining

Dissection of adult brains and immunostaining was done as previously described (Matthews et al., 2019). 6-14 day-old mosquitoes were anesthetized on ice. Heads were carefully removed from the body by pinching at the neck with sharp forceps. Heads were placed in a 1.5 mL tube for fixation with 4% paraformaldehyde, 0.1 M Millonig's Phosphate Buffer (pH 7.4), 0.25% Triton X-100, and nutated for 3 hr. Brains were then dissected out of the head capsule in ice-cold Ca²⁺-, Mg²⁺-free phosphate buffered saline (PBS, Lonza 17-517Q) and transferred to a 24-well plate. All subsequent steps were done on a low-speed orbital shaker. Brains were washed in PBS containing 0.25% Triton X-100 (PBT) at room temperature 6 times for 15 min. Brains were permeabilized with PBS, 4% Triton X-100, 2% normal goat serum (Jackson ImmunoResearch #005-000-121) for ~48 hr (2 nights) at 4°C. Brains were rinsed once and then washed with PBT at room temperature 6 times for 15 min. Primary antibodies were diluted in PBS, 0.25% Triton X-100, 2% normal goat serum for ~48 hr (2 nights) at 4°C. Brains were rinsed once then washed in PBT at room temperature 6 times for 15 min. Secondary antibodies were diluted in PBS, 0.25% Triton X-100, 2% normal goat serum for ~48 hr (2 nights) at 4°C. Brains were rinsed once then washed in PBT at room temperature 6 times for 15 min. Brains were equilibrated overnight in Vectashield (Vector Laboratories H-1000) and were mounted in Vectashield.

The following primary antibodies were used: anti-Brp/nc82 (mouse; 1:500, Developmental Studies Hybridoma Bank – see below) and/or anti-GFP (rabbit: 1:10,000; Life Technologies A11122).

The secondary antibodies used in all experiments except Figure S1 and Figure S6 were anti-mouse-Cy5 (1:250; Life Technologies A-10524) and anti-rabbit-Alexa Fluor 488 (1:500; Life Technologies A-11034). In Figure S1, the secondary antibody was anti-mouse-Alexa Fluor 488 (1:500; Life Technologies A-11001) and in Figure S6, the secondary antibodies were anti-mouse-Alexa Fluor 594 (1:500; Life Technologies A-11005) and anti-rabbit-Alexa Fluor 488 (1:500; Life Technologies A-11034).

Purification of nc82/Brp monoclonal antibody

Hybridoma cells expressing monoclonal antibody nc82 (Antibody Registry ID: AB_2314866), which recognizes the *Drosophila melanogaster* Brp protein (Wagh et al., 2006)

developed by Erich Buchner were obtained from the Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa, Department of Biology, Iowa City, IA 52242. Frances Weis-Garcia and the members of the MSKCC Antibody and Bioresource Core Facility subsequently used these hybridoma cells to purify this monoclonal antibody. The hybridoma was adapted to Gibco™ Hybridoma-SFM (Cat # 12045084) and 1% fetal bovine serum prescreened for ultra-low levels of bovine Ig. Antibody expression was confirmed and the adapted hybridoma was inoculated into the cell compartment of the Corning™ CELLLine Disposable Bioreactor (Cat # 353137) in 15 ml of Hybridoma-SFM + 0.5% fetal bovine serum (production media) at 3 million viable cells / ml. The media compartment of the flask contained 350 ml of production media. The bioreactor was incubated at 37°C with 7% CO₂ for 3 days, at which time the cells and media containing nc82 were harvested. 30 million viable cells from the harvest were re-inoculated back into the cell compartment in 30 ml fresh production media. The media in the media compartment was replaced the following day with 650 ml production media.

Three days later, the media in the media compartment was replaced with 1,000 ml production media, with the next harvest 3 days later (7 days after the previous harvest). Cells were harvested weekly and fed bi-weekly until the desired amount of monoclonal antibody was reached. After the first harvest, each one contained about 3 mg of monoclonal antibody nc82/ml production media. The harvests to be purified were pooled, centrifuged at 12,855 x g for 15 min. 6.5 mg / run were loaded onto a Cytiva (formerly GE Life Sciences) 1 ml HiTrap Protein G HP antibody purification column (Cat # 29048581) at 1 ml / min. The column was then washed with 0.02 M Sodium Phosphate (pH 7.0) before the monoclonal antibody was eluted with 0.1 M Glycine-HCl (pH 2.7). One ml fractions were collected and immediately neutralized with 60 ml of 1.0 M Tris-HCl (pH 9.0). The harvest, flow through and fractions from the peak were run on an a 10% SDS-PAGE (Bio-Rad Cat # 345-0010) to confirm purity and determine which should be pooled. The pooled fractions of monoclonal antibody were dialyzed into PBS overnight using dialysis tubing (Spectrum™ 132544) with a 50 kDa MWCO. Another 10% SDS-PAGE was run and the concentration determined using the absorbance at 280 using an extinction coefficient of 1.43.

Antennal lobe confocal imaging

All brains were imaged using a Zeiss Inverted LSM 880 laser scanning confocal microscope with a 25x / 0.8 NA immersion-corrected objective unless otherwise noted. Glycerol was used as the immersion medium to most closely match the refractive index of the mounting medium Vectashield. Antennal lobes in [Figure 1, 2, 5, S2-S8](#) were imaged at either 1024 x 1024 or 2048 x 2048 pixel resolution in X and Y with 0.5 μ m Z-steps for a final voxel size of either 0.0615 x 0.0615 x 0.5 μ m³ or 0.1230 x 0.1230 x 0.5 μ m³. Both conditions oversampled relative to the objective resolution and no differences were noted between imaging conditions. The laser intensity and gain were adjusted along the Z-axis to account for a loss of intensity due to depth and care was taken to avoid saturation and ensure that the deepest glomeruli were visible for segmentation.

3xP3 was used as a promoter to express fluorescent proteins as markers for the knock-ins and QUAS transgenes used in this study, and care was taken to distinguish expression derived from the 3XP3 promoter from the expression of the QF2 driver and QUAS effector lines under investigation. 3xP3 drives expression in the optic lobes, as well as some cells in the dorsal brain. Neither area overlaps with the antennal lobes, and as reported previously (Matthews et al., 2019), we saw no 3xP3-driven expression in the antennal lobes in the reporter lines alone (data not shown). Representative antennal lobe images presented in the figures were cropped to remove 3xP3-driven expression elsewhere in the brain.

Antennal lobe glomerulus quantification

Confocal images of the antennal lobes in [Figure 1, 2, 5, S2-S8](#) were processed in ImageJ/FIJI (NIH). The number of glomeruli was quantified as follows: a single region of interest (ROI) was manually drawn around each glomerulus at a section approximately central along the Z-axis. Every glomerulus was outlined and an ROI set was collected that contained the outlines of all glomeruli. Glomeruli were then separated into two groups, GFP-positive and GFP-negative glomeruli. A count of each was made to determine the number of glomeruli labeled by each line as well as the total number of glomeruli. The ROIs were flattened along the Z-axis to enable representation of the data in two dimensions in [Figure 1, 2, S2-S5, S7](#). The left antennal lobe in 3 brains was analyzed for each genotype in [Figure 1](#) except for Gr3, for which the left antennal lobe was analyzed in 1 brain, and both left and right antennal lobes were analyzed in an additional 4 brains in [Figure S6](#).

Although we were able to recognize general regions of the antennal lobe, the interindividual variability made it impossible to identify most glomeruli by anatomy alone. We therefore have not attempted to name and number every glomerulus in *Ae. aegypti* as

has been done in previous studies (Ignell et al., 2005; Shankar and McMeniman, 2020). As noted by Ito et al. (Ito et al., 2014), there is considerable confusion about the use of coordinate axes in the brains of animals in general and insects in particular. The glomeruli in the antennal lobe of *Ae. aegypti* were originally named by Ignell et al. (Ignell et al., 2005) using a set of coordinate axes that differ from those consistently used in *Drosophila melanogaster* (Couto et al., 2005; Fishilevich and Vosshall, 2005; Grabe et al., 2015; Laissue et al., 1999; Stocker et al., 1990). A recent study of the antennal lobe of *Ae. aegypti* renamed glomeruli to account for this discrepancy in coordinate axes (Shankar and McMeniman, 2020), and throughout this paper we use the same coordinate axes they have implemented. While Shankar and McMeniman renamed most antennal lobe regions and glomeruli, they chose not to rename the MD (Medio-Dorsal) cluster of glomeruli comprising MD1, MD2, and MD3 whose sensory input derives from the maxillary palp. We have observed in our study that the MD glomeruli are medial, but they are not notably dorsal, and therefore refer to them as Glomerulus 1, Glomerulus 2, and Glomerulus 3 in this paper for simplicity. While there is utility in naming glomeruli, we suspect that the *Ae. aegypti* mosquito antennal lobe atlas will be refined in the future with the advent of new genetic tools that will unambiguously allow the field to distinguish and name genetically identifiable glomeruli. We found that the size, shape, and number of antennal lobe glomeruli in *Ae. aegypti* was variable from animal to animal. It is possible that the boundaries between glomeruli are not easily distinguished by synaptic staining and that specific glomeruli will become identifiable once there are genetic tools available that label smaller populations of olfactory sensory neurons. The anatomical variability we see is consistent with both the original map that identified 50 glomeruli (Ignell et al., 2005), which divided glomeruli into 3 classes based on their variability in location, as well as a recent study that looked specifically at the size and shape of glomeruli across animals (Shankar and McMeniman, 2020) and revised the original map to a count of ~80 glomeruli. Shankar and McMeniman named and numbered these glomeruli across animals, but they noted that they were only able to consistently identify 63 glomeruli. This is similar to the ~65 glomeruli we observed in our work. While there is not yet a clear consensus on the exact number of antennal lobe glomeruli in *Ae. aegypti*, the number of chemosensory receptors expressed in the antenna and maxillary palp is at least twice as large as any of the estimates of glomerulus number.

The variability in antennal lobe structure appears at first to contrast with *Drosophila melanogaster*, where each glomerulus can be clearly identified and named. However, we note that the antennal lobe map in *Drosophila melanogaster* has been refined with the advent of new genetic techniques, starting with 35 glomeruli in the original atlas (Stocker et al., 1990), then modified to 40 glomeruli (Laissue et al., 1999), and further refined in numerous studies (Couto et al., 2005; Fishilevich and Vosshall, 2005; Tanaka et al., 2012) including a recent count of 54 (Grabe et al., 2015) and 58 (Task et al., 2020) glomeruli. We have refrained from naming glomeruli in *Ae. aegypti* at this time because we believe that a more stereotyped anatomy will emerge as new genetic lines are generated that allow cell-type-specific labelling.

A recent study in the mosquito *Anopheles gambiae* using mosquitoes that label *Orco*-expressing olfactory neurons also noted that the antennal lobe was variable between animals relative to *Drosophila melanogaster* (Riabinina et al., 2016). It is therefore possible that mosquito antennal lobes are more variable than *Drosophila* (Grabe et al., 2015; Prieto-Godino et al., 2017). Variability in olfactory bulb structure is seen even in the mouse, *Mus musculus*, where the principles of olfactory organization were first established (Schaefer et al., 2001; Strotmann et al., 2000; Zou et al., 2009). The exact size and location of glomeruli can vary between animals more than initially appreciated and appears to be determined by both genetic factors and activity in olfactory sensory

neurons during the early life of the animal. In *Drosophila melanogaster*, glomerulus size is highly genetically determined and correlates strongly with the number of olfactory sensory neurons that innervates each glomerulus (Grabe et al., 2015). Whether the variability in glomerulus size in the mosquito is due to activity-dependent changes in structure or other factors remains to be seen.

Additional technical notes on expression and projection patterns of chemosensory receptor knock-in strains

Orco-QF2>QUAS-mCD8:GFP: We noted that the intensity of GFP varies between glomeruli in this driver line, with some bright and others comparably dim. We speculate that this is due to a combination of the variability in Orco expression levels in individual neurons and variability in the density of innervation in individual glomeruli. A large region of the anterior ventral antennal lobe was previously referred to as the Johnston's organ center and was thought to comprise a single large glomerulus (Ignell et al., 2005). In other insect species, Johnston's organ mediates detection of auditory cues. Consistent with a recent study (Shankar and McMeniman, 2020), we segmented this region into multiple glomeruli based on anatomical boundaries revealed with Brp immunofluorescence. Glomeruli in this region are innervated by Orco-expressing neurons, calling into doubt the original report that these glomeruli process auditory stimuli and suggesting instead that they serve an olfactory function. In support of this hypothesis, the analogous area of the *Anopheles coluzzii* antennal lobe has been shown to receive projections from Orco-expressing olfactory sensory neurons (Riabinina et al., 2016). We also observed GFP projections into the subesophageal zone in Orco-QF2>QUAS-mCD8:GFP animals, which appear to derive from expression in the proboscis, the primary taste organ in insects. This is consistent with similar expression in *Anopheles coluzzii* (Riabinina et al., 2016) and functional data in *Anopheles gambiae* showing that olfactory responses are detected in this gustatory organ (Kwon et al., 2006).

Ir25a-QF2>QUAS-mCD8:GFP: The intensity of GFP projections varies between glomeruli in this driver line, with some bright and other comparably dim, as noted for Orco-QF2. The brightest glomeruli are primarily medial and anterior. We see the dimmest innervation in the area previously described as Johnston's organ center as well as in the central antennal lobe. Labeling was also seen in other areas of the brain, most notably the subesophageal zone and anterior mechanosensory motor center.

Ir8a-QF2>QUAS-mCD8:GFP: Depending on the brain being analyzed there were either 2 or 3 medial glomeruli labelled in this line. In the cases where there were 3 medial glomeruli, this third medial glomerulus was innervated by a few large-diameter axons. These were larger and sparser than the smaller axons that densely innervated most other glomeruli in this line. We also note that there are 2-3 cell bodies that express GFP located in the cell body rind lateral to the antennal lobe (rALI). We are unable to definitively describe where these cells project without genetic reagents that selectively label these cells, but they appear to send bilateral processes that cross the midline within what appears to be the saddle to innervate the anterior mechanosensory motor center outside the antennal lobe. All naming is in accordance with the new insect brain nomenclature presented in Ito et al. (Ito et al., 2014).

Ir76b-QF2>QUAS-mCD8:GFP: In addition to projections to the antennal lobe, this line shows innervation of the subesophageal zone of the brain.

Gr3-QF2>QUAS-mCD8:GFP: All antennal lobes in this line show innervation of a single glomerulus (also referred to as "MD1" and here referred to as "Glomerulus 1"; (Ignell et al., 2005; Shankar and McMeniman, 2020). In several brains, we saw a second small medial glomerulus that derives its innervation from the antenna and is in a small medial cluster of landmark glomeruli midway down the anterior-posterior axis closest to the center of the brain. Innervation appears to come from only a few axons.

This low and variable reporter expression is consistent with the low level of expression of Gr3 in the antennal transcriptome (Matthews et al., 2016). Because this line only shows innervation of these 1-2 glomeruli, we analyzed all glomeruli only in the single brain in Figure 11, and additionally analyzed 8 more antennal lobes in 4 brains for the presence or absence of labelling in these two glomeruli.

We analyzed both left and right antennal lobes from 4 brains and found that in 3 of the 4 brains there was a second glomerulus in one or both antennal lobes (Figure S6). The presence of the second glomerulus was not specific within a single animal as we found all variations of presence and absence of this glomerulus across both antennal lobes in these 4 animals. In some Gr3-QF2>QUAS-mCD8:GFP animals, we detected a small number of processes that extended beyond the antennal lobe and into the higher brain, although the exact termination site varied. We never saw CO₂-evoked activity in the variable second glomerulus or these projections outside the antennal lobe. Images in Figure S6 were taken as described above with the following changes: Secondary antibodies used were anti-mouse-Alexa Fluor 594 (1:500; Life Technologies A-11005) and anti-Rabbit-Alexa Fluor 488 (1:500; Life Technologies A-11034). Images were taken using a Zeiss Inverted LSM 880 laser scanning confocal microscope with a Plan-Apochromat 40x/1.4 Oil DIC objective. Images were taken at 1024 x 1024 in XY to generate images with a final voxel size of 0.1384 x 0.1384 x 0.5 μm³. Images were scored as containing GFP in one or two glomeruli.

Additional technical notes on expression and projection patterns of Split-QF2 strains

Ir25a-QF2-AD, Orco-QF2-DBD > QUAS-dTomato-T2A-GCaMP6s: All antennal lobe immunostaining in Figure 2, 5, S7, S8 was carried out as described above with slight modifications to utilize the 15xQUAS-dTomato-T2A-GCaMP6s effector line. The same primary antibodies were used because of the structural similarity between GCaMP6s and GFP. Intrinsic dTomato was detected without antibody amplification, as it retained fluorescence throughout fixation and staining. Brp (Cy5), dTomato, and GCaMP6s (Alexa Fluor 488) were imaged as three separate confocal channels as described above. Glomeruli labelled by dTomato completely overlapped with those labelled by GCaMP6s immunofluorescence, so both channels were used during the quantification of positive and negative glomeruli. dTomato labeling was used to generate sample images. There was no staining in the antennal lobes of the individual split effector lines crossed to 15xQUAS-dTomato-T2A-GCaMP6s (n=3 / genotype, Figure 2, S7).

Due to an initial observation that certain glomeruli were consistently unlabeled in the Ir25a-AD;Orco-DBD line but were labelled in the Orco-QF2 and Ir25a-QF2 lines, particularly in an anterior region of the brain, we conducted a second experiment to compare glomeruli labelled by the Orco-QF2, Ir25a-QF2 and Ir25a-AD;Orco-DBD drivers side-by-side. All drivers were crossed to the 15xQUAS-dTomato-T2A-GCaMP6s effector (Figure 2J). 3 brains per genotype were immunostained and imaged as described above.

Antennal lobe anterograde dye fill

For images in Figure S1, mosquitoes were anesthetized on ice until immobile and then transferred to a cold dissection dish. A single antenna or maxillary palp was loaded with Texas-red conjugated dextran (Molecular Probes D3328) diluted 10 mg in 100 μL external saline (103 mM NaCl, 3 mM KCl, 5 mM 2-[Tris(hydroxymethyl)methyl]-2-aminoethanesulfonic acid (TES), 1.5 mM CaCl₂, 4 mM MgCl₂, 26 mM NaHCO₃, 1 mM NaH₂PO₄, 10 mM trehalose, 10 mM glucose, pH 7.3, osmolality adjusted to 275 mOsm/kg). To load the dye a small drop (approximately 0.5-1 μL) of dye was placed onto the surface of the dish and the animal was moved such that the intended cut-site on a single antenna or maxillary palp was placed in the drop of dye. The antenna or maxillary palp was then removed with sharp forceps and a fine scalpel (F.S.T 10315-12) while it was

submerged in the dye. Care was taken to remove the maxillary palp proximal to the 4th segment, to include all the basiconic sensilla, and to remove the antenna near the base but to leave the antennal pedicel completely intact. The animal remained immobile on ice with the antenna or maxillary palp submerged and the dye was loaded for 2-5 min. After this time the animal was placed in a small soup cup with access to 10% sucrose and returned to standard rearing conditions overnight to give the dye time to diffuse throughout the neurons and fill the length of the axon. The next morning dissection of adult brains and immunostaining was carried out as described above.

Antennal lobe 3-D reconstructions

In an attempt to develop a map of the *Ae. aegypti* antennal lobe, 3 brains from the Liverpool strain were immunolabeled with Brp to identify the boundaries between antennal lobe glomeruli. The left antennal lobe in each brain was independently reconstructed from confocal sections taken with a Plan-Apochromat 63x/1.40NA oil immersion objective, at 1024 x 1024 pixel resolution in X and Y with 0.5 μ m Z-steps for a final voxel size of either 0.1318 x 0.1318 x 0.5 μ m³ using the software Imapis (Bitplane). Although the area previously termed Johnston's organ center was considered a single glomerulus in a previous study (Ignell et al., 2005), we noted anatomical boundaries in this region, suggesting that it contains multiple glomeruli. This observation is consistent with recently published work (Shankar and McMeniman, 2020) and this area was segmented by an individual researcher to generate the final reconstructions. Two of these are shown in [Figure S1](#). Each glomerulus was manually segmented into an individual surface using Surpass View. We were consistently able to identify the three glomeruli innervated by the maxillary palp, previously termed MD1, MD2 and MD3 (Ignell et al., 2005) which we refer to in this study as Glomerulus 1, Glomerulus 2, and Glomerulus 3 ([Figure 1](#), [Figure 5](#)). The overall structure of the antennal lobe varied considerably from animal to animal and although we were able to identify certain regions and certain landmark glomeruli including those that are targeted by the maxillary palp, we were unable to assign an unambiguous identity to every glomerulus, as is possible in *Drosophila melanogaster* (Couto et al., 2005; Fishilevich and Vosshall, 2005). This variability makes it essentially impossible to identify a given glomerulus between animals and we therefore have decided to avoid referring to glomeruli by previous naming schemes, including MD1, MD2, MD3. An authoritative atlas of the *Ae. aegypti* antennal lobe awaits genetic reagents that label subpopulations of sensory neurons that will permit the field to refer to glomeruli by their molecular identity.

Antennal whole mount immunofluorescence

Whole-mount immunostaining of adult antennae was performed as described (Riabinina et al., 2016) with modifications. 7-11 day-old Liverpool mosquitoes were immobilized on ice, decapitated and heads and placed in 1 mL ZnFA fixative solution (0.25% ZnCl₂, 2% paraformaldehyde, 135 mM NaCl, 1.2% sucrose and 0.03% Triton X-100) for 20–24 h at room temperature in the dark. Next, the heads were washed three times for 30 min each with HBS buffer (150 mM NaCl, 5 mM KCl, 25 mM sucrose, 10 mM HEPES, 5 mM CaCl₂ and 0.03% Triton X-100). Antennae were carefully removed in HBS on ice and placed in 400 μ L HBS in 0.5 mL Eppendorf tubes. After a brief wash in HBS, the tissue was incubated in 400 μ L 80% methanol/20% dimethyl sulfoxide (DMSO) solution for 1 h at room temperature, washed for 5 min in 400 μ L 0.1 M Tris pH 7.4, 0.03% Triton X-100 solution and incubated in 400 μ L blocking solution (PBS, 5% normal goat serum (Jackson 005-000-121), 1% DMSO and 0.3% Triton X-100) for at least 3 h at room temperature or overnight at 4°C. Next, the tissue was placed in a 0.5 mL Eppendorf tubes containing 400 μ L blocking solution with primary antibodies [rabbit anti-Orco EC2 (Larsson et al., 2004), 1:50, Vosshall lab; chicken anti-GFP, 1:200, Aves GFP-1020] and submerged and held in a water bath sonicator (Branson m1800) for 30 sec at the high setting. Next, the tubes were placed on a rotator for 2 days at

4°C in the dark, after which the sonication procedure was repeated. The tubes were placed on a rotator for 2 additional days (for a total of 4 days) at 4°C in the dark. Next, the tissue was washed 5X 30 min each at room temperature in PBS, 1% DMSO and 0.3% Triton X-100. Secondary antibodies (anti-rabbit Alexa Fluor 555 Plus, 1:200, Thermo Fisher A32732, anti-chicken Alexa Fluor 488, 1:200, Thermo Fisher A-11039) and nuclear dye (TO PRO 3 Iodide, 1:400, Thermo Fisher T3605) were added to the blocking solution, and tubes were sonicated as described above and incubated for 4 days at 4°C in the dark with the sonication repeated after 2 days of incubation. The tissue was then washed 5X 30 min at room temperature in PBS, 1% DMSO and 0.3% Triton X-100, rinsed in PBS and mounted in Slow Fade Diamond for confocal imaging.

Whole mount antennal and maxillary palp RNA in situ hybridization

RNA was detected in whole mount antenna and maxillary palp using the hybridization chain reaction (HCR) technique as previously described (Choi et al., 2018) with modifications. Probes, amplifiers, Probe Hybridization Buffer, Amplification Buffer, and Probe Wash Buffer were purchased from Molecular Instruments. Full list of probe lot numbers can be found in [Date File 1](#). 5-8 day-old Liverpool mosquitoes were cold anesthetized, manually decapitated with forceps, and heads with antennae and the proboscis were digested in a chitinase-chymotrypsin solution (119 mM NaCl, 48 mM KCl, 2 mM CaCl₂, 2 mM MgCl₂, 25 mM HEPES, 5 U/mL chitinase (Sigma-Aldrich C6137-50UN), 100 U/mL alpha-chymotrypsin (Sigma-Aldrich CHY5S-10VL), 2% DMSO) (Manning and Doe, 2017) at 37°C for 30 min (antennae) or 1 hr (maxillary palps) in a Fisher Isotemp oven and subsequently fixed in 4% paraformaldehyde, 1X PBS, 0.03% Triton X-100 on a rotator at 4°C overnight. Heads were washed 4 times on ice for 10 min each in 0.1% PBS-Tween-20. Antennae or maxillary palps were dissected in 0.1% PBS-Tween-20 on ice and dehydrated with a graded series of methanol/0.1% PBS-Tween: 25% methanol in 0.1% PBS-Tween-20 for 10 min on ice, 50% methanol in 0.1% PBS-Tween-20 for 10 min on ice, 75% methanol in 0.1% PBS-Tween-20 for 10 min on ice, and two washes of 100% methanol for 10 min on ice. Tissues were incubated overnight in 100% methanol at -20°C and were subsequently rehydrated with a series of graded methanol/0.1% PBS-Tween-20: 75% methanol in 0.1% PBS-Tween-20 for 10 min on ice, 50% methanol in 0.1% PBS-Tween-20 for 10 min on ice, 25% methanol in 0.1% PBS-Tween-20 for 10 min on ice, and two washes of 0.1% PBS-Tween-20 for 10 min each on ice. Tissue was digested in 20 μ g/mL Proteinase-K (Thermo Fisher AM2548) in 0.1% PBS-Tween for 30 min at room temperature and washed twice with 0.1% PBS-Tween-20 for 10 min each at room temperature. Tissue was fixed in 4% paraformaldehyde in 0.1% PBS-Tween-20 for 20 min at room temperature and washed 3 times for 10 min each in 0.1% PBS-Tween-20 at room temperature.

Tissue was incubated in Probe Hybridization Buffer at room temperature for 5 min and then in 37°C pre-warmed Probe Hybridization Buffer rotating in a hybridization oven for 30 min. 8 pmol of each probe set was prepared in 37°C pre-warmed Probe Hybridization Buffer and tissue was incubated in probe solution at 37°C in a hybridization oven for 2 nights. Tissues were washed in 37°C pre-warmed Probe Wash Buffer 5 times for 10 min each at 37°C. Tissues were washed twice in 5X SSC 0.1% Tween-20 at room temperature for 10 min each. Tissues were pre-amplified in room temperature Amplification Buffer for 10 min. 18 pmol hairpins were separately prepared by heating 6 μ L of 3 μ M stock of hairpins H1 and H2 at 95°C for 90 sec on an Eppendorf Mastercycler and allowing to cool to room temperature in a dark drawer for 30 min. Hairpins were resuspended in 100 μ L amplification buffer and tissues were incubated in this hairpin solution in the dark on a rotator at room temperature overnight. Tissues were washed 5 times for 10 min each in 5X SSC 0.1% Tween-20 and mounted in SlowFade

Diamond (Thermo Fisher S36972) on glass slides with coverslips for confocal imaging.

Whole mount antennal and maxillary palp dTomato visualization

7-14 day-old *Ir25a-QF2*, *Orco-QF2*, *Ir25a-QF2AD*, *Orco-QFDBD*, and *Ir25a-QF2AD,Orco-QFDBD > 15XQUAS-dTomato-T2A-GCaMP6s* mosquitoes were cold anesthetized, manually decapitated with forceps and heads with antennae and maxillary palps were immediately fixed in 1 mL 4% paraformaldehyde, 1X PBS, 0.03% Triton X-100, on a rotator in the dark at 4°C overnight. Heads were washed 3X 30 min each in 1X PBS, 0.03% Triton X-100 at room temperature, then antennae and maxillary palps were carefully removed and placed in 1X PBS, 0.03% Triton X-100. Next, antennae and maxillary palps were placed in a solution of 1X PBS, 0.03% Triton X-100, 1% DMSO, and a 1:400 dilution of TO PRO 3 (Thermo Fisher T3605) for 24 hr at 4°C in the dark. Antennae and maxillary palps were then washed 5X 30 min each in 1X PBS, 0.03% Triton X-100 at room temperature in the dark, washed once with 1X PBS, transferred to a well of SlowFade diamond to remove excess PBS, and mounted in SlowFade Diamond for confocal imaging.

Antennal and maxillary palp confocal imaging and cell quantification

Images of peripheral tissues were acquired with a Zeiss Axio Observer Z1 Inverted LSM 880 NLO laser scanning confocal microscope (Zeiss) with a 25x/0.8 NA or 63x/1.4 NA immersion-corrected objective at a resolution of 3096 x 3096 pixels or 2048 x 2048 pixels. When comparing dTomato fluorescence across genotypes, image acquisition parameters were kept consistent. When necessary, tiled images were stitched with 20% overlap. Confocal images were processed in ImageJ (NIH). Because the antenna is a cylindrical structure, when whole antennal segments are mounted on a slide and imaged on a confocal microscope, signal can be easily detected from the region closest to the coverslip and confocal objective, but signal is weaker when imaging the side further from the coverslip and objective. For the purposes of consistent quantification, we only quantified cell numbers from the region closest to the coverslip (orange region in Figure S9A). For quantifying expression in the maxillary palp, only the dorso-lateral region of the 4th maxillary palp segment was analyzed. (yellow region in Figure S9B). Quantification of co-expression in antennae and maxillary palps was done in ImageJ (NIH) using the Cell Counter plugin. Cells in each channel were manually marked independently of the signal in the other channels. After cells in each channel are marked, and markers were then merged. Cells that were labeled with multiple markers (co-expressing cells) were then marked with a third marker (Figure S9C-H). Cell counts were then imported into Microsoft Excel and R for analysis.

In vivo 2-photon GCaMP calcium imaging preparation

Calcium imaging was performed on an Ultima IV two-photon laser-scanning microscope (Bruker Nanosystems) equipped with galvanometers and illuminated by a Chameleon Ultra II Ti:Sapphire laser (Coherent). GaAsP photomultiplier tubes (Hamamatsu) were used to collect emitted fluorescence. Images were acquired with a 60X/1.0N.A. Long Working Distance Water-Immersion Objective (Olympus) at a resolution of 256 x 256 pixels. GCaMP6s was expressed via the *Ir25a-QF2* knock-in which labels all three maxillary palp glomeruli. Preliminary imaging studies were conducted in animals that were wild-type at the *Gr3* locus. In Figure 5 all *Gr3*^{+/+} and *Gr3*^{-/-} animals were reared and imaged in parallel. Because the *Gr3* mutants were generated in the Orlando wild-type background and all Q-system lines were generated in the Liverpool wild-type background, care was taken to compensate for background effects in the following way. A single stock was created containing the *15x-QUAS-dTomato-T2A-GCaMP6s-3xP3-ECFP* transgene (Matthews et al., 2019), the *Ir25a-QF2-3xP3-dsRed* gene-sparing knock-in, and the *Gr3^{ECFP}* knock-out allele that is marked with

ECFP (McMeniman et al., 2014). This parental line was then crossed to either the unmarked *Gr3^{Δ4}* mutant that is in the Orlando background (McMeniman et al., 2014) to generate *Gr3*^{-/-} animals, or to wild-type Orlando to generate *Gr3*^{+/+} animals. The *Gr3* mutation is recessive (McMeniman et al., 2014), so this comparison of heteroallelic and heterozygous animals both controls for genetic background and makes it possible to assay the *Gr3* mutant phenotype.

Calcium imaging experiments were performed on 7-14 day-old female mosquitoes. Mosquitoes were fed on 10% sucrose and switched to water overnight prior to imaging. Mosquitoes were anesthetized at 4°C for dissection. The mosquito was fixed to a custom Delrin plastic holder with UV-curable glue (Bondic). The mosquito was inserted into a hole in the holder, such that the head and thorax were exposed above the surface of the holder, with the rest of the mosquito below. The mosquito was secured with a few points of glue (Bondic) on the thorax and head. The antennae and maxillary palp remained below the plate and were kept free of saline to prevent damage to the tissue. The top of the dish was then filled with external saline, which is based on *Drosophila melanogaster* imaging saline (Fişek and Wilson, 2014) with the following composition: 103 mM NaCl, 3 mM KCl, 5 mM 2-[Tris(hydroxymethyl)methyl]-2-aminoethanesulfonic acid (TES), 1.5 mM CaCl₂, 4 mM MgCl₂, 26 mM NaHCO₃, 1 mM NaH₂PO₄, 10 mM trehalose, 10 mM glucose, pH 7.3, osmolality adjusted to 275 mOsm/kg). A small window was gently opened in the head and the cuticle was removed anterior to the antennal pedicels and the tissue was secured with dental wax to expose the antennal lobes. Opaque non-neural tissue, primarily fat cells, was removed if they obstructed the antennal lobes. Great care was taken not to damage the antennal nerves. The preparation was placed on a custom laser cut acrylic holder that was secured to the stage and placed the mosquito in close proximity to the output port of the olfactometer. Plans for the fabrication of the imaging holder are available on Github: https://github.com/VosshallLab/Younger_Herre_Vosshall2020. On rare occasions mosquitoes showed no movement or odor responses and were discarded.

Image volumes were taken at higher resolution to identify glomeruli innervated by the maxillary palp, including the nerves that innervate these glomeruli. Glomerulus 1 was always easily identifiable by eye. A single plane through the center of each glomerulus was scanned at 4.22 frames per second with a 920 nm excitation wavelength imaged through a 680 nm shortpass infrared (IR) blocking filter, a 565 nm longpass dichroic and 595/50 nm or 525/70 nm bandpass filters. GCaMP6s and dTomato emission was collected simultaneously for at least 70 frames per trial.

In vivo 2-photon GCaMP calcium imaging stimulus delivery

Odor stimulation was achieved by directing a continuous stream (400-800 mL/min) of clean air, originating from a breathing air tank (Praxair), through 0.125 inch inner diameter/0.250 inch outer diameter perfluoroalkoxy alkaline (PFA) tubing. The antenna and maxillary palp of the mosquitoes were positioned close to the tube such that they were exposed to constant air flow (carrier stream). 4-60% of the total air stream was diverted through the headspace of a 20 mL borosilicate glass vial (EnviroWare #03-339-14E) containing the odorant diluted in either paraffin oil or water (odor stream). At a trigger, a custom-built solenoid valve controller system redirected the odor stream from a blank vial to a vial containing various odorants diluted in paraffin oil or water to a final volume of 3 mL. Unless otherwise noted, 40% of the air stream was used for odor delivery. Prior to conducting experiments, a mini-PID (Aurora Scientific, model 201A) was used to measure odorant waveforms and ensure the consistency of odor presentations across trials.

CO₂ was introduced into this system from a tank of 5% CO₂ in a 20.9% oxygen/ 79.1% nitrogen balance (Praxair). At a trigger, CO₂ was diverted through the headspace of a second clean 20 mL borosilicate glass vial (CO₂ stream) that was redirected from a bypass

stream into the system. This was coupled to a clean air stream with a matched flow rate that was simultaneously removed from the system. In experiments where odor concentration of CO₂ concentration was varied, the fraction of odor stream directed to the mosquito was adjusted. CO₂ was delivered at 4-60% of the total air stream for total concentrations of 0.2- 3.0% CO₂. In **Figure 5**, a final concentration of 2% CO₂ was used in all experiments. Prior to conducting experiments, a CO₂ meter (Vaisala GM70) was used to measure CO₂ concentration in the final air stream and we found that the expected concentration was delivered. Odor and CO₂ delivery was controlled using custom software written in Matlab R2020a (MathWorks). The odor delivery system and software was based on a previously used system and updated for CO₂ delivery (Cohn et al., 2015).

After the mosquito was transferred to the microscope and placed within the air stream, there was a 10-min acclimation period before any imaging began. For each trial, 5 sec of GCaMP6s fluorescence at baseline was imaged and then a one second odor or CO₂ pulse was delivered while imaging continued for at least 10 more sec (at least 70 frames total). There was a one-min delay between imaging trials so that the mosquito could reacclimate to the air stream after exposure to each odor. In an initial experiment we did not see any change in the amplitude of the CO₂ response across multiple trials within an animal. Imaging remained stable during the duration of the imaging session in all animals that were included in this study. We did not notice a decrease in the response to stimuli over time. Odorants were selected for the highest purity available for purchase (greater than 98%) and were either diluted in MilliQ ultrapure water (resistance 18 megaohm) or paraffin oil (EMD Millipore #PX0045-3) depending on solubility. All odorants were diluted into a large stock solution that was used throughout each entire experiment to avoid variability in concentrations.

The odorants used in **Figure 5** were: R-(-)-1-octen-3-ol (PubChem CID: 6992244, Penta Manufacturing 15-18900) diluted 1:10 in paraffin oil; 3-octanone (PubChem CID: 246782, Sigma 136913) diluted 1:10 in paraffin oil; acetone (PubChem CID: 180 Sigma A4206) diluted to 5%v/v in MilliQ ultrapure water; ammonia (PubChem CID: 222, Suprapur, Supelco 105428) diluted to 2.5% v/v in MilliQ ultrapure water; hexyl amine (PubChem CID: 8102, Sigma 219703) diluted to 1% v/v in MilliQ ultrapure water; triethyl amine (PubChem CID: 1146, Sigma T0886) diluted to 1% v/v in MilliQ ultrapure water.

In vivo 2-photon GCaMP calcium imaging data analysis

All image processing was done using FIJI/ImageJ (NIH). Data analysis used Matlab R2020a (MathWorks), Excel (Microsoft), and Prism (GraphPad). Regions of interest were selected based on the dTomato fluorescence intensity and GCaMP6s was normalized to dTomato intensity for analysis. A Gaussian blur with a sigma value of 1 was performed on the GCaMP6s signal. In the calculation of $\Delta F/F$, six frames were averaged before stimulus presentation to determine the baseline fluorescence. To determine maximum $\Delta F/F$, the average of the maximum 3 frames at the peak after stimulus delivery was determined for each sweep.

Quattroport olfactometer behavioral assays

Mosquitoes that were blood fed for behavior were fed on warmed defibrinated sheep's blood (Hemostat Laboratories DSB100) through a parafilm membrane. Animals were sorted by eye to determine if they were blood fed by looking at engorgement of the abdomen, and non-fed animals were not used in subsequent behavior assays. Blood-fed animals were returned to the same rearing conditions as their non-blood-fed counterparts, and behavior was assayed ~48 hr after they had consumed a blood meal. Quattroport olfactometer experiments were conducted as previously

described (Basrur et al., 2020) with modifications to enable the delivery of monomolecular odorants. Details of fabrication and operation of the Quattroport are available at https://github.com/VosshallLab/Basrur_Vosshall2020.

All behavioral experiments were carried out in an environmental room set to 25 – 28°C, 70-80% relative humidity, and were carried out between ZT3-14. Trials with identical stimuli were interspersed throughout experimental days to control for circadian rhythm fluctuations. Mosquitoes in the Quattroport began each trial in a start canister, flew through a flight tube, and, if they were attracted to a stimulus, into a trap proximal to the stimulus. Filtered air, with or without additional CO₂, was pumped through the stimulus box and into the flying tube. Mosquitoes were prevented from contacting the stimulus by a mesh barrier, and adjustable gates between each chamber allow the experimenter to count the number of mosquitoes in each compartment.

The night before each experiment, approximately 20 7-22 days old mosquitoes were cold-anesthetized and placed in each canister. They were sugar-starved with access to water overnight. The next day, at the start of each trial a set of 4 canisters was attached to the Quattroport and mosquitoes were given 10 min to acclimate with filtered air flowing at the 25 mm setting on the flowmeter throughout the assay (see https://github.com/VosshallLab/Basrur_Vosshall2020 for details on filtered air and CO₂ flowmeter settings). They were then exposed to the stimulus with or without added CO₂ flowing at the 50 mm flowmeter setting for 30 sec, after which they were given access to the flying tube and attraction trap for 5 min in the continued presence of filtered air with or without added CO₂ and the stimulus. At the end of the trial, the gates enclosing the start canister and the attraction trap were closed, and mosquitoes that remained in the start canister, entered the flying tube, or flew into the attraction trap were counted.

For live human host-seeking assays, a single human volunteer placed their forearm in the stimulus box. For the delivery of lactic acid and triethyl amine, the odorant solutions were delivered using 35 mm Petri dishes (Fisher Scientific #08-757-100A) placed within the air stream in the stimulus box. The odorants used in this assay were L-(+)-Lactic acid solution, 88-92% (PubChem CID: 107689, Sigma-Aldrich #27714-500mL) and Triethyl amine (PubChem CID: 1146, Sigma T0886) diluted to 1% v/v in MilliQ ultrapure water. In cases where both odorants were used as a stimulus, care was taken not to mix the liquids when they were placed in the stimulus box. Different volumes of each liquid odorant were delivered to generate the dose response curves in **Figure 6F,H** and **Data File 1**. Depending on the experiment, between 1 μ L and 2 mL of liquid odorant was used. When less than 1 mL of liquid was used, the lid of the Petri dish was inverted, and the liquid odorant was pipetted onto the center of the inverted lid. The inverted lid was then placed on the bottom of the Petri dish, which elevated it slightly, placing the odorants in the path of the air stream that ran through the stimulus box. When 1 mL or greater of liquid was used, the liquid odorant was pipetted into the base of the Petri dish, which was set atop the inverted lid of the Petri dish to elevate it. If the volume of odor was 1 mL or greater for either odorant, they were delivered using 2 separate Petri dishes to avoid possible mixing. If the volume of each liquid was less than 1 mL, both liquid odorants were delivered as separated drops of liquid on a single inverted Petri dish lid, and the drops remained separate because of surface tension. When neither lactic acid nor triethyl amine was delivered, an empty Petri dish was placed inside the stimulus box to serve as a control for any odors that might be emitted by the dish itself and/or any perturbations in air flow introduced by the presence of the Petri dish lid and base

SUPPLEMENTARY FIGURES

Figure S1 Organization of *Ae. aegypti* antennal lobe glomeruli (Related to Figure 1)

(A-B) Maximum-intensity projections of confocal Z-stacks of a brain after anterograde dye fill of a single ipsilateral antenna (A) or ipsilateral maxillary palp (B) using a dextran-conjugated fluorophore (green) with immunofluorescent labeling of Brp (synaptic marker, magenta). (C) Approximate number of antennal lobe glomeruli per brain hemisphere innervated by the indicated sensory structure, derived from quantification of the left antennal lobe in 12 brains presented in Figure 1I-J, S2-S5. (D-K) 3-D reconstruction of a single left antennal lobe with 61 (D-G) or 66 (H-K) glomeruli shown at 4 different angles. Glomeruli are colored according to innervation by the indicated sensory appendage. Panel (G) is reprinted in Figure 5B. Scale bars: 50 μ m (A-B), 20 μ m (D-K). Orientation: d=dorsal, m=medial, p=posterior.

Figure S2. Projections of *Orco-QF2*-expressing neurons in the antennal lobe (Related to Figure 1)

(A) *Orco* locus with exons (grey boxes), introns (grey lines) and CRISPR-Cas9 gRNA site (arrowhead) used to insert T2A-QF2 (light blue). (B) Quantification of the number of glomeruli that are GFP positive (green), GFP negative (magenta), and total number of glomeruli (black). Analysis based on brains in (C-D) and Figure 11,J. (C) Maximum-intensity projections of confocal Z-stacks of left antennal lobes from two different brains of the indicated genotype with immunofluorescent labeling of GFP (green) and Brp (synaptic marker, magenta) (top) and 2-D representation of the boundary of each glomerulus that is GFP positive and GFP negative (bottom). (D) Single confocal sections taken from the maximum-intensity projection confocal Z-stack of the left antennal lobe shown in Figure 11 with immunofluorescent labeling of GFP (green) and Brp (synaptic marker, magenta). A single plane is shown every 5 μ m in Z to capture each glomerulus. Scale bar (C-D): 50 μ m. Orientation: d=dorsal, m=medial.

Figure S3. Projections of *Ir25a-QF2*-expressing neurons in the antennal lobe (Related to Figure 1)

(A) *Ir25a* locus with exons (grey boxes), introns (grey lines) and CRISPR-Cas9 gRNA site (arrowhead) used to insert T2A-QF2 (light blue). (B) Quantification of the number of glomeruli that are GFP positive (green), GFP negative (magenta), and total number of glomeruli (black). Analysis based on brains in (C-D) and Figure 11J. (C) Maximum-intensity projections of confocal Z-stacks of left antennal lobes from two different brains of the indicated genotype with immunofluorescent labeling of GFP (green) and Brp (synaptic marker, magenta) (top) and 2-D representation of the boundary of each glomerulus that is GFP positive and GFP negative (bottom)

(D) Single confocal sections taken from the maximum-intensity projection confocal Z-stack of the left antennal lobe shown in Figure 11J with immunofluorescent labeling of GFP (green) and Brp (synaptic marker, magenta). A single plane is shown every 5 μm in Z to capture each glomerulus. Scale bar (C-D): 50 μm. Orientation: d=dorsal, m=medial.

Figure S4. Projections of *Ir8a*-QF2-expressing neurons in the antennal lobe (Related to Figure 1)

(A) *Ir8a* locus with exons (grey boxes), introns (grey lines) and CRISPR-Cas9 gRNA site (arrowhead) used to insert T2A-QF2 (light blue). (B) Quantification of the number of glomeruli that are GFP positive (green), GFP negative (magenta), and total number of glomeruli (black). Analysis based on brains in (C-D) and Figure 11,J. (C) Maximum-intensity projections of confocal Z-stacks of left antennal lobes from two different brains of the indicated genotype with immunofluorescent labeling of GFP (green) and Brp (synaptic marker, magenta) (top) and 2-D representation of the boundary of each glomerulus that is GFP positive and GFP negative (bottom). (D) Single confocal sections taken from the maximum-intensity projection confocal Z-stack of the left antennal lobe shown in Figure 11 with immunofluorescent labeling of GFP (green) and Brp (synaptic marker, magenta). A single plane is shown every 5 μ m in Z to capture each glomerulus. Scale bar (C-D): 50 μ m. Orientation: d=dorsal, m=medial.

Figure S5. Projections of *Ir76b*-QF2-expressing neurons in the antennal lobe (Related to Figure 1)

(A) *Ir76b* locus with exons (grey boxes), introns (grey lines) and CRISPR-Cas9 gRNA site (arrowhead) used to insert T2A-QF2 (light blue). (B) Quantification of the number of glomeruli that are GFP positive (green), GFP negative (magenta), and total number of glomeruli (black). Analysis based on brains in (C-D) and Figure 11,J. (C) Maximum-intensity projections of confocal Z-stacks of left antennal lobes from two different brains of the indicated genotype with immunofluorescent labeling of GFP (green) and Brp (synaptic marker, magenta) (top) and 2-D representation of the boundary of each glomerulus that is GFP positive and GFP negative (bottom). (D) Single confocal sections taken from the maximum-intensity projection confocal Z-stack of the left antennal lobe shown in Figure 11 with immunofluorescent labeling of GFP (green) and Brp (synaptic marker, magenta). A single plane is shown every 5 μ m in Z to capture each glomerulus. Scale bar (C-D): 50 μ m. Orientation: d=dorsal, m=medial.

Figure S6. Projections of *GR3-QF2*-expressing neurons in the antennal lobe (Related to Figure 1)

(A) *Gr3* locus with exons (grey boxes), introns (grey lines) and CRISPR-Cas9 gRNA site (arrowhead) used to insert T2A-QF2 (light blue). (B) Maximum-intensity projection confocal Z-stack through the medial antennal lobes of 4 brains with immunofluorescent labeling of GFP (green) and Brp (synaptic marker, magenta). Scale bar: 25 μm. Orientation: d=dorsal, m=medial.

Figure S7. Specificity of Split-QF2 reagents (Related to Figure 2)

(A-B) Maximum-intensity projections of confocal Z-stacks of antennae (A) and maxillary palps (B) of the indicated genotypes showing intrinsic dTomato fluorescence and stained with the nuclear dye TO-PRO-3, with transmitted light overlay. (C) Maximum-intensity projections of confocal Z-stacks of left antennal lobes from two different brains of the indicated genotype with immunofluorescent labeling of dTomato (green) and Brp (synaptic marker, magenta) (top) and 2-D representation of the boundary of each glomerulus that is GFP positive and GFP negative (bottom). Scale bars: 50 μ m. Orientation (C,D): d=dorsal, m=medial.

Figure S9. Quantification of antennal and maxillary palp cell populations (Related to Figure 3 and Figure 4)

(A-H) Workflow for cell quantification. Schematic of antennal region imaged on a confocal microscope (A) and image of maxillary palp with imaged area indicated with the yellow square (B). Whole-mount maxillary palp RNA *in situ*, yellow region from (C). Cells are manually marked independently as *Orco*+, *Or49*+, or *Or8*+ (red inset from B) using FIJI Cell Counter (D-F) and markers from each channel are merged (G). Cells with markers 1 and 2 are then scored as *Orco*+*Or49*+ with marker 4, and cells with markers 1 and 3 are then scored as *Orco*+*Or8*+ with marker 5 (H). Counts from each marker for each image are exported into Excel and R for further analysis. (I-N) Total cell counts from whole mount maxillary palp RNA *in situ* in Figure 4. Mean with range, n=5. (O) RNA *in situ* of whole-mount maxillary palp with the indicated probe and cell identity schematic. (P-Q) Whole-mount maxillary palp immunostaining showing *Ir25a* expression in "small" and "large" cells (P) and *Gr3* expression in "large" cells and *Orco* protein in "small" cells (Q). Scale bars: 25 μm except (C): 50 μm. Orientation (B, P): distal right.