Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Synonymization of the male-based ant genus *Phaulomyrma* (Hymenoptera, Formicidae) with *Leptanilla* based upon Bayesian total-evidence phylogenetic inference

Zachary H. Griebenow

Abstract. Although molecular data have proven indispensable in confidently resolving the phylogeny of many clades across the tree of life, these data may be inaccessible for certain taxa. The resolution of taxonomy in the ant subfamily Leptanillinae is made problematic by the absence of DNA sequence data for leptanilline taxa that are known only from male specimens, including the monotypic genus *Phaulomyrma* Wheeler & Wheeler. Focusing upon the considerable diversity of undescribed male leptanilline morphospecies, the phylogeny of 35 putative morphospecies sampled from across the Leptanillinae, plus an outgroup, is inferred from 11 nuclear loci and 41 discrete male morphological characters using a Bayesian total-evidence framework, with *Phaulomyrma* represented by morphological data only. Based upon the results of this analysis *Phaulomyrma* is synonymized with *Leptanilla* Emery, and male-based diagnoses for *Leptanilla* that are grounded in phylogeny are provided, under both broad and narrow circumscriptions of that genus. This demonstrates the potential utility of a total-evidence approach in inferring the phylogeny of rare extant taxa for which molecular data are unavailable and begins a long-overdue systematic revision of the Leptanillinae that is focused on male material.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Introduction

Over the past three decades, DNA sequences have provided great insight into phylogenetic relationships across the Metazoa, including the insects (Kjer *et al.* 2018). The application of maximum-likelihood (ML) and Bayesian statistical methods to analysis of genetic data has robustly resolved many problems that were intractable when using morphological data alone (e.g. Niehuis *et al.* 2012; Wipfler *et al.* 2019). However, DNA sequences may be unavailable for some taxa, necessitating the integration of morphological and molecular data under the same inferential framework. Fossils are the most obvious example of this: these are valuable for calibration of phylogenies in absolute time under a Bayesian approach, preferably with their topological position being inferred from the data (Ronquist *et al.* 2012; O'Reilly *et al.* 2015; Bapst *et al.* 2016; Matzke & Wright 2016). Although the inclusion of fossils for the purposes of “tip-dating” has received the bulk of attention in Bayesian total-evidence phylogenetic inference, the lack of molecular data may afflict rare extant taxa as well (Sánchez *et al.* 2016; Robertson & Moore 2016). This is problematic if the affinities of these taxa are not immediately clear from morphology alone.

The ant subfamily Leptanillinae (Hymenoptera: Formicidae) is an apt test case for methods to resolve this problem. A group of small, hypogaeic ants largely restricted to the Old World tropics and subtropics, the Leptanillinae are understood to be one of the earliest-diverging lineages in the ant crown-group (Rabeling *et al.* 2008; Kück *et al.* 2011; Borowiec *et al.* 2019; Boudinot *et al.* submitted). Three out of eight described genera are known from both workers and males: *Opamyrma* Yamane, Bui & Eguchi, 2008 (Yamada *et al.* 2020), *Protanilla* Taylor, 1990 (Griebenow, in press), and *Leptanilla* Emery, 1870 (e.g. Ogata *et al.* 1995). Males of *Anomalomyrma* Taylor, 1990 are unknown. Four leptanilline genera—*Scyphodon* Brues, 1925;

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Phaulomyrma Wheeler & Wheeler, 1930; *Noonilla* Petersen, 1968; and *Yavnella* Kugler, 1986—have been described solely from males, as have many species of *Leptanilla* (cf. Bolton 1990). Recent molecular data indicate that the type species of *Yavnella* and a specimen provisionally assigned to *Phaulomyrma* are nested within a clade of putative *Leptanilla* morphospecies (Borowiec *et al.* 2019). Moreover, although *Scyphodon anomalum* Brues, 1925 and *Noonilla copiosa* Petersen, 1968 exhibit bizarre autapomorphies such as hypertrophied mandibles (Brues 1925) and a ventromedian genital “trigger” (Petersen 1968), respectively, these ants are otherwise similar to males attributed to *Leptanilla* (Boudinot 2015).

This indicates a need for a systematic revision of the Leptanillinae, but almost all published taxonomic studies of the group have been descriptive without recourse to molecular phylogeny, with the exceptions being revisions to our concept of the subfamily. Multi-locus DNA datasets demonstrated that the enigmatic Afrotropical genus *Apomyrma* Gotwald, Brown & Lévieux, 1971 is closely related to the Amblyoponinae rather than the Leptanillinae (Brady *et al.* 2006; Moreau *et al.* 2006), and that the superficially similar Asian genus *Opamyрма* is in fact sister to the remaining Leptanillinae (Ward & Fisher 2016). None of these studies focused upon the Leptanillinae or the internal phylogeny of this clade. Such a study must confront two challenges: first, the lack of DNA sequences for certain critical taxa across the Leptanillinae (e.g., *Scyphodon*), which hampers any attempt to confidently resolve relationships among these; second, the definition of genera based only upon males, which prevents an integrated phylogenetic classification of the Leptanillinae, since phenotypes of only one sex are considered.

The dissociation of leptanilline castes results from collecting bias. Subterranean workers have been largely collected with *lavage de terre* methodology (López *et al.* 1994; Wong and Guénard 2016), Winkler trapping (Belshaw & Bolton 1994; Leong *et al.* 2018), and subterranean pitfall

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

traps (Wong & Guénard 2016; Man *et al.* 2017); whereas male leptanillines are typically collected by sweeping foliage or by deploying Malaise or pan traps (Robertson 2000). None of these methods are likely to collect males in association with workers, nor is the queen caste often collected in association with conspecifics. Contrasting with the alate condition observed in most ants, queens described from the tribe Leptanillini are completely wingless and blind (Emery 1870; Kutter 1948; Masuko 1990; López *et al.* 1994; Ogata *et al.* 1995), meaning that these are no more likely to be collected than corresponding workers. Queens belonging to other leptanilline lineages (*Opamyrma* and the Anomalomyrmini) are alate so far as is known (Bolton 1990; Baroni Urbani & de Andrade 2008; Borowiec *et al.* 2011; Chen *et al.* 2017; Hsu *et al.* 2017; Man *et al.* 2017), save for an apparent record of queen polyphenism in an undescribed *Protanilla* (Billen *et al.* 2013), but are infrequently collected.

Therefore, the bulk of known leptanilline diversity, most of it undescribed, is represented by exclusively male material. In some cases, molecular data are inaccessible for male morphotaxa due to paucity of suitably recent specimens, obliging a total-evidence approach to infer the phylogeny of these lineages. This study uses such an approach to resolve the position of the male-based species *Phaulomyrma javana* Wheeler & Wheeler, 1930, the sole species included in this genus. Here, the phylogeny of the Leptanillinae is inferred jointly from 10 protein-coding genes, 28S rDNA, and 41 discrete male morphological characters under a Bayesian statistical framework. This is the first combined-evidence Bayesian analysis to include the Leptanillinae and is novel among studies of ant phylogeny in its inclusion of exclusively male morphological characters (Barden *et al.* [2017] used both worker and male morphology in their Bayesian total-evidence inference). Despite the absence of nucleotide sequences for *P. javana* a Bayesian total-evidence approach facilitates the inclusion of this terminal and its confident phylogenetic

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

placement. Based upon the results of these joint molecular and morphological phylogenetic analyses, a revised male-based definition of *Leptanilla* is provided, and *Phaulomyrma* is synonymized with that genus.

Materials and Methods

Taxon Sampling

Thirty-five terminals were included in total (Tables 1-2). Discrete morphological data were scored for those 33 terminals for which male material was known. *Anomalomyrma boltoni* Borowiec, Schulz, Alpert & Bañar, 2011 and *Leptanilla revelierii* Emery, 1870 were represented in this study by workers alone. The latter was included on account of its status as the type species of that genus: regardless of future systematic revision to the Leptanillinae, the concept of the genus *Leptanilla* will not exclude this species. DNA sequences for the outgroup *M. heureka* were obtained from a worker ant, as published in Borowiec *et al.* (2019). Most putative morphospecies were represented by singletons (Table 1), but phenotypic variation within those morphospecies for which material was abundant (e.g., *Leptanilla zhg-my02*) is minimal, and so gives no reason to suspect heterospecificity among the specimens referred to these morphospecies.

Representatives of all male-based genera were included in total-evidence analyses, except for *Scyphodon*. These include both *Yavnella argamani* Kugler, 1986 and *Yavnella* cf. *indica*, along with two undescribed *Yavnella* morphospecies from Bhutan and Thailand, respectively; *Phaulomyrma javana*; and two morphospecies of *Noonilla* identified as such according to the definition given by Petersen (1968). *Leptanilla* TH02-6 and -08, along with *Phaulomyrma* MM01, were placed in those genera by Borowiec *et al.* (2019) and/or Boudinot (2015) but are here identified as *Yavnella* (Table 1) according to the definition of Kugler (1986).

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Material is deposited in the following repositories: the Bohart Museum of Entomology, University of California, Davis, CA, USA (UCDC); the California Academy of Sciences, San Francisco, CA, USA (CASC); the California State Collection of Arthropods, Sacramento, CA, USA (CSCA); the Lund Museum of Zoology, Lund, Sweden (MZLU); and the Australian National Insect Collection, Canberra, Australia (ANIC).

Molecular Dataset

Total-evidence phylogenetic inference was based upon 11 nuclear loci: *28S ribosomal DNA* (28S), *abdominal-A (abdA)*, *arginine kinase (argK)*, *antennapedia (Antp)*, *elongation factor 1-alpha F2 copy (EF1αF2)*, *long wavelength rhodopsin (LW Rh)*, *NaK ATPase (NaK)*, *DNA polymerase delta (POLDI)*, *topoisomerase I (TopI)*, *ultrabithorax (Ubx)*, and *wingless (Wg)*. I derived these “legacy loci” for 19 terminals from the alignment of Borowiec *et al.* (2019) (doi: 10.5281/zenodo.2549806) but expanded to include autapomorphic indels and introns, and constituting 11,090 bp. Legacy loci for *Leptanilla* GR03 were derived from Ward & Sumnicht (2012). For further detail on the protocols for the extraction and amplification of these genetic data, refer to Ward *et al.* (2010) and Ward & Fisher (2016). I added 14 terminals to this “legacy-locus” intron-inclusive dataset by retrieving orthologous loci from phylogenomic data acquired with the ultra-conserved element (UCE) probe set *hym-v2* (Branstetter *et al.* 2017). *P. javana* was the only terminal for which molecular data were not obtained: this species is known only from two slide-mounted syntypes collected in 1907.

DNA was extracted non-destructively using a DNeasy Blood and Tissue Kit (Qiagen Inc., Valencia, CA) according to manufacturer instructions. DNA was quantified for each sample with a Qubit 2.0 fluorometer (Life Technologies Inc., Carlsbad, CA). Phylogenomic data were obtained from these taxa using the *hym-v2* probe set, with libraries being prepared and target loci

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

enriched using the protocol of Branstetter *et al.* (2017). Enrichment success and size-adjusted DNA concentrations of pools were assessed using the SYBR FAST qPCR kit (Kapa Biosystems, Wilmington, MA) and all pools were combined into an equimolar final pool. The contents of this final pool were sequenced by an Illumina HiSeq 2500 at the University of Utah's High Throughput Genomics Facility or an Illumina HiSeq 4000 at Novogene, Sacramento, CA. The FASTQ output was demultiplexed and cleansed of adapter contamination and low-quality reads using *illumiprocessor* (Faircloth 2013) in the PHYLUCE package. Raw reads were then assembled with *trinity* v. 2013-02-25 (Grabherr *et al.* 2011) or SPAdes v. 3.12.0 (Bankevich *et al.* 2013). The possibility of genetic contamination and/or mis-assembly in the UCE samples was tested by inferring a phylogeny from a concatenated UCE alignment, unpartitioned, using IQ-Tree 1.6.10 (Nguyen *et al.* 2015) on the CIPRES Science Gateway (v. 3.3) (Miller *et al.* 2010) with the GTR+G model of substitution for 1,000 ultrafast bootstrap replicates (Hoang *et al.* 2018); this phylogeny was plausible given preliminary hypotheses, providing no positive evidence of sequence contamination or mis-assembly. Summary statistics for these UCE assemblies were computed using *statswrapper.sh* in *BBMap* (Bushnell 2014), and provided in Supplemental Table 1.

In the cases of the 14 terminals not included in Ward & Sumnicht (2012) or Borowiec *et al.* (2019) for which molecular data could be obtained, legacy loci orthologous with those used by Borowiec *et al.* (2019) were then recovered from genome-scale data as follows using PHYLUCE (Faircloth 2016). I derived sequences representing each locus for *Leptanilla* GR02 from the alignment ANT-exon-sequences-40-taxa-reduced.fasta published by Branstetter *et al.* (2017), given the comparative completeness of the matrix for that species, and its phylogenetic position nested well within the Leptanillinae. These sequences were then used analogously to probes.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Species-specific contig assemblies were obtained using *phyluce_assembly_match_contigs_to_probes.py* (min_coverage = 50, min_identity = 85), a list of legacy loci shared across all taxa was generated using *phyluce_assembly_get_match_counts.py*, and separate FASTA files for each locus were created using these outputs. Sequences were aligned separately by locus using MAFFT (Katoh *et al.* 2009) implemented with the command *phyluce_assembly_seqcap_align.py*, and these sequences were then trimmed with Gblocks (Castresana 2000) as implemented by the wrapper script *phyluce_assembly_get_gblocks_trimmed_alignment_from_untrimmed.py* (settings: b1 = 0.5, b2 = 0.5, b3 = 12, b4 = 7). Alignment statistics for the output FASTA files were calculated with *phyluce_align_get_align_summary_data.py*. Finally, a matrix that was 80% complete with respect to locus coverage was generated using the script *phyluce_align_get_only_loci_with_min_taxa.py*. This contained 7 out of the 10 protein-coding loci that I attempted to recover using the exon-based bioinformatic protocol of Branstetter *et al.* (2017), in addition to 28S rDNA. Legacy loci recovered from UCE assemblies often included non-coding sequences adjacent to the regions included in Borowiec *et al.* (2019), which were trimmed manually in AliView. In whichever cases those loci had been recovered, sequences for the taxa represented only in the dataset of Borowiec *et al.* (2019) were then aligned with the recovered legacy loci using the online MAFFT interface (Katoh *et al.* 2019) with default settings. In cases where legacy loci were not successfully recovered or were incomplete relative to preexisting Sanger-derived sequences, these loci were derived from the datasets of Borowiec *et al.* (2019) or, in the case of *Leptanilla* GR03, Ward & Sumnicht (2012). These data were concatenated with UCE-derived sequences across all FASTA files, inasmuch all sequences for each morphospecies were derived from the same specimen; and all loci were concatenated to

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

produce a final alignment, which was 9,351 bp in length. Further summary statistics for this final alignment are provided in Table 1 and Supplemental Table 2. Alignment was unambiguous once all loci were brought into their respective reading frames. GenBank accession numbers for all loci used in this study are provided in Table 2.

Those terminals for which loci were obtained using the 11 nuclear loci from *Leptanilla* GR02 as “probes” according to the modified PHYLUCE protocol cited above (“Molecular Dataset”) (Faircloth 2016), with 80% locus coverage implemented in *phyluce_align_get_only_loci_with_min_taxa.py*, exhibit low coverage relative to those that were sequenced prior to this study (Table 1). Therefore, a 9,062-bp legacy-locus alignment was created that includes only those data published prior to this study (Ward & Sumnicht 2012; Borowiec *et al.* 2019), with 20 terminals. These sequences can be used to test the possibility that missing data would have an appreciable effect on phylogenetic inference phylogenetic analyses.

Morphological Dataset

Forty-one discrete binary morphological characters were coded for all 33 morphospecies known from males. All these specimens were examined with a Leica MZ75 compound microscope or by reference to images on AntWeb, except for the male of *M. heureka* and *O. hungvuong*, in the cases of which observations were derived from Boudinot (2015: Figs. 11-12) and Yamada *et al.* (2020: Figs. 11-13), respectively, or from the textual descriptions by those authors. I imaged specimens when necessary using a JVC KY-F75 digital camera and compiled color photographs from these with the Syncroscopy AutoMontage Program. Scanning electron microscopy was undertaken using a Hitachi TM4000 tabletop microscope. Morphological terminology follows the Hymenoptera Anatomy Ontology (Yoder *et al.* 2010), with some exceptions being derived from Bolton (2003) and Boudinot (2018). The character coding scheme was binary and non-

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

additive (Pleijel 1995). Missing data were scored as ‘?’. Autapomorphic characters were included. Numerical scores for all morphological characters are presented in the Supplemental Table 3.

Non-additive binary coding has been criticized for its susceptibility to redundancy (Strong & Lipscomb 1999), stipulation of compound characters, and the inadvertent conflation of morphological absences that are not hierarchically equivalent (Brazeau 2011). These problems largely result from careless character delimitation. I compensated for these potential flaws by defining and using only characters that do not logically depend upon other characters. Definitions of morphological character states are provided in the Appendix.

Phylogenetic Analyses

For the two legacy-locus molecular datasets, the partitioning scheme was inferred with PartitionFinder2 v. 2.1.1 (Guindon *et al.* 2010; Lanfear *et al.* 2012, 2017) on the CIPRES Science Gateway, with subsets being asserted *a priori* according to locus and codon position. Introns were included. Models with I+G extensions were excluded from consideration due to undesirable behavior in a model-based framework (Yang 1996). As an alternative *ad hoc* partitioning scheme for the 9,351-bp alignment, I respectively partitioned all exonic loci so that 1st-2nd codon positions were placed in their own partition separate from the 3rd, and modeled nucleotide substitution in all partitions under GTR+G. Using AMAS (Borowiec 2016), the full 9,351-bp and 9,062-bp molecular alignments were respectively split according to partition scheme(s) for partitioned Bayesian total-evidence inference.

In total-evidence and morphology-only Bayesian phylogenetic analyses, the Mkv model (Lewis 2001) was used to model substitution of morphological character states, albeit with stationary

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

frequencies of character states treated as free parameters (Felsenstein 1981) in order to accommodate asymmetry in character state frequencies. Variation in evolutionary rate among characters was accommodated by drawing rates from a gamma-distributed prior probability distribution (+G), approximated with 8 discrete categories k .

All phylogenetic analyses were performed in a Bayesian statistical framework using RevBayes v. 1.0.11 (Höhna *et al.* 2017) compiled on Ubuntu Linux v. 13.04. The following phylogenetic analyses were implemented: one using the 41-character male morphological dataset alone; one using the 9,351-bp molecular dataset alone; two total-evidence analyses using the 9,351-bp molecular alignment, respectively with algorithmic or *ad hoc* partitioning schemes as described above; and a total-evidence analysis using the 9,062-bp molecular alignment, partitioned algorithmically as described above with PartitionFinder2. Each analysis consisted of four independent Markov chain Monte Carlo (MCMC) chains, each run for 50,000 generations. Trees were sampled every 10 generations, with the first 25% of the run being discarded as burn-in. MCMCs with respect to all continuous parameters were considered converged if the effective sample sizes as given in Tracer v. 1.7.1 (Rambaut *et al.* 2018) were ≥ 200 , with sufficiency of MCMC mixing across posterior probability landscapes being qualitatively assessed using traces of the respective log-likelihoods of each parameter across the course of the analysis. Maximum *a posteriori* trees were compiled from this sample of each run, with node support expressed as Bayesian posterior probability (BPP).

Data Availability and Nomenclature

All nucleotide and morphological data along with PartitionFinder2 configuration files, RevBayes scripts, and output of all phylogenetic analyses, are available at the Dryad Digital Repository

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

(doi:10.25338/B8GP7C). Sequence Read Archives (SRAs) of raw UCE reads, and UCE assemblies, are publicly available on NCBI (Table 2).

This article has been registered in Zoobank (www.zoobank.org). The LSID number is 5F3BECF6-3715-47B3-8D0F-DE7D66E1DA0A.

Results

Bayesian total-evidence inference of leptanilline phylogeny using the 9,351-bp legacy-locus dataset under the two partitioning schemes resulted in similar topologies, with none of the differences affecting the composition or interrelationship of major clades. All Bayesian total-evidence phylogenies inferred under the *ad hoc* partitioning schemes are provided on Dryad. Most nodes in these phylogenies were supported with $BPP \geq 0.95$. Those nodes supported with $BPP \leq 0.95$ were scattered and shallow (Fig. 1), meaning that the interrelationships among all major leptanilline clades are well-resolved. Although the sampling of the Leptanillinae was more extensive than that of Borowiec *et al.* (2019), our inferences were largely congruent. Bayesian total-evidence inference from the 9,062-bp alignment also drew a consistent conclusion (Fig. 2), indicating that the taxonomically biased distribution of missing data in the 9,351-bp legacy-locus dataset does not have an appreciable effect on the backbone of inferred leptanilline phylogeny. Phylogenetic inference from the 9,351-bp alignment alone, and therefore excluding *P. javana*, fully corroborates the conclusions of total-evidence Bayesian phylogenetic inference with high Bayesian posterior probabilities overall, while inference from the morphological dataset alone was insufficient to resolve the phylogeny of the Leptanillinae (see Dryad). All discussion from here on refers to the phylogeny inferred under the partitioning scheme derived with PartitionFinder2 (Fig. 1) for the 9,351-bp molecular alignment, unless otherwise noted.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

269 The clade corresponding to the tribe Anomalomyrmini (labeled as *Protanilla sensu lato* in Fig.
270 1) is recovered with maximal support (BPP = 1), with *A. boltoni* sister to all sampled *Protanilla*
271 save *Protanilla* TH03—thus rendering *Protanilla* paraphyletic—well-supported (BPP = 0.9577).
272 This same topology was recovered by Bayesian total-evidence analysis from the 9,062-bp
273 alignment, with higher support (BPP = 0.9947) (Fig. 2). Borowiec *et al.* (2019) recovered *A.*
274 *boltoni* as sister to *Protanilla* TH03 with weak support irrespective of statistical framework,
275 albeit with more extensive sampling within the Anomalomyrmini, as did total-evidence inference
276 from the 9,351-bp dataset under the *ad hoc* partitioning scheme (BPP = 0.6535). However, the
277 internal topology of the Anomalomyrmini does not have any bearing upon the status of
278 *Phaulomyrma* relative to other male-based leptanilline genera, nor its status relative to
279 *Leptanilla*.

280 *Noonilla*, *Yavnella argamani* and *Yavnella cf. indica*, *Leptanilla revelierii*, and *Phaulomyrma*
281 *javana* were firmly recovered within a clade corresponding to the Leptanillini (BPP = 1). As in
282 Borowiec *et al.* (2019) the Leptanillini bifurcate robustly, with *Y. argamani* (and *Yavnella cf.*
283 *indica*, which was not included in Borowiec *et al.* [2019]) recovered in a clade otherwise without
284 described representatives, which is hereinafter designated *Yavnella sensu lato* (BPP = 1).
285 Although morphologically diverse (Fig. 3), the male morphospecies that comprise the sister-
286 group to *Yavnella s. l.* are distinguished from that clade by 1) clypeus with a medial axis no
287 longer than the diameter of the torulus, where the epistomal sulcus is distinct; and 2) pronotum
288 and mesoscutum that are not extended posteriorly in profile view. Since *L. revelierii* is recovered
289 within this clade, it is hereinafter referred to as *Leptanilla sensu lato* (BPP = 0.9964). *Leptanilla*
290 *s. l.* bifurcates into two well-supported clades: one is broadly Eurasian and Australian in its
291 representation (with a single Afrotropical representative), including *L. revelierii* and *P. javana*

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

(BPP = 0.9531); the other is Indo-Malayan, and includes *Noonilla* (BPP = 0.9839) (Figs. 1, 4). Since *L. revelierii* is included within the Eurasian-Australian clade, this clade is hereinafter referred to as *Leptanilla sensu stricto*. The two circumscriptions of the name *Leptanilla* presented here are supported by male morphology (see Discussion).

Noonilla (BPP = 0.9999) is sister to a clade represented by highly distinctive male morphospecies, recovered with maximal support (BPP = 1) (Figs. 1, 4), that are immediately recognizable by bizarre metasomal processes (heretofore hypothesized to be extensions of the gonocoxae *sensu* Boudinot [2018] [Boudinot 2015: Fig. 10D]) and a comb-like row of robust bristles on the protibia (Fig. 5) in combination with a putatively grasping profemur. These morphospecies remain undescribed. Boudinot (2015: p. 33) adduced the grasping profemur of *Noonilla* as an autapomorphy of that genus, which could justify terming the undescribed clade as *Noonilla*_cf; but this profemoral condition is more widespread across male Leptanillini than Boudinot (2015) was aware, and better sampling is required to infer whether the grasping profemur is a synapomorphy of *Noonilla* and this undescribed clade. I therefore provisionally refer to said clade as the “Bornean morphospecies-group”: while present sampling is too sparse to judge whether this clade is precinctive to Borneo, available material exclusively originates on that island. Of the 9 terminals recovered in the Indo-Malayan subclade, only *Leptanilla* TH01 was included in Borowiec *et al.* (2019) or in the 9,062-bp legacy-locus alignment. The rather disparate morphospecies *Leptanilla* TH01 and *Leptanilla* zhg-th01 are recovered as a clade with high support (BPP = 0.9961), and this clade is in turn sister to *Noonilla* + Bornean morphospecies-group (Figs. 1, 4). *Leptanilla* zhg-th01 is unique among the Leptanillinae in possessing a recurved mesoscutellar horn (Fig. 6B).

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

The support values of internal nodes within *Leptanilla s. str.* are generally poor under Bayesian total-evidence inference from the 9,351-bp legacy-locus alignment, with the placement of *Leptanilla* ZA01 and *Leptanilla* zhg-bt01 differing according to partitioning scheme. The position of *P. javana* cannot be confidently resolved within this clade, but the basalmost node of *Leptanilla s. str.* is well-supported, whether inferred under an algorithmic (BPP = 0.9531) or *ad hoc* (BPP = 0.9585) partitioning scheme. While the internal phylogeny of *Leptanilla s. str.* cannot be resolved with Bayesian total-evidence inference, the monophyly of this clade is probable under the model and partitioning schemes used. The topology of *Leptanilla s. str.* is likely subject to strong stochastic error due to the inclusion of *P. javana*, for which molecular data are entirely absent. This is supported by Bayesian phylogenetic inference from molecular data alone, which with only one exception recovers the internal phylogeny of *Leptanilla s. str.* with $BPP \geq 0.95$ (see Dryad).

Bayesian total-evidence inference from the 9,062-bp alignment (which does not include *Leptanilla revelierii*, zhg-au02 or zhg-bt01) gives mediocre support to *Leptanilla s. str.* (BPP = 0.9237) inclusive of *P. javana*, but provides a phylogeny consistent with the results of other phylogenetic analyses (Fig. 2). The recovery of *P. javana* within *Leptanilla s. str.* is therefore supported by Bayesian total-evidence inference. Qualitatively, male morphological characters support *Leptanilla s. str.* (see Discussion).

P. javana and the taxon dubbed *Phaulomyrma* MM01 by Boudinot (2015) and Borowiec *et al.* (2019) were recovered distant from one another in the leptanilline phylogeny (Figs. 1-2, 4). Total-evidence phylogenetic inference recovered the latter terminal within *Yavnella s. l.*, indicating that it was incorrectly assigned to *Phaulomyrma* by these authorities, corroborating morphological evidence (see Discussion). An undescribed male morphospecies referred to as

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Phaulomyrma by Boudinot (2015: Fig. 4F) was not sequenced in this study but also conforms morphologically to *Yavnella s. l.*, and so likewise was incorrectly identified as *Phaulomyrma*. Conversely, *P. javana* is here recovered within *Leptanilla s. l.*, and moreover within *Leptanilla s. str.* (BPP = 0.9531).

Discussion

Delimitation of Subclades in Leptanillinae using Male Morphology

Male morphological characters corroborate inferred phylogeny at nodes of variable depth. *O. hungvuong* and the four male representatives of the Anomalomyrmini included in the present study can easily be distinguished from male Leptanillini by the presence of a pterostigma (although wing venation may be inaccessible due to deciduous wings in some male Leptanillini [pers. obs.]) and the absence of an ocellar tubercle. Griebenow (in press) provides a formal description of female-associated male *Protanilla* and a male-based definition of the leptanilline tribes, as well as *O. hungvuong*. *Yavnella s. l.* is likewise well-supported (Figs. 1-2), as is *Leptanilla s. l.*, with the former clade diagnosed almost entirely by morphological symplesiomorphies: the only putative autapomorphy of *Yavnella s. l.* is concavity of the propodeum in profile view (Fig. 7A), which was previously noted by Kugler (1986) as being distinctive to *Yavnella*.

Leptanilla s. str. is identifiable relative to other subclades of *Leptanilla s. l.* based upon the following combination of male morphological characters: absence of posterior mesoscutellar prolongation (observed in *Leptanilla zhg-th01* and *Leptanilla TH01*); propodeum convex and without distinct dorsal face (Fig. 7C); gonopodites articulated (otherwise among the Leptanillini articulated only in *Leptanilla zhg-th01* and some *Noonilla*); gonocoxae fully separated ventrally (this character state [Fig. 8A] elsewhere observed among sampled Leptanillini in all *Yavnella s.*

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

360 *l.* except for *Yavnella* TH03; and *Leptanilla* zhg-th01); and penial sclerites dorsoventrally
361 compressed along their entire length, entire, and lacking sculpture (Fig. 9A), this character state
362 elsewhere observed in Leptanillini only among *Yavnella s. l.* (excluding *Yavnella* TH03) and
363 *Leptanilla* zhg-th01.

364 *Leptanilla* TH09 is weakly recovered as sister to remaining *Leptanilla s. str.*, including *P.*
365 *javana*, under all Bayesian total-evidence analyses (Figs. 1-2, 4). Therefore, the phylogeny of *P.*
366 *javana* relative to other *Leptanilla s. str.* would not be resolved if that clade were delimited to
367 exclude *Leptanilla* TH09. However, *Leptanilla* TH09 conforms fully to the diagnosis of
368 *Leptanilla s. str.* given above, and aside from apomorphies of the foreleg (a perhaps opposable
369 calcar and apical probasitarsal seta; Fig. 10A) is not a phenotypic outlier among the terminals
370 representing *Leptanilla s. str.* Nor given the weak BPP of *Leptanilla* TH09 as sister to the
371 remainder of *Leptanilla s. str.* is there probabilistic support for qualitatively defining that clade to
372 exclude *Leptanilla* TH09. Therefore, *P. javana* can be confidently placed within *Leptanilla s.*
373 *str.*, despite the inability of Bayesian total-evidence inference from these data and under these
374 models to resolve its position within that clade.

375 Unlike *Scyphodon*, *Noonilla*, and even the male-based species *Leptanilla palauensis* Smith, 1953
376 (Petersen, 1968: p. 593), the status of *Phaulomyrma* as a leptanilline—and as an ant—has never
377 been debated. Wheeler & Wheeler (1930) established the genus based upon the presence of wing
378 veins and “unusually large genitalia” (Wheeler & Wheeler 1930: p. 193), transferring also
379 *Leptanilla tanit* Santschi, 1907 to *Phaulomyrma*. Their argument regarding wing venation has no
380 merit, given that the forewing venation of *P. javana* falls within the range of variation observed
381 in putative *Leptanilla* morphospecies (Petersen 1968: pp. 594-595), with all leptanilline males
382 examined by Boudinot (2015) exhibiting at least one compound abscissa on the forewing.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Petersen (1968: p. 597) even referred to *Leptanilla* and *Phaulomyrma* as “nearly identical” (when comparing these taxa to *L. palauensis*), and returned *L. tanit* to *Leptanilla*, but refrained from synonymizing *Leptanilla* and *Phaulomyrma* on account of the apparent uniqueness of the genitalia of *P. javana* as illustrated by Wheeler and Wheeler (1930: Figs. 2A, C). In passing, Taylor (1965: p. 365) also mentioned *Phaulomyrma* as being “possibly synonymous” with *Leptanilla*.

Examination of a syntype of *P. javana* (lectotypified below) demonstrates that its genitalia are consistent with other sampled male *Leptanilla s. str.* to the exclusion of males within the Indo-Malayan sister-group of *Leptanilla s. str.* (Fig. 11). Although the preservation of this specimen on a slide prevents us from directly confirming stylar articulation, the sharply recurved styli are consistent with the syndrome seen in dried male leptanillines with articulated gonopodites (Kugler 1986; Ward and Sumnicht 2012), indicating that the gonopodites are articulated in *P. javana*. *Contra* Fig. 2C of Wheeler and Wheeler (1930), the volsellae of *P. javana* are not discernible *in situ* (Fig. 11D). If their condition is truly “plate-like” as described by Wheeler and Wheeler (1930: p. 196), the volsellae of *P. javana* resemble those observed in undescribed Sicilian male morphospecies attributed to *Leptanilla* (Scupola & Ballarin 2009). Dissection of Anatolian *Leptanilla* GR03, and Spanish material that closely resembles sequenced males of *Leptanilla s. str.*, demonstrates that the volsellae are likewise lamellate in these morphospecies, having much the same condition as in *Leptanilla africana* (Baroni Urbani 1977: Fig. 37) (not included in this study). Therefore, given the phylogeny of *P. javana* and its morphological conformity to *Leptanilla s. str.* there is no justification for maintaining the genus *Phaulomyrma*. It is a nomenclatural irony that Wheeler & Wheeler (1930: p. 193) note that the derivation of the

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

genus name is from the Greek *phaulus*, which they translate as “trifling or paltry”: the justification for establishing *Phaulomyrma* as a genus was trifling indeed.

A complete male-based diagnosis of *Leptanilla* relative to other Leptanillinae under both broad and strict circumscriptions of *Leptanilla* is provided below, with putative synapomorphies for the two circumscriptions represented in italics. Only genital characters could be scored for *Leptanilla* ZA01.

Leptanilla javana (Wheeler & Wheeler, 1930) comb. nov.

Figs. 11A-D.

Phaulomyrma javana Wheeler & Wheeler 1930: 193. Figs. 1, 2C.

Phaulomyrma javana – Petersen 1968: 293. Figs. 16A-C.

Lectotype

INDONESIA • ♂; Jawa Barat, “Buitenzorg” [Bogor]; Mar. 1907; F.A.G. Muir leg.; MCZ 31142.

Paralectotype

Same data as for lectotype (no accession code).

Genus *Leptanilla* Emery, 1870

Type species: *Leptanilla revelierii* Emery, 1870: 196.

= *Leptomesites* Kutter, 1948 (286). Synonymy by Baroni Urbani, 1977 (433). Holotype deposited at MHNG (Muséum d’Histoire Naturelle, Geneva).

= *Phaulomyrma* Wheeler & Wheeler, 1930 (193); **syn. nov.** Lectotype and paralectotype deposited at MCZC (Museum of Comparative Zoology, Cambridge, Massachusetts).

Male diagnosis of Leptanilla s. l. relative to other Leptanillinae

1. Mandibles articulated to gena (Fig. 12B).

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

- 427 2. *Medial axis of clypeus no longer than diameter of torulus, when epistomal sulcus is*
428 *distinct.*
- 429 3. Antennomere 3 shorter than scape.
- 430 4. Ocelli present and set on tubercle (Fig. 13) (with exception of *Leptanilla* [Bornean
431 morphospecies-group] zhg-my05).
- 432 5. *Pronotum and mesoscutum posteriorly extended (Fig. 14B-C).*
- 433 6. Notauli absent.
- 434 7. Pterostigma absent.
- 435 8. Propodeum not concave in profile view.
- 436 *Male Diagnosis of Leptanilla s. str. relative to other Leptanilla s. l.*
- 437 9. *Anteromedian ocellus and compound eye not intersecting line parallel to dorsoventral*
438 *axis of cranium.*
- 439 10. Profemoral ventral cuticular hooks absent.
- 440 11. Ventromedian protibial comb-like row of setae absent.
- 441 12. Infuscation at juncture of Rf and 2s-rs+Rs+4-6 absent.
- 442 13. *Antero-admedian line absent (HAO: 0000128).*
- 443 14. Mesoscutellum not posteriorly prolonged.
- 444 15. Propodeum convex in profile view, without distinct dorsal face.
- 445 16. Abdominal sternite IX without posterolateral filiform processes.
- 446 17. Abdominal tergite VIII broader than long.
- 447 18. Gonocoxae medially separated*.
- 448 19. Gonopodites articulated.
- 449 20. *Volsella lamellate, entire distally, without denticles*.*

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

21. Penial sclerites dorsoventrally compressed, dorsomedian carina absent, ventromedian carina sometimes present.

22. Phallotreme situated at penial apex, without vestiture.

*These character states observed so far as is possible with available specimens.

Notes.

1. The mandibles are fused to the gena (Fig. 12A) in sampled *Yavnella s. l.* except for *Yavnella* TH04: this character state is seen elsewhere in male ants only among some *Simopelta* spp. (Ponerinae) (Brendon Boudinot, pers. comm.).
2. The epistomal sulcus is often difficult to distinguish in *Leptanilla s. l.*, but the anteroposterior reduction of the clypeus can be inferred by the situation of the toruli at the anterior-most margin of the head (cf. Boudinot 2015: p. 30).
3. Antennomere 3 is longer than the scape in all sampled *Yavnella s. l.* except for *Yavnella* TH05.
4. Ocelli are entirely absent in *Yavnella* TH03 and *Yavnella* zhg-bt01. The ocellar tubercle is absent in the Anomalomyrmini and *O. hungvuong*; within *Leptanilla s. l.* it is absent in *Leptanilla* zhg-my05, which is here inferred to be a secondary loss.
5. As noted by Petersen (1968: p. 87), *N. copiosa* contrasts with other described male Leptanillinae by the lack of an “elongated, laterally compressed” mesosoma. *Yavnella* was described by Kugler (1986) as sharing this condition, which Petersen (1968) adduced as plesiomorphic for the Leptanillinae. While the relative modification of the mesosoma—here approximated by the proportions of the pronotum and mesoscutum—forms a morphocline across the male Leptanillinae, this morphocline is discontinuous, with a gap between the morphospace occupied by *Leptanilla s. l.* (Fig. 14B-C) and that

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

occupied by *O. hungvuong*, the Anomalomyrmini, and *Yavnella s. l.* (Fig. 14A). Future sampling of male Leptanillinae may close this gap in morphospace, which would limit the diagnostic utility of pronotal and mesonotal length.

6. The absence of notauli is a synapomorphy of the tribe Leptanillini. The notauli in *Protanilla* TH01 and *Protanilla* zhg-vn01, in the tribe Anomalomyrmini, are homoplastically absent.
7. The absence of the pterostigma (Figs. 15A, C) is a synapomorphy of the Leptanillini.
8. The convexity of the propodeum in profile view is plesiomorphic for the Leptanillinae. Its concave condition in *Yavnella* (Kugler 1986) is apomorphic for that genus.
9. The anteromedian ocellus is not situated orthogonally to the compound eye in profile view in *Leptanilla s. str.* *Leptanilla* TH01 and zhg-th01, the Bornean morphospecies-group, and all examined *Noonilla*. The concomitant prognathy of the male cranium is unique among male Leptanillinae to *Leptanilla s. l.*, and as adduced by Petersen (1968), this condition appears apomorphic among the Leptanillinae.
10. A profemoral ventral cuticular hook (Fig. 16B) is unique among the morphospecies sampled herein to *Leptanilla* (“Bornean morphospecies-group”) zhg-my02 and -05.
11. The ventromedian comb-like row of setae on the protibia is an autapomorphy of the Bornean morphospecies-group.
12. The infuscation observed in the Bornean morphospecies-group at the juncture of Rf and 2s-rs+Rs+4-6 (Fig. 15C) is not enclosed anteriorly by an abscissa and appears to be homoplasious with the pterostigma observed in male Anomalomyrmini. Infuscation of the forewing is otherwise absent in the Leptanillini.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

13. The antero-admedian line is present among sampled Leptanillini only among some *Yavnella s. l.*
14. The mesoscutellum is posteriorly prolonged in *Leptanilla* TH01 and *Leptanilla* zhg-th01 (Fig. 6B). The differences in mesoscutellar shape between these morphospecies (see Appendix) are such that the homology of posterior mesoscutellar prolongation is uncertain.
15. The propodeum has a distinct planar to depressed dorsal face in the Bornean morphospecies-group (Fig. 7B). This condition is an autapomorphy of that clade.
16. The posterior margin of abdominal sternite IX is variously emarginate to entire in male Leptanillinae or with a posteromedian process (e.g., *Protanilla* zhg-vn01, *Yavnella* TH03), but posterolateral filiform processes of abdominal sternite IX are an autapomorphy of the Bornean morphospecies-group.
17. Abdominal tergite VIII is longer than broad only in *Noonilla* (Fig. 17B), *Scyphodon* and a bizarre male morphospecies from Côte d'Ivoire (CASENT0102373) for which molecular data are unavailable.
18. The gonocoxae exhibit partial (Fig. 8B) to full (Fig. 8C) medial fusion at least in ventral view in *Yavnella* TH03, *Noonilla*, and all sampled members of the Bornean morphospecies-group. Within *Leptanilla s. l.*, complete lack of medial gonocoxal fusion (Fig. 8A) is a symplesiomorphy of *Leptanilla s. str.*, *Leptanilla* TH01, and *Leptanilla* zhg-th01.
19. Articulation of the gonopodites encompasses both cases in which conjunctival membrane is visible between the gonocoxa and stylus, and those in which the stylus is recurved relative to the gonocoxa without apparent conjunctival membrane. This character state is

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

a symplesiomorphy of *Leptanilla s. str.*, and among *Leptanilla s. l.* included in this study is observed in *Noonilla* zhg-my02 and -6, and *Leptanilla* zhg-th01.

20. The volsellae cannot be observed without dissection in many male Leptanillinae (e.g., *Noonilla*), limiting my ability to assess their condition. However, *Leptanilla s. str.* contrast with the Anomalomyrmini, *Yavnella s. l.*, and the Bornean morphospecies-group in that the volsellae (where visible) are dorsoventrally flattened, entire, and lacking sculpture (Fig. 18). This is one of only two synapomorphies of *Leptanilla s. str.* relative to other *Leptanilla s. l.*

21. Dorsoventral compression at the penial apex is also observed in *Yavnella s. l.* (except for *Yavnella* TH03). In the Indo-Malayan sister clade of *Leptanilla s. str.* the penial sclerites are lateromedially compressed to subcircular, at least basally. *Leptanilla* zhg-th01 exhibits an intermediate condition, with the penial apex being lateromedially compressed and this condition less pronounced towards the base.

22. Position of the phallotreme with distal margin adjoining the penial apex appears to be ancestral for the Leptanillini. The phallotreme is shifted basally in *Leptanilla* zhg-my02 and -5 (Fig. 19B), *Noonilla*, and *Scyphodon*. The outline of the phallotreme is subcircular in these morphotaxa. Setae surrounding the phallotreme are observed in *Noonilla* and *Scyphodon*; this character state is likely a synapomorphy of these genera.

Goals of Future Research

Two described male-based species of *Leptanilla* are worth noting here as requiring further study and acquisition of fresh material: *L. palauensis*, which was transferred with some reservation to *Leptanilla* from *Probolomyrmex* (Formicidae: Proceratiinae) by Taylor (1965); and *Leptanilla astylina*. Examination of the holotype of *L. palauensis* demonstrates that according to the

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

morphological hypotheses made herein this species can be confidently referred to *Leptanilla s. l.*, but beyond that its affinities are unclear. Based upon available illustrations (Petersen 1968: Fig. 1) *L. astylina* likewise can be placed in *Leptanilla s. l.*, and closely resembles *Leptanilla s. str.*, excluding its genitalia, which to judge from Petersen (1968) are unlike those of any specimen that was examined in this study, and exclude it from the definition of *Leptanilla s. str.* given herein.

The case of *Scyphodon* must also be briefly addressed here. Examination of a specimen attributable to this monotypic male-based genus shows that it can be placed in *Leptanilla s. l.* As reported by Petersen (1968), the genitalia of *Scyphodon* conspicuously resemble those of *Noonilla*: there is no reason to conclude that *Scyphodon* belongs within *Leptanilla s. str.*, and I predict that *Scyphodon* is either sister to, or nested within, *Noonilla*. Future total-evidence Bayesian phylogenetic inference will resolve the relation of *Scyphodon* to other *Leptanilla s. l.*

Future acquisition and examination of novel material may necessitate revision of the male diagnosis of *Leptanilla* provided here, but this diagnosis is robust to all morphological observations made with sequenced material. As *Yavnella s. l.*, *Noonilla* and the Bornean morphospecies-group are known only from males, and *L. revelierii* is known only from female castes, no argument can yet be made regarding the ranking of the former clades relative to *Leptanilla*. *Yavnella* is here ranked as a genus, but the description of *Yavnella* workers may reveal a morphological basis for subjective arguments for the subsumption of *Yavnella* within *Leptanilla*. The delimitation of genera within the Leptanillini—including the status of *Noonilla* and undescribed male morphospecies more closely related to that genus than to *L. revelierii*—therefore depends not only upon phylogenetic resolution of the many lineages known only from male material, but upon the morphology of corresponding workers. Future molecular sequencing

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

will be needed to associate workers and/or gynes to leptanilline lineages that are known only from males: such an effort has successfully linked *Protanilla lini* (Anomalomyrmini) with previously unassociated males (Griebenow, in press).

Conclusions

I have here demonstrated the utility of discrete morphological data within a total-evidence framework that includes molecular data in inferring the phylogeny of an ant taxon known only from male morphology. Using probabilistic models, the phylogenetic position of *P. javana* is robustly inferred in conjunction with taxa for which only molecular data, or both these and male morphological data, are available. In that phylogeny, *P. javana* and *L. revelierii* are confidently recovered within a subclade easily diagnosed by male morphological characters; disregarding future retrieval of worker material and/or novel male specimens, *Phaulomyrma* can be synonymized with *Leptanilla* despite continued uncertainty in the bounds of the latter genus. Future work will employ this Bayesian total-evidence approach to infer the affinity of other, more peculiar leptanilline taxa for which molecular data are unavailable. With a robust phylogeny inferred for the Leptanillinae that is congruent with male morphology, the parallel taxonomy that bedevils this little-understood group of ants can begin to be resolved.

Conflict of Interest

The author declares no conflict of interest.

Acknowledgments

First and foremost, I must thank Ziad Khouri for his generosity in providing indispensable assistance and conceptual advice on the writing of scripts for the Bayesian phylogenetic analyses upon which this project relied. I thank Jadranka Rota (MZLU), Debbie Jennings (ANIC), Kevin

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Williams (CSCA), and Brian Fisher (CASC) for loans of material for examination and non-destructive DNA extraction; Stefan Cover was of great help in facilitating access to a syntype of *Leptanilla javana*. I also thank my present and past lab-mates Jill Oberski and Matt Prebus for their diligent and painstaking work enriching UCEs from the specimens that were used in this study (sometimes with my help, sometimes not). In the realm of data collection, I am grateful to Eli Sarnat, Steve Heydon and Lynn Kimsey for allowing me the usage of equipment for this study; the aid of Michael Branstetter in providing advice on the retrieval of legacy loci from UCE datasets; Ziv Lieberman in advising me to use *BBMap*; and my labmate Brendon Boudinot, who was an enduring source of informative feedback on the coding of morphological observation into discrete character states. Lastly, I must thank my adviser Phil Ward for his invaluable feedback on the construction and finer details of this manuscript, along with past work in acquiring the Sanger-sequenced data that I utilized in this study, and for tutoring me in the delimitation of protein-coding loci from flanking introns. This research was supported by the University of California, Davis and by NSF grant DEB-1932405 to P. S. Ward.

Appendix

Definition of binary presence/absence morphological characters. Note that all non-genital morphological data are missing in *Leptanilla* ZA01, since all that remained of this specimen after destructive DNA extraction was the male genitalia. Missing observations are noted for other terminals where relevant. Males of *Protanilla lini* Terayama, 2009 were identified as such by molecular data (Griebenow, in press).

1. Mesal protibial margins carinate: a sclerotized carina (Fig. 20A) is present (1) on the mesal margin of the ventral protibial surface in *Noonilla*. This character could not be

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

scored in *Leptanilla* TH01. Under the alternative character state (0) the mesal protibial face is convex (Fig. 20B) to carinate.

2. Ventral cuticular hook present on profemur: the lateral margin of the ventral profemoral surface is ventrally produced into a hook-like structure (Figs. 16B, 21B) (1) in *Leptanilla* (Bornean morphospecies-group) zhg-my02 and -my05. The morphospecies imaged in Figs. 16B, 21B is closely related to these (Griebenow, in press) but was not sequenced in this study. Under the alternative character state (0) there are no cuticular extensions of the profemur (Fig. 16A). This character could not be scored in *Leptanilla* TH01.
3. Row of ventral protibial bristles present: a single medial row of parallel-sided setae is present (1) on the ventral protibial surface only in the “Bornean morphospecies-group” (Fig. 21B). These are robust by comparison to adjacent unmodified setae. Under the alternative character state (0) setae on the protibial venter are not robust, parallel-sided, and arranged in a single medial row (Fig. 21A). This character could not be scored in *Leptanilla* TH01.
4. Head inclusive of compound eyes wider than long: this character state is observed (1) in *O. hungvuong*; all male *Anomalomyrmini* sampled herein; all *Yavnella s. l.* except for *Yavnella* TH05, -8, and *Yavnella* MM01; *Leptanilla* (Bornean morphospecies-group) zhg-my04; and *Noonilla* zhg-my06 (Fig. 22B). Under the alternative character state (0) the head inclusive of the compound eyes is narrower than long in full-face view (Fig. 22A).
5. Clypeus broader than torular diameter along medial axis: this character state is observed (1) in *M. heureka*; *O. hungvuong*; in all *Anomalomyrmini* sampled herein; and in all *Yavnella s. l.* for which observations are available (Fig. 23B). Clypeus narrower than

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

torular diameter along medial axis (0) (Fig. 23A) may therefore be diagnostic for

Leptanilla s. l. This character could not be scored in *Yavnella* TH03, -5 and -8; *Yavnella* zhg-bt01; *Phaulomyrma* MM01; *Leptanilla* zhg-th01; and *Leptanilla* GR01-3, zhg-au02 and zhg-bt01.

6. Anterior tentorial pits situated directly anterior to toruli: the anterior tentorial pits are situated directly anterior to the toruli, in whole (Fig. 24B) or in part (1) so that at least some portion of the anterior tentorial pit intersects an anteroposterior axis drawn through the torulus, in *M. heureka*, *O. hungvuong*, all Anomalomyrmini save *Protanilla* TH01, and all *Yavnella s. l.* save *Yavnella* TH05 and MM01. Under the alternative character state (0), the anterior tentorial pits are situated anterolaterad the toruli or may not be readily discernible (Fig. 24A), so that no part of the anterior tentorial pit intersects an anteroposterior axis drawn through the torulus. This character could not be scored in *Yavnella* TH03 and -8, *Leptanilla* (Bornean morphospecies-group) zhg-my02, *Leptanilla* zhg-au02, and *Leptanilla javana*.

7. Antennomere 3 longer than scape: this character state (1) (Fig. 25B) is observed in *Protanilla* TH03 and all *Yavnella s. l.* except for *Yavnella* TH05. Under the alternative character state (0) the scape is shorter than (Fig. 25A) or subequal in length to antennomere 3. This character could not be scored in *O. hungvuong* or *Leptanilla* zhg-au02.

8. Mandible articulated to gena: the base of the male mandible is visibly fused to the gena (0) in all *Yavnella s. l.* for which observations are available (Fig. 12A), except for *Yavnella* TH04. In all other terminals in which this character can be assessed a complete

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

- point of articulation to the gena is visible (1) (Fig. 12B). This character could not be scored in *Yavnella* TH03 and MM01; and *Leptanilla* zhg-au02, -TH09 and *javana*.
9. Occipital margin angularly emarginate in dorsal view: the occiput is coded as angularly emarginate in dorsal view (1) if the posterolateral corners of the occipital margin are produced; this character state is observed in *Leptanilla* TH01 (Fig. 26B), *Leptanilla* zhg-th01, and the Bornean morphospecies-group except for *Leptanilla* zhg-my03. Under the alternative character state (0) the occiput is linear to shallowly emarginate (Fig. 26A).
10. Mesoscutum convex in profile view: the mesoscutum is scored as convex (1) if not planar to shallowly convex (0) (Fig. 27A). Mesoscutal convexity (1) (Fig. 27B) is present in *M. heureka*, *O. hungvuong*, the Anomalomyrmini, *Yavnella s. l.*, and *Leptanilla* (Bornean morphospecies-group) zhg-my04.
11. Notauli present: the presence (1) or absence (0) (Fig. 28A) of notauli is always unambiguous. These are observed only in *M. heureka*, *Protanilla* TH01, and -03 (Fig. 28B).
12. Parapsidal signa present: the presence (1) (Fig. 29B) or absence (0) (Fig. 29A) of the parapsidal signa can be difficult to discern, varying from a distinct impressed signum to a stripe of glabrous cuticle. Some form of parapsidal signum is present in *M. heureka*; *O. hungvuong*; *Protanilla* zhg-vn01; *Yavnella* zhg-th01, cf. *indica* and *argamani*, *Yavnella* TH02, -4 and -6; *Yavnella* MM01; *Yavnella* TH01; *Noonilla* zhg-my06; the Bornean morphospecies-group; and *Leptanilla* GR01.
13. Oblique mesopleural sulcus adjoining posterior mesopectal margin: this character state is observed (1) in *O. hungvuong*, all Anomalomyrmini (Fig. 30B) and most *Leptanilla s. str.* for which this character can be scored, except for *Leptanilla* GR01, -03, TH09, and

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

- Leptanilla* zhg-bt01. Complete bisection of the mesospectus by the oblique mesopleural sulcus is seen in the Anomalomyrmini. The alternative character state (0) encompasses a morphocline from the near-complete loss of the oblique mesopleural sulcus (as in *Leptanilla* zhg-bt01) to the termination of this feature immediately anterior to the upper metapleuron (e.g., *Yavnella* TH02: Fig. 30A) or propodeum (as in the Bornean morphospecies-group). This character could not be scored in *Leptanilla* zhg-au02.
14. Pterostigma present: this character state is observed (1) only in *M. heureka*, *O. hungvuong*, and the Anomalomyrmini (Fig. 31B). Rf and 2s-rs+Rs+4-6 are confluent in the Bornean morphospecies-group and in *Noonilla* zhg-my06, producing an infuscation of the wing membrane that resembles a pterostigma (0). No infuscation or pterostigma (0) is observed in all other terminals scored (Fig. 31A). Wings are lost in all available specimens of *Noonilla* zhg-my02, *Leptanilla* zhg-th01, and *Leptanilla* GR03; therefore, this character could not be scored in these terminals.
15. Mesoscutellum densely pubescent: the mesoscutellum is covered with sparse setae (0) in all leptanilline males sampled herein except for *Leptanilla* TH01 and zhg-th01, and the Bornean morphospecies-group (Fig. 32B); in these cases, the mesoscutellar vestiture is densely pubescent (1) (Fig. 32A). This character could not be scored in *Yavnella* TH04.
16. Mesoscutellum projecting posteriorly in profile view: this character state is observed (1) either as a dorsoventrally robust cuneiform process (*Leptanilla* TH01) or as a recurved spine (*Leptanilla* zhg-th01) (Fig. 6B). Under the alternative character state, the posterior margin of the mesoscutellum is rounded (0) (Fig. 6A). This character could not be scored in *Yavnella* TH02.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

17. Propodeum concave in profile view: this character state (1) (Fig. 7A) is an autapomorphy of *Yavnella s. l.* Under the alternative character state (0) the propodeum is convex in profile view (Fig. 7C) or produced into a right angle, with largely planar dorsal and posterior faces (the Bornean morphospecies-group; Fig. 7B).
18. Abdominal tergite II produced into distinct node: there is a shallow to pronounced dorsal node (Fig. 33B) present on the petiole (1) in *O. hungvuong*, *Protanilla zhg-vn01* and TH01-2, *Yavnella* TH08, *Leptanilla zhg-th01*, the Bornean morphospecies-group, and *Leptanilla s. str.* except for *Leptanilla zhg-au02*. Under the alternative character state (0) the dorsal surface of the petiole is slightly convex (Fig. 33A), or planar without any supra-axial projection (as in *Leptanilla zhg-au02*).
19. Abdominal sternite II with ventral process: a ventral rounded to angular process (1), shallow or well-produced, is present on abdominal sternite II in *Protanilla zhg-vn01* and TH02, *Leptanilla zhg-my02* (Fig. 34C) and -5, and *Leptanilla s. str.* except for *Leptanilla zhg-au02*, and *javana*. Under the alternative character state (0) there is no ventrally projecting process on abdominal sternite II (Fig. 34A). A moderate ventral bulge without a distinct anterior and/or posterior face may be present under this character state (Fig. 34B). This character could not be scored in *Protanilla* TH01.
20. Petiole higher than long including peduncle: this character state (Fig. 35B) is observed in profile view (1) in *Protanilla zhg-vn01* and TH01-2, *Yavnella* MM01, *Yavnella* TH05, *Yavnella cf. indica* and zhg-th01, *Leptanilla* TH01, the Bornean morphospecies-group, and *Noonilla*. This includes cases in which there is no distinct dorsal node. Under the alternative character state (0) the distance between two lines drawn tangential to the

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

- dorsal- and ventral-most points of the petiole in profile view is no greater than petiole length in profile view (Fig. 35A). This character could not be scored in *Yavnella* TH02.
21. Cinctus present on abdominal segment III: the corollary of this character state (1) is the existence of a petiole (Fig. 36B), which has been secondarily lost (0) in *Yavnella* zhg-th01 (Fig. 36A), *Yavnella* TH02 (as noted by Boudinot 2015: p. 14), and *Noonilla* zhg-my02. There is a tendency towards petiolar reduction in *Yavnella* s. l. and *Noonilla*, but in many cases a cinctus on abdominal segment III is still discernible.
22. Cinctus present on abdominal segment IV: the corollary of this character state (1) is the presence of a post-petiole. This character state is unique to *Protanilla* TH03 (Fig. 37B), although the anterior margin of abdominal segment IV may be slightly constricted relative to more posterior abdominal segments (0); otherwise, there is no constriction whatsoever (Fig. 37A).
23. Abdominal sternite IX with posteromedian filiform process: while a posteromedian process of abdominal sternite IX is present in all male Anomalomyrmini and *Opamyрма hungvuong* (0), its filiform condition (1) is unique to *Yavnella* TH03. Abdominal sternite IX is not thus produced medially in all other male leptanillines sampled herein (0).
24. Abdominal sternite IX with posterolateral filiform processes: these “bizarre, elongate, filamentous extensions” of the metasoma were noted by Boudinot (2015: Fig. 10D) as being extensions of the gonocoxae *sensu* Boudinot (2018). Detailed examination and micro-CT segmentation (Griebenow, Fischer and Economo in prep.) demonstrate that these processes are in fact extensions of abdominal sternite IX (Fig. 38B). This character state is unique to the Bornean morphospecies-group. Under the alternative character state (0) the posterior margin of abdominal sternite IX may be medially indented (Fig. 38A),

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

entire, or with a posteromedian process, as noted above. This character could not be scored in *Leptanilla* zhg-au02.

25. Abdominal tergite XIII broader than long: this character state is observed (1) in all male Leptanillinae scored (Fig. 17A) except for *Noonilla*, to which elongation of abdominal tergite XIII (0) is unique (Fig. 17B). This character could not be scored in *Yavnella* MM01 and *L. javana*.

26. Gonocoxae ventromedially fused along entire length: this character state (Fig. 8C) is observed in *O. hungvuong*, *Yavnella* TH03, and in all terminals within the Bornean morphospecies-group that could be scored (1). The alternative character state (0) encompasses partial (Fig. 8B) to complete (Fig. 8A) ventromedian fusion of the gonocoxae. This character could not be scored in *Protanilla* TH01, *Yavnella* MM01, *Leptanilla* TH01, *Leptanilla* (Bornean morphospecies-group) zhg-my05, and *L. javana*.

27. Gonocoxae dorsomedially fused along entire length: this character state is observed (1) in *O. hungvuong*, *Yavnella* TH03 and the Bornean morphospecies-group (Fig. 39B). Under the alternative character state (0) the gonocoxae are fully (Fig. 39A) to partly separate medially. This character could not be scored without dissection in *Noonilla* (in which abdominal tergite XIII conceals the gonocoxal dorsum) or in *Leptanilla* zhg-au02, -bt01, ZA01, and *javana*.

28. Gonocoxa with ventral lamina: a ventral laminate margin, variably produced and shaped, is present (1) on the gonocoxa (Fig. 40B), or on the basal part of the gonopodite in those cases in which the gonocoxa and stylus are insensibly fused, in *Yavnella* cf. *indica*; *Leptanilla* zhg-th01; *Leptanilla* (Bornean morphospecies-group) zhg-my02 and -5; and *Leptanilla* TH09, GR01-2, ZA01 (Fig. 40B), and zhg-au02. Under the alternative

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

character state (0) no lamina is discernible whatsoever on the gonocoxa (Fig. 40A). I do not assert primary homology *sensu* de Pinna (1991) of laminate portions of the gonopodites in *Leptanilla* zhg-my02 and -5 with styli, since this does not meet the criterion of conjunction (Patterson 1982; de Pinna 1991) in CASENT0178838, a heterospecific member of the Bornean morphospecies-group (misattributed to *Protanilla* by Boudinot [2015]). This character could not be scored in *Protanilla* TH01 or in *Leptanilla* zhg-bt01 and *javana*.

29. Stylus articulated to gonocoxa: this character state (1) includes cases in which the stylus is sharply deflexed relative to the gonocoxa (Fig. 41B) or a conjunctiva is visible between the gonopodital sclerites. Under the alternative character state (0) a suture might be visible (as in many *Yavnella* *s. l.*) or the gonocoxa and stylus insensibly fused (as in the Bornean morphospecies-group; Fig. 41A). Gonopodital articulation is fully present in *O. hungvuong*, *Protanilla* zhg-vn01, *Yavnella* zhg-bt01, *Leptanilla* zhg-th01, all *Leptanilla* *s. str.* for which this character can be scored and both *Noonilla* included in this study. This character could not be scored in *Leptanilla* zhg-bt01.

30. Gonopodital apex with vestiture: this character and the next are so termed in order to encompass cases in which the stylus is insensibly fused to the gonocoxa (figs. 41A, 42A-B). The only terminals sampled here in which setal vestiture is not present on the gonopodital apex (0) are the Bornean morphospecies-group (Fig. 42A) except for *Leptanilla* zhg-my04. Otherwise (1) there are at least some setae present on the gonopodital apex (Fig. 42B). This character could not be scored in *Leptanilla* zhg-bt01.

31. Gonopodital apex bifurcated: this character state is observed (1) only in *Yavnella* TH08 (Fig. 43B), *Leptanilla* ZA01 and GR02. Under the alternative character state (0) the

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

stylus may be entire (Fig. 43A) or may have a subapical tooth. This character could not be scored in *Leptanilla* zhg-bt01.

32. Penial sclerites enclosed dorsally by gonopodites at base: in this character state (1) the gonopodites may completely enclose (Fig. 44C) or partially overlap with (Fig. 44B) the penial sclerites. This character state is observed in *M. heureka*, *Yavnella* TH03, *Yavnella* zhg-th01 and zhg-bt01, the Bornean morphospecies-group, and *Leptanilla* zhg-au02. Under the alternative character state (Fig. 44A) (0), the penial sclerites are never dorsally surmounted by any portion of the gonopodites. This character could not be scored in *Noonilla* zhg-my06 and *L. javana*.

33. Penial sclerites dorsally recurved at base in profile view: among the terminals sampled here, this bizarre character state (1) is only present in *Leptanilla* (Bornean morphospecies-group) zhg-my02 and -5 (Fig. 45B). In these cases the penial sclerites are curved at the base so that in preserved specimens the apex is situated dorsally of the gonocoxae. Otherwise (0) in profile view the penial sclerites are slightly curved at the base towards the venter of the genital anteroposterior axis (Fig. 45A) or are parallel to that axis.

34. Penial sclerites dorsoventrally compressed at base: this character state is observed (1) in *M. heureka*, and all *Yavnella s. l.* (Fig. 46B) and *Leptanilla s. str.* for which this character can be scored. Under the alternative character state (0) the penial sclerites are basally wider along the dorsoventral axis, exclusive of any ventromedian processes, than along the lateromedial axis (Fig. 46A). This character could not be scored in *O. hungvuong*, *Protanilla* TH02-3, *Yavnella* TH03-4 and zhg-bt01, *Leptanilla* TH01, *Noonilla* zhg-my06, and *L. javana*.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

35. Penial sclerites dorsoventrally compressed at apex: this character state is observed (1) in *M. heureka*, *Yavnella s. l.* except for *Yavnella* TH03, *Leptanilla s. str.*, and *Leptanilla* zhg-my03 (Fig. 47B). Under the alternative character state (0) the penial sclerites are apically wider along the dorsoventral axis, exclusive of any ventromedian processes, than along the lateromedial axis (Fig. 47A). The alternative character state (0) encompasses cases in which the penial sclerites are lateromedially compressed to varying extents (e.g., *Anomalomyrmini*) or are subcircular in cross-section (e.g., *Noonilla*). This character could not be scored in *O. hungvuong* or *Protanilla* TH02.
36. Lateral margins of penial sclerites laminate: this character state is observed (1) in *Yavnella* TH02-5, *Yavnella* MM01, *Yavnella* zhg-th01, cf. *indica* and *argamani*; *Leptanilla* zhg-my02 (Bornean morphospecies-group) and *Leptanilla* zhg-my05; and all *Leptanilla s. str.* (Fig. 48B) for which this character can be scored, except for *Leptanilla* zhg-bt01. In the Bornean morphospecies-group the lateral laminae, when present, are strongly produced ventrally relative to the remainder of the penial sclerites. Under the alternative character state (0) (Fig. 48A) lateral flanges may be present or absent, but when present are not laminate. This character could not be scored in *M. heureka* and *Leptanilla* zhg-au02.
37. Penial sclerites with dorsomedian carina: this character state is observed (1) only in *Leptanilla* TH01 and *Leptanilla* (Bornean morphospecies-group) zhg-my04 (Fig. 47A). In both cases the penial sclerites are strongly lateromedially compressed. Under the alternative character state (0) there is no dorsomedian penial carina, such that the dorsum of the penial sclerite(s) is/are rounded in cross-section (Fig. 47B). This character could not be scored in *Protanilla* TH03.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

38. Penial sclerites with ventromedian projection: this character state is observed (1) in *Leptanilla* zhg-my03-4; *Leptanilla* zhg-th01; and *Leptanilla* zhg-bt01, GR01-2, and zhg-au02. When present and discernible, the volsellae flank this projection, which can be rounded (as in *Leptanilla* zhg-my03-4; Fig. 49B) or carinate. Under the alternative character state (0) the penial sclerites are entirely separated, or if fused then lacking any ventromedian process (Fig. 49A). This character could not be scored in *M. heureka*, the *Anomalomyrmini*, *Leptanilla* TH01 and *Yavnella* TH03-8 and *Yavnella* MM01.
39. Phallotreme flanked with vestiture: this character state (1) occurs only in *Noonilla* (Fig. 50B). Under the alternative character state (0) the phallotrematic rim is visibly bare of any setae (Fig. 50A). This character could not be scored in *Yavnella* MM01.
40. Phallotreme preapical: under the alternative character state (0) the phallotreme is situated adjoining the posterior penial margin or, if the penial sclerites are lateromedially compressed, at the penial apex (Fig. 19A). This includes cases in which the phallotreme is situated well basal to the penial apex but has a distal margin that extends to the penial apex. The phallotreme is therefore preapical (1) in *Leptanilla* (Bornean morphospecies-group) zhg-my02-3 and -5 (Fig. 19B), and in *Noonilla* zhg-my06. This character could not be scored in *Protanilla* TH02-3; *Yavnella* TH04, -8 and *Yavnella* zhg-bt01; and *Yavnella* MM01, and *Leptanilla* zhg-bt01 and *L. javana*.
41. Penial apex entire: the alternative (0) to this character state encompasses cases in which the penial sclerites are medially separated at the apex (as in *Protanilla* TH01-2), strongly bifurcated (Fig. 51A). Under this character state (1) none of these observations apply (Fig. 51B), encompassing cases in which the distal phallotrematic margin forms a narrow slit-like indentation in the penial sclerites (e.g., *Yavnella* cf. *indica*: Fig. 44A). The penial

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

apex is entire in *M. heureka*; *Protanilla* TH03 and zhg-vn01; *Yavnella* TH02, -5-8, *Yavnella* cf. *indica*, zhg-bt01, and zhg-th01; *Leptanilla* TH01; *Leptanilla* zhg-th01; the Bornean morphospecies-group; and *Leptanilla* s. str. except for *Leptanilla* ZA01.

References

- Agosti, D. (1991). Revision of the oriental ant genus *Cladomyrma*, with an outline of the higher classification of the Formicinae (Hymenoptera: Formicidae). *Systematic Entomology* **16**, 293–310.
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M., Nikolenko, S. I., Pham, S., Pribelski, A. D., Pyshkin, A. V., Sirotkin A. V., Vyahhi, N., Tesler, G., Alekseyev, M. A., & Pevzner, P. A. (2012). SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* **19**(5), <https://doi.org/10.1089/cmb.2012.0021>
- Bapst, D. W., Wright, A. M., Matzke, N. J., and Lloyd, G. T. (2016). Topology, divergence dates, and macroevolutionary inferences vary between different tip-dating approaches applied to fossil theropods (Dinosauria). *Biology Letters* **12**, 20160237.
- Barden, P., Boudinot, B. E., and Lucky, A. (2017). Where fossils dare and males matter: combined morphological and molecular analysis untangles the evolutionary history of the spider ant genus *Leptomyrme* Mayr (Hymenoptera: Dolichoderinae). *Invertebrate Systematics* **31**, 765–780.
- Baroni Urbani, C. (1977). Materiali per una revision della sottofamiglia Leptanillinae Emery (Hymenoptera: Formicidae). *Entomologica Basiliensia* **2**, 427–488.
- Baroni Urbani C. and de Andrade, M.L. (2006). A new *Protanilla* Taylor, 1990 (Hymenoptera: Formicidae: Leptanillinae) from Sri Lanka. *Myrmecologische Nachrichten* **8**, 45–47.
- Belshaw, R. and Bolton, B. (1994). A survey of the leaf litter ant fauna in Ghana, West Africa (Hymenoptera: Formicidae). *Journal of Hymenoptera Research* **3**, 5–16.
- Billen, J., Bauwleers, E., Hashim, R., and Ito, F. (2013). Survey of the exocrine system in *Protanilla wallacei* (Hymenoptera, Formicidae). *Arthropod Structure and Development* **42**(3), 173–183.
- Bolton, B. (1990). The higher classification of the ant subfamily Leptanillinae (Hymenoptera: Formicidae). *Systematic Entomology* **15**, 267–282.
- Bolton, B. (2003). Synopsis and classification of Formicidae. *Memoirs of the American Entomological Institute* **71**, 1–370.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

- 890 Borowiec M. L., Schulz, A., Alpert, G.D., and Bañar, P. (2011). Discovery of the worker caste
891 and description of two new species of *Anomalomyrma* (Hymenoptera: Formicidae: Leptanillinae)
892 with unique abdominal morphology. *Zootaxa* **2810**, 1–14.
- 893 Borowiec, M. L., C. Rabeling, S. G. Brady, B. L. Fisher, T. R. Schultz, and P. S. Ward. (2019).
894 Compositional heterogeneity and outgroup choice influence the internal phylogeny of the ants.
895 *Molecular Phylogenetics & Evolution* **134**, 111–121.
- 896 Boudinot, B. E. (2015). Contributions to the knowledge of Formicidae (Hymenoptera, Aculeata):
897 a new diagnosis of the family, the first global male-based key to subfamilies, and a treatment of
898 early-branching lineages. *European Journal of Taxonomy* **120**, 1–62.
- 899 Boudinot, B. E. (2018). A general theory of genital homologies for the Hexapoda (Pancrustacea)
900 derived from skeletomuscular correspondences, with emphasis on the Endopterygota. *Arthropod*
901 *Structure and Development* **47**(6), 563–613.
- 902 Brady, S. G., Schultz, T. R., Fisher, B. L., and Ward, P. S. (2006). Evaluating alternative
903 hypotheses for the early evolution and diversification of ants. *Proceedings of the National*
904 *Academy of Sciences of the United States of America* **103**, 18172–18177
- 905 Branstetter, M. G., Longino, J. T., Ward, P. S., and Faircloth, B.C. (2017). Enriching the ant tree
906 of life: enhanced UCE bait set for genome-scale phylogenetics of ants and other Hymenoptera.
907 *Methods in Ecology & Evolution* **8**(1), 768–776.
- 908 Brazeau, M. D. (2011). Problematic character coding methods in morphology and their effects.
909 *Biological Journal of the Linnean Society* **104**, 489–498.
- 910 Brues, C. T. (1925). *Scyphodon*, an anomalous genus of Hymenoptera of doubtful affinities.
911 *Treubia* **6**, 93–96.
- 912 Bushnell, B. (2014). ‘BBMap: a fast, accurate, splice-aware aligner.’ Available at
913 <https://sourceforge.net/projects/bbmap/>.
- 914 Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in
915 phylogenetic analysis. *Molecular Biology & Evolution* **17**, 540–552.
- 916 Chen Z.-L., Shi, F.-M., and Zhou, S.-Y. (2017). First record of the monotypic genus *Opamyрма*
917 (Hymenoptera: Formicidae) from China. *Far Eastern Entomologist* **335**, 7–11.
- 918 Emery, C. (1870). Studi mirmecologici. *Bullettino della Società Entomologica Italiana* **2**, 193–
919 201.
- 920 Faircloth, B. C. (2013). ‘Illumiprocessor: a trimmomatic wrapper for parallel adapter and quality
921 trimming.’ Available at <http://dx.doi.org/10.6079/J6079ILL> Faircloth, B. C. (2016). PHYLUCE
922 is a software package for the analysis of conserved genomic loci. *Bioinformatics* **32**(5), 786–788.
923 <https://doi.org/10.1093/bioinformatics/btv646>
- 924 Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood
925 approach. *Journal of Molecular Evolution* **17**(6), 368–376.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

- 926 Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010).
927 New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
928 performance of PhyML 3.0. *Systematic Biology* **59**(3), 307–321.
- 929 Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X.,
930 ... Regev, A. (2011). Full-length transcriptome assembly from RNA-Seq data without a
931 reference genome. *Nature Biotechnology* **29**(7), 644–652. Published 2011 May 15.
932 doi:10.1038/nbt.1883
- 933 Griebenow, Z. H. In press. Delimitation of tribes in the subfamily Leptanillinae (Hymenoptera:
934 Formicidae), with a description of the male of *Protanilla lini* TERAYAMA, 2009. *Myrmecological*
935 *News* **30**, 229-xxx.
- 936 Hsu, P.-W., Hsu, F.-C., Hsiao, Y., and Lin, C.-C. (2017). Taxonomic notes on the genus
937 *Protanilla* (Hymenoptera: Formicidae: Leptanillinae) from Taiwan. *Zootaxa* **4268**(1), 117–130.
938 <http://dx.doi.org/10.11646/zootaxa.4268.1.7>
- 939 Höhna, S., Landis, M. J., and Heath, T. A. (2017). Phylogenetic inference using RevBayes.
940 *Current Protocols in Bioinformatics*, <https://doi.org/10.1002/cpbi.22>
- 941 Hoang, D. P., Chernomor, O., von Haeseler, A., Minh, B. Q., and Vinh, L. S. (2018). UFBoot2:
942 improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **35**(2), 518–
943 522.
- 944 Katoh, K., Asimenos, G., and Toh, H. (2009). Multiple alignment of DNA sequences with
945 MAFFT. In 'Bioinformatics for DNA Sequence Analysis'. (Ed. D. Posada.) pp. 39–64.
946 (Springer: New York City, NY.)
- 947 Katoh, K., Rozewicki, J., and Yamada, K. D. (2019). MAFFT online service: multiple sequence
948 alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* **20**(4),
949 1160–1166.
- 950 Kjer, K., Borowiec, M. L., Frandsen, P. B., Ware, J. and Wiegmann, B. (2018). Advances using
951 molecular data in insect systematics. *Current Opinions in Insect Science* **18**, 40–47.
- 952 Kück, P., Hita Garcia, F., Misof, B., and Meusemann, K. (2011). Improved phylogenetic
953 analyses corroborate a plausible position of *Martialis heureka* in the ant tree of life. *PLOS One* **6**,
954 e21031.
- 955 Kugler, J. (1986). The Leptanillinae (Hymenoptera: Formicidae) of Israel and a description of a
956 new species from India. *Israel Journal of Entomology* **20**, 45–57.
- 957 Kutter, H. (1948). Beitrag zur Kenntnis der Leptanillinae (Hym. Formicidae): eine neue
958 Ameisengattung aus Süd-Indien. *Mitteilungen der Schweizerischen Entomologischen*
959 *Gesellschaft* **11**, 286–295.
- 960 Lanfear, R., Calcott, B., Ho, S. Y., and Guindon, S. (2012). PartitionFinder: combined selection
961 of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology*
962 *and Evolution* **29**(6), 1695–1701.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

- 963 Lanfear, R., P. B. Frandsen, A. M. Wright, T. Senfeld, and B. Calcott. (2016). PartitionFinder 2:
964 New Methods for Selecting Partitioned Models of Evolution for Molecular and Morphological
965 Phylogenetic Analyses. *Molecular Biology and Evolution* **34**(3), 772–773.
- 966 Leong, C.–M., Yamane, S., and Guénard, B. (2018). Lost in the city: discovery of the rare ant
967 genus *Leptanilla* (Hymenoptera: Formicidae) in Macau with description of *Leptanilla*
968 *macauensis* sp. nov. *Asian Myrmecology* **10**, e010001. DOI: 10.20362/am.010001
- 969 Lewis, P. O. (2001). A likelihood approach to estimating phylogeny from discrete morphological
970 character data. *Systematic Biology* **50**(6), 913–925.
- 971 López, F., Martinez, M. D., and Barandica, J. M. (1994). Four new species of the genus
972 *Leptanilla* (Hymenoptera: Formicidae) from Spain – relationships to other species and ecological
973 issues. *Sociobiology* **24**, 179–212.
- 974 Man, P.; Ran, H., Chen, Z., and Xu, Z. (2017). The northernmost record of Leptanillinae in
975 China with description of *Protanilla beijingensis* sp. nov. (Hymenoptera: Formicidae). *Asian*
976 *Myrmecology* **9**, e009008. DOI: 0.20362/am.009008
- 977 Masuko, K. (1990). Behavior and ecology of the enigmatic ant *Leptanilla japonica* Baroni
978 Urbani (Hymenoptera: Formicidae: Leptanillinae). *Insectes Sociaux* **37**(1), 31–57.
- 979 Matzke, N. J. and Wright, A. (2016). Inferring node dates from tip dates in fossil Canidae: the
980 importance of tree priors. *Biology Letters*, **12**, 20160328. <https://doi.org/10.1098/rsbl.2016.0328>
- 981 Miller, M. A., Pfeiffer, W., and Schwartz, T. (2010). Creating the CIPRES Science Gateway. In
982 ‘Proceedings, the Gateway Computing Environments Workshop (GCE)’, 14 November 2010,
983 New Orleans, LA. pp. 1–8. (Institute of Electrical and Electronics Engineers: Piscataway, NJ.)
- 984 Moreau, C. S., Bell, C. D., Vila, R., Archibald, S. B., and Pierce, N. E. (2006). Phylogeny of the
985 ants: diversification in the age of angiosperms. *Science* (Washington, D. C.) **312**, 101–104.
- 986 Nguyen, L.–T., Schmidt, H. A., von Haeseler, A., and Minh, B. Q. (2015). IQ–TREE: A Fast and
987 Effective Stochastic Algorithm for Estimating Maximum–Likelihood Phylogenies. *Molecular*
988 *Biology and Evolution* **32**(1), 268–274.
- 989 Niehuis, O., Hartig, G., Grath, S., Pohl, H., Lehmann, J., Tafer, H., ... Misof, B. (2012).
990 Genomic and morphological evidence converge to resolve the enigma of Strepsiptera. *Current*
991 *Biology* **22**(14), 1309–1313.
- 992 Ogata, K., Terayama, M., and Masuko, K. (1995). The ant genus *Leptanilla*: discovery of the
993 worker–associated male of *Leptanilla japonica*, and a description of a new species from Taiwan
994 (Hymenoptera: Formicidae: Leptanillinae). *Systematic Entomology* **20**, 27–34.
- 995 O’Reilly, J. E., dos Reis, M., and Donoghue, P. C. J. (2015). Dating tips for divergence–time
996 estimation. *Trends in Genetics* **31**(11), 637–650.
- 997 Patterson, C. (1982). Morphological characters and homology. In ‘Problems of Phylogenetic
998 Reconstruction’. (Eds. K. A. Joysey and A. E. Friday.) pp. 21–74. (Academic Press: London,
999 UK.)

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

- 1000 Petersen, B. (1968). Some novelties in presumed males of Leptanillinae (Hym., Formicidae).
1001 *Entomologiske Meddelelser* **36**, 577–598.
- 1002 de Pinna, M. C. C. (1991). Concepts and tests of homology in the cladistic paradigm. *Cladistics*
1003 **7**, 367–394.
- 1004 Pleijel, P. (1995). On character coding for phylogeny reconstruction. *Cladistics* **11**, 309–315.
- 1005 Rabeling, C., Brown, J., and Verhaagh, M. (2008). Newly discovered sister lineage sheds light
1006 on early ant evolution. *Proceedings of the National Academy of Sciences of the United States of*
1007 *America* **105**(39), 14913–14917.
- 1008 Rambaut, A., Drummond, A. J., Xie, D., Baele, D., and Suchard, M. A. (2018). Posterior
1009 summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **1;67**(5), 901–
1010 904.
- 1011 Robertson, H. G. (2000). Formicidae (Hymenoptera: Vespoidea). In ‘Dâures Biodiversity of the
1012 Brandberg Massif, Namibia’. (Eds. A. H. Kirk–Spriggs and E. Marais.) pp. 371–382.
1013 Cimbabesia Memoir 9. (National Museum of Namibia: Windhoek.)
- 1014 Robertson, J. A. and Moore, W. (2016). Phylogeny of *Paussus* L. (Carabidae: Paussinae):
1015 unravelling morphological convergence associated with myrmecophilous life histories.
1016 *Systematic Entomology* **42**(1), 124–170. <https://doi.org/10.1111/syen.12205>
- 1017 Ronquist, F., Klopfstein, S., Vilhelmsen, L., Schulmeister, S., Murray, D. L., and Rasnitsyn, A.
1018 P. (2012). A total–evidence approach to dating with fossils, applied to the early radiation of the
1019 Hymenoptera. *Systematic Biology* **61**(6), 973–999.
- 1020 Scupola, A. and Ballarin, R. (2009). The genus *Leptanilla* Emery, 1870 in Sicily (Hymenoptera:
1021 Formicidae). *Myrmecological News* **12**, 129–132.
- 1022 Sánchez, N., Yamasaki, H., Pardos, F., Sørensen, M. V., and Martínez, A. (2016). Morphology
1023 disentangles the systematics of a ubiquitous but elusive meiofaunal group (Kinorhyncha:
1024 Pyncophyidae). *Cladistics* **32**(5), 479–505. <https://doi.org/10.1111/cla.12143>
- 1025 Strong, E. E. and Lipscomb, D. (1999). Character coding and inapplicable data. *Cladistics* **15**,
1026 363–371.
- 1027 Taylor, R. W. (1965). A monographic revision of the rare tropicopolitan ant genus
1028 *Probolomyrmex* Mayr (Hymenoptera: Formicidae). *Transactions of the Royal Entomological*
1029 *Society of London* **117**, 345–365.
- 1030 Taylor, R. W. (1980). Notes on the Russian endemic ant genus *Aulacopone* Arnoldi
1031 (Hymenoptera: Formicidae). *Psyche* **86**(4), 353–361.
- 1032 Ward, P. S. and Sumnicht, T. P. (2012). Molecular and morphological evidence for three
1033 sympatric species of *Leptanilla* (Hymenoptera: Formicidae). *Myrmecological News* **17**, 5–11.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

- 1034 Ward, P. S., Brady, S. G., Fisher, B. L., and Schultz, T. R. (2010). Phylogeny and biogeography
1035 of dolichoderine ants: effects of data partitioning and relict taxa on historical inference.
1036 *Systematic Biology* **59**, 342–362.
- 1037 Wheeler, G. C. and Wheeler, E. W. (1930). Two new ants from Java. *Psyche* **37**(3), 193–201.
- 1038 Wipfler, B., Letsch, H., Frandsen, P. B., Kapli, P., Mayer, C., Buckley, T. R., Donath, A. ...
1039 Simon, S. (2019). Evolutionary history of Polyneoptera and its implications for our
1040 understanding of early winged insects. *Proceedings of the National Academy of Sciences of the*
1041 *United States of America* **116**(8), 3024–3029.
- 1042 Wong, M. K. L. and Guénard, B. (2016). *Leptanilla hypodracos* sp. n., a new species of the
1043 cryptic ant genus *Leptanilla* (Hymenoptera, Formicidae) from Singapore, with new distribution
1044 data and an updated key to Oriental *Leptanilla* species. *ZooKeys* **551**, 129–144.
- 1045 Yamada, A., Nguyen, D. D., and Eguchi, K. (2020). Unveiling the morphology of the Oriental
1046 rare monotypic ant genus *Opamyрма* Yamane, Bui & Eguchi, 2008 (Hymenoptera: Formicidae:
1047 Leptanillinae) and its evolutionary implications, with first descriptions of the male, larva,
1048 tentorium, and sting apparatus. *Myrmecological News* **30**, 27–52.
- 1049 Yang, Z. (1996). Among-site rate variation and its impact on phylogenetic analyses. *Trends in*
1050 *Ecology and Evolution* **11**, 367–372.
- 1051 Yoder, M. J., I. Miko, K. C. Seltmann, M. A. Bertone, and A. R. Deans. (2010). A gross anatomy
1052 ontology for Hymenoptera. *PLOS One* **5**, e15991.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 1. Bayesian total-evidence phylogeny of the Leptanillinae under partitioning scheme inferred with PartitionFinder2 for 9,351-bp legacy-locus alignment. Phylogeny was rooted *a posteriori* on *Martialis heureka*. Nodes with BPP $\geq$ 0.95 marked in red. (A) *Protanilla zhg-vn01* (CASENT0842613); (B) *Yavnella* TH08 (CASENT022755; Shannon Hartman); (C) *Leptanilla zhg-my02* (CASENT0106416).

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 2. Bayesian total-evidence phylogeny of the Leptanillinae under partitioning scheme inferred with PartitionFinder2 for 9,062-bp legacy-locus alignment. Phylogeny was rooted *a posteriori* on *Martialis heureka*. Nodes with BPP $\geq$ 0.95 marked in red.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 3. Selected diversity of male *Leptanilla s. l.* (A) *Leptanilla* TH01 (CASENT0119792; April Nobile); (B) *Leptanilla* zhg-bt02 (CASENT084612; not sequenced in this study); (C) *Noonilla* zhg-my04 (CASENT0842610; not sequenced in this study); (D) *Leptanilla* (Bornean morphospecies-group) zhg-my05 (CASENT0842571). Scale bar A=0.2 mm.; B=1 mm.; C-D=0.5 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 4. Bayesian total-evidence phylogeny of *Leptanilla s. l.* under partitioning scheme inferred with PartitionFinder2 for 9,351-bp legacy-locus alignment. Phylogeny was rooted *a posteriori* on *Martialis heureka*. Nodes with BPP $\geq$ 0.95 marked in red.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1075 Fig. 5. Protibia of *Leptanilla* zhg-my04 (CASENT0842555). Scale bar = 0.2 mm.

1076

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1077 Fig. 6. Presence (B: *Leptanilla* zhg-th01; CASENT0842619) versus absence (A: *Yavnella* zhg-
1078 th01; CASENT0842621) of the posterior prolongation of the mesoscutellum in male Leptanillini.
1079 Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1081 Fig. 7. Conditions of the propodeum in the Leptanillinae. (A) Concave (*Yavnella* zhg-bt01;
1082 CASENT0106384); (B) convex with distinct dorsal face (*Leptanilla* zhg-my02;
1083 CASENT0106456); (C) convex without distinct dorsal face (*Protanilla* lini [OKENT0011097];
1084 male described by Griebenow, in press) (not sequenced in this study).

1085

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1086 Fig. 8. Ventral view of male genitalia across the Leptanillini. (A) *Leptanilla* ZA01
1087 (CASENT0106354), (B) *Noonilla* zhg-my02 (CASENT0842595); (C) *Leptanilla* zhg-my04
1088 (CASENT0842553). Scale bar A = 0.1 mm.; B = 0.3 mm.; and C = 0.5 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1090 Fig. 9. Dorsoventral (A) (*Yavnella* zhg-th01; CASENT0842620) vs. lateromedial (B) (*Leptanilla*
1091 zhg-my04; CASENT0842553) compression of the penial sclerites in posterodorsal view.
1092 Dorsomedian carina marked with arrow. Scale bar = 0.3 mm.

1093

1094

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1095 Fig. 10. Profemur, protibia, and basal protarsomeres of (A) *Leptanilla* TH09 (CASENT0842664)
1096 and (B) *Leptanilla* zhg-bt01. Abbreviations: clc = calcar; pbs = probasitarsal seta. Not to scale.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1098 Fig. 11. Lectotype of *Phaulomyrma javana* as designated by this study (MCZ:Ent:31142). (A)
1099 Full-face view; (B) profile view of mesosoma; (C) forewing; (D) metasoma and genitalia. Scale
1100 bar = 0.2 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 12. Mandible of (A) *Yavnella* cf. *indica* (CASENT0106377) and (B) *Leptanilla* zhg-my03 (CASENT0842618). Scale bar A=0.03 mm.; B=0.05 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1107 Fig. 13. Full-face view of *Yavnella* TH02 (CASENT0119531; Michele Esposito), with ocellar
1108 tubercle marked. Scale bar=0.1 mm.

1109
1110

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1111 Fig. 14. Profile of pronotum (black) and mesoscutum (blue) in male Leptanillini. (A) *Yavnella*
1112 zhg-bt01 (CASENT0106384); (B) *Noonilla* zhg-my04 (CASENT0842610; not sequenced in this
1113 study); (C) *Leptanilla* (Bornean morphospecies-group) zhg-my03 (CASENT0106416).

1114

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1115 Fig. 15. Examples of male forewing venation across the Leptanillinae. (A) *Yavnella* zhg-bt01
1116 (CASENT0106384); (B) *Protanilla* zhg-vn01 (CASENT0842613); (C) *Leptanilla* (Bornean
1117 morphospecies-group) zhg-my05 (CASENT0842571).

1118

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1119 Fig. 16. Foreleg of (A) *Yavnella argamani* (CASENT0235253) and (B) *Leptanilla zhg-id01*
1120 (CASENT0842626; not sequenced in this study). Scale bar=0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1122 Fig. 17. Posterior view of abdominal tergite VIII in male Leptanillini. (A) *Yavnella* zhg-th01
1123 (CASENT0842620) and (B) *Noonilla* zhg-my02 (CASENT0842592). Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1125 Fig. 18. Gonopodite and volsella (*vol*) of *Leptanilla africana*, sketched after Baroni Urbani
 1126 (1977: Fig. 37) by M. K. Lippey. Top of image is distal to body.

1127

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1128 Fig. 19. Dorsoposterior view of phallotreme in (A) *Leptanilla* zhg-my04 (CASENT0842553);
 1129 ventroposterior view of phallotreme in (B) *Leptanilla* zhg-my05 (CASENT0106432). Scale bar
 1130 A = 0.3 mm.; B = 0.4 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 20. Mesal view of protibia of (A) *Protanilla* zhg-vn01 (CASENT0106382) and (B) *Noonilla* zhg-my01 (CASENT0842587; not sequenced in this study). Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 21. Foreleg of (A) *Yavnella argamani* (CASENT0235253) and (B) *Leptanilla zhg-id01* (CASENT0842626; not sequenced in this study). Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 22. Full-face views of (A) *Yavnella* TH08 (CASENT0119531; Michele Esposito) and (B) *Yavnella* TH02 (CASENT0227555; Shannon Hartman). Scale bar = 0.1 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 23. Full-face views of (A) *Leptanilla* zhg-my04 (CASENT0842558) and (B) *Protanilla* TH01 (CASENT0119776; Michele Esposito). Scale bar A = 0.1 mm.; B = 0.2 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1145 Fig. 24. Full-face views of (A) *Leptanilla* zhg-my03 (CASENT084545) and (B) *Protanilla* TH01
1146 (CASENT0119776; Michele Esposito). Scale bar A = 0.1 mm.; B = 0.2 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 25. Full-face views of (A) *Leptanilla* zhg-my04 (CASENT0842548) and (B) *Yavnella* argamani (CASENT0235253; Shannon Hartman). Scale bar A = 0.3 mm.; B = 0.1 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 26. Dorsal view of occipital margin in (A) *Leptanilla* TH09 (CASENT0842664) and (B) *Leptanilla* TH01 (CASENT0119792; April Nobile). Scale bar A = 0.3 mm.; B = 0.2 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 27. Profile view of mesosoma in (A) *Leptanilla* zhg-my02 (CASENT0106416) and (B) *Yavnella* zhg-th01 (CASENT0842621). Scale bar A = 0.5 mm.; B = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 28. Dorsal view of (A) *Leptanilla* TH01 (CASENT0119776; April Nobile) and (B) *Protanilla* TH03 (CASENT0119791; Erin Prado). Scale bar A = 0.2 mm.; B = 1 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1164 Fig. 29. Dorsal view of (A) *Leptanilla* zhg-au01 (CASENT0758873; not sequenced in this study)
1165 and (B) *Leptanilla* zhg-my04 (CASENT0842558). Scale bar A = 1 mm.; B = 0.5 mm.

1166

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 30. Profile view of mesosoma in (A) *Yavnella* TH02 (CASENT0119531; Michele Esposito) and (B) *Protanilla* zhg-vn01 (CASENT0842656). Scale bar A = 0.2 mm.; B = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 31. Forewing of (A) *Phaulomyrma javana* (MCZ:Ent:31142) and (B) *Protanilla zhg-vn01* (CASENT0842613). Scale bar A = 0.2 mm.; B = 0.5 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 32. Dorsal view of mesosoma in (A) *Protanilla* zhg-vn01 (CASENT0842613) and (B) *Leptanilla* zhg-my04 (CASENT0842548). Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 33. Profile view of petiole in (A) *Yavnella* zhg-bt01 (CASENT0106384) and (B) *Protanilla* *lini* (OKENT0011097; male described by Griebenow, in press) (not sequenced in this study). Scale bar A = 0.3 mm.; B = 0.5 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 34. Profile view of petiole in (A) *Leptanilla* zhg-my04 (CASENT0842553), (B) *Yavnella* TH08 (CASENT0227555) and (C) *Leptanilla* zhg-my02 (CASENT0106417). Scale bars A, C = 0.5 mm.; scale bar B = 0.2 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 35. Profile view of petiole in (A) *Yavnella* zhg-bt01 (CASENT0106384) and (B) *Protanilla* *lini* (OKENT0011097; male described by Griebenow, in press) (not sequenced in this study). Scale bar A = 0.3 mm.; B = 0.4 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 36. Profile view of petiole in (A) *Yavnella* zhg-th01 (CASENT0842621) and (B) *Protanilla* *lini* (OKENT0011097; male described by Griebenow, in press) (not sequenced in this study). Scale bar A = 0.3 mm.; B = 0.4 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 37. Profile view of (A) *Protanilla* TH02 (CASENT0128922) and (B) *Protanilla* TH03 (CASENT0119791). Scale bar A = 0.5 mm.; B = 1 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 38. Ventral view of abdominal sternite IX in (A) *Leptanilla* zhg-th01 (CASENT0842619) and (B) *Leptanilla* zhg-my04 (CASENT0842553). Scale bar A = 0.3 mm.; B = 0.2 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 39. Dorsal view of genitalia in (A) *Yavnella* zhg-th01 (CASENT0842620) and (B) *Leptanilla* zhg-my04 (CASENT0842565). Scale bar A = 0.3 mm.; B = 0.4 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 40. Ventral view of genitalia (A) *Protanilla lini* (OKENT0018456; male described by Griebenow, in press) (not sequenced in this study) and (B) *Leptanilla* ZA01 (CASENT0106354). Scale bar A = 0.3 mm.; B = 0.1 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 41. Profile view of genitalia in (A) *Leptanilla* zhg-my04 (CASENT0842558) and (B) *Leptanilla* ZA01 (CASENT0106354). Scale bar A = 0.2 mm.; B = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 42. Profile view of genitalia in (A) *Leptanilla* zhg-my03 (CASENT0842545) and (B) *Leptanilla* zhg-my04 (CASENT0842558). Scale bar = 0.2 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 43. Posterolateral view of gonopodite in (A) *Yavnella argamani* (CASENT0235253) and (B) *Yavnella* TH08 (CASENT0227555) (both images by Shannon Hartman). Scale bar A = 0.2 mm.; B = 0.1 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 44. Posterior view of genitalia in (A) *Yavnella* cf. *indica* (CASENT0106378), (B) *Yavnella* zhg-th01 (CASENT0842620) and (C) *Leptanilla* zhg-my04 (CASENT0842565). Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 45. Profile view of penial sclerites in (A) *Leptanilla* zhg-my04 (CASENT0842550) and (B) *Leptanilla* zhg-my05 (CASENT0842571). Scale bar A = 0.3 mm.; B = 0.2 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 46. Posterior view of penial sclerites in (A) *Leptanilla* zhg-my04 (CASENT0842553) and (B) *Yavnella* zhg-th01 (CASENT0842620). Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 47. Posterior view of penial sclerites in (A) *Leptanilla* zhg-my04 (CASENT0842553) and (B) *Yavnella* zhg-th01 (CASENT0842620). Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 48. Posterior view of penial sclerites in (A) *Yavnella* TH06 (CASENT0129609; Erin Prado) and (B) *Leptanilla* GR02 (CASENT0106068). Scale bar A = 0.1 mm.; B = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 49. Posteroventral view of penial sclerites in (A) *Noonilla* zhg-my02 (CASENT0842595) and (B) *Leptanilla* zhg-my05 (CASENT0106432). Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

1243 Fig. 50. Posterior view of phallotreme in (A) *Leptanilla* zhg-my05 (CASENT0106432) and (B)
1244 *Noonilla* zhg-my02 (CASENT0842599). Scale bar A = 0.4 mm.; B = 0.3 mm.

1245

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Fig. 51. Posterodorsal view of the penial sclerites in (A) *Yavnella argamani* (CASENT0235253) and (B) *Leptanilla* GR02 (CASENT0106068). Scale bar = 0.3 mm.

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Table 1. Summary statistics for full 9,351-bp DNA legacy-locus alignment. “–” = absent base; “?” = unknown base. χ^2 test of nucleotide homogeneity executed with IQ-Tree 1.6.10 (Nguyen *et al.* 2015) on the CIPRES Science Gateway (v. 3.3) (Miller *et al.* 2010).

Taxon	Designation in Borowiec <i>et al.</i> (2019)	Caste/Sex	Identifier	Percent missing	AT content	χ^2 test of nucleotide homogeneity, % <i>p</i> -value	-	?	# of specimens physically examined
<i>Anomalomyrma boltoni</i>	<i>Anomalomyrma boltoni</i>	Worker	CASENT0217032	18.682	0.526	57.33, passed	1747	0	N/A
<i>Leptanilla</i> GR01	<i>Leptanilla</i> GR01	Male	CASENT0106060	19.474	0.536	2.23, failed	1821	0	5
<i>Leptanilla</i> GR02	<i>Leptanilla</i> GR02	Male	CASENT0106236	18.939	0.539	0.95, failed	1771	0	9
<i>Leptanilla</i> GR03	N/A	Male	CASENT0106058	9.817	0.529	10.55, passed	918	0	9
<i>Leptanilla</i> TH01	<i>Leptanilla</i> TH01	Male	CASENT0119792	19.731	0.523	59.55, passed	1845	0	1
<i>Yavnella</i> TH02	<i>Leptanilla</i> TH02	Male	CASENT0119531	18.629	0.512	42.16, passed	1742	0	1
<i>Yavnella</i> TH03	<i>Leptanilla</i> TH03	Male	CASENT0129721	18.747	0.521	73.57, passed	1753	0	1
<i>Yavnella</i> TH04	<i>Leptanilla</i> TH04	Male	CASENT0129695	18.768	0.512	46.39, passed	1755	0	1
<i>Yavnella</i> TH05	<i>Leptanilla</i> TH05	Male	CASENT0134656	18.811	0.516	58.31, passed	1759	0	1
<i>Yavnella</i> TH06	<i>Leptanilla</i> TH06	Male	CASENT0179537	18.918	0.511	41.26, passed	1769	0	1
<i>Yavnella</i> TH08	<i>Leptanilla</i> TH08	Male	CASENT0227775	18.961	0.514	60.05, passed	1773	0	1
<i>Leptanilla</i> TH09	<i>Leptanilla</i> TH09	Male	CASENT0227556	18.822	0.542	0.01, failed	1760	0	1
<i>Leptanilla</i> ZA01	<i>Leptanilla</i> ZA01	Male	CASENT0106354	18.886	0.545	0.01, failed	1766	0	1
<i>Leptanilla</i> zhg-au02	N/A	Male	CASENT0758864	18.682	0.545	0.51, failed	777	4920	1
<i>Leptanilla</i> zhg-th01	N/A	Male	CASENT0842614	19.474	0.552	34.54, passed	518	4337	2
<i>Leptanilla</i> revelierii	N/A	Worker	CASENT0842627	18.939	0.517	25.57, passed	928	5217	4
<i>Leptanilla</i> zhg-bt01	N/A	Male	CASENT0842617	60.924	0.538	0.00, failed	511	4337	1
<i>Leptanilla</i> zhg-my02	N/A	Male	CASENT0106451	51.92	0.532	2.89, failed	1980	2133	49
<i>Leptanilla</i> zhg-my03	N/A	Male	CASENT0842618	65.715	0.526	1.04, failed	1999	2133	4
<i>Leptanilla</i> zhg-my04	N/A	Male	CASENT0842553	51.845	0.518	11.36, passed	1942	2452	21
<i>Leptanilla</i> zhg-my05	N/A	Male	CASENT0842568	43.985	0.474	19.01, passed	1525	0	7
<i>Martialis</i> heureka	<i>Martialis</i> heureka	Worker	CASENT0106181	44.188	0.523	0.00, failed	2216	3297	N/A
<i>Noonilla</i> zhg-my02	N/A	Male	CASENT0842599	16.308	0.477	6.76, passed	1723	0	12
<i>Noonilla</i> zhg-my06	N/A	Male	CASENT0106373	42.552	0.536	14.61, passed	1715	0	3
<i>Opamyrra hungvuong</i>	<i>Opamyrra hungvuong</i>	Worker	CASENT0178347	18.426	0.539	0.00, failed	1717	0	N/A
<i>Yavnella</i> MM01	<i>Phaulomyrma</i> MM01	Male	CASENT0179537	19.014	0.514	72.37, passed	1772	0	1
<i>Phaulomyrma javana</i>	N/A	Male	MCZ:Ent:31142	100	N/A	N/A	N/A	N/A	1
<i>Protanilla</i> TH01	<i>Protanilla</i> TH01	Male	CASENT0119776	18.34	0.529	29.55, passed	1715	0	1
<i>Protanilla</i> TH02	<i>Protanilla</i> TH02	Male	CASENT0128922	18.362	0.529	30.49, passed	1717	0	1
<i>Protanilla</i> TH03	<i>Protanilla</i> TH03	Male	CASENT0119791	18.95	0.497	0.08, failed	1772	0	1
<i>Protanilla</i> zhg-vn01	N/A	Male	CASENT0842613	58.208	0.512	3.27, failed	2146	3297	5
<i>Yavnella</i> argamani	<i>Yavnella</i> argamani	Male	CASENT0235253	18.789	0.52	61.37, passed	1757	0	1
<i>Yavnella</i> cf. <i>indica</i>	N/A	Male	CASENT0106375	52.989	0.502	6.42, passed	2503	2452	8
<i>Yavnella</i> zhg-bt01	N/A	Male	CASENT0842616	49.054	0.514	47.75, passed	2454	2133	5

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Taxon	Designation in Borowiec <i>et al.</i> (2019)	Caste/Sex	Identifier	Percent missing	AT content	χ^2 test of nucleotide homogeneity, % <i>p</i>-value	-	?	# of specimens physically examined
<i>Yavnella</i> zhg-th01	N/A	Male	CASENT0842615	51.706	0.522	7.42, passed	498	4337	2

1253

1254

Phylogeny of the Male-Based Ant Genus *Phaulomyrma*

Table 2. NCBI and SRA accession numbers for DNA sequences used in Bayesian total-evidence inference.

Taxon	CASENT #	SRA	28S	AbdA	EF2	LwRh	Wg	AP	ArgK	NaK	POLD1	Top1	Ubx
<i>Anomalomyrma boltoni</i>	CASENT0217032	SRR11742957	KU671445	KU672069	KU671496	KU671547	KU671598	KU671848	KU671656	KU672002	KU671925	KU671719	KU671782
<i>Leptanilla GR01</i>	CASENT0106236	SRR11881502	EF012999	JN967847	JN967830	JN967890	JN967854	MF625736	JN967880	MF626276	MF625821	JN967820	JN967809
<i>Leptanilla GR02</i>	CASENT0106060	SRR11881501	JN967864	JN967868	JN967831	JN967891	JN967856	MF625737*	JN967883	MF626277*	MF625822*	JN967823	JN967812
<i>Leptanilla GR03</i>	CASENT0106058	SRR11793843	JN967868	JN967851	JN967834	JN967894	JN967859	XXXXXXX	JN967885	MT603718	MT526730	JN967826	JN967815
<i>Leptanilla TH01</i>	CASENT0119792	SRR11881509	KU671447	JN967845	JN967836	KU671549	JN967853	KU671856	KU671660	KU672010	KU671933	KU671723	KU671786
<i>Leptanilla TH09</i>	CASENT0227556	N/A	MF626114	MF625683	MF625896	MF626223	MF626005	MF625744	MF626167	MF626234	MF625829	MF626058	MF625949
<i>Leptanilla ZA01</i>	CASENT0106354	N/A	AY867452	AY867468	EF013432	AY867483	AY867421	MF625745	JN967878	MF626285	MF625830	JN967818	JN967807
<i>Leptanilla zbg-a02</i>	CASENT0758864	SRR11793848	N/A	N/A	N/A	N/A	N/A	XXXXXXX	MT526686	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX
<i>Leptanilla zbg-th01</i>	CASENT0842614	SRR11793854	N/A	N/A	N/A	N/A	N/A	XXXXXXX	MT526687	XXXXXXX	MT526731	XXXXXXX	XXXXXXX
<i>Leptanilla revelieri</i>	CASENT0842627	SRR11881510	XXXXXXX	N/A	N/A	N/A	N/A	XXXXXXX	N/A	MW197678	MW197679	XXXXXXX	XXXXXXX
<i>Leptanilla zbg-b01</i>	CASENT0842617	SRR11793849	XXXXXXX	N/A	N/A	N/A	N/A	XXXXXXX	MT526684	XXXXXXX	MT526729	XXXXXXX	XXXXXXX
<i>Leptanilla zbg-my02</i>	CASENT0106451	SRR11793840	N/A	N/A	N/A	N/A	N/A	XXXXXXX	MT526688	XXXXXXX	MT526732	XXXXXXX	XXXXXXX
<i>Leptanilla zbg-my03</i>	CASENT0842618	SRR11793851	N/A	N/A	N/A	N/A	N/A	XXXXXXX	XXXXXXX	XXXXXXX	MT526734	XXXXXXX	XXXXXXX
<i>Leptanilla zbg-my04</i>	CASENT0842553	SRR11793838	XXXXXXX	N/A	N/A	N/A	N/A	XXXXXXX	XXXXXXX	XXXXXXX	MT526735	XXXXXXX	XXXXXXX
<i>Leptanilla zbg-my05</i>	CASENT0842568	SRR11793837	XXXXXXX	N/A	N/A	N/A	N/A	XXXXXXX	MT526687	XXXXXXX	MT526733	XXXXXXX	XXXXXXX
<i>Martialis heureka</i>	CASENT0106181	SRR11881511	KU671448	KU672072	KU671499	KU671550	KU671601	KU671858	KU671661	KU672012	KU671935	KU671724	KU671787
<i>Noonilla zbg-my02</i>	CASENT0842599	SRR11793856	XXXXXXX	N/A	N/A	N/A	N/A	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX
<i>Noonilla zbg-my06</i>	CASENT0842610	SRR11793842	XXXXXXX	N/A	N/A	N/A	N/A	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX
<i>Opomyrma longicornis</i>	CASENT0178347	SRR11742960	KU671407	KU672031	KU671458	KU671509	KU671560	KU671806	KU671616	KU671960	KU671883	KU671679	KU671742
<i>Proatantilla TH01</i>	CASENT0119776	SRR12006305	MF626129	MF625698	MF625911	MF626238	MF626020	MF625776	MF626191	MF626316	MF625861	MF626082	MF625973
<i>Proatantilla TH02</i>	CASENT0128922	SRR11742959	MF626130	MF625699	MF625912	MF626239	MF626021	MF625777	MF626192	MF626317	MF625862	MF626083	MF625974
<i>Proatantilla TH03</i>	CASENT0119791	SRR11742954	MF626131	MF625700	MF625913	MF626240	MF626022	MF625778	MF626193	MF626318	MF625863	MF626084	MF625975
<i>Yavonella argamasi</i>	CASENT0235253	SRR11793861	KU671449	KU672073	KU671500	KU671551	KU671602	KU671868	KU671665	KU672022	KU671945	KU671728	KU671791
<i>Yavonella cf. indica</i>	CASENT0106375	SRR11793841	XXXXXXX	N/A	N/A	N/A	N/A	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX
<i>Yavonella MM01</i>	CASENT0179537	SRR11742953	MF626120	MF625689	MF625902	MF626229	MF626011	MF625762	MF626178	MF626302	MF625847	MF626049	MF625960
<i>Yavonella TH02</i>	CASENT0119531	SRR11881508	MF626108	MF625677	MF625890	MF626217	MF625999	MF625728	MF626161	MF626278	MF625823	MF626052	MF625941
<i>Yavonella TH03</i>	CASENT0129721	SRR11742956	MF626109	MF625678	MF625891	MF626218	MF626000	MF625739	MF626162	MF626279	MF625824	MF626053	MF625944
<i>Yavonella TH04</i>	CASENT0129695	SRR11742958	MF626110	MF625679	MF625892	MF626219	MF626001	MF625740	MF626163	MF626280	MF625825	MF626054	MF625945
<i>Yavonella TH05</i>	CASENT0134656	SRR11881507	MF626111	MF625680	MF625893	MF626220	MF626002	MF625741	MF626164	MF626281	MF625826	MF626055	MF625946
<i>Yavonella TH06</i>	CASENT0179537	SRR11742955	MF626112	MF625681	MF625894	MF626221	MF626003	MF625742	MF626165	MF626282	MF625827	MF626056	MF625947
<i>Yavonella TH08</i>	CASENT0227775	SRR11881506	MF626113	MF625682	MF625895	MF626222	MF626004	MF625743	MF626166	MF626283	MF625828	MF626057	MF625948
<i>Yavonella zbg-b01</i>	CASENT0842616	SRR11793849	XXXXXXX	N/A	N/A	N/A	N/A	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX
<i>Yavonella zbg-b01</i>	CASENT0842615	SRR11793853	XXXXXXX	N/A	N/A	N/A	N/A	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX	XXXXXXX

*These accession numbers are erroneously attributed to CASENT0106067 (*Leptanilla GR02*) on GenBank.