Negative density-dependent parasitism in a group-living carnivore

Gregory F Albery^{*1}, Chris Newman², Julius Bright Ross², Shweta Bansal¹,
Christina Buesching²

1. Department of Biology, Georgetown University, Washington, DC, USA

2. Wildlife Conservation Research Unit, The Recanati-Kaplan Centre, Department of
Zoology, University of Oxford, Oxford, U.K.

*Corresponding author: gfalbery@gmail.com

Abstract

Animals living at high population densities commonly experience greater exposure to disease, leading to increased parasite burdens. However, social animals can benefit immunologically and hygienically from cooperation, and individuals may alter their socio-spatial behaviour in response to infection, both of which could counteract density-related increases in exposure. Consequently, the costs and benefits of sociality for disease are often uncertain. Here, we use a long-term study of a wild European badger population (*Meles meles*) to investigate how within-population variation in host density determines infection with multiple parasites. Four out of five parasite taxa exhibited consistent spatial hotspots of infection, which peaked among badgers living in areas of low local population density. Combined movement, survival, spatial, and social network analyses revealed that parasite avoidance was the likely cause of this negative density dependence, with possible roles for localised mortality, encounter-dilution effects, and micronutrient-enhanced immunity. These findings demonstrate that animals can organise their societies in space to minimise parasite infection, with important implications for badger behavioural ecology and for the control of badger-associated diseases.

25 Introduction

26 A wild animal's infectious disease burden is determined by its exposure and susceptibility to
 27 infective pathogens. Typically, higher population density results in increased contact rates
 28 and thus greater exposure [1,2]; however, sociality carries immunological and hygienic
 29 benefits that can counteract this exposure-exacerbating effect [3,4]. For instance, cooperative
 30 foraging improves nutrition, alleviating infection costs [5]; mutual grooming removes
 31 ectoparasites [6–8]; and “social immune responses” maintain collective group health [9,10],
 32 e.g. through extirpating sick individuals [11]. Additionally, the environmental distribution of
 33 resources can influence spatial behaviour [12–14] and determine local population densities
 34 [15], while altering susceptibility [16] and transmission efficiency [17]. Sociality and disease
 35 may thus be confounded through shared causal origins rather than being mechanistically
 36 linked [17]. Finally, individuals can minimise exposure by avoiding environmental cues
 37 [18,19] or infected conspecifics [8,20], creating a population-level “landscape of disgust”
 38 analogous to predatory “landscapes of fear” [21,22]. These processes could produce negative
 39 density effects, sometimes depending on parasite transmission mode [1,17,23], but their role
 40 in defining observed density-infection relationships is poorly understood.

41 Uncertainty concerning density-infection relationships largely arises because studies of social
 42 effects are often carried out across discrete groups [1,24], between populations [25,26], or
 43 between species (e.g. [1,2,24,27]). Continuous density measures are rarely linked to parasite
 44 burden within one contiguous host population, and studies that do so generally use purely
 45 social metrics rather than spatial density gradients (e.g. [28–30]). Continuous spatial density
 46 measures are advantageous because: density is a continuous, spatially explicit variable [31–
 47 33]; for the many parasites that are spread through the environment, spatial density measures
 48 better represent transmission than direct (social network) metrics [17,34–36]; and between-
 49 population and cross-species comparisons are fraught with confounding factors such as
 50 shared environmental causality or compensatory evolutionary changes [1,3], which within-
 51 population studies can more easily avoid or overcome. Furthermore, accounting for spatial
 52 effects may help anticipate or test the caveats of cooperation, environmental causality, and
 53 behavioural avoidance outlined above.

54 European badgers (*Meles meles*) are nocturnal, ground-dwelling mustelids with a pan-
 55 European distribution, eating a varied diet composed largely of earthworms [37]. Badgers are
 56 of particular interest for cattle health in the United Kingdom because of their role in

maintaining and spreading bovine tuberculosis [37,38]. Although intended as a control measure, intense culling perturbs badger social systems, causing stressed survivors to disperse, thereby spreading bovine tuberculosis to neighbouring farms [39–41]. Understanding badgers’ socio-spatial behaviours and their epidemiological implications is therefore an important research priority (e.g. [38,42–44]). The demography, behaviour, and parasitism of badgers in Wytham Woods, Oxfordshire have been studied continuously since 1987 [37,45]. Dens (termed “setts”) are situated dependent on soil composition and landscape topography [46]. These badgers reside in cohesive mixed-sex social groups, with around 35% consistent philopatry; however, these groups are also fluid: 16.4% of individuals are trapped at a different social group to their previous capture history, and 19.8% away from their natal group [47]. Furthermore, from genetic pedigree, 48% of cubs have extra-group paternity [48]. Badgers host several arthropod parasites, including badger-specific fleas (*Paraceras melis*) and lice (*Trichodectes melis*) and generalist ticks (*Ixodes* sp.) [49]. They also carry two gastrointestinal protozoans: *Eimeria melis* and *Isospora melis*. *E. melis* predominantly infects young individuals, causing substantial mortality [50,51]. Although studies have examined social grooming effects on ectoparasite burden [6,52] and roles of denning behaviour in parasite transmission [49], the within-population spatial-social parasite epidemiology has yet to be investigated.

Here, we investigate parasite burdens in the Wytham badger population and their associations with spatial and social behaviour. We establish parasite distributions using spatial autocorrelation models. We quantify social drivers using both spatial density gradients and social network metrics, postulating that greater badger densities would drive higher parasite burdens through increased exposure. Finally, we examine survival effects, investigating whether parasite-linked mortality might alter spatial patterns of badger population density. We consider a range of potential social/spatial covariates of parasite burden, including density-related exposure changes, benefits of cohabitation, condition/susceptibility effects, encounter-dilution effects, and parasite avoidance behaviours.

Methods

Data collection

Badgers were sampled as part of a long-term study in Wytham Woods, Oxfordshire, UK (51.778°N, 1.313°W), established in 1987 and recording one of the highest local badger densities ever reported [37,45]. Badgers were trapped overnight at their setts in steel-mesh

89 cages baited with peanuts, collected the next morning, transported to a handling facility and
 90 sedated. Individuals were identifiable by tattoo, applied on first trapping (typically as cubs).
 91 Measures were taken of head-body length (mm) and weight (to the nearest 0.1kg). We
 92 calculated a standardised index of body condition dividing $\log(\text{weight})$ by $\log(\text{body length})$.
 93 The population was trapped seasonally (winter: Jan/Feb/March; spring: April/May/June;
 94 summer: July/Aug/Sept; autumn: Oct/Nov/Dec) in 4 quadrants, for 3 days per quadrant. Our
 95 dataset included 9016 captures of 1369 badgers spanning 29 years (1990-2018).

96 The population currently comprises 23 social groups (mean group size=11; range 2-29
 97 individuals), each using more than one sett, with sett/social group affiliation established using
 98 baitmarking [46,53]. Badgers were assigned to groups using established residency rules [48].
 99 We computed social networks based on co-trapping using a “gambit of the group” approach ,
 100 where individuals trapped in the same sett in the same year were assumed to have interacted
 101 [54].

102 Fleas were counted during a stereotyped 20-second inspection of the badger’s full body, and
 103 ticks (*Ixodes* sp.) and lice (*Trichodectes melis*) were counted within a 4x4cm square of
 104 preferentially-parasitised skin near the groin (per Cox *et al.* 1999). Faecal sampling for two
 105 protozoan endoparasites, *Eimeria melis* and *Isospora melis*, was carried out through 1993-
 106 1997 and 2009-2017 (N=1287 counts). Endoparasite were counted using salt flotation and
 107 microscopy [51,55]. Each count was duplicated, and data were reduced to a binary
 108 infected/uninfected status (rather than counts) because of their highly overdispersed
 109 distribution.

110 **Statistical Analysis**

111 **Covariates**

112 Statistical analysis and data manipulation used R version 3.6.0 [56]. All data and code are
 113 available at <https://github.com/gfalbery/Badgeworks>. Models were constructed using the
 114 `inla` package [57,58]. Response variables included counts of fleas and lice (negative
 115 binomial distribution) and prevalence of ticks, *Eimeria*, and *Isospora*. Explanatory covariates
 116 included: Year (continuous); Month (categorical); Age Category (categorical: cub, yearling,
 117 and adult); Sex (male and female); and Body Condition (continuous). Continuous covariates
 118 were standardised (mean=0; standard deviation=1). Individual Identity and Year were fitted
 119 throughout as categorical random effects.

1. Spatial autocorrelation models

To identify generalised spatial patterns we first fitted models accounting for continuous spatial autocorrelation across the population [33,57,58]. We fitted a base model using only Year, Month, Age Category, and Sex, with individual and annual random effects. We then added an SPDE random effect controlling for spatial autocorrelation in the response variable in space, based on individuals' point locations. We then compared these models using Deviance Information Criterion (DIC) to establish whether accounting for spatial autocorrelation improved model fit; models within $2\Delta\text{DIC}$ were judged competitive.

2. Density models

We then fitted models that replaced the SPDE effects with individual measures of local population density. Models included a range of social/spatial density measures, calculated using various methods. Spatial density metrics represent numbers of individuals in a given space, and therefore tread the line between social and spatial behavioural traits [17]. We computed these by creating space use distribution kernels with the `adehabitatR` package in R [59]. We rasterised the usage distribution, assigning each individual a local density based on the raster value for their map location (Figure 1A-B). Measures included: 1) **Lifetime density**: the density of individuals' centroids across the study period. 2) **Trapping density**: the density of trapping events, incorporating multiple captures of the same individual per year. 3) **Annual density**: the density of individuals' centroids in the sampling year, calculated from annual density kernels. Only one spatial density measure was allowed in a given model, as the measures correlate ($R > 0.5$) and co-fitting several measures produced spurious, unexpected correlations in the opposite direction expected from data exploration. We also used two social metrics based on co-trapping patterns. **Degree centrality** was the number of unique badgers with whom each individual was trapped in the same sett per year. **Group size** was the total number of individuals trapped in a given social group per year. Following spatial autocorrelation model procedures, we conducted a model selection procedure for behavioural metrics, using $2\Delta\text{DIC}$ as the cutoff, and only including the best-fitting metric. We conducted these models for: the overall dataset ($N=9016$); adults only (age 2+, fitting age in years as a fixed covariate; $N=6159$); and juveniles only ($N=1639$). For these models, we display only the best-fitting metric, and for the overall dataset if it exhibited a significant trend (and a subset-only model otherwise).

3. Multi-response models

To investigate whether parasites covaried with badger density at the within- or between-individual level, we constructed multi-response models in MCMCglmm with an unstructured covariance matrix [60,61]. We fitted parasites and local density as response variables, estimating their covariance when accounting for other fixed effects and decomposing this covariance at the within- and between-individual level. A negative within-individual (residual) correlation would imply that higher burden/prevalence individuals (compared to their baseline) moved to lower density areas during their lives, supporting either social ostracism or sickness behaviours. In contrast, negative between-individual covariance would imply that individuals inhabiting lower densities generally had inherently greater parasitism. We only constructed these models for parasite-host age category combinations that demonstrated density effects in the INLA models.

4. Survival models

To investigate parasites' mortality effects, we fitted models with survival as a binary response (1=animal seen in any subsequent year; 0 otherwise). These models included the same covariates as the parasite models, plus parasite prevalence/count and badger density measures. Parasite measures were either log(x+1)-transformed integers (fleas, lice) or binary (ticks, *Eimeria*, and *Isospora*). We conducted another model selection procedure, adding parasite metrics successively if they improved model fit. This was carried out for adults and juveniles separately.

Results

INLA autocorrelation models revealed considerable spatial structuring of parasite burdens. Models for 4/5 parasites were substantially improved by incorporating spatial effects ($\Delta\text{DIC}>8$; Table 1). Only *Isospora*'s model was not improved by spatial autocorrelation (Table 1).

Density models provided substantial support for negative associations between badger density and parasite infection. Including at least one density measure as a covariate improved the model for at least one age category for 4/5 parasites; all had significant negative slopes for density effects (Figure 2; Figure SI1-4; Table SI1-3). Slopes were relatively shallow, but extremely significant and robust (Figure 2). Flea counts were best described by Lifetime Density for the overall model ($\Delta\text{DIC}=-16.31$; $P<0.0001$; Figure 2A), while lice burdens decreased with Trapping Density in the juveniles-only model ($\Delta\text{DIC}=-8.77$; $P=0.0017$; Figure

2B). Tick prevalence was lower in areas of higher Lifetime Density ($\Delta\text{DIC}=-2.47$; $P=0.0038$; Figure 2C), and *Eimeria* prevalence was lower with greater Trapping Density ($\Delta\text{DIC}=-5.83$; $P=0.0053$; Figure 2D). Only one positive social effect was detected: adults with greater degree had slightly more lice ($\Delta\text{DIC}=-2.39$; $P=0.027$; Figure SI2). All DIC changes associated with the model addition procedure are presented in Table SI1-3.

MCMCglmm multi-response models revealed similar trends to INLA univariate models (Figure SI1-4), and allowed us to decompose density-parasite correlations into within- and between-individual changes (Figure 3B). Correlations were greater and much more significant between- than within-individuals in all cases (Figure 3B). This demonstrates that movement between high- and low-density areas was unlikely to produce changes in parasite burden over an individual's lifetime, but rather that individuals' home ranges exhibited inherently different local density-parasite relationships. Only louse infection predicted survival probability; the effect was relatively small, minimally significant, and specific to juveniles ($P=0.038$; Figure 3A).

Our models also revealed various other effects (Figures SI1-4). Briefly, in the overall models, juveniles had fewer fleas, more lice, and greater *Eimeria* prevalence than did yearlings and adults, and lower *Isospora* prevalence than adults (Figure SI1). Males had more lice than did females, with substantial monthly variation in all parasites (Figure SI1). Additionally, body condition was negatively associated with fleas, lice, and *Eimeria* infection in all age/sex classes (Figure SI1). The adults-only and juveniles-only models were very similar to the overall models (Figures SI1-3); notably, in adults, lice burden increased with age in years, whereas *Eimeria* and *Isospora* prevalence decreased (Figure SI2).

Discussion

Using a combination of spatial and social network analysis, we uncovered negative associations between local population density and multiple parasites in this wild group-living carnivore. We found strong but contrasting spatial structuring of fleas, lice, ticks, and *Eimeria*, and, contrary to expectations, all four were most prevalent or abundant in areas of lowest badger density. Co-trapping network and grouping metrics were not predictive of parasitism, implying that "direct" social behaviours, such as mutual allo-grooming, were unlikely to explain the negative density effects. Additionally, badger density effects manifested independent of survival and body condition effects, implying that spatial structuring of the host population did not originate from: 1) localised host mortality, 2)

reduced susceptibility arising from co-habitation and implied cooperation benefits [3,5], or 3) greater local resource availability influencing susceptibility [16,62,63]. Additionally, multi-response models revealed low within-individual covariance between density and parasites, providing little support for heavily-parasitised individuals becoming ostracised during the course of their lives [11]. Taking spatial structuring together with setts' propensity to harbour parasites [49,64], the most parsimonious interpretation is that badgers avoid infection behaviourally, preferring to inhabit areas poorly disposed to parasite transmission (Table 2). These individual-level avoidance behaviours amplify at the population level to produce patterns of badger density inversely related to parasite distributions in the environment, as expected under a "landscape of disgust" [19,21,22]. As well as providing rare evidence of non-consumptive effects of parasites in a wild carnivore, these results imply that animals can arrange their society in space according to parasite transmission.

Our study demonstrates observationally that parasites can determine a society's structure in the environment, and that this phenomenon may counteract the more conventional prediction that host density exacerbates parasitism through increased exposure (e.g. [1,2,65]).

Consequently, studies aiming to quantify social covariates of disease should explicitly seek to investigate individual movement and avoidance behaviours, socio-spatial structuring, and feedbacks between sociality and space use [17]. Previous studies on other mammals have revealed avoidance of infected conspecifics [8] and faeces [18,66], but it remained unclear whether animals avoid spatial hotspots of transmission themselves, and whether these could have population-level consequences [21,22]. Badgers respond to social scent cues and may use these to detect and avoid infested or infectious individuals [67]; furthermore, they move between setts regularly to avoid accumulating parasites [64,68], abandoning highly infested setts and chambers [49]; we posit that this behaviour has emergent population-level consequences. Wytham badgers preferentially establish their setts on northwest-facing slopes in areas with sandy soils [46], and variation in internal sett temperature and humidity are associated with reproductive success [69,70]. Our data suggest additional selection for sett traits and sites resistant to parasite infestation and transmission, which produce an emergent trend for fewer setts, with fewer occupants, in more highly parasitised areas. Quantifying parasites in the environment or inside setts and correlating them with badger behaviour could provide support for this hypothesis. Environmental (sett chamber-based) data on parasite presence would be needed to determine whether badgers avoid infection *actively* (pre-

infection), using environmental cues that predict parasitism, or *reflexively* (post-infection), by moving away from areas of high parasitism due to irritation or sickness.

Four parasites exhibited negative density effects, which could impose tradeoffs on avoidance behaviour. The parasites' distributions were highly contrasting, likely driven by different environmental factors: for example, sett microclimates favouring flea survival will facilitate their transmission [49,64]; similarly, *Eimeria* is transmitted faecal-orally, and oocysts are likely acquired from warm, moist sett chambers, which may explain the gradual decline moving away from the Thames river in the Northwest toward drier parts of the woods (Figure 1F). Ultimately, in combination with other socio-ecological factors (e.g., finding suitable mates), badgers may be incapable of completely avoiding all parasites via denning decisions, which could mediate co-infection and promote diverse parasite communities, giving rise to local hotspots across the population. This may lead to parasite avoidance being traded off against foraging, reproductive success and survival [22,71]. If badgers exhibit between-individual variation in movement and foraging specialisation [72], avoidance of multiple co-infecting parasites could maintain between-individual immune heterogeneity [55]. Notably, only cub density was negatively associated with lice. This observation could be linked to the detectable mortality effects in cubs, driving local cub mortality and/or motivating reproducing females to avoid spatial hotspots of lice.

All of the parasites we examined employ some degree of indirect transmission, likely yielding different relationships with host density than more directly transmitted parasites [1,17]. Ultimately, only avoiding conspecifics can help avoid direct parasite transmission (e.g. [8]), which may not be achievable through purely spatial structuring. Future studies could examine sexually transmitted infections or aerosol-transmitted viruses to investigate whether individuals living in areas of lower density gain any benefit in terms of direct parasite transmission. This may be of particular importance for bovine tuberculosis (bTB), which has a complex, nonlinear relationship with badger sociality [73–75]. Notably, previous bTB studies have generally used social group size as a metric of sociality; given that bTB has an environmental transmission component, particularly between badgers and cattle [76], spatial density metrics, such as those employed here, could be revealing.

We note two potential sources of negative density dependence untested by our modelling approach: the encounter-dilution effect [77,78] and micronutrient impacts on immunity [79]. For the former, where parasites exhibit a constant attack rate in space, greater host densities actually drive a lower per-individual burden of disease because the same number of parasites

is divided among more available hosts [77,78]. Because this requires that the spatial distributions of parasites are not tightly linked to the host distribution (e.g. through reproducing on the host), it is unlikely to apply for any of the badger-specific parasites studied here. However, non-host-specific ticks (*Ixodes* sp.) could transmit from other species to badgers in a given area, producing an encounter-dilution effect; therefore, we are unable to rule out this mechanism (Table 2). Regarding nutritional status, we used body condition index as a coarse predictor of the ability to mount a healthy immune response. However, micronutrients are essential to effective immune function [79] and would not be detected from body condition indices; therefore, it is plausible that badgers might congregate in areas of high micronutrient availability. If this behavioural tendency functions to combat pathogens specifically, it amounts to a “landscape of disgust” acting through reduced susceptibility rather than reduced exposure. This possibility could be tested in this or similar systems by comparing the spatial distributions of high-resolution individual-level immune measures with host density distributions.

Our findings have important implications for the socio-spatial dynamics of this system and its resilience to pathogens and ecological change. If badger populations are organised optimally to occupy areas of least parasite transmission, even small disturbances (e.g. setts lost to forestry) could disrupt its socio-spatial structure and force individuals into unfavourable, more highly parasitised areas, exacerbating their disease burden. Therefore, disruptive anthropogenic activities such as culling-linked perturbation could have unseen consequences for badger disease beyond larger-scale movement impacts (e.g. [39,40]). These findings further inform our understanding of the drivers of badger spatial behaviour, offering insights that may be invaluable for their conservation and disease control [38,46].

Acknowledgements

We would like to thank two anonymous reviewers, and Dan Becker, Amy Sweeny, and Quinn Webber for comments on the manuscript. We also warmly acknowledge the many students and co-workers who helped to collect this substantial data set. GFA and SB were supported by NSF grant number 1414296.

Figure 1: Spatial distributions of badger population density and parasites in Wytham Woods, Oxfordshire, between 1989 and 2018. A: badger population density distributed across Wytham Woods, calculated based on a space use kernel for individuals' annual centroids. B: Individual badgers trapped at setts (represented by points) were assigned a local density value based on their location on the rasterised space use kernel. Darker blue colours in A and B correspond to greater population density. C-F: the spatial distribution of the four spatially distributed parasites, estimated using the INLA SPDE effect. Darker colours correspond to increased parasitism. The density values in B were fitted as covariates in linear models to explain individual parasite burdens, revealing a negative correlation between density and parasitism (see Fig. 2). All axes are in kilometres, with the 0,0 point at the bottom left of the study area. We examined a fifth parasite, *Isospora melis*, but it was not significantly spatially distributed (see Table 1).

Figure 2: Negative associations between population density and parasite infection. Badgers living at higher densities had fewer fleas (A), fewer lice (B), lower tick prevalence (C), and lower *Eimeria melis* prevalence (D). B represents juveniles only. Points represent individual samples, with colours randomised along a colour palette for plotting clarity. Opaque black lines are derived from linear model fits, taking the mean of the posterior distribution. Transparent grey lines represent 100 fits drawn randomly from the posterior estimate distributions of each model, to demonstrate error in the slope and intercepts. Text at the top of the Figures communicates slope estimates, with lower and upper 95% credibility intervals in brackets, and P values. NB: in panel C, the y axis scale and the location of the positive points have been altered for plotting clarity and to better visualise the slope, due to low tick prevalence.

Figure 3: Associations of parasitism with survival and density-parasitism covariance partitioning. A) Louse burden was negatively associated with individual survival probability in juveniles. Opaque black line=linear model fit. Transparent grey lines=100 fits drawn randomly from the posterior estimate distribution, to demonstrate error in the slope and intercept. Text at the top of panel A communicates slope estimate, with lower and upper credibility interval in brackets and P value. B) Estimates for within- and between-individual covariance between parasitism and density, taken from MCMCglmm multivariate models. Points=posterior mean effect size estimates; error bars=95% credibility intervals.

	Fleas	Lice	Ticks	Eimeria	Isospora
Base	153.03	36.57	49.25	8.28	0
SPDE	0	0	0	0	2.25

Table 1: DIC changes associated with spatial autocorrelation terms for the five examined parasites. Lower numbers denote better models; best-fitting models Δ DIC=0. All models except *Isospora* were improved by addition of spatial fields (“SPDE”).

Potential mechanism	Conclusion	Reason for conclusion
Co-habitation/ cooperation benefits	No	No direct social effects or competition with body condition effects (Model Set 2; Figure SI1-3)
Allo-grooming	No	Spatial rather than direct social effects; grooming not possible for endoparasites (Model Set 2; Figure SI1-3; Table SI1-3)
Nutrition-associated immune benefits	No	No competition with body condition effects (Model Set 2; Figure SI1-3)
Social ostracism/self-isolation	No	Low within-individual covariance of density and parasitism (Model Set 3; Figure 3B)
Local host die-offs	~No	Possible for lice, in juveniles, but no other mortality effects were evident (Model Set 4; Figure 3A)
Encounter-Dilution	~No	Possible for generalist ticks, but not for the other (badger-specific, non-mobile) parasites [77,78]
Avoidance	Yes	All other possibilities eliminated; consistent with individual-level behavioural responses [64]

Table 2: Reasoning surrounding our hypothesis testing. We rejected most hypotheses for our observed negative density effects other than parasite avoidance.

References

1. Cote IM, Poulin R. 1995 Parasitism and group-size in social animals: a meta-analysis. *Behav. Ecol.* **6**, 159–165. (doi:10.1093/beheco/6.2.159)
2. Altizer S *et al.* 2003 Social Organization and Parasite Risk in Mammals: Integrating Theory and Empirical Studies. *Annu. Rev. Ecol. Evol. Syst.* **34**, 517–547. (doi:10.1146/annurev.ecolsys.34.030102.151725)
3. Ezenwa VO, Ghai RR, McKay AF, Williams AE. 2016 Group living and pathogen infection revisited. *Curr. Opin. Behav. Sci.* **12**, 66–72. (doi:10.1016/j.cobeha.2016.09.006)
4. Kappeler PM, Cremer S, Nunn CL. 2015 Sociality and health: impacts of sociality on disease susceptibility and transmission in animal and human societies. *Philos. Trans.*

- 367 *R. Soc. B Biol. Sci.* **370**, 20140116–20140116. (doi:10.1098/rstb.2014.0116)
- 368 5. Almberg ES, Cross PC, Dobson AP, Smith DW, Metz MC, Stahler DR, Hudson PJ.
369 2015 Social living mitigates the costs of a chronic illness in a cooperative carnivore.
370 *Ecol. Lett.* **18**, 660–667. (doi:10.1111/ele.12444)
- 371 6. Stewart PD, Macdonald DW. 2003 Badgers and Badger Fleas: Strategies and Counter-
372 Strategies. *Ethology* **109**, 751–764. (doi:10.1046/j.1439-0310.2003.00910.x)
- 373 7. Madden JR, Clutton-Brock TH. 2009 Manipulating grooming by decreasing
374 ectoparasite load causes unpredicted changes in antagonism. *Proc. R. Soc. B Biol. Sci.*
375 **276**, 1263–1268. (doi:10.1098/rspb.2008.1661)
- 376 8. Poirotte C, Massol F, Herbert A, Willaume E, Bomo PM, Kappeler PM, Charpentier
377 MJE. 2017 Mandrills use olfaction to socially avoid parasitized conspecifics. *Sci. Adv.*
378 **3**. (doi:10.1126/sciadv.1601721)
- 379 9. Meunier J. 2015 Social immunity and the evolution of group living in insects. *Philos.*
380 *Trans. R. Soc. B Biol. Sci.* **370**, 19–21. (doi:10.1098/rstb.2014.0102)
- 381 10. Cremer S, Armitage SAO, Schmid-Hempel P. 2007 Social Immunity. *Curr. Biol.* **17**,
382 693–702. (doi:10.1016/j.cub.2007.06.008)
- 383 11. Baracchi D, Fadda A, Turillazzi S. 2012 Evidence for antiseptic behaviour towards
384 sick adult bees in honey bee colonies. *J. Insect Physiol.* **58**, 1589–1596.
385 (doi:10.1016/j.jinsphys.2012.09.014)
- 386 12. Bachorec E *et al.* 2020 Spatial networks differ when food supply changes□: Foraging
387 strategy of Egyptian fruit bats. , 1–15. (doi:10.1371/journal.pone.0229110)
- 388 13. Methion S, Díaz López B. 2019 Individual foraging variation drives social
389 organization in bottlenose dolphins. *Behav. Ecol.* , 1–10. (doi:10.1093/beheco/arz160)
- 390 14. Webber QMR, Laforge MP, Bonar M, Robitaille AL, Hart C, Zabihi-Seissan S, Wal E
391 Vander. 2020 The ecology of individual differences empirically applied to space-use
392 and movement tactics. *The American Naturalist. Am. Nat.* , 1–4. (doi:10.1086/708721)
- 393 15. Webber QMR, Vander Wal E. 2018 An evolutionary framework outlining the
394 integration of individual social and spatial ecology. *J. Anim. Ecol.* **87**, 113–127.
395 (doi:10.1111/1365-2656.12773)
- 396 16. Becker DJ, Albery GF, Kessler MK, Lunn TJ, Falvo CA, Czirják GÁ, Martin LB,

- 397 Plowright RK. 2019 Macroimmunology: the drivers and consequences of spatial
398 patterns in wildlife immune defense. *J. Anim. Ecol.* **89**, 972–995. (doi:10.1111/1365-
399 2656.13166)
- 400 17. Albery GF, Kirkpatrick L, Firth JA, Bansal S. 2020 Unifying spatial and social
401 network analysis in disease ecology. *J. Anim. Ecol.* , 1–17. (doi:10.1111/1365-
402 2656.13356)
- 403 18. Weinstein SB, Moura CW, Mendez JF, Lafferty KD. 2017 Fear of feces? Tradeoffs
404 between disease risk and foraging drive animal activity around raccoon latrines. *Oikos*
405 **00**, 1–8. (doi:10.1111/oik.04866)
- 406 19. Curtis VA. 2014 Infection-avoidance behaviour in humans and other animals. *Trends*
407 *Immunol.* **35**, 457–464. (doi:10.1016/j.it.2014.08.006)
- 408 20. Stockmaier S, Bolnick DI, Page RA, Carter GG. 2018 An immune challenge reduces
409 social grooming in vampire bats. *Anim. Behav.* **140**, 141–149.
410 (doi:10.1016/j.anbehav.2018.04.021)
- 411 21. Weinstein SB, Buck JC, Young HS. 2018 A landscape of disgust. *Science (80-.).* **359**,
412 1213–1215.
- 413 22. Buck JC, Weinstein SB, Young HS. 2018 Ecological and Evolutionary Consequences
414 of Parasite Avoidance. *Trends Ecol. Evol.* **xx**, 1–14. (doi:10.1016/j.tree.2018.05.001)
- 415 23. Hopkins SR, Fleming-Davies AE, Belden LK, Wojdak JM, Fleming□ Davies AE,
416 Belden LK, Wojdak JM. 2020 Systematic review of modeling assumptions and
417 empirical evidence: does parasite transmission increase nonlinearly with host density?
418 *Methods Ecol. Evol.* **11**, 0–2. (doi:10.1111/2041-210x.13361)
- 419 24. Ezenwa VO. 2004 Host social behavior and parasitic infection: A multifactorial
420 approach. *Behav. Ecol.* **15**, 446–454. (doi:10.1093/beheco/arh028)
- 421 25. Downs CJ, Stewart KM, Dick BL. 2015 Investment in constitutive immune function
422 by north American elk experimentally maintained at two different population densities.
423 *PLoS One* **10**, 1–17. (doi:10.1371/journal.pone.0125586)
- 424 26. Brown CR, Brown MB. 1986 Ectoparasitism as a cost of coloniality in cliff swallows (
425 *Hirundo pyrrhonota*). *Ecology* **67**, 1206–1218. (doi:10.2307/1938676)
- 426 27. Rifkin JL, Nunn CL, Garamszegi LZ. 2012 Do Animals Living in Larger Groups

- 427 Experience Greater Parasitism? A Meta-Analysis. *Am. Nat.* **180**, 70–82.
428 (doi:10.1086/666081)
- 429 28. MacIntosh AJJ, Jacobs A, Garcia C, Shimizu K, Mouri K, Huffman MA, Hernandez
430 AD. 2012 Monkeys in the Middle: Parasite Transmission through the Social Network
431 of a Wild Primate. *PLoS One* **7**, 15–21. (doi:10.1371/journal.pone.0051144)
- 432 29. VanderWaal KL, Atwill ER, Isbell LA, McCowan B. 2014 Linking social and
433 pathogen transmission networks using microbial genetics in giraffe (*Giraffa*
434 *camelopardalis*). *J. Anim. Ecol.* **83**, 406–414. (doi:10.1111/1365-2656.12137)
- 435 30. Drewe JA. 2010 Who infects whom? Social networks and tuberculosis transmission in
436 wild meerkats. *Proc. Biol. Sci.* **277**, 633–42. (doi:10.1098/rspb.2009.1775)
- 437 31. Pullan RL, Sturrock HJW, Soares Magalhaes RJ, Clements ACA, Brooker SJ. 2012
438 Spatial parasite ecology and epidemiology: a review of methods and applications.
439 *Parasitology* **139**, 1870–1887. (doi:10.1017/S0031182012000698)
- 440 32. Kirby RS, Delmelle E, Eberth JM. 2017 Advances in spatial epidemiology and
441 geographic information systems. *Ann. Epidemiol.* **27**, 1–9.
442 (doi:10.1016/j.annepidem.2016.12.001)
- 443 33. Albery GF, Becker DJ, Kenyon F, Nussey DH, Pemberton JM. 2019 The fine-scale
444 landscape of immunity and parasitism in a wild ungulate population. *Integr. Comp.*
445 *Biol.* **ic2016**, 1–11. (doi:10.1093/icb/ic2016)
- 446 34. Manlove K, Aiello C, Sah P, Cummins B, Hudson PJ, Cross PC. 2018 The ecology of
447 movement and behaviour: a saturated tripartite network for describing animal contacts.
448 *Proceedings. Biol. Sci.* **285**, 20180670. (doi:10.1098/rspb.2018.0670)
- 449 35. Richardson TO, Gorochowski TE. 2015 Beyond contact-based transmission networks:
450 The role of spatial coincidence. *J. R. Soc. Interface* **12**. (doi:10.1098/rsif.2015.0705)
- 451 36. Leu ST, Sah P, Krzyszczyk E, Jacoby A, Mann J. 2020 Sex, synchrony, and skin
452 contact: integrating multiple behaviors to assess pathogen transmission risk. *Behav.*
453 *Ecol.* , 1–10. (doi:10.1093/beheco/araa002)
- 454 37. Macdonald DW, Newman C, Buesching CD. 2015 Badgers in the rural landscape—
455 conservation paragon or farmland pariah? Lessons from the Wytham Badger Project.
456 In *Wildlife Conservation on Farmland Volume 2*, pp. 65–95.
457 (doi:10.1093/acprof:oso/9780198745501.003.0004)

- 458 38. McDonald JL, Robertson A, Silk MJ. 2018 Wildlife disease ecology from the
459 individual to the population: Insights from a long-term study of a naturally infected
460 European badger population. *J. Anim. Ecol.* **87**, 101–112. (doi:10.1111/1365-
461 2656.12743)
- 462 39. Ham C, Donnelly CA, Astley KL, Jackson SYB, Woodroffe R. 2019 Effect of culling
463 on individual badger *Meles meles* behaviour: Potential implications for bovine
464 tuberculosis transmission. *J. Appl. Ecol.* **56**, 2390–2399. (doi:10.1111/1365-
465 2664.13512)
- 466 40. Tuytens F a M, Delahay RJ, MacDonald DW, Cheeseman CL, Long B, Donnelly C a.
467 2000 Spatial perturbation caused by a badger (*Meles meles*) culling operation:
468 implication for the function of territoriality and the control of bovine tuberculosis
469 (*Mycobacterium bovis*. *J. Appl. Ecol.* **69**, 815–828.
- 470 41. Donnelly CA *et al.* 2007 Impacts of widespread badger culling on cattle tuberculosis:
471 concluding analyses from a large-scale field trial. *Int. J. Infect. Dis.* **11**, 300–8.
472 (doi:10.1016/j.ijid.2007.04.001)
- 473 42. Rozins C, Silk MJ, Croft DP, Delahay RJ, Hodgson DJ, McDonald RA, Weber N,
474 Boots M. 2018 Social structure contains epidemics and regulates individual roles in
475 disease transmission in a group-living mammal. *Ecol. Evol.* , 1–12.
476 (doi:10.1002/ece3.4664)
- 477 43. Silk MJ, Weber NL, Steward LC, Hodgson DJ, Boots M, Croft DP, Delahay RJ,
478 McDonald RA. 2018 Contact networks structured by sex underpin sex-specific
479 epidemiology of infection. *Ecol. Lett.* **21**, 309–318. (doi:10.1111/ele.12898)
- 480 44. Weber N, Carter SP, Dall SRX, Delahay RJ, McDonald JL, Bearhop S, McDonald RA.
481 2013 Badger social networks correlate with tuberculosis infection. *Curr. Biol.* **23**,
482 R915–R916. (doi:10.1016/j.cub.2013.09.011)
- 483 45. Macdonald DW, Newman C. 2002 Population dynamics of badgers (*Meles meles*) in
484 Oxfordshire, U.K.: Numbers, density and cohort life histories, and a possible role of
485 climate change in population growth. *J. Zool.* **256**, 121–138.
486 (doi:10.1017/S0952836902000158)
- 487 46. Macdonald DW, Newman C, Dean J, Buesching CD, Johnson PJ. 2004 The
488 distribution of Eurasian badger, *Meles meles*, setts in a high-density area: Field

- 489 observations contradict the sett dispersion hypothesis. *Oikos* **106**, 295–307.
490 (doi:10.1111/j.0030-1299.2004.12879.x)
- 491 47. Macdonald DW, Newman C, Buesching CD, Johnson PJ. 2008 Male-biased
492 Movement in a High-density Population of the Eurasian Badger (*Meles meles*). *J.*
493 *Mammal.* **89**, 1077–1086. (doi:10.1644/07-mamm-a-185.1)
- 494 48. Annavi G, Newman C, Dugdale HL, Buesching CD, Sin YW, Burke T, Macdonald
495 DW. 2014 Neighbouring-group composition and within-group relatedness drive extra-
496 group paternity rate in the European badger (*Meles meles*). *J. Evol. Biol.* , 2191–2203.
497 (doi:10.1111/jeb.12473)
- 498 49. Cox R, Stewart PD, Macdonald DW. 1999 The ectoparasites of the european badger,
499 *Meles meles*, and the behavior of the host-specific flea, *Paraceras melis*. *J. Insect*
500 *Behav.* **12**, 245–265.
- 501 50. Anwar MA, Newman C, Macdonald DW, Woolhouse MEJ, Kelly DW. 2000
502 Coccidiosis in the European badger (*Meles meles*) from England, an epidemiological
503 study. *Parasitology* **120**, 255–260. (doi:10.1017/S0031182099005491)
- 504 51. Newman C, Macdonald DW, Anwar MA. 2001 Coccidiosis in the European badger,
505 *Meles meles* in Wytham Woods: Infection and consequences for growth and survival.
506 *Parasitology* **123**, 133–142. (doi:10.1017/S0031182001008265)
- 507 52. Johnson DDP, Stopka P, Macdonald DW. 2004 Ideal flea constraints on group living:
508 Unwanted public goods and the emergence of cooperation. *Behav. Ecol.* **15**, 181–186.
509 (doi:10.1093/beheco/arg093)
- 510 53. Buesching CD, Newman C, Service K, Macdonald DW, Riordan P. 2016 Latrine
511 marking patterns of badgers (*Meles meles*) with respect to population density and
512 range size. **7**, 1–11.
- 513 54. Franks DW, Ruxton GD, James R. 2010 Sampling animal association networks with
514 the gambit of the group. *Behav. Ecol. Sociobiol.* **64**, 493–503. (doi:10.1007/s00265-
515 009-0865-8)
- 516 55. Sin YW, Annavi G, Dugdale HL, Newman C, Burke T, MacDonald DW. 2014
517 Pathogen burden, co-infection and major histocompatibility complex variability in the
518 European badger (*Meles meles*). *Mol. Ecol.* **23**, 5072–5088. (doi:10.1111/mec.12917)
- 519 56. R Core Team. 2020 R: A language and environment for statistical computing. R

- 520 Foundation for Statistical Computing, Vienna, Austria.
- 521 57. Lindgren F, Rue H. 2015 Bayesian Spatial Modelling with R-INLA. *J. Stat. Softw.* **63**,
522 1–25. (doi:10.18637/jss.v063.i19)
- 523 58. Lindgren F, Rue H, Lindstrom J. 2011 An explicit link between Gaussian fields and
524 Gaussian Markov random fields: the stochastic partial differential equation approach.
525 *J. R. Stat. Soc. B* **73**, 423–498.
- 526 59. Calenge C. 2011 Home range estimation in R: the adehabitatHR package. See
527 <https://cran.r-project.org/web/packages/adehabitatHR/index.html> (accessed on 10
528 March 2020).
- 529 60. Hadfield JD. 2010 MCMC methods for multi-response generalized linear mixed
530 models: the MCMCglmm R package. *J. Stat. Softw.* **33**, 1–22. (doi:10.1002/ana.22635)
- 531 61. Brommer JE, Covarrubias-Pazaran G. 2019 Multivariate mixed models in ecology and
532 evolutionary biology: inference and implementation in R. *EcoEvoRxiv Prepr.*
533 (doi:10.1017/CBO9781107415324.004)
- 534 62. Strandin T, Babayan SA, Forbes KM. 2018 Reviewing the effects of food provisioning
535 on wildlife immunity. *Philos. Trans. R. Soc. B Biol. Sci.* **373**.
536 (doi:10.1098/rstb.2017.0088)
- 537 63. Sweeny AR, Clerc M, Pontifes PA, Venkatesan S, Babayan SA, Pedersen AB. 2019
538 Supplemented nutrition increases immunity and drug efficacy in a natural host-
539 helminth system. , 1–43.
- 540 64. Butler JM, Roper TJ. 1996 Ectoparasites and sett use in European badgers. *Anim.*
541 *Behav.* **52**, 621–629. (doi:10.1006/anbe.1996.0203)
- 542 65. Ezenwa VO. 2016 Helminth-microparasite co-infection in wildlife: lessons from
543 ruminants, rodents, and rabbits. *Parasite Immunol.* **38**, 527–534.
544 (doi:10.1111/pim.12348)
- 545 66. Poirotte C, Kappeler PM. 2019 Hygienic personalities in wild grey mouse lemurs vary
546 adaptively with sex. *Proc. R. Soc. B Biol. Sci.* **286**, 20190863.
547 (doi:10.1098/rspb.2019.0863)
- 548 67. Newman C, Buesching CD. 2019 Detecting the Smell of Disease and Injury: Scoping
549 Evolutionary and Ecological Implications. In *Chemical Signals in Vertebrates 14*, pp.

- 238–250. Cham: Springer International Publishing. (doi:10.1007/978-3-030-17616-7_17)
68. Brøseth H, Bevanger K, Knutsen B. 1997 Function of multiple badger *Meles meles* setts: distribution and utilisation . *Wildlife Biol.* **3**, 89–96. (doi:10.2981/wlb.1997.011)
69. Tsunoda M, Newman C, Buesching CD, Macdonald DW, Kaneko Y. 2018 Badger setts provide thermal refugia, buffering changeable surface weather conditions. *J. Therm. Biol.* **74**, 226–233. (doi:10.1016/j.jtherbio.2018.04.005)
70. Kaneko Y, Newman C, Buesching CD, MacDonald DW. 2010 Variations in badger (*meles meles*) sett microclimate: Differential cub survival between main and subsidiary setts, with implications for artificial sett construction. *Int. J. Ecol.* **2010**. (doi:10.1155/2010/859586)
71. Hutchings MR, Judge J, Gordon IJ, Athanasiadou S, Kyriazakis I. 2006 Use of trade-off theory to advance understanding of herbivore-parasite interactions. *Mamm. Rev.* **36**, 1–16. (doi:10.1111/j.1365-2907.2006.00080.x)
72. Robertson A, McDonald RA, Delahay RJ, Kelly SD, Bearhop S. 2014 Individual foraging specialisation in a social mammal: the European badger (*Meles meles*). *Oecologia* **176**, 409–421. (doi:10.1007/s00442-014-3019-2)
73. Benton CH, Delahay RJ, Robertson A, McDonald RA, Wilson AJ, Burke TA, Hodgson D. 2016 Blood thicker than water: Kinship, disease prevalence and group size drive divergent patterns of infection risk in a social mammal. *Proc. R. Soc. B Biol. Sci.* **283**. (doi:10.1098/rspb.2016.0798)
74. Delahay RJ, Walker N, Smith GS, Wilkinson D, Clifton-Hadley RS, Cheeseman CL, Tomlinson AJ, Chambers MA. 2013 Long-term temporal trends and estimated transmission rates for *Mycobacterium bovis* infection in an undisturbed high-density badger (*Meles meles*) population. *Epidemiol. Infect.* **141**, 1445–1456. (doi:10.1017/S0950268813000721)
75. Woodroffe R *et al.* 2009 Social group size affects *Mycobacterium bovis* infection in European badgers (*Meles meles*). *J. Anim. Ecol.* **78**, 818–827. (doi:10.1111/j.1365-2656.2009.01545.x)
76. Woodroffe R, Donnelly CA, Ham C, Jackson SYB, Moyes K. 2016 Badgers prefer

cattle pasture but avoid cattle□: implications for bovine tuberculosis control. *Ecol. Lett.* **19**, 1201–1208. (doi:10.1111/ele.12654)

77. Buck JC *et al.* 2017 Host density increases parasite recruitment but decreases host risk in a snail — trematode system Published by□: Wiley on behalf of the Ecological Society of America Stable URL□: <http://www.jstor.com/stable/26601143> Host density increases parasite recruit. *Ecology* **98**, 2029–2038.

78. Mooring MS, Hart BL. 1992 Animal Grouping for Protection from Parasites: Selfish Herd and Encounter-Dilution Effects. *Behaviour* **123**, 173–193. (doi:10.2307/4535069)

79. Calder PC, Jackson AA. 2000 Undernutrition, infection and immune function. *Nutr. Res. Rev.* **13**, 3–29. (doi:10.1079/095442200108728981)

Supplementary figures and tables: Negative density-dependent parasitism in a group-living carnivore

Figure SI1: Model effect outputs for the full dataset INLA density models. Points represent modes of the posterior distribution of the effect estimates; error bars represent the 95% credibility intervals. Asterisks denote significant results, whose 95% credibility intervals did not overlap with zero. Only the behavioural traits that improved the model fit ($\Delta\text{DIC} < -2$) and were kept in the final model are shown.

604

605 Figure SI2: Model effect outputs for the adults-only dataset INLA density models. Points represent
606 modes of the posterior distribution of the effect estimates; error bars represent the 95% credibility
607 intervals. Asterisks denote significant results, whose 95% credibility intervals did not overlap with
608 zero. Only the behavioural traits that improved the model fit ($\Delta\text{DIC} < -2$) and were kept in the final
609 model are shown.

610

611 Figure SI3: Model effect outputs for the juveniles-only dataset INLA density models. Points represent
612 modes of the posterior distribution of the effect estimates; error bars represent the 95% credibility
613 intervals. Asterisks denote significant results, whose 95% credibility intervals did not overlap with
614 zero. Only the behavioural traits that improved the model fit ($\Delta\text{DIC} < -2$) and were kept in the final
615 model are shown.

616

617 Figure SI4: Model effect outputs for the multivariate MCMCglmm models closely represent the INLA
 618 univariate models. Points represent modes of the posterior distribution of the effect estimates; error
 619 bars represent the 95% credibility intervals. Asterisks denote significant results, whose 95%
 620 credibility intervals did not overlap with zero. Only the behavioural traits that improved the model fit
 621 ($\Delta\text{DIC} < -2$) and were kept in the final model are shown.

622

623

PARASITE	ROUND	VARIABLE	DELTA	KEPT
FLEAS	1	Degree	2.08	0
	1	GroupSize	2.055	0
	1	**LifetimeDensity**	-16.001	1
	1	TrappingDensity	1.115	0
	1	AnnualDensity	-6.889	0
LICE	1	Degree	1.796	0
	1	GroupSize	1.3	0
	1	LifetimeDensity	1.694	0
	1	TrappingDensity	0.166	0
	1	AnnualDensity	1.117	0
TICKS	1	Degree	0.773	0
	1	GroupSize	3.644	0
	1	**LifetimeDensity**	-2.192	1
	1	TrappingDensity	1.007	0
	1	AnnualDensity	-0.266	0
EIMERIA	1	Degree	1.134	0
	1	GroupSize	1.79	0
	1	LifetimeDensity	1.084	0
	1	**TrappingDensity**	-6.035	1
	1	AnnualDensity	1.851	0
ISOSPORA	1	Degree	1.897	0
	1	GroupSize	1.861	0

	1	LifetimeDensity	1.873	0
	1	TrappingDensity	1.507	0
	1	AnnualDensity	1.717	0

624

625 Table SII: DIC Changes (Δ DIC) associated with density measure model addition for the full
626 models. Negative Δ DIC values correspond to increased model fit. In each round, the variable
627 that increased model the most was retained, and then the process was repeated, until no
628 variables improved the model by decreasing Δ DIC by more than -2. Variables that were
629 retained in each round are highlighted by asterisks.

630

631

PARASITE	ROUND	VARIABLE	DELTA	KEPT
FLEAS	1	Degree	1.835	0
	1	GroupSize	1.251	0
	1	**LifetimeDensity**	-8.444	1
	1	TrappingDensity	1.509	0
	1	AnnualDensity	-4.549	0
LICE	1	**Degree**	-2.388	1
	1	GroupSize	1.747	0
	1	LifetimeDensity	1.332	0
	1	TrappingDensity	1.943	0
	1	AnnualDensity	0.944	0
TICKS	1	Degree	1.93	0
	1	GroupSize	-12.573	0
	1	LifetimeDensity	-5.864	0
	1	TrappingDensity	1.2	0
	1	**AnnualDensity**	-14.067	1
EIMERIA	1	Degree	14.117	0
	1	GroupSize	1.964	0
	1	LifetimeDensity	1.817	0
	1	TrappingDensity	-0.283	0
	1	AnnualDensity	1.237	0
ISOSPORA	1	Degree	1.569	0
	1	GroupSize	0.882	0

	1	LifetimeDensity	1.9	0
	1	TrappingDensity	1.65	0
	1	AnnualDensity	1.8	0

632

633 Table SI2: DIC Changes (Δ DIC) associated with density measure model addition for the
634 adults-only models. Negative Δ DIC values correspond to increased model fit. In each round,
635 the variable that increased model the most was retained, and then the process was repeated,
636 until no variables improved the model by decreasing Δ DIC by more than -2. Variables that
637 were retained in each round are highlighted by asterisks.

638

639

PARASITE	ROUND	VARIABLE	DELTA	KEPT
FLEAS	1	Degree	0.985	0
	1	GroupSize	-2.362	0
	1	**LifetimeDensity**	-15.209	1
	1	TrappingDensity	1.337	0
	1	AnnualDensity	-10.016	0
LICE	1	Degree	-2.862	0
	1	GroupSize	1.747	0
	1	LifetimeDensity	-3.907	0
	1	**TrappingDensity**	-8.087	1
	1	AnnualDensity	-2.572	0
TICKS	1	Degree	1.602	0
	1	GroupSize	0.462	0
	1	LifetimeDensity	-0.244	0
	1	TrappingDensity	0.918	0
	1	AnnualDensity	-0.459	0
EIMERIA	1	Degree	1.42	0
	1	GroupSize	1.067	0
	1	LifetimeDensity	-0.127	0
	1	**TrappingDensity**	-7.465	1
	1	AnnualDensity	-0.945	0
ISOSPORA	1	Degree	1.73	0
	1	GroupSize	1.096	0

	1	LifetimeDensity	1.775	0
	1	TrappingDensity	1.567	0
	1	AnnualDensity	1.709	0

640

641 Table SI3: DIC Changes (Δ DIC) associated with density measure model addition for the
642 juvenile-only models. Negative Δ DIC values correspond to increased model fit. In each
643 round, the variable that increased model the most was retained, and then the process was
644 repeated, until no variables improved the model by decreasing Δ DIC by more than -2.
645 Variables that were retained in each round are highlighted by asterisks.

646

647