Title: Genome-wide analysis of long noncoding RNAs, 24-nt siRNAs, DNA methylation and H3K27me3 marks in *Brassica rapa* and related species

Hasan Mehraj¹, Daniel J. Shea², Satoshi Takahashi³, Naomi Miyaji¹, Ayasha Akter¹, Motoaki Seki^{3,4,5}, Elizabeth S. Dennis^{6,7}, Fujimoto Ryo¹, Kenji Osabe^{8*}

Affiliation:

¹Graduate School of Agricultural Science, Kobe University, Rokkodai, Nada-ku, Kobe 657-8501, Japan

² Bloomberg LP, Chiyoda-ku, Tokyo 100-0005, Japan

³Riken center for Sustainable Resource Science, Yokohama, Kanagawa, 230-0045, Japan

⁴RIKEN Cluster for Pioneering Research, 2-1 Hirosawa, Wako, Saitama, 351-0198, Japan

⁵Kihara Institute for Biological Research, Yokohama City University, 641-12 Maioka-cho, Totsuka-ku, Yokohama, Kanagawa 244-0813, Japan

⁶CSIRO Agriculture and Food, Canberra, ACT 2601, Australia

⁷ University of Technology, Sydney, PO Box 123, Broadway, NSW 2007, Australia

⁸Institute of Scientific and Industrial Research, Osaka University, Mihogaoka, Ibaraki-city, Osaka, Japan

*Corresponding author: o3b@sanken.osaka-u.ac.jp

Abstract

Long noncoding RNAs (lncRNAs) are RNA fragments that generally do not code for a protein but are involved in epigenetic gene regulation. In this study, lncRNAs of *Brassica rapa* were classified into long intergenic noncoding RNAs, natural antisense RNAs, and intronic noncoding RNAs and their expression were analyzed in relation to genome-wide 24-nt small interfering RNAs (siRNAs), DNA methylation, and histone H3 lysine 27 trimethylation marks (H3K27me3). More than 65% of the lncRNAs analyzed consisted of one exon, and more than 55% overlapped with inverted repeat regions (IRRs). Overlap of lncRNAs with IRRs or genomic regions encoding for 24-nt siRNAs resulted in increased DNA methylation levels and when both were present, there were further increase in DNA methylation levels. LncRNA did not overlap greatly with H3K27me3 marks, but the expression level of intronic noncoding RNAs that did coincide with H3K27me3 marks was higher than without H3K27me3 marks. The *Brassica* genus comprises important vegetables and oil seed crops grown across the world. *Brassica rapa* is a diploid (AA genome) thought to be one of the ancestral species of both *B. juncea* (AABB genome) and *B. napus* (AACC) through genome merging (allotetrapolyploidization). Complex genome restructuring and epigenetic alterations are thought to be involved in these allotetrapolyploidization events, but the detailed mechanism is not known. Comparison of lncRNAs between *B. rapa* and *B. nigra*, *B. oleracea*, *B. juncea*, and *B. napus* showed the highest conservation with *B.*

oleracea. This study presents a comprehensive analysis of the epigenome structure of the *B. rapa* at multi-epigenetic levels (siRNAs, DNA methylation, H3K27me3, and lncRNAs) and offers insights into the function of lncRNA in the Brassica genus.

Key words: Brassica rapa, noncoding RNA, siRNA, DNA methylation, H3K27 methylation, epigenome, Brassica nigra, Brassica oleracea, Brassica juncea, Brassica napus

Introduction

The Brassica genus comprises vegetable and oil seed crops. The “Triangle of U” proposed the genomic relationship among six major species of the Brassica genus. Three allotetraploid species, each of which contains two complete diploid genomes derived from two different parental species, *Brassica juncea* L. (AABB genome, $2n = 4x = 36$), *B. napus* L. (AACC, $2n = 4x = 38$), and *B. carinata* L. (BBCC, $2n = 4x = 34$) are derived from the natural hybridization of the diploid species, *B. rapa* L. (AA, $2n = 2x = 20$), *B. nigra* L. (BB, $2n = 2x = 16$), and *B. oleracea* L. (CC, $2n = 2x = 18$) [1]. Some species in the Brassica genus show morphological divergence (termed morphotype). *B. rapa* includes leafy vegetables including Chinese cabbage (var. *pekinensis*), pak choi (var. *chinensis*), and komatsuna (var. *perviridis*), root vegetables including turnip (var. *rapa*), and oilseed crops (var. *oleifera*) [2]. The first whole

genome sequence determined in the genus Brassica was that of *B. rapa* [3]. Later the whole genome sequences of *B. oleracea*, *B. nigra*, *B. napus*, and *B. juncea* were determined [4–7].

Plant transcriptome analyses have revealed transcribed RNAs devoid of protein-coding potential, which are called noncoding RNAs (ncRNAs). Some ncRNAs contain exons, which potentially code for short proteins or peptides, however, experimental investigation is required to validate their functions. Two families of ncRNAs are known; long noncoding RNAs (lncRNAs) that are longer than 200 nucleotides (nt), and small RNAs (sRNAs) that are ~18-30 nt in length [8–12]. LncRNAs are classified by their position and orientation of transcription; long intergenic noncoding RNAs (lincRNAs), intronic noncoding RNAs (incRNAs) derived from introns, and natural antisense transcripts (NATs) transcribed from the complementary DNA strand of their associated genes [8-12]. Recent studies have shown that lncRNAs play crucial roles in various physiological processes such as vernalization [13-15], photoperiod-sensitive male sterility [16], red-light-mediated seedling photomorphogenesis [17], seed dormancy [18], and the transcriptional regulation of plant innate immunity [19].

Epigenetics has been defined as “the study of changes in genome expression that are mitotically and/or meiotically heritable and that do not entail a change in DNA sequence” [20]. DNA methylation and histone modification are well-known epigenetic

modifications, and lncRNAs are considered to be involved in epigenetic regulation [21]. In plants, DNA methylation is established through RNA-directed DNA methylation (RdDM) [22,23]. Plant-specific RNA POLYMERASE IV (Pol IV) and Pol V are involved in RdDM. Pol IV-transcribed ncRNAs are cleaved into 24-nucleotide small interfering RNAs (24-nt siRNAs) by DICER-LIKE 3 (DCL3), and 24-nt siRNAs are loaded into ARGONAUTE 4 (AGO4). PolV-transcribed lncRNAs act as scaffold molecules of AGO-siRNA complexes, which recruit DOMAINS REARRANGED METHYLTRANSFERASE 2 (DRM2) for catalyzing *de novo* DNA methylation. In *Arabidopsis thaliana* two cold-induced lncRNAs (COLDAIR and COLDWRAP) regulate histone methylation by recruiting polycomb repressive complex 2 (PRC2), which catalyzes the tri-methylation of histone H3 lysine 27 (H3K27me3), to the chromatin region of *FLOWERING LOCUS C* (*FLC*) [14,24-26].

Previous studies have identified lncRNAs associated with different lines, tissues or environmental changes in Brassica [27–33]. However, there are few reports of the association between lncRNAs and epigenetic modifications in *B. rapa*. In this study, we examined the DNA methylation levels and H3K27me3 levels in the region covering lncRNAs, and found an association between lncRNAs and inverted repeat regions (IRRs) or 24-nt siRNAs, but association with H3K27me3 to a specific type of lncRNA.

Materials and methods

Plant materials and growth conditions

Six lines of *B. rapa*, inbred lines of RJKB-T24 (var. *pekinensis*) [34] and Yellow sarson (var. *trilocularis*), doubled haploid lines of BRA2209 (var. *rapa*), Homei (var. *pekinensis*), and Osome (var. *perviridis*), and a commercial F₁ hybrid cultivar, ‘Harunosaiten’ (var. *pekinensis*) (Watanabe Seed Co., Ltd.) were used. Three *B. oleracea* cabbage (var. *capitata*) F₁ hybrid cultivars, ‘Reiho’, ‘Matsunami’ (Ishii Seed Growers CO., LTD), and ‘Kinkei 201’ (Sakata Seed Co., Ltd.) were also used.

Seeds were surface sterilized and grown on agar solidified Murashige and Skoog (MS) plates with 1 % (w/v) sucrose under long day (LD) condition (16h light / 8h dark) at 21 °C. Fourteen-day first and second leaves of *B. rapa* and 19-day first and second leaves of *B. oleracea* were harvested for isolation of genomic DNA or total RNA.

RJKB-T24 was used for RNA-sequencing (RNA-seq) for detection of lncRNAs. Briefly, after performing QC of the sequenced reads, putative mRNAs were identified by aligning the sequence reads to the *B. rapa* reference genome v1.5 using HISAT2 and then assembling transcripts with Stringtie. Assembled transcripts with a mapping code of 'u', indicating they are intergenic but not part of the annotated reference genome, were then compared to the SwissProt database using blastx (e-value 1e-10). Transcripts with hits were classified as putative mRNAs, while transcripts with

no hits were classified as putative lincRNAs. A more detailed description of this method can be found in Shea et. al, 2018 [27].

DNA extraction and PCR

Genomic DNA was isolated by the Cetyl trimethyl ammonium bromide (CTAB) method [35]. The PCR reaction was performed using the following conditions; 1 cycle of 94 °C for 3 min, 35 cycles of 94 °C for 30 s, 55 °C for 30 s, and 72 °C for 1 min, and final extension at 72 °C for 3 min. The PCR products were electrophoresed on 1.0% agarose gel. Primer sequences used in this study are shown in S1 Table.

RNA extraction and RT-PCR

Total RNA from the first and second leaves were isolated by SV Total RNA Isolation System (Promega Co., WI, USA). To analyze lincRNA expression, cDNA was synthesized from 500 ng total RNA using PrimeScript RT reagent Kit (Takara Bio., Shiga, JAPAN). The absence of genomic DNA contamination was confirmed by PCR using a control without reverse transcriptase. The PCR conditions were 94 °C for 2 min followed by 35 cycles of 94 °C for 30 s, 55 °C for 30 s, and 68 °C for 30 s. The primers used in this study are listed in S1 Table.

Detection of epigenetic states in lincRNA regions

To examine the epigenetic states (DNA methylation levels, H3K27me3, and 24-nt siRNA levels) of lncRNA encoding regions in *B. rapa*, we used previous sequence data of whole genome bisulfite sequencing (WGBS) [36], chromatin immunoprecipitation sequencing (ChIP-seq) [37], and small RNA-sequencing (sRNA-seq) data [36], which were generated using samples from the same line, tissue, and developmental stages but harvested independently.

The reads of WGBS were mapped to the *B. rapa* reference genome v.1.5 using Bowtie2 version 2.2.3 and Bismark v0.14.3, and data covering genomic regions encoding lncRNA in chromosomes A01 to A10 were extracted. In order to estimate the methylation levels of CG, CHG, and CHH contexts, the numbers of methylated and unmethylated reads were extracted for each cytosine position using bismark methylation extractor script with the paired-end parameter. The methylation level at each cytosine position was calculated by dividing the number of methylated cytosines (mC) reads by the total number of reads.

The reads of sRNA-seq were mapped to the *B. rapa* reference genome v.1.5 using Bowtie2 version 2.2.3. We classified the alignment reads by length, and the 24-nt aligned reads covering the lncRNA regions in chromosomes A01 to A10 were extracted.

The reads of ChIP-seq using anti-H3K27me3 (Millipore, 07-449) antibodies were mapped to the *B. rapa* reference genome v.1.5 using Bowtie2 version 2.2.3, and

data covering genomic regions encoding for lncRNA in chromosomes A01 to A10 were extracted.

Comparison of putative mRNA and lncRNAs in *B. rapa* to other related species of *Brassica*

The putative mRNAs and lncRNAs were first compared to the reference genomes of *B. nigra*, *B. oleracea*, *B. juncea*, and *B. napus* by best-hit blastn (e-value 1e-10) to identify homologous regions in closely related Brassica species [6,7,38]. In order to parse the local High-scoring segment pair (HSP) alignments produced by blastn, genBlastA was used to produce a representative putative gene that is homologous to the query [39]. The parsed local alignments were then analyzed by a custom python script (available at <http://www.github.com/danshea/lncRNA>) to examine the overall alignment length and computed coverage of the aligned homologous region to the putative mRNA or lncRNA query transcript sequence. These results were then imported into R and plotted to assess overall relative coverage of the sequences among the Brassica species.

Results

Characterization of lncRNA in *B. rapa*

We identified 1,444 lincRNAs, 551 NATs, and 93 incRNAs using RNA-seq data of 14-day first and second leaves with and without four weeks of cold treatment

[27]. In order to investigate the relationship between lncRNAs and epigenetic modifications or the species specificity of lncRNAs, we analyzed in more detail the RNA-seq data of the 14-day first and second leaves without cold treatment. There was no strong bias in the chromosomal distribution of expressed lncRNAs (S1 Fig). More than 65% of lncRNAs contained one exon (lincRNAs, 65.7%; NATs, 72.4%; incRNAs, 71.0%), whereas the proportion of mRNAs containing only one exon is 15.1% (Fig 1A). The mean transcript lengths of lincRNAs (725 nt) and incRNAs (779 nt) were shorter than that of NATs (1,271 nt) and mRNAs (1,305 nt, Fig 1B). About 40% of lincRNAs were located within 2 kb of the genic regions, and about 10% of lincRNAs were located more than 20kb from the genic region (Fig 1C). In this study, we focused on lncRNAs mapped to the chromosomes A01 to A10 as previous WGBS, ChIP-seq, and sRNA-seq have omitted the placed scaffolds for their analyses [36,37]. 763 of 1,173 (65.0%) lincRNAs, 291 of 529 (55.0%) NATs, and 66 of 92 (71.7%) incRNAs overlapped with IRRs such as transposable elements (TEs) detected by RepeatMasker, suggesting that IRRs are the source of more than half of lncRNAs in *B. rapa* (Fig 2).

Fig 1. Analysis of lncRNAs of 14-day first and second leaves of *B. rapa*. (A) Number of exons in lncRNA. (B) Nucleotide length of lncRNA, NAT, incRNA, putative mRNA and mRNA. “a” and “b” represent significant differences by one-way ANOVA test (“*”, $p < 0.05$; “***”, $p < 0.001$). (C) The proportion lncRNA distances to the nearest gene.

Fig 2. Proportion of each type of lncRNA and putative mRNAs that overlap with IRRs and siRNA. “+” indicates with and “-” indicated without overlapping IRRs or genomic regions encoding siRNAs.

Relationship between lncRNAs and siRNA or DNA methylation

We have performed sRNA-seq using 14-day first and second leaves, which are identical developmental stages and tissues to previously analyzed samples [36], but independently harvested for this study. We identified the lncRNAs having perfect sequence identity to genomic regions encoding for 24-nt siRNAs. 219 of 1,173 (18.7 %) lincRNAs, 74 of 529 (14.0%) NATs, and 16 of 92 (17.4%) incRNAs in A01-A10 overlapped with unique-mapped genomic regions encoding for 24-nt siRNAs, and more than 80% of the lncRNAs overlapping with genomic regions encoding for 24-nt siRNAs were from regions that harbored IRRs (Fig 2). 24-nt siRNAs were mapped in a similar way to lncRNA and its 5’ and 3’ flanking regions; this mapping pattern is different from those of genic regions or IRRs (Fig 3).

Fig 3. 24nt-siRNAs mapped to genic region, IRRs, putative mRNAs and each type of lncRNAs. The x-axis represents the target region (genic region, IRRs, putative mRNAs, or lncRNAs) and the flanking 5' and 3' regions. The y-axis represents the reads per million (RPM) value of 24-nt siRNA that overlaps with the corresponding target region.

We examined the whole genome DNA methylation state by WGBS using the same 14-day first and second leaves [36]. The average DNA methylation levels in regions covering lncRNAs were similar to those of the whole genome (S2 Fig). DNA methylation level in regions producing NATs was lower than those producing any of the three types of lncRNAs. Overlap of lncRNAs with IRRs or genomic regions

encoding for 24-nt siRNAs was associated with increased DNA methylation levels and overlapping with both caused further increases in DNA methylation levels (Fig 4).

Fig 4. DNA methylation level of genomic regions encoding each RNA type with siRNAs or IRRs. “+” indicates with and “-” indicates without overlapping genomic regions encoding for siRNAs or IRRs.

The DNA methylation were similar in lincRNA and NAT encoding regions, and DNA methylation levels in the 5’ and 3’ flanking regions were higher than in the body regions encoding lincRNAs and NATs (S3 Fig). The DNA methylation levels over the incRNA encoding regions and the flanking regions did not change (S3 Fig). Overall, mapping of DNA methylation levels to lncRNA encoding regions were different from those of genic regions or IRRs (S3 Fig).

We examined the influence on the lncRNA expression level when IRRs or siRNA clusters on the genome spanned the lncRNA regions. IRRs did not affect the expression levels of lncRNAs (Fig 5). The expression levels of lncRNAs covered with siRNA clusters were higher than those not covered by siRNA clusters (Fig 5)

Fig 5. Expression level of each type of RNAs with (+) or without (-) overlapping IRRs, 24nt-siRNAs, or H3K27me3. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$ (Student t-test)

Relationship between lncRNAs and H3K27me3

We have examined the H3K27me3 distribution using the 14-day first and second leaf samples [37] and identified the genomic regions encoding for lncRNAs that have H3K27me3 marks. 127 of 1,173 (10.8 %) lncRNAs, 83 of 529 (15.7%) NATs,

and 15 of 92 (16.3%) lncRNAs encoding genomic regions had H3K27me3 marks (S4 Fig). H3K27me3 was enriched in the transcribed region of lncRNAs, similar to the genomic regions, but H3K27me3 levels in lncRNA encoding genomic regions were lower than in regions coding for mRNAs (S5 Fig). The expression level of lncRNAs was higher when the encoding regions had H3K27me3 than without H3K27me3 (Fig 5). In NATs and lincRNAs, there was no difference of expression level with and without H3K27me3 on their encoding regions (Fig 5).

Characterization of un-annotated genes

In 2,052 lncRNAs mapped to intergenic regions of the genome, 608 transcripts had hits (e-value < 1.0e-10) against the Swissport database using BLASTX, indicating they could be un-annotated genes. The expression levels of the putative mRNAs from these regions were similar to that of mRNA [27]. Putative mRNAs tended to have fewer exons compared with annotated mRNAs (Fig 1A). The mean transcript length of putative mRNA genes was similar to that of mRNAs (Fig 1B).

314 of 490 (64.1%) putative mRNAs in A01-A10 overlapped with IRRs (Fig 2) but the expression level of putative mRNAs overlapping with IRRs was similar to those not-overlapping with IRRs (Fig 5). Mapping of 24-nt siRNAs and DNA methylation levels to genomic regions encoding putative mRNAs were similar to that of genomic regions (Fig 3, S3 Fig). The average expression level of putative mRNAs with

overlapping 24-nt siRNAs was higher than that without any overlap (Fig 5). The pattern of the average of DNA methylation level in the genomic region encoding putative mRNAs and their flanking regions was similar to that to the average of the genic regions (S3 Fig). Overlapping IRRs or 24-nt siRNAs resulted in an increase in DNA methylation levels and overlapping with both causes further increase in DNA methylation levels (Fig 4).

53 of 490 (10.8%) genomic regions corresponding to putative mRNAs had H3K27me3. H3K27me3 was enriched in the genomic regions encoding for lncRNAs, especially around the transcription start site, and this was similar to the pattern in the genic region (S5 Fig). The expression levels of putative mRNAs with corresponding genomic H3K27me3 were lower than those without H3K27me3, but there was no significant difference (Fig 5).

Examination of the conservation of lncRNAs among the Brassica genus

Using the sequences of the 2,088 lncRNAs of *B. rapa*, a best-blast hit search against the *B. nigra*, *B. oleracea*, *B. juncea*, and *B. napus* reference genomes was conducted using GenBlastA (e-value = 1e-10) to examine the conservation of lncRNAs. *B. rapa* lncRNAs were most conserved in *B. oleracea* and moderate conservation was

297 observed in *B. nigra* and *B. napus*. LncRNAs were less conserved in *B. juncea*,
298 especially in the incRNAs (Fig 6).

299

300

301 **Fig 6. BLAST search of each type of lncRNA from *B. rapa* against *B. nigra*, *B.***
302 ***oleracea*, *B. juncea*, and *B. napus*.**

303

304 We selected twelve lncRNAs that showed high sequence similarity with the *B.*
305 *oleracea* reference genome. We tested whether these lncRNA coding genomic
306 sequences were conserved in three commercial cultivars of cabbage (*B. oleracea*) by
307 PCR using genomic DNA as template. PCR amplification of genomic regions
308 corresponding to all twelve lncRNAs was confirmed in all three *B. oleracea* lines. Next,
309 we tested by RT-PCR whether these putative lncRNAs in *B. oleracea* are expressed in
310 the first and second leaves. Six of twelve lncRNAs were expressed in all three cultivars.
311 One lncRNA was expressed in two of three lines, and one lncRNA was expressed in

one of three lines. The remaining four lncRNAs were not expressed or were slightly expressed in all three lines (Table 1). We also examined the variation within *B. rapa* species using six lines. Seven of the twelve lncRNAs were expressed in all six lines, and the remaining five lncRNA showed line-specificity (Table 1).

Table 1. Conservation and diversity of the expression of lncRNAs within *B. rapa* species or between species. RT-PCR results of each lncRNAs “+++” indicates strong PCR band, “+” indicates weak PCR band, and “-” indicates no PCR band.

	<i>B. rapa</i>						<i>B. oleracea</i>		
lncRNAs	RJKB-T23	Homei	Harunosaiten	BRA2209	Osome	Yellow Sarson	Reiho	Matsunami	Kinkei
M15784	+++	-	+++	-	+++	+++	-	-	-
M26919	+++	+++	+++	+++	+++	+++	+++	+++	+++
M3316	+++	+++	+++	+++	+++	+++	+++	+++	-
M491	+++	-	+++	+++	+++	-	+	+	-
M17356	+++	+	-	-	-	-	-	-	-
M17153	+++	-	-	-	-	+	+++	+++	+++
M25534	+++	+++	+++	+++	+++	+++	+++	+++	+++
M259	+++	+++	+++	+++	+++	+++	+++	+++	+++
M4317	+++	+++	+++	+++	+++	+++	+++	+++	+++
M26796	+++	+++	+++	+++	+++	+++	+++	+++	+++
M4921	+++	+++	+	+	+++	+++	-	+++	-
M24531	+++	+++	+++	+++	+++	+++	-	+	-

Discussion

We analyzed 2,088 lncRNAs in leaves of *B. rapa*; their characteristics such as number of exons, length of transcripts (except for NATs), and lower expression levels were similar to those reported in other plant species [40–45]. The total number of lncRNAs seems to be similar among species [46,47], but there tends to be low sequence conservation between different species [48]. Lower conservation of lncRNAs than mRNAs was observed between *B. napus* and its two ancestral species, *B. rapa* and *B. oleracea* [29,42], even though the hybridization was relatively recent event, ~1,910–7,180 years ago [49]. We also found a lower sequence conservation of *B. rapa* lncRNAs among other species of the genus Brassica (*B. nigra*, *B. oleracea*, *B. juncea*, and *B. napus*). In 12 selected highly conserved lncRNAs, we found conservation of the lncRNAs at both the sequence and transcriptional level between *B. rapa* and *B. oleracea*, but there was transcriptional variation within species agreeing with the IRR diversification of each species after the allotetraploidization of *B. napus* [50]. The analysis of sequence homology of lncRNAs in *B. rapa* with other members of Brassica genus may deepen the understanding of the evolutionary dynamics of lncRNAs in the genus Brassica.

We found over 55% of each lncRNAs (65.0% lincRNAs, 55.0% NATs, and 71.7% incRNAs) overlapped with IRRs including TEs throughout the *B. rapa* genome. TEs are considered a source of siRNAs [51], and lincRNAs identified in *A. thaliana*,

rice, and maize, were associated with 22.2%, 49.7%, and 51.5% of TEs, respectively [52]. In this study, we found 18.7% lincRNAs, 14.0% NATs, and 17.4% of incRNAs covered unique-mapped 24-nt siRNAs, and over 80% of lincRNAs covering the genomic regions encoding for 24-nt siRNAs overlapped with IRRs with perfect sequence match, suggesting that lincRNAs covering IRRs are also the likely source of siRNAs in *B. rapa*. For mRNA, the average expression levels of lincRNAs having 24-nt siRNAs was higher than that of lincRNAs not having 24-nt siRNAs [36]. LncRNAs and 24-nt siRNAs are known to be involved in the increase of DNA methylation, which agrees with reports of increased gene expression when DNA methylation overlaps with an exonic region [53]. The detailed mechanism is not clear, but our results also agrees that lincRNAs may be involved in this gene regulation mechanism.

The epigenetic functions of different lincRNAs have been identified in diverse organisms [21,22], but there is limited information in *B. rapa*. LncRNAs show a close association with DNA methylation of the genomic region encoding the lincRNA [54], but studies have focused more on the model plant species, *A. thaliana* [11,55]. In this study, we found a similar level of DNA methylation in regions encoding for lincRNA compared to previously described DNA methylation in the *B. rapa* genome [36]. Levels of DNA methylation can be positively regulated by siRNAs through the RdDM pathway [56], and Pol IV is important for biogenesis of 24-nt siRNAs [57]. Identification of long Pol IV-dependent transcripts is difficult because these transcripts

can be rapidly processed to produce 24-nt siRNAs. There are several reports that detected Pol IV-dependent siRNA-precursor transcripts in *A. thaliana* of different lengths [57–59]. Most Pol IV-dependent transcript regions overlapped with Pol IV-dependent siRNA loci, and CHH methylation depends on the production of Pol IV-dependent transcripts [59]. However, if Pol IV-dependent transcripts are short as 30-40 nucleotides [57,58], the lncRNAs that we identified are of much longer length and not likely to be Pol-IV dependent transcripts that act as siRNA-precursors.

Pol V transcripts accumulate at very low levels and have been difficult to identify. However, using RNA immunoprecipitation, Pol V-dependent lncRNAs were identified, and CHH methylation and 24-nt siRNA accumulation were shown to be restricted to Pol V transcribed regions [60]. In this study, about 13 % of lncRNAs overlapped with genomic regions encoding for 24-nt siRNAs, and the DNA methylation level of those regions was higher than the average. It has been reported that the median length of Pol V transcripts was 689 nt [60]. The length of the lncRNAs overlapping genomic regions encoding for 24-nt siRNAs identified in this study resembles the length of the Pol V transcript precursors.

The level of H3K27me3 plays a role in tissue specific gene expression [61]. LncRNAs are also considered to be involved in regulating histone modifications. Binding of lncRNAs to PRC2 has been observed in human/animal [62,63], and COLDAIR and COLDWRAP have been shown to recruit the PRC2 complex to *FLC*

381 during vernalization in *A. thaliana* [13,14,64,65]. EMF2B is a component of the PRC2
382 complex and in rice, a mutant of *emf2b* was reported to lose H3K27me3 and derepress
383 some of the lincRNAs, suggesting that expression of these lincRNAs is regulated by
384 PRC2-mediated histone methylation of the region encoding the lincRNA [66]. We found
385 that about 10-16% of lincRNAs overlapped with H3K27me3 regions, which is lower
386 than the overlap with mRNAs. The pattern of H3K27me3 in lincRNA overlapping
387 regions was similar to mRNAs; H3K27me3 accumulated in the body region of
388 transcripts, especially around the transcription start site. However, there was no
389 negative relationship between the presence of H3K27me3 and lincRNA expression
390 levels, and lincRNA coding regions having H3K27me3 marks showed higher expression
391 levels than lincRNA coding regions without H3K27me3 marks. H3K27me3 over
392 lincRNA coding regions was also found in different tissues of maize, which was
393 responsible for the regulation of tissue-specific lincRNAs expression [67].
394 Coincidentally, DNA methylation in intronic regions is known to influence splicing
395 patterns [68], which may reflect epigenetic tissue specific gene regulation involving
396 RdDM and H3K27me3. However, as our study focused on the analyses in a single
397 tissue/developmental stage, we could not identify the lincRNAs that are transcriptionally
398 silenced by H3K27me3, leading to the underestimation of lincRNAs with H3K27me3
399 marks. Further study will be required to examine the transcriptional regulation of
400 lincRNAs by H3K27me3.

This study revealed that a small proportion of lncRNAs in *B. rapa* are conserved with other Brassica species. The majority of lncRNAs in *B. rapa* overlap with IRRs and there is some overlap with DNA methylation and 24-nt siRNAs, hinting at regulation of lncRNA expression through the RdDM pathway. Interestingly, some of the lncRNAs that overlapped with the genomic regions having H3K27me3 marks were more highly expressed, which was unexpected because of the known gene suppression activity of H3K27me3. This may indicate an unknown regulatory mechanism of lncRNAs by H3K27me3. However, the current study focuses on the genome-wide quantification of lncRNAs, 24-nt siRNAs, DNA methylation, and H3K27me3, and further quantitative analysis at the locus specific level may reveal a clearer relationship between these marks. Epigenetic regulation of the genome including lncRNA, 24-nt siRNAs, DNA methylation and H3K27me3 depend on the stage of development, and abiotic stress conditions. Further exploration of lncRNAs under different tissues and conditions may reveal lncRNAs with specific functions.

Acknowledgement

We are grateful to Ms. Tomoko Kusumi for her technical assistance throughout this project.

References

1. U N. Genome analysis in Brassica with special reference to the experimental formation of *B. napus* and peculiar mode of fertilization. Japanese J Bot. 1935;7: 389–452.
2. Lv H, Miyaji N, Osabe K, Akter A, Mehraj H, Shea DJ, et al. The Importance of genetic and epigenetic research in the *Brassica* vegetables in the face of climate change. In: Kole C, editor. Genomic designing of climate-smart vegetable crops. Springer; 2020. pp. 161-255.
3. Wang X, Wang H, Wang J, Sun R, Wu J, Liu S, et al. The genome of the mesopolyploid crop species *Brassica rapa*. Nat Genet. 2011;43: 1035–1040. doi:10.1038/ng.919
4. Liu S, Liu Y, Yang X, Tong C, Edwards D, Parkin IAP, et al. The *Brassica oleracea* genome reveals the asymmetrical evolution of polyploid genomes. Nat Commun. 2014;5: 3930. doi:10.1038/ncomms4930
5. Parkin IAP, Koh C, Tang H, Robinson SJ, Kagale S, Clarke WE, et al. Transcriptome and methylome profiling reveals relics of genome dominance in the

- 436 mesopolyploid *Brassica oleracea*. Genome Biol. 2014;15: R77.
437 doi:10.1186/gb-2014-15-6-r77
- 438 6. Yang J, Liu D, Wang X, Ji C, Cheng F, Liu B, et al. The genome sequence of
439 allopolyploid *Brassica juncea* and analysis of differential homoeolog gene
440 expression influencing selection. Nat Genet. 2016;48: 1225–1232.
441 doi:10.1038/ng.3657
- 442 7. Chalhoub B, Denoeud F, Liu S, Parkin IAP, Tang H, Wang X, et al. Early
443 allopolyploid evolution in the post-Neolithic *Brassica napus* oilseed genome.
444 Science. 2014;345: 950–953. doi:10.1126/science.1253435
- 445 8. Karlik E, Ari S, Gozukirmizi N. LncRNAs: genetic and epigenetic effects in plants.
446 Biotechnol Biotechnol Equip. 2019;33: 429–439.
447 doi:10.1080/13102818.2019.1581085
- 448 9. Rai MI, Alam M, Lightfoot DA, Gurha P, Afzal AJ. Classification and experimental
449 identification of plant long non-coding RNAs. Genomics. 2019;111: 997–1005.
450 doi:10.1016/j.ygeno.2018.04.014
- 451 10. Mattick JS, Rinn JL. Discovery and annotation of long noncoding RNAs. Nat Struct
452 Mol Biol. 2015;22: 5–7. doi:10.1038/nsmb.2942
- 453 11. Chekanova JA. Long non-coding RNAs and their functions in plants. Curr Opin
454 Plant Biol. 2015;27: 207–216. doi:10.1016/j.pbi.2015.08.003

- 455 12. Liu J, Wang H, Chua NH. Long noncoding RNA transcriptome of plants. Plant
456 Biotechnol J. 2015;13: 319–328. doi:10.1111/pbi.12336
- 457 13. Heo JB, Sung S. Vernalization-mediated epigenetic silencing by a long intronic
458 noncoding RNA. Science 2011;331: 76–79. doi:10.1126/science.1197349
- 459 14. Kim DH, Sung S. Vernalization-triggered intragenic chromatin loop formation by
460 long noncoding RNAs. Dev Cell. 2017;40: 302-312.e4.
461 doi:10.1016/j.devcel.2016.12.021
- 462 15. Swiezewski S, Liu F, Magusin A, Dean C. Cold-induced silencing by long antisense
463 transcripts of an *Arabidopsis* Polycomb target. Nature. 2009;462: 799–802.
464 doi:10.1038/nature08618
- 465 16. Ding J, Lu Q, Ouyang Y, Mao H, Zhang P, Yao J, et al. A long noncoding RNA
466 regulates photoperiod-sensitive male sterility, an essential component of hybrid rice.
467 Proc Natl Acad Sci U S A. 2012;109: 2654–2659. doi:10.1073/pnas.1121374109
- 468 17. Wang Y, Fan X, Lin F, He G, Terzaghi W, Zhu D, et al. *Arabidopsis* noncoding
469 RNA mediates control of photomorphogenesis by red light. Proc Natl Acad Sci U S
470 A. 2014;111: 10359–10364. doi:10.1073/pnas.1409457111
- 471 18. Fedak H, Palusinska M, Krzyczmonik K, Brzezniak L, Yatusovich R, Pietras Z, et al.
472 Control of seed dormancy in *Arabidopsis* by a *cis*-acting noncoding antisense
473 transcript. Proc Natl Acad Sci U S A. 2016;113: E7846–E7855.
474 doi:10.1073/pnas.1608827113

- 475 19. Seo JS, Sun HX, Park BS, Huang CH, Yeh SD, Jung C, et al. ELF18-INDUCED
476 LONG-NONCODING RNA associates with mediator to enhance expression of
477 innate immune response genes in Arabidopsis. *Plant Cell*. 2017;29: 1024–1038.
478 doi:10.1105/tpc.16.00886
- 479 20. Wu CT, Morris JR. Genes, genetics, and epigenetics: A correspondence. *Science*.
480 2001;293: 1103–1105. doi:10.1126/science.293.5532.1103
- 481 21. Wang C, Wang L, Ding Y, Lu X, Zhang G, Yang J, et al. LncRNA structural
482 characteristics in epigenetic regulation. *Int J Mol Sci*. 2017;18: 2659.
483 doi:10.3390/ijms18122659
- 484 22. Matzke MA, Kanno T, Matzke AJM. RNA-directed DNA methylation: The
485 evolution of a complex epigenetic pathway in flowering plants. *Annu Rev Plant*
486 *Biol*. 2015;66: 243–267. doi:10.1146/annurev-arplant-043014-114633
- 487 23. Fujimoto R, Sasaki T, Ishikawa R, Osabe K, Kawanabe T, Dennis ES. Molecular
488 mechanisms of epigenetic variation in plants. *Int J Mol Sci*. 2012;13: 9900–9922.
489 doi:10.3390/ijms13089900
- 490 24. Liu ZW, Zhao N, Su YN, Chen SS, He XJ. Exogenously overexpressed intronic
491 long noncoding RNAs activate host gene expression by affecting histone
492 modification in Arabidopsis. *Sci Rep*. 2020;10: 3094.
493 doi:10.1038/s41598-020-59697-7

25. Kim DH, Xi Y, Sung S. Modular function of long noncoding RNA, COLDAIR, in the vernalization response. PLoS Genet. 2017;13: e1006939. doi:10.1371/journal.pgen.1006939
26. Itabashi E, Osabe K, Fujimoto R, Kakizaki T. Epigenetic regulation of agronomical traits in Brassicaceae. Plant Cell Rep. 2018;27: 87-101. doi.org/10.1007/s00299-017-2223-z
27. Shea DJ, Nishida N, Takada S, Itabashi E, Takahashi S, Akter A, et al. Long noncoding RNAs in *Brassica rapa* L. following vernalization. Sci Rep. 2019;9: 9302. doi:10.1038/s41598-019-45650-w
28. Wang A, Hu J, Gao C, Chen G, Wang B, Lin C, et al. Genome-wide analysis of long non-coding RNAs unveils the regulatory roles in the heat tolerance of Chinese cabbage (*Brassica rapa* ssp.chinensis). Sci Rep. 2019;9: 5002. doi:10.1038/s41598-019-41428-2
29. Zhang J, Wei L, Jiang J, Mason AS, Li H, Cui C, et al. Genome-wide identification, putative functionality and interactions between lncRNAs and miRNAs in Brassica species. Sci Rep. 2018;8: 4960. doi:10.1038/s41598-018-23334-1
30. Huang L, Dong H, Zhou D, Li M, Liu Y, Zhang F, et al. Systematic identification of long non-coding RNAs during pollen development and fertilization in *Brassica rapa*. Plant J. 2018;96: 203–222. doi:10.1111/tpj.14016

- 513 31. Song X, Liu G, Huang Z, Duan W, Tan H, Li Y, et al. Temperature expression
514 patterns of genes and their coexpression with LncRNAs revealed by RNA-Seq in
515 non-heading Chinese cabbage. BMC Genomics. 2016;17: 297.
516 doi:10.1186/s12864-016-2625-2
- 517 32. Liu X, Hao L, Li D, Zhu L, Hu S. Long non-coding RNAs and their biological roles
518 in plants. Genomics, Proteomics Bioinformatics. 2015;13: 137–147.
519 doi:10.1016/j.gpb.2015.02.003
- 520 33. Yu X, Yang J, Li X, Liu X, Sun C, Wu F, et al. Global analysis of cis-natural
521 antisense transcripts and their heat-responsive nat-siRNAs in *Brassica rapa*. BMC
522 Plant Biol. 2013;13: 208. doi:10.1186/1471-2229-13-208
- 523 34. Kawamura K, Kawanabe T, Shimizu M, Nagano AJ, Saeki N, Okazaki K, et al.
524 Genetic distance of inbred lines of Chinese cabbage and its relationship to heterosis.
525 Plant Gene. 2016;5: 1-7. doi:10.1016/j.plgene.2015.10.003
- 526 35. Murray MG, Thompson WF. Rapid isolation of high molecular weight plant DNA.
527 Nucleic Acids Res. 1980;8: 4321–4326. doi:10.1093/nar/8.19.4321
- 528 36. Takahashi S, Osabe K, Fukushima N, Takuno S, Miyaji N, Shimizu M, et al.
529 Genome-wide characterization of DNA methylation, small RNA expression, and
530 histone H3 lysine nine di-methylation in *Brassica rapa* L. DNA Res. 2018;25:
531 511–520. doi:10.1093/dnares/dsy021

37. Akter A, Takahashi S, Deng W, Shea DJ, Itabashi E, Shimizu M, et al. The histone modification H3 lysine 27 tri-methylation has conserved gene regulatory roles in the triplicated genome of *Brassica rapa* L. DNA Res. 2019;26: 433–443. doi:10.1093/dnares/dsz021
38. Yu J, Zhao M, Wang X, Tong C, Huang S, Tehrim S, et al. Bolbase: A comprehensive genomics database for *Brassica oleracea*. BMC Genomics. 2013;14: 664. doi:10.1186/1471-2164-14-664
39. She R, Chu JSC, Wang K, Pei J, Chen N. genBlastA: Enabling BLAST to identify homologous gene sequences. Genome Res. 2009;19: 143–149. doi:10.1101/gr.082081.108
40. Wang R, Zou J, Meng J, Wang J. Integrative analysis of genome-wide lncRNA and mRNA expression in newly synthesized Brassica hexaploids. Ecol Evol. 2018;8: 6034–6052. doi:10.1002/ece3.4152
41. Wang H, Chung PJ, Liu J, Jang IC, Kean MJ, Xu J, et al. Genome-wide identification of long noncoding natural antisense transcripts and their responses to light in *Arabidopsis*. Genome Res. 2014;24: 444–453. doi:10.1101/gr.165555.113
42. Shen E, Zhu X, Hua S, Chen H, Ye C, Zhou L, et al. Genome-wide identification of oil biosynthesis-related long non-coding RNAs in allopolyploid *Brassica napus*. BMC Genomics. 2018;19: 745. doi:10.1186/s12864-018-5117-8

551 43. Liu J, Jung C, Xu J, Wang H, Deng S, Bernad L, et al. Genome-wide analysis
552 uncovers regulation of long intergenic noncoding RNAs in *Arabidopsis*. *Plant Cell*.
553 2012;24: 4333–4345. doi:10.1105/tpc.112.102855

554 44. Joshi RK, Megha S, Basu U, Rahman MH, Kav NNV. Genome wide identification
555 and functional prediction of long non-coding RNAs responsive to *Sclerotinia*
556 *sclerotiorum* infection in *Brassica napus*. *PLoS One*. 2016;11: e0158784.
557 doi:10.1371/journal.pone.0158784

558 45. Chen R, Li M, Zhang H, Duan L, Sun X, Jiang Q, et al. Continuous salt
559 stress-induced long non-coding RNAs and DNA methylation patterns in soybean
560 roots. *BMC Genomics*. 2019;20: 730. doi:10.1186/s12864-019-6101-7

561 46. Kopp F, Mendell JT. Functional classification and experimental dissection of long
562 noncoding RNAs. *Cell*. 2018;172: 393–407. doi:10.1016/j.cell.2018.01.011

563 47. Kung JTY, Colognori D, Lee JT. Long noncoding RNAs: Past, present, and future.
564 *Genetics*. 2013;193: 651–669. doi:10.1534/genetics.112.146704

565 48. Vandivier LE, Anderson SJ, Foley SW, Gregory BD. The conservation and function
566 of RNA secondary structure in plants. *Annu Rev Plant Biol*. 2016;67: 463–488.
567 doi:10.1146/annurev-arplant-043015-111754

568 49. Lu K, Wei L, Li X, Wang Y, Wu J, Liu M, et al. Whole-genome resequencing
569 reveals *Brassica napus* origin and genetic loci involved in its improvement. *Nat*
570 *Commun*. 2019;10: 1154. doi:10.1038/s41467-019-09134-9

- 571 50. Sampath P, Murukarthick J, Izzah NK, Lee J, Choi H-I, Shirasawa K, et al.
572 Genome-wide comparative analysis of 20 miniature inverted-repeat transposable
573 element families in *Brassica rapa* and *B. oleracea*. PLoS One. 2014;9: e94499.
574 doi.org/10.1371/journal.pone.0094499
- 575 51. Cho J. Transposon-derived non-coding RNAs and their function in plants. Front
576 Plant Sci. 2018;9: 600. doi:10.3389/fpls.2018.00600
- 577 52. Wang D, Qu Z, Yang L, Zhang Q, Liu ZH, Do T, et al. Transposable elements (TEs)
578 contribute to stress-related long intergenic noncoding RNAs in plants. Plant J.
579 2017;90: 133–146. doi:10.1111/tpj.13481
- 580 53. Bewick AJ, Schmitz RJ. Gene body DNA methylation in plants. Curr Opin Plant
581 Biol. 2017;36: 103–110. doi:10.1016/j.pbi.2016.12.007
- 582 54. Di Ruscio A, Ebralidze AK, Benoukraf T, Amabile G, Goff LA, Terragni J, et al.
583 DNMT1-interacting RNAs block gene-specific DNA methylation. Nature.
584 2013;503: 371–376. doi:10.1038/nature12598
- 585 55. Au PCK, Dennis ES, Wang MB. Analysis of argonaute 4-associated long
586 non-coding RNA in *Arabidopsis thaliana* sheds novel insights into gene regulation
587 through RNA-directed DNA methylation. Genes (Basel). 2017;8: 198.
588 doi:10.3390/genes8080198
- 589 56. Yan H, Bombarely A, Xu B, Frazier TP, Wang C, Chen P, et al. siRNAs regulate
590 DNA methylation and interfere with gene and lncRNA expression in the

heterozygous polyploid switchgrass. *Biotechnol Biofuels*. 2018;11: 208.
doi:10.1186/s13068-018-1202-0

57. Zhai J, Bischof S, Wang H, Feng S, Lee TF, Teng C, et al. A one precursor one siRNA model for Pol IV-dependent siRNA biogenesis. *Cell*. 2015;163: 445–455.
doi:10.1016/j.cell.2015.09.032

58. Blevins T, Podicheti R, Mishra V, Marasco M, Wang J, Rusch D, et al. Identification of Pol IV and RDR2-dependent precursors of 24 nt siRNAs guiding de novo DNA methylation in *Arabidopsis*. *Elife*. 2015;4: e09591.
doi:10.7554/eLife.09591

59. Li S, Vandivier LE, Tu B, Gao L, Won SY, Li S, et al. Detection of Pol IV/RDR2-dependent transcripts at the genomic scale in *Arabidopsis* reveals features and regulation of siRNA biogenesis. *Genome Res*. 2015;25: 235–245.
doi:10.1101/gr.182238.114

60. Böhmendorfer G, Sethuraman S, Jordan Rowley M, Krzyszton M, Hafiz Rothi M, Bouzit L, et al. Long non-coding RNA produced by RNA polymerase V determines boundaries of heterochromatin. *Elife*. 2016;5: e19092. doi:10.7554/eLife.19092

61. Makarevitch I, Eichten SR, Briskine R, Waters AJ, Danilevskaya ON, Meeley RB, et al. Genomic distribution of maize facultative heterochromatin marked by trimethylation of H3K27. *Plant Cell*. 2013;25: 780–793.
doi:10.1105/tpc.112.106427

62. Tu S, Yuan GC, Shao Z. The PRC2-binding long non-coding RNAs in human and mouse genomes are associated with predictive sequence features. *Sci Rep.* 2017;7: 41669. doi:10.1038/srep41669
63. Gaiti F, Degnan BM, Tanurdžić M. Long non-coding regulatory RNAs in sponges and insights into the origin of animal multicellularity. *RNA Biol.* 2018;15: 696–702. doi:10.1080/15476286.2018.1460166
64. Tian Y, Zheng H, Zhang F, Wang S, Ji X, Xu C, et al. PRC2 recruitment and H3K27me3 deposition at *FLC* require FCA binding of *COOLAIR*. *Sci Adv.* 2019;5: eaau7246. doi:10.1126/sciadv.aau7246
65. Yu T, Zhu H. Long non-coding RNAs: Rising regulators of plant reproductive development. *Agronomy.* 2019;9: 53. doi:10.3390/agronomy9020053
66. Johnson C, Conrad LJ, Patel R, Anderson S, Li C, Pereira A, et al. Reproductive long intergenic noncoding RNAs exhibit male gamete specificity and polycomb repressive complex 2-mediated repression. *Plant Physiol.* 2018;177: 1198–1217. doi:10.1104/pp.17.01269
67. Li L, Eichten SR, Shimizu R, Petsch K, Yeh CT, Wu W, et al. Genome-wide discovery and characterization of maize long non-coding RNAs. *Genome Biol.* 2014;15: R40. doi:10.1186/gb-2014-15-2-r40
68. Regulski M, Lu Z, Kendall J, Donoghue MTA, Reinders J, Llaca V, et al. The maize methylome influences mRNA splice sites and reveals widespread paramutation-like

631 switches guided by small RNA. Genome Res. 2013;23: 1651–1662.

632 doi:10.1101/gr.153510.112

633

634

Supporting information

S1 Fig. Chromosomal (A01-A10) distribution of lncRNA and mRNAs. Log₂ FPKM

of mRNAs represented in blue (positive values) and orange (negative values), and lncRNAs in green (positive values) and pink (negative values).

S2 Fig. DNA methylation level of each type of lincRNAs, putative mRNA, total, and IRRs. “Total” represents the methylation level of all regions.

S3 Fig. DNA methylation of the genic region, IRRs, or regions coding for putative mRNA and lincRNAs.

S4 Fig. Proportion of lincRNAs, putative mRNA, mRNA with (K27+) or without (K27-) H3K27me3 on encoding regions.

S5 Fig. H3K27me3 mapped to the genic region, IRRs, or encoding regions of putative mRNA or lincRNAs.