Visual effort moderates postural cascade dynamics

Madhur Mangalam^{a,*}, I-Chieh Lee^b, Karl M. Newell^c, and Damian G. Kelty-Stephen^{d,**}

^a*Department of Physical Therapy, Movement and Rehabilitation Sciences, Northeastern University, Boston, MA 02115, USA*

^b*UNC-NC State Joint Department of Biomedical Engineering, UNC-Chapel Hill, Chapel Hill, NC 27514, USA*

^c*Department of Kinesiology, University of Georgia, Athens, GA 30602, USA*

^d*Department of Psychology, Grinnell College, Grinnell, IA 50112, USA*

*Corresponding author at: Department of Physical Therapy, Movement and Rehabilitation Sciences, Northeastern University, Boston, MA 02115, USA.

Email address: m.mangalam@northeastern.edu (M. Mangalam.).

**Corresponding author at: Department of Psychology, Grinnell College, Grinnell, IA 50112, USA.

Email address: keltysda@grinnell.edu (D. G. Kelty-Stephen).

Author contributions: M.M. I.-C.L., K.M.N., and D.G.K-S. conceived and designed research; I.-C.L. performed experiments; M.M. and D.G.K-S. analyzed data; M.M. and D.G.K-S. interpreted results of experiments; M.M. prepared figures; M.M. and D.G.K-S. drafted manuscript; M.M., I.-C.L., K.M.N., and D.G.K-S. edited and revised manuscript; M.M., I.-C.L., K.M.N., and D.G.K-S. approved final version of manuscript.

The authors declare no competing interest.

26 **Abstract**

27 Standing still and focusing on a visible target in front of us is a preamble to many
 28 coordinated behaviors (e.g., reaching an object). Hiding behind its apparent simplicity is a
 29 deep layering of texture at many scales. The task of standing still laces together activities at
 30 multiple scales: from ensuring that a few photoreceptors on the retina cover the target in the
 31 visual field on an extremely fine scale to synergies spanning the limbs and joints at smaller
 32 scales to the mechanical layout of the ground underfoot and optic flow in the visual field on
 33 the coarser scales. Here, we used multiscale probability density function (PDF) analysis to
 34 show that postural fluctuations exhibit similar statistical signatures of cascade dynamics as
 35 found in fluid flow. In participants asked to stand quietly, the oculomotor strain of visually
 36 fixating at different distances moderated postural cascade dynamics. Visually fixating at a
 37 comfortable viewing distance elicited posture with a similar cascade dynamics as posture with
 38 eyes closed. Greater viewing distances known to stabilize posture showed more diminished
 39 cascade dynamics. In contrast, nearest and farthest viewing distances requiring greater
 40 oculomotor strain to focus on targets elicited a dramatic strengthening of postural cascade
 41 dynamics, reflecting active postural adjustments. Critically, these findings suggest that vision
 42 stabilizes posture by reconfiguring the prestressed poise that prepares the body to interact
 43 with different spatial layouts.

44 **Keywords:** center of pressure; ergonomics; inverse power law; multiscale PDF; postural
 45 control; quiet stance

1. Introduction

Standing still and visually fixating on a target in front of us is a seemingly simple task. However, hiding within this apparent simplicity is a deep layering of texture at many scales. On an extremely fine-scale, task completion depends only on ensuring that a few photoreceptors on the retina cover the entire target in the visual field. Standing still and visually fixating at much coarser scales rests on the mechanical layout of the ground underfoot and optic flow in the visual field before you. Between optic flow and mechanical layout, standing still relies on synergies spanning the limbs and joints at smaller scales and subtle postural perturbations, triggering reflex arcs and muscle twitches, at yet smaller scales. The task of standing still and visually fixating laces together activities at these multiple scales, entailing a close connection among events of widely varying scales, from whole-body postural fluctuations at large scales, to head sway and eye movements at medium scales, and photoreceptor activity at the finest scales [1–3]. Consequently, the postural center of pressure (CoP) fluctuations—which directly relate to sway—exhibit similar statistical signatures of cascade dynamics as found in fluid flow [4]. The metaphor of ‘cascades,’ as in tumbling water that accelerates and splashes down a rockface, captures some critical aspects of the dynamics of emergent behavior [5–7]. Destabilizing posture emphasizes the postural cascade dynamics [8,9]. Here, we propose that perturbing the visual system has a similar effect of strengthening postural cascade dynamics, in which the optical flow varies with fixation distance, as a meandering stream joins other streams and gathers momentum into rushing rapids, becoming more turbulent along the way. Whereas multiscale entropy addresses the difference of variation across different scales [10,11], cascade dynamics explicitly implicates nonlinear interactions lacing these scales together.

Visually guided posture presents a curious case in which light entails mechanical change in bodily center of mass [12–17]. Beyond minuscule photoreceptor responses, large-scale layout of visible surfaces supports optic flow with bodily sway. Optic flow is the expansion, contraction, or rotation of visible surfaces as the body moves forward or backward, or turns to the side, respectively. Relative accelerations of visible surfaces provide the standing participant with rich information about the layout of objects. Placing a fixation target at different viewing distances perturbs the optic flow: changing the relative accelerations of visible surfaces as well as imposing retinal-specific constraints of oculomotor convergence (i.e., higher oculomotor strain in specific or visual effort in general) [18,19], prompting new head- and torso-sway (Fig. 1). Indeed, optic flow relies on nesting of behaviors across the aforementioned range of scales, including the organism nested in the task environment at the largest scale right down to the hierarchical organization of photoreceptor response of optic flow at the finest scale [3]. Comfortable viewing distances (50-100cm) requiring least oculomotor strain should most resemble posture with eyes closed [20,21]. Medium viewing distances beyond most-comfortable viewing in the range of 200 cm stabilize posture [22]. Extremely distant or extremely near targets will be harder to focus on and have a much slower or faster optic flow, respectively, requiring more sway to resolve the image [23].

Testing these hypotheses requires examining the divergence of postural fluctuations from Gaussian form. Gaussian distributions are classically the result of adding very many independent (i.e., non-interacting) variables. Interactivity amongst the participating variables promotes persistent if also intermittent, uneven growth of variance with timescale. Under the central limit theorem, homogeneity of variance keeps measurements within short, thin tails of a Gaussian distribution. But intermittent growth in variance manifests in distributions with longer, heavier tails specifically due to nonlinear temporal correlations [24,25]. Testing these

tails is even more difficult because they are by definition only populated by extreme and relatively rare values. However, fortunately, hydrodynamic-cascade research has yielded an analytical method for estimating non-Gaussianity based on querying better-populated regions of the distribution [26]. Applying this method to physiological measures produces estimates of non-Gaussianity that predict clinical outcomes [27–31] and that change with variation in both endogenous postural control (i.e., unmanipulated factors that respond to but also exert effects upon other variables of the postural system) and exogenous postural demands [8,9]. Here, we used this method to investigate how constraints imposed by visual effort affect postural cascade dynamics. Participants were asked to minimize postural sway while maintaining quiet stance for 120 s in six different viewing conditions: eyes closed and eyes fixated on a point on a screen at a distance of 25, 50, 135, 220, and 305 cm from the eyes. We predicted that postural cascade dynamics while fixating at a 50-cm distance would resemble eyes-closed posture (Hypothesis-1), and that postural cascade dynamics would be weakest while fixating at medium (i.e., 130- and 220-cm) distances (Hypothesis-2) but strongest at extremely near (i.e., 25-cm) and extremely far (i.e., 305-cm) distances (Hypothesis-3).

2. Materials and methods

The present study is a reanalysis of data from a previous study examining postural control in different viewing conditions [32].

2.1. Participants

Seven men and eight women (18–40 years old) participated after providing institutionally-approved informed consent.

2.2. Experimental setup and procedure

Each participant stood barefoot on two force plates (AMTI Inc., Watertown, MA) recording 3D moments and ground reaction forces at 100 Hz—one foot on each plate, 25 cm apart, before a white background spanning his/her visual field (Fig. 1A). A square white screen (5×5 cm) mounted on a tripod was placed at specific distances in front of the participant. From behind the participant, a laser pen projected a static light point on the screen's center.

Each participant experienced six different viewing conditions (eyes-closed and eyes fixated on the screen's light point at a distance of 25, 50, 135, 220, and 305 cm in the front), each three times in randomized order over a single 90-min session. In each trial, each participant was instructed to minimize the postural sway by maintaining visual fixation at the light point projected on the screen for 120 s. Each participant had a 5-min break for rest after every six trials.

2.3. Data processing

All data processing was performed in MATLAB 2019b (Matlab Inc., Natick, MA). Trial-by-trial ground reaction forces yielded a 2D center of pressure (CoP) series, each dimension describing anterior-posterior (AP) and medial-lateral (ML) axes. Each 60-s trial yielded 12000-sample 2D CoP series and 11999-sample 2D CoP displacement series. Finally, a 1D CoP sample-to-sample planar Euclidean displacement (PED) series described postural fluctuations along the transverse plane. Iterated Amplitude Adjusted Fourier Transformation (IAAFT) provided phase-randomized surrogates using original series' spectral amplitudes to preserve only linear temporal correlations [33].

137 2.4. Canonical indices of endogenous postural fluctuations

138 Three linear indices were computed for each CoP PED series: (i) Mean of all
139 fluctuations (CoP_PED_Mean), (ii) Standard deviation (CoP_PED_SD), and (iii) Root mean
140 square (CoP_PED_RMSE).

141 Four nonlinear indices were computed for each CoP PED series: (i) Sample entropy
142 (CoP_PED_SampEn), indexing the extent of complexity in the signal [34], using $m = 2$ and $r =$
143 0.2 [35]. (ii) Multiscale sample entropy (CoP_PED_MSE), indexing signal complexity over
144 multiple timescales [10], using $m = 2$, $r = 0.2$, and $\tau = 20$ [35]. (iii) We used detrended
145 fluctuation analysis [36] to compute Hurst's exponent, H_{fGn} , indexing temporal correlations in
146 original CoP PED series (CoP_PED_ $H_{fGn_Original}$) and shuffled versions of each series
147 (CoP_PED_ $H_{fGn_Shuffled}$) using scaling region: 4, 8, 12,... 1024 [37]. Shuffling destroys the
148 temporal structure of a signal. (iv) Multifractal spectrum width, $\Delta\alpha$, indexing the extent of
149 multifractal temporal correlations in the signal. Chhabra and Jensen's direct method [38] was
150 used to compute $\Delta\alpha$ for original CoP series (CoP_PED_ $\Delta\alpha_{Original}$) and IAAFT surrogate
151 (CoP_PED_ $\Delta\alpha_{Surrogate}$) [6].

152 2.5. Multiscale probability density function (PDF) analysis

153 Multiscale PDF analysis characterizes the distribution of abrupt changes in CoP PED
154 series $\{ b(t) \}$ using the PDF tail. The first step is to generate $\{ B(t) \}$ by integrating $\{$
155 $b(t) \}$ after centering by the mean b_{ave} (Fig. S1A):

$$156 \quad B(t) = \sum_{i=1}^{t/\Delta t} b(i\Delta t) - b_{ave} \quad (1)$$

157 A 3rd order polynomial detrends $\{ B(t) \}$ within k overlapping windows of length $2s$,
158 s being the timescale (Fig. S1B). Intermittent deviation $\Delta_s B(t)$ in k^{th} window from
159 $1+s(k-1)$ to sk in the detrended time series $\{ B^d(t) = B(t) - f_{fit}(t) \}$ is computed as
160 $\Delta_s B^d(t) = B^d(t+s) - B^d(t)$, where $1+s(k-1) \leq t \leq sk$ and $f_{fit}(t)$ is the polynomial
161 representing the local trend of $\{ B(t) \}$, of which the elimination assures the zero-mean
162 probability density function in the next step (Fig. S1C). Finally, $\Delta_s B$ is normalized by the
163 SD (i.e., variance is set to one) to quantify the PDF.

164 To quantify the non-Gaussianity of $\Delta_s B$ at timescale s , the standardized PDF
165 constructed from all the $\Delta_s B(t)$ values is approximated by the Castaing model [26], with
166 λ_s as a single parameter characterizing the non-Gaussianity of the PDF. λ_s is estimated
167 as

$$168 \quad \lambda_s^2 = \frac{2}{q(q-2)} [\ln(\sqrt{\pi} \langle |\Delta_s B|^q \rangle) + \ln \Gamma(\frac{q+1}{2}) - \frac{q}{2} \ln 2] \quad (2)$$

169 where $\langle |\Delta_s B|^q \rangle$ denotes an estimated value of q^{th} order absolute moment of $\{ \Delta_s B \}$. As
170 λ_s^2 increases, the PDF becomes increasingly peaked and fat-tailed (Fig. S2A). λ_s^2 can
171 be estimated by Eq. (2) based on q^{th} order absolute moment of a time series independent
172 of q . Estimating λ_s^2 based on 0.2th moment ($q = 0.2$) emphasizes the center part of
173 the PDF, reducing the effects of extreme deviations due to heavy-tails and kurtosis. We used
174 0.2th moment because estimates of λ_s^2 for a time series of ~ 12000 samples are more
175 accurate at lower values of q [39].

Cascade-type multiplicative processes yield the inverse relationship $\lambda_s^2 \sim -\log s$ (Fig. S2B) [29]. For the present purposes, we quantified λ_s^2 for each original CoP PED series and corresponding IAAFT surrogate at timescales 5 to 1000 samples (i.e., 50 ms to 10 s) at steps of 5 samples (50 ms).

We used Akaike Information Criterion (AIC) weights obtained via the maximum likelihood estimation (MLE) to determine whether the PDF at each timescale was a power law, lognormal, exponential, or gamma distribution. We include this analysis also to show how AIC-based MLE sensitivity to lognormal-like heavy tails differs from multiscale PDF sensitivity to the bulk of the distribution.

2.6. Statistical analysis

A linear mixed-effect (LME) model using lmer [40] in R package *lme4* tested λ_s^2 vs. log-timescale curves for orthogonal linear and quadratic polynomials, for interactions with grouping variables (Viewing condition $\times$ Original, where Original encoded differences in original series from surrogates) and with indices of endogenous postural fluctuations (Section 2.4.). Statistical significance was assumed at the alpha level of 0.05 using R package *lmerTest* [41]. To test how lognormality changed with log-timescale, a generalized linear mixed-effect (GLME) fit changes in Lognormality as a dichotomous variable using orthogonal linear, quadratic, and cubic polynomials and tested interaction effects of grouping variables (Viewing condition $\times$ Original) with those polynomials using glmer [42] in *lme4*.

3. Results

The eyes-closed condition elicited nonlinearity in non-Gaussianity with log-timescale, showing a stronger linear increase in λ_s^2 up to the timescale of roughly 75 samples or 750 ms ($B = 0.95$, $P = 0.031$) and a stronger negative quadratic, downward-facing parabolic form ($B = -3.78$, $P < 0.001$) for the original CoP PED series than the surrogates (Table S1; Fig. 3A). Fixating at 25 and 305 cm elicited a stronger linear increase ($B = 2.43$, $P < 0.001$; and $B = 3.09$, $P < 0.001$, respectively) in the 25- and 305- cm viewing conditions while reversing the downward-facing parabolic form of the eyes-closed condition more weakly ($B = 4.42$, $P < 0.001$; and $B = 1.85$, $P = 0.003$, respectively) of λ_s^2 vs. log-timescale curves (Figs. 3B and 3F). Fixating the eyes at 135 and 220 cm elicited a reversal of linear ($B = -2.98$, $P < 0.001$; and $B = -2.32$, $P < 0.001$, respectively) and downward-facing parabolic forms ($B = 1.81$, $P = 0.004$; and $B = 2.00$, $P = 0.001$, respectively) of λ_s^2 vs. log-timescale curves in Figs. 3D and 3E. In contrast, fixating at 50 cm neither affected the linear form ($B = -0.09$, $P = 0.879$) nor the downward-facing parabolic form ($B = 0.73$, $P = 0.244$; Fig. 3C). Thus, the λ_s^2 vs. log-timescale curves differ between the eye-closed condition and all the eyes-open conditions except the eyes-fixated-at-50-cm condition. Furthermore, whereas the 50- and 305-cm viewing conditions show significant growth of the λ_s^2 vs. log-timescale curves, the 135- and 220-cm viewing conditions exhibit significant decay of this curve with scale.

All indices, except sample entropy of postural fluctuations, showed predictive effects on λ_s^2 . λ_s^2 decreased with increases in root mean square, and increased with increases in mean, standard deviation, and multiscale entropy of postural fluctuations (Table S2). Nonetheless, greater 'original-minus-shuffled/surrogate' differences in fractality and

multifractality diminished λ_s^2 . Hence, evidence of multifractality with weak difference from the surrogates suggests heavy-tailed distributions and thus non-Gaussianity [30,31].

Interactions of these indices with linear and quadratic terms in the first model significantly improved the model fit ($\chi^2_{315} = 153961$, $P < 0.001$; Table S2). Most variability in the quadratic form of the λ_s^2 vs. log-timescale curves depended on these indices, as all but two of 41 significant ($P < 0.05$) quadratic effects relate to interactions of these indices with log-timescale. That is, the parabolic tempering of non-Gaussianity across viewing conditions depended on endogenous postural control. A cubic polynomial improved model fit ($\chi^2(120) = 2061.8$, $P < 0.001$), reflecting 17 significant interactions with the cubic effect, but left all lower-order terms unchanged in coefficients and significance (Table S3).

Postural instability typically weakens the growth of lognormality in postural fluctuations with log-timescale [8,9]. The original CoP PED series exhibited less lognormality than the surrogates ($B = -4.28$, $P < 0.001$; Table S4; Fig. 4). Lognormality grew with log-timescale: a linear increase ($B = 2219.40$, $P = 0.016$), quadratic decrease ($B = -1678.38$, $P = 0.052$), and cubic increase ($B = 559.57$, $P = 0.051$) in the original CoP PED series than the surrogates. However, polynomial profiles of the growth of lognormality with log-timescale did not differ among the conditions. Hence, observed differences in the λ_s^2 vs. log-timescale curves reflect differences in the bulk of the postural-fluctuations distributions rather than in the tail.

4. Discussion

Visual effort moderates postural cascade dynamics, given that standing still and visually fixating at shorter and longer distances unleashed stronger cascade dynamics than standing still (Hypothesis-3) and visually fixating at an optimal distance or standing still without visually fixating (i.e., with eyes closed; Hypotheses-1 & 2). Importantly, these effects are not reducible to the mere availability of visual information, as standing still and visually fixating at an optimal distance and standing still with eyes closed showed similar, if not identical cascade dynamics. Nor is this effect reducible to postural stability, as the growth of lognormality in postural fluctuations with timescale did not depend on the viewing condition. Instead, these effects reflect endogenous postural control, as known indices of endogenous postural fluctuations moderated the effect of viewing condition on postural cascade dynamics. In short, the moderating of postural cascade dynamics by visual effort is an endogenous phenomenon that is difficult to counteract.

Past work has shown that variability in postural fluctuations is reduced when visually fixating at near as opposed to distant targets [12,13,22], an effect that has been directly attributed to the precision demands of viewing distance [32]. The present finding significantly adds to this work by showing that this effect is not linear. Visually fixating at too-close and too-far targets may introduce ambiguity between sway and optic flow: too-close targets making head-sway more likely to destabilize the posture, and too-far targets might recruit more torso-sway. However, visually fixating at an optimally placed target may resolve this ambiguity as the body courses with fluctuations ferrying information across the body. Furthermore, the stark similarities in the cascade dynamics between standing still and visually fixating at an optimal distance and standing still with eyes closed adds a novel dimension to the family of perception-action couplings in postural control. Postural control does not depend on vision per se. Instead, it depends on how visual information is laced together with activities at multiple

262 scales: subtle fluctuations of the head, torso, and the whole body and photoreceptor support
263 of the target's retinal image.

264 Postural cascade dynamics is rooted in the known physiology. Indeed, the biophysical
265 substrate supporting postural control is imagined as a neurally-tunable bodywide multifractal
266 tensegrity (MFT) network consisting of the connective-tissue net and extra-cellular matrix
267 (ECM) in which components hang together by tensional and compression forces at multiple
268 scales, from micro (e.g., cells and membranes) to macro (e.g., muscles and bones) [43–45].
269 This network cooperates directly and uninterruptedly with the nervous system and exhibits
270 similar cascade-like dynamics as in “neuronal avalanches” [46,47]. It is no small coincidence
271 that tensegrity and avalanche models all converge on multiplicativity [48,49] that inspired
272 multiscale quantification of non-Gaussianity [27–29]. Hence, MFT approaches suggest
273 multiple points of entry through which stimulation elicits cascades to spread across scales—
274 from the surface underfoot [8] to the loads at hand [9] and now from the incoming visual
275 information. As the cascade metaphor entails, multiple scales are not merely coexisting but
276 explicitly interacting, with influences of one spreading to others. If mechanically and visually
277 perturbing posture engender cascade dynamics similarly [8,9], then postural control may
278 emerge from the perceptuomotor sensitivity of specifically multiplicative fluctuations to the
279 environment. The methods introduced here would enable future investigations into how
280 different suprapostural tasks lace together deep texture at very many scales.

281 In short, the present findings suggest that visual perception by no means is a passive
282 function proceeding in parallel to still posture—the poise carried by the body as it stands and
283 looks suggests that moving visual targets nearer or farther brings a reorganization of postural
284 interactions across scales. This standing-and-looking preamble to action brims with dynamic
285 redistributions of tensional and compression forces. Indeed, the participants completed no
286 task beyond standing still and maintaining visual fixation. But these cascades affecting a
287 dynamic balance of forces across the body seem to be essential foundations for how the body
288 reaches out beyond itself to engage with the environment [7,50–52]. So, the present findings
289 show how vision plays not only upon the photoreceptors in the eyes but also across the full-
290 body postural system.

Supplementary material

Table S1. Coefficients of linear mixed-effect (LME) model examining the effect of Viewing condition on λ_s^2 vs. log-timescale curves.

Table S2. Coefficients of LME model examining the effects of Viewing condition on λ_s^2 vs. log-timescale curves for orthogonal linear and quadratic polynomials, for interactions with indices of endogenous postural fluctuations.

Table S3. Coefficients of LME model examining the effects of Viewing condition on λ_s^2 vs. log-timescale curves for orthogonal linear, quadratic, and cubic polynomials, for interactions with indices of endogenous postural fluctuations.

Table S4. Coefficients of generalized linear mixed-effect (GLME) model examining the effect of Viewing condition on the growth of lognormality in postural fluctuations with log-timescale.

References

- [1] J.S. Matthis, K.S. Muller, K. Bonnen, M.M. Hayhoe, Retinal optic flow during natural locomotion, *BioRxiv.* (2020) 2020.07.23.217893. <https://doi.org/10.1101/2020.07.23.217893>.
- [2] J.K. O'Regan, Solving the “real” mysteries of visual perception: The world as an outside memory., *Can. J. Psychol. Can. Psychol.* 46 (1992) 461–488. <https://doi.org/10.1037/h0084327>.
- [3] A.S. Mauss, A. Borst, Optic flow-based course control in insects, *Curr. Opin. Neurobiol.* 60 (2020) 21–27. <https://doi.org/10.1016/j.conb.2019.10.007>.
- [4] B. Manor, M.D. Costa, K. Hu, E. Newton, O. Starobinets, H.G. Kang, C.K. Peng, V. Novak, L.A. Lipsitz, Physiological complexity and system adaptability: Evidence from postural control dynamics of older adults, *J. Appl. Physiol.* 109 (2010) 1786–1791. <https://doi.org/10.1152/japplphysiol.00390.2010>.
- [5] D.G. Kelty-Stephen, K. Palatinus, E. Saltzman, J.A. Dixon, A tutorial on multifractality, cascades, and interactivity for empirical time series in ecological science, *Ecol. Psychol.* 25 (2013) 1–62. <https://doi.org/10.1080/10407413.2013.753804>.
- [6] M. Mangalam, D.G. Kelty-Stephen, Multiplicative-cascade dynamics supports whole-body coordination for perception via effortful touch, *Hum. Mov. Sci.* 70 (2020) 102595. <https://doi.org/10.1016/j.humov.2020.102595>.
- [7] M. Mangalam, N.S. Carver, D.G. Kelty-Stephen, Multifractal signatures of perceptual processing on anatomical sleeves of the human body, *J. R. Soc. Interface.* 17 (2020) 20200328. <https://doi.org/10.1098/rsif.2020.0328>.
- [8] M. Mangalam, D.G. Kelty-Stephen, Hypothetical control of postural sway, *BioRxiv.* (2020) 104760. <https://doi.org/10.1101/2020.05.19.104760>.

- 327 [9] P.F. Mariusz, M. Mangalam, D.G. Kelty-Stephen, G. Juras, Postural instability recruits
328 shorter-timescale processes into the non-Gaussian cascade processes, *BioRxiv.* (2020)
329 136895. <https://doi.org/10.1101/2020.06.05.136895>.
- 330 [10] M. Costa, A.L. Goldberger, C.-K. Peng, Multiscale entropy analysis of complex
331 physiologic time series, *Phys. Rev. Lett.* 89 (2002) 68102.
332 <https://doi.org/10.1103/PhysRevLett.89.068102>.
- 333 [11] M. Costa, A.L. Goldberger, C.-K. Peng, Multiscale entropy analysis of biological signals,
334 *Phys. Rev. E.* 71 (2005) 21906. <https://doi.org/10.1103/PhysRevE.71.021906>.
- 335 [12] T.A. Stoffregen, R.J. Pagulayan, B.G. Bardy, L.J. Hettinger, Modulating postural control
336 to facilitate visual performance, *Hum. Mov. Sci.* 19 (2000) 203–220.
337 [https://doi.org/10.1016/S0167-9457\(00\)00009-9](https://doi.org/10.1016/S0167-9457(00)00009-9).
- 338 [13] T.A. Stoffregen, L.J. Smart, B.G. Bardy, R.J. Pagulayan, Postural stabilization of
339 looking, *J. Exp. Psychol. Hum. Percept. Perform.* 25 (1999) 1641–1658. <https://doi.org/10.1037/0096-1523.25.6.1641>.
- 341 [14] M.C. Dault, A.C.H. Geurts, T.W. Mulder, J. Duysens, Postural control and cognitive task
342 performance in healthy participants while balancing on different support-surface
343 configurations, *Gait Posture.* 14 (2001) 248–255. [https://doi.org/10.1016/S0966-6362\(01\)00130-8](https://doi.org/10.1016/S0966-6362(01)00130-8).
- 345 [15] O. Oullier, B.G. Bardy, T.A. Stoffregen, R.J. Bootsma, Postural coordination in looking
346 and tracking tasks, *Hum. Mov. Sci.* 21 (2002) 147–167. [https://doi.org/10.1016/S0167-9457\(02\)00093-3](https://doi.org/10.1016/S0167-9457(02)00093-3).
- 348 [16] M.R. Givens, K. Yoshida, B. Bardy, M. Riley, T.A. Stoffregen, Postural sway and the
349 amplitude of horizontal eye movements, *Ecol. Psychol.* 23 (2011) 247–266.
350 <https://doi.org/10.1080/10407413.2011.617215>.
- 351 [17] T.A. Stoffregen, B.G. Bardy, C.T. Bonnet, R.J. Pagulayan, Postural stabilization of
352 visually guided eye movements, *Ecol. Psychol.* 18 (2006) 191–222.
353 https://doi.org/10.1207/s15326969eco1803_3.
- 354 [18] C.M. Fiorelli, P.F. Polastri, S.T. Rodrigues, A.M. Baptista, T. Penedo, V.A.I. Pereira, L.
355 Simieli, F.A. Barbieri, Gaze position interferes in body sway in young adults, *Neurosci.*
356 *Lett.* 660 (2017) 130–134. <https://doi.org/10.1016/j.neulet.2017.09.008>.
- 357 [19] T.-T. Lê, Z. Kapoula, Role of ocular convergence in the Romberg quotient, *Gait Posture.*
358 27 (2008) 493–500. <https://doi.org/10.1016/j.gaitpost.2007.06.003>.
- 359 [20] M.C. Musolino, P.J. Loughlin, P.J. Sparto, M.S. Redfern, Spectrally similar periodic and
360 non-periodic optic flows evoke different postural sway responses, *Gait Posture.* 23
361 (2006) 180–188. <https://doi.org/10.1016/j.gaitpost.2005.02.008>.

- 362 [21] T.A. Stoffregen, P. Hove, J. Schmit, B.G. Bardy, Voluntary and involuntary postural
363 responses to imposed optic flow, *Motor Control*. 10 (2006) 24–33.
364 <https://doi.org/10.1123/mcj.10.1.24>.
- 365 [22] Z. Kapoula, T.-T. Lê, Effects of distance and gaze position on postural stability in young
366 and old subjects, *Exp. Brain Res.* 173 (2006) 438–445. [https://doi.org/10.1007/s00221-](https://doi.org/10.1007/s00221-006-0382-1)
367 [006-0382-1](https://doi.org/10.1007/s00221-006-0382-1).
- 368 [23] J. Munafo, C. Curry, M.G. Wade, T.A. Stoffregen, The distance of visual targets affects
369 the spatial magnitude and multifractal scaling of standing body sway in younger and
370 older adults, *Exp. Brain Res.* 234 (2016) 2721–2730. [https://doi.org/10.1007/s00221-](https://doi.org/10.1007/s00221-016-4676-7)
371 [016-4676-7](https://doi.org/10.1007/s00221-016-4676-7).
- 372 [24] E.A. Codling, M.J. Plank, S. Benhamou, Random walk models in biology, *J. R. Soc.*
373 *Interface*. 5 (2008) 813–834. <https://doi.org/10.1098/rsif.2008.0014>.
- 374 [25] M.A. Lomholt, K. Tal, R. Metzler, K. Joseph, Lévy strategies in intermittent search
375 processes are advantageous, *Proc. Natl. Acad. Sci.* 105 (2008) 11055–11059.
376 <https://doi.org/10.1073/pnas.0803117105>.
- 377 [26] B. Castaing, Y. Gagne, E.J. Hopfinger, Velocity probability density functions of high
378 Reynolds number turbulence, *Phys. D Nonlinear Phenom.* 46 (1990) 177–200.
379 [https://doi.org/10.1016/0167-2789\(90\)90035-N](https://doi.org/10.1016/0167-2789(90)90035-N).
- 380 [27] B.J. West, M. Turlaska, Hypothetical control of heart rate variability, *Front. Physiol.* 10
381 (2019) 1078. <https://doi.org/10.3389/fphys.2019.01078>.
- 382 [28] K. Kiyono, Z.R. Struzik, N. Aoyagi, S. Sakata, J. Hayano, Y. Yamamoto, Critical scale
383 invariance in a healthy human heart rate, *Phys. Rev. Lett.* 93 (2004) 178103.
384 <https://doi.org/10.1103/PhysRevLett.93.178103>.
- 385 [29] K. Kiyono, J. Hayano, E. Watanabe, Z.R. Struzik, Y. Yamamoto, Non-Gaussian heart
386 rate as an independent predictor of mortality in patients with chronic heart failure, *Hear.*
387 *Rhythm*. 5 (2008) 261–268. <https://doi.org/10.1016/j.hrthm.2007.10.030>.
- 388 [30] K. Kiyono, J. Hayano, S. Kwak, E. Watanabe, Y. Yamamoto, Non-gaussianity of low
389 frequency heart rate variability and sympathetic activation: Lack of increases in multiple
390 system atrophy and parkinson disease, *Front. Physiol.* 3 (2012) 34.
391 <https://doi.org/10.3389/fphys.2012.00034>.
- 392 [31] J. Hayano, K. Kiyono, Z. Struzik, Y. Yamamoto, E. Watanabe, P. Stein, L. Watkins, J.
393 Blumenthal, R. Carney, Increased non-Gaussianity of heart rate variability predicts
394 cardiac mortality after an acute myocardial infarction, *Front. Physiol.* 2 (2011) 65.
395 <https://doi.org/10.3389/fphys.2011.00065>.

- 396 [32] I.-C. Lee, M.M. Pacheco, K.M. Newell, The precision demands of viewing distance
397 modulate postural coordination and control, *Hum. Mov. Sci.* 66 (2019) 425–439. <https://doi.org/10.1016/j.humov.2019.05.019>.
398
- 399 [33] T. Schreiber, A. Schmitz, Improved surrogate data for nonlinearity tests, *Phys. Rev. Lett.*
400 77 (1996) 635–638. <https://doi.org/10.1103/PhysRevLett.77.635>.
- 401 [34] J.S. Richman, J.R. Moorman, Physiological time-series analysis using approximate
402 entropy and sample entropy, *Am. J. Physiol. Circ. Physiol.* 278 (2000) H2039–H2049.
403 <https://doi.org/10.1152/ajpheart.2000.278.6.H2039>.
- 404 [35] J.-H. Ko, K.M. Newell, Aging and the complexity of center of pressure in static and
405 dynamic postural tasks, *Neurosci. Lett.* 610 (2016) 104–109.
406 <https://doi.org/10.1016/j.neulet.2015.10.069>.
- 407 [36] C.-K. Peng, S. V Buldyrev, S. Havlin, M. Simons, H.E. Stanley, A.L. Goldberger, Mosaic
408 organization of DNA nucleotides, *Phys. Rev. E.* 49 (1994) 1685–1689.
409 <https://doi.org/10.1103/PhysRevE.49.1685>.
- 410 [37] M. Mangalam, R. Chen, T.R. McHugh, T. Singh, D.G. Kelty-Stephen, Bodywide
411 fluctuations support manual exploration: Fractal fluctuations in posture predict
412 perception of heaviness and length via effortful touch by the hand, *Hum. Mov. Sci.* 69
413 (2020) 102543. <https://doi.org/10.1016/j.humov.2019.102543>.
- 414 [38] A. Chhabra, R. V Jensen, Direct determination of the $f(\alpha)$ singularity spectrum, *Phys.*
415 *Rev. Lett.* 62 (1989) 1327–1330. <https://doi.org/10.1103/PhysRevLett.62.1327>.
- 416 [39] K. Kiyono, Z.R. Struzik, Y. Yamamoto, Estimator of a non-Gaussian parameter in
417 multiplicative log-normal models, *Phys. Rev. E.* 76 (2007) 41113.
418 <https://doi.org/10.1103/PhysRevE.76.041113>.
- 419 [40] D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting linear mixed-effects models using
420 lme4, *ArXiv.* (2014) 5823. <https://doi.org/arXiv:1406.5823>.
- 421 [41] A. Kuznetsova, P.B. Brockhoff, R.H.B. Christensen, lmerTest package: Tests in linear
422 mixed effects models, *J. Stat. Softw.* 82 (2017) 1548–7660.
423 <https://doi.org/10.18637/jss.v082.i13>.
- 424 [42] W. Lee, K.J. Grimm, Generalized linear mixed-effects modeling programs in R for
425 binary outcomes, *Struct. Equ. Model. A Multidiscip. J.* 25 (2018) 824–828.
426 <https://doi.org/10.1080/10705511.2018.1500141>.
- 427 [43] R. Schleip, F. Mechsner, A. Zorn, W. Klingler, The bodywide fascial network as a
428 sensory organ for haptic perception, *J. Mot. Behav.* 46 (2014) 191–193.
429 <https://doi.org/10.1080/00222895.2014.880306>.
- 430 [44] M.T. Turvey, S.T. Fonseca, The medium of haptic perception: A tensegrity hypothesis, *J.*
431 *Mot. Behav.* 46 (2014) 143–187. <https://doi.org/10.1080/00222895.2013.798252>.

- 432 [45] D.E. Ingber, From cellular mechanotransduction to biologically inspired engineering,
433 Ann. Biomed. Eng. 38 (2010) 1148–1161. <https://doi.org/10.1007/s10439-010-9946-0>.
- 434 [46] S.R. Miller, S. Yu, D. Plenz, The scale-invariant, temporal profile of neuronal avalanches
435 in relation to cortical γ -oscillations, Sci. Rep. 9 (2019) 16403.
436 <https://doi.org/10.1038/s41598-019-52326-y>.
- 437 [47] D.R. Chialvo, Emergent complex neural dynamics, Nat. Phys. 6 (2010) 744–750.
438 <https://doi.org/10.1038/nphys1803>.
- 439 [48] T. Zorick, J. Landers, A. Leuchter, M.A. Mandelkern, EEG multifractal analysis
440 correlates with cognitive testing scores and clinical staging in mild cognitive impairment,
441 J. Clin. Neurosci. 76 (2020) 195–200. <https://doi.org/10.1016/j.jocn.2020.04.003>.
- 442 [49] A. Das, A. Levina, Critical neuronal models with relaxed timescale separation, Phys.
443 Rev. X. 9 (2019) 21062. <https://doi.org/10.1103/PhysRevX.9.021062>.
- 444 [50] M. Mangalam, N.S. Carver, D.G. Kelty-Stephen, Global broadcasting of local fractal
445 fluctuations in a bodywide distributed system supports perception via effortful touch,
446 Chaos, Solitons & Fractals. 135 (2020) 109740.
447 <https://doi.org/10.1016/j.chaos.2020.109740>.
- 448 [51] N.S. Carver, D. Bojovic, D.G. Kelty-Stephen, Multifractal foundations of visually-guided
449 aiming and adaptation to prismatic perturbation, Hum. Mov. Sci. 55 (2017) 61–72.
450 <https://doi.org/10.1016/j.humov.2017.07.005>.
- 451 [52] C. Bell, N. Carver, J. Zbaracki, D. Kelty-Stephen, Nonlinear amplification of variability
452 through interaction across scales supports greater accuracy in manual aiming:
453 Evidence from a multifractal analysis with comparisons to linear surrogates in the
454 Fitts task, Front. Physiol. 10 (2019) 998.
455 <https://doi.org/10.3389/fphys.2019.00998>.

Fig. 1. Schematic of the task and effects of eye-to-target distance on postural sway. (A) The suprapostural viewing task of standing quietly with the eyes fixated at a distant visual element. (B and C) Visual angle gain for short vs. long eye-to-target distances along the anterior-posterior (AP) and medial-lateral (ML) axes. (D and E) Visual angle gain as a function of eye-to-target distance for different sway magnitudes. Closer targets increase AP sway, whereas farther targets increase ML sway.

Fig. 2. Multiscale PDF characterization of postural fluctuations in a representative participant maintaining quiet stance for 120 s with eyes closed, and eyes fixated at a point at a distance of 25, 50, 135, 220, and 305 cm from the eyes. From to bottom: CoP trajectories along the anterior-posterior (AP) and medial-lateral (ML) axes. CoP PED series. $\{ \Delta_s B(i) \}$ for $s = 0.5, 2, \text{ and } 8$ s. Standardized PDFs (in logarithmic scale) of $\{ \Delta_s B(i) \}$ for $s = 0.5, 2, \text{ and } 8$ s (from top to bottom), where σ_s denotes the SD of $\{ \Delta_s B(i) \}$ for $s = 0.5, 2, \text{ and } 8$ s (from top to bottom), where σ_s denotes the SD of $\{ \Delta_s B(i) \}$.

Fig. 3. Log-timescale dependence of the non-Gaussianity index λ_s^2 . Mean values of λ_s^2 for the participants standing quietly for 120 s in different viewing conditions. (A) Eyes closed. (B) Eyes fixated at 25 cm. (C) Eyes fixated at 50 cm. (D) Eyes fixated at 135 cm. (E) Eyes fixated at 220 cm. (F) Eyes fixated at 305 cm. Vertical bars indicate ± 1 SEM (N = 15). Notice that the λ_s^2 vs. log-timescale curves differ between the eye-closed condition all the eyes-open conditions except the eyes-fixated-at-50-cm condition.

475 **Fig. 4.** Log-timescale dependence of the mean proportion of trials with a lognormal
476 distribution for the participants standing quietly for 120 s in different viewing conditions. (A)
477 Eyes closed. (B) Eyes fixated at 25 cm. (C) Eyes fixated at 50 cm. (D) Eyes fixated at 135 cm.
478 (E) Eyes fixated at 220 cm. (F) Eyes fixated at 305 cm. Vertical bars indicate $\pm 1 SEM$ ($N = 15$).
479 Notice that the polynomial profiles of the growth of lognormality with log-timescale do not
480 differ between the eyes-closed condition and any of the eyes-open conditions.

Fig. S1. Schematic illustration of multiscale probability density function (PDF) analysis. (A) The first step is to generate $\{ B(t) \}$ by integrating $\{ b(t) \}$ after centering by the mean b_{ave} : $B(t) = \sum_{i=1}^{t/\Delta t} b(i\Delta t) - b_{\text{ave}}$. (B) A 3rd order polynomial detrends $\{ B(t) \}$ within k overlapping windows of length $2s$, s being the timescale. (C) Intermittent deviation $\Delta_s B(t)$ in k^{th} window from $1+s(k-1)$ to sk in the detrended time series $\{ B^d(t) = B(t) - f_{\text{fit}}(t) \}$ is computed as $\Delta_s B^d(t) = B^d(t+s) - B^d(t)$, where $1+s(k-1) \leq t \leq sk$ and $f_{\text{fit}}(t)$ is the polynomial representing the local trend of $\{ B(t) \}$, of which the elimination assures the zero-mean probability density function in the next step.

489

490 **Fig. S2.** Schematic illustration of non-Gaussianity index λ_s^2 . (A) The relationship between
 491 λ_s^2 and shapes of PDF plotted in linear-log coordinates for $\lambda_s^2 = 0.9, 0.33$, and 0.13 (from
 492 top to bottom). For clarity, the PDFs have been shifted vertically. As λ_s^2 increases, the PDF
 493 becomes increasingly peaked and fat-tailed. As λ_s^2 decreases, the PDF increasingly
 494 resembles the Gaussian (dashed line), assuming a perfect Gaussian as λ_s^2 approaches 0.
 495 (B) Cascade-type multiplicative processes yield the inverse relationship $\lambda_s^2 \sim -\log s$.