**Chemical composition and insecticidal activity of
Aeollanthus pubescens Benth leaf essential oil on the
malaria vector *Anopheles gambiae* (Diptera: Culicidae).
Bioinsecticidal properties of *Aeollanthus pubescens* leaf
essential oil on *Anopheles gambiae*.**

Roméo Barnabé Bohounton^{1,2}, Luc Salako Djogbénou^{2,3*}, Oswald Yédjinnavênan Djihinto²,
Oronce Sedjro-Ludolphe Dedome¹, Pierre Marie Sovegnon², Bruno Barea⁵, Aristide Adomou⁴,
Pierre Villeneuve⁵, Fidèle Paul Tchobo¹

¹ Laboratory of Study and Research of Applied Chemistry, Polytechnic School of Abomey-Calavi, 01 BP 2009, Cotonou, Benin.

² Laboratory of Vector-Borne Infectious Diseases, Institut Régional de Santé Publique/ University of Abomey Calavi, BP 384, Ouidah, Benin.

³ Department of Vector Biology, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool L3 5QA, United Kingdom.

⁴ Laboratoire de Botanique et Écologie Végétale (LaBEV), Faculté des Sciences et Techniques (FAST), University of Abomey-Calavi, Benin, 01 BP 4521 Cotonou.

⁵ Cirad, Persyst, UMR IATE, 34398 Montpellier, France.

***Corresponding author**

Mail: ldjogbenou22002@yahoo.fr (LSD)

Abstract

The use of synthetic insecticides is responsible for many cases of resistance in insects. Therefore, the use of natural molecules of ecological interest with insecticidal properties turns out to be an alternative approach to the use of synthetic insecticides. This study aims at investigating the larvicidal, adulticidal activity and the composition of the essential oil of *Aeollanthus pubescens* Benth on the major malaria vector *Anopheles gambiae*.

The leaves of *Aeollanthus pubescens* were collected in the South of the Republic of Benin. Three reference strains of *Anopheles gambiae* s.s. such as Kisumu, Kiskdr and Acerkis were used. The chemical composition of the essential oil was analysed by gas chromatography coupled to mass spectrometry. Larvae were exposed to the essential oil extract for 24 h. Adult mosquitoes were exposed to the fragment nets coated with the essential oil for 3 min. Larval mortality and adult survivorship were monitored.

Fourteen components were identified representing 98.31% of the total of oil. The major components were carvacrol (51.06 %), thymyle acetate (14.01 %) and γ -terpinene (10.60 %). The essential oil has remarkable larvicidal properties with LC₅₀ of 29.26, 22.65, and 28.37 ppm respectively on Kisumu, Acerkis and Kiskdr strains. With the fragment net treated at 165 $\mu\text{g}/\text{cm}^2$, the KDT₅₀ of both Acerkis (1.71 s, $p < 0.001$) and Kiskdr (2.67 s, $p < 0.001$) individuals were significantly lower than that of Kisumu (3.77 s). The lifespan of the three mosquito strains decreased respectively to one day for Kisumu ($p < 0.001$), two days for Acerkis ($p < 0.001$) and three days for Kiskdr ($p < 0.001$) compared to their control.

Our findings show that the *Aeollanthus pubescens* essential oil is an efficient larvicide and adulticide against malaria vector *Anopheles gambiae*. This bioinsecticidal activity is a promising discovery for the control of the resistant malaria-transmitting vectors.

Keywords: *Aeollanthus pubescens*, *Anopheles gambiae*, essential oil, bioinsecticidal activity.

Introduction

Vector-borne diseases remain the major causes of death in many tropical countries. The most important vector-borne diseases are malaria, lymphatic filariasis, dengue fever and yellow fever; the pathogens responsible for these diseases are transmitted by the arthropods vectors including mosquitoes (Diptera: Culicidae) which are the major agents [1,2]. Among the mosquito-borne infectious diseases, malaria is the most dreadful and the major public health concern in terms of the number of incidence, prevalence, morbidity and mortality in low-income countries of Africa, Asia, Latin America and beyond [3,4]. Despite the national malaria control programmes efforts, nearly 85% of malaria deaths occurred in 21 sub-Saharan African countries including the Republic of Benin [4]. Malaria is transmitted by the bites of parasite-infected *Anopheles* female mosquitoes that carry out the infection to humans [5].

So far, most malaria control programs have mainly relied on Artemisinin-based Combination Therapies (ACTs) for the treatment of diagnosed patients and the use of chemical compounds through Insecticides Treated Nets (ITNs) and Indoor Residual Spraying (IRS) for the prevention of human-vector contacts [6]. Nowadays, 14 insecticides belonging to four major classes of synthetic chemical insecticides are recommended by WHO Pesticide Evaluation Scheme (WHOPES) for IRS [7] and four insecticides, all from the pyrethroids group, are recommended for ITNs [8]. However, the overuse of these synthetic chemical insecticides has led to the emergence of resistant malaria vectors; and the frequency of insecticide resistance is widespread especially in African regions [9,10]. Also, these synthetic insecticides have been recognized to have adverse effects on non-targeted species and affect animal and plant biodiversity [11–13]. The indiscriminate use of these chemicals has also been shown to have severe effects on the environment and impacts on human health [14].

These facts combined with multidrug-resistance in malaria parasite [15] and the absence of an effective vaccine led scientists to focus on searching for environmentally-friendly vector

control alternatives with the aim of decreasing the selection pressure for insecticide resistance [16]. This kind of eco-friendly vector control alternatives could be achieved especially with insecticide from botanical sources. Indeed, they are potentially safer for human and the environment and have a minimal residual effect. They are more target-specific; less toxic to vertebrates, and more sustainable than their synthetic counterparts [17,18].

Beninese traditional medicine and pharmacopoeia medications are richly bio-diversified, which could be a great source for natural insecticides for malaria control [19]. Therefore, it appeared of interest to learn more about Beninese flora for its insecticidal activities. However, to our knowledge, only few studies have been conducted regarding the mosquitocidal activity of the extracts from Beninese plants species [20–22].

Aeollanthus pubescens (*A. pubescens*) Benth is an annual herbaceous plant distributed in many West African countries as well as in Benin [23]. It is commonly used by local populations as food and also for its medicinal properties [24,25]. The current study aims at determining the insecticidal potential of *A. pubescens* Benth leaf essential oil against Afrotropical malaria vector *Anopheles gambiae* under laboratory conditions by investigating safer alternatives for the management of malaria vector resistance compared to existing synthetic insecticides.

Material and methods

Plant material and extraction

The leaves of *A. pubescens* Benth (**Fig 1**) were collected in July 2014 at Covè in Benin and authenticated at the National Herbarium of University of Abomey-Calavi (UAC) where it was kept under voucher AAC 188/HNB.

The leaves were dried at 25 °C ±2 °C for 72 hours. Three (03) batches of 200 g of dried leaves were submitted to hydro-distillation in Clevenger apparatus at 100 °C for 2 hours. The distilled oil was dried using anhydrous sodium sulphate and transferred into an airtight amber-coloured

vial and stored at 4 °C until further use. The yields were averaged over the three experiments of the plant materials.

Fig 1: Areal part of *Aeollanthus pubescens* Benth harvested in Benin.

Chemical analysis of *Aeollanthus pubescens* Benth leaves essential oil

GC-FID analysis

The essential oil constituents were analysed by capillary gas chromatography coupled with flame ionization detection (GC-FID), equipped with a Supelco SPB-1 column (30 m×0.32 mm i.d, 0.25 µm film thickness). One (1) µL of the essential oil diluted in chloroform were directly injected into the GC. Helium was used as carrier gas with the flow rate of 6 mL/min and the splitting ratio of 1/17. The inlet temperature was 250 °C/min, 200-310 °C at 20 °C/min, and then maintained at 310 °C for 2 min.

GC-MS analysis

Capillary gas chromatography coupled with a mass spectrometry detector (GC-MS) was used on a TR-1MS column (30 m x 0.25 mm i.d., 0.25 µm film thickness). An electron impact was used with ionization energy of 70 eV. Helium was used as the carrier gas at a flow rate of 0.6 mL/min, and the splitting ratio 1/17. The temperature settings were as follows: 70-200 °C at 10 °C/min, 200-300 °C at 20 °C/min, and then maintained at 300°C for 1 min. Inlet and MS transfer line temperatures were set at 250 and 320 °C, respectively. All apparatus and accessories were from Thermo Scientific (Courtaboeuf, France) and software controlled data processing (Chromocard and XCalibur). The identification of the essential oil constituents was based on the comparison of their retention times and their Kovats retention indexes relative to (C₈-C₂₀)

n-alkanes. Whenever possible, identifications were based on mass spectra of the authentic standard compounds. Otherwise, identifications were performed using published data [26] and comparison with the NIST mass spectral library.

Mosquito strains

Three *Anopheles gambiae* s.s (*An. gambiae*) laboratory strains that were already available at the insectarium of the laboratory of Vector-Borne Infectious Diseases at the Institut Régional de Santé Publique Alfred Quenum (IRSP-AQ) of the University of Abomey-Calavi in Ouidah (Benin) were used in this study. Kisumu strain originating from Kenya is a reference strain susceptible to all insecticides [27]. Acerkis, which is resistant towards both organophosphate and carbamate based insecticides and is homozygous for (G119S) mutation [28]. Kiskdr, which is homozygous for *kdr^R* (L1014F) and confers resistance to pyrethroids and DDT [29]. Both AcerKis and Kiskdr were supposed to share the same genetic background as the Kisumu strain but differ by the presence of resistance alleles.

The colonies of each strain were maintained at the insectarium under optimum conditions (25-27°C temperature and 70-80% relative humidity). The third instar larvae, as well as adult females of 3-5 days old of each mosquito strains were used for the bioassays.

Bioassays

Larval bioassay

The larvicidal properties of the essential oil were conducted according to the standard method recommended by the World Health Organization [30] with slight modifications. Since the essential oil does not dissolve in water, six different concentrations (1000, 2000, 3000, 4000, 5000 and 6000 ppm) of the essential oil were prepared in ethanol 96%. Twenty-five (25) third instar larvae of each strain were gently transferred gently into plastic beaker containing 99 mL

of water and 1 mL of each prepared concentration was added to obtain test solutions of 10, 20, 30, 40, 50 and 60 ppm respectively. In each round of the experiment, larvae were exposed for 24 hours at $26 \pm 2^\circ\text{C}$ without any food provided to them. After exposure, the larval mortality was recorded. Larvae were considered dead when they were not able to move or swim actively when touched. For each strain, four replicates were performed for a total of 100 larvae per concentration. The control group consisted of batches of larvae exposed to water and the solvent alone (ethanol). In total, three different experiments were conducted on three different days.

Adult Bioassay

Impregnation of mosquito nets with the essential

oil

Fragments of the non-insecticide treated net (13 cm x 13 cm; 169 cm^2) corresponding to the surface necessary for the base of the cone were cut and coated with the essential oil. Concentration ranges corresponding to 10, 20 and 30 fold of that of the standard PermaNet 2.0 (55 mg/m^2 of deltamethrin) were chosen for the impregnation after preliminary doses screening. Then, the essential oil which was yellow with a distinct and sharp odour, was incorporated into the net fragment fibres at 55, 110 and $165 \mu\text{g/cm}^2$.

The masses of essential oil proportional to the net area (169 cm^2) per concentration were determined: 9.3, 18.6 and 27.9 mg for the impregnation at 55, 110 and 165 ng/cm^2 respectively. A volume of 1.5 mL of ethanol HPLC grade was poured into a Petri dish containing the mass of essential oil corresponding to a given concentration. After complete dissolution, the fragment of the mosquito net was soaked in the mixture. The impregnated fragment nets were left to dry at room temperature for 5 minutes to allow the essential oil to adhere to the mosquito net and to evaporate the ethanol completely. After drying treated fragment nets were stored in the dark and used 2 to 4 hours later. All coated fragment nets used during the day were treated in the

morning at the same time. The net of the same size were also treated with 1.5 mL of solvent and was used as control.

Cone test

The cone test was used to assess the effectiveness of the essential oil on the adult mosquitoes. The cone test is an adaptation of the WHO cone bioassay with the following modification: During the assay, the test operator holds a forearm behind the cone to provide a host (**Fig 2**). Unfed 3-5 days old female mosquitoes of Kisumu, Acerkis and Kiskdr strains were used in the test. On the day of testing, female were starved for 4 hours before testing. Groups of five female mosquitoes were placed into plastic cups and moved into the testing room one hour before testing begins to allow the mosquitoes to acclimatise to room conditions. The fragment nets for test or control were placed over a dedicated hole on the Perspex boards and secured using a clear tape. A second Perspex board was laid on the first board creating a test/control net “sandwich” between the two boards. The cone was placed over the net and plugged above with a piece of parafilm. A batch of 5 mosquitoes was transferred into the cone with the operator’s forearm in position. Mosquitoes were then exposed to the fragments for three (03) minutes. Ten (10) replicates of batches of 5 mosquitoes of each strain were run per concentration of impregnated nets.

Fig 2. Cone test equipment.

Monitoring of the lethal effect of mosquito exposure to the essential oil.

After exposure, mosquitoes were removed from the cone, transferred into a recovery cups and provided with 10% of honey solution soaked on a cotton pad. Mosquito knockdown was

recorded at 60 minutes post-test. Mosquito mortality was then recorded every day until the death of the last female of each mosquito strain.

Data analysis

The analysis of dose-mortality responses in larval bioassays was performed using the BioRssay script version 6.2 [31] in R software Version 3.0 [32]. This script computes the lethal concentrations inducing 50% (LC₅₀) and 95% (LC₉₅) mortality recorded in each strain and the associated confidence intervals; the resistance ratios, i.e. RR₅₀ or RR₉₅ (LC₅₀ or LC₉₅ in each strain divided respectively by the LC₅₀ or LC₉₅ of the reference strain) and their 95% confidence intervals. The times at which 50% or 95% of mosquitoes fell on their back or their side, i.e. knockdown time (KDT₅₀ or KDT₉₅) and their 95% confidence intervals were estimated after probit regression in R software using the package ‘ecotox’ [33] based on the method described by Finney [34]. The mosquito survival after exposure to the essential oil impregnated net was analysed by Kaplan–Meier survival curves using GraphPad Prism 8.0.2 software (San Diego, California USA). The Log-rank test was performed to evaluate the difference in survival between the strains. All statistical analyses were set at a significance threshold of $p < 0.05$.

Results

Extract yield (dry weight %) and Chemical composition of *Aeollanthus pubescens* Benth leaf essential

The percentage yields of essential oil obtained from the hydro-distillation of the leaves of *A. pubescens* was 0.30 ± 0.02 (w/w based on fresh leaves). From the chemical composition of the essential oil of *A. pubescens* Benth (**Table 1**), fourteen compounds were identified, accounting for 98.31% of the crude essential oil’s mass. The essential oil of *A. pubescens* aerial part had higher oxygenated monoterpenes (60.4%) than monoterpene hydrocarbons (22.39%) and sesquiterpene hydrocarbons (15.52%) (**Table 1**). The major component of the essential oil was

carvacrol (51.06%), followed by other components thymol acetate (14.01%), γ -terpinene (10.60%), O-cymene (8.40%) and thymol (5.46%). The percentage of composition of the remaining nine compounds ranged from 0.19 to 2.02% (**Table 1**).

Table 1: Chemical composition of the *Aeollanthus pubescens* Benth essential oil.

Peak No	RI	Components	Area (%)
1	940	α -Pinene	0.58
2	986	Myrcene	2.02
3	1005	Lumicolchicine	0.19
4	1020	o-Cymene	8.40
5	1031	1,8-cineole	0.60
6	1057	γ -Terpinene	10.60
7	1088	Linalool	0.88
8	1162	Borneol	1.40
9	1173	Terpin-4-ol	1.60
10	1273	Thymol	5.46
11	1284	Carvacrol	51.06
12	1359	Thymol acetate	14.01
13	1488	β -Cubebene	0.24
14	1503	Acid [(2,4,6-triethylbenzoyl) thio]acetic	1.26
		Total identified (%)	98.31
		Sesquiterpenes hydrocarbons	15.52
		Monoterpenes hydrocarbons	22.39
		Oxygenated monoterpenes	60.4

RI: relative retention indices as determined on an HP-1 column using the homologous series of n-alkanes.

Toxicity of the essential oil on *Anopheles gambiae* s.s larvae

We carried out a total of 24 larval bioassays with four replicates for each of the three mosquito strains. The strains Acerkis larvae ($LC_{50} = 22.65$ ppm) were significantly more susceptible to the essential oil compared to Kiskdr ($LC_{50} = 28.37$ ppm, $p < 0.001$) and Kisumu ($LC_{50} = 29.26$ ppm, $p < 0.001$) (**Table 2**). However, Kisumu and Kiskdr larvae susceptibility was not significantly different ($p = 0.69$). All resistance ratio (RRs) calculated were not significantly different to 1 (the RR_{50} was 0.77 and 0.96 in Acerkis and Kiskdr respectively).

Table 2: Toxicity against *Anopheles gambiae* larvae after 24 h exposure.

Mosquito strains	LC ₅₀ (ppm)	95% C.I	RR ₅₀	95% C.I	LC ₉₅ (ppm)	95% C.I	Slope ±S.E
Kisumu	29.26	27.73 - 30.73	-	-	47.29	44.12 - 51.71	7.88 ± 0.58
Acerkis	22.65	20.18 - 24.97	0.77	0.63-0.96	44.80	39.48 - 53.49	5.55 ± 0.57
Kiskdr	28.37	26.84 - 29.83	0.96	0.79-1.19	48.00	44.67 - 52.59	7.19 ± 0.51

LC_{50/95}: lethal concentrations; S.E: standard error; C.I: Confidence interval; RR₅₀ is resistance ratio at LC₅₀: LC₅₀ (resistant strain)/ LC₅₀ (Kisumu).

Adulticidal activity of the essential oil

Knockdown time of essential oil impregnated net on *Anopheles gambiae* strains

The average time estimated for knockdown 50% (KDT₅₀) or 95% (KDT₉₅) of adult *An. gambiae* females of each strain decreased with the increasing treatment concentration. The KDT₅₀ was less than 4 seconds for all mosquito strains in contact with fragment net treated at 165 µg/cm² (3.77 s for Kisumu; 1.71 s for Acerkis and 2.67 s for Kiskdr), which were significantly lower than that recorded with the lowest essential oil treatment (55 µg/cm²) (Kisumu: 22.06 s, $p < 0.001$; Acerkis: 291.72 s, $p < 0.001$; Kiskdr: 591.63s, $p = 0$) (Table 3). At the highest treatment concentration (165 µg/cm²), both Acerkis and Kiskdr mosquitoes were quickly knocked down (KDT₅₀ of 1.71 s, $p < 0.001$ and 2.67 s, $p < 0.001$ respectively) than Kisumu individuals (KDT₅₀: 3.77 s).

However, the highest knocked down times values were observed for the Kiskdr (KDT₅₀ > 597s) and Acerkis (KDT₅₀ > 291s) females exposed to the essential oil at 55 µg/cm².

Table 3: Probable time for 50 and 95% knockdown of *Anopheles gambiae* s.s. strains per fragment net treatment.

Strain	Essential oil treatment (µg/cm ²)	KDT ₅₀ (s)	95% CI	KDT ₉₅ (s)	95% CI
--------	---	-----------------------	--------	-----------------------	--------

Kisumu	55	22.06	[20.03-23.82]	45.31	[40.51-53.26]
	110	4.74	[4.54 - 4.93]	6.27	[5.92 - 6.8]
	165	3.77	[3.55 - 3.98]	5.65	[5.21 - 6.36]
Acerkis	55	291.72	[280.64 - 302.38]	373.70	[355.49 - 400.70]
	110	4.63	[4.44 - 4.82]	6.17	[5.79 - 6.77]
	165	1.71	[1.52 - 1.89]	3.52	[3.06 - 4.29]
Kiskdr	55	591.63	[576.42-607.56]	813.01	[777.61-859.02]
	110	197.44	[185.02-209.26]	308.68	[285.79-341.92]
	165	2.67	[2.46-2.86]	4.40	[3.99-5.09]

(s): second; **KDT₅₀** and **KDT₉₅**: Knock-down times for 50 and 95% of adult mosquitoes after three minutes of exposure to impregnated fragment net with the essential oil in cone test; **CI**: Confidence Interval.

Mortality induced by essential oil the impregnated net on *Anopheles gambiae* strains

Overall, the three essential oil treatments had decreased significantly the survival of all mosquito strains after exposure. For the essential oil coating at 165 µg/cm², the longevity of the three mosquito strains decreased significantly from twenty-four (24) days for Kisumu, twenty-five (25) days for Acerkis and twenty-six (26) days for Kiskdr in control groups to respectively one (01) days for Kisumu ($p < 0.001$), two (02) days for Acerkis ($p < 0.001$) and three (03) days for Kiskdr ($p < 0.001$) in exposed groups (**Fig 3C**). With the net treated at 110 µg/cm², Kisumu females longevity was significantly reduced by 21 days compared to that recorded with the 55 µg/cm² treatment (by 14 days; $p < 0.001$) (**Figure 3A and 3B**). With each of these two treatments, no significant effect was observed on the longevity of Kiskdr ($p = 0.8$).

Fig 3. Survivorship of mosquitoes post exposure. Each of mosquito strains was followed up after exposure to fragment net impregnated at 55 µg/cm² (**A**); 110 µg/cm² (**B**) and 165 µg/cm² (**C**).

Discussion

The increasing number of reports of natural mosquito resistance to the existing synthetic insecticide has strengthened the focus on searching for environmentally-friendly insecticide compounds for vector control strategy. This is a beneficial alternative as essential oils represent a rich source of bioactive compounds that are biodegradable into non-toxic products and due to their natural synergism, they reduce the risk of the development of resistance in the vectors [35]. Besides, essential oils are known as nucleophilic in nature and hinder efficiently with a range of biological processes (metabolic, physiological, biochemical and behavioural) in insects [36–38]. This study is the first report of the larvicidal and adulticidal activity of the *A. pubescens* Benth leaves essential oil on the major African malaria vector *An. gambiae* s.s. The insecticidal properties of the essential oil of *A. pubescens* leaves were carried out in the laboratory using immature and adult stages of *An. gambiae* mosquitoes.

The essential oil isolated by hydro-distillation process from the aerial parts of *A. pubescens* collected during this study was found to be a pale yellow liquid, with a yield of $0.30 \pm 0.02\%$. The same oil yield (0.3%) was reported for the fresh aerial material of *A. pubescens* collected from Togo with the same isolation technique [39]. Agbodan et al. [40] have also reported similar oil yield (0.20%) by steam distillation of the fresh leaves of *A. pubescens* collected from Togo. However, a slight increase of the oil yield (1.2 to 1.5%) was reported for the fresh leaves collected in other regions in Benin [25]. These observations suggest that *A. pubescens* oil yield does not depend on the drying step, is neither specific to the extraction technique but may depend on the oil constituents (chemotype). However, the increase in yield of extracted oil could be achieved using the supercritical fluid extraction which will also improve the quality of the extract [41].

The chemical composition analysis of the essential oil of *A. pubescens* aerial parts showed a quantitative variation of all and the major compounds identified. The chemical analysis has

displayed the presence of 14 compounds. Carvacrol was the major compound representing 51.06% of the constituents, followed by the thymol acetate (14.01%) and γ -terpinene (10.60%). This oil composition is characteristic of the carvacrol chemotype. Overall, five (05) different chemotypes were identified from the essential oil of *A. pubescens* aerial part from Togo: i) the thymol chemotype: containing 46.3-58% of thymol; ii) the carvacrol chemotype: 58.21% of Carvacrol; iii) the carvacrol and thymol chemotype: 41% of Carvacrol and 27% of thymol, iv) the carvacrol and thymol acetate chemotype: 55.36% of Carvacrol and 35.05% of thymol acetate, and v) the D-fenchone chemotype: 83.69 % of D-fenchone [39,40,42]. However, for the same plant material collected in the central regions of Benin, Alitonou et al. [25] carried out only the thymol (63%) and carvacrol (51.1%) chemotypes. This variability of the chemical composition of essential oils of the same plant material could be due to many factors including: the bioclimate, soil composition, the harvesting period, the geographical location, the degree of maturity of the plant, the seasonal variation and even the plant genetic background [43].

The results from larvae bioassays using the laboratory colonies of *An. gambiae* showed that the essential oil of *A. pubescens* aerial parts is highly active ($LC_{50} < 50$ ppm) on the specimens of these strains according to the classification of Komalamisra et al. [44]. Surprisingly, the **RR**₅₀ for Acerkis (0.77) was less than one (01), indicating that Acerkis strain is more susceptible to the oil than Kisumu strain while no apparent difference was found between Kiskdr and Kisumu (**RR**₅₀ very close to 1). These results suggest that the essential oil had a promising larvicidal property with low LC_{50} values. The significant activity on the resistant strain over the reference susceptible Kisumu mosquitoes indicates that the essential oil does not affect one of the former target sites (Ace-1^R and Kdr^R (L1014F) alleles) represented in the corresponding strain. Therefore, it could be implied that its mode of action is different from that of pyrethroids, organophosphates and carbamates. However, in the context of the increasing insecticide

resistance in natural mosquito populations, it will be interesting to investigate the bioactivity of this essential oil on field-collected larvae and field caught-adults.

Other biological activities of *A. pubescens* essential oil were reported including antioxidant [25,40], antibacterial [45], but no report was made on its mosquitocidal activity so far. In addition, there is still a lack of information on the bioinsecticidal property of the other plant species belonging to the genus *Aeollanthus*. However, previous studies have investigated the larvicidal activity of the essential oil of plants belonging to the same family (Lamiaceae). Indeed, Tchoumboungang et al. [46] showed that *Ocimum canum*, *Ocimum gratissimum* and *Thymus vulgaris* displayed respectively LC₅₀ value of 201, 180 and 119 ppm on field-collected *An. gambiae* larvae. These values are higher than those recorded in our study. This variation could be due to the difference in the oils chemical composition and the genetic background of the larvae strains used. Moreover, essential oils from other *Lamiaceae* species (*Plectranthus amboinicus* and *Plectranthus mollis*) were found to be active against *Anopheles stephensi* larvae with LC₅₀ value less than 50 ppm [47,48]. These findings suggest that essential oils from *Lamiaceae* plant species could be a potential source of environmental eco-friendly mosquitocidal agents. In our study, *A. pubescens* essential oil was dominated by monoterpenes (oxygenated and hydrocarbons) which represented 82.79% of the oil. Other plants species with similar major constituents have been reported to be active against *An. gambiae* larvae. Ollengo et al. [49] reported that *Clausena anisata* containing 56.7% of monoterpenes revealed a potential larvicidal activity against *An. gambiae* (LC₅₀ = 75.96 ppm) [50]. Also, Wangrawa et al. [51] demonstrated that *Lantana camara* essential oil with 70.5% of monoterpenes showed differential larval mortalities on both the laboratory and the field strains of *An. gambiae*. The high proportion of monoterpenes in the essential oil could be correlated to the observed bioactivity. Indeed, many studies reported the larvicidal effect of monoterpenes against mosquitoes strains [52–54]. Therefore, it would be interesting to evaluate further the toxicity

of the monoterpenes isolated from *A. pubescens* areal part on mosquito larvae in both laboratory and field trials.

We assume that Carvacrol, the main compound (51.06%) found within the monoterpenes in our *A. pubescens* oil extract, might be the main contributor of the observed activity against *An. gambiae* larvae. Carvacrol is well known for its larvicidal property against both *Anophelinae* (*Anopheles stephensi*, *Anopheles subpictus*) [55] and *Culicidae* (*Aedes aegypti*, *Culex quinquefasciatus* and *Culex tritaeniorhynchus*) [55–58].

Essential oils are mixtures of volatile compounds and due to the antagonistic or synergistic phenomena, the bioactivity of the crude oil extract in some cases is lower or higher than those of purified compounds. For instance, Evergetis et al. [59] demonstrated that larvicidal activity of the essential oil of *Origanum vulgare* ssp against *Aedes albopictus* ($LC_{50} = 30.1$ ppm) is lower than that of its major component, the pure Carvacrol ($LC_{50} = 13.1$ ppm), accounted for 88.7% of the oil. The same trend was noticed with the leaf essential oil of *Coleus aromaticus* which displayed lower toxicity than its major component Carvacrol against *Anopheles stephensi* larvae [60]. This opens perspectives for further investigations to evaluate the larvicidal efficacy of the Carvacrol in comparison to that of the crude oil extract. It is well known that monoterpenes from essential oils could act by absorption through the cuticle, or via the respiratory tract or by ingestion via the gastrointestinal tract [61–64]. Besides, several monoterpenes were reported to target primarily the cholinergic, octopamenergic and GABA neurosystems in insects [65]. One or a combination of these mechanisms might be the pathway of mortality induction by the *A. pubescens* oil. In this study, the reference resistant strain *Acerkis* larvae harboring *ace-1^R* allele coding for the insensitive acetylcholinesterase enzyme was the most susceptible to our essential oil. This indicates that the essential oil overcomes the target site modification resistance mechanism and therefore appears as a hopeful alternative tool for vector control programs.

Among the vector life-history traits, mosquito survival is strongly associated with the malaria transmission intensity [66]. Thus, this study also investigated the effect of exposure to various doses of *A. pubescens* oil on adults *An. gambiae* survival. The *A. pubescens* essential oil reduced significantly the lifespan of the three mosquitoes strains exposed to the fragment net at 165 $\mu\text{g}/\text{cm}^2$. None of the three mosquito strains was able to survive after 72 hours. This observation suggests that even the resistance mosquitoes (Acerkis and Kiskdr) could not survive long enough to allow the extrinsic incubation period of the *Plasmodium* parasites if they ingested a gametocyte infected blood meal. Overall, the drastic reduction in daily survival of mosquitoes observed with the oil treatment at 165 $\mu\text{g}/\text{cm}^2$ would contribute to a reduction in vectorial capacity in a typical endemic setting and therefore will lead to a reduction in parasite transmission according to the Ross-MacDonald model [67]. This is a promising finding for the management of the resistant malaria-transmitting vectors. Spray-type solution formulations could be made towards the development of botanical insecticides for the use of an integrated approach with the existing conventional vector control strategies. However, the results were obtained using mosquito strains in which only one resistance mechanism is present. It will be interesting to evaluate further the survivorship of the natural mosquito populations where several resistance mechanisms could coexist. The susceptibility of *Plasmodium* infection following essential oil exposure is also another promising parameter to be evaluated.

During the experiment, we observed that the legs of mosquitoes were detached from their bodies when exposed to the net coated at 165 $\mu\text{g}/\text{cm}^2$. To our knowledge, this phenomenon has not been observed yet. This effect of the essential oil may be caused by its main constituent, the Carvacrol. However, such mechanism is so far unexplored. Possible neurotoxicity in insects could have been easily overlooked. Thus, investigations around the mechanisms used by this compound is urgently sought. The mosquito legs loss suggests that the essential oil could

interfere with the insect locomotor system and even the nervous system leading to the death in the following days.

At the doses 55 and 110 $\mu\text{g}/\text{cm}^2$ of the essential oil, the lifespan of mosquitoes is fourteen (14) days maximum for both resistant Acerkis and Kiskdr strains. During this time period, mosquitoes might still be able to reproduce. Therefore, further studies are needed to assess the blood-feeding success, the fecundity and the fertility of mosquitoes following exposure. This could lead to highlight putative detrimental effects of the essential oil exposure that could also hamper the vectorial competence of the mosquitoes.

Conclusion

The present study paves the way to develop new and safer natural insecticide and repellent against malaria mosquito vectors. The *A. pubescens* Benth essential oil is proved to be an efficient larvicide and adulticide against malaria vector *An. gambiae*. This opens the perspectives for implementing sustainable control of mosquito populations that are resistant to the current existing synthetic insecticides. The larval and adult vector control with the essential oil could be considered in an integrated fashion to the existing malaria control strategies. Further studies are needed to help in designing *A. pubescens* essential oil formulation that would potentially increase its efficacy on *An. gambiae* and its cost-effectiveness.

Acknowledgments

The authors thank Geraldine Foster from Liverpool School of Tropical Medicine (LSTM) for the material support. We are grateful to Ms Marie Joelle Fanou for her assistance in insectary work; Mrs Laurette Djossou for her help during data acquisition; Mrs Emilienne Fiogbe and Mr Jean-Louis Amoussou for their helpful proofreading of the manuscript. The authors also acknowledge Wellcome Trust for financial support to LSD (grant 109917/Z/15/Z).

Author Contributions

Conceived and designed the experiments: Roméo Barnabé Bohounton, Luc Salako Djogbénou, Pierre Villeneuve, Fidèle Paul Tchobo.

Performed the experiments: Roméo Barnabé Bohounton, Pierre Marie Sovegnon, Bruno Barea.

Data analysis: Roméo Barnabé Bohounton, Oswald Yédjinnavênan Djihinto, Oronce Sedjro-Ludolphe Dedome.

Writing – original draft: Roméo Barnabé Bohounton, Oswald Yédjinnavênan Djihinto.

Writing – review & editing: Roméo Barnabé Bohounton, Oswald Yédjinnavênan Djihinto, Aristide Adomou, Luc Salako Djogbénou, Fidèle Paul Tchobo.

References

1. Leitner WW, Wali T, Kincaid R, Costero-Saint Denis A. Arthropod Vectors and Disease Transmission: Translational Aspects. PLoS Negl Trop Dis. 2015;9. doi:10.1371/journal.pntd.0004107
2. World Health Organization. Vector-borne diseases. 2020 [cited 26 Aug 2020]. Available: <https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases>
3. Karunamoorthi K. Global malaria eradication: is it still achievable and practicable? <http://www.ncbi.nlm.nih.gov/nlmcatalog/101566132>. In: Peterson AM, Calamandrei GE, editors. Malaria: etiology, pathogenesis and treatments. <http://www.ncbi.nlm.nih.gov/nlmcatalog/101566132>. New York: Nova Science Publishers; 2012.
4. World Health Organization. World malaria report 2019. 2019. Available: <https://www.who.int/publications/i/item/world-malaria-report-2019>
5. Institute of Medicine (US) Committee on the Economics of Antimalarial Drugs, Arrow KJ, Panosian C, Gelband H. The Parasite, the Mosquito, and the Disease. Saving Lives, Buying Time: Economics of Malaria Drugs in an Age of Resistance. National Academies Press (US); 2004. Available: <https://www.ncbi.nlm.nih.gov/books/NBK215619/>
6. Karunamoorthi K. Vector control: a cornerstone in the malaria elimination campaign. Clin Microbiol Infect. 2011;17: 1608–1616. doi:10.1111/j.1469-0691.2011.03664.x
7. World Health Organization. WHO Recommended Insecticides for Indoor Residual Spraying against Malaria Vectors. 21 Sep 2018 [cited 27 Aug 2020]. Available:

https://www.who.int/neglected_diseases/vector_ecology/vector-control/Insecticides_IRS_22_September_2018.pdf?ua=1

8. Nogueira Barradas T, Perdiz Senna J, Ricci Júnior E, Regina Elias Mansur C. Polymer-based Drug Delivery Systems Applied to Insects Repellents Devices: A Review. CDD. 2016;13: 221–235. doi:10.2174/1567201813666151207110515
9. Ranson H, Abdallah H, Badolo A, Guelbeogo WM, Kera-Hinzoumbé C, Yangalbé-Kalnoné E, et al. Insecticide resistance in *Anopheles gambiae*: data from the first year of a multi-country study highlight the extent of the problem. Malaria Journal. 2009;8: 299. doi:10.1186/1475-2875-8-299
10. Mouhamadou CS, de Souza SS, Fodjo BK, Zoh MG, Bli NK, Koudou BG. Evidence of insecticide resistance selection in wild *Anopheles coluzzii* mosquitoes due to agricultural pesticide use. Infectious Diseases of Poverty. 2019;8: 64. doi:10.1186/s40249-019-0572-2
11. Newsom LD. Consequences of Insecticide Use on Nontarget Organisms. Annual Review of Entomology. 1967;12: 257–286. doi:10.1146/annurev.en.12.010167.001353
12. Smith TM, Stratton GW. Effects of synthetic pyrethroid insecticides on nontarget organisms. In: Gunther FA, editor. Residue Reviews. New York, NY: Springer New York; 1986. pp. 93–120. doi:10.1007/978-1-4612-4934-4_4
13. Ware GW. Effects of pesticides on nontarget organisms. In: Gunther FA, Gunther JD, editors. Residue Reviews. New York, NY: Springer New York; 1980. pp. 173–201. doi:10.1007/978-1-4612-6107-0_9
14. Mahmood I, Imadi SR, Shazadi K, Gul A, Hakeem KR. Effects of Pesticides on Environment. In: Hakeem KR, Akhtar MS, Abdullah SNA, editors. Plant, Soil and Microbes. Cham: Springer International Publishing; 2016. pp. 253–269. doi:10.1007/978-3-319-27455-3_13
15. Rout S, Mahapatra RK. Plasmodium falciparum: Multidrug resistance. Chem Biol Drug Des. 2019;93: 737–759. doi:10.1111/cbdd.13484
16. Benelli G, Jeffries CL, Walker T. Biological Control of Mosquito Vectors: Past, Present, and Future. Insects. 2016;7. doi:10.3390/insects7040052
17. Shaalan EA-S, Canyon D, Younes MWF, Abdel-Wahab H, Mansour A-H. A review of botanical phytochemicals with mosquitocidal potential. Environment International. 2005;31: 1149–1166. doi:10.1016/j.envint.2005.03.003
18. Thomson M, Vijan A. Environmental Friendly Bio-pesticides: A Review. Research & Reviews: Journal of Agriculture and Allied Sciences. 2016;5: 31–39.
19. Hermans M, Akoègninou A, van der Maesen J. Medicinal plants used to treat malaria in Southern Benin. Econ Bot. 2004;58: S239–S252. doi:10.1663/0013-0001(2004)58[S239:MPUTTM]2.0.CO;2
20. Guevara N, Wotto V, Jean-Pierre N, Avlessi F, Martin A, Sohounhloue D. Insecticidal activities of essential oils extracted from three species of poaceae on *Anopheles*

- 473 gambiaespp, major vector of malaria. Scientific Study & Research Chemistry & Chemical
474 Engineering, Biotechnology, Food Industry. 2010;11.
- 475 21. Bossou AD, Mangelinckx S, Yedomonhan H, Boko PM, Akogbeto MC, De Kimpe N, et
476 al. Chemical composition and insecticidal activity of plant essential oils from Benin
477 against *Anopheles gambiae* (Giles). *Parasites Vectors*. 2013;6: 337. doi:10.1186/1756-
478 3305-6-337
- 479 22. Akpo AA, Chougourou DC, Djènantin A, Dossou J, Anagonou R, Akogbéto M. Etude De
480 L'efficacité Du Cashew Nut Shell Liquid (CNSL) De *Anacardium Occidentale* L. Extrait
481 À Froid Sur Le Contrôle De *Anopheles Gambiae* S.L Résistant Aux Pyréthrinoides.
482 *European Scientific Journal*, ESJ. 2017;13: 249–249. doi:10.19044/esj.2017.v13n24p249
- 483 23. Adjanohoun É, Agence de Coopération Culturelle et Technique, editors. Contribution aux
484 études ethnobotaniques et floristiques au Togo. Paris: Agence de Coopération Culturelle
485 et Technique; 1987.
- 486 24. Kafoutchoni KM, Idohou R, Egeru A, Salako KV, Agbangla C, Adomou AC, et al.
487 Species richness, cultural importance, and prioritization of wild spices for conservation in
488 the Sudano-Guinean zone of Benin (West Africa). *J Ethnobiol Ethnomed*. 2018;14.
489 doi:10.1186/s13002-018-0267-y
- 490 25. Alitonou G, Tchobo F, Avlessi F, Sohounhloue DK, Menut C. *Aeollanthus pubescens*
491 Benth. from Benin: A Potential Source of Essential Oil with High Antiradical Efficiency.
492 *Journal of Essential Oil Bearing Plants*. 2013;16: 308–314.
493 doi:10.1080/0972060X.2013.813203
- 494 26. Adams RP, Adams RP. Identification of essential oil components by gas
495 chromatography/quadrupole mass spectroscopy. Carol Stream, Ill.: Allured Pub. Corp.;
496 2004.
- 497 27. Shute GT. A Method of Maintaining Colonies of East African Strains of *Anopheles*
498 *Gambiae*. *Annals of Tropical Medicine & Parasitology*. 1956;50: 92–94.
499 doi:10.1080/00034983.1956.11685743
- 500 28. Djogbénou L, Weill M, Hougard JM, Raymond M, Akogbéto M, Chandre F.
501 Characterization of insensitive acetylcholinesterase (ace-1R) in *Anopheles gambiae*
502 (Diptera: Culicidae): resistance levels and dominance. *J Med Entomol*. 2007;44: 805–810.
503 doi:10.1603/0022-2585(2007)44[805:coiaai]2.0.co;2
- 504 29. Alout H, Ndam NT, Sandeu MM, Djégbe I, Chandre F, Dabiré RK, et al. Insecticide
505 Resistance Alleles Affect Vector Competence of *Anopheles gambiae* s.s. for *Plasmodium*
506 *falciparum* Field Isolates. *PLOS ONE*. 2013;8: e63849.
507 doi:10.1371/journal.pone.0063849
- 508 30. World Health Organization. Guidelines for laboratory and field testing of mosquito
509 larvicides. 2005 [cited 28 Aug 2020]. Available:
510 <https://apps.who.int/iris/handle/10665/69101>
- 511 31. Milesi P, Labbé P. BioRssay: A R Script for Bioassay Analyses v. 6.2. Montpellier,
512 France; 2015. Available:

- 513 <http://www.wisem.univ-montp2.fr/recherche/equipes/genomiquedeladaptation/personnel/lab>
514 be-pierrick/
- 515 32. R Core Team. R: A language and environment for statistical computing. Vienna, Austria:
516 R Foundation for Statistical Computing; 2017. Available: <https://www.R-project.org/>
- 517 33. Hlina BL. Package “ecotox.” Cambridge [Eng.]: University Press; 2020.
- 518 34. Finney DJ, Tattersfield F. Review of Probit Analysis: A Statistical Treatment of the
519 Sigmoid Response Curve. *Journal of the Royal Statistical Society*. 1947;110: 263–266.
520 doi:10.2307/2981407
- 521 35. Park H-M, Kim J, Chang K-S, Kim B-S, Yang Y-J, Kim G-H, et al. Larvicidal activity of
522 Myrtaceae essential oils and their components against *Aedes aegypti*, acute toxicity on
523 *Daphnia magna*, and aqueous residue. *J Med Entomol*. 2011;48: 405–410.
524 doi:10.1603/me10108
- 525 36. Benelli G. Plant-borne ovicides in the fight against mosquito vectors of medical and
526 veterinary importance: a systematic review. *Parasitol Res*. 2015;114: 3201–3212.
527 doi:10.1007/s00436-015-4656-z
- 528 37. Pavela R. Essential oils for the development of eco-friendly mosquito larvicides: A
529 review. *Industrial Crops and Products*. 2015;76: 174–187.
530 doi:10.1016/j.indcrop.2015.06.050
- 531 38. Brattsten LB. Cytochrome P-450 Involvement in the Interactions Between Plant Terpenes
532 and Insect Herbivores. *Plant Resistance to Insects*. AMERICAN CHEMICAL SOCIETY;
533 1983. pp. 173–195. doi:10.1021/bk-1983-0208.ch010
- 534 39. Sanda K, Koba K, Walla BA, Akpagana K, Garneau F-X, Gagnon H, et al. *Aeollanthus*
535 *pubescens* Benth. A Potential Source of Thymol-rich Essential Oil. *Journal of Essential*
536 *Oil Research*. 1999;11: 257–258. doi:10.1080/10412905.1999.9701126
- 537 40. Agbodan KA, Dotse K, Koumaglo KH. Activités antioxydantes des huiles essentielles de
538 trois plantes aromatiques acclimatées au Togo. *International Journal of Biological and*
539 *Chemical Sciences*. 2014;8: 1103–1110. doi:10.4314/ijbcs.v8i3.23
- 540 41. Akram A, Younis A, Akhtar G, Ameer K, Farooq A, Hanif MA, et al. Comparative
541 Efficacy of Various Essential Oil Extraction Techniques on Oil Yield and Quality of
542 *Jasminum sambac* L. *Science International*. 2017;5: 84–95.
543 doi:10.17311/sciintl.2017.84.95
- 544 42. Koba K, Sanda K, Raynaud C, Millet J, Chaumont J-P. Composition chimique des huiles
545 essentielles d’*Aeollanthus pubescens* Benth. acclimatée au Togo. *Comptes Rendus*
546 *Chimie*. 2004;7: 1107–1111. doi:10.1016/j.crci.2003.12.038
- 547 43. Dhifi W, Bellili S, Jazi S, Bahloul N, Mnif W. Essential Oils’ Chemical Characterization
548 and Investigation of Some Biological Activities: A Critical Review. *Medicines (Basel)*.
549 2016;3. doi:10.3390/medicines3040025

- 550 44. Komalamisra N, Trongtokit Y, Rongsriyam Y, Apiwathnasorn C. Screening for larvicidal
551 activity in some Thai plants against four mosquito vector species. Southeast Asian J Trop
552 Med Public Health. 2005;36: 1412–1422.
- 553 45. Sessou P, Yaovi BA, Yovo M, Gamedjo J, Dossa F, Aguidissou ON, et al. Phytochemistry
554 and antibacterial activity of plants extracts compared with two commercial antibiotics
555 against E coli responsible for avian colibacillosis in Benin. ijpm. 2018;10: 168.
556 doi:10.5138/09750185.2259
- 557 46. Tchoumboungang F, Dongmo PMJ, Lambert M, Mbanjo EGN, Fotso GBT, Henri P, et
558 al. Activité larvicide sur *Anopheles gambiae* Giles et composition chimique des huiles
559 essentielles extraites de quatre plantes cultivées au Cameroun. Biotechnol Agron Soc
560 Environ. 2009; 8.
- 561 47. Senthilkumar A, Venkatesalu V. Chemical composition and larvicidal activity of the
562 essential oil of *Plectranthus amboinicus* (Lour.) Spreng against *Anopheles stephensi*: a
563 malarial vector mosquito. Parasitol Res. 2010;107: 1275–1278. doi:10.1007/s00436-010-
564 1996-6
- 565 48. Kulkarni RR, Pawar PV, Joseph MP, Akulwad AK, Sen A, Joshi SP. *Lavandula gibsoni*
566 and *Plectranthus mollis* essential oils: chemical analysis and insect control activities
567 against *Aedes aegypti*, *Anopheles sfttephensi* and *Culex quinquefasciatus*. J Pest Sci.
568 2013;86: 713–718. doi:10.1007/s10340-013-0502-1
- 569 49. Ollengo MA, Vulule JM, Matasyoh JC. Larvicidal Activity of *Clausena Anisata* Oils and
570 Extracts *Anopheles Gambie* Larvea. International Journal of Research in Pharmacy and
571 Pharmaceutical Sciences. 2016;1: 13–20.
- 572 50. Ollengo MA. Larvicidal Activity of *Clausena Anisata* Oils and Extracts *Anopheles*
573 *Gambie* Larvea. [cited 22 Sep 2020]. Available:
574 [https://www.academia.edu/29553647/Larvicidal_Activity_of_Clausena_Anisata_Oils_a](https://www.academia.edu/29553647/Larvicidal_Activity_of_Clausena_Anisata_Oils_and_Extracts_Anopheles_Gambie_Larvea)
575 [nd_Extracts_Anopheles_Gambie_Larvea](https://www.academia.edu/29553647/Larvicidal_Activity_of_Clausena_Anisata_Oils_and_Extracts_Anopheles_Gambie_Larvea)
- 576 51. Wangrawa DW, Badolo A, Ilboudo Z, Guelbéogo WM, Kiendrébeogo M, Nèbié RCH, et
577 al. Insecticidal Activity of Local Plants Essential Oils Against Laboratory and Field
578 Strains of *Anopheles gambiae* s. l. (Diptera: Culicidae) From Burkina Faso. Journal of
579 Economic Entomology. 2018 [cited 6 Oct 2020]. doi:10.1093/jee/toy276
- 580 52. Sanei-Dehkordi A, Soleimani-Ahmadi M, Akbarzadeh K, Abadi YS, Paksa A, Gorouhi
581 MA, et al. Chemical Composition and Mosquito Larvicidal Properties of Essential Oil
582 from Leaves of an Iranian Indigenous Plant *Zhumeria majdae*. Journal of Essential Oil
583 Bearing Plants. 2016;19: 1454–1461. doi:10.1080/0972060X.2016.1222886
- 584 53. Govindarajan M, Benelli G. α -Humulene and β -elemene from *Syzygium zeylanicum*
585 (Myrtaceae) essential oil: highly effective and eco-friendly larvicides against *Anopheles*
586 *subpictus*, *Aedes albopictus*, and *Culex tritaeniorhynchus* (Diptera: Culicidae). Parasitol
587 Res. 2016;115: 2771–2778. doi:10.1007/s00436-016-5025-2
- 588 54. Kweka EJ, Lima TC, Marciale CM, de Sousa DP. Larvicidal efficacy of monoterpenes
589 against the larvae of *Anopheles gambiae*. Asian Pacific Journal of Tropical Biomedicine.
590 2016;6: 290–294. doi:10.1016/j.apjtb.2016.03.001

55. Govindarajan M, Rajeswary M, Hoti SL, Benelli G. Larvicidal potential of carvacrol and terpinen-4-ol from the essential oil of *Origanum vulgare* (Lamiaceae) against *Anopheles stephensi*, *Anopheles subpictus*, *Culex quinquefasciatus* and *Culex tritaeniorhynchus* (Diptera: Culicidae). *Research in Veterinary Science*. 2016;104: 77–82. doi:10.1016/j.rvsc.2015.11.011
56. Lima MAA, Oliveira FFM de, Gomes GA, Lavor PL, Gilv, Santiago ete MP, et al. Evaluation of larvicidal activity of the essential oils of plants species from Brazil against *Aedes aegypti* (Diptera: Culicidae). *AJB*. 2011;10: 11716–11720. doi:10.5897/AJB.9000241
57. Santos SRL, Silva VB, Melo MA, Barbosa JDF, Santos RLC, de Sousa DP, et al. Toxic Effects on and Structure-Toxicity Relationships of Phenylpropanoids, Terpenes, and Related Compounds in *Aedes aegypti* Larvae. *Vector-Borne and Zoonotic Diseases*. 2010;10: 1049–1054. doi:10.1089/vbz.2009.0158
58. Silva WJ, Dória GAA, Maia RT, Nunes RS, Carvalho GA, Blank AF, et al. Effects of essential oils on *Aedes aegypti* larvae: Alternatives to environmentally safe insecticides. *Bioresource Technology*. 2008;99: 3251–3255. doi:10.1016/j.biortech.2007.05.064
59. Evergetis E, Bellini R, Balatsos G, Michaelakis A, Carrieri M, Veronesi R, et al. From Bio-Prospecting to Field Assessment: The Case of Carvacrol Rich Essential Oil as a Potent Mosquito Larvicidal and Repellent Agent. *Front Ecol Evol*. 2018;6. doi:10.3389/fevo.2018.00204
60. Govindaraju S, Karthik C, Arulselvi PI. Evaluation of chemical composition and larvicidal activity of *Coleus aromaticus* essential oil, its major compound carvacrol against *Aedes aegypti*, *Culex quinquefasciatus* and *Anopheles stephensi* (Diptera: Culicidae). *Int J Mosq Res*. 2016;3: 06–11.
61. Cantrell CL, Pridgeon JW, Fronczek FR, Becnel JJ. Structure–Activity Relationship Studies on Derivatives of Eudesmanolides from *Inula helenium* as Toxicants against *Aedes aegypti* Larvae and Adults. *Chemistry & Biodiversity*. 2010;7: 1681–1697. doi:10.1002/cbdv.201000031
62. Gnankiné O, Bassolé IHN. Essential Oils as an Alternative to Pyrethroids’ Resistance against *Anopheles* Species Complex Giles (Diptera: Culicidae). *Molecules*. 2017;22. doi:10.3390/molecules22101321
63. Prates HT, Santos JP, Waquil JM, Fabris JD, Oliveira AB, Foster JE. Insecticidal activity of monoterpenes against *Rhyzopertha dominica* (F.) and *Tribolium castaneum* (Herbst). *Journal of Stored Products Research*. 1998;34: 243–249. doi:10.1016/S0022-474X(98)00005-8
64. Regnault-Roger C. The potential of botanical essential oils for insect pest control. *Integrated Pest Management Reviews*. 1997;2: 25–34. doi:10.1023/A:1018472227889
65. Rattan RS. Mechanism of action of insecticidal secondary metabolites of plant origin. *Crop Protection*. 2010;29: 913–920. doi:10.1016/j.cropro.2010.05.008

- 630 66. Ferguson H, Maire N, Takken W, Lyimo I, Briet O, Lindsay S, et al. Selection of mosquito
631 life-histories: A hidden weapon against malaria? Malaria journal. 2012;11: 106.
632 doi:10.1186/1475-2875-11-106
- 633 67. Smith DL, Battle KE, Hay SI, Barker CM, Scott TW, McKenzie FE. Ross, Macdonald,
634 and a Theory for the Dynamics and Control of Mosquito-Transmitted Pathogens. PLOS
635 Pathogens. 2012;8: e1002588. doi:10.1371/journal.ppat.1002588

Fig 1: Areal part of *Aeollanthus pubescens* Benth harvested in Be

Fig 2. Cone test equipment.

Fig 3. Survivorship of mosquitoes post exposure.