1 **The role of maternal effects on offspring performance** 2 **in familiar and novel environments**

3 **Authors:** Milan Vrtílek^{1*}, Pierre J. C. Chuard^{2*}, Maider Iglesias-Carrasco³,
4 Michael D. Jennions³ and Megan L. Head³

6 **Affiliations:**

7 ¹The Czech Academy of Sciences, Institute of Vertebrate Biology, Brno, Czech
8 Republic

9 ²Department of Biological Sciences, Bishop's University, Sherbrooke, Canada

10 ³Research School of Biology, The Australian National University, Canberra,
11 Australia

12 * These authors share joint first authorship on this work

14 **Correspondence:** Milan Vrtílek, The Czech Academy of Sciences, Institute of
15 Vertebrate Biology, Květná 8, 603 65 Brno, Czech Republic; phone: +420 543
16 422 557, fax: +420 543 211 346, e-mail: vrtilék@ivb.cz

18 **Conflict of interest:** We declare that we do not have any conflict of interest.

20 **Running title:** Environment-specific maternal effects

22 **Version:** November 26, 2020

23 **Figures:** 1

24 **Tables:** 4 (+1 Supplementary Table)

25 **Word count:** 6 736 words

27 **Abstract**

28 Maternal effects are an important evolutionary force that may either
 29 facilitate adaptation to a new environment or buffer against unfavourable
 30 conditions. The degree of variation in traits expressed by siblings from
 31 different mothers is frequently sensitive to environmental conditions. This
 32 could generate a Maternal-by-Environment interaction ($M \times E$) that could
 33 inflate estimates of Genotype-by-Environment effects ($G \times E$). We aimed to
 34 test for environment-specific maternal effects ($M \times E$) using a paternal full-
 35 sib/half-sib breeding design in the seed beetle *Callosobruchus maculatus*,
 36 where we split and reared offspring from the same mother on two different
 37 bean host types – original and novel. Our quantitative genetic analysis
 38 indicated that maternal effects were very small on both host types for all the
 39 measured life-history traits. There was also little evidence that maternal
 40 oviposition preference for a particular host type predicted her offspring's
 41 performance on that host. Further, additive genetic variance for most traits
 42 was relatively high on both hosts. While there was higher heritability for
 43 offspring reared in the novel host, there was no evidence for $G \times E$ s, and most
 44 cross-host genetic correlations were positive. This suggests that offspring
 45 from the same family ranked similarly for performance on both host types.
 46 Our results point to a genetic basis of host adaptation in the seed beetle,
 47 rather than maternal effects. Even so, we encourage researchers to test for

48 potential M×Es because, due to a lack of testing, it remains unclear how

49 often they arise.

50

51 **Keywords:** genetic correlation, host adaptation, maternal effects, seed

52 beetle, life history

53

54 **Introduction**

55 Maternal effects modify phenotypes that undergo selection and are
 56 therefore a potentially important evolutionary force (Mousseau and Fox,
 57 1998; Räsänen and Kruuk, 2007; Moore *et al.*, 2019). They may either
 58 facilitate adaptation to novel environments (Fox and Savalli, 2000; Leftwich
 59 *et al.*, 2019), or buffer against changing conditions through trans-
 60 generational phenotypic plasticity (Shama *et al.*, 2014). Maternal effects
 61 have also been shown to have a range of other implications, such as
 62 moderating population dynamics (Plaistow and Benton, 2009), and
 63 increasing niche breadth (Van Asch *et al.*, 2010). Maternal effects, which can
 64 arise due to both genetic and environmental variation among mothers, can
 65 be defined as ‘the causal influence of the maternal genotype or phenotype
 66 on the offspring phenotype’ (Wolf and Wade, 2009). Genetic maternal
 67 effects might increase total heritability and facilitate evolution (Wilson *et al.*,
 68 2005; Räsänen and Kruuk, 2007), as opposed to phenotypic (environmental)
 69 maternal effects. Teasing these two sources of maternal effects apart is
 70 often challenging, however (Kruuk and Hadfield, 2007). Irrespective of their
 71 genetic basis, maternal effects increase the degree of similarity between
 72 siblings and may thus bias estimates of direct genetic effects upward if
 73 ignored (Räsänen and Kruuk, 2007; McAdam *et al.*, 2014). This can lead to

74 an overestimation of the likely response to selection (McGlothlin and
75 Galloway, 2014).

76 Maternal effects are often environment sensitive (Rossiter, 1998;
77 Galloway, 2005; Räsänen and Kruuk, 2007; McAdam *et al.*, 2014). It is
78 unclear, however, whether challenging or favourable conditions will result
79 in stronger maternal effects (Charmantier and Garant, 2005; Rowiński and
80 Rogell, 2017). A harsher environment is typically assumed to increase
81 variation among offspring from different mothers (Charmantier and Garant,
82 2005; Räsänen and Kruuk, 2007). This may, for example, stem from
83 differences in maternal ability to provision offspring (Parichy and Kaplan,
84 1992). On the other hand, marine sticklebacks showed weaker maternal
85 effects under stressful conditions (warmer temperature) (Shama *et al.*,
86 2014). Similarly higher population densities reduced offspring differences
87 attributable to variation in maternal age in soil mites (Plaistow and Benton,
88 2009). Environment-dependent variation in maternal effects like this
89 represents a form of Maternal-by-Environment interaction, M×E (Figure 1B).
90 Another form of M×E occurs, irrespective of change in variance across
91 environments, if individual mothers differ in how they affect their offspring
92 in an environment-specific manner (Figure 1C). In an experiment with great
93 tits for instance (Berthouly *et al.*, 2008), mothers supplemented with
94 carotenoids raised smaller nestlings than the (un-supplemented) control

95 mothers. When the brood size was increased, however, the result was
 96 opposite and the offspring of carotenoid-supplemented mothers grew faster
 97 compared to controls (Berthouly *et al.*, 2008). A similar interaction was also
 98 observed for offspring immune response (Berthouly *et al.*, 2008). The level
 99 of maternal carotenoids thus affects the fate and relative performance of
 100 the great tit offspring depending on the brood size. This interaction would
 101 cause cross-over of the maternal effects reaction norms because of the
 102 change in rank of maternal siblings between the two environments (Figure
 103 1C). In general, a quantitative genetic analysis of potential M×E can help us
 104 to identify the mechanisms that increase similarity between siblings within
 105 the same environment beyond genetic relatedness. That analysis would be
 106 especially useful if such mechanisms themselves are undescribed, or their
 107 role is not fully understood. Many cases probably involve a mixture of
 108 changes in ranking and variances (as in Figure 1D), but the change in relative
 109 phenotype of offspring (i.e. shift in the rank) due to maternal effects is less
 110 often accounted for. In general, ignoring significant levels of M×E might bias
 111 estimates of additive genetic effects in a host-specific way and inflate G×E
 112 as shown by Vega-Trejo *et al.* (2018). This could partially explain
 113 inconsistency in G×E estimation across different studies (Saltz *et al.*, 2018).

114 Phytophagous insects are a valuable model group to study life-history
 115 evolution, especially when interested in trade-offs arising from

116 specialization to a preferred host (Via and Lande, 1985; Agrawal, 2000). At
 117 the same time, phytophagous insects exhibit a variety of maternal effects
 118 that influence key life-history traits (Fox and Dingle, 1994; Rossiter, 1996;
 119 Fox and Czesak, 2000; Van Asch *et al.*, 2010), including host-specific
 120 maternal effects (Fox *et al.*, 1997; Cahenzli and Erhardt, 2013). In the current
 121 study, we investigated the presence of maternal effects, and the potential
 122 for an interaction between maternal effects and offspring rearing host
 123 environment, on general offspring life-history traits, as well as daughters'
 124 egg laying preferences for different host types in the seed beetle,
 125 *Callosobruchus maculatus* (Coleoptera: Chrysomelidae). This beetle is a
 126 common pest of stored legumes worldwide. The larvae feed on multiple
 127 legume species (family Fabaceae) with varying degrees of success (Gompert
 128 and Messina, 2016; Price *et al.*, 2017; Messina *et al.*, 2018). Females lay eggs
 129 on host beans shortly after copulation (Mitchell, 1975). Host choice is crucial
 130 as larvae cannot move between beans. Incorrect decisions about where a
 131 female chooses to lay her eggs inevitably lead to lower quality, or even dead,
 132 offspring (Mitchell, 1975; Messina and Fry, 2003; Messina *et al.*, 2007).
 133 Larvae hatch 4-5 days after oviposition and burrow into the bean to feed on
 134 the endosperm. A single seed commonly harbours multiple larvae. The
 135 variable level of larval competition within seeds results in different sized
 136 adult beetles, which affects their condition, female reproductive

137 performance and egg size (Fox and Savalli, 1998). Depending on the
138 temperature and host species, adults typically emerge within 26-36 days of
139 oviposition (Fox *et al.*, 2003; Messina, 2004a). *Callosobruchus maculatus*
140 shows large sexual dimorphism in life-history traits with males emerging
141 earlier, being smaller and having a shorter lifespan than females (Guntrip *et*
142 *al.*, 1997; Fox, Bush, *et al.*, 2004; Iglesias-Carrasco *et al.*, 2020).

143 We used a full-sib/half-sib split brood design to tease apart the role of
144 additive genetic and maternal effects on *C. maculatus* life-history traits
145 when larvae develop on two host types – original (cowpea) and novel (mung
146 bean). We hypothesised that:

147 1) There are strong maternal effects on offspring life-history traits
148 (Fox, 1993; Messina and Fry, 2003). This led us to predict that a considerable
149 component of offspring phenotypic variation, beyond that due to additive
150 genetic effects, would be due to offspring having different mothers.

151 2) Maternal effects are environment-specific (M×E). Specifically, we
152 predicted that the maternal effects variance would differ between offspring
153 host environments (Figure 1B), and/or that the rank of offspring from the
154 same mother will change between the two host types decreasing the cross-
155 environmental maternal correlation (Figure 1C, D).

156 3) Novel host type will be more challenging for offspring. Hence,
157 offspring developing in the novel host would suffer reduced performance
158 compared to those developing in the original host type (i.e. lower larval
159 survival, longer larval development, lower body mass at emergence, and
160 shorter adult lifespan).

161 4) Maternal host preference will predict offspring performance. We
162 predicted that offspring would perform better on the host type preferred by
163 their mother when she laid her eggs.

164

165

166 **Methods**

167 **Experiment overview.** We aimed to estimate genetic and maternal effects
168 for four life-history traits and host egg laying preference. To do this we used
169 a half-sib/full-sib split brood design with offspring reared on two different
170 host types - original or novel. We created the parental generation for our
171 experiment by mating beetles from our stock population. We mated parents
172 and tested mother egg laying preference for the two host types. We then let
173 them lay eggs on both host types and measured offspring life-history traits
174 and the daughters' host preference. We used animal mixed-effects models
175 to test our predictions.

176

177 **Origin and maintenance of stock beetles.** Beetles used as the parental
178 generation for our experiment were obtained from a large stock population
179 originally sourced from the University of Western Australia (Perth, Australia)
180 in 2017 where they had been bred on cowpea (*Vigna unguiculata*, Fabaceae)
181 for at least 90 generations. We maintained this stock in our lab at the
182 Australian National University on cowpea for another 9 generations before
183 we began this experiment. Our stock was maintained in four, regularly mixed
184 (every 5 generations) populations of over 500 individuals, each kept on
185 cowpea at 25-26 °C.

186

187 **Establishing the parental generation (P).** To obtain virgin males and females
188 for the parental generation (P), we exposed approximately 2000 un-infested
189 cowpea beans to stock beetles for a period of 48 h. Each bean, which had 5-
190 8 eggs on its surface (a density which is usual for our stock), was then placed
191 in an individual Eppendorf tube (with a pinhole for airflow). Once isolated,
192 we monitored these beans until adults began to emerge. We collected virgin
193 beetles each morning and used them in our experiment on the same day.
194 We knew that beetles were virgins as they were either the only beetle to
195 have emerged that day, or all beetles that had emerged were of the same

196 sex. Every evening, we discarded any extra beetles that had emerged during
 197 that day to aid in the collection of virgins the following day. Using the beetles
 198 that emerged each morning, we mated males and females according to a
 199 full-sib/half-sib breeding design: each male was sequentially mated with
 200 four random females over the day and the mating order was noted (similar
 201 to Fox *et al.* (2003)). Pairs that did not copulate within 30 min were
 202 separated for half an hour before being placed together for another mating
 203 attempt. Each female was weighed (to the nearest 0.001 mg) prior to mating
 204 (Cubis Ultra-Micro balance, Sartorius Lab Instruments GmbH., Goettingen,
 205 Germany). All matings took place over six days ('day mated', range 1-6, see
 206 *Model comparison and partitioning of phenotypic variance*). There is
 207 therefore a positive correlation between parental developmental duration
 208 (egg to adulthood) and 'day mated', but it is imperfect because parents
 209 emerged from eggs initially laid over two days (i.e. 48 hours; see above).
 210 There is no correlation between 'day mated' and adult parental age because
 211 all matings occurred on the day that the parents emerged.

212

213 **Maternal host preference.** We used mung bean (*V. radiata*) as the novel
 214 host type. We define the host as novel in that it was 'unlikely to have been
 215 experienced by the study population within an evolutionary timescale'

216 (Rowiński and Rogell, 2017), approximately 100 generations in our case (see
217 also Kawecki (1995). Cowpea and mung are both suitable hosts for *C.*
218 *maculatus* (Messina, 2004a; Fox and Messina, 2018), but populations kept
219 on a specific host for multiple generations usually show better performance
220 on their usual host than a novel host (Messina, 2004a).

221 Once females had mated, we conducted choice trials to determine
222 their preferred host type for egg laying. We considered females ready to lay
223 eggs once the pair dismounted (Wilson and Hill, 1989). For the host
224 preference trials, we mixed cowpeas (original host) and mung beans (novel
225 host) in covered Petri dish (ø 5.5 cm). We used proportionally more mung
226 than cowpea beans (8:4) to roughly equalise the available surface area of
227 each bean type (similar to 10:5 used by Messina and Slade (1997)). Cowpea
228 beans have an approximately 1.6-times larger surface area (Paukku and
229 Kotiaho, 2008) and are 4-times heavier than mung beans (in our study the
230 average $\pm$ SD bean mass was 294 ± 45 mg for cowpea, and 72 ± 8 mg for mung
231 bean). Females were left to lay eggs for two hours, after which they were
232 removed and the number of eggs on each bean was counted. Relative
233 preference was calculated as the proportion of the eggs that a female laid
234 on the original host type (cowpea). The values for relative host preference
235 therefore ranged between 0 and 1.

236

237 **Generation of offspring (F1).** Directly after the host preference trial, we
 238 transferred individual dams to plastic containers (ø 4 cm, height 6 cm) with
 239 10-13 mung beans and left them to lay eggs for 14-18 h. We then moved
 240 dams to new containers with 10-13 cowpea beans for 9-10 h. This difference
 241 in laying time was required because initial trials showed that more time was
 242 necessary for females to lay a sufficient number of eggs on the novel host
 243 (mung beans), than on cowpea (the original host). Females were presented
 244 with mung beans first to prevent them from laying all their eggs on what we
 245 expected to be the preferred host type (cowpea) (Messina and Slade, 1997).
 246 Maternal age and/or laying order effects are unlikely to have influenced
 247 differences between mung and cowpea-reared offspring due to the short
 248 period of time we allowed females to lay eggs (i.e. <24 h). Previous studies
 249 have shown no effect of female age (hence laying order) on offspring quality
 250 during the first days of laying after emergence (Wasserman and Asami, 1985;
 251 Fox, 1993; Fox and Dingle, 1994; Iglesias-Carrasco *et al.*, 2018). Further, one
 252 of the authors (MLH) found, in a separate experiment looking at oviposition
 253 preferences and offspring traits over the same time frame, that the order in
 254 which host beans are presented does not have any observable effect on
 255 offspring survival, development duration, or body mass at emergence
 256 (Zhang *et al.*, in preparation).

257 Once females had laid eggs on both host types, we collected up to 10
258 beans of each host type for each female. If a female had laid eggs on fewer
259 than 10 beans of a given type, we used them all. We ensured that each bean
260 had only one egg laid on it by scraping off surplus eggs with a scalpel. We
261 then weighed beans, to the nearest 0.001 mg, within 24 h of oviposition to
262 measure the resources available to the larvae. Beans with an egg were then
263 placed individually in Eppendorf tubes with perforated lids and incubated at
264 26 °C.

265 We started regular monitoring of F1 emergence on day 24 post-
266 oviposition. However, we missed the onset of emergence of 40 beetles on
267 mung (1.3% of all emerged offspring) by approximately 1 day as they
268 emerged sooner than expected. Uncertainty in the date of their emergence
269 could introduce noise into the estimates of larval development duration,
270 body mass at emergence and adult lifespan, so we removed these beetles
271 from the final dataset. Including these early emerging beetles did not
272 qualitatively change our results.

273

274 **Measurement of offspring (F1) traits.** We recorded the date of offspring
275 emergence from the bean, sex and body mass at emergence (to the nearest
276 0.001 mg), and then removed the bean from the tube. After weighing, the

277 beetle was returned to its tube and checked daily for survival. We also tested
 278 the host preference of two newly emerged female offspring per dam raised
 279 on each host type (total N = 4/dam). To do this we mated each daughter with
 280 a randomly selected male from the stock and then ran an egg laying
 281 preference trial identical to that described above for their mothers. We
 282 monitored survival until the death of all emerged offspring (day 79 post-
 283 oviposition). At that time, we also censused larval (egg-to-emergence)
 284 survival, assuming that larvae in beans from which a beetle had not yet
 285 emerged had died.

286

287 **Sample sizes.** Our target sample size was determined *a priori* based on Lynch
 288 and Walsh (1998). We created 89 families (89 sires with 356 dams) and
 289 aimed for 10 offspring per dam, per host type (i.e. 7 120 offspring). The final
 290 sample size (N = 3 431) is lower due to unsuccessful matings or too few eggs
 291 being laid. Despite not reaching our target sample size our final sample size
 292 is still higher than that in other similar studies of maternal effects (the
 293 studies in a meta-analysis by Moore *et al.* (2019), for example, have a mean
 294 sample size of 1 370, and a much lower median of 555). We analysed data
 295 for sons and daughters separately due to their large sexual dimorphism
 296 which invariably leads to strong interactions between sex and other effects.

297 We excluded offspring of dams that did not lay eggs during host preference
 298 trials (285 individuals) and data from 13-24 daughters and 12-27 sons with
 299 missing, or extreme values (>3 standard deviations (SD) from the mean),
 300 with the exact number depending on the focal trait. We analysed four
 301 offspring life-history traits: (egg-to-emergence) 'larval survival' (dead or
 302 emerged; N = 3 146), 'duration of larval development' (number of days
 303 between oviposition and offspring emergence; N = 1 420 daughters and 1
 304 384 sons), 'body mass' (weight at emergence in mg; N = 1 431 daughters and
 305 1 399 sons) and 'adult lifespan' (number of days between offspring
 306 emergence and death; N = 1 425 daughters and 1 389 sons). In addition, we
 307 measured the host egg laying preferences of 665 daughters. The analysed
 308 offspring came from 82 sires and 245-249 dams for the life-history traits, and
 309 239 dams for host preference analysis. The proportion of offspring reared
 310 on the original host type was 52-57 % (see Supplementary Table S1). Prior to
 311 any statistical analysis, but after all data had been collected, we registered
 312 this project on the Open Science Forum webpage:
 313 https://osf.io/ft7eq/?view_only=0bab0a33bb4246adb64c919601a72757
 314 The final analytical approach we used was different, however, from the
 315 original plan – partly due to feedback from reviewers. We explain our
 316 changes in the *Annotated registration* section in the Supplementary
 317 Information.

318

319 **Overview of data analysis approach.** Our main objective was to test for
 320 maternal effects on the offspring traits (Prediction 1) and their potential
 321 interaction with offspring rearing environment (Prediction 2). First, we built
 322 a ‘minimal model’ containing only fixed effects directly attributable to the
 323 experimental design. To test whether maternal effects are important, we
 324 then created a set of candidate models with different random effects
 325 structures (Table 1). We used the best-selected model for partitioning of
 326 phenotypic variance (Table 2). Finally, we assess the effect of host type
 327 (Prediction 3) and dam host egg laying preference (Prediction 4) on offspring
 328 traits with a ‘full’ model containing additional fixed effects (Table 3).

329

330 **Model specification, model comparison and partitioning of phenotypic**
 331 **variance.** We started data analysis with a minimal mixed-effects model that
 332 contained, in the fixed-effects part, only variables ensuing from the
 333 experimental design. These were ‘host type’ (original/novel), ‘bean mass’
 334 (standardized to zero mean and 1 SD variance, within each host type, to look
 335 at the effect of size within each host type), ‘dam mating order’ (i.e. dam
 336 position in a sire’s mating sequence, integer 1 to 4), and ‘day mated’ (integer
 337 1-6). We did not fit ‘day mated’ as a random effect because of the low

338 number of levels (Bolker *et al.*, 2009). We specified the random effects
339 structure using an animal model approach (Kruuk, 2004; Wilson *et al.*, 2010),
340 where variance-covariance estimates between relatives are computed from
341 a pedigree.

342 We fitted the models using Bayesian framework with package
343 MCMCglmm (ver. 2.29) (Hadfield, 2010) in R software (ver. 4.0.0; R Core
344 Team, 2020). The continuous response variables (duration of larval
345 development, body mass and adult lifespan) were fitted using Gaussian
346 family and identity link, while larval survival (dead or emerged) was fitted
347 using ‘categorical’ family (binomial distribution with logit link). Daughter
348 host preference was treated as a bivariate vector with the number of eggs
349 on the original host and the total number of eggs collected in the egg laying
350 preference trial using ‘multinomial2’ family (binomial distribution and log-
351 odds ratio link function) (Hadfield, 2010).

352 To test for the best random effects structure, we performed model
353 comparison on a set of 7 candidate models (Table 1) for each trait using the
354 Deviance Information Criterion (DIC) (Spiegelhalter *et al.*, 2002). DIC is a
355 Bayesian alternative to Akaike Information Criterion (AIC) that takes into
356 account the fit of the model but penalizes for model complexity, with lower
357 DIC values indicating a better model (Hadfield, 2010; Wilson *et al.*, 2010).

358 Model comparison with DIC should be performed cautiously for non-
359 Gaussian response variables (Hadfield, 2010; Wilson *et al.*, 2010), so we did
360 not conduct model selection for larval survival or host preference.

361 The simplest model ‘G’ included only additive genetic effects specified
362 by the random effect of ‘animal’. We then also included dam identity to
363 estimate maternal effects in addition to additive genetic effects (‘G+M’
364 model). These two models do not, however, take into account possible host-
365 specific differences in respective variance components (this is shown in
366 Figure 1A). We thus formulated more realistic models that estimated the
367 variance component (including residual variance) separately for each host
368 type (as in Figure 1B). These were specified through random-effect
369 interactions (Hadfield, 2010) with host type ‘idh(HOST):animal’ for ‘G_{sep}’, or
370 with ‘idh(HOST):dam’ for ‘M_{sep}’ models (Table 1). This is similar to fitting two
371 separate host-specific models. To account for potential change in the
372 ranking of additive genetic or maternal effects (Figure 1C), we fitted models
373 that also estimated cross-environmental covariance (‘G_{cov}’ or ‘M_{cov}’). When
374 specifying variance-covariance matrix of the random-effect interaction we
375 used ‘unstructured’ form, e.g. ‘us(HOST):animal’ (Hadfield, 2010). As a
376 result, we not only obtained separate variance estimates for each host type,
377 but also the corresponding covariance to compute the cross-environmental
378 correlations (Figure 1D) – the proxies for Genotype-by-Environment (G×E),

379 or Maternal-by-Environment (M×E) interactions (Lynch and Walsh, 1998). In
380 the models estimating separate variances per host type, we always specified
381 host-specific residual variance (as 'rcov=~idh(HOST):units') (Hadfield, 2010).

382 We then performed phenotypic variance partitioning based on the
383 outputs of the best-selected minimal model for each trait. We calculated the
384 maternal-effects proportion (m^2), narrow-sense heritability (h^2), evolvability
385 (coefficient of additive genetic variation, CV_A) and the cross-environmental
386 correlation for additive genetic (r_G – potential G×E interaction) and for
387 maternal effects (r_M – potential M×E interaction). We used maternal effects

388 proportion, $m^2 = \frac{V_M}{V_P} = \frac{V_{dam}}{(V_{animal} + V_{dam} + V_{residual})}$, to quantify the

389 importance of maternal effects (as in e.g. Messina and Fry (2003); Vega-
390 Trejo *et al.* (2018); Moore *et al.* (2019)). The V_M contains both genetic and
391 non-genetic maternal effects, and potentially also genetic dominance, as we
392 did not separate them with our experimental design. We calculated

393 heritability as $h^2 = \frac{V_A}{V_P} = \frac{V_{animal}}{(V_{animal} + V_{dam} + V_{residual})}$ (Lynch and Walsh, 1998),

394 assuming epistatic effects were negligible. Evolvability was defined by
395 (Houle, 1992) as $CV_A = \frac{\sqrt{V_A}}{\mu} = \frac{\sqrt{V_{animal}}}{\mu}$, where μ is the mean of a trait.

396 Evolvability standardizes additive genetic variance over the trait mean and
397 is therefore useful to compare the potential evolutionary response among
398 traits, unlike heritability which is conditional on the amount of residual

phenotypic variation (Hansen *et al.*, 2011). The correlation between additive genetic effects on the two host types (G×E interaction) was $r_G = \frac{COV_{animal(original-novel)}}{\sqrt{V_{animal(original)}V_{animal(novel)}}}$. The same formula was used for maternal effects correlation between the two hosts (potential M×E interaction) using the dam-related covariance across the two host types. Estimates of V_A , V_P and h^2 for larval survival and host preference were back-transformed to the original data scale using the function ‘QGparams’ (from QGglmm package, ver.0.7.4; de Villemereuil *et al.* (2016)).

Testing for host effect and dam host egg laying preference on offspring phenotype. To determine the effect of novel host type on offspring performance (Prediction 3) and to test whether offspring performed better on the host type that was preferred by their mother (Prediction 4), we built a ‘full model’. The full mixed-effects model always included all the terms outlined above for the minimal model with the random effects from the best-selected minimal model (Table 1). We then included ‘dam host preference’ (i.e. the relative preference ratio: 0-1 values) and the interaction between preference and host type ‘HOST:dam host preference’ to test if the effect of the strength of the preference for the original host on offspring traits differs depending on the host type. We also included interactions

419 between host type and bean mass 'HOST:bean mass', as well as host type
420 and day mated 'HOST:day mated' as fixed effects to test for potential host-
421 specific effects.

422

423 **Model fitting.** To fit the MCMC models, we used non-informative priors for
424 the random-effects variance (G) with expected variance-covariance matrix
425 at limit 'V' = 1, and the degree of freedom 'nu' = 0.002 for each random term
426 (with nu = 1.002 in models with the random effects estimated separately per
427 host type) (Wilson *et al.*, 2010). To further improve mixing and effective
428 sample size, we specified parameter-expanded priors as ' $\alpha \mu$ ' = 0 and ' αV '
429 = 1 000 (Hadfield, 2010). Priors for residual variance (R) were set as V = 1 and
430 nu = 0.002 (nu = 1.002 when residual variance was estimated separately for
431 each host type). In the analysis of survival, the residual variance had to be
432 fixed to 1, so that the binomial mixed-effects model could be estimated
433 (Hadfield, 2010). For fixed effects (B), we retained default priors with mean
434 = 0 and variance = 10^{10} . To generate posterior distributions from the minimal
435 and full models, we ran 360 000 iterations with burn-in of 60 000 and
436 thinning interval of 300, so that we obtained an effective sample size of 1
437 000 for each estimated model term. For larval survival, we increased the
438 iterations to 5 500 000 with a burn-in of 500 000 and thinning interval of

439 5000 as we were aiming for an autocorrelation <0.1 at the lag corresponding
440 to the thinning interval (Wilson *et al.*, 2010). Model convergence was
441 evaluated by visual examination of the traces using function
442 'plot(mcmc.model)'. We assessed the importance of individual model terms
443 based on their credible intervals (CrI).

444

445

446 **Results**

447 **Prediction 1 and 2 – testing for maternal effects and M×E interaction.** The
448 model comparison suggests very weak maternal effects for all the measured
449 traits. The models incorporating maternal effects performed poorly (Table
450 1), and the best model contained only additive genetic effects estimated for
451 each host type and the cross-environmental covariance (the 'G_{cov}' model).
452 This shows that either additive genetic variance, residual variance or both
453 differed between host types and that the cross-environmental genetic
454 covariance differed from zero. Estimate of the cross-environmental
455 covariance has to be examined directly to see whether the correlation differs
456 from +1 indicating a potential Genotype-by-Environment interaction (see
457 below).

Table 1. Model comparison.

458

459 **Components of phenotypic variance related to additive genetic effects -**

460 **heritability, evolvability and G×E.** We observed higher heritability for

461 offspring life-history traits in the novel host type ($h^2 = 0.365-0.649$), than on

462 the original host ($h^2 = 0.210-0.459$) (Table 2). However, the additive genetic

463 variance was substantially higher in the novel host only for adult lifespan of

464 sons. For the other traits, we recorded a similar magnitude of additive

465 genetic variance in the two hosts, but higher residual phenotypic variance in

466 the original host type, thereby reducing trait heritability (Table 2). The

467 presence of similar levels of additive genetic variance in both hosts is also

468 apparent in the estimates of evolvability (variance due to additive genetic

469 effects standardised over trait means) which were comparable for the two

470 host types (Table 2). When comparing evolvability among traits (across sexes

471 and host types) we found that it was highest for adult lifespan ($CV_A = 16.8-$

472 20.8% , Table 2). For the non-Gaussian traits (larval survival and host egg

473 laying preference), estimates of additive genetic variance and heritability

474 obtained from the G_{cov} model were very low ($h^2 = 0.001-0.060$, Table 4A)

475 when compared to the Gaussian life-history traits.

476 We recorded highly positive correlations between offspring
477 performance on the two host types for additive genetic effects (r_G) (Table 2).
478 The only exception was for the development duration of sons, which
479 exhibited only a weak genetic correlation between hosts (Table 2). The
480 credible interval (-0.279, 0.533) suggests a Genotype-by-Environment
481 interaction ($G \times E$).

Table 2. Phenotypic variance partitioning.
--

482

483 **Predictions 3 and 4 – testing for host type suitability and host preference-**
484 **performance.** Unexpectedly, given our predictions, offspring generally
485 performed better on the novel host (mung) than on the original host
486 (cowpea) (Table 3; Supplementary Table S1). We recorded very high larval
487 survival overall, however, so that survival on mung beans was only
488 marginally better than on cowpea (Table 4B). We did not find a positive
489 relationship between the strength of the dam host preference for egg laying
490 and the performance of their offspring on either host type. The one
491 exception was for daughter body mass: daughters of dams that more
492 strongly preferred the cowpea emerged heavier when raised on cowpea, but
493 not when raised on mung beans (Table 3). Daughters reared on the original
494 cowpea host also exhibited a higher egg laying preference for cowpeas than

495 those reared on mung beans. Raw average host preference ratio of
496 daughters reared on cowpea was 0.681 versus 0.517 for those from mung
497 (where 1 is a strict preference for cowpeas). The host preference ratio of
498 their mothers (reared on cowpea) was 0.724.

Table 3. Fixed-effects estimates from the full models.

499

500 **Effect of experimental variables on offspring traits - bean size, mating date**
501 **and mating order.** Host-specific relative bean size (zero-centered per host
502 type) and mating date were important predictors of offspring traits. For both
503 host types, offspring that developed in larger beans had a poorer outcome:
504 increased development duration in daughters, as well as reduced body mass
505 and lower adult lifespan in both sexes. The effects of mating day on offspring
506 performance were, however, mixed (Table 3). Offspring of later mated
507 families took longer to develop, but sons emerged at a larger size and lived
508 longer. Daughters from later mated families also showed a modest decrease
509 in the preference for the original host type (Table 4B). Finally, if a dam was
510 later in the sire's mating sequence this marginally extended the
511 development duration of her daughters.

Table 4. Phenotypic variance partitioning and full model inference for the non-Gaussian traits.

512

513

514 **Discussion**

515 **Summary.** Our main aim was to test for environment-specific maternal
 516 effects on offspring life-history traits and their correlation, that is, Maternal-
 517 by-Environment interactions (M×E). We used a full-sib/half-sib split brood
 518 design with seed beetles (*C. maculatus*) reared on two types of host: original
 519 – cowpea; and novel – mung bean. Maternal effects, however, proved to be
 520 negligible for all of the measured traits, on both host types. Instead, we
 521 found that additive genetic variance played an important role in determining
 522 phenotypic variation. Contrary to our predictions, offspring reared on the
 523 novel host type performed better for all the measured traits. Daughters
 524 whose mothers preferred to lay eggs on the original host were heavier on
 525 this host than those of mothers that preferred the novel host type. There
 526 was, however, no equivalent effect of host preference on body mass in the
 527 novel host. Below we discuss our findings in more detail.

528

529 **Weak maternal effects.** We used an original and a novel host type to
 530 examine whether mothers affect their offspring's phenotype in a host-

531 specific way. Previous studies have shown that environmental stress can
 532 either increase or decrease the magnitude of maternal effects (Charmantier
 533 and Garant, 2005; Rowiński and Rogell, 2017). Maternal effects variation is
 534 sometimes not evident in resource rich conditions as differences among
 535 mothers are reduced compared to a poor environment where differences
 536 are exacerbated (e.g. Parichy and Kaplan (1992)). If both host types in our
 537 study are resource rich (especially when few eggs are laid per bean) this
 538 could explain the negligible maternal effects that we see here. Previous
 539 quantitative genetic studies employing a similar approach to ours have
 540 reported significant maternal effects in *C. maculatus*. This has been,
 541 however, in other hosts (adzuki (*Vigna angularis*) in Fox (1993)) or when
 542 testing the effect of bean removal after beetle emergence on adult traits
 543 (Messina and Fry, 2003). The host species we used are both thought to be
 544 highly favourable to *C. maculatus* (Messina, 2004a; Fox and Messina, 2018)
 545 (discussed further below). This, in addition to the fact we prevented larval
 546 competition with our experimental design, may have substantially reduced
 547 the role of maternal effects in the current experiment.

548

549 **Heritability was higher in the novel host type, but evolutionary potential**
 550 **was similar in both hosts.** Heritability of fitness-related traits is traditionally
 551 assumed to be low, as additive genetic variation is expected to be depleted

552 by natural selection (Mousseau and Roff, 1987; Hill, 2010). Egg size, body
 553 size, fecundity or lifespan were nevertheless previously shown to have
 554 intermediate to high heritability in *C. maculatus* (h^2 from 0.27 to 0.74) (Fox,
 555 1993; Messina, 1993; Fox *et al.*, 2003; Fox and Messina, 2018). Here, we
 556 recorded considerable heritability in all life-history traits (except larval
 557 survival), with the highest values occurring for daughter body mass and son
 558 adult lifespan (both $h^2 > 0.65$). Heritability was notably higher in the novel
 559 host environment (mung bean), with the one exception of adult lifespan of
 560 daughters. Holloway *et al.* (1990) proposed that a novel environment should
 561 yield higher additive genetic variance due to new genes being expressed
 562 thereby uncovering cryptic genetic variation (Hoffmann and Merilä, 1999).
 563 Indeed, Messina and Fry (2003) found that *C. maculatus* exhibited increased
 564 additive genetic variance for longevity under a novel environment (absence
 565 of host beans after emergence). Our results indicate, however, that it was
 566 lower residual variance in the novel host (mung bean), rather than increased
 567 additive genetic variation, that was responsible for higher heritability in the
 568 novel environment. Residual phenotypic variation may entail non-additive
 569 genetic, environmental variance and developmental noise (Rowiński and
 570 Rogell, 2017) and the influence of at least some of these factors apparently
 571 declined in the novel host type. More elaborated breeding design would be
 572 required (see e.g. Tucić and Šešlija (2007), or Bilde *et al.* (2008)), however,

573 to examine their respective effects in *C. maculatus*. Interpretation of
574 heritability as the proportion of additive genetic versus total variance can
575 therefore become problematic (Hansen *et al.*, 2011) and evolvability offers
576 a more conclusive comparison of evolutionary potential (Houle, 1992;
577 Hansen *et al.*, 2011). Similar values of trait evolvability indicated that the
578 traits measured in our study possess similar evolutionary potential in the
579 two host types.

580 Genetic trade-offs are thought to prevent populations of parasites or
581 herbivores from becoming ‘masters of all trades’ and performing well on
582 multiple hosts (Joshi and Thompson, 1995; Agrawal *et al.*, 2010). In
583 phytophagous insects, performance trade-offs on different hosts are,
584 however, rarely reported (Ueno *et al.*, 2003; Scheirs *et al.*, 2005; Messina
585 and Durham, 2015). This includes studies of *C. maculatus* where offspring
586 from the same family usually perform similarly across hosts without
587 exhibiting a G×E interaction (Fox, 1993; Guntrip and Sibly, 1998; Messina,
588 2004a). We also found strong positive cross-environmental genetic
589 correlations indicating parallel reaction norms for offspring from different
590 families when reared on the two host types. Only the development duration
591 of sons showed a low genetic correlation between the two host types
592 indicating that it could evolve differently on cowpea and mung (Via and
593 Lande, 1985). It is, however, important to bear in mind that breeding design

594 studies cannot definitively conclude there are no genetic trade-offs in
595 adaptation to different hosts because they lack mechanistic explanation.
596 This can be obtained through analysis of long-term selected lines (Agrawal
597 *et al.*, 2010) or data on the potential genes involved (Gompert and Messina,
598 2016).

599

600 **Offspring performed better on the novel host type.** We treated mung bean
601 as a novel host type. This was despite mung bean regularly being used to
602 rear other strains of *C. maculatus* (e.g. the Southern India strain (Fox, Bush,
603 *et al.*, 2004)), and regarded with cowpea as a favourable host species for *C.*
604 *maculatus* (Fox and Messina, 2018). Our stock population was kept on
605 cowpea for at least 100 generations. In addition, cowpea strains are overall
606 more viable than those from mung bean (Messina, 2004a) and the cowpea
607 is generally considered to be the ancestral host for *C. maculatus* (Kébé *et al.*,
608 2017). Due to these reasons and the fact that a previous study showed that
609 long-term maintenance on a specific host yields decreased viability on other
610 hosts (Messina, 2004a), we predicted a lower suitability of the novel host
611 type (mung bean). Our finding that offspring performed better on the novel
612 host type (mung bean) was therefore unexpected. While offspring survival
613 was almost 90 % on cowpea, it was even higher on the novel host (93 % on
614 mung bean). There was no obvious cost to the offspring of rearing in the

615 novel host, as the duration of larval development was shorter on mung bean.
 616 However, beetles that emerged from mung beans were larger and lived
 617 longer as adults than those that emerged from cowpeas. It is not
 618 straightforward, however, to tell which of the two host species is generally
 619 better for *C. maculatus*, because it depends on other factors such as the level
 620 of competition (Messina, 2004a; Pauku and Kotiaho, 2008; Fox and
 621 Messina, 2018), temperature (Stillwell *et al.*, 2007) and the population's
 622 evolutionary history (Messina, 2004a). In the present study, we eliminated
 623 effects of larval competition by rearing one larva per bean. We therefore
 624 removed the fitness advantage that cowpea may provide due to its larger
 625 size when there is larval competition (Fox and Savalli, 1998). Surprisingly,
 626 relative bean size within a given host type also negatively affected body mass
 627 and lifespan as well as extending the developmental duration of daughters.

628 We cannot reject the hypothesis that the host effect recorded in our
 629 experiment was confounded by egg laying order, as eggs were laid first on
 630 mung bean (see Methods for the experimental justification). The lower
 631 performance of offspring on cowpea might have been because, following
 632 the preference trial, females were first provided with mung beans to lay
 633 eggs, and only after that with cowpeas. There are, however, several reasons
 634 to think that an effect of laying order in our experiment is unlikely. The
 635 lifetime fecundity of *C. maculatus* is 60-80 eggs (Mitchell, 1975; Credland

636 and Wright, 1989; Wilson and Hill, 1989; Messina and Fry, 2003). The dams
637 in our experiment laid around 30 eggs per 24 h and probably still possessed
638 more than half of their lifetime fecundity after they had finishing laying in
639 our experiment. Crucially, past studies suggest there is no negative effect of
640 maternal age on offspring performance on the first day of egg laying (i.e. the
641 relevant time period in our study) (Wasserman and Asami, 1985; Fox, 1993;
642 Fox and Dingle, 1994; Iglesias-Carrasco *et al.*, 2018; Zhang *et al.*, in
643 preparation). We therefore think that the sequence of host types available
644 for egg laying is unlikely to explain the poorer offspring performance on
645 cowpea.

646

647 **Host preference of *C. maculatus* increased daughter body mass and rearing**
648 **host affected host preference.** An egg laying preference for a particular host
649 often predicts offspring performance on that host in phytophagous insects
650 (Gripenberg *et al.*, 2010). However, despite the critical role of oviposition
651 choice for larval success in *C. maculatus*, of the five offspring traits we
652 measured, the strength of dam preference for cowpea only correlated with
653 daughter body mass. Dams that showed a stronger preference for cowpea
654 produced heavier-than-average daughters on cowpea, but not when laying
655 on mung. This contributes to mother fitness through higher daughter
656 reproductive potential in cowpea as body size is a good proximate measure

657 for fecundity in *C. maculatus* (Messina and Fry, 2003). In mung bean,
658 however, daughters' body mass was not affected by their mothers' host
659 preference and the preference-performance relationship thus only holds for
660 daughter body mass in cowpea. The weak overall relationship between
661 mother's host preference for one host type and the performance of her
662 offspring on that host suggests that *C. maculatus* females might aim to
663 maximize their own fitness and not that of individual offspring (i.e. they
664 prioritise offspring number over offspring quality), perhaps through saving
665 searching time for the best host (Wood *et al.*, 2018).

666 Previous studies show that strains of *C. maculatus* maintained for > 40
667 generations on mung bean or cowpea exhibit a strong preference for their
668 recent host type (Messina, 2004b). Fox, Stillwell, *et al.* (2004) used different
669 host strains to demonstrate that switching a strain to another host for just a
670 single generation has no effect on host preference. We found that while
671 daughters reared on cowpea preferred cowpea over the novel host (mung
672 bean), those reared on mung beans showed no preference for either host
673 type. This suggests that there might be a G×E interaction affecting host egg
674 laying preferences, but tracking of more generations is necessary to test
675 whether the host preference evolves after the host switch (as in Fox,
676 Stillwell, *et al.* (2004), or Messina (2004b)).

677

678 **Conclusion.** Maternal effects were low for all measured traits in our study
 679 for offspring reared on either a novel or a familiar host. A next step in the
 680 exploration of whether M×E interactions affect estimates of genetic
 681 variation, and of G×E interactions specifically, would be to conduct
 682 experiments using less favourable novel hosts of *C. maculatus*, for example
 683 adzuki bean (Fox, 1993) or lentil (Gompert and Messina, 2016). The host
 684 species we used are both considered to be of high quality (Fox and Messina,
 685 2018). By creating more challenging conditions for offspring, maternal
 686 effects might emerge as an important host-specific factor that affects
 687 variation in offspring performance (Fox *et al.*, 1997; Räsänen and Kruuk,
 688 2007). Despite the fact that maternal effects were negligible in our study,
 689 we still argue that it is important testing for M×E interactions. The M×E
 690 effects have potential to bias measurements of G×E and the lack of research
 691 on M×E interaction is an obstacle towards understanding their role in
 692 adaptive processes.

693

694

695 **Acknowledgements**

696 We appreciate detailed comments by three anonymous referees that helped
 697 us to improve the manuscript. We thank Timothée Bonnet and Loeske E. B.

698 Kruuk (RSB ANU) for very helpful discussions about phenotypic variation
 699 partitioning and Bayesian mixed-effect models. We are thankful to Carolina
 700 Dolan, Krish S. Sanghvi and Elroy Kwan Au (RSB ANU) for their help with the
 701 experiment. MV and PJCC were both supported by 6-month Endeavour
 702 fellowships from the Australian Government.

703

704 **Author contribution**

705 MV, PJCC, MLH and MDJ conceived and designed the study; MV, PJCC and
 706 MIC collected data; MV analysed the data and drafted the manuscript; and
 707 PJCC, MIC, MLH and MDJ revised it. All authors approved the final version of
 708 the manuscript.

709

710 **Conflict of interest**

711 We declare that we do not have any conflict of interest.

712

713 **Data archiving**

714 Data with accompanying R code will be made available after manuscript
 715 acceptance.

716

717 **Supplementary Information**

718 Table S1 and Annotated Registration.

719

720 **References**

721 Agrawal AA (2000). Host-range evolution: Adaptation and trade-offs in
722 fitness of mites on alternative hosts. *Ecology* **81**: 500–508.

723 Agrawal AA, Conner JK, Rasmann S (2010). Tradeoffs and negative
724 correlations in evolutionary ecology. In: Futuyma D, Levinton J, Eanes
725 W (eds) *Evolution since Darwin: the first 150 years*, Sinauer, pp. 243–
726 268.

727 Van Asch M, Julkunen-Tiito R, Visser ME (2010). Maternal effects in an
728 insect herbivore as a mechanism to adapt to host plant phenology.
729 *Funct Ecol* **24**: 1103–1109.

730 Berthouly A, Cassier A, Richner H (2008). Carotenoid-induced maternal
731 effects interact with ectoparasite burden and brood size to shape the
732 trade-off between growth and immunity in nestling great tits. *Funct*
733 *Ecol* **22**: 854–863.

734 Bilde T, Friberg U, Maklakov AA, Fry JD, Arnqvist G (2008). The genetic
735 architecture of fitness in a seed beetle: Assessing the potential for
736 indirect genetic benefits of female choice. *BMC Evol Biol* **8**.

- 737 Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, et
738 *al.* (2009). Generalized linear mixed models: a practical guide for
739 ecology and evolution. *Trends Ecol Evol* **24**: 127–35.
- 740 Cahenzli F, Erhardt A (2013). Transgenerational acclimatization in an
741 herbivore-host plant relationship. *Proc R Soc B Biol Sci* **280**.
- 742 Charmantier A, Garant D (2005). Environmental quality and evolutionary
743 potential: Lessons from wild populations. *Proc R Soc B Biol Sci* **272**:
744 1415–1425.
- 745 Credland PF, Wright AW (1989). Factors affecting female fecundity in the
746 cowpea seed beetle, *Callosobruchus maculatus* (Coleoptera:
747 Bruchidae). *J Stored Prod Res* **25**: 125–136.
- 748 Fox CW (1993). Maternal and genetic influences on egg size and larval
749 performance in a seed beetle. *Heredity* **73**: 509–517.
- 750 Fox CW, Bush ML, Roff DA, Wallin WG (2004). Evolutionary genetics of
751 lifespan and mortality rates in two populations of the seed beetle,
752 *Callosobruchus maculatus*. *Heredity* **92**: 170–181.
- 753 Fox CW, Bush ML, Wallin WG (2003). Maternal age affects offspring
754 lifespan of the seed beetle. *Funct Ecol* **17**: 811–820.
- 755 Fox CW, Czesak ME (2000). Evolutionary ecology of progeny size in
756 arthropods. *Annu Rev Entomol* **45**: 341–369.

- 757 Fox CW, Dingle H (1994). Dietary mediation of maternal age effects on
758 offspring performance in a seed beetle (Coleoptera: Bruchidae). *Funct*
759 *Ecol* **8**: 600–606.
- 760 Fox CW, Messina FJ (2018). Evolution of larval competitiveness and
761 associated life-history traits in response to host shifts in a seed beetle.
762 *J Evol Biol* **31**: 302–313.
- 763 Fox CW, Savalli UM (1998). Inheritance of environmental variation in body
764 size: Superparasitism of seeds affects progeny and grandprogeny body
765 size via a nongenetic maternal effect. *Evolution* **52**: 172.
- 766 Fox CW, Savalli UM (2000). Maternal effects mediate host expansion in a
767 seed-feeding beetle. *Ecology* **81**: 3–7.
- 768 Fox CW, Stillwell RC, Amarillo-Suárez AR, Czesak ME, Messina FJ (2004).
769 Genetic architecture of population differences in oviposition
770 behaviour of the seed beetle *Callosobruchus maculatus*. *J Evol Biol* **17**:
771 1141–1151.
- 772 Fox CW, Thakar MS, Mousseau TA (1997). Egg size plasticity in a seed
773 beetle: An adaptive maternal effect. *Am Nat* **149**: 149–163.
- 774 Galloway LF (2005). Maternal effects provide phenotypic adaptation to
775 local environmental conditions. *New Phytol* **166**: 93–100.
- 776 Gompert Z, Messina FJ (2016). Genomic evidence that resource-based

- 777 trade-offs limit host-range expansion in a seed beetle. *Evolution* **70**:
778 1249–1264.
- 779 Gripenberg S, Mayhew PJ, Parnell M, Roslin T (2010). A meta-analysis of
780 preference-performance relationships in phytophagous insects. *Ecol*
781 *Lett* **13**: 383–393.
- 782 Guntrip J, Sibly RM (1998). Phenotypic plasticity, genotype-by-environment
783 interaction and the analysis of generalism and specialization in
784 *Callosobruchus maculatus*. *Heredity* **81**: 198–204.
- 785 Guntrip J, Sibly RM, Holloway GJ (1997). The effect of novel environment
786 and sex on the additive genetic variation and covariation in and
787 between emergence body weight and development period in the
788 cowpea weevil, *Callosobruchus maculatus* (Coleoptera, Bruchidae).
789 *Heredity* **78**: 158–165.
- 790 Hadfield JD (2010). MCMC methods for multi-response generalized linear
791 mixed models: The MCMCglmm R package. *J Stat Softw* **33**.
- 792 Hansen TF, Pélabon C, Houle D (2011). Heritability is not evolvability. *Evol*
793 *Biol* **38**: 258–277.
- 794 Hill WG (2010). Understanding and using quantitative genetic variation.
795 *Philos Trans R Soc B Biol Sci* **365**: 73–85.
- 796 Hoffmann AA, Merilä J (1999). Heritable variation and evolution under

- 797 favourable and unfavourable conditions. *Trends Ecol Evol* **14**: 96–101.
- 798 Holloway GJ, Povey SR, Sibly RM (1990). The effect of new environment on
799 adapted genetic architecture. *Heredity* **64**: 323–330.
- 800 Houle D (1992). Comparing evolvability and variability of quantitative traits.
801 *Genetics* **130**: 195–204.
- 802 Iglesias-Carrasco M, Brookes S, Kruuk LEB, Head ML (2020). The effects of
803 competition on fitness depend on the sex of both competitors. *Ecol*
804 *Evol* **10**: 9808–9826.
- 805 Iglesias-Carrasco M, Jennions MD, Zajitschek SRK, Head ML (2018). Are
806 females in good condition better able to cope with costly males?
807 *Behav Ecol* **29**: 876–884.
- 808 Joshi A, Thompson JN (1995). Trade-offs and the evolution of host
809 specialization. *Evol Ecol* **9**: 82–92.
- 810 Kawecki TJ (1995). Expression of genetic and environmental variation for
811 life history characters on the usual and novel hosts in *Callosobruchus*
812 *maculatus* (Coleoptera: Bruchidae). *Heredity* **75**: 70–76.
- 813 Kébé K, Alvarez N, Tuda M, Arnqvist G, Fox CW, Sembène M, *et al.* (2017).
814 Global phylogeography of the insect pest *Callosobruchus maculatus*
815 (Coleoptera: Bruchinae) relates to the history of its main host, *Vigna*
816 *unguiculata*. *J Biogeogr* **44**: 2515–2526.

- 817 Kruuk LEB (2004). Estimating genetic parameters in natural populations
818 using the ‘animal model’. *Philos Trans R Soc B Biol Sci* **359**: 873–890.
- 819 Kruuk LEB, Hadfield JD (2007). How to separate genetic and environmental
820 causes of similarity between relatives. *J Evol Biol* **20**: 1890–1903.
- 821 Leftwich PT, Nash WJ, Friend LA, Chapman T (2019). Contribution of
822 maternal effects to dietary selection in Mediterranean fruit flies.
823 *Evolution* **73**: 278–292.
- 824 Lynch M, Walsh B (1998). *Genetics and Analysis of Quantitative Traits*.
825 Sinauer.
- 826 McAdam AG, Garant D, Wilson AJ (2014) The effects of others' genes:
827 maternal and other indirect genetic effects. In: Charmantier A, Garant
828 D, Kruuk LEB (eds) *Quantitative Genetics in the Wild*. Oxford University
829 Press, pp. 81-104.
- 830 McGlothlin JW, Galloway LF (2014). The contribution of maternal effects to
831 selection response: An empirical test of competing models. *Evolution*
832 **68**: 549–558.
- 833 Messina FJ (1993). Heritability and ‘evolvability’ of fitness components in
834 *Callosobruchus maculatus*. *Heredity* (**71**): 623–629.
- 835 Messina FJ (2004a). Predictable modification of body size and competitive
836 ability. *Evolution* **58**: 2788–2797.

- 837 Messina FJ (2004b). How labile are the egg-laying preferences of seed
838 beetles? *Ecol Entomol* **29**: 318–326.
- 839 Messina FJ, Durham SL (2015). Loss of adaptation following reversion
840 suggests trade-offs in host use by a seed beetle. *J Evol Biol* **28**: 1882–
841 1891.
- 842 Messina FJ, Fry JD (2003). Environment-dependent reversal of a life history
843 trade-off in the seed beetle *Callosobruchus maculatus*. *J Evol Biol* **16**:
844 501–509.
- 845 Messina FJ, Lish AM, Gompert Z (2018). Variable responses to novel hosts
846 by populations of the seed beetle *Callosobruchus maculatus*
847 (Coleoptera: Chrysomelidae: Bruchinae). *Environ Entomol* **47**: 1194–
848 1202.
- 849 Messina FJ, Morrey JL, Mendenhall M (2007). Why do host-deprived seed
850 beetles ‘dump’ their eggs? *Physiol Entomol* **32**: 259–267.
- 851 Messina FJ, Slade AF (1997). Inheritance of host-plant choice in the seed
852 beetle *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Ann*
853 *Entomol Soc Am* **90**: 848–855.
- 854 Mitchell R (1975). The evolution of oviposition tactics in the bean weevil,
855 *Callosobruchus maculatus* (F.). *Ecology* **56**: 696–702.
- 856 Moore MP, Whiteman HH, Martin RA (2019). A mother’s legacy: the

857 strength of maternal effects in animal populations. *Ecol Lett* **22**: 1620–
858 1628.

859 Mousseau TA, Fox CW (1998). The adaptive significance of maternal
860 effects. *Trends Ecol Evol* **13**: 403–407.

861 Mousseau TA, Roff DA (1987). Natural selection and the heritability of
862 fitness components. *Heredity* **59**: 181–197.

863 Parichy DM, Kaplan RH (1992). Maternal effects on offspring growth and
864 development depend on environmental quality in the frog *Bombina*
865 *orientalis*. *Oecologia* **91**: 579–586.

866 Paukku S, Kotiaho JS (2008). Female oviposition decisions and their impact
867 on progeny life-history traits. *J Insect Behav* **21**: 505–520.

868 Plaistow SJ, Benton TG (2009). The influence of context-dependent
869 maternal effects on population dynamics: An experimental test. *Philos*
870 *Trans R Soc B Biol Sci* **364**: 1049–1058.

871 Price TN, Leonard A, Lancaster LT (2017). Warp-speed adaptation to novel
872 hosts after 300 generations of enforced dietary specialisation in the
873 seed beetle *Callosobruchus maculatus* (Coleoptera: Chrysomelidae:
874 Bruchinae). *Eur J Entomol* **114**: 257–266.

875 R Core Team (2020) R: A language and environment for statistical
876 computing. v. 4.0.0. R Foundation for Statistical Computing, Vienna,

- 877 Austria.
- 878 Räsänen K, Kruuk LEB (2007). Maternal effects and evolution at ecological
879 time-scales. *Funct Ecol* **21**: 408–421.
- 880 Roff DA, Wilson AJ (2014) Quantifying Genotype-by-Environment
881 interactions in laboratory systems. In: Hunt J, Hosken D (eds)
882 *Genotype-by-Environment Interactions and Sexual Selection*. John
883 Wiley and Sons, Ltd., pp. 101-136.
- 884 Rossiter M (1996). Incidence and consequences of inherited environmental
885 effects. *Annu Rev Ecol Syst* **27**: 451–476.
- 886 Rossiter M (1998) The role of environmental variation in parental effects
887 expression. In: Mousseau TA, Fox CW (eds) *Maternal Effects as*
888 *Addaptations*. Oxford University Press, pp. 112-134.
- 889 Rowiński PK, Rogell B (2017). Environmental stress correlates with
890 increases in both genetic and residual variances: A meta-analysis of
891 animal studies. *Evolution* **71**: 1339–1351.
- 892 Saltz JB, Bell AM, Flint J, Gomulkiewicz R, Hughes KA, Keagy J (2018). Why
893 does the magnitude of genotype-by-environment interaction vary?
894 *Ecol Evol* **8**: 6342–6353.
- 895 Scheirs J, Jordaens K, De Bruyn L (2005). Have genetic trade-offs in host use
896 been overlooked in arthropods? *Evol Ecol* **19**: 551–561.

- 897 Shama LNS, Strobel A, Mark FC, Wegner KM (2014). Transgenerational
898 plasticity in marine sticklebacks: Maternal effects mediate impacts of
899 a warming ocean. *Funct Ecol* **28**: 1482–1493.
- 900 Spiegelhalter DJ, Best NG, Carlin BP, Van Der Linde A (2002). Bayesian
901 measures of model complexity and fit. *J R Stat Soc Ser B Stat Methodol*
902 **64**: 583–616.
- 903 Stillwell RC, Wallin WG, Hitchcock LJ, Fox CW (2007). Phenotypic plasticity
904 in a complex world: Interactive effects of food and temperature on
905 fitness components of a seed beetle. *Oecologia* **153**: 309–321.
- 906 Tucić N, Šešlija D (2007). Genetic architecture of differences in oviposition
907 preference between ancestral and derived populations of the seed
908 beetle *Acanthoscelides obtectus*. *Heredity* **98**: 268–273.
- 909 Ueno H, Fujiyama N, Yao I, Sato Y, Katakura H (2003). Genetic architecture
910 for normal and novel host-plant use in two local populations of the
911 herbivorous ladybird beetle, *Epilachna pustulosa*. *J Evol Biol* **16**: 883–
912 895.
- 913 Vega-Trejo R, Head ML, Jennions MD, Kruuk LEB (2018). Maternal-by-
914 environment but not genotype-by-environment interactions in a fish
915 without parental care. *Heredity* **120**: 154–167.
- 916 Via S, Lande R (1985). Genotype-environment interaction and the evolution

917 of phenotypic plasticity. *Evolution* **39**: 505–522.

918 de Villemereuil P, Schielzeth H, Nakagawa S, Morrissey MB (2016). General
919 methods for evolutionary quantitative genetic inference from
920 generalized mixed models. *Genetics* **204**: 1281–1294.

921 Wasserman SS, Asami T (1985). The effect of maternal age upon fitness of
922 progeny in the southern cowpea weevil, *Callosobruchus maculatus*.
923 *Oikos* **45**: 191–196.

924 Wilson AJ, Coltman DW, Pemberton JM, Overall ADJ, Byrne KA, Kruuk LEB
925 (2005). Maternal genetic effects set the potential for evolution in a
926 free-living vertebrate population. *J Evol Biol* **18**: 405–414.

927 Wilson K, Hill L (1989). Factors affecting egg maturation in the bean weevil
928 *Callosobruchus maculatus*. *Physiol Entomol* **14**: 115–126.

929 Wilson AJ, Réale D, Clements MN, Morrissey MB, Postma E, Walling CA, et
930 al. (2010). An ecologist's guide to the animal model. *J Anim Ecol* **79**:
931 13–26.

932 Wolf JB, Wade MJ (2009). What are maternal effects (and what are they
933 not)? *Philos Trans R Soc London B* **364**: 1107–1115.

934 Wood CW, Wice EW, del Sol J, Paul S, Sanderson BJ, Brodie ED (2018).
935 Constraints imposed by a natural landscape override offspring fitness
936 effects to shape oviposition decisions in wild forked fungus beetles.

937 *Am Nat* **191**: 524–538.

938 **TABLES**

939 **Table 1.** Model comparison to estimate the best random effects structure for minimal animal MCMC models. The table shows the difference
940 between Deviance Information Criterion (DIC) for each candidate model and the model with the lowest DIC (denoted as '-'). Model 'G' contained
941 additive genetic variance only. The model with 'M' included also maternal effects variance. Models where variance was estimated separately
942 'sep' for each host type are 'G_{sep}' or 'M_{sep}'. Models 'G_{cov}' (or 'M_{cov}') estimated covariance 'cov' between the two host types in addition to the host-
943 specific variance. When the host-specific interaction was fitted as a random effect (models 'sep' and 'cov'), we also estimate residual variance
944 separately for each host type. Further details on model fitting are in the Methods. Please note that the low DIC for model 'G_{cov}' means that the
945 genetic covariance between the host types is different from zero. It is, however, inconclusive about Genotype-by-Environment interaction as
946 the covariance can still be highly positive (i.e. the rank order of genotypes does not change). So we also examined the cross-environmental
947 genetic correlation (r_G) estimates from the model given in Table 2. DIC is not reliable for model comparison in non-Gaussian traits (Hadfield,
948 2010; Wilson *et al.*, 2010) and we thus did not perform model selection for larval survival and host egg laying preference of daughters and used
949 model G_{cov} for inference.

950

951

Model	Random effects	Development duration		Body mass		Adult lifespan	
		daughters	sons	daughters	sons	daughters	sons
G	animal	137.6	225.4	82	78.8	70.4	125.6
G+M	animal + dam	151.8	233.3	136.3	115.9	79.4	153.2
G _{sep}	idh(HOST):animal	10.4	-	18.1	46.3	15.6	72.8
G _{cov}	us(HOST):animal	-	0.4	-	-	-	-
G _{sep} +M _{sep}	idh(HOST):animal + idh(HOST):dam	42.5	29.6	97.5	108.9	39.2	133.1
G _{cov} +M _{sep}	us(HOST):animal + idh(HOST):dam	23.7	33.5	31.3	22.1	16.3	17.6
G _{cov} +M _{cov}	us(HOST):animal + us(HOST):dam	30.1	31.3	53.3	63.6	17.2	34.6

952

953

954 **Table 2.** Phenotypic variance partitioning based on the best model from the model comparison. We used model ‘G_{cov}’ for all the traits (Table 1)
955 as it estimates cross-environmental genetic covariance and models with DIC difference <2 are usually considered equally supported. We
956 estimated host-specific additive genetic effects (V_A), proportion of additive genetic variance in total phenotypic variance (V_P) - heritability (h²),
957 additive genetic variance standardized over trait mean - evolvability (CV_A), and cross-environmental genetic correlation (r_G). The estimates and
958 their credible intervals (in brackets) come from posterior distribution of sex-specific animal MCMC models.

Host type	Par.	Development duration		Body mass		Adult lifespan	
		daughters (N = 1 420)	sons (N = 1 384)	daughters (N = 1 431)	sons (N = 1 399)	daughters (N = 1 425)	sons (N = 1 389)
original	V _A	0.430 (0.105,0.678)	0.448 (0.208,0.876)	0.215 (0.147,0.317)	0.070 (0.034,0.114)	19.981 (11.423,26.851)	9.694 (5.578,14.841)
	V _P	1.768 (1.578,1.948)	2.167 (1.979,2.409)	0.501 (0.446,0.555)	0.310 (0.282,0.343)	39.267 (35.595,45.099)	28.298 (24.908,31.138)
	h ²	0.229 (0.084,0.387)	0.210 (0.082,0.374)	0.447 (0.327,0.615)	0.225 (0.122,0.358)	0.459 (0.301,0.624)	0.333 (0.218,0.505)
	CV _A	0.020 (0.012,0.027)	0.021 (0.014,0.030)	0.084 (0.066,0.097)	0.075 (0.058,0.100)	0.195 (0.148,0.226)	0.184 (0.146,0.234)
novel	V _A	0.349 (0.213,0.622)	0.527 (0.355,0.892)	0.250 (0.176,0.319)	0.103 (0.068,0.148)	16.664 (10.973,27.715)	16.304 (11.261,24.817)
	V _P	0.956 (0.853,1.083)	1.154 (1.028,1.339)	0.374 (0.340,0.426)	0.211 (0.186,0.236)	37.645 (33.595,43.241)	26.675 (23.763,31.265)
	h ²	0.365 (0.241,0.606)	0.495 (0.348,0.708)	0.649 (0.501,0.786)	0.513 (0.355,0.669)	0.507 (0.334,0.684)	0.627 (0.480,0.854)
	CV _A	0.020 (0.016,0.026)	0.026 (0.020,0.031)	0.077 (0.065,0.087)	0.082 (0.067,0.099)	0.168 (0.131,0.204)	0.208 (0.176,0.260)
	r _G	0.706 (0.277,0.981)	0.147 (-0.279,0.533)	0.901 (0.670,0.995)	0.996 (0.750,1.000)	0.790 (0.615,0.982)	0.967 (0.777,0.999)

959

960 **Table 3.** The effect of host type, other experimental variables and dam host preference on three offspring life-history traits (fixed-effects
961 estimates from the full animal models). The intercept always corresponds to the trait average on the original host type and the ‘novel HOST
962 effect’ is the effect of the novel host type (i.e. difference between the two means). The same applies to interaction terms, where the
963 interaction gives the estimate for the difference between the effect on the novel and the original host type. We embolden any credible
964 intervals (in brackets) that do not overlap zero. The estimates and their credible intervals come from posterior distribution of sex-specific
965 animal MCMC models.

Term	Development duration [days]		Body mass [mg]		Adult lifespan [days]	
	daughters (N = 1 420)	sons (N = 1 384)	daughters (N = 1 431)	sons (N = 1 399)	daughters (N = 1 425)	sons (N = 1 389)
Intercept (original HOST)	31.899 (31.713,32.103)	31.442 (31.187,31.607)	5.784 (5.660,5.904)	3.466 (3.414,3.593)	23.62 (22.432,24.739)	16.298 (15.338,17.259)
novel HOST effect	-1.285 (-1.388,-1.115)	-1.148 (-1.296,-0.935)	0.728 (0.650,0.792)	0.349 (0.303,0.408)	3.338 (2.595,3.975)	2.355 (1.792,2.885)
bean mass (original)	0.145 (0.035,0.229)	0.105 (-0.017,0.189)	-0.083 (-0.130,-0.032)	-0.039 (-0.089,-0.011)	-0.499 (-0.930,-0.111)	-0.365 (-0.742,-0.038)
dam mating order	0.060 (-0.002,0.136)	0.054 (-0.013,0.134)	0.007 (-0.037,0.052)	0.006 (-0.017,0.047)	-0.306 (-0.766,0.069)	0.210 (-0.116,0.587)
day mated (original)	0.326 (0.206,0.454)	0.122 (0.028,0.291)	0.058 (-0.006,0.149)	0.074 (0.037,0.139)	0.443 (-0.113,1.273)	0.797 (0.199,1.262)
dam preference for cowpea (original)	-0.073 (-0.180,0.031)	-0.025 (-0.174,0.068)	0.086 (0.027,0.144)	0.000 (-0.042,0.046)	0.219 (-0.366,0.733)	0.128 (-0.267,0.572)
Interaction: bean mass (novel)	-0.092 (-0.196,0.032)	0.031 (-0.131,0.138)	0.047 (-0.022,0.102)	0.040 (-0.011,0.091)	0.456 (-0.052,1.124)	-0.060 (-0.463,0.560)
Interaction: day mated (novel)	-0.017 (-0.141,0.142)	0.112 (-0.026,0.326)	-0.057 (-0.127,0.020)	-0.018 (-0.072,0.035)	-0.625 (-1.272,0.117)	-0.360 (-0.910,0.237)
Interaction: dam preference (novel)	0.051 (-0.056,0.206)	0.051 (-0.084,0.228)	-0.081 (-0.146,-0.015)	0.008 (-0.038,0.063)	-0.369 (-0.916,0.402)	-0.109 (-0.590,0.470)

966

Table 4. Phenotypic variance partitioning (A) and full model inference (B) based on G_{cov} model for the non-Gaussian traits – larval survival and host egg laying preference. The estimates for the variance parameters (A) are back-transformed to original data scale. The effect of host type, the additional experimental variables and dam host preference (B) for larval survival is on logit scale and for host preference on log scale. We embolden any credible intervals (in brackets) that do not overlap zero. The intercept always corresponds to the trait average on the original host type and the ‘novel HOST effect’ is the effect of the novel host type (i.e. difference between the two means). The same applies to interaction terms, where the interaction gives the estimate for the difference between the effect on the novel and the original host type. The estimates and their credible intervals come from posterior distribution of animal MCMC models.

A)

Host type	Par.	Larval survival	Host preference
		juveniles (N = 3 146)	daughters (N = 665)
original	V_A	0.006	<0.001
	V_P	0.094	0.671
	h^2	0.060	0.001
	CV_A	0.084	0.033
novel	V_A	0.003	0.001
	V_P	0.065	0.522
	h^2	0.040	0.003
	CV_A	0.055	0.071
r_G		0.286* (-0.469,0.985)*	-0.046* (-0.905,0.948)*

* these estimates are on the transformed (latent) scale

981 B)

Term	Larval survival	Host preference
	juveniles	daughters
	(N = 3 146)	(N = 665)
Intercept (original HOST)	2.868 (2.341,3.441)	-0.409 (-0.528,-0.338)
novel HOST effect	0.512 (-0.094,1.447)	-0.247 (-0.324,-0.180)
bean mass (original)	-0.161 (-0.353,0.050)	0.010 (-0.046,0.049)
dam mating order	-0.021 (-0.206,0.124)	-0.001 (-0.027,0.047)
day mated (original)	0.095 (-0.153,0.350)	-0.053 (-0.094,0.004)
dam preference for cowpea (original)	-0.042 (-0.258,0.183)	0.014 (-0.041,0.057)
Interaction: bean mass (novel)	-0.035 (-0.376,0.221)	0.017 (-0.075,0.076)
Interaction: day mated (novel)	-0.122 (-0.501,0.240)	-0.016 (-0.066,0.083)
Interaction: dam preference (novel)	0.242 (-0.106,0.549)	-0.037 (-0.096,0.049)

982

983

Figure caption

Figure 1. Illustration of how maternal effects might differ between environments (host beans in our study). Panel A) shows that ignoring the potential for environment-specific maternal effects, as indicated by the colour of points representing maternal effects in the original (black points) and the novel host (white points), could be misleading. Panel B) represents a scenario where variance due to maternal effects (V_M) differs between the environments, i.e. the first type of Maternal-by-Environment interaction ($M \times E$). Panel C) illustrates the other type of $M \times E$ where the rank of maternal effects differs between the hosts. The evidence for the shift in ranking comes from the low cross-environmental correlation in maternal effects (r_M) despite the maternal effects variance (V_M) being comparable between the hosts. Panel D) represents a combination of B) and C).