Association between the extent of DNA methylation at the CpG sites of *HIF3A* and parameters of obesity in the general Japanese population

Genki Mizuno ^{1,2}, Hiroya Yamada ³, Eiji Munetsuna ⁴, Mirai Yamazaki ⁵, Yoshitaka Ando ^{1,6}, Ryosuke Fujii ¹, Yoshiaki Tsuboi ¹, Atsushi Teshigawara ⁶, Itsuki Kageyama ⁶, Keisuke Osakabe ⁷, Keiko Sugimoto ⁷, Hiroaki Ishikawa ⁶, Naohiro Ichino ⁷, Yoshiji Ohta ⁸, Koji Ohashi ⁶, Shuji Hashimoto ³, and Koji Suzuki ^{1*}

¹ Department of Preventive Medical Sciences, Fujita Health University School of Medical Sciences, Toyoake, Aichi, Japan

² Department of Joint Research Laboratory of Clinical Medicine, Fujita Health University Hospital, Toyoake, Aichi, Japan

³ Department of Hygiene, Fujita Health University School of Medicine, Toyoake, Aichi, Japan

⁴ Department of Biochemistry, Fujita Health University School of Medicine, Toyoake, Aichi, Japan

⁵ Department of Medical Technology, Kagawa Prefectural University of Health Sciences, Takamatsu, Kagawa, Japan

16 ⁶ Department of Clinical Biochemistry, Fujita Health University School of Medical Sciences, Toyoake, Aichi,

17 Japan

18 ⁷ Department of Clinical Physiology, Fujita Health University School of Medical Sciences, Toyoake, Aichi,

19 Japan

20 ⁸ Department of Chemistry, Fujita Health University School of Medicine, Toyoake, Aichi, Japan

21

22 *Corresponding author

23 E-mail:ksuzuki@fujita-hu.ac.jp (KS)

24

25 Short Title

26 Association between DNA methylation of *HIF3A* and parameters of obesity

27

28

29

Abstract

Obesity is a major public health problem worldwide owing to the substantial increase in risk of metabolic diseases. Hypoxia-inducible factors (HIFs) regulate transcriptional responses to hypoxic stress. DNA methylation in the CpG sites of intron 1 of *HIF3A* is associated with body mass index in the whole blood and adipose tissue. This study investigates the correlation between DNA methylation of *HIF3A* and parameters of obesity, including thickness of visceral (VAT) and subcutaneous adipose tissues, in the general Japanese population. Participants (220 men and 253 women) who underwent medical examination were enrolled in this cross-sectional study. We used pyrosequencing to quantify DNA methylation (CpG sites of cg16672562, cg22891070, and cg27146050) in *HIF3A*. DNA methylation of *HIF3A* was only different in women. Multiple regression analysis showed that DNA methylation level at cg27146050 was associated with thickness of VAT in women. DNA methylation level at cg27146050 also correlated with body mass index and percentage of body fat in women after excluding smokers and non-smokers who quit smoking with the last 5 years. DNA methylation in the CpG site (cg27146050) of *HIF3A* correlated with parameters of obesity in Japanese women.

Introduction

Obesity is a major public health concern worldwide. In obese individuals, non-esterified fatty acids, adipokines, and other factors are extensively released from adipose tissues, thereby leading to abnormalities in obesity-related cell functions¹. Consequently, obesity induces various diseases, such as insulin resistance, type 2 diabetes, and cardiovascular disease^{2,3}. Thus, obesity is a risk factor for various metabolic diseases, and preventing obesity results in the prevention of metabolic diseases. Recent years have seen the diversification of lifestyle and eating habits that have increased the number of obese individuals globally⁴. Lifestyle, environmental factors, and genetic factors trigger obesity⁵.⁶. Lifestyle and/or environmental factors cause epigenetic alterations in several health conditions, such as obesity and metabolic disease⁷⁻¹⁰.

DNA methylation is an epigenetic mechanism that regulates gene expression by adding a methyl donor to cytosine to enable the regulation of transcription¹¹. Lifestyle factors, including dietary habits, modulate DNA methylation¹². Several animals¹³⁻¹⁵ and epidemiological studies¹⁶⁻¹⁸ have shown that environmental factors, including food intake, tobacco smoking, and alcohol consumption, cause DNA methylation in the blood or tissues. Moreover, global DNA hypermethylation in

leukocytes is associated with increased risk of cardiovascular diseases in the general Japanese population¹⁹. Thus, DNA methylation may be a novel biomarker for metabolic diseases caused by environmental factors and lifestyles.

Dick et al.²⁰ conducted two epigenetic genome-wide analyses to show the increase in DNA methylation at three CpG sites (cg16672562, cg22891070, and cg27146050) in intron 1 of *HIF3A* in the blood was associated with body mass index (BMI). Similarly, Main et al.²¹ and Wang et al.²² demonstrated that DNA methylation in *HIF3A* in the blood is associated with BMI in patients with type 2 diabetes and childhood obesity, respectively. Isoforms of HIF are constitutively expressed in mammalian cells and regulate transcriptional response to hypoxic stress^{23, 24}. HIFs are unstable at normal oxygen levels in mammalian cells. The reduction in normal cellular oxygen levels caused by environmental factors, diseases, effusion of blood, and adiposity stabilize HIFs, thereby enabling its nucleocytoplasmic translocation and binding to the hypoxia response element in the promoter of target genes and regulating target gene transcription and expression. Pfeiffer et al.²⁵ have shown that methylation of *HIF3A* in the adipose tissue correlates with dysfunctional human subcutaneous adipose tissue (SAT) and visceral adipose tissue (VAT). These studies indicate that DNA methylation of

HIF3A is associated with the development of obesity, and may be an obesity-related factor worldwide. Furthermore, DNA methylation of *HIF3A* in the blood is associated with insulin resistance in patients with type 2 or gestational diabetes ^{21,26}. There are only a few reports on the association between DNA methylation of *HIF3A* and BMI in humans. The thickness of adipose tissues is a more reliable parameter of obesity as compared to BMI that is an indirect parameter. To the best of our knowledge, there is no study on the correlation between DNA methylation of *HIF3A* and thickness of adipose tissues, such as VAT and SAT, that directly reflects obesity.

In this study, we attempted to verify whether DNA methylation of *HIF3a* (CpG sites of cg16672562, cg22891070, and cg27146050) in the blood associated with the thickness of VAT and SAT in the general Japanese population. We further determined whether DNA methylation in *HIF3A* in the blood correlated with the thickness of VAT and SAT in Japanese non-smokers ^{27,28}.

Materials and methods

Participants

This cross-sectional study was approved by the Ethics Review Committee of Fujita Health

University (Approval number: HG19-069). We enrolled 473 participants (220 men and 253 women) who took part in the medical examination of the general (middle-aged) population in Yakumo town, Hokkaido, Japan, in August 2015 ^{29,30}. We obtained written informed consent from all the participants for the use of individual genome samples. Information on lifestyle habits was obtained from questionnaires.

Measurements of obesity parameters

Parameters of obesity were measured as described previously ³¹. Percentage of body fat (% body fat) was measured using bioelectrical impedance analysis with the Tanita MC780 multifrequency segmental body composition analyzer (Tokyo, Japan). The thicknesses of VAT and SAT were assessed using ultrasound with ProSound a7 and UST-9130 convex probe (Hitachi Aloka Medical, Ltd, Tokyo, Japan). Thickness of VAT and SAT were defined as the distance (cm) from the peritoneum to the vertebral bodies and depth (cm) from the skin to the linea alba, respectively.

Blood test and determination of DNA methylation

105 Blood was collected during the medical examination of the general population, and the
106 serum was separated from the blood by centrifugation at 2,000×g for 10 min at room temperature. For
107 biochemical analysis of the blood, enzymes and components in the serum were assayed using an auto-
108 analyzer (JCS-BM1650, Nihon Denshi Co., Tokyo, Japan) at Yakumo General Hospital.

109 DNA methylation was analyzed using the buffy coat obtained upon centrifugation of the
110 blood collected in ethylenediaminetetraacetic acid (EDTA)-2Na-containing tubes under the same
111 conditions as those used for blood biochemical tests. Genomic DNA was extracted from the buffy coat
112 using the NucleoSpin Tissue kit (Takara, Shiga, Japan). Bisulfite conversion was performed using the
113 EpiTect Bisulfite Kit (Qiagen, Valencia, CA, USA). Polymerase chain reaction (PCR) was used to
114 amplify the intron 1 of *HIF3A* using EpiTaq™ HS (for bisulfite-treated DNA; Takara, Shiga, Japan).
115 Levels of DNA methylation were quantified using pyrosequencing with the PyroMark Q24 Advanced
116 kit (Qiagen, Valencia, CA, USA) and analyzed using the parameters previously described ²⁰⁻²²,
117 including three CpG sites (Fig 1). Table 1 lists the sequences of primers used for PCR and
118 pyrosequencing. The primers used for pyrosequencing were designed based on a previous study ²⁵
119 using PyroMark Assay Design 2.0 (Qiagen, Valencia, CA, USA).

Fig 1. A target sequence of intron 1 region in *HIF3A* gene

The target region of *HIF3A* gene DNA methylation analyzed by pyrosequence was decided based on previous studies. It has reported that the 3 CpG sites (cg16672562, cg22891070 and cg27146050) of intron 1 in *HIF3A* gene in the blood are associated with BMI in EWAS study.

Table 1. Sequences of primers used for PCR and Pyrosequence

Primer	Sequence (5'-3')
Forward	TGTTGAAGGGTATTTAGGG
Reverse	Biotin-ACTCTATCCCACCCCTTTT
Sequence 1	TTTAGGGGGTGTAGG
Sequence 2	GGTGAGATGATTTTATAGGAA

Statistical analysis

All statistical analyses were performed using JMP version 14.0 (SAS Institute, Cary, NC, USA). Serum aspartate transaminase (AST), alanine transaminase (ALT), triglyceride, and high-density lipoprotein (HDL) cholesterol levels have been represented by the geometric means and

interquartile ranges owing to log-normal distribution. Other characteristics (including DNA methylation) have been represented as mean±standard deviation (SD). We analyzed the association of DNA methylation level at each CpG site of intron 1 in *HIF3A* with the parameters of obesity using single correlation and multiple linear regression and adjusted for age, systolic blood pressure, hemoglobin A1c, %neutrophil, smoking habit and exercise habit. For multiple testing, the Bonferroni method was used to counteract the problem of multiple comparisons. $P<0.05$ was considered statistically significant.

Results

Table 2 lists the characteristics of the participants in this study. There were significant differences in various parameters of obesity between men and women, such as smoking habit and blood biochemical test, but not hemoglobin A1c and blood pressure. Moreover, DNA methylation levels at three CpG sites in intron 1 of *HIF3A* were significantly different between the sexes (Table 3).

Table 2. Characteristics of participants in this study

	Men	Women	P-value
n	220	253	
Age (years)	66.3 ± 8.28	64.5 ± 8.00	0.017 ^a
Blood glucose (mg/dL)	93.8 ± 14.5	87.6 ± 17.0	<0.001 ^a
Hemoglobin A1c (%)	5.80 ± 0.54	5.72 ± 0.55	0.190
AST (IU/L)	23.4 (19.0-26.8)	21.6 (18.0-24.5)	0.003 ^b
ALT (IU/L)	23.1 (17.3-31.0)	18.9 (14.0-24.0)	<0.001 ^b
Triglyceride (mg/dL)	102.5 (72.0-144.8)	86.3 (64.5-117.0)	<0.001 ^b
Total cholesterol (mg/dL)	203.5 ± 32.4	217.5 ± 35.0	<0.001 ^a
HDL cholesterol (mg/dL)	52.4 (44.0-61.0)	61.8 (53.0-72.0)	<0.001 ^b
LDL cholesterol (mg/dL)	121.3 ± 30.2	127.0 ± 31.1	0.042 ^a
Systolic blood pressure (mmHg)	134.8 ± 20.6	128.8 ± 19.2	0.001 ^a
Diastolic blood pressure (mmHg)	79.8 ± 13.0	72.9 ± 12.6	<0.001 ^a
Various parameters of obesity			

BMI (kg/m²)	24.1 ± 2.78	23.0 ± 3.60	<0.001 ^a
VAT thickness (cm)	64.6 ± 14.7	50.4 ± 11.7	<0.001 ^a
SAT thickness (cm)	14.3 ± 3.81	13.2 ± 4.62	0.006 ^a
% body fat	23.7 ± 4.16	32.8 ± 6.13	<0.001 ^a
Smoking habit, n (%)			
Never	45 (21)	193 (77)	<0.001 ^c
Ever	126 (57)	40 (16)	
Current	49 (22)	19 (7)	
Exercise habit, n (%)			
few	102 (47)	139 (55)	0.366
sometimes	43 (20)	44 (17)	
1 time/week	24 (11)	23 (9)	
> 2 times/week	49 (22)	47 (19)	

145 Values are mean ± SD, geometric mean (25-75th parentheses), or n (%). *P* < 0.05 was considered

146 statistically significant. a: student t test, b: Wilcoxon test, c: Pearson's chi-square test

147

148 Correlations between DNA methylation levels at the CpG sites in intron 1 of *HIF3A* and

149 parameters of obesity were analyzed using single linear regression owing to the differences in DNA

150 methylation of *HIF3A* in men and women (Tables 3 and 4). There was no significant correlation

151 between DNA methylation level at each CpG site and the parameters of obesity in men and women.

152 **Table 3. DNA methylation levels (%) at *HIF3A* gene sites by pyrosequence analysis**

CpG site	Men	Women	<i>P</i> -value
cg16672562	17.3 ± 5.12	20.1 ± 5.96	<0.001 ^a
cg22891070	21.5 ± 7.37	24.9 ± 8.03	<0.001 ^a
cg27146050	14.3 ± 4.74	17.1 ± 5.00	<0.001 ^a
mean	17.6 ± 5.00	20.6 ± 5.57	<0.001 ^a

153 Values are mean ± SD. *P* < 0.05 was considered statistically significant. a: student t test

154 **Table 4. Single correlation analysis between *HIF3A* gene DNA methylation levels and obesity**

155 **parameters in Japanese men and women**

Men								
CpG site	BMI		VAT		SAT		% Body fat	
	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>
cg16672562	0.087	0.210	0.082	0.244	0.029	0.682	0.074	0.295
cg22891070	0.114	0.111	0.028	0.689	0.045	0.525	0.054	0.444
cg27146050	0.013	0.854	-0.006	0.937	0.050	0.478	-0.021	0.766
mean	0.089	0.202	0.049	0.565	0.048	0.496	0.046	0.520
Women								
CpG site	BMI		VAT		SAT		% Body fat	
	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>
cg16672562	-0.029	0.661	-0.066	0.317	0.025	0.704	-0.026	0.690
cg22891070	0.024	0.720	-0.042	0.530	0.047	0.476	0.049	0.461
cg27146050	-0.038	0.562	-0.109	0.099	-0.028	0.679	-0.045	0.497
mean	0.176	0.791	0.075	0.263	0.019	0.771	0.006	0.926

157

158 Table 5 shows the results of multiple linear regression analysis for the correlation between

159 DNA methylation levels at CpG sites in intron 1 of *HIF3A* and the parameters of obesity. In men, there

160 was no significant correlation between DNA methylation level at each CpG site and the parameters of

161 obesity. In women, significant correlations were observed between DNA methylation level at

162 cg27146050 and VAT thickness ($P < 0.05$). However, DNA methylation levels at cg16672562 and

163 cg22891070 did not significantly correlate with any parameter of obesity in women.

164 **Table 5. Multiple linear regression analysis for correlations between *HIF3A* gene DNA**

165 **methylation levels and obesity parameters**

Men								
CpG site	BMI		VAT		SAT		% Body fat	
	β	P	β	P	β	P	β	P
cg16672562	0.060	0.391	0.092	0.175	-0.024	0.736	0.052	0.462
cg22891070	0.081	0.241	0.030	0.656	-0.007	0.922	0.040	0.570

cg27146050	-0.012	0.868	0.005	0.938	-0.016	0.819	-0.038	0.594
mean	0.059	0.402	0.050	0.467	-0.016	0.818	0.027	0.706
Women								
CpG site	BMI		VAT		SAT		% Body fat	
	β	P	β	P	β	P	β	P
cg16672562	-0.071	0.299	-0.084	0.220	-0.027	0.682	-0.062	0.369
cg22891070	-0.002	0.974	-0.060	0.374	0.004	0.946	0.021	0.755
cg27146050	-0.085	0.215	-0.161	0.029	-0.095	0.156	-0.095	0.165
mean	-0.059	0.400	-0.104	0.140	-0.039	0.569	-0.047	0.505

166 Adjusted for age, systolic blood pressure, hemoglobin A1c, %neutrophil, smoking habit and exercise

167 habit. $P < 0.05$ was considered statistically significant.

168

169 Smoking habits alter the status of DNA methylation^{27, 28}. Therefore, we examined whether

170 DNA methylation of the different regions of *HIF3A* were associated with the parameters of obesity in

171 non-smokers (i.e., the participants excluding current smokers and non-smokers who stopped smoking

within the last 5 years; Table 6). There was no significant correlation between DNA methylation level

at each CpG site and the parameters of obesity in men. In women, there were significant correlations

between DNA methylation level at cg27146050 and BMI, VAT thickness, and % body fat ($P < 0.05$).

Table 6. Multiple linear regression analysis for correlations between HIF3A gene DNA

methylation and obesity parameters in non-smokers

Men								
CpG site	BMI		VAT		SAT		% Body fat	
	β	P	β	P	β	P	β	P
cg16672562	0.086	0.306	0.098	0.228	-0.046	0.581	0.068	0.426
cg22891070	0.101	0.233	0.040	0.627	0.004	0.963	0.030	0.727
cg27146050	-0.017	0.847	0.012	0.883	-0.027	0.754	-0.067	0.449
mean	0.077	0.368	0.059	0.475	-0.228	0.790	0.020	0.821
Women								
	BMI		VAT		SAT		% Body fat	

CpG site	β	P	β	P	β	P	β	P
cg16672562	-0.009	0.103	-0.075	0.311	-0.050	0.492	-0.110	0.133
cg22891070	-0.059	0.413	-0.068	0.345	-0.007	0.924	-0.019	0.791
cg27146050	-0.152	0.040	-0.179	0.016	-0.110	0.132	-0.162	0.029
mean	-0.124	0.096	-0.114	0.129	-0.056	0.449	-0.102	0.170

Adjusted for age, systolic blood pressure, hemoglobin A1c, %neutrophil and exercise habit, and excluded for current smoker (include stopped smoking less than 5 years). $P < 0.05$ was considered statistically significant.

Discussion

We determined the association between DNA methylation at three CpG sites (cg16672562, cg22891070, and cg27146050) of the intron 1 of *HIF3A* and parameters of obesity in the general Japanese population. There was a significant difference in the DNA methylation of *HIF3A* between the sexes. Multiple linear regression analysis showed a correlation between DNA methylation at cg27146050 in *HIF3A* and thickness of VAT in women. Excluding current smokers and non-smokers

who stopped smoking within the last 5 years, there correlations between DNA methylation at cg27146050 in *HIF3A* and thickness of VAT thickness, BMI, and % body fat in women.

In this study, DNA methylation at each CpG site of intron 1 of *HIF3A* was higher in women than those in men. This was consistent with that reported by Main et al.²¹. This difference between men and women suggests that women exhibit lower expression of *HIF3A* during hypoxia than the expression in men under similar conditions owing to differential capacities of gene regulation. Women exhibit a relatively less pronounced physiological response to hypoxic stress than that in men³². This can be attributed to the increase in DNA methylation of *HIF3A* in women.

Dietary factors, such as nutrition, cause a change in DNA methylation¹². We have recently demonstrated that the intake of dietary vitamin affects lipid profiles via the modulation of DNA methylation within lipid-related genes¹⁶. DNA methylation variants of *HIF3A* are associated with alterations in BMI based on the consumption of total vitamins or supplemental vitamin B³³. Therefore, it is possible that the intake of vitamins or other nutrients causes a change in DNA methylation in *HIF3A*, thereby resulting in the development of obesity.

Tobacco smoking is another environmental factor that affects the incidence of obesity³⁴.

DNA methylation positively correlates with smoking habits^{27,28}. Thus, smoking habits may influence the association between *HIF3A* DNA methylation and the parameters of obesity. We determined whether DNA methylation in *HIF3A* associated with parameters of obesity in non-smokers (excluding current smokers and non-smokers who stopped smoking within the last 5 years). The correlation between DNA methylation at various sites of *HIF3A* and parameters of obesity increased in this population than that including non-smokers and smokers. To the best of our knowledge, this is the first report on the correlation between DNA methylation levels at the CpG sites in *HIF3A* and parameters of obesity, such as thickness of VAT and smoking habits, in the general Japanese population. However, smokers were not excluded from the group of non-smokers. Therefore, future studies should focus on determining the association between DNA methylation in *HIF3A* and parameters of obesity within non-smokers.

Dick et al.²⁰ demonstrated the correlation between DNA methylation levels at three CpG sites (cg16672562, cg22891070, and cg27146050) in intron 1 of *HIF3A* in the blood and BMI; this has also been confirmed by other studies^{21,34}. In women, DNA methylation at cg27146050 correlated with the thickness of VAT (based on multiple linear regression analysis). However, this association

has been reported in men²⁰. This discrepancy may be explained attributed to the differences in DNA methylation of *HIF3A* and smoking habits of men and women. Thus, future studies are warranted to elucidate the extent of DNA methylation in *HIF3A* between men and women.

Dick et al.²⁰ used a microarray to demonstrate the association between *HIF3A* DNA methylation and BMI in humans. Microarrays are useful in understanding global DNA methylation. However, this technique cannot measure methylation using immobilized methylated probes and exhibits poor quantification. Thus, this study employed pyrosequencing to analyze DNA methylation in *HIF3A*. This method is excellent in quantifying the extent of DNA methylation at selected CpG sites in specific target genes. Thus, pyrosequencing provides a more reliable scenario of the association between DNA methylation at CpG sites of intron 1 of *HIF3A* and parameters of obesity in the general Japanese population than the correlation reported by Dick et al.²⁰.

Finally, Hatanaka et al.³⁵ showed that the ectopic expression of *HIF3A* induces the expression of several adiposity-associated genes in 3T3-L1 cells. This suggests that low levels of DNA methylation in *HIF3A* upregulates *HIF3A*, thereby resulting in adiposity. Accordingly, we observed a positive correlation between DNA methylation in *HIF3A* and thickness of VAT (that directly reflects

obesity as compared to BMI) in women. Therefore, the thickness of VAT has important clinical implications in obesity-related diseases.

Taken together, the DNA methylation level at cg27146050 of intron 1 of *HIF3A* correlated well with parameters of obesity in non-smokers of the general Japanese women. This study has some limitations. First, the data do not show a causal relationship between DNA methylation at different sites of *HIF3A* and parameters of obesity since this was a cross-sectional study. Second, this study analyzed a small sample size. Finally, we did not determine alterations in the mRNA levels of *HIF3A*. Thus, future studies should focus on analyzing the association between DNA methylation level in *HIF3A* and the parameters of obesity over a longer period using a larger sample size. Furthermore, we will attempt to analyze the mRNA and protein levels of *HIF3A* in the blood of participants.

Conclusion

This is the first study to report the correlation between DNA methylation at CpG site in *HIF3A* and parameters of obesity, such as thickness of visceral adipose tissue and smoking habit, in the general Japanese population. DNA methylation of the CpG sites of *HIF3A* may be associated with

247 body mass index.

248

249 **Acknowledgments**

250 We thank the participants and staff of the Health Examination Program for Residents of

251 Yakumo, Hokkaido, Japan.

252

253 **Conflict of interest**

254 There is no conflict of interest.

255

256 **Reference**

257 1. Blüher M. Adipose tissue dysfunction in obesity. *Exp Clin Endocrinol Diabetes* 2009;117(6): 241-

258 250. doi: 10.1055/s-0029-1192044. PubMed PMID: 19358089.

259 2. Bastien M, Poirier P, Lemieux I, Després JP. Overview of epidemiology and contribution of obesity

260 to cardiovascular disease. *Prog Cardiovasc Dis* 2014;56: 369-381. doi: 10.1016/j.pcad.2013.10.016.

261 PubMed PMID: 24438728.

262 3. Kahn SE, Hull RL, Utzschneider KM. Mechanisms linking obesity to insulin resistance and type 2
263 diabetes. *Nature* 2006;444(7121): 840-846. doi: 10.1038/nature05482. PubMed PMID: 17167471.

264 4. Blüher M. Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol* 2019;15(5): 288-
265 298. doi: 10.1038/s41574-019-0176-8. PubMed PMID: 30814686.

266 5. Hruby A, Manson JE, Qi L, Malik VS, Rimm EB, Sun Q, et al. Determinants and Consequences of
267 Obesity. *Am J Public Health* 2016;106(9): 1656-1662. doi: 10.2105/AJPH.2016.303326. PubMed
268 PMID: 27459460.

269 6. Temelkova-Kurktschiev T, Stefanov T. Lifestyle and genetics in obesity and type 2 diabetes. *Exp*
270 *Clin Endocrinol Diabetes* 2012;120(1): 1-6. doi: 10.1055/s-0031-1285832. PubMed PMID: 21915815.

271 7. Yamada H, Ohashi K, Suzuki K, Munetsuna E, Ando Y, Yamazaki M, et al. Longitudinal study of
272 circulating miR-122 in a rat model of non-alcoholic fatty liver disease. *Clin Chim Acta* 2015;446: 267-
273 271. doi: 10.1016/j.cca.2015.05.002. PubMed PMID: 25958847.

274 8. Hiratsuka I, Yamada H, Munetsuna E, Hashimoto S, Itoh M. Circulating MicroRNAs in Graves'
275 Disease in Relation to Clinical Activity. *Thyroid* 2016;26(10): 1431-1440. doi: 10.1089/thy.2016.0062.
276 PubMed PMID: 27610819.

- 277 9. van Dijk SJ, Molloy PL, Varinli H, Morrison JL, Muhlhausler BS. Epigenetics and human obesity.
278 *Int J Obes (Lond)* 2015;39(1): 85-97. doi: 10.1038/ijo.2014.34. PubMed PMID: 24566855.
- 279 10. van Dijk SJ, Tellam RL, Morrison JL, Muhlhausler BS, Molloy PL. Recent developments on the
280 role of epigenetics in obesity and metabolic disease. *Clin Epigenetics* 2015;7: 66. doi:
281 10.1186/s13148-015-0101-5. PubMed PMID: 27408648.
- 282 11. Moore LD, Le T, Fan G. DNA methylation and its basic function. *Neuropsychopharmacology*
283 2013;38(1): 23-38. doi: 10.1038/npp.2012.112. PubMed PMID: 22781841.
- 284 12. Lim U, Song MA. Dietary and lifestyle factors of DNA methylation. *Methods Mol Biol* 2012;863:
285 359-376. doi: 10.1007/978-1-61779-612-8_23. PubMed PMID: 22359306.
- 286 13. Munetsuna E, Yamada H, Yamazaki M, Ando Y, Mizuno G, Hattori Y, et al. Maternal high-
287 fructose intake increases circulating corticosterone levels via decreased adrenal corticosterone
288 clearance in adult offspring. *J Nutr Biochem* 2019;67: 44-50. doi: 10.1016/j.jnutbio.2019.01.016.
289 PubMed PMID: 30856463.
- 290 14. Ohashi K, Munetsuna E, Yamada H, Ando Y, Yamazaki M, Taromaru N, et al. High fructose
291 consumption induces DNA methylation at PPAR α and CPT1A promoter regions in the rat liver.

292 *Biochem Biophys Res Commun* 2015;468: 185-189. doi: 10.1016/j.bbrc.2015.10.134. PubMed PMID:
293 26519879.

294 15. Yamazaki M, Munetsuna E, Yamada H, Ando Y, Mizuno G, Murase Y, et al. Fructose
295 consumption induces hypomethylation of hepatic mitochondrial DNA in rats. *Life Sci* 2016;149: 146-
296 152. doi: 10.1016/j.lfs.2016.02.020. PubMed PMID: 26869391.

297 16. Fujii R, Yamada H, Munetsuna E, Yamazaki M, Ando Y, Mizuno G, et al. Associations between
298 dietary vitamin intake, ABCA1 gene promoter DNA methylation, and lipid profiles in a Japanese
299 population. *Am J Clin Nutr* 2019;110(5): 1213-1219. doi: 10.1093/ajcn/nqz181. PubMed PMID:
300 31504085.

301 17. Fujii R, Yamada H, Munetsuna E, Yamazaki M, Mizuno G, Tsuboi Y, et al. Dietary vegetable
302 intake is inversely associated with ATP-binding cassette protein A1 (ABCA1) DNA methylation
303 levels among Japanese women. *Nutrition* 2019;65: 1-5. doi: 10.1016/j.nut.2019.02.010. PubMed
304 PMID: 31029915.

305 18. Gao X, Zhang Y, Breitling LP, Brenner H. Tobacco smoking and methylation of genes related to
306 lung cancer development. *Oncotarget* 2016;7(37): 59017-59028. doi: 10.18632/oncotarget.10007.

307 PubMed PMID: 27323854.

308 19. Tsuboi Y, Yamada H, Munetsuna E, Yamazaki M, Mizuno G, Murase Y, et al. Relationship
309 between Long Interspersed Nuclear Element-1 DNA Methylation in Leukocytes and Dyslipidemia in
310 the Japanese General Population. *J Atheroscler Thromb* 2018;25(12): 1231-1239. doi:
311 10.5551/jat.43570. PubMed PMID: 29628482.

312 20. Dick KJ, Nelson CP, Tsaprouni L, Sandling JK, Aïssi D, Wahl S, et al. DNA methylation and
313 body-mass index: a genome-wide analysis. *Lancet* 2014;383(9933): 1990-1998. doi: 10.1016/S0140-
314 6736(13)62674-4. PubMed PMID: 24630777.

315 21. Main AM, Gillberg L, Jacobsen AL, Nilsson E, Gjesing AP, Hansen T, et al. DNA methylation
316 and gene expression of HIF3A: cross-tissue validation and associations with BMI and insulin
317 resistance. *Clin Epigenetics* 2016;8(1): 89. doi: 10.1186/s13148-016-0258-6. PubMed PMID:
318 27594926.

319 22. Wang S, Song J, Yang Y, Zhang Y, Wang H, Ma J. HIF3A DNA Methylation Is Associated with
320 Childhood Obesity and ALT. *PLoS One* 2015;10(12): e0145944. doi: 10.1371/journal.pone.0145944.
321 PubMed PMID: 26717317.

- 322 23. Duan C. Hypoxia-inducible factor 3 biology: complexities and emerging themes. *Am J Physiol*
323 *Cell Physiol* 2016;310(4): C260-269. doi: 10.1152/ajpcell.00315.2015. PubMed PMID: 26561641.
- 324 24. Ravenna L, Salvatori L, Russo MA. HIF3 α : the little we know. *Febs j* 2016;283(6): 993-1003. doi:
325 10.1111/febs.13572. PubMed PMID: 26507580.
- 326 25. Pfeiffer S, Krüger J, Maierhofer A, Böttcher Y, Klötting N, El Hajj N, et al. Hypoxia-inducible
327 factor 3A gene expression and methylation in adipose tissue is related to adipose tissue dysfunction.
328 *Sci Rep* 2016;6: 27969. doi: 10.1038/srep27969. PubMed PMID: 27346320.
- 329 26. Zhang Y, Chen Y, Qu H, Wang Y. Methylation of HIF3A promoter CpG islands contributes to
330 insulin resistance in gestational diabetes mellitus. *Mol Genet Genomic Med* 2019;7(4): e00583. doi:
331 10.1002/mgg3.583. PubMed PMID: 30743315.
- 332 27. Maeda K, Yamada H, Munetsuna E, Fujii R, Yamazaki M, Ando Y, et al. Association of smoking
333 habits with TXNIP DNA methylation levels in leukocytes among general Japanese population. *PLoS*
334 *One* 2020;15(7): e0235486. doi: 10.1371/journal.pone.0235486. PubMed PMID: 32609762.
- 335 28. Silva CP, Kamens HM. Cigarette smoke-induced alterations in blood: A review of research on
336 DNA methylation and gene expression. *Exp Clin Psychopharmacol*. Epub ahead of print 13 July 2020.

- DOI: 10.1037/pha0000382. PubMed PMID: 32658533.
29. Ando Y, Yamazaki M, Yamada H, Munetsuna E, Fujii R, Mizuno G, et al. Association of circulating miR-20a, miR-27a, and miR-126 with non-alcoholic fatty liver disease in general population. *Sci Rep* 2019;9(1): 18856. doi: 10.1038/s41598-019-55076-z. PubMed PMID: 31827150.
30. Yamada H, Suzuki K, Ichino N, Ando Y, Sawada A, Osakabe K, et al. Associations between circulating microRNAs (miR-21, miR-34a, miR-122 and miR-451) and non-alcoholic fatty liver. *Clin Chim Acta* 2013;424: 99-103. doi: 10.1016/j.cca.2013.05.021. PubMed PMID: 23727030.
31. Munetsuna E, Yamada H, Ando Y, Yamazaki M, Tsuboi Y, Kondo M, et al. Association of subcutaneous and visceral fat with circulating microRNAs in a middle-aged Japanese population. *Ann Clin Biochem* 2018;55: 437-445. doi: 10.1177/0004563217735124. PubMed PMID: 28920467.
32. Botek M, Krejčí J, McKune A. Sex Differences in Autonomic Cardiac Control and Oxygen Saturation Response to Short-Term Normobaric Hypoxia and Following Recovery: Effect of Aerobic Fitness. *Front Endocrinol (Lausanne)* 2018;23(9): 697. doi: 10.3389/fendo.2018.00697. PubMed PMID: 30532736.
33. Huang T, Zheng Y, Qi Q, Xu M, Ley SH, Li Y, et al. DNA Methylation Variants at HIF3A Locus,

- 352 B-Vitamin Intake, and Long-term Weight Change: Gene-Diet Interactions in Two U.S. Cohorts.
- 353 *Diabetes* 2015;64(9): 3146-3154. doi: 10.2337/db15-0264. PubMed PMID: 26001398.
- 354 34. Watanabe T, Tsujino I, Konno S, Ito YM, Takashina C, Sato T, et al. Association between Smoking
- 355 Status and Obesity in a Nationwide Survey of Japanese Adults. *PLoS One* 2016;11(3): e0148926. doi:
- 356 10.1371/journal.pone.0148926. PubMed PMID: 27007232.
- 357 35. Hatanaka M, Shimba S, Sakaue M, Kondo Y, Kagechika H, Kokame K, et al. Hypoxia-inducible
- 358 factor-3alpha functions as an accelerator of 3T3-L1 adipose differentiation. *Biol Pharm Bull*
- 359 2009;32(7): 1166-1172. doi: 10.1248/bpb.32.1166. PubMed PMID: 19571379.

Sequence 1→

5' – GTAGGAGGGGATGCGGTGTAGTTAGGATTCGGGGTGCGAGTTACGAGT

Sequence 2→

GGGTGCGTACGGCGGTGAGATGATTTTATAGGAAAGGGTCTGGTTTTG

cg16672562

cg22891070

GGTGGGGAGGGGGGGGTATTCGAGTTTAGTTAAGAGGGGGTTTTTATT

AGTTAGGAGGGGGGCGTTGAGAGGGGGCGGAACGATAGTTGGTTTAAAA – 3'

cg27146050

Figure