Genome-wide identification, characterization, and expression analysis of the monovalent cation-proton antiporter superfamily in maize, and functional analysis of its role in salt tolerance

Mengsi Kong^{1,2†}, Meijie Luo^{2†}, Jingna Li², Zhen Feng^{2,3}, Yunxia Zhang², Wei Song², Ruyang Zhang², Ronghuan Wang², Yuandong Wang², Jiuran Zhao^{2*}, Yongsheng Tao^{1*}, Yanxin Zhao^{2*}

¹ College of Agronomy, Hebei Agricultural University, Baoding 071001, Hebei, China.

² Beijing Key Laboratory of Maize DNA Fingerprinting and Molecular Breeding, Maize Research Center, Beijing Academy of Agriculture and Forestry Sciences (BAAFS), Beijing 100079, China.

³ Plant Science and Technology College, Beijing University of Agriculture, Beijing 102206, China.

[†] Mengsi Kong and Meijie Luo contributed equally to this work.

* Correspondence: rentlang2003@163.com (Yanxin Zhao); maizezhao@126.com (Jiuran Zhao); yshtao@126.com (Yongsheng Tao)

ABSTRACT

Na⁺, K⁺ and pH homeostasis are important for plant life and they are controlled by the monovalent cation proton antiporter (CPA) superfamily. The roles of *ZmCPAs* in salt tolerance are not fully elucidated. In this study, we identified 35 *ZmCPAs* comprising 13 Na⁺/H⁺ exchangers (*ZmNHXs*), 16 cation/H⁺ exchanger (*ZmCHXs*), and 6 K⁺ efflux antiporters (*ZmKEAs*). All *ZmCPAs* have transmembrane domains and maize protoplasts transient expression showed that *ZmNHX5* and *ZmCHX17* localized to cell membrane system. *ZmCHXs* were specifically highly expressed in anthers, while *ZmNHXs* and *ZmKEAs* showed high expression in various tissues. *ZmNHX5* and *ZmKEA2* were up-regulated in maize seedlings under both NaCl and KCl stresses. Yeast complementation experiments revealed the roles of *ZmNHX5*, *ZmKEA2* in NaCl tolerance. Analyses of the maize mutants further validated the salt tolerance functions of *ZmNHX5* and *ZmKEA2*. Our study highlights comprehensive information of *ZmCPAs* and provides new gene targets for salt tolerance maize breeding.

Keywords: CPA, Maize, Salinity, *ZmNHX*, *ZmCHX*, *ZmKEA*

1. Introduction

High salinity is a significant abiotic stress factor restraining crop production worldwide. High concentration of sodium chloride (NaCl) or potassium chloride (KCl) affects the osmotic potential of the soil solution, thus inhibiting water absorption by plant. In addition, Na⁺ and Cl⁻ ions are easily absorbed to toxic levels by plant roots, which cause ion toxicity to plant cells [1, 2]. Under the above effects, salt stress eventually leads to plant metabolism disorder, oxidation-reduction system damage, photosynthesis limitation, and finally plant growth inhibition [3].

Many monovalent cation channels/transporters have been reported to play key roles in regulating salt tolerance in plants. In *Arabidopsis*, *AtNHX1*–6 and *AtNHX7/AtSOS1* were shown to be involved in maintaining Na⁺, K⁺ and pH homeostasis, and are critical for *Arabidopsis* salt tolerance [4, 5]. *AtKEA1*, *AtKEA3* and *AtKEA4* can enhance the tolerance of *Arabidopsis* to high concentration of K⁺, and *AtKEA2* overexpression enhanced the resistance of yeast mutants to Na⁺ and K⁺ stress. Functional characterization of *AtCHX17*, *AtCHX21* and *AtCHX23* knockout *Arabidopsis* mutants suggested their roles in K⁺ acquisition and in *Arabidopsis* salt tolerance. In rice, the expression of *OsNHX1*–5 and *OsNHX7/OsSOS1* were increased by salt stress and they can suppress sensitivity of rice to high Na⁺ and K⁺ concentration [6, 7]. The K⁺/Rb⁺ competition assay of *OsCHX14* in *Xenopus laevis* oocytes verified its function in K⁺ homeostasis. In maize, the expression of *ZmNHX2* was up-regulated under NaCl stress in both shoot and root of maize seedling and it was presumed to be responsible for Na⁺ sequestration into vacuoles [8]; *ZmNHX7* putatively encodes a Na⁺ efflux plasma membrane protein and its expression increased in maize plants exposed to salt stress [9]. All these Na⁺/H⁺ exchangers (NHXs), K⁺ efflux antiporters (KEAs) and cation/H⁺ exchangers (CHXs) carried a conserved Na⁺/H⁺ exchanger protein domain (Pfam00999), and are members of the monovalent cation/proton antiporter (CPA) superfamily. However, the CPA gene family in maize has not been comprehensively studied, and the role of its members in salt tolerance have not been fully elucidated.

The CAP superfamily is grouped into the CPA1 and CPA2 families based on the Transporter Classification database (<http://www.tcdb.org/>) [10, 11] and the CPA1 family

comprises the *NHXs* subfamily, while the CPA2 family is comprised of the *KEAs* and the *CHXs* subfamilies [4, 12]. The *NHXs* is predicted to contain 10–12 transmembrane (TM) domains. The *AtNHXs* consist of 8 members in *Arabidopsis*, and the *AtNHX1–4* and *AtNHX5–6* are localized inside cell, while *AtNHX7/AtSOS1* and *AtNHX8* are localized to the plasma membrane (PM) [4, 5]. The different subcellular location and tissue expression specificity between members are consistent with their different biological functions. For instance, the *AtNHX1–2* on vacuolar membrane are responsible for cell expansion in rapidly elongating organs; the *AtNHX5* and *AtNHX6* located at the trans-Golgi network are important for vesicle trafficking to the vacuole in different organs throughout plant development, while *AtNHX7* on PM affect membrane trafficking and vacuolar biogenesis in root cells [4, 5]. Compared with *Arabidopsis* and rice, the function of *ZmNHXs* in maize have not been well characterized.

The CPA2-type transporters have 8–14 putative TM domains [4]. The *Arabidopsis* had 6 members in the KEA subfamily [13]. The *AtKEA1* and *AtKEA2* are anchored in the inner envelope membrane of chloroplasts and are involved in chloroplasts division and membrane formation. The *AtKEA3* is located in the thylakoid membrane and it benefits photosynthesis efficiency [14, 15]. The *AtKEA4–6* were found mainly at the Golgi and they have a dramatic impact on K^+ , pH homeostasis in endomembrane compartments and in plant growth [4]. The function of *OsKEAs* in rice is rarely known. For the CHX subfamily, the *Arabidopsis* had 28 members. Among them, 18 *AtCHX* genes are pollen-specific or pollen-enhanced, and 6 are highly expressed in vegetative tissues, revealing the crucial role of *AtCHXs* in pollen development [4, 16]. In rice, *OsCHX14* was demonstrated to locate at Endoplasmic reticulum and regulate K^+ homeostasis in flowering organs [17]. In maize, the members of *ZmKEAs* and *ZmCHXs* have not been investigated and their roles in salt tolerance need to be determined.

In this study, we conducted genome-wide identification of *ZmCPA* genes in maize. Their encoded proteins were classified into *ZmNHX*, *ZmKEA* and *ZmCHX* subfamilies, and their protein properties comprising molecular weight (MW), isoelectric point (pI), evolutionary relationships, number of TM domains, sub-cellular localization, and motifs were determined. The splice variants, exon–intron structure, and intron phases of identified genes were

investigated. The transcriptional patterns of *ZmCPA* genes were analyzed during numerous developmental stages and under salinity stress. Six *ZmCPA* genes were functionally characterized by expression in a salt-sensitive yeast mutant. Analyses of two maize mutants, *zmnhx5* and *zmkea2*, further verified the functions of *ZmNHX5* and *ZmKEA2* in salt tolerance.

2. Results

2.1. Genome-wide identification and characterization of maize CPA superfamily genes

Extensive BLAST searches based on known *AtCPAs* identified a total of 35 *ZmCPA* superfamily proteins encoded in the genome of maize (Table S1). These were further confirmed through searching the Pfam database (<http://pfam.xfam.org/search>) for the existence of the signature Na⁺/H⁺ exchanger (PF00999) domain. The 35 genes were classified into three subfamilies: *ZmNHX* (13 genes), *ZmCHX* (16 genes) and *ZmKEA* (6 genes). The identified CPA genes were named *ZmNHX1–ZmNHX13*, *ZmCHX1–ZmCHX12*, *ZmCHX14–ZmCHX17*, and *ZmKEA1–ZmKEA6* according to previous reports or based on their degree of similarity with the family members in other species. The *ZmCPAs* were distributed on all chromosomes of maize, with 5, 3, 3, 4, 1, 3, 5, 4, 3, and 4 genes on chromosome 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10, respectively (Fig. S1).

As most of the *ZmCPA* genes had more than one transcript annotated in the MaizeGDB database (Table S1 and Fig. S2). The transcripts with a conserved gene structure similar to that of their orthologs (*Arabidopsis*, rice, *Sorghum*, *Brachypodium distachyon*, and *Setaria italica*) were selected for further analyses. The transcript of each *ZmCPA* gene model was further validated using reverse transcription polymerase chain reaction (RT-PCR) assays with the gene-specific primers listed in Table S2. Notably, the transcripts of five *ZmCPA* genes verified by Sanger sequencing of cDNA were different from their annotation in MaizeGDB (Fig. S3).

The gene structures of the *ZmCPA* genes were determined by aligning their genomic sequences with cDNA sequences obtained by RT-PCR. In general, the *ZmCPA* genes in the same subfamily shared a conserved gene structure and the majority of introns in each family were in 0 phase (Fig. 1). The number of introns ranged from 11 to 25, from 0 to 5, and from

146 16 to 19 in *ZmNHXs*, *ZmCHXs* and *ZmKEAs*, respectively (Table 1). The average length of
147 *ZmNHX*, *ZmCHX*, and *ZmKEA* proteins was 640, 824, and 815 amino acid (aa) residues,
148 respectively. The average molecular weight (MW) of *ZmNHX*, *ZmCHX*, and *ZmKEA*
149 proteins was 70.8, 88.4, and 87.4 kDa, respectively. The isoelectric point (pI) value of
150 *ZmNHX*, *ZmCHX*, and *ZmKEA* proteins ranged from 5.28 to 9.07, from 5.95 to 9.88, and
151 from 5.22 to 6.01, respectively (Table 1).

152

153 **2.2. Phylogenetic analysis of maize CPA genes**

154 To analyze the evolutionary relationships among the identified *ZmCPA* genes, we aligned
155 their protein sequences with 26 *OsCPA* and 42 *AtCPA* proteins to construct a neighbor-joining
156 (NJ) phylogenetic tree. Detailed information for *AtCPA* and *OsCPA* genes is shown in Table
157 S3. Based on the topology of the phylogenetic tree, the *CPA* gene superfamily in plants can be
158 subdivided into three subfamilies; *NHX*, *CHX*, and *KEA*. The *NHX*, *CHX*, *KEA* subfamilies
159 can be further divided into groups N1–3, C1–3, and K1–2, respectively (Fig. 2). All groups
160 contained genes from the three species of *Arabidopsis*, rice, and maize, indicating similar
161 functions of orthologous genes in each group. We also performed a phylogenetic analysis
162 using *ZmCPAs* only. As shown in Fig. 3, *ZmCPA* genes were grouped into *ZmNHX*, *ZmCHX*,
163 and *ZmKEA* three subfamilies, which was consistent with the classification shown in Fig. 1
164 (Fig. 3).

165

166 **2.3. Motif and subcellular localization analysis of ZmCPA proteins**

167 Known conserved domains of *ZmCPA* proteins were identified by screening the Pfam
168 database (<http://pfam.xfam.org/>). All proteins contained the Na⁺/H⁺ exchanger domain
169 (PF00999), and three *ZmKEAs* and five *ZmNHXs* also contained the Trka_N domain
170 (PF02254) and the TatD_DNase domain (PF01026), respectively (Fig. S4). Putative motifs of
171 the *ZmCPA* proteins were mined using tools at the MEME server
172 (<http://meme-suite.org/tools/meme>). The 15 motif sequences identified are listed in Table S4.
173 Generally, *ZmNHX* proteins contained five to eight conserved motifs, the *ZmCHX* proteins
174 had six to seven conserved motifs and the *ZmKEA* proteins had two to three conserved motifs

(Fig. S4). Most closely related members in the same subfamily had common motifs, indicative of functional similarities among ZmCPA proteins.

The conserved domains of ZmCPA proteins corresponded to partial motifs and covered the TM domain (Figure S4–5). The TM helices within ZmCPA proteins were predicted using tools at the TMHMM Server v. 2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>). All ZmCPA proteins had TM domains, and each subfamily had its own characteristics (Fig. S5). A maximum of 12 TM regions were predicted in ZmNHX5, -7, ZmCHX7, -16, -17, and ZmKEA1. The average occurrence of TM regions was 8, 10, and 10 in ZmNHX, ZmCHX and ZmKEA proteins, respectively (Table S1). The TM domain was mainly located at the C-terminal in ZmCHX subfamily, while at the N-terminal in ZmKEAs.

We performed transient expression assays in maize protoplasts to determine the subcellular localization of ZmCHX17 and ZmNHX5 proteins. The ZmNHX5-GFP and ZmCHX17-GFP fusion proteins co-localized with OsSCMP1-RFP protein (rice SECRETORY CARRIER MEMBRANE PROTEIN 1 fused with RFP as a membrane protein control [18]) suggesting that ZmNHX5 and ZmCHX17 localize to the membrane system (Fig. 4).

2.4. Gene expression analysis in different tissues and developmental stages

NimbleGen maize microarray data [19] (ZM37) including 60 tissues representing 11 major organs and various developmental stages of the B73 maize inbred line were analyzed to detect the expression patterns of *ZmCPA* genes. Gene expression data for 31 *ZmCPA* genes including 10 *ZmNHXs*, 16 *ZmCHXs* and 5 *ZmKEAs* were used for the cluster analysis. As revealed by the heatmap, all *ZmNHX* genes except *ZmNHX4*, *ZmNHX3* and *ZmNHX9* were highly expressed in nearly all 60 tissues. All five *ZmKEA* genes were highly expressed in all 60 tissues. All *ZmCHX* genes were expressed at lower levels than *ZmNHX* and *ZmKEA* genes, and *ZmCHX* genes were only expressed at high levels in anther (Fig. 5).

To confirm the tissue-specific expression of *ZmCPA* genes as revealed by the microarray data, semi-quantitative reverse transcription polymerase chain reaction (semi qRT-PCR) analyses of nine *ZmCPA* genes were performed with total RNA isolated from the roots, leaves,

ears, immature tassels, pollen, anthers, silk, and whole seeds (20 days after pollination) of the B73 inbred line. The primers for semi qRT-PCR are listed in Table S5. Generally, the expression patterns of nine *ZmCPA* genes were consistent with the results from the microarray data. The *ZmNHX5*, *ZmNHX6* and *ZmNHX8* were highly expressed in root and leaf of maize, while the *ZmKEA1*, *ZmKEA2* and *ZmKEA5* were highly expressed in most of 8 tissues. The *ZmCHX8* showed low expression in all tissues, and the *ZmCHX5* together with *ZmCHX17* were specifically highly expressed in anther and pollen of maize (Fig. 5 and Fig. 6).

211

2.5. Expression analysis of *ZmCPA* genes under 100 mM NaCl and 100 mM KCl stress

To understand the expressional responses of *ZmCPA* genes to salt stress, genes in two gene subfamilies, *ZmNHX* (four genes) and *ZmKEA* (four genes), were selected for analysis by real-time quantitative reverse transcription polymerase chain reaction (qRT-PCR) analyses. The primers are listed in Table S6. We detected the gene expression response at 0h, 1 h, 2 h, 4 h, and 24 h of salt stress. In the leaf of plants treated with 100 mM NaCl, *ZmNHX2*, *ZmNHX5*, *ZmNHX6*, *ZmKEA1* and *ZmKEA2* were up-regulated, while other genes were down-regulated. In the root, *ZmNHX5* and *ZmKEA2* were up-regulated after 4 h, while other genes were down-regulated or did not change much. Under 100 mM KCl stress treatment, all all eight *ZmCPAs* (*ZmNHX2*, *ZmNHX5*, *ZmNHX6*, *ZmNHX8*, *ZmKEA1*, *ZmKEA2*, *ZmKEA5* and *ZmKEA6*) had a tendency of up-regulation in leaf. In the root, only *ZmNHX5* was up-regulated while other genes were down-regulated or showed fluctuation (Fig. 7). These results implied that *ZmCPAs* might play a role in salinity stress tolerance through expression regulation.

226

2.6. Functional analysis of *ZmCPA* genes in a yeast mutant under NaCl stress

To test the functions of *ZmCPAs* in salt tolerance, the coding sequences of *ZmNHX5*, *ZmCHX2*, *ZmCHX3*, *ZmCHX5*, and *ZmCHX17* and *ZmKEA2* were each cloned into the yeast expression vector pDR196 under the control of the PMA1 promoter. Each vector was introduced into the *Saccharomyces cerevisiae* mutant strain AXT3K, which lacks the functions of four plasma membrane Na⁺-ATPases (ScENA1–4), the plasma membrane Na⁺,

233 K^+/H^+ antiporter ScNHA1, as well as the vacuolar Na^+ , K^+/H^+ antiporter ScNHX1 [20].
 234 Therefore, this mutant is very sensitive to Na^+ stress. Each transformed yeast strain was
 235 grown on Arg phosphate (AP) medium with or without 20 mM NaCl. Results showed that the
 236 transformed AXT3K lines expressing these genes grew much better than the AXT3K mutant
 237 as well as the AXT3K mutant transformed with the empty pDR196 vector under 20 mM NaCl
 238 stress (Fig. 8). These results confirmed the functions of *ZmNHX5*, *ZmCHX2*, *ZmCHX3*,
 239 *ZmCHX5*, and *ZmCHX17* and *ZmKEA2* in salt tolerance.

240

241 **2.7. Functional analysis of *ZmNHX5* and *ZmKEA2* under high concentration salt stress**

242 We obtained two maize mutants, *zmnhx5* (EMS4-0a18d8) and *zmkea2* (EMS4-02c2af),
 243 which were produced by EMS mutagenesis of the B73 inbred line, from the Maize
 244 EMS-induced Mutant Database (MEMD) [21]. The *zmnhx5* and *zmkea2* mutants have
 245 pre-termination mutations in *ZmNHX5* (*Zm00001d022504*) and *ZmKEA2* (*Zm00001d026645*),
 246 respectively, resulting in the production of truncated proteins (Fig. 9A, E). The phenotypes of
 247 inbred B73 lines and the two maize mutants were analyzed after 4 days of growth under salt
 248 stress. Under normal conditions, the growth status of the wild type and mutants was not
 249 significantly different (Fig. 9). However, under 100 mM NaCl treatment, the shoot length of
 250 the *zmkea2* mutant was significantly lower than that of the wild type ($P<0.05$). The shoot dry
 251 weight of *zmkea2* was reduced greater than that of the wild type (Fig. 9B–D). Similarly, under
 252 100 mM KCl condition, the shoot length of the *zmnhx5* mutant were significantly lower than
 253 that of the wild type ($P<0.05$). The shoot dry weight of *zmnhx5* was reduced greater than that
 254 of the wild type. ((Fig. 9F–H)). These results further verified that the *ZmNHX5* and *ZmKEA2*
 255 are important salt tolerance-related genes.

256

257 **3. Discussion**

258 The cation/proton antiporter superfamily is responsible for monovalent cations exchange
 259 in fungi, bacteria, animals, and plants [12, 16]. Functional analyses of CPA genes have been
 260 conducted in *Arabidopsis* [22, 23], rice [17], wheat [24] and *Arachis hypogea* [25]. Most of
 261 them acted on maintaining the pH, Na^+ and K^+ homeostasis in plants under high salinity [26].

262 In maize, the *ZmNHX7* is homologous to *AtNHX7/SOS1*, which is critical for excluding Na^+
 263 from plant cell. The expression of *ZmNHX7* was up-regulated upon NaCl stress, and it is
 264 associated with the major quantitative trait locus (QTL), *qST1*, which confers salt tolerance on
 265 maize plants [27, 28]. Till now, there are few studies on maize *CPAs* gene family, and the
 266 functional analysis of *ZmCPAs* are rarely performed.

267 In this study, 35 *ZmCPAs* were identified in maize, and they were named according to
 268 previous reports [29] or based on their sequence similarity to known plant *CPA* genes.
 269 Analysis of chromosomal distribution revealed that *ZmCPAs* are evenly distributed on the 10
 270 chromosomes of maize. Similarly, *CPA* genes are located on all chromosomes in wheat [24],
 271 and 15 out of 17 chromosomes in pear [30]. A phylogenetic tree using *CPA* proteins from
 272 *Arabidopsis*, rice and maize revealed that the homologous proteins from maize, *Arabidopsis*
 273 and rice were tightly clustered. For example, the *ZmNHX7*, *AtNHX7* and *OsNHX7* clustered
 274 together; the *ZmKEA3*, *AtKEA3* and *OsKEA3* gathered together; the *ZmCHX1*, *AtCHX1* and
 275 *OsCHX1* were in one clade. In *Arabidopsis*, genes with similar functions are clustered nearby,
 276 e.g., the *AtNHX1–4* localized on vacuolar membrane, *AtNHX5–6* for vesicle trafficking and
 277 *AtNHX7–8* localized on PM were distributed together, respectively. Thus it was speculated
 278 that the clustered homologous genes from maize and *Arabidopsis* or clustered maize genes
 279 had similar functions. Consistent with previous studies in *Arabidopsis* and pear, we also
 280 classified the *CPA* superfamily proteins into NHX, KEA and CHX subfamilies and into the
 281 N1–N3, K1–K2, and C1–C3 groups [12, 30].

282 The predicted sub-cellular localization of these proteins in different species is consistent
 283 to some extent. The *AtNHXs* exhibit vacuole, endosome, and PM localization [4] and the
 284 *ZmNHXs* have similar predicted localizations. Like *AtKEA1*, *AtKEA2* and *AtKEA3*, the
 285 *ZmKEA4* was predicted to localize to the chloroplast. All *ZmCHX* proteins were predicted to
 286 localize to the PM, like *AtCHX13* and *AtCHX14* [31]. The *ZmNHX5* and *ZmCHX17* were
 287 selected for transient expression in maize protoplast cells. As predicted, they were localized to
 288 the membrane system.

289 We analyzed the expression patterns of *ZmCPA* genes according to the NimbleGen maize
 290 microarray data. The *ZmNHX* and *ZmKEA* genes showed high expression at multiple

developmental stages, suggesting that they are important for maize growth and development. The *ZmCHXs* were specifically expressed in anthers, indicating that they participate in reproductive organ development. Similar expression trends of *CPA* genes have been found in other plant species. For example, in wheat, *TaNHX2*, *TaNHX5* and *TaNHX8* are highly expressed at multiple developmental stages, and most *TaCHX* family genes are relatively highly expressed in anthers [24]. The expression patterns of selected genes in maize were verified by semi-qRT-PCR in this study.

The differential expression of *ZmCPAs* between the control and salinity treatments were analyzed using qRT-PCR. Under 100 mM NaCl treatment, the *ZmNHX5* and *ZmKEA2* were up-regulated in both leaf and root, providing evidence of their roles in NaCl stress in maize. Under 100 mM KCl treatment, the *ZmNHX2*, *ZmNHX5*, *ZmNHX6*, *ZmNHX8*, *ZmKEA1*, *ZmKEA2*, *ZmKEA5* and *ZmKEA6* were up-regulated in the leave. The up-regulated expression of them to K⁺ stress indicates that they might be involved in maintaining K⁺ homeostasis in maize. The *ZmNHX5*, *ZmKEA2*, *ZmCHX2*, *ZmCHX3*, *ZmCHX5* and *ZmCHX17* and were respectively introduced into the yeast mutant strain AXT3K. All constructs restored Na⁺-resistance of the AXT3K mutant. Furthermore, the growth reduction of maize EMS mutants under salt stress validated the functions of *zmnhx5* and *zmkea2* in salt tolerance. These analyses provided evidence for the role of CHX and KEA subfamilies in salt tolerance in maize. Similarly, *AtNHX5* and *AtNHX6* recovered salt tolerance in yeast [23], indicating that *ZmNHXs* have a common mode of action as other plant species. Unlike *ZmCHX* and *ZmKEA* from maize, neither *AtKEAs* nor *AtCHXs* from *Arabidopsis* are shown to improve yeast growth under high salt condition [4, 13].

It is worth mentioning that both *ZmNHX5* and *ZmKEA2* played an important role in salt tolerance of maize according to the results obtained in this study. The expression of *ZmNHX5* and *ZmKEA2* were up-regulated under salt stress (NaCl and KCl), and they can restore the Na⁺-resistance of yeast mutant. Moreover, the salt tolerance function (NaCl) of *ZmNHX5* and *ZmKEA2* were further verified by maize mutant experiment. The functional analysis of these two genes needs further study, and they are likely to be new gene targets for breeding salt tolerance maize.

320 4. Conclusion

321 We identified and characterized the *ZmCPA* superfamily, consisting of *ZmNHX*, *ZmKEA*
322 and *ZmCHX* subfamily genes, from the maize genome. Gene and protein sequences analyses
323 demonstrated high conservation of evolutionarily related members in each subfamily, while
324 members differed among different groups. The existence of many helices, coils and TM
325 regions in protein structure are consistent with the hydrophobic and membrane-bound nature
326 of these proteins. Expression analyses revealed diverse expression patterns in various tissues
327 and under different abiotic stresses, indicating that these genes are important for maize growth,
328 development, and salt stress responses. Characterization of *ZmNHX5*, *ZmCHX2*, -3, -5, -17
329 and *ZmKEA2* in yeast confirmed their roles in maintaining monovalent cation homeostasis
330 and in salt stress tolerance. The functions of *ZmNHX5* and *ZmKEA2* were verified by
331 phenotypic analyses of EMS mutants. This study not only describes numerous features of
332 *ZmCPA* genes, but also provides the foundation for functional validation of these genes in
333 further studies.

334

335 5. Materials and methods

336 5.1. Identification and bioinformatic analysis of *ZmCPA* gene superfamily

337 The known *CPA* genes in *Arabidopsis* were used to query the maize AGPv4 gene set
338 (<https://download.maizegdb.org/Zm-B73-REFERENCE-GRAMENE-4.0/>) using a local
339 BLASTP program with an E-value <1e-10. The putatively identified sequences were further
340 confirmed by an HMMER search for the presence of signature Na⁺/H⁺ exchanger (PF00999)
341 domain.

342 The phylogenetic tree was constructed using full-length CPA protein sequences of maize,
343 *Arabidopsis*, and rice. Sequences were aligned using MUSCLE v3.8.31 [32, 33], and a
344 phylogenetic tree was built using the neighbor-joining (NJ) method with MEGA 6.0 [34] with
345 a bootstrap value of 1000 and amino acid substitution according to the Poisson model. The
346 TM domains were predicted using tools at the TMHMM Server v. 2.0
347 (<http://www.cbs.dtu.dk/services/TMHMM/>). Predotar [35], TargetP [36], and WoLF PSORT
348 (<https://www.genscript.com/wolf-psort.html>), three *in silico* programs, were used to predict

the organellar localization of ZmCPA proteins. Putative conserved motifs in maize CPA proteins were identified using the MEME Suite 5.1.1 (<http://meme-suite.org/tools/meme>) with the following parameters: motif length, 10–50 aa; maximum number of motifs, 15.

5.2. Gene structure analysis

The DNA and transcript sequences of *ZmCPA* genes obtained from the maize sequence annotation database MaizeGDB were used to design gene-specific PCR primers with Primer3 (<http://primer3.ut.ee/>). The DNA and cDNA sequences were validated using PCR and RT-PCR with B73 genomic DNA and total RNA as templates. The gene-specific primers shown in Table S2. Alignment of validated DNA and cDNA sequences of each maize *CPA* gene revealed the structure of *ZmCPA* genes. Gene structure display server (GSDS 2.0) was used to display the exon–intron structure, and intron phases [37].

5.3. Plant materials and treatments

The maize B73 inbred line was used in this study. For qRT-PCR, the sterilized seeds were grown in a hydroponic system as described previously [38] with sterile water in a greenhouse at 27/23°C under a 12-h light/12-h dark photoperiod. Four days later, the plants were incubated in 1× Hoagland’s solution (PhytoTechnologies Ltd., Shawnee Mission, KS, USA) until the first leaves were fully expanded. Maize plants were grown with 1 × Hoagland’s solution as the control (CK) and with 1 × Hoagland’s solution containing 100 mM NaCl for the NaCl stress treatment or 100 mM KCl for the KCl stress treatment. These concentrations were maintained until the end of the experiments.

For maize EMS mutants experiment, the sterilized seeds were cultured hydroponically in a greenhouse at 27/23°C under a 12-h light/12-h dark photoperiod as described above. The 1× Hoagland’s solution was exchanged every 2 days. Ten days later, 100 mM KCl and 100 mM NaCl were added. After 4 days of salt stress, phenotypic analyses were performed.

5.4. Expression analysis of *ZmCPA* genes in different tissues

To investigate the spatiotemporal expression patterns of *ZmCPA* genes, the

log2-transformed and RMA-normalized data for *ZmCPA* genes were downloaded from PLEXdb (<http://www.plexdb.org/>) [39]. A heat map was produced using the Lianchuan Bio Cloud Platform (<https://www.lc-bio.cn/overview>).

5.5. RNA isolation and cDNA synthesis

Total RNA was isolated from different tissues of the B73 inbred line, including seedling roots, leaves, 5-cm ears, immature tassels, anthers, pollens, silks, and seeds at 20 days after pollination, using Trizol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol. All RNA was purified using DNase I (Thermo Scientific, Waltham, MA, USA). First-strand cDNA was synthesized from 1 µg total RNA (20 µL reaction volume) using a PrimeScript™ 1st Strand cDNA Synthesis Kit (Takara, Dalian, China) according to the manufacturer's protocol.

5.6. Semi-quantitative reverse transcription PCR

All gene-specific primers were designed as shown in Table S5. Specific primers were used to amplify maize *Actin1* (*GRMZM2G126010*) as an internal control. Reactions were performed with 2 × Taq Master Mix (Vazyme, Nanjing, China) on a Bio-Rad Thermal Cycler (Bio-Rad, Hercules, CA, USA) using the following procedure: 5 min at 94 °C to start; 33 cycles of 30 s at 94 °C, 30 s at 59 °C and 2 min at 72 °C; and a final extension step of 72 °C for 10 min to complete the reaction. *Actin1* was amplified with 29 PCR cycles. Each PCR was performed in triplicate, mixtures without a template were employed as negative controls, and the maize *Actin1* amplicon served as an internal control for each gene investigated.

5.7. Real-time PCR

Real-time PCR was performed using TB Green Premix Ex Taq™ II (Takara, Dalian, China). An ABI 7500 real-time PCR system (Applied Biosystems, Foster City, CA, USA) was used with the following thermal cycling conditions: 95 °C for 5 min, followed by 40 cycles of 95 °C for 15 s, 60 °C for 5 s, and 72 °C for 34 s. The maize *Actin1* gene (*GRMZM2G126010*) was used as an endogenous control to normalize the samples. Based on the cDNA sequences

of *ZmCPA* genes, real-time PCR primers (Table S6) were designed with Primer 3 (<http://primer3.ut.ee/>). The experiment was performed with three technical replicates for each sample. The specificity of each PCR reaction was confirmed by melting curve analysis of the amplicons. The comparative $2^{-\Delta\Delta CT}$ method was used to calculate the relative transcript levels in the samples [40].

5.8. Subcellular localization

The full-length cDNAs of *ZmCHX17* and *ZmNHX5* were amplified using the primers listed in Table S7 and then were introduced into the pM999-EGFP vector to construct the GFP fusion proteins *ZmCHX17*-GFP and *ZmNHX5*-GFP, respectively, with the ClonExpress II One Step Cloning Kit (Vazyme). Constitutive expression of the fused constructs, *ZmCHX17*-GFP and *ZmNHX5*-GFP, was driven by the cauliflower mosaic virus 35S promoter. Maize mesophyll protoplasts were isolated and prepared from etiolated leaves according to established protocols [41]. The plasmids harboring the *ZmCHX17*-GFP and *ZmNHX5*-GFP fusion constructs were co-transfected with the OsSCAMP1-RFP construct into protoplast cells, respectively. OsSCAMP1 is a known rice secretory carrier membrane protein and was used as a membrane protein control [18]. The transformed protoplast cells were cultured at room temperature overnight and then observed under a Leica SP8 confocal microscope (Leica, Wetzlar, Germany).

5.9. Functional expression of *ZmCPAs* in yeast

The coding sequences of *ZmNHX5*, *ZmCHX2*, *ZmCHX3*, *ZmCHX5*, *ZmCHX17*, and *ZmKEA2* were each cloned into the PDR196 vector, and then transformed into the yeast strain AXT3K (*ena1-4::HIS3, nha1::LEU2, nhx1::KanMX*). The transformed yeast cells were cultured overnight at 29 °C in YPDA medium containing 1 mM KCl. Cell cultures were diluted in water to an A_{600} of 0.8. To test cation tolerance, 5- μ L aliquots from yeast cultures or 10-fold serial dilutions were spotted onto AP [42] plates containing 1 mM KCl and supplemented with or without 20 mM NaCl.

436 **List of abbreviations**

437 CPA: monovalent cation proton antiporter; NHX: Na⁺/H⁺ exchanger; KEA: K⁺ efflux
438 antiporter; CHX: cation/H⁺ exchanger; Na⁺: sodium; Cl⁻: chloride; TM: transmembrane; K⁺:
439 potassium; H⁺: proton; MW: molecular weight; pI: isoelectric point; RT-PCR: reverse
440 transcription polymerase chain reaction; DNA: deoxyribonucleic acid; cDNA:
441 complementary DNA; NJ: neighbor-joining; aa: amino acid; kDa: kilodalton; TF:
442 transcription factor; semi qRT-PCR: semi-quantitative reverse transcription polymerase chain
443 reaction; AP: Arg phosphate; EMS: ethyl methanesulfonate; MEMD: Maize EMS-induced
444 Mutant Database; QTL: quantitative trait locus; GFP: green fluorescent protein; RFP: red
445 fluorescent protein.

447 **Authors' contributions**

448 Y.T., Y.Z., (Yanxin Zhao) and J.Z. conceived the experiment. M.K, M.L, J.L, Y.Z.
449 (Yunxia Zhang), Z.F, W.S, R.Z, R.W., and Y.W. performed bioinformatic analysis and data
450 acquisition. M.K, M.L, and Y.Z (Yanxin Zhao) analyzed the data and wrote the manuscript.
451 All authors revised and approved the final manuscript, and agreed to be accountable for this
452 work.

454 **Funding**

455 This research was financially supported by the Beijing Municipal Natural Science
456 Foundation (6204041), the Construction of Collaborative Innovation Center of Beijing
457 Academy of Agricultural and Forestry Sciences (Collaborative Innovation Center of Crop
458 Phenomics, KJCX201917), Youth Research Fund of Beijing Academy of Agriculture and
459 Forestry Sciences (QNJJ202028), and the Beijing Scholars Program (BSP041). The funding
460 body does not play roles in the design of the study and collection, analysis, and interpretation
461 of data and in writing the manuscript.

463 **Declaration of Competing Interest**

464 The authors declare that they have no competing interests.

465 Acknowledgements

466 We thank Prof. Liwen Jiang (Chinese University of Hong Kong) for kindly providing
467 pM999-EGFP and membrane protein marker (OsSCAMP1-RFP) plasmids, Prof. Xiaoduo Lu
468 (Qilu Normal University) for identifying and providing maize EMS mutants *zmkea2* and
469 *zmnhx5*, and Profs. Yi Wang and Zhenxian Zhang (China Agricultural University) for kindly
470 sharing yeast mutant AXT3K and yeast expression vector pDR196.

471

472 References

- 473 [1] Zhu JK. Plant salt tolerance. Trends Plant Sci. 2001, 6:66–71.
- 474 [2] Gupta B, Huang B. Mechanism of salinity tolerance in plants: physiological, biochemical,
475 and molecular characterization. Int J Genom. 2014, 1:701596.
- 476 [3] Farooq M, Hussain M, Wakeel A, Siddique KHM. Salt stress in maize: effects, resistance
477 mechanisms, and management. A review. Agron Sustain Dev. 2015(35), 35:461–481.
- 478 [4] Chanroj S, Wang G, Venema K, Zhang MW, Delwiche CF, Sze H. Conserved and
479 diversified gene families of monovalent Cation/H⁺ Antiporters from algae to flowering
480 plants. Front Plant Sci. 2012, 3:25.
- 481 [5] Tian FX, Chang E, Li Y, Sun P, Hu JJ, Zhang J. Expression and integrated network
482 analyses revealed functional divergence of NHX-type Na⁺/H⁺ exchanger genes in poplar.
483 Sci Rep. 2017, 7:2607.
- 484 [6] Fukuda A, Nakamura A, Hara N, Toki S, Tanaka Y. Molecular and functional analyses of
485 rice NHX-type Na⁺/H⁺ antiporter genes. Planta. 2011. 233(1):175–188.
- 486 [7] Sellamuthu G, Jegadeeson V, Sajeevan RS, Rajakani R, Parthasarathy P, Raju K, Shabala
487 L, Chen ZH, Zhou MX, Sowdhamini R, Shabala S, Venkataraman G. Distinct
488 evolutionary origins of intron retention splicing events in *NHX1* antiporter transcripts
489 relate to sequence specific distinctions in *Oryza* species. Front Plant Sci. 2020, 11:267.
- 490 [8] Huanca-Mamani W, Ortiz MV, Cárdenas-Ninasivincha S, Acosta-Garcia G, Bastías E.
491 Gene expression analysis in response to combined salt and boron (B) stresses in a
492 tolerant maize landrace. Plant Omics. 2018, 11(2):80–88.
- 493 [9] Bosnic P, Bosnic D, Jasnic J, Nikolic M. Silicon mediates sodium transport and
494 partitioning in maize under moderate salt stress. Environ Exp Bot. 2018, 155:681–687.
- 495 [10] Pires IS, Negrão S, Pentony MM, Abreu IA, Oliveira MM, Purugganan MD. Different
496 evolutionary histories of two cation/proton exchanger gene families in plants. BMC Plant
497 Biol. 2013, 13(1): 97.
- 498 [11] Saier MH, Reddy VS, Tamang DG, Västermark A. The transporter classification database.
499 Nucleic Acids Res. 2014, 42(D1):D251–D258.
- 500 [12] Ye CY, Yang X, Xia X, Yin W. Comparative analysis of cation/proton antiporter
501 superfamily in plants. Gene. 2013, 521(2):245–251.
- 502 [13] Zheng S, Pan T, Fan L, Qiu Q. A novel *AtKEA* gene family, homolog of bacterial K⁺/H⁺
503 antiporters, plays potential roles in K⁺ homeostasis and osmotic adjustment in
504 *Arabidopsis*. PLoS One. 2013, 8(11):e81463.

- 505 [14] Kunz HH, Gierth M, Herdean A, Satoh-Cruz M, Kramer DM, Spetea C, Schroeder JJ.
506 Plastidial transporters KEA1, -2, and -3 are essential for chloroplast osmoregulation,
507 integrity, and pH regulation in *Arabidopsis*. Proc Natl Acad Sci USA. 2014,
508 111(20):7480–7485.
- 509 [15] Aranda-Sicilia MN, Aboukila A, Armbruster U, Cagnac O, Schumann T, Kunz HH,
510 Jahns P, Rodríguez-Rosales MP, Sze H, Venema K. Envelope K⁺/H⁺ antiporters AtKEA1
511 and AtKEA2 function in plastid development. Plant Physiol. 2016, 172(1):441–449.
- 512 [16] Isayenkov SV, Dabravolski SA, Pan T, Shabala S. Phylogenetic diversity and
513 physiological roles of plant monovalent cation/H⁺ antiporters. Front Plant Sci. 2020,
514 11:573564.
- 515 [17] Chen Y, Ma J, Miller AJ, Luo B, Wang M, Zhu Z, Ouwerkerk PBF. OsCHX14 is
516 involved in the K⁺ homeostasis in rice (*Oryza sativa*) flowers. Plant Cell Physiol. 2016,
517 57(7):1530–1543.
- 518 [18] Lam SK, Siu CL, Hillmer S, Jang S, An G, Robinson DG, Jiang L. Rice SCAMP1
519 defines clathrin-coated, *trans*-golgi-located tubular-vesicular structures as an early
520 endosome in tobacco BY-2 cells. Plant Cell. 2007, 19(1):296–319.
- 521 [19] Sekhon RS, Lin H, Childs KL Hansey CN, Buell CR, de Leon N, Kaeppler SM.
522 Genome-wide atlas of transcription during maize development. Plant J. 2011,
523 66(4):553–563.
- 524 [20] Quintero FJ, Blatt MR, Pardo JM. Functional conservation between yeast and plant
525 endosomal Na⁺/H⁺ antiporters. FEBS Lett. 2000, 471(2-3):224–228.
- 526 [21] Lu X, Liu J, Ren W, Yang Q, Chai Z, Chen R, Wang L, Zhao J, Lang Z, Wang H, Fan Y,
527 Zhao J, Zhang C. Gene-indexed mutations in maize. Mol Plant. 2018, 11(3):496–504.
- 528 [22] Apse MP, Sottosanto JB, Blumwald E. Vacuolar cation/H⁺ exchange, ion homeostasis,
529 and leaf development are altered in a T-DNA insertional mutant of *AtNHX1*, the
530 *Arabidopsis* vacuolar Na⁺/H⁺ antiporter. Plant J. 2003;36:229–239.
- 531 [23] Bassil E, Ohto M, Esumi T, Tajima H, Zhu Z, Cagnac O, Belmonte M, Peleg Z,
532 Yamaguchi T, Blumwald E. The *Arabidopsis* intracellular Na⁺/H⁺ antiporters NHX5 and
533 NHX6 are endosome associated and necessary for plant growth and development. Plant
534 Cell. 2011, 23(1):224–239.
- 535 [24] Sharma H, Taneja M, Upadhyay SK. Identification, characterization and expression
536 profiling of cation-proton antiporter superfamily in *Triticum aestivum* L. and functional
537 analysis of *TaNHX4-B*. Genomics. 2020, 112(1):356–370.
- 538 [25] Zhang WW, Meng JJ, Xing JY, Yang S, Guo F, Li XG, Wan SB. The K⁺/H⁺ antiporter
539 *AhNHX1* improved tobacco tolerance to NaCl stress by enhancing K⁺ retention. J Plant
540 Biol. 2017, 60(3):259–267.
- 541 [26] Jia Q, Zheng C, Sun S, Amjad H, Liang K, Lin W. The role of plant cation/proton
542 antiporter gene family in salt tolerance. Biol Plantarum. 2018, 62(4):617–629.
- 543 [27] Luo M, Zhao Y, Zhang R, Xing J, Duan M, Li J, Wang N, Wang W, Zhang S, Chen Z,
544 Zhang H, Shi Z, Song W, Zhao J. Mapping of a major QTL for salt tolerance of mature
545 field-grown maize plants based on SNP markers. BMC Plant Biol. 2017, 17(1):140.
- 546 [28] Luo M, Zhang Y, Chen K, Kong M, Song W, Lu B, Shi Y, Zhao Y, Zhao J. Mapping of
547 quantitative trait loci for seedling salt tolerance in maize. Mol Breeding. 2019, 39:64.
- 548 [29] Zörb C, Noll A, Karl S, Leib K, Yan F, Schubert S. Molecular characterization of Na⁺/H⁺

antiporters (ZmNHX) of maize (*Zea mays* L.) and their expression under salt stress. J Plant Physiol. 2005, 162(1):55–66.

[30] Zhou H, Qi K, Liu X, Yin H, Wang P, Chen J, Wu J, Zhang S. Genome-wide identification and comparative analysis of the cation proton antiporters family in pear and four other Rosaceae species. Mol Genet Genomics. 2016, 291(4):1727–1742.

[31] Zhao J, Li P, Motes CM, Park S, Hirschi KD, CHX14 is a plasma membrane K-efflux transporter that regulates K⁺ redistribution in *Arabidopsis thaliana*. Plant Cell Environ. 2015, 38(11):2223–2238.

[32] Luo MJ, Shi YX, Yang Y, Zhao YX, Zhang YX, Shi YM, Kong MS, Li CH, Feng Z, Fan YL, Xu L, Xi SL, Lu BS, Zhao JR. Sequence polymorphism of the *waxy* gene in waxy maize accessions and characterization of a new *waxy* allele. Sci Rep. 2020, 10:15851.

[33] Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. BMC Bioinformatics. 2004, 5:113.

[34] Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: molecular evolutionary genetics analysis version 6.0. Mol Biol Evol. 2013, 30(12):2725–2729.

[35] Small I, Peeters N, Legeai F, Lurin C. Predotar: A tool for rapidly screening proteomes for N-terminal targeting sequences. Proteomics. 2004, 4(6):1581–1590.

[36] Emanuelsson O, Brunak S, von Heijne G, Nielsen H. Locating proteins in the cell using TargetP, SignalP and related tools. Nat Protoc. 2007, 2(4):953–971.

[37] Hu B, Jin J, Guo AY, Zhang H, Luo J, Gao G. GSDS 2.0: an upgraded gene feature visualization server. Bioinformatic. 2015, 31:1296–1297.

[38] Luo M, Zhao Y, Wang Y, Shi Z, Zhang P, Zhang Y, Song W, Zhao J. Comparative proteomics of contrasting maize genotypes provides insights into salt-stress tolerance mechanisms. J Proteome Res. 2018, 17(1):141–153.

[39] Dash S, Van Hemert J, Hong L, Wise RP, Dickerson JA. PLEXdb: gene expression resources for plants and plant pathogens. Nucleic Acids Res. 2010, 40(Database issue):D1194–1201.

[40] Thomas S, Kenneth D, Livak J. Analyzing real-time PCR data by comparative CT method. Nat Protoc. 2008, 3(6):1101–1108.

[41] Yoo SD, Cho YH, Sheen J. *Arabidopsis* mesophyll protoplasts: a versatile cell system for transient gene expression analysis. Nat Protoc. 2007, 2(7):1565–1572.

[42] Rodríguez-Navarro A, Ramos J. Dual system for potassium transport in *Saccharomyces cerevisiae*. J Bacteriol. 1984, 159(3):940–945.

589 Figure legends

590 Fig. 1. Gene structures of *ZmCPA* genes.

591 Neighbor-joining tree of maize *CPA* genes at left is based on Fig. 3. Exon–intron organization
592 of *ZmCPA* genes (right) as determined by aligning DNA and cDNA sequences of *CPA* genes
593 at GSDS 2.0 server (<http://gsds.gao-lab.org/index.php>). Introns and exons are shown as thin
594 lines and yellow boxes, respectively. Numbers 0, 1, and 2 represent different intron phases.
595 Blue boxes at 5' and 3' ends represent untranslated regions (UTRs).

596

597 Fig. 2. Phylogenetic tree of *CPA* genes in maize, *Arabidopsis*, and rice.

598 Neighbor-joining tree constructed using MEGA 6.0 based on alignment of 103 CPA proteins
599 including 35 ZmCPAs, 42 AtCPAs, and 26 OsCPAs. NHX, KEA, and CHX subfamilies are
600 divided into three (N1, N2 and N3), three (C1, C2, and C3) and two (K1 and K2) groups,
601 respectively.

602

603 Fig. 3. Phylogenetic tree of *ZmCPA* genes.

604 All 35 ZmCPA proteins were aligned using MUSCLE v3.8.31 and alignment was used to
605 construct neighbor-joining tree with MEGA 6.0. *ZmCPA* genes are divided into three
606 subfamilies; *ZmNHXs*, *ZmKEAs*, and *ZmCHXs*.

607

608 Fig. 4. Subcellular localization of *ZmCHX17* and *ZmNHX5* proteins.

609 *ZmCHX17*-GFP (upper panel) and *ZmNHX5*-GFP (lower panel) fusion proteins transiently
610 expressed in maize protoplast cells co-transformed with OsSCAMP1-RFP fusion construct
611 (membrane protein marker). GFP and RFP signals were detected by confocal fluorescent
612 microscopy (Leica SP8). Scale bar = 5 μm.

613

614 Fig. 5. Gene expression profiles of *ZmCPA* genes in different tissues.

615 Normalized microarray expression data of *ZmCPA* genes were downloaded from PLEXdb
616 (<http://www.plexdb.org/>). Cluster analysis was performed and heatmap was generated using
617 tools in the Lianchuan BioCloud Platform (<https://www.lc-bio.cn/overview>).

618

619 Fig. 6. Expression analysis of *ZmCPA* genes in eight tissues of B73 inbred line by semi 620 qRT-PCR.

621 Total RNA was isolated from eight tissues: seedling roots, seedling leaves, 5-cm immature
622 ears, tassels, anthers, pollen, silk and whole seeds (20 days after pollination, DAP), and then
623 used for semi qRT-PCR analyses of *ZmCPA* genes. Maize *Actin1* gene was the internal
624 control.

625

626 Fig. 7. Expression analysis of *ZmCPA* genes under salt stress conditions by qRT-PCR.

627 Expression analysis of *ZmCPA* genes under 100 mM NaCl or 100 mM KCl as determined by
628 quantitative reverse transcription-polymerase chain reaction (qRT-PCR). Y-axis represents
629 relative transcript levels of *ZmCPA* genes compared with that of *actin1*. X-axis represents
630 different time points after salt treatment in each group. Error bars represent standard
631 deviations for three replicates.

Fig. 8. *ZmCPA* genes facilitate growth of yeast mutant strain AXT3K under salt stress.

Full-length CDS sequences of *ZmNHX5*, *ZmCHX2*, *ZmCHX3*, *ZmCHX5*, *ZmCHX17*, *ZmKEA2* were each cloned into PDR196 vector, then transformed into yeast strain AXT3K (*ena1-4::HIS3*, *nha1::LEU2*, *nhx1::KanMX*). Yeast transformant cells were cultured overnight, then were diluted with water to A_{600} of 0.8. Aliquots (5 μ L) of 10-fold serial dilutions were spotted on AP plates supplemented with 20 mM NaCl at pH 7.5 (right panel) and normal AP plates as the control (left panel). Strains were grown at 29°C for 5 days and photographed.

Fig. 9. Growth performance of *zmnhx5* and *zmkea2* mutant seedlings under salt stress.

Point mutation sites of EMS mutants *zmnhx5* and *zmkea2* were identified by PCR sequencing. Stop codons induced by EMS resulted in truncated proteins (a, e). Growth status (b, f), seedling length (c, g) and dry weight (d, h) of mutant seedlings shoot under 100 mM NaCl or under 100 mM KCl treatment for 4 days ($n = 6$, mean $\pm$ SEM). *, Significant difference (Student's t-test; $P < 0.05$). NS, non significant.

Table 1. Information of maize CPA genes identified in this study.

Gene Name	Gene Model	Chromosomal Location	CDS (bp)	Protein Length (aa)
<i>ZmNHX1</i>	<i>Zm00001d028330</i>	Chr1: 31041489-31049951	1599	532
<i>ZmNHX7</i>	<i>Zm00001d031232</i>	Chr1: 182925112-182945175	3411	1136
<i>ZmNHX11</i>	<i>Zm00001d003728</i>	Chr2: 56632216-56643296	2367	788
<i>ZmNHX6</i>	<i>Zm00001d048732</i>	Chr4: 4469601-4475393	1641	546
<i>ZmNHX8</i>	<i>Zm00001d019978</i>	Chr7: 81486383-81509317	1611	536
<i>ZmNHX3</i>	<i>Zm00001d020892</i>	Chr7: 135679272-135687679	1491	496
<i>ZmNHX10</i>	<i>Zm00001d021844</i>	Chr7: 164344371-164361391	2820	939
<i>ZmNHX5</i>	<i>Zm00001d022504</i>	Chr7: 179047417-179052457	1620	539
<i>ZmNHX4</i>	<i>Zm00001d045883</i>	Chr9: 45473833-45477106	1638	545
<i>ZmNHX12</i>	<i>Zm00001d048459</i>	Chr9: 156705048-156709982	1440	479
<i>ZmNHX2</i>	<i>Zm00001d024832</i>	Chr10: 90134129-90141743	1680	559
<i>ZmNHX13</i>	<i>Zm00001d025052</i>	Chr10: 101802712-101807654	1563	520
<i>ZmNHX9</i>	<i>Zm00001d026118</i>	Chr10: 138832710-138862665	2115	704
<i>ZmCHX11</i>	<i>Zm00001d031077</i>	Chr1: 176192733-176196049	2388	795
<i>ZmCHX15</i>	<i>Zm00001d031078</i>	Chr1: 176200721-176205899	2061	686
<i>ZmCHX7</i>	<i>Zm00001d005032</i>	Chr2: 155039785-155042337	2553	850
<i>ZmCHX4</i>	<i>Zm00001d041198</i>	Chr3: 105028259-105030854	2529	842
<i>ZmCHX9</i>	<i>Zm00001d044623</i>	Chr3: 233503244-233505658	2415	804
<i>ZmCHX8</i>	<i>Zm00001d049663</i>	Chr4: 39074035-39076527	2493	830
<i>ZmCHX6</i>	<i>Zm00001d050509</i>	Chr4: 94925858-94928987	2232	743
<i>ZmCHX2</i>	<i>Zm00001d053237</i>	Chr4: 221575337-221578465	2478	825
<i>ZmCHX1</i>	<i>Zm00001d017805</i>	Chr5: 207302222-207308368	2484	827
<i>ZmCHX12</i>	<i>Zm00001d035631</i>	Chr6: 37793506-37814621	2571	856
<i>ZmCHX16</i>	<i>Zm00001d038517</i>	Chr6: 159098609-159101687	2598	865
<i>ZmCHX3</i>	<i>Zm00001d021461</i>	Chr7: 152741608-152744549	2424	807
<i>ZmCHX14</i>	<i>Zm00001d009889</i>	Chr8: 87464523-87467367	2391	796
<i>ZmCHX5</i>	<i>Zm00001d010601</i>	Chr8: 121526743-121529490	2475	824
<i>ZmCHX10</i>	<i>Zm00001d010629</i>	Chr8: 122278283-122281316	2874	957
<i>ZmCHX17</i>	<i>Zm00001d012521</i>	Chr8: 176021482-176024817	2628	875
<i>ZmKEA5</i>	<i>Zm00001d027466</i>	Chr1: 5893819-5906194	1791	596
<i>ZmKEA4</i>	<i>Zm00001d001788</i>	Chr2: 935639-948444	3480	1159
<i>ZmKEA3</i>	<i>Zm00001d041308</i>	Chr3: 111434917-111440097	2397	798
<i>ZmKEA1</i>	<i>Zm00001d036981</i>	Chr6: 108202268-108211871	1848	615
<i>ZmKEA6</i>	<i>Zm00001d046231</i>	Chr9: 73621310-73635421	1872	623
<i>ZmKEA2</i>	<i>Zm00001d026645</i>	Chr10: 149326566-149340249	3315	1104

[illegible]

AP

AP+20mM NaCl

AXT3K

PDR196

ZmNHX5

ZmCHX2

ZmCHX3

ZmCHX5

ZmCHX17

ZmKEA2