To the canopy and beyond: Air samples reveal wind dispersal as a driver of ubiquitous protistan pathogen assembly in tree canopies

Robin-Tobias Jauss^{a#[co]} & Anne Nowack^{a[co]}, Susanne Walden^b, Ronny Wolf^a, Stefan Schaffer^a, Barbara Schellbach^c, Michael Bonkowski^b, Martin Schlegel^{a,d#}

^a University of Leipzig, Institute of Biology, Molecular Evolution & Animal Systematics, Talstraße 33, 04103 Leipzig

^b University of Cologne, Institute of Zoology, Terrestrial Ecology, Zùlpicher Straße 47b, 50674 Köln

^c Max Planck Institute for Evolutionary Anthropology, Department of Evolutionary Genetics, Deutscher Platz 6, 04103 Leipzig

^d German Centre for Integrative Biodiversity Research (iDiv) Halle Jena Leipzig, Deutscher Platz 5e, 04103 Leipzig

[co] These authors contributed equally and should be considered joint first author. Author order was determined in order of decreasing seniority

To whom correspondence should be addressed: jauss@uni-leipzig.de, schlegel@uni-leipzig.de

Keywords: Airborne Microorganisms, Protists, Forest Ecosystems, Canopies, Pathogens

Abstract

We analyzed air dispersal of the protistan phyla Cercozoa and Oomycota with an air sampler near the ground (~2 m) and in tree crowns (~25 m) of three tree species (oak, linden and ash) in a temperate floodplain forest in March (before leafing) and May (after leaf unfolding) with a cultivation-independent high throughput metabarcoding approach. Both, Cercozoa and Oomycota, contain important pathogens of forest trees and other vegetation. We found a high diversity of Cercozoa and Oomycota in air samples with 122 and 81 OTUs, respectively. Especially oomycetes showed a high temporal variation in beta diversity between both sampling dates. Differences in community composition between air samples

in tree canopies and close to the ground were however negligible, and also tree species identity did not affect communities in air samples, indicating that the distribution of protistan propagules through the air was not spatially restricted in the forest ecosystem. OTUs of plant pathogens, whose host species that did not occur in the forest, demonstrate wind dispersal of propagules from outside the forest biome. Overall, our results lead to a better understanding of the stochastic processes of wind dispersal of protists and protistan pathogens, a prerequisite to understand the mechanisms of their community assembly in forest ecosystems.

Importance

Wind dispersal has been shown to play a crucial role in protistan community assembly. The protistan taxa Cercozoa and Oomycota contain important plant parasites with a major ecologic and economic impact. However, comprehensive assessments of cercozoan and oomycete diversity in forest air samples were lacking. Using a cultivation-independent high throughput metabarcoding approach, we analyzed cercozoan and oomycete air dispersal in forest floors and the canopy region – a potential filter for microbial propagules. Our study provides insights into the diversity and community assembly of protists within the air, contributing to a better understanding which factors drive the distribution of plant pathogens within forest ecosystems.

1. Introduction

The air is an effective means of long-distance propagation for a wide range of microbial organisms (Foissner & Hawksworth, 2009; Pepper & Dowd, 2009). The phyllosphere – and especially the crowns of trees – are the largest biological interface between the soil and the atmosphere (Ozanne et al., 2003; Ellwood & Foster, 2004), which therefore may act as a huge natural filter for airborne microbial propagules, including unicellular Eukaryotes (Protists). Within the paraphyletic taxon of protists, the group of Cercozoa (Rhizaria) are highly diverse in morphology and physiology and show a high functional and ecological

variety (Bass et al., 2009; Harder et al., 2016). They dominate terrestrial habitats (Urich et al., 2008; Voss et al., 2019) and harbor important plant pathogens, such as the Endomyxa, which have recently been elevated from the Cercozoa into a separate phylum (Cavalier-Smith et al., 2018). Another protistan taxon, the Oomycetes (Stramenopiles), contain important parasites of forest trees, and many lineages produce caducous sporangia for dissemination (Goheen & Frankel, 2009; Robideau et al., 2011; Lang-Yona et al., 2018). With almost 800 described species, Oomycota are reported to have a broad distribution and a wide variety of ecological roles (Robideau et al., 2011; Thines, 2014; Judelson, 2017). Further, it is one of the eukaryotic groups with a great impact on ecosystems, as well as on economics and human health: the most famous species is *Phytophthora infestans*, which causes the potato blight. In the 1840s it led to the great famine in Ireland followed by massive emigration (Lara & Belbahri, 2011; Robideau et al., 2011).

Protists can be passively disseminated over long distances by viable propagules, mostly as resting stages (cysts), while some groups, especially pathogens with more complex life cycles, also form sporangia for dispersal (Cowling, 1994; Kageyama & Asano, 2009). Cysts are formed under unfavorable conditions, e.g. due to dryness, lack of food, or microbial antibiotics (Petz & Foissner, 1988; Adl & Gupta, 2006; Jousset et al., 2006), and it has been assumed that the cyst bank plays an important role for the resilience of protists and their functions in terrestrial environments (Geisen et al., 2017). Viable protist cysts can be retrieved from soils even after decades (Moon-van der Staay et al., 2006; Kageyama & Asano, 2009), leading to the long-standing question of how cosmopolitan protists are (Finlay, 2002; Fenchel & Finlay, 2004; Foissner, 2009).

Finlay et al. (2001) proposed that the spatial distribution of protistan propagules is influenced by several randomizing factors, such as soil particles dispersed by wind, convective transport, percolating rainwater, fog or animals. Rogerson and Detwiler (1999) determined that on average $0.25 \text{ cysts m}^{-3}$ are contained in the air depending on wind speed

82 and time since last precipitation. Using a molecular approach, Genitsaris et al. (2014) came
83 to generally similar conclusions, while they further detected operational taxonomic units
84 (OTUs) with constant presence as well as OTUs exhibiting seasonal variation. High humidity
85 increases the chance of survival of transported microbes and promotes their deposition
86 (Fuzzi et al., 1997; Evans et al., 2019) and airborne microorganisms can be transported in
87 fog droplets by atmospheric turbulence over long distances (Fuzzi et al., 1997; Amato et al.,
88 2005).

89 Recently, Jauss et al. (2020) confirmed a ubiquitous distribution of Cercozoa and Oomycota
90 in a floodplain forest, despite strong differences in community composition of different
91 microhabitats related to differences in the relative abundance of taxa. This led to the
92 conclusion that within forest ecosystems both cercozoans and oomycetes can colonize most
93 habitats, in which they then however do not perform similarly well due to habitat filtering.
94 One reason for this ubiquitous presence of these protists could be wind dispersal.

95 Here, we studied the air dispersal of Cercozoa, Endomyxa and Oomycota by a cultivation-
96 independent high throughput metabarcoding approach to analyze protistan diversity in the
97 air surrounding tree canopies and near the ground of a temperate floodplain forest, to gain
98 a deeper insight into the mechanisms how protists and their pathogenic lineages are
99 distributed in the environment. These examinations tackled three hypotheses: (1) Wind
100 dispersal explains the ubiquitous presence of these protists in the floodplain forest. (2) There
101 are differences in the distribution in the vertical plane as a strong discrepancy between
102 canopy and ground habitats was previously described. (3) Temporal variation of wind
103 dispersed propagules further drives the community and pathogen assembly in forest
104 ecosystems.

2. Material and Methods

2.1. Sampling and DNA extraction

Air samples were taken in a temperate deciduous floodplain forest in the northwest of the city of Leipzig, Germany (51.3657 N, 12.3094 E) with a MicroBio MB2 Bioaerosol Sampler (Cantium Scientific, Dartford, UK) containing 1% agar plates. The samples were collected under defined conditions drawing 100 l/min of air for ten minutes in two strata: (1) near the ground (~2m) and (2) in ~25m height in tree canopies with the help of the Leipzig Canopy Crane (LCC) facility. Two samplings were carried out – one in March and one in May 2019. For each sampling, three tree species with three replicates each were chosen (*Quercus robur*, *Tilia cordata* and *Fraxinus excelsior*). As non-arboreal control, samples were also taken on the crane tracks near the ground and at canopy height. Two plates per stratum and of each replicated tree species were collected, yielding 40 plates per sampling. After air sampling, the agar plates were taken out of the instrument, sealed with parafilm to prevent contaminations and frozen until the DNA was extracted with the DNeasy PowerSoil® Kit according to the instructions supplied by the manufacturer. Weather conditions were tracked with a WebVIS data logger attached to the crane (Umweltanalytische Produkte GmbH, Ibbenbüren, Germany) (Table 1).

Table 1: Weather conditions at the sampling days in March and May 2019.

	March	May
Average temperature [°C]	6.5	14.3
Average humidity [%]	59.4	75.0
Average wind speed [m/s]	4.01	3.23
Last precipitation event	Immediately before sampling	2 days before sampling

2.2. PCR amplification and sequencing

DNA was amplified in duplicates with tagged oomycete- and cercozoan-specific primers (Fiore-Donno & Bonkowski, 2020; Jauss et al., 2020; Fiore-Donno et al., 2020) (Supplementary Tables 1-2). PCR-products were purified following the directions of the NucleoSpin® PCR clean-up protocol. Afterwards, DNA concentrations were measured with the Qubit™4 fluorometer in combination with the Qubit™ dsDNA HS Assay Kit. For consecutive Illumina MiSeq Sequencing, a library was prepared following the Meyer and Kircher (2010) protocol. DNA concentrations were checked repeatedly before and after Illumina sequencing by utilization of DNA chips analyzed with the Agilent 2100 Bioanalyzer. Between the steps of library preparation, reaction clean-up was performed with the AMPure XP System using carboxyl coated magnetic beads (SPRI beads). Subsequent steps and the Illumina MiSeq sequencing itself were performed by the sequencing team of the Max Planck Institute for Evolutionary Anthropology in Leipzig, Germany.

2.3. Sequence processing and statistical analyses

Bioinformatic and statistical analyses followed the pipeline described in Jauss et al. (2020). Briefly, resulting reads were merged and clustered into operational taxonomic units (OTUs) using a custom pipeline utilizing cutadapt v1.18 (Martin, 2011), Swarm v2.2.2 (Mahé et al., 2015) and VSEARCH v2.10.3 (Rognes et al., 2016). OTUs were then annotated using NCBI's non-redundant nucleotide database and the Protist Ribosomal Reference Database (Guillou et al., 2013) for oomycete and cercozoan OTUs, respectively (Supplementary Tables 3-4). OTUs resembling non-oomycete or non-cercozoan sequences were excluded. Samples with less than 5 OTUs or with a sequencing depth lower than 20617 reads (Oomycota) and 16922 reads (Cercozoa) were omitted. Statistical analyses of alpha and beta diversity and final visualizations were performed in R v3.5.3 (R Core Team, 2019) with the packages vegan (Oksanen et al., 2019), ggplot2 (Wickham, 2016) and ggraph (Pedersen, 2020).

3. Results

3.1. Amplification, sequencing and bioinformatic pipeline

After DNA isolation, all oomycete samples were amplified successfully whereas nine out of 20 cercozoan samples had to be excluded due to the failure of successful amplification of duplicates. Further, samples containing less than 1ng/μl DNA were excluded from subsequent processing, as well as samples with a low sequencing depth (see 2.3), yielding 9 cercozoan samples from March and 4 from May, as well as 13 oomycete samples from March and 20 from May (Supplementary Table 5). Of cercozoan sequences, 94.4% could be merged with a mean length of 370 ± 35 bp resulting in 122 OTUs in total. Of oomycete sequences, 92.6% of derived 300 bp long paired-end sequences could be merged and the mean fragment length accounted for 285 ± 38 bp, which were finally clustered into 81 OTUs.

3.2. Alpha diversity

Neither species richness nor Shannon-diversity nor evenness of Cercozoa or Oomycota differed between tree species, ground and canopy or non-arboreal controls, although variation of canopy samples was much lower than of ground samples in Cercozoa (e.g. species richness $CV_{\text{Ground}} = 42.3\%$, $CV_{\text{Canopy}} = 5.6\%$). However, Shannon-diversity and evenness of both protistan groups and species richness of oomycetes were higher in May than in March, indicating that the tree foliage in May did not restrict protistan distribution (Figure 1).

Figure 1: Boxplot of alpha diversity indices of cercozoan (A) and oomycete (B) samples. Pairwise comparisons of March and May samples and canopy and ground samples, respectively are shown. Significance was tested with Wilcoxon Sign test and is indicated by asterisks (ns = $p > 0.05$, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, **** = $p < 0.0001$).

3.3. Beta diversity

For Cercozoa, β -diversity of air samples did not differ between tree species (permANOVA $R^2=0.165$, $p=0.514$), ground vs. canopy stratum (permANOVA $R^2=0.093$, $p=0.296$) nor sampling season (permANOVA $R^2=0.101$, $p=0.168$). However, variation of β -diversity was much lower in May compared to March, and lower in canopy samples compared to ground samples (Figure 2A). Oomycete communities differed between sampling seasons (permANOVA $R^2=0.170$, $p=0.001$), but not between tree species (permANOVA $R^2=0.080$, $p=0.719$) or the strata ground and canopy (permANOVA $R^2=0.037$, $p=0.259$) (Figure 2B).

Figure 2: Non-metric multidimensional scaling (NMDS) plot of cercozoan (A) and oomycete (B) samples. Canopy and ground samples show a large overlap, while in oomycetes the March and May samples show a strong separation.

3.4. Taxonomic diversity

Cercozoan OTUs were dominated by the orders Cryomonadida and Glissomonadida, whereas the least abundant ones were Marimonadida and an unspecified order named Cercozoa_XX, comprising undescribed cercozoan lineages (Figure 3A). We detected no OTUs assigned to the plant parasitic group of Endomyxa. Oomycete OTUs were almost exclusively dominated by Peronosporales (Figure 3B), with only few members of the Pythiales, and the Albuginales being the least abundant order.

Figure 3: Taxonomic annotation of cercozoan (A) and oomycete (B) OTUs. Labels give the detected orders and the ten most abundant species with their corresponding genus.

The number of shared OTUs indicated a temporal variation in air dispersal of both protistan taxa (Figure 4), and dispersal of cercozoan OTUs varied also spatially between canopy and ground at the incidence level.

Figure 4: Venn diagram giving the number and proportion of shared OTUs between March and May samples and Canopy and Ground samples, respectively, for Cercozoa (A) and Oomycota (B).

190

191 Partitioning of the taxonomy into the two sampling seasons revealed similar patterns (Figure
192 5), yet, the cercozoan order Euglyphida was exclusively present in March samples, and
193 oomycete Pythiales showed a higher abundance in March than in May.

Figure 5: Sankey distribution diagram of cercozoan (A) and oomycete (B) orders in March and May samples. Orders represented by less than 1% of all reads were removed from the visualisation for the sake of clarity.

4. Discussion

In a recent study, Jauss et al. (2020) quantified the diversity of Oomycota and Cercozoa in canopy microhabitats and in litter and soil on the ground of the same floodplain forest. This allows for a direct comparison of the total diversity of these protists in the forest stand with the potential diversity of taxa distributed by air during two time points. We detected 122 and 81 OTUs of Cercozoa and Oomycota in air samples, respectively, which corresponds to 22 and 24% of the former reported total diversity of these protistan phyla. The high temporal variation, also reflected by the number of shared OTUs (Figure 4), suggests protistan distribution to be not restricted by dispersal-limitation, but rather indicates a continuous propagule rain of potentially invasive species and their accumulating resting stages occupying vacant niches. The vertical distribution of protists in air samples was rather homogeneous and did not differ between tree canopy and ground. In contrast, Jauss et al.

(2020) found clear spatial patterns of oomycetes and cercozoans in tree canopies compared to the forest floor, suggesting that only part of the wind-borne propagule rain finds suitable conditions for survival in tree crowns due to habitat filtering. The temporal variation could be either related to temporal variations in the activity and distribution of protists, or more likely due to a dependency on the weather during the sampling. In March, the conditions were less favorable, with comparatively low temperatures and humidity with immediate previous precipitation events that have to be taken into account (Table 1). The remaining ground moisture might have prevented the lofting of protists through wind currents, while in May the atmospheric conditions were more preferable with higher average temperatures and a higher humidity. The conditions in May probably favored the lofting of protists into the atmosphere and their long-distance dispersal, leading to a higher protistan diversity and OTU richness (Figure 1), even though the wind speed was slower compared to March (Table 1). As wind speed was determined to be an important factor governing the species richness of microorganisms in air samples (Rogerson & Detwiler, 1999; Genitsaris et al., 2014), faster wind speeds in May probably could have revealed more protists. This suggests not a single factor, but rather the interplay between atmospheric conditions driving the species richness and community assembly in the air, while our samples possibly only represent the lower counts of what can be dispersed by air.

Wind dispersal is an important means for the distribution of microbial plant pathogens, and oomycetes are no exception (Fawke et al., 2015; Lang-Yona et al., 2018). Yet, comprehensive assessments of their abundances within the forest air were lacking. The presence of ~54% of all oomycete OTUs in both sampling events (Figure 4B) indicates a continuous presence of both peronosporan and pythialean oomycete spores and consequently a high proportion of potentially physiologically active oomycetes, including potential pathogens within forest ecosystems. Oomycetes pose a serious threat to forest health and functioning, it is therefore crucial to better understand their diversity and

233 distribution patterns of the total forest ecosystem, including air samples (Derevnina et al.,
 234 2016; Ajchler et al., 2017; Jung et al., 2018; Lang-Yona et al., 2018). We detected no OTUs
 235 assigned to the orders Saprolegniales, Lagenidiales or Myzocytiosporales, even though
 236 Jauss et al. (2020) found them in canopy and ground samples. All three orders are capable
 237 of forming dispersal stages, while their absence in our air samples could be due to a different
 238 timing of their sporulation, as our samples can only represent a snapshot of aerobic diversity.

239 Three dominant cercozoan orders were detected in air samples, but surprisingly no plant
 240 parasites of the Endomyxa. Testate amoebae from the orders Cryomonadida and
 241 Euglyphida occurred in high numbers. Cryomonadida (Thecofilosea) are filose amoeba with
 242 a robust extracellular organic tests (Adl et al., 2019). OTUs assigned to the Rhogostoma-
 243 lineage within the Cryomonadida dominated the samples. *Rhogostoma* species form resting
 244 stages resistant against desiccation for up to three months, although they form no cysts or
 245 zoospores (Mylnikova & Mylnikov, 2012; Öztoprak et al., 2020). *Assulina seminulum*, has a
 246 silica test with a remarkable size of 60-90 μm (Lara et al., 2010), and dominated the air
 247 dispersed Euglyphids, demonstrating that protists of this size can be still easily dispersed
 248 by air (Finlay, 2002). Not surprising was the dominance of Glissomonadida, represented by
 249 small flagellates of the families Sandonidae and Allapsidae. Their high abundance is
 250 consistent with observations Ploch et al (2016) and Jauss et al (2020). All these orders are
 251 an integral part of the protist phyllosphere microbiome (Agler et al., 2016; Dumack et al.,
 252 2017; Flues et al., 2018). Overall, their presence in the microbiome as well as their high
 253 abundance in air samples indicates canopies and their phyllosphere to be a potential filter
 254 not only for dust and particles (Weber et al., 2014; Chen et al., 2017), but also for
 255 microorganisms and potential plant pathogens.

256 Conclusion

257 A significant temporal variation in oomycetes indicates protistan community and,
 258 correspondingly, pathogen assembly to be driven by random factors and neutral processes,

259 while spatial differences in the vertical distribution of cercozoans and oomycetes were not
260 found. Accordingly, wind dispersal alone may well explain the ubiquitous presence of
261 Cercozoa and Oomycota (and likely of other protistan taxa) in the floodplain forest. Our
262 results further contribute to the understanding of how protists disperse, and which factors
263 drive the distribution of plant pathogens within forest ecosystems.

264 Acknowledgements

265 This paper was supported by the Priority Program SPP 1991: Taxon-omics – New
266 Approaches for Discovering and Naming Biodiversity of the German Research Foundation
267 (DFG) with funding to MB (1907/19-1) and MS (Schl 229/20-1). The authors would like to
268 thank Rolf Engelmann for his assistance with the field work by operating the canopy crane,
269 and the German Centre for Integrative Biodiversity Research (iDiv) for providing the site
270 access.

271 Conflict of Interest

272 None declared

273 Author Contributions

274 MS and MB conceived the study, RW and StS designed the sampling. BS assisted the library
275 preparation. SW assisted the sampling and contributed valuably to the discussion. AN
276 performed the sampling, laboratory work, bioinformatics analyses and outline of the
277 manuscript. R-TJ supervised the study, sampling and bioinformatic analyses, and wrote the
278 manuscript. All authors contributed to the manuscript and approved the final version.

279 Data availability

280 Raw sequence data have been submitted to the European Nucleotide Archive (ENA)
281 database under the Bioproject number PRJEB37525, with accession numbers ERS5388855
282 (Cercozoa) and ERS5388854 (Oomycota) respectively.

283 All figures, codes and detailed bioinformatic/statistical methods used in this study are
284 available at <https://github.com/RJauss/ToTheCanopyAndBeyond>.

285 References

- 286
287 Adl, S. M., Bass, D., Lane, C. E., Lukeš, J., Schoch, C. L., Smirnov, A., Agatha, S., Berney, C.,
288 Brown, M. W., Burki, F., Cárdenas, P., Čepička, I., Chistyakova, L., del Campo, J., Dunthorn,
289 M., Edvardsen, B., Eglit, Y., Guillou, L., Hampl, V., ... Zhang, Q. (2019). Revisions to the
290 Classification, Nomenclature, and Diversity of Eukaryotes. *Journal of Eukaryotic*
291 *Microbiology*, 66(1), 4–119. <https://doi.org/10.1111/jeu.12691>
292 Adl, S. M., & Gupta, V. S. R. (2006). Protists in soil ecology and forest nutrient cycling. In
293 *Canadian Journal of Forest Research* (Vol. 36, Issue 7, pp. 1805–1817).
294 <https://doi.org/10.1139/X06-056>
295 Agler, M. T., Ruhe, J., Kroll, S., Morhenn, C., Kim, S. T., Weigel, D., & Kemen, E. M. (2016).
296 Microbial Hub Taxa Link Host and Abiotic Factors to Plant Microbiome Variation. *PLoS*
297 *Biology*, 14(1), e1002352. <https://doi.org/10.1371/journal.pbio.1002352>
298 Ajchler, M., Łobocka, M., & Oszako, T. (2017). Pathogenic oomycetes of Phytophthora genus - a
299 new threat to forests in Europe. *Sylvan*, 161(10), 870–880. <https://sylvan.lasy.gov.pl/>
300 Amato, P., Ménager, M., Sancelme, M., Laj, P., Mailhot, G., & Delort, A.-M. (2005). Microbial
301 population in cloud water at the Puy de Dôme: Implications for the chemistry of clouds.
302 *Atmospheric Environment*, 39(22), 4143–4153. <https://doi.org/10.1016/j.atmosenv.2005.04.002>
303 Bass, D., Howe, A. T., Mylnikov, A. P., Vickerman, K., Chao, E. E., Edwards Smallbone, J., Snell,
304 J., Cabral Jr, C., & Cavalier-Smith, T. (2009). Phylogeny and Classification of Cercomonadida
305 (Protozoa, Cercozoa): Cercomonas, Eocercomonas, Paracercomonas, and Cavernomonas gen.
306 nov. *Protist*, 160, 483–521. <https://doi.org/10.1016/j.protis.2009.01.004>
307 Cavalier-Smith, T., Chao, E. E., & Lewis, R. (2018). Multigene phylogeny and cell evolution of
308 chromist infrakingdom Rhizaria: contrasting cell organisation of sister phyla Cercozoa and
309 Retaria. *Protoplasma*, 255, 1517–1574. <https://doi.org/10.1007/s00709-018-1241-1>
310 Chen, L., Liu, C., Zhang, L., Zou, R., & Zhang, Z. (2017). Variation in Tree Species Ability to
311 Capture and Retain Airborne Fine Particulate Matter (PM2.5). *Scientific Reports*, 7(1), 3206.
312 <https://doi.org/10.1038/s41598-017-03360-1>
313 Cowling, A. J. (1994). Protozoan distribution and adaptation. In J. Darbyshire (Ed.), *Soil Protozoa*
314 (pp. 5–42). CAB International.
315 Derevnina, L., Petre, B., Kellner, R., Dagdas, Y. F., Sarowar, M. N., Giannakopoulou, A., de la
316 Concepcion, J. C., Chaparro-Garcia, A., Pennington, H. G., van West, P., & Kamoun, S.
317 (2016). Emerging oomycete threats to plants and animals. In *Philosophical Transactions of the*
318 *Royal Society B: Biological Sciences* (Vol. 371, Issue 1709). Royal Society of London.
319 <https://doi.org/10.1098/rstb.2015.0459>
320 Dumack, K., Flues, S., Hermanns, K., & Bonkowski, M. (2017). Rhogostomidae (Cercozoa) from
321 soils, roots and plant leaves (*Arabidopsis thaliana*): Description of *Rhogostoma epiphylla* sp.
322 nov. and *R. cylindrica* sp. nov. *European Journal of Protistology*, 60, 76–86.
323 <https://doi.org/10.1016/j.ejop.2017.06.001>
324 Ellwood, M. D. F., & Foster, W. A. (2004). Doubling the estimate of invertebrate biomass in a
325 rainforest canopy. *Nature*, 429(6991), 549–551. <https://doi.org/10.1038/nature02560>
326 Evans, S. E., Dueker, M. E., Logan, J. R., & Weathers, K. C. (2019). The biology of fog: results
327 from coastal Maine and Namib Desert reveal common drivers of fog microbial composition.
328 *Science of The Total Environment*, 647, 1547–1556.
329 <https://doi.org/10.1016/j.scitotenv.2018.08.045>
330 Fawke, S., Doumane, M., & Schornack, S. (2015). Oomycete Interactions with Plants: Infection
331 Strategies and Resistance Principles. *Microbiology and Molecular Biology Reviews*, 79(3),

- 263–280. <https://doi.org/10.1128/mmbr.00010-15>
- Fenchel, T., & Finlay, B. J. (2004). The Ubiquity of Small Species: Patterns of Local and Global Diversity. *BioScience*, 54(8), 777–784. [https://doi.org/10.1641/0006-3568\(2004\)054\[0777:TUOSSP\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0777:TUOSSP]2.0.CO;2)
- Finlay, B. J. (2002). Global Dispersal of Free-Living Microbial Eukaryote Species. *Science*, 296, 1061–1063. <https://doi.org/10.1126/science.1070710>
- Finlay, B. J., Esteban, G. F., Clarke, K. J., & Olmo, J. L. (2001). Biodiversity of Terrestrial Protozoa Appears Homogeneous across Local and Global Spatial Scales. *Protist*, 152, 355–366. <https://doi.org/10.1078/1434-4610-00073>
- Fiore-Donno, A. M., & Bonkowski, M. (2020). Different community compositions between obligate and facultative oomycete plant parasites in a landscape-scale metabarcoding survey. *Biology and Fertility of Soils*. <https://doi.org/10.1007/s00374-020-01519-z>
- Fiore-Donno, A. M., Richter-Heitmann, T., & Bonkowski, M. (2020). Contrasting Responses of Protistan Plant Parasites and Phagotrophs to Ecosystems, Land Management and Soil Properties. *Frontiers in Microbiology*, 11, 1823. <https://doi.org/10.3389/fmicb.2020.01823>
- Flues, S., Blokker, M., Dumack, K., & Bonkowski, M. (2018). Diversity of Cercomonad Species in the Phyllosphere and Rhizosphere of Different Plant Species with a Description of *Neocercomonas epiphylla* (Cercozoa, Rhizaria) a Leaf-Associated Protist. *Journal of Eukaryotic Microbiology*, 65, 587–599. <https://doi.org/10.1111/jeu.12503>
- Foissner, W. (2009). Protist diversity and distribution: some basic considerations. In W. Foissner & D. L. Hawksworth (Eds.), *Protist Diversity and Geographical Distribution* (pp. 1–8). https://doi.org/10.1007/978-90-481-2801-3_1
- Foissner, W., & Hawksworth, D. L. (Eds.). (2009). *Protist Diversity and Geographical Distribution* (Vol. 8). Springer Netherlands. <https://doi.org/10.1007/978-90-481-2801-3>
- Fuzzi, S., Mandrioli, P., & Perfetto, A. (1997). Fog droplets—an atmospheric source of secondary biological aerosol particles. *Atmospheric Environment*, 31(2), 287–290. [https://doi.org/10.1016/1352-2310\(96\)00160-4](https://doi.org/10.1016/1352-2310(96)00160-4)
- Geisen, S., Mitchell, E. A. D., Wilkinson, D. M., Adl, S., Bonkowski, M., Brown, M. W., Fiore-Donno, A. M., Heger, T. J., Jassey, V. E. J., Krashevskaya, V., Lahr, D. J. G., Marcisz, K., Mulot, M., Payne, R., Singer, D., Anderson, O. R., Charman, D. J., Ekelund, F., Griffiths, B. S., ... Lara, E. (2017). Soil protistology rebooted: 30 fundamental questions to start with. In *Soil Biology and Biochemistry* (Vol. 111, pp. 94–103). <https://doi.org/10.1016/j.soilbio.2017.04.001>
- Genitsaris, S., Kormas, K. A., Christaki, U., Monchy, S., & Moustaka-Gouni, M. (2014). Molecular diversity reveals previously undetected air-dispersed protist colonists in a Mediterranean area. *Science of the Total Environment*, 478, 70–79. <https://doi.org/10.1016/j.scitotenv.2014.01.071>
- Goheen, E. M., & Frankel, S. J. (2009). Proceedings of the fourth meeting of the International Union of Forest Research Organizations (IUFRO) Working Party S07.02.09: Phytophthoras in forests and natural ecosystems. In *Gen. Tech. Rep. PSW-GTR-221*. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 334 p (Vol. 221). <https://doi.org/10.2737/PSW-GTR-221>
- Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., De Vargas, C., Decelle, J., Del Campo, J., Dolan, J. R., Dunthorn, M., Edvardsen, B., Holzmann, M., Kooistra, W. H. C. F., Lara, E., Le Bescot, N., Logares, R., ... Christen, R. (2013). The Protist Ribosomal Reference database (PR2): A catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Research*, 41(D1), D597–D604. <https://doi.org/10.1093/nar/gks1160>
- Harder, C. B., Rønn, R., Brejnrod, A., Bass, D., Al-Soud, W. A., & Ekelund, F. (2016). Local diversity of heathland Cercozoa explored by in-depth sequencing. *ISME Journal*, 10(10), 2488–2497. <https://doi.org/10.1038/ismej.2016.31>
- Jauss, R.-T., Walden, S., Fiore-Donno, A.-M., Dumack, K., Schaffer, S., Wolf, R., Schlegel, M., & Bonkowski, M. (2020). From forest soil to the canopy: increased habitat diversity does not increase species richness of Cercozoa and Oomycota in tree canopies. *Authorea Preprints*.

- 384 <https://doi.org/10.22541/AU.158679920.02842084>
- 385 Jousset, A., Lara, E., Wall, L. G., & Valverde, C. (2006). Secondary metabolites help biocontrol
- 386 strain *Pseudomonas fluorescens* CHA0 to escape protozoan grazing. *Applied and*
- 387 *Environmental Microbiology*, 72(11), 7083–7090. <https://doi.org/10.1128/AEM.00557-06>
- 388 Judelson, H. S. (2017). Metabolic Diversity and Novelty in the Oomycetes. *Annual Review of*
- 389 *Microbiology*, 71, 21–39. <https://doi.org/10.1146/annurev-micro-090816-093609>
- 390 Jung, T., Pérez-Sierra, A., Durán, A., Jung, M. H., Balci, Y., & Scanu, B. (2018). Canker and decline
- 391 diseases caused by soil- and airborne *Phytophthora* species in forests and woodlands. In
- 392 *Persoonia: Molecular Phylogeny and Evolution of Fungi* (Vol. 40, pp. 182–220). Nationaal
- 393 Herbarium Nederland. <https://doi.org/10.3767/persoonia.2018.40.08>
- 394 Kageyama, K., & Asano, T. (2009). Life cycle of plasmodiophora brassicae. *Journal of Plant*
- 395 *Growth Regulation*, 28(3), 203–211. <https://doi.org/10.1007/s00344-009-9101-z>
- 396 Lang-Yona, N., Pickersgill, D. A., Maurus, I., Teschner, D., Wehking, J., Thines, E., Pöschl, U.,
- 397 Després, V. R., & Fröhlich-Nowoisky, J. (2018). Species Richness, rRNA Gene Abundance,
- 398 and Seasonal Dynamics of Airborne Plant-Pathogenic Oomycetes. *Frontiers in Microbiology*,
- 399 9(NOV), 2673. <https://doi.org/10.3389/fmicb.2018.02673>
- 400 Lara, E., & Belbahri, L. (2011). SSU rRNA reveals major trends in oomycete evolution. *Fungal*
- 401 *Diversity*, 49, 93–100. <https://doi.org/10.1007/s13225-011-0098-9>
- 402 Lara, E., Heger, T. J., Scheihing, R., & Mitchell, E. A. D. (2010). COI gene and ecological data
- 403 suggest size-dependent high dispersal and low intra-specific diversity in free-living terrestrial
- 404 protists (Euglyphida: Assulina): High dispersal in testate amoebae. *Journal of Biogeography*,
- 405 38(4), 640–650. <https://doi.org/10.1111/j.1365-2699.2010.02426.x>
- 406 Mahé, F., Rognes, T., Quince, C., de Vargas, C., & Dunthorn, M. (2015). Swarm v2: highly-scalable
- 407 and high-resolution amplicon clustering. *PeerJ*, 3, 1–12. <https://doi.org/10.7717/peerj.1420>
- 408 Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
- 409 *EMBnet.Journal*, 17(1), 10. <https://doi.org/10.14806/ej.17.1.200>
- 410 Meyer, M., & Kircher, M. (2010). Illumina Sequencing Library Preparation for Highly Multiplexed
- 411 Target Capture and Sequencing. *Cold Spring Harbor Protocols*, 2010(6), pdb.prot5448--
- 412 pdb.prot5448. <https://doi.org/10.1101/pdb.prot5448>
- 413 Moon-van der Staay, S. Y., Tzeneva, V. A., Van Der Staay, G. W. M., De Vos, W. M., Smidt, H., &
- 414 Hackstein, J. H. P. (2006). Eukaryotic diversity in historical soil samples. *FEMS Microbiology*
- 415 *Ecology*, 57(3), 420–428. <https://doi.org/10.1111/j.1574-6941.2006.00130.x>
- 416 Mylnikova, Z. M., & Mylnikov, A. P. (2012). Structure of filose amoeba *Rhagostoma minus* Belar
- 417 1921 (Cryomonadida, Cercozoa) cell. *Inland Water Biology*, 5(3), 236–240.
- 418 <https://doi.org/10.1134/S1995082912020101>
- 419 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,
- 420 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H.
- 421 (2019). *vegan: Community Ecology Package*. <https://cran.r-project.org/package=vegan>
- 422 Ozanne, C. H. P., Anhuf, D., Boulter, S. L., Keller, H., Kitching, R. L., Körner, C., Meinzer, F. C.,
- 423 Mitchell, A. W., Nakashizuka, T., Silva Dias, P. L., Stork, N. E., Wright, S. J., & Yoshimura,
- 424 M. (2003). Biodiversity meets the atmosphere: A global view of forest canopies. In *Science*
- 425 (Vol. 301, Issue 5630, pp. 183–186). American Association for the Advancement of Science.
- 426 <https://doi.org/10.1126/science.1084507>
- 427 Öztoprak, H., Walden, S., Heger, T., Bonkowski, M., & Dumack, K. (2020). What drives the
- 428 diversity of the most abundant terrestrial cercozoan family (Rhagostomidae, cercozoa,
- 429 rhizaria)? *Microorganisms*, 8(8), 1–16. <https://doi.org/10.3390/microorganisms8081123>
- 430 Pedersen, T. L. (2020). *ggraph: An Implementation of Grammar of Graphics for Graphs and*
- 431 *Networks* (2.0.3). <https://cran.r-project.org/package=ggraph>
- 432 Pepper, I. L., & Dowd, S. E. (2009). Aeromicrobiology. In R. M. Maier, I. L. Pepper, & C. P. Gerba
- 433 (Eds.), *Environmental Microbiology (Second Edition)* (Second Edi, pp. 83–102). Academic
- 434 Press. <https://doi.org/https://doi.org/10.1016/B978-0-12-370519-8.00005-5>
- 435 Petz, W., & Foissner, W. (1988). Spatial separation of terrestrial ciliates and testaceans (Protozoa): a

- contribution to soil ciliatostasis. *Acta Protozoologica*, 27(3), 249–258.
- Ploch, S., Rose, L. E., Bass, D., & Bonkowski, M. (2016). High Diversity Revealed in Leaf-Associated Protists (Rhizaria: Cercozoa) of Brassicaceae. *The Journal of Eukaryotic Microbiology*, 63(5), 635–641. <https://doi.org/10.1111/jeu.12314>
- R Core Team. (2019). *R: A Language and Environment for Statistical Computing*. <https://www.r-project.org/>
- Robideau, G. P., De Cock, A. W. A. M., Coffey, M. D., Voglmayr, H., Brouwer, H., Bala, K., Chitty, D. W., Désaulniers, N., Eggertson, Q. A., Gachon, C. M. M., Hu, C. H., Küpper, F. C., Rintoul, T. L., Sarhan, E., Verstappen, E. C. P., Zhang, Y., Bonants, P. J. M., Ristaino, J. B., & André Lévesque, C. (2011). DNA barcoding of oomycetes with cytochrome c oxidase subunit I and internal transcribed spacer. *Molecular Ecology Resources*, 11(6), 1002–1011. <https://doi.org/10.1111/j.1755-0998.2011.03041.x>
- Rogerson, A., & Detwiler, A. (1999). Abundance of airborne heterotrophic protists in ground level air of South Dakota. *Atmospheric Research*, 51, 35–44. [https://doi.org/10.1016/S0169-8095\(98\)00109-4](https://doi.org/10.1016/S0169-8095(98)00109-4)
- Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: A versatile open source tool for metagenomics. *PeerJ*, 2016(10). <https://doi.org/10.7717/peerj.2584>
- Thines, M. (2014). Phylogeny and evolution of plant pathogenic oomycetes-a global overview. *European Journal of Plant Pathology*, 138(3), 431–447. <https://doi.org/10.1007/s10658-013-0366-5>
- Urich, T., Lanzén, A., Qi, J., Huson, D. H., Schleper, C., & Schuster, S. C. (2008). Simultaneous Assessment of Soil Microbial Community Structure and Function through Analysis of the Meta-Transcriptome. *PLoS ONE*, 3(6), e2527. <https://doi.org/10.1371/journal.pone.0002527>
- Voss, C., Fiore-Donno, A. M., Guerreiro, M. A., Peršoh, D., & Bonkowski, M. (2019). Metatranscriptomics reveals unsuspected protistan diversity in leaf litter across temperate beech forests, with Amoebozoa the dominating lineage. *FEMS Microbiology Ecology*, 95(10), 142. <https://doi.org/10.1093/femsec/fiz142>
- Weber, F., Kowarik, I., & Säumel, I. (2014). Herbaceous plants as filters: Immobilization of particulates along urban street corridors. *Environmental Pollution*, 186, 234–240. <https://doi.org/https://doi.org/10.1016/j.envpol.2013.12.011>
- Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. <https://ggplot2.tidyverse.org>