An acute dose of intranasal oxytocin rapidly increases maternal communication and maintains maternal care in primiparous postpartum California mice

Caleigh D. Guoynes and Catherine A. Marler

Department of Psychology, University of Wisconsin, Madison, WI 53706, USA

Abstract

Maternal-offspring communication and care are essential for offspring survival. Oxytocin (OXT) is known for its role in initiation of maternal care, but whether OXT can rapidly influence maternal behavior or ultrasonic vocalizations (USVs; above 50 kHz) has not been examined. To test for rapid effects of OXT, California mouse mothers were administered an acute intranasal (IN) dose of OXT (0.8 IU/kg) followed by a separation test with three phases: habituation with pups in a new testing chamber, separation via a wire mesh, and finally reunion with pups. We measured maternal care, maternal USVs, and pup USVs. In mothers, we primarily observed simple sweep USVs, a short downward sweeping call around 50 kHz, and in pups we only observed pup whines, a long call with multiple harmonics ranging from 20 kHz to 50 kHz. We found that IN OXT rapidly and selectively enhanced the normal increase in maternal simple sweep USVs when mothers had physical access to pups (habituation and reunion), but not when mothers were physically separated from pups. Maternal-pup USVs were correlated upon reunion, but IN OXT did not influence this correlation. Finally, mothers given IN OXT showed a more positive change in retrievals/carrying and greater total maternal care upon reunion. Behavioral changes were specific to maternal behaviors (e.g. retrievals) as mothers given IN OXT did not differ from controls in stress-related behaviors (e.g. freezing). Overall, these findings highlight the rapid effects and context-dependent effect a single dose of IN OXT has on both maternal USV production and offspring care.

Key words: *oxytocin, maternal care, ultrasonic vocalizations (USVs)*

Introduction

Quality of maternal care has significant impacts on offspring survival outcomes across many mammalian species (Fairbanks & McGuire, 1995; Nowak et al., 2000; Blomquist, 2013; Lerch-Haner et al., 2008; Francis et al., 1999). These studies underscore the importance of maternal behavior from an evolutionary perspective. However, the proximate mechanisms that reinforce maternal care remain more elusive. Several studies in rodents illustrate that pup whines quickly and reliably elicit maternal care (Allin & Banks, 1972; Hennessy et al., 1980; Shiavo et al., 2020; Portfors, 2007; White et al., 1992; Bell, 2018). Other studies, however, show that mothers are more apt to exhibit care in response to stressful events or disturbances and that pup calls do not influence their care above and beyond the disturbance (Brewster & Leon, 1980). It has also been shown that female house mice prefer calls from their own pups and can locate their own faster than alien pups (Mogi et al., 2017). These studies show that pup whines can elicit changes in maternal behavior. However, the role of maternal vocalizations in this relationship has not been studied. Adult rats, including mothers, make spontaneous vocalizations that typically occur above 50 kHz when presented with drug and social rewards (Knutson et al., 2002; Simola et al., 2012). The association of reward with these calls in rats is interpreted as an indicator of a positive internal state. In mothers, USVs may indicate maternal motivation but could also reinforce maternal care and the mother-offspring bond (Stern, 1989; Cheng, 1992) or reduce maternal anxiety (Arnon et al., 2014).

Complementing the stimulus of maternal and pup vocalizations, neural substrates also play an important role in processing and producing behaviors that support maternal care. One neuropeptide hormone likely involved in this process is oxytocin (OXT); OXT modulates many

social behaviors including bonding and parental care (Bartz & Hollander, 2006; Lim & Young, 2006; Maynard et al., 2018; Taylor et al., 2015; Guoynes & Marler 2020). During mammalian birth, plasma OXT increases to stimulate parturition and milk let-down in mothers; this increase was likely co-opted during evolution to also facilitate maternal care (Rosenblatt, 1969; Moltz et al., 1970; Pederson & Prange, 1979; Feldman et al., 2007). Immediately after parturition, rising levels of peripheral estrogen (Siegel & Rosenblatt, 1975) prime OXT to initiate maternal behaviors in rats (Pedersen & Prange, 1979). Additionally, OXT knockout mice have greater latency to onset of maternal behaviors (Takayanagi et al., 2005). In the brain, OXT antagonists blocked maternal behavior after natural delivery in pregnant rats (Van Leengoed et al., 1987). This reveals that central OXT is important for initiating maternal care in many species.

In addition to differences in OXT levels in mothers influencing maternal care, receptor densities are also clearly important. In rats genetically selected for differences in maternal care, high grooming compared to low grooming had more OXT receptors in the bed nucleus of the stria terminalis, medial preoptic area, central nucleus of the amygdala, and these differences were observed in both non-lactating and lactating females (Francis et al., 2000; Champagne et al., 2001). Acute activation of maternal care by OXT is indirectly supported by an optogenetics study in which dopamine neurons were activated in the anteroventral periventricular nucleus (AVPV) that monosynaptically connect to and activate OXT neurons in the paraventricular nucleus (PVN), resulting in enhanced maternal care (Scott et al., 2015). Collectively, these studies provide strong evidence that OXT plays an important role in activating maternal care and improving its efficiency.

OXT also plays a role in the production and perception of vocalizations. In mice and other rodents, the majority of their vocalizations occur above 50 kHz and are called ultrasonic

vocalizations (USVs) (Mulsolf et al., 2010; Scattoni et al., 2009; Kalcounis-Ruepell et al., 2010; Arriaga & Jarvis, 2013). In OXT knockout mice, OXT null pups emit fewer USVs in response to separation from their mother compared to wildtype mice (Winslow & Insel, 2002). This suggests that OXT may enhance pup communication with their mother. OXT can also improve the signal-to-noise ratio in mothers responding to pup calls via mediation of temporal inhibition and excitation in the left auditory cortex of female mice, leading to increased pup retrievals (Marlin et al., 2015). These data provide evidence that OXT is changing the perception and social salience of pup calls, leading to increased maternal care. Furthermore, in humans, the OXT receptor has a polymorphism (rs53576) with functional significance. The genotype GG (produces more OXTRs compared to AG or AA genotypes) is associated with better ability to discriminate content of language under noisy conditions (Tops et al., 2011). This suggests that across taxa, OXT may play an important neuromodulatory role in promoting sensory processing and behavioral response to social auditory information. A key social behavior that has previously not been measured is maternal vocalizations. We speculated that mothers modulate vocalization quantity or type when interacting with their offspring and that maternal vocalizations would be associated with maternal care. Moreover, we predicted that OXT would modulating these vocalizations.

The California mouse is a strictly monogamous, biparental rodent species uniquely suited to examine how OXT modulates auditory sensory processing, vocal production, and social behavior. California mice have a diverse, well-characterized repertoire of ultrasonic vocalizations (USVs) including simple sweeps, complex sweeps, syllable vocalizations, barks, and pup whines (Briggs et al. 2011; Kalcounis-Ruepell et al., 2006; Pultorak et al., 2015; Rieger

& Marler, 2018). Preliminary data indicate that maternal simple sweeps and pup whines are the primary maternal USVs used in interactions between mothers and pups.

In the current study, we aimed to address the gaps in our understanding of proximate mechanisms that contribute to maternal care by determining the association between maternal vocalizations and maternal care and whether an acute dose of IN OXT in primiparous female California mice could rapidly increase both maternal vocalizations and care. We hypothesized that 1) maternal care would be associated with maternal USVs and that 2) IN OXT would have a positive effect on maternal care. Specifically, we predicted that during our behavioral paradigm, IN OXT would increase maternal care, maternal USV production, and enhance the correlation between pup USVs and maternal USVs.

Methods

Animals

We used 24 primiparous postpartum female *P. californicus* aged 5–10 months because of ages of previously unpaired females available within our colony. Females across this age range also show equivalent corticosterone responses to corticotrophin releasing hormone and dexamethasone stress (Harris & Saltzman, 2013), suggesting that females between this age range have comparable neuroendocrine responses. Females were pair-housed (1 female, 1 male, and 1–3 pups per cage; 48 × 27 × 16 cm) under a 14L: 10D light cycle. Animals were maintained in accordance with the National Institute of Health Guide for the Care and Use of Laboratory Animals. Female and male pairs were randomly assigned, and pairs produced offspring in their home cage. After females were visibly pregnant, cages were checked once daily for pups. Female mothers were randomly assigned to either the saline control group (N=11) or the IN OXT group (N=13).

Intranasal Oxytocin Preparation

Treatments were infused intranasally with either sterile saline control or IN OXT (0.8 IU/kg) (Bachem, Torrance, California). The IN OXT dose was equivalent to doses used in other rodent species (Bales et al. 2014; Guoynes et al. 2018; Murgatroyd et al. 2016) and similar to weight-adjusted doses used in clinical studies examining the effects of IN OXT on social deficits in autism (Bales et al. 2013). IN OXT was dissolved in saline and prepared in one batch that was aliquoted into small plastic tubes and frozen at 20°C. IN OXT was defrosted just prior to administration. A blunt cannula needle (33-gauge, 2.8 mm length; Plastics One, Roanoke, Virginia) was attached to cannula tubing, flushed, and filled with the compound, then attached to an airtight Hamilton syringe (Bachem, Torrance, California). The animal was scruffed and 25 uL of compound was expelled dropwise through the cannula needle, alternating between left and right nostrils. Mice are obligate nose-breathers and thus the solution was quickly absorbed into the nasal mucosa (~10seconds). One person conducted all IN OXT administrations throughout the entire procedure to maintain consistency in handling and IN OXT infusion.

We chose to use the method of intranasal administration of IN OXT for two primary reasons. (1) IN OXT is used in clinical studies and is less invasive, does not require special transporters for the molecule, and is less stressful compared to intracerebroventricular or intraperitoneal injections (Talegaonkar & Mishra 2004). (2) IN OXT shows similar behavioral effects as centrally administered OXT, increases CSF plasma concentrations of OXT, and reaches the relevant brain areas in both human and animal models (Neumann et al., 2013; Striepens et al., 2013; Lee et al. 2018; Oppong-Damoah et al., 2019; Lee et al., 2020). Several studies have also shown changes in plasma OXT concentrations that peak between 15 to 30 min

post-administration (Freeman et al., 2016; Gossen et al., 2012). These results suggest that IN OXT passes through the blood-brain barrier to exert central effects with minimal stress on the animal. Other studies that suggest that some of the effects of IN OXT are acting through peripheral mechanisms (Churchland & Winkielman, 2012; Quintana et al., 2015; Leng & Ludwig, 2016). Regardless of whether IN OXT is directly targeting the brain, is acting through peripheral mechanisms, or a combination of both, IN OXT has been shown to rapidly alter social behavior in adult California mice (Steinman et al., 2016).

Behavioral Testing and USV Recording

In order to test the effects of IN OXT on acute maternal care, we conducted this experiment in a novel chamber where mothers and pups could be briefly isolated from the father. Separation from the mate may be a mild stressor, but would occur in natural populations in response to competing demands such as pup care, foraging and defending territories. Moreover, California mice do not exhibit a change in short- or long-term paternal care in response to corticosterone (Harris et al., 2017), show limited correlations between individual baseline corticosterone levels and behavior (Dlugosz et al., 2012), parents exhibit blunted behavioral response to predator odor stress (Chauke et al., 2011) and diel corticosterone cycle between single mothers and paired mothers does not differ (Zhao et al., 2019). While we can not exclude an effect of a baseline level of stress, it is both a normal experience for these mice in the wild and we do not expect corticosterone to influence the results above and beyond our IN OXT manipulation.

On postnatal day (PND) two to three, fathers were temporarily removed from the home cage, and the home cage with the mother and her pups was transferred to a behavioral testing

room. In the behavioral testing room, mothers were randomly selected to receive 25 microliters of either 0.8 IU/kg IN OXT or saline control intranasally. Immediately after dosing, mothers and pups were placed into one side of a partitioned two-chambered apparatus (45.0 cm × 30.0 cm × 30.0 cm) that contained a circular opening (3.8 cm in diameter, center of opening 7 cm from the side wall) covered by a wire mesh (**Fig. 1A**). This apparatus, like their home cages, had approximately 1/2 inch of aspen shavings covering the entire floor. Microphones sensitive to ultrasonic frequencies (described below) were placed on each side of the divider, such that the microphones were far enough apart to identify the source (chamber) of the calls (Pultorak et al., 2018) (**Fig. 1A**).

Figure 1. (A) Schematic for experimental design. Mother and pup groups were temporarily removed from their home cage and placed in the right side of two-chambered apparatus (five min) to habituate to the testing arena. Next, pups were moved from the right chamber and placed into the left chamber (three min). Lastly, the researchers lifted the mesh gate separating the right and left chambers, allowing mother-pup interactions (five min). Animals not to scale in diagram. **(B) Ultrasonic vocalizations (USVs) on a spectrogram.** Pup whines have multiple harmonics, a peak frequency around 20 kHz, and downward modulation at the end of the call that distinguish these calls from adult syllable vocalizations. Maternal simple sweeps have short downward-sweeping vocalizations that sweep through multiple frequencies, typically between 80 kHz and 40 kHz.

For each test, there were always two researchers present who coordinated activation of the audio software and the video camera at the same time using visual cues. This coordination allowed us to subsequently compare USVs and behavior with temporal precision. The test consisted of three phases that occurred in immediate succession: habituation, separation, and reunion. During the five-min habituation phase mothers and pups were placed together on the right side of the chamber with the partition down and allowed to freely interact with each other. During the three-minute separation phase, the pups were removed from the mother and placed on the other side of the partitioned divider. Mothers remained in the right-most chamber and pups

were isolated in the left-most chamber. This setup allowed visual, auditory, and olfactory communication between pups and their mother, but restricted physical contact between individuals until the mesh wire was removed. Lastly, during the five-minute reunion phase, the mesh divider was lifted, and mothers could retrieve and interact with their pups. USVs and video were recorded for the entire 13-minute period. We chose a five-minute initial mother-pup interaction time to allow mothers time to adjust to the chamber and to mirror the time in the reunion phase where we measured latency to enter the chamber. This time period is important because it is the first time that the mothers and her pups are removed from the home cage and the father, so this time period served as an initial measure of behavior. Results from a pilot study measuring maternal retrievals indicates that five-minutes allowed most mothers to enter the chamber, approach the pups, and engage in maternal behaviors. We shortened the separation phase to three minutes because it was still sufficient to see signs of maternal distress but minimized the time that the pups were away from their mother.

Behavior Quantification

All behaviors from the videos were scored in random order and by an independent, blind observer. During each video, maternal behaviors (licking and grooming, huddling, and retrieving/carrying) were quantified. Of note, unlike house mice and rats that show several different types of nursing and huddling behavior, California mouse mothers do not show arched back nursing (Bester-Meredith et al., 2017; Johnson et al., 2015). The definitions of behaviors measured are detailed in an ethogram (**S. Fig. 1**). To gain insights into the correlations between maternal behavior and maternal and pup USVs, videos were coded by the exact time that mothers were huddling, licking and grooming, and retrieving/carrying. Huddling was counted

when mothers were physically over their pups' bodies (Bester-Meredith & Marler, 2003). Retrieving/carrying was counted when mothers picked up their pups and transferred them to a different location. Using the precise times that mothers engaged in different types of maternal care (or none at all) throughout the 13-minute testing window, we counted the maternal USVs within those windows. This allowed us to determine how maternal behavior was related to maternal USV production.

S. Figure 1. Ethogram with description of behaviors measured.

Ultrasonic Vocalization Analysis

Techniques used for recording were similar to those previously used in our laboratory (Pultorak et al. 2017; Rieger & Marler 2018). USVs were collected using two Emkay/Knowles FG series microphones capable of detecting broadband sound (10–120 kHz). Microphones were placed at the far ends of each of the two chambers. Microphone channels were calibrated to equal gain (– 60 dB noise floor). We used RECORDER software (Avisoft Bioacoustics) to produce triggered WAV file recordings (each with a duration of 0.5 s) upon the onset of a sound event that surpassed a set threshold of 5% energy change (Kalcounis-Rueppell et al., 2010). Recordings were collected at a 250 kHz sampling rate with a 16-bit resolution. Spectrograms were produced with a 512 FFT (Fast Fourier Transform) using Avisoft-SASLab Pro sound analysis software (Avisoft Bioacoustics). The only USVs found in these recordings were pup whines and maternal simple sweeps. Pup whines have a peak frequency around 20 kHz (Johnson et al., 2017; Kalcounis-Rueppell et al., 2018a) and the typical downward modulation at the end of the call often distinguishes these calls from adult syllable vocalizations (Nathaniel Rieger, Jose Hernandez, & Catherine Marler, unpublished) (**Figure 1B**). Maternal simple sweeps were

categorized by short downward-sweeping vocalizations that sweep through multiple frequencies, typically between 80 kHz and 40 kHz (Kalcounis-Rueppell et al., 2018b) (**Figure 1B**). It is extremely rare for pups to produce simple sweep USVs during PND 0-4 (Rieger, N. S., Hernandez, J. B., and Marler, C. M., unpublished). When young pups do produce simple sweeps, they are produced much quicker, and present completely vertical on the spectrogram (Johnson et al., 2017). This makes these rare pup simple sweeps easy to distinguish from the slower adult simple sweep USVs (**Fig. 1B**). Because of their different spectrogram and acoustic properties, all USVs could be categorized and counted by combined visual and auditory inspections of the WAV files (sampling rate reduced to 11,025 kHz, corresponding to 4% of real-time playback speed).

Data Analysis

Statistical analyses were conducted using the program R. Significance level was set at $p < 0.05$ for all analyses and all tests were two-tailed. All reported p-values were corrected using Bonferroni adjustments to control for multiple comparisons. Grubb's outlier test was performed, and outliers for maternal vocalizations and freezing were removed from all analyses. Two mice (one control and one IN OXT-treated mouse) were Grubb's outliers ($p < 0.05$) for freezing (likely due to train noise during the test). One control mouse was a Grubb's outlier ($p < 0.05$) for both maternal and pup USVs. All analyses used a generalized linear mixed model (GLMM). Treatment condition was used in all models as each female was given one treatment (between-subjects design). Thus, when a relationship between two variables was significant, treatment was left as a moderator in the model even if not significant. All reported p-values have been corrected using Bonferroni adjustments to control for multiple comparisons.

Effects of IN OXT on vocalizations

To assess differences in total USV production across testing conditions, a within-subjects two-way ANOVA was used. To correct for differences in total time of the three testing phases, average number of USVs/second during each phase for each animal was calculated and used for comparison. To assess main effects of treatment, Student's t-test was used in each of the three time points.

To examine the effects of IN OXT in the relationships between maternal simple sweep USVs and pup whine USVs, an interactive multivariate model was used (e.g. [Maternal behavior] ~ [Maternal USV] + [treatment]). Factor included in all models was treatment condition.

Effects of IN OXT on maternal and non-maternal behaviors

For maternal and non-maternal behavioral analysis, behavioral changes after the separation event, were calculated to examine reunion behaviors with and without OXT administration. To examine changes in behavior over time and after the separation challenge with and without OXT, the scores from total duration of each behavior in the reunion phase were subtracted from total duration of each behavior in the habituation phase. Thus, positive scores indicate more of the behavior was observed during the reunion phase and negative scores indicate more of the behavior was observed during the habituation phase. To compare main effects of IN OXT and saline control on maternal behavior, Students t-tests were used to assess behavioral outcomes. To calculate total maternal behavior, amount of time spent huddling and amount of time spent retrieving were summed. To calculate total non-maternal behavior, amount of time spent autogrooming, rearing, and freezing were summed.

Correlations between maternal care and maternal USVs and maternal care and pup USVs

To assess for mediation by IN OXT in the relationships between (a) maternal USVs and maternal behavior and (b) maternal behavior and pup USVs, a multivariate comparison was used. Factors included in the model were treatment condition and the interaction between treatment and maternal behavior.

Results

Effects of IN OXT on vocalizations

To determine how testing conditions affected vocal production in mothers and pups, we first assessed number of maternal and pup USVs per second across the habituation, separation, and reunion phases, and in response to IN OXT versus saline. Controlling for within subject analyses and treatment effects, mothers made fewer simple sweeps/second during the separation phase compared to the habituation or reunion phases ($F_{2,20}=13.00$, $p<0.00001$). (**Fig. 2A**). In the habituation phase, IN OXT mothers produced more simple sweeps than control mothers ($F_{2,20}=5.83$, $p<0.03$, $\Delta R^2=0.23$) (**Fig. 2A**). In the separation phase, IN OXT and control mothers did not differ in number of simple sweeps produced ($F_{2,20}=1.86$, $p=0.19$, $\Delta R^2=0.09$) (**Fig. 2A**). Similar to the habituation phase, in the reunion phase, IN OXT mothers showed a nonsignificant trend for producing more simple sweeps than control mothers ($F_{2,20}=3.13$, $p=0.08$, $\Delta R^2=0.15$) (**Fig. 2A**).

Figure 2. Rapid effects of IN OXT on USV production in mothers and pups. (A) All mothers made more simple sweeps when given free access to their pups (during the habituation and reunion phases). Importantly, IN OXT mothers made more simple sweeps when given free access to their pups during habituation and reunion, but not when they were physically apart from pups during separation. **(B)** All pups made more whines when first placed into the chamber during the habituation phase. There were no effects of maternal IN OXT on pup USVs. pups made more whines during the habituation phase than the separation and reunion phases. There was no effect of IN OXT treatment of pup whines during habituation, separation or reunion. **(C)** Mediation analysis of the relationship between maternal USVs, pup USVs and treatment in the habituation phase showed no simple effects of maternal USVs, no simple effects of

treatment, but IN OXT showed a nonsignificant trend for the two-way interaction between maternal simple sweep USVs and treatment. **(D)** The separation phase showed no simple effects of maternal USVs, no simple effects of treatment, and no effect of interaction. **(E)** The reunion phase showed a significant positive correlation between maternal simple sweeps and pup whines during, no simple effect of treatment, and no effect of interaction. Correlation line collapsing across treatment groups. * $p < 0.05$; # $p < 0.10$.

Across the three phases, pup USVs showed both consistency and differences in relationships with maternal USVs but showed no effect of IN OXT. Controlling for within-subject analyses, pups made more whines during the habituation phase than the separation and reunion phases ($F_{2,20} = 25.26$, $p < 0.0000001$) (**Fig. 2B**). Pups with IN OXT and control mothers did not differ in number of USVs produced in the habituation phase ($F_{1,20} = 0.35$, $p = 0.56$, $\Delta R^2 = 0.02$), separation phase ($F_{1,20} = 1.64$, $p = 0.22$, $\Delta R^2 = 0.08$) or the reunion phase ($F_{1,20} = 0.68$, $p = 0.42$, $\Delta R^2 = 0.04$) (**Fig. 2B**).

Next, we examined the relationship between maternal simple sweeps and pup whines using a model with treatment as a covariate. There were no significant correlations between maternal and pup USVs in either the habituation ($F_{1,20} = 1.63$, $p = 0.22$) (**Fig. 2C**) or the separation phase ($F_{1,20} = 0.03$, $p = 0.86$) (**Fig. 2D**). However, during the reunion phase, maternal simple sweeps and pup whines positively correlated ($F_{1,20} = 7.51$, $p < 0.02$, $\Delta R^2 = 0.21$). For each pup whine, there were approximately 1.17 maternal simple sweeps (**Fig. 2E**). There were no simple effects of OXT on the correlation between maternal-pup USVs: habituation ($F_{1,20} = 0.25$, $p = 0.62$) (**Fig. 2C**), separation ($F_{1,20} = 1.27$, $p = 0.27$) (**Fig. 2D**), reunion ($F_{1,20} = 0.15$, $p = 0.69$) (**Fig. 2E**). Finally, our model also assessed the two-way interaction between maternal simple sweep USVs and treatment. IN OXT showed a nonsignificant trend for the two-way interaction between maternal simple sweep USVs and treatment, with IN OXT animals having a more positive slope (non-significant trend) than controls ($F_{1,20} = 3.71$, $p = 0.069$) (**Fig. 2C**). Slope for IN OXT-treated

mothers did not differ from controls in either the separation ($F_{1,20}=0.07, p=0.79$) (**Fig. 2D**) or reunion ($F_{1,20}=2.90, p=0.10$) (**Fig. 2E**) phases.

Effects of IN OXT on maternal and non-maternal behaviors

To assess the impact of IN OXT on maternal care following separation from pups, we calculated change in maternal care from habituation to reunion. This allowed us to determine how IN OXT affected response to pup separation above and beyond any initial effects of IN OXT observed in the habituation phase. IN OXT mothers showed a non-significant trend for a shorter latency to approach pups ($F_{1,20}=3.63, p=0.072, \Delta R^2=0.16$) (**Fig. 3A**). We also tested for main effects of IN OXT in several maternal and non-maternal behaviors. Negative scores mean that the behavior occurred more frequently during habituation and positive scores mean that the behavior occurred more frequently during reunion. Mothers tended to retrieve/carry more during the habituation phase and huddle more during the reunion phase. Mothers given IN OXT did not show any differences in huddling from controls ($F_{1,20}=0.62, p=0.44$) (**Fig. 3B**). However, mothers given IN OXT showed a significantly more positive change in retrieval/carrying behavior from the habituation to the reunion phase ($F_{1,20}=6.71, p<0.04, \Delta R^2=0.26$) (**Fig. 3C**). This is driven by high rates of retrieval in control mothers during habituation. High rates of retrieval are associated with less efficient maternal care (Beach & Jaynes, 1956; Smart & Preece, 1973; Dunlap et al., 2020; Aguggia et al., 2013; Liu et al., 2006). Thus, the IN OXT mothers are likely more efficient at maintaining offspring care during this disruption. While mothers from both groups increased huddling behavior post-separation from pups, control mothers decreased retrieval/carrying behavior while IN OXT mothers maintained a consistent level of retrieval/carrying behavior. The net increase in maternal care for IN OXT mothers from

habituation to reunion ($F_{1,20}=6.6$, $p<0.02$, $\Delta R^2=0.26$) (**Fig. 3D**) is therefore largely due to changes in retrieval behavior.

Figure 3. Rapid effects of OXT on change in maternal and non-maternal behavior from habituation to reunion. *Maternal behaviors:* (A) There was non-significant trend for mothers given IN OXT having a shorter pup approach latency. (B) There were no treatment differences for maternal huddling. (C) Mothers given IN OXT showed a significantly greater decrease in retrieval/carrying behavior from the habituation to the reunion phase. (D) IN OXT had a net positive effect on total maternal care from the habituation to reunion phases. *Non-maternal behaviors:* (E) There were no treatment differences for change in autogrooming, (F) rearing, or (G) freezing. (H) There was no net change on non-maternal behaviors from the habituation to reunion. Correlation line collapsing across treatment groups. * $p<0.05$; # $p<0.10$.

In order to determine if IN OXT was acting on neural systems that were specific to maternal care, we also tested for main effects of IN OXT on measures of activity (autogrooming, rearing) and stress/anxiety (freezing). From habituation to reunion, there were no changes in autogrooming ($F_{1,20}=0.32$, $p=0.58$) (**Fig. 3E**), rearing ($F_{1,20}=1.23$, $p=0.28$) (**Fig. 3F**), or freezing ($F_{1,20}=0.03$, $p=0.85$) (**Fig. 3G**). When summing all activity and stress-related behaviors, there was no net effect on non-maternal behaviors from the habituation to reunion ($F_{1,20}=1.94$, $p=0.18$) (**Fig. 3H**). In this context, IN OXT is not influencing general, non-maternal behaviors in response to an offspring separation event. The individual means and SEMs for each behavior in each phase are also reported in **S. Table 1**.

S. Table 1. Means and SEMs of behaviors measured.

Correlations between maternal care and maternal USVs and maternal care and pup USVs

To determine whether maternal simple sweeps were associated with specific maternal and investigative behaviors during each of the three testing phases, we correlated maternal USVs, which were always simple sweeps, during each phase with the corresponding behavior while controlling for moderation by IN OXT administration. In the habituation phase, maternal

simple sweeps positively correlated with licking behavior ($F_{2,20}=12.04$, $p<0.007$, $\Delta R^2=0.34$) (**Fig. 4B**). There was also a nonsignificant trend for a correlation between maternal simple sweeps and huddling ($F_{2,20}=4.48$, $p=0.096$, $\Delta R^2=0.18$) (**Fig. 4A**), and no effect associated with retrieving/carrying ($F_{2,20}=0.44$, $p=0.51$, $\Delta R^2=0.02$) (**Fig. 4C**). There was also, however, a significant moderation by OXT in the relationship between maternal retrievals and maternal simple sweeps. During habituation, mothers given IN OXT carried/retrieved pups less than mothers given saline ($F_{2,20}=9.95$, $p<0.005$, $\Delta R^2=0.33$) (**Fig. 4C**) but note that overall IN OXT mothers had consistent retrieval levels across the test (**Fig. 2B**). In the separation phase, there was no correlation between time the mother spent at the mesh divider and maternal simple sweep USVs ($F_{2,20}=1.2$, $p=0.58$, $\Delta R^2=0.058$) (**Fig. 4D**). Other maternal behaviors could not be assessed because of the mesh divider between mothers and pups. In the reunion phase, maternal simple sweeps positively correlated with maternal retrievals/carrying ($F_{1,20}=7.65$, $p<0.037$, $\Delta R^2=0.26$) (**Fig. 4G**). Other maternal behaviors did not correlate with maternal simple sweeps in this phase: huddling ($F_{1,20}=0.48$, $p=0.99$, $\Delta R^2=0.02$) (**Fig. 4E**) and licking ($F_{1,20}=0.001$, $p=0.98$, $\Delta R^2=0.00$) (**Fig. 4F**). Thus, overall there were associations between maternal simple sweeps and maternal care, but the maternal behavior that correlated with maternal simple sweeps varied depending on context.

Fig. 4. Correlations between maternal simple sweep USVs and maternal care. (A-C) Habituation. (A) There was a nonsignificant trend correlation between maternal simple sweeps and huddling. **(B)** Maternal simple sweeps positively correlated with licking behavior. **(C)** There was no effect between maternal simple sweeps and retrieving/carrying. Mothers given IN OXT carried/retrieved pups less than mothers given saline. **(D) Separation.** There was no correlation between time mothers spent at the mesh divider and maternal simple sweep USVs. **(E-G) Reunion. (E)** Maternal simple sweeps did not correlate with huddling or **(F)** licking. **(G)** Maternal simple sweeps positively correlated with maternal retrievals. Correlation line collapsing across treatment groups. * $p<0.05$; # $p<0.10$.

Next, we correlated pup whine USVs with specific types of maternal behavior to see if these pup calls were associated with a specific maternal response. During the habituation phase, there was a significant positive correlation between pup whines and maternal huddling ($F_{1,20}=8.93$, $p<0.024$, $\Delta R^2=0.30$) (**Fig. 5A**), but not with maternal licking ($F_{1,20}=3.26$, $p=0.174$, $\Delta R^2=0.13$) (**Fig. 5B**) or retrieval/carrying behavior ($F_{1,20}=1.14$, $p=0.26$, $\Delta R^2=0.04$) (**Fig. 5C**). During the separation phase, pup whines did not correlate with time that the mother spent at the mesh gate ($F_{1,20}=0.89$, $p=0.36$, $\Delta R^2=0.04$) (**Fig. 5D**). Lastly, during the reunion phase, pup USVs positively correlated with retrieving/carrying ($F_{1,20}=9.94$, $p=0.016$, $\Delta R^2=0.34$) (**Fig. 5G**) but had no effect on either huddling ($F_{1,20}=0.05$, $p=0.823$, $\Delta R^2=0.003$) (**Fig. 5E**) or licking ($F_{1,20}=0.893$, $p=0.36$, $\Delta R^2=0.043$) (**Fig. 5F**). Across all correlations of USVs and maternal care, significant correlations occurred, but neither maternal simple sweep USVs nor pup whine USVs consistently correlated with a specific type of maternal care.

Fig. 5. Correlations between pup whine USVs and maternal care. (A-C) Habituation. (A) There was a significant positive correlation between pup whines and maternal huddling. **(B)** There was no correlation between pup whines and maternal licking or **(C)** pup whines and maternal retrieval/carrying behavior. **(D) Separation.** Pup whines did not correlate with time that the mother spent at the mesh gate. **(E-G) Reunion. (E)** There was no correlation between pup whines and maternal huddling or **(F)** pup whines and maternal licking. **(G)** Pup USVs positively correlated with maternal retrieving/carrying. * $p<0.05$.

Discussion

Maternal care and communication have lifelong consequences for offspring later in life (Champagne & Meaney, 2006; Champagne & Curley, 2009; Cutuli et al., 2015; Champagne & Curley, 2016). Therefore, it is important to elucidate the proximate hormonal mechanisms that increase maternal care and communication. OXT is known for its potent role in maternal physiology, neurophysiology, and social behavior, but whether OXT could rapidly change vocal production and behavior in mothers remained unknown. We aimed to fill these knowledge gaps

by testing maternal response and effects of IN OXT during a potentially challenging and stressful pup separation paradigm.

We predicted that maternal care activity would be associated with vocal production and that IN OXT would amplify this effect. We found support for this hypothesis as all mothers produced more simple sweep USVs per second during both the habituation phase and reunion phase when mothers had access to multisensory pup experience (including touch). In support of the social salience theory (Shamay-Tsoory & Abu-Akel, 2016; Kemp & Guastella, 2010; Egito et al., 2020; Bartz et al., 2011; Ramsey et al., 2019), IN OXT amplified the effect of the multisensory experience, leading to an increase above and beyond the control increase in the habituation phase (and a trend in reunion phase). To our knowledge, this is the first study reporting context-dependent IN OXT-moderated changes in USV production that were associated with physical access to a social stimulus (review by Marler & Monari, 2020).

There are several possible functions for simple sweep USV production in the context of maternal care. Increased simple sweep USVs during maternal care may result from a higher state of arousal that is regulated by the autonomic nervous system (Porges, 2009). This is supported by evidence in prairie voles, where vocal features covary with heart rate—longer vocalizations were associated with increased vagal tone and more calm behavior whereas shorter vocalizations were associated with decreased vagal tone and more anxious behavior (Stewart et al., 2015). Alternatively, increases in simple sweep USVs may be associated with a positive affective state (review by Simola & Granon, 2019). In rats, 50 kHz USVs (similar kHz as California mouse simple sweeps) have been associated with positive affective state (Portfors, 2007), and in California mice, simple sweeps typically occur during affiliative contexts (Pultorak et al., 2018). This adds to a growing body of literature that aims to elucidate California mouse call type with

function. Complex sweeps predict pair mate social behavior (Pultorak et al. 2017); syllable USVs are associated with aggression (shorter calls) (Rieger & Marler, 2018) and female preference (longer calls) (Pultorak et al., 2017); both syllable USVs (shorter calls) and barks are associated with increased aggression (Rieger et al., 2019); pup whines can also elicit paternal retrievals (Chary et al., 2015). As expected, we did not observe any syllable vocalizations or barks and only a handful of complex sweeps across all tests as these calls tend to be associated with aggression or adult interactions. Because IN OXT increased maternal simple sweep USVs, and in other studies, IN OXT has increased both vagal tone (Higa et al., 2002) and positive affective state (Dolen et al., 2013; Hung et al., 2017; Baracz & Cornish, 2013), we suggest that maternal simple sweeps are most likely to be associated with changes in affective state.

In contrast to mothers, we did not expect to see a treatment effect with regards to pup vocalizations because only the mothers were treated and the pups are likely deaf due to their ear canals not being open yet (Fox, 1965). As expected, we did not observe differences in pup vocalizations between pup with control mothers and IN OXT mothers. This suggests that effects of IN OXT on the mother do not directly and rapidly influence pup behavior. Overall, pups called the most when first placed into the new chamber with their mothers at the rate of 0.94 whines per second, and then called at a steady rate of 0.33 whines per second during separation and reunion. This suggests that pup vocal production does not vary by social contact in the same way as maternal vocalizations. Instead, pup vocal production may be based on the degree of the thermal challenge, as indicted by previous studies on pup USVs demonstrating pups increase USVs when separated from their nest and given thermal challenges (Blumberg & Alberts 1990, Blumberg & Sokoloff 2001).

We predicted a correlation between maternal simple sweeps and pup whines. We expected that the correlation between mother-pup USVs would be driven by the mother, as the pups' ear canals have not opened at PND 2-3, likely rendering them deaf (Fox, 1965). Mother and pup USVs did not correlate during habituation, possibly because pup whine USVs were highest during this phase (**Fig. 2B**) or possibly because the removal from the home cage at the start of the test disrupted the coordination between mothers and pups. There was also a nonsignificant trend for IN OXT to improve the correlation between mother and pup USVs. If the removal from the home cage at the start of the test disrupted the coordination between mothers and pups, IN OXT may be mediating this negative effect by increasing the salience of pup whine stimuli (Valtcheva & Froemke, 2019), allowing mothers to more effectively and efficiently cope with the challenge. During reunion, we found support for our initial prediction: maternal simple sweeps and pup whines positively correlated. Mothers may be more responsive to pup whines during the reunion phase because the pups have been without care for a longer period of time. An alternative explanation is that this synchrony occurred out of necessity because, with the second chamber open, the mice had double the space in the reunion phase compared to the habituation phase. In sheep and ewes, mother-offspring vocalizations have been shown to be important for recognition and location purposes (Sèbe et al., 2007; Searby & Jouventin, 2003) with young ewes only being able to distinguish their mother via low frequency calls but using high frequency calls to during vocal exchanges (Sèbe et al., 2010). Mother-offspring vocalizations that are contingent on each other's vocalizations have also been observed across cultures in humans (Bornstein et al., 2015). To our knowledge, this is the first reporting of correlations between maternal USVs and offspring USVs in an animal model.

We also wanted to explore whether maternal simple sweeps or pup whines are correlated with specific types of maternal behavior. During habituation, USVs from both mothers and pups were associated with greater maternal care but were not associated with the same maternal behavior; maternal simple sweeps positively correlated with maternal licking behavior and pup whines positively correlated with maternal huddling behavior. During reunion, we saw a different relationship between USVs and maternal behavior, suggesting that if pup calls and not maternal responsiveness are driving these correlations, the pup whines are not eliciting a specific type of maternal care. During reunion, both maternal simple sweeps and pup whines positively correlated with maternal retrieval/carrying behavior. This suggests that after a separation event, pup whines may drive maternal retrieval/carrying behavior, similar to the finding previously reported in fathers (Chary et al., 2015) and that maternal simple sweeps may be a reliable signal supporting maternal responsiveness. Studies in the literature show support for this effect being driven by maternal responsiveness (D'Amato et al., 2005; D'Amato & Populin, 1987) and other studies show that the effect can also be driven by the pups (Curry et al., 2013; Wöhr, et al., 2010).

Finally, based on the prosocial effects of OXT, we predicted that an acute dose of IN OXT would increase maternal care. Our results show that in the context of our paradigm, IN OXT maintains maternal care rather than overtly increasing it. Mothers given IN OXT showed consistent amount of retrievals pre- and post-separation while control mothers significantly decreased amount of retrievals performed post-separation, leading to a significant increase in total maternal care for IN OXT mothers. These findings are consistent with other studies in the literature (Keverne & Kendrick, 1992; Francis et al., 2002; Feldman et al., 2010; Bosch & Neumann, 2012; Rich et al., 2014) but highlight that OXT can also increase maternal behavior

within minutes of administration. We did not find that IN OXT led to a significant decrease in latency to approach pups after separation, but we found a non-significant trend. This supports previous findings in the literature that OXT has been associated with a reduction in the latency to retrieve or start maternal behavior (Pedersen et al., 2006; Liu et al., 2019) though several other studies have not reported an effect of OXT on latency to retrieve pups (Watarai et al., 2020; Rich et al., 2014; Macbeth et al., 2010). Notably, we did not find any simple effects of IN OXT on nonmaternal behaviors during the habituation, separation, or reunion phases. This suggests that IN OXT specifically influences maternal behavior and not general activity or anxiety in female California mice. If IN OXT is dampening the stress/anxiety response, it is specific to maternal anxiety. This is important to note because one hypothesis on the effects of IN OXT is that it primarily functions as anxiolytic agent across a variety of contexts and species (Yoshida et al., 2009; Waldherr & Neumann, 2007; de Oliveira et al., 2012). Our results suggest that in this context, IN OXT has a specific effect on maternal care behavior that is not explained by differences in the non-maternal activities measured in the IN OXT mothers.

In summary, these data are consistent with the idea that IN OXT rapidly and selectively increases maternal vocalizations and maternal care. This data also highlights the importance of social contact for normal communication and care and enhancement by IN OXT. Overall, we propose that higher levels of IN OXT in mothers function to increase efficiency and maintain maternal care, particularly during challenges.

Declaration of competing interest

We know of no conflicts of interest and state that no financial support for this work has been given that could have influenced its outcome.

Acknowledgments

We would like to thank NSF grant 1355163 for generously funding this research, undergraduate student support for their work during implementation of the experiment and quantification of behavior, and UW Madison animal care staff for their excellent care of the animals.

Aguggia JP, Suárez MM, Rivarola MA. Early maternal separation: neurobehavioral consequences in mother rats. *Behavioural Brain Research*. 2013 Jul 1;248:25-31.

Allin JT, Banks EM. Functional aspects of ultrasound production by infant albino rats (*Rattus norvegicus*). *Animal Behaviour*. 1972 Feb 1;20(1):175-85.

Arriaga G, Jarvis ED. Mouse vocal communication system: Are ultrasounds learned or innate?. *Brain and Language*. 2013 Jan 1;124(1):96-116.

Arnon S, Diamant C, Bauer S, Regev R, Sirota G, Litmanovitz I. Maternal singing during kangaroo care led to autonomic stability in preterm infants and reduced maternal anxiety. *Acta Paediatrica*. 2014 Oct;103(10):1039-44.

Bales KL, Perkeybile AM, Conley OG, Lee MH, Guoynes CD, Downing GM, Yun CR, Solomon M, Jacob S, Mendoza SP. Chronic intranasal oxytocin causes long-term impairments in partner preference formation in male prairie voles. *Biological Psychiatry*. 2013 Aug 1;74(3):180-8.

Bales KL, Solomon M, Jacob S, Crawley JN, Silverman JL, Larke RH, Sahagun E, Puhger KR, Pride MC, Mendoza SP. Long-term exposure to intranasal oxytocin in a mouse autism model. *Translational Psychiatry*. 2014 Nov;4(11):e480-.

Baracz SJ, Cornish JL. Oxytocin modulates dopamine-mediated reward in the rat subthalamic nucleus. *Hormones and Behavior*. 2013 Feb 1;63(2):370-5.

Bartz JA, Hollander E. The neuroscience of affiliation: forging links between basic and clinical research on neuropeptides and social behavior. *Hormones and Behavior*. 2006 Nov 1;50(4):518-28.

Bartz JA, Zaki J, Bolger N, Ochsner KN. Social effects of oxytocin in humans: context and person matter. *Trends in Cognitive Sciences*. 2011 Jul 1;15(7):301-9.

Beach FA, Jaynes J. Studies of Maternal Retrieving in Rats. Iii. Sensory cues involved in the lactating female's response to her young 1. *Behaviour*. 1956 Jan 1;10(1):104-24.

Bell MR. Comparing postnatal development of gonadal hormones and associated social behaviors in rats, mice, and humans. *Endocrinology*. 2018 Jul;159(7):2596-613.

Bester-Meredith JK, Burns JN, Conley MF, Mammarella GE, Ng ND. *Peromyscus* as a model system for understanding the regulation of maternal behavior. In *Seminars in Cell & Developmental Biology* 2017 Jan 1 (Vol. 61, pp. 99-106). Academic Press.

Bester-Meredith JK, Marler CA. The association between male offspring aggression and paternal and maternal behavior of *Peromyscus* mice. *Ethology*. 2003 Oct;109(10):797-808.

Blomquist GE. Maternal effects on offspring mortality in rhesus macaques (*Macaca mulatta*). *American journal of primatology*. 2013 Mar;75(3):238-51.

Blumberg MS, Alberts JR. Ultrasonic vocalizations by rat pups in the cold: an acoustic by-product of laryngeal braking?. *Behavioral Neuroscience*. 1990 Oct;104(5):808.

Blumberg MS, Sokoloff G. Do infant rats cry?. *Psychological Review*. 2001 Jan;108(1):83.

Bornstein MH, Putnick DL, Cote LR, Haynes OM, Suwalsky JT. Mother-infant contingent vocalizations in 11 countries. *Psychological Science*. 2015 Aug;26(8):1272-84.

Bosch OJ, Neumann ID. Both oxytocin and vasopressin are mediators of maternal care and aggression in rodents: from central release to sites of action. *Hormones and Behavior*. 2012 Mar 1;61(3):293-303.

Brewster J, Leon M. Relocation of the site of mother-young contact: Maternal transport behavior in Norway rats. *Journal of Comparative and Physiological Psychology*. 1980 Feb;94(1):69.

Briggs JR, Kalcounis-Rueppell MC. Similar acoustic structure and behavioural context of vocalizations produced by male and female California mice in the wild. *Animal Behaviour*. 2011 Dec 1;82(6):1263-73.

Champagne FA, Curley JP. Epigenetic mechanisms mediating the long-term effects of maternal care on development. *Neuroscience & Biobehavioral Reviews*. 2009 Apr 1;33(4):593-600.

Champagne FA, Curley JP. Plasticity of the maternal brain across the lifespan. *New Directions for Child and Adolescent Development*. 2016 Sep;2016(153):9-21.

Champagne F, Diorio J, Sharma S, Meaney MJ. Naturally occurring variations in maternal behavior in the rat are associated with differences in estrogen-inducible central oxytocin receptors. *Proceedings of the National Academy of Sciences*. 2001 Oct 23;98(22):12736-41.

Champagne FA, Meaney MJ. Stress during gestation alters postpartum maternal care and the development of the offspring in a rodent model. *Biological Psychiatry*. 2006 Jun 15;59(12):1227-35.

Chard T, Boyd N, Forsling ML, McNeilly AS, Landon J. The development of a radioimmunoassay for oxytocin: the extraction of oxytocin from plasma, and its measurement during parturition in human and goat blood. *Journal of Endocrinology*. 1970 Oct 1;48(2):223-34.

Chary MC, Cruz JP, Bardi M, Becker EA. Paternal retrievals increase testosterone levels in both male and female California mouse (*Peromyscus californicus*) offspring. *Hormones and Behavior*. 2015 Jul 1;73:23-9.

Chauke M, Malisch JL, Robinson C, de Jong TR, Saltzman W. Effects of reproductive status on behavioral and endocrine responses to acute stress in a biparental rodent, the California mouse (*Peromyscus californicus*). *Hormones and Behavior*. 2011 Jun 1;60(1):128-38.

Cheng MF. For whom does the female dove coo? A case for the role of vocal self-stimulation. *Animal Behaviour*. 1992 Jun 1;43(6):1035-44.

Churchland PS, Winkielman P. Modulating social behavior with oxytocin: how does it work? What does it mean?. *Hormones and Behavior*. 2012 Mar 1;61(3):392-9.

Curley JP, Champagne FA. Influence of maternal care on the developing brain: Mechanisms, temporal dynamics and sensitive periods. *Frontiers in Neuroendocrinology*. 2016 Jan 1;40:52-66.

Curry T, Egeto P, Wang H, Podnos A, Wasserman D, Yeomans J. Dopamine receptor D2 deficiency reduces mouse pup ultrasonic vocalizations and maternal responsiveness. *Genes, Brain and Behavior*. 2013 Jun;12(4):397-404.

Cutuli D, Caporali P, Gelfo F, Angelucci F, Laricchiuta D, Foti F, De Bartolo P, Bisicchia E, Molinari M, Farioli Vecchioli S, Petrosini L. Pre-reproductive maternal enrichment influences rat maternal care and offspring developmental trajectories: behavioral performances and neuroplasticity correlates. *Frontiers in Behavioral Neuroscience*. 2015 Mar 12;9:66.

D'Amato FR, Populin R. Mother-offspring interaction and pup development in genetically deaf mice. *Behavior Genetics*. 1987 Sep 1;17(5):465-75.

D'Amato FR, Scalera E, Sarli C, Moles A. Pups call, mothers rush: does maternal responsiveness affect the amount of ultrasonic vocalizations in mouse pups?. *Behavior genetics*. 2005 Jan 1;35(1):103-12.

Davis EP, Stout SA, Molet J, Vegetabile B, Glynn LM, Sandman CA, Heins K, Stern H, Baram TZ. Exposure to unpredictable maternal sensory signals influences cognitive development across species. *Proceedings of the National Academy of Sciences*. 2017 Sep 26;114(39):10390-5.

de Oliveira DC, Zuairi AW, Graeff FG, Queiroz RH, Crippa JA. Anxiolytic-like effect of oxytocin in the simulated public speaking test. *Journal of Psychopharmacology*. 2012 Apr;26(4):497-504.

- Dölen G, Darvishzadeh A, Huang KW, Malenka RC. Social reward requires coordinated activity of nucleus accumbens oxytocin and serotonin. *Nature*. 2013 Sep;501(7466):179-84.
- Dunlap AG, Besosa C, Pascual LM, Chong KK, Walum H, Kacsoh DB, Tankeu BB, Lu K, Liu RC. Becoming a better parent: Mice learn sounds that improve a stereotyped maternal behavior. *Hormones and Behavior*. 2020 Aug 1;124:104779.
- Egito JH, Nevat M, Shamay-Tsoory SG, Osório AA. Oxytocin increases the social salience of the outgroup in potential threat contexts. *Hormones and Behavior*. 2020 Jun 1;122:104733.
- Fairbanks LA, McGuire MT. Maternal condition and the quality of maternal care in vervet monkeys. *Behaviour*. 1995 Jan 1;132(9-10):733-54.
- Feldman R, Gordon I, Schneiderman I, Weisman O, Zagoory-Sharon O. Natural variations in maternal and paternal care are associated with systematic changes in oxytocin following parent–infant contact. *Psychoneuroendocrinology*. 2010 Sep 1;35(8):1133-41.
- Feldman R, Weller A, Zagoory-Sharon O, Levine A. Evidence for a neuroendocrinological foundation of human affiliation: plasma oxytocin levels across pregnancy and the postpartum period predict mother-infant bonding. *Psychological Science*. 2007 Nov;18(11):965-70.
- Fox WM. Reflex-ontogeny and behavioural development of the mouse. *Animal Behaviour*. 1965 Apr 1;13(2-3):234-IN5.
- Francis DD, Champagne FC, Meaney MJ. Variations in maternal behaviour are associated with differences in oxytocin receptor levels in the rat. *Journal of Neuroendocrinology*. 2000 Dec;12(12):1145-8.
- Francis D, Diorio J, Liu D, Meaney MJ. Nongenomic transmission across generations of maternal behavior and stress responses in the rat. *Science*. 1999 Nov 5;286(5442):1155-8.
- Francis DD, Young LJ, Meaney MJ, Insel TR. Naturally occurring differences in maternal care are associated with the expression of oxytocin and vasopressin (V1a) receptors: gender differences. *Journal of Neuroendocrinology*. 2002 May;14(5):349-53.
- Freeman SM, Samineni S, Allen PC, Stockinger D, Bales KL, Hwa GG, Roberts JA. Plasma and CSF oxytocin levels after intranasal and intravenous oxytocin in awake macaques. *Psychoneuroendocrinology*. 2016 Apr 1;66:185-94.
- Fuchs AR. Oxytocin and the onset of labour in rabbits. *Journal of Endocrinology*. 1964 Sep 1;30(2):217-24.
- Gossen A, Hahn A, Westphal L, Prinz S, Schultz RT, Gründer G, Spreckelmeyer KN. Oxytocin plasma concentrations after single intranasal oxytocin administration—a study in healthy men. *Neuropeptides*. 2012 Oct 1;46(5):211-5.

Guoynes C, Marler C. Paternal Behavior from a Neuroendocrine Perspective. In Oxford Research Encyclopedia of Neuroscience 2020 Mar 31.

Guoynes CD, Simmons TC, Downing GM, Jacob S, Solomon M, Bales KL. Chronic intranasal oxytocin has dose-dependent effects on central oxytocin and vasopressin systems in prairie voles (*Microtus ochrogaster*). *Neuroscience*. 2018 Jan 15;369:292-302.

Harris BN, Saltzman W. Effects of aging on hypothalamic-pituitary-adrenal (HPA) axis activity and reactivity in virgin male and female California mice (*Peromyscus californicus*). *General and Comparative Endocrinology*. 2013 Jun 1;186:41-9.

Hennessy MB, Li J, Lowe EL, Levine S. Maternal behavior, pup vocalizations, and pup temperature changes following handling in mice of 2 inbred strains. *Developmental Psychobiology: The Journal of the International Society for Developmental Psychobiology*. 1980 Nov;13(6):573-84.

Higa KT, Mori E, Viana FF, Morris M, Michelini LC. Baroreflex control of heart rate by oxytocin in the solitary-vagal complex. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2002 Feb 1;282(2):R537-45.

Hung LW, Neuner S, Polepalli JS, Beier KT, Wright M, Walsh JJ, Lewis EM, Luo L, Deisseroth K, Dölen G, Malenka RC. Gating of social reward by oxytocin in the ventral tegmental area. *Science*. 2017 Sep 29;357(6358):1406-11.

Johnson SA, Javurek AB, Painter MS, Peritore MP, Ellersieck MR, Roberts RM, Rosenfeld CS. Disruption of parenting behaviors in California Mice, a monogamous rodent species, by endocrine disrupting chemicals. *PloS One*. 2015 Jun 3;10(6):e0126284.

Johnson SA, Painter MS, Javurek AB, Murphy CR, Howald EC, Khan ZZ, Conard CM, Gant KL, Ellersieck MR, Hoffmann F, Schenk AK. Characterization of vocalizations emitted in isolation by California mouse (*Peromyscus californicus*) pups throughout the postnatal period. *Journal of Comparative Psychology*. 2017 Feb;131(1):30.

Kalcounis-Rueppell MC, Metheny JD, Vonhof MJ. Production of ultrasonic vocalizations by *Peromyscus* mice in the wild. *Frontiers in Zoology*. 2006 Dec 1;3(1):3.

Kalcounis-Rueppell MC, Petric R, Briggs JR, Carney C, Marshall MM, Willse JT, Rueppell O, Ribble DO, Crossland JP. Differences in ultrasonic vocalizations between wild and laboratory California mice (*Peromyscus californicus*). *PloS One*. 2010 Apr 1;5(4):e9705.

Kalcounis-Rueppell MC, Pultorak JD, Blake BH, Marler CA. Ultrasonic vocalizations of young mice in the genus *Peromyscus*. In *Handbook of Behavioral Neuroscience* 2018 Jan 1 (Vol. 25, pp. 149-156). Elsevier.

- Kalcounis-Rueppell MC, Pultorak JD, Marler CA. Ultrasonic vocalizations of mice in the genus *Peromyscus*. In *Handbook of Behavioral Neuroscience* 2018 Jan 1 (Vol. 25, pp. 227-235). Elsevier.
- Kemp AH, Guastella AJ. Oxytocin: prosocial behavior, social salience, or approach-related behavior?. *Biological Psychiatry*. 2010 Mar 15;67(6):e33-4.
- Keverne EB, Kendrick KM. Oxytocin Facilitation of Maternal Behavior in Sheep a. *Annals of the New York Academy of Sciences*. 1992 Jun;652(1):83-101.
- Knutson B, Burgdorf J, Panksepp J. Ultrasonic vocalizations as indices of affective states in rats. *Psychological Bulletin*. 2002 Nov;128(6):961.
- Lee MR, Scheidweiler KB, Diao XX, Akhlaghi F, Cummins A, Huestis MA, Leggio L, Averbeck BB. Oxytocin by intranasal and intravenous routes reaches the cerebrospinal fluid in rhesus macaques: determination using a novel oxytocin assay. *Molecular Psychiatry*. 2018 Jan;23(1):115-22.
- Lee MR, Shnitko TA, Blue SW, Kaucher AV, Winchell AJ, Erikson DW, Grant KA, Leggio L. Labeled oxytocin administered via the intranasal route reaches the brain in rhesus macaques. *Nature Communications*. 2020 Jun 3;11(1):1-0.
- Leng G, Ludwig M. Intranasal oxytocin: myths and delusions. *Biological Psychiatry*. 2016 Feb 1;79(3):243-50.
- Lerch-Haner JK, Frierson D, Crawford LK, Beck SG, Deneris ES. Serotonergic transcriptional programming determines maternal behavior and offspring survival. *Nature Neuroscience*. 2008 Sep;11(9):1001.
- Lim MM, Young LJ. Neuropeptidergic regulation of affiliative behavior and social bonding in animals. *Hormones and Behavior*. 2006 Nov 1;50(4):506-17.
- Liu RC, Linden JF, Schreiner CE. Improved cortical entrainment to infant communication calls in mothers compared with virgin mice. *European Journal of Neuroscience*. 2006 Jun;23(11):3087-97.
- Liu XY, Li D, Li T, Liu H, Cui D, Liu Y, Jia S, Wang X, Jiao R, Zhu H, Zhang F. Effects of Intranasal Oxytocin on Pup Deprivation-Evoked Aberrant Maternal Behavior and Hypogalactia in Rat Dams and the Underlying Mechanisms. *Frontiers in Neuroscience*. 2019 Feb 26;13:122.
- Macbeth AH, Stepp JE, Lee HJ, Young III WS, Caldwell HK. Normal maternal behavior, but increased pup mortality, in conditional oxytocin receptor knockout females. *Behavioral Neuroscience*. 2010 Oct;124(5):677.

- Malisch JL, Saltzman W, Gomes FR, Rezende EL, Jeske DR, Garland Jr T. Baseline and stress-induced plasma corticosterone concentrations of mice selectively bred for high voluntary wheel running. *Physiological and Biochemical Zoology*. 2007 Jan;80(1):146-56.
- Marlin BJ, Mitre M, D'amour JA, Chao MV, Froemke RC. Oxytocin enables maternal behaviour by balancing cortical inhibition. *Nature*. 2015 Apr;520(7548):499-504.
- Maynard KR, Hobbs JW, Phan BN, Gupta A, Rajpurohit S, Williams C, Rajpurohit A, Shin JH, Jaffe AE, Martinowich K. BDNF-TrkB signaling in oxytocin neurons contributes to maternal behavior. *Elife*. 2018 Sep 7;7:e33676.
- Meites J, Hopkins TF. Mechanism of action of oxytocin in retarding mammary involution: study in hypophysectomized rats. *Journal of Endocrinology*. 1961 Jun 1;22(3):207-13.
- Mogi K, Takakuda A, Tsukamoto C, Ooyama R, Okabe S, Koshida N, Nagasawa M, Kikusui T. Mutual mother-infant recognition in mice: The role of pup ultrasonic vocalizations. *Behavioural Brain Research*. 2017 May 15;325:138-46.
- Moltz H, Lubin M, Leon M, Numan M. Hormonal induction of maternal behavior in the ovariectomized nulliparous rat. *Physiology & Behavior*. 1970 Dec 1;5(12):1373-7.
- Musolf K, Hoffmann F, Penn DJ. Ultrasonic courtship vocalizations in wild house mice, *Mus musculus musculus*. *Animal Behaviour*. 2010 Mar 1;79(3):757-64.
- Nagasawa M, Okabe S, Mogi K, Kikusui T. Oxytocin and mutual communication in mother-infant bonding. *Frontiers in Human Neuroscience*. 2012 Feb 28;6:31.
- Neumann I, Douglas AJ, Pittman QJ, Russell JA, Landgraf R. Oxytocin released within the supraoptic nucleus of the rat brain by positive feedback action is involved in parturition-related events. *Journal of Neuroendocrinology*. 1996 Mar;8(3):227-33.
- Neumann IN, Koehler EL, Landgraf RA, Summy-Long JO. An oxytocin receptor antagonist infused into the supraoptic nucleus attenuates intranuclear and peripheral release of oxytocin during suckling in conscious rats. *Endocrinology*. 1994 Jan 1;134(1):141-8.
- Neumann ID, Maloumy R, Beiderbeck DI, Lukas M, Landgraf R. Increased brain and plasma oxytocin after nasal and peripheral administration in rats and mice. *Psychoneuroendocrinology*. 2013 Oct 1;38(10):1985-93.
- Nowak R, Porter RH, Lévy F, Orgeur P, Schaal B. Role of mother-young interactions in the survival of offspring in domestic mammals. *Reviews of Reproduction*. 2000 Sep 1;5(3):153-63.
- Oppong-Damoah A, Zaman RU, D'Souza MJ, Murnane KS. Nanoparticle encapsulation increases the brain penetrance and duration of action of intranasal oxytocin. *Hormones and Behavior*. 2019 Feb 1;108:20-9.

Pedersen CA, Ascher JA, Monroe YL, Prange AJ. Oxytocin induces maternal behavior in virgin female rats. *Science*. 1982 May 7;216(4546):648-50.

Pedersen CA, Prange AJ. Induction of maternal behavior in virgin rats after intracerebroventricular administration of oxytocin. *Proceedings of the National Academy of Sciences*. 1979 Dec 1;76(12):6661-5.

Pedersen CA, Vadlamudi SV, Boccia ML, Amico JA. Maternal behavior deficits in nulliparous oxytocin knockout mice. *Genes, Brain and Behavior*. 2006 Apr;5(3):274-81.

Porges SW. The polyvagal theory: new insights into adaptive reactions of the autonomic nervous system. *Cleveland Clinic Journal of Medicine*. 2009 Apr;76(Suppl 2):S86.

Portfors CV. Types and functions of ultrasonic vocalizations in laboratory rats and mice. *Journal of the American Association for Laboratory Animal Science*. 2007 Jan 1;46(1):28-34.

Pultorak JD, Alger SJ, Loria SO, Johnson AM, Marler CA. Changes in behavior and ultrasonic vocalizations during pair bonding and in response to an infidelity challenge in monogamous California mice. *Frontiers in Ecology and Evolution*. 2018 Aug 28;6:125.

Pultorak JD, Fuxjager MJ, Kalcounis-Rueppell MC, Marler CA. Male fidelity expressed through rapid testosterone suppression of ultrasonic vocalizations to novel females in the monogamous California mouse. *Hormones and Behavior*. 2015 Apr 1;70:47-56.

Pultorak JD, Matusinec KR, Miller ZK, Marler CA. Ultrasonic vocalization production and playback predicts intrapair and extrapair social behaviour in a monogamous mouse. *Animal Behaviour*. 2017 Mar 1;125:13-23.

Quintana DS, Alvares GA, Hickie IB, Guastella AJ. Do delivery routes of intranasally administered oxytocin account for observed effects on social cognition and behavior? A two-level model. *Neuroscience & Biobehavioral Reviews*. 2015 Feb 1;49:182-92.

Ramsey ME, Fry D, Cummings ME. Isotocin increases female avoidance of males in a coercive mating system: assessing the social salience hypothesis of oxytocin in a fish species. *Hormones and Behavior*. 2019 Jun 1;112:1-9.

Rich ME, deCárdenas EJ, Lee HJ, Caldwell HK. Impairments in the initiation of maternal behavior in oxytocin receptor knockout mice. *PloS One*. 2014 Jun 3;9(6):e98839.

Rieger NS, Marler CA. The function of ultrasonic vocalizations during territorial defence by pair-bonded male and female California mice. *Animal Behaviour*. 2018 Jan 1;135:97-108.

Rieger NS, Stanton EH, Marler CA. Division of labour in territorial defence and pup retrieval by pair-bonded California mice, *Peromyscus californicus*. *Animal Behaviour*. 2019 Oct 1;156:67-78.

Rosenblatt JS. The development of maternal responsiveness in the rat. *American Journal of Orthopsychiatry*. 1969 Jan;39(1):36.

Rossoni E, Feng J, Tirozzi B, Brown D, Leng G, Moos F. Emergent synchronous bursting of oxytocin neuronal network. *PLoS Comput Biol*. 2008 Jul 18;4(7):e1000123.

Scattoni ML, Crawley J, Ricceri L. Ultrasonic vocalizations: a tool for behavioural phenotyping of mouse models of neurodevelopmental disorders. *Neuroscience & Biobehavioral Reviews*. 2009 Apr 1;33(4):508-15.

Schiavo JK, Valtcheva S, Bair-Marshall C, Song SC, Martin KA, Froemke RC. Innate sensitivity and plastic mechanisms in auditory cortex for reliable maternal behavior. *bioRxiv*. 2020 Jan 1.

Scott N, Prigge M, Yizhar O, Kimchi T. A sexually dimorphic hypothalamic circuit controls maternal care and oxytocin secretion. *Nature*. 2015 Sep;525(7570):519-22.

Searby A, Jouventin P. Mother-lamb acoustic recognition in sheep: a frequency coding. *Proceedings of the Royal Society of London. Series B: Biological Sciences*. 2003 Sep 7;270(1526):1765-71.

Sèbe F, Duboscq J, Aubin T, Ligout S, Poindron P. Early vocal recognition of mother by lambs: contribution of low-and high-frequency vocalizations. *Animal Behaviour*. 2010 May 1;79(5):1055-66.

Sebe F, Nowak R, Poindron P, Aubin T. Establishment of vocal communication and discrimination between ewes and their lamb in the first two days after parturition. *Developmental Psychobiology: The Journal of the International Society for Developmental Psychobiology*. 2007 May;49(4):375-86.

Shamay-Tsoory SG, Abu-Akel A. The social salience hypothesis of oxytocin. *Biological Psychiatry*. 2016 Feb 1;79(3):194-202.

Siegel HI, Rosenblatt JS. Estrogen-induced maternal behavior in hysterectomized-ovariectomized virgin rats. *Physiology & Behavior*. 1975 Apr 1;14(4):465-71.

Simola N, Granon S. Ultrasonic vocalizations as a tool in studying emotional states in rodent models of social behavior and brain disease. *Neuropharmacology*. 2019 Nov 15;159:107420.

Simola N, Fenu S, Costa G, Pinna A, Plumitallo A, Morelli M. Pharmacological characterization of 50-kHz ultrasonic vocalizations in rats: comparison of the effects of different psychoactive drugs and relevance in drug-induced reward. *Neuropharmacology*. 2012 Aug 1;63(2):224-34.

Simola N, Granon S. Ultrasonic vocalizations as a tool in studying emotional states in rodent models of social behavior and brain disease. *Neuropharmacology*. 2019 Nov 15;159:107420.

Smart JL, Preece J. Maternal behaviour of undernourished mother rats. *Animal Behaviour*. 1973 Aug 1;21(3):613-9.

Steinman MQ, Duque-Wilckens N, Greenberg GD, Hao R, Campi KL, Laredo SA, Laman-Maharg A, Manning CE, Doig IE, Lopez EM, Walch K. Sex-specific effects of stress on oxytocin neurons correspond with responses to intranasal oxytocin. *Biological Psychiatry*. 2016 Sep 1;80(5):406-14.

Stewart AM, Lewis GF, Yee JR, Kenkel WM, Davila MI, Carter CS, Porges SW. Acoustic features of prairie vole (*Microtus ochrogaster*) ultrasonic vocalizations covary with heart rate. *Physiology & Behavior*. 2015 Jan 1;138:94-100.

Striepens N, Kendrick KM, Hanking V, Landgraf R, Wüllner U, Maier W, Hurlemann R. Elevated cerebrospinal fluid and blood concentrations of oxytocin following its intranasal administration in humans. *Scientific Reports*. 2013 Dec 6;3:3440.

Stern JM. Maternal behavior: Sensory, hormonal, and neural determinants. In *Psychoendocrinology* 1989 Jan 1 (pp. 105-226). Academic Press.

Takayanagi Y, Yoshida M, Bielsky IF, Ross HE, Kawamata M, Onaka T, Yanagisawa T, Kimura T, Matzuk MM, Young LJ, Nishimori K. Pervasive social deficits, but normal parturition, in oxytocin receptor-deficient mice. *Proceedings of the National Academy of Sciences*. 2005 Nov 1;102(44):16096-101.

Talegaonkar S, Mishra PR. Intranasal delivery: An approach to bypass the blood brain barrier. *Indian Journal of Pharmacology*. 2004 May 1;36(3):140.

Taylor JH, French JA. Oxytocin and vasopressin enhance responsiveness to infant stimuli in adult marmosets. *Hormones and Behavior*. 2015 Sep 1;75:154-9.

Tops M, Van IJzendoorn MH, Riem MM, Boksem MA, Bakermans-Kranenburg MJ. Oxytocin receptor gene associated with the efficiency of social auditory processing. *Frontiers in Psychiatry*. 2011 Nov 4;2:60.

Valtcheva S, Froemke RC. Neuromodulation of maternal circuits by oxytocin. *Cell and Tissue Research*. 2019 Jan 28;375(1):57-68.

Van Leengoed E, Kerker E, Swanson HH. Inhibition of post-partum maternal behaviour in the rat by injecting an oxytocin antagonist into the cerebral ventricles. *Journal of Endocrinology*. 1987 Feb 1;112(2):275-82.

Waldherr M, Neumann ID. Centrally released oxytocin mediates mating-induced anxiolysis in male rats. *Proceedings of the National Academy of Sciences*. 2007 Oct 16;104(42):16681-4.

- 1004 Watarai A, Tsutaki S, Nishimori K, Okuyama T, Mogi K, Kikusui T. The blockade of oxytocin
- 1005 receptors in the paraventricular thalamus reduces maternal crouching behavior over pups in
- 1006 lactating mice. *Neuroscience Letters*. 2020 Feb 16;720:134761.
- 1007
- 1008 White NR, Adox R, Reddy A, Barfield RJ. Regulation of rat maternal behavior by broadband
- 1009 pup vocalizations. *Behavioral and Neural Biology*. 1992 Sep 1;58(2):131-7.
- 1010
- 1011 Winslow JT, Insel TR. The social deficits of the oxytocin knockout mouse. *Neuropeptides*. 2002
- 1012 Apr 1;36(2-3):221-9.
- 1013
- 1014 Wöhr M, Oddi D, D'Amato FR. Effect of altricial pup ultrasonic vocalization on maternal
- 1015 behavior. In *Handbook of Behavioral Neuroscience* 2010 Jan 1 (Vol. 19, pp. 159-166). Elsevier.
- 1016
- 1017 Yoshida M, Takayanagi Y, Inoue K, Kimura T, Young LJ, Onaka T, Nishimori K. Evidence that
- 1018 oxytocin exerts anxiolytic effects via oxytocin receptor expressed in serotonergic neurons in
- 1019 mice. *Journal of Neuroscience*. 2009 Feb 18;29(7):2259-71.
- 1020
- 1021 Zhao M, Harris BN, Nguyen CT, Saltzman W. Effects of single parenthood on mothers'
- 1022 behavior, morphology, and endocrine function in the biparental California mouse. *Hormones and*
- 1023 *Behavior*. 2019 Aug 1;114:104536.
- 1024

(A)

(B)

bioRxiv preprint doi: <https://doi.org/10.1101/2020.12.03.409557>; this version posted December 3, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Figure 1

Figure 2

bioRxiv preprint doi: <https://doi.org/10.1101/2020.12.03.409557>; this version posted December 3, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Figure 3

bioRxiv preprint doi: <https://doi.org/10.1101/2020.12.03.409557>; this version posted December 3, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Figure 4

bioRxiv preprint doi: <https://doi.org/10.1101/2020.12.03.409557>; this version posted December 3, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Figure 5

Behavior	Maternal/Non-maternal	Description
<i>Huddling</i>	Maternal	Amount of time that the mother that the mother spends over the pups. At least 50% of the body of pup must be covered by the mother to count.
<i>Licking and grooming</i>	Maternal	Amount of time that the mother that the mother spends licking and grooming pups.
<i>Retrieving/carrying</i>	Maternal	Amount of time that the mother with a pup in her mouth while locomoting in the chamber. Is counted regardless of whether locomotion is toward or away from home bedding and cotton ball.
<i>Time spent at mesh</i>	Maternal	Amount of time that the mother spends in front of (within a whisker's length distance) or climbing on the mesh door that partitions her chamber and the pup chamber.
<i>Latency to approach pups</i>	Maternal	Amount of time that it takes the mother to first reach her pups after the partitioned mesh door is opened.
<i>Autogrooming</i>	Non-maternal	Licking, biting, or scratching their own body, at any location.
<i>Freezing</i>	Non-maternal	When the mouse does not move at all but has eyes alert/open. Usually their ears are up, and they do not make any other movements. Considered a second event once the mouse moves for at least 2 seconds and before freezing again.
<i>Rearing</i>	Non-maternal	Both front paws lift up towards the wall of the cage. Counted as an additional event if front paws touch the ground again and lift back up.

S. Figure 1. Ethogram with description of behaviors measured.

Behavior	Phase	Treatment	Mean	SEM
<i>Huddling</i>	Habituation	CTRL	77.50	± 10.69
		OXT	101.33	± 13.23
	Reunion	CTRL	92.22	± 13.78
		OXT	142.67	± 22.57
<i>Licking and grooming</i>	Habituation	CTRL	46.83	± 5.95
		OXT	62.76	± 7.50
	Reunion	CTRL	60.30	± 8.08
		OXT	84.67	± 13.43
<i>Retrieving/carrying</i>	Habituation	CTRL	61.00	± 15.69
		OXT	21.33	± 6.73
	Reunion	CTRL	17.44	± 9.00
		OXT	14.33	± 6.00
<i>Time spent at mesh</i>	Separation	CTRL	31.52	± 3.25
		OXT	29.48	± 3.80
	Reunion	CTRL	38.00	± 12.96
		OXT	14.42	± 2.58
<i>Autogrooming</i>	Habituation	CTRL	7.81	± 1.84
		OXT	7.72	± 1.54
	Separation	CTRL	0.41	± 0.30
		OXT	2.34	± 1.83
	Reunion	CTRL	12.81	± 6.51
		OXT	9.60	± 2.49
	Habituation	CTRL	0.11	± 0.11
		OXT	0.71	± 0.58
<i>Freezing</i>	Separation	CTRL	1.26	± 0.87
		OXT	0.62	± 0.39
	Reunion	CTRL	9.64	± 6.20
		OXT	8.27	± 8.09
	Habituation	CTRL	52.44	± 4.48
		OXT	48.33	± 5.74
<i>Rearing</i>	Separation	CTRL	45.89	± 4.30
		OXT	42.96	± 5.04
	Reunion	CTRL	37.61	± 7.27
		OXT	27.96	± 7.67

S. Table 1. Means and SEMs of behaviors measured.