1 **A protease-mediated mechanism regulates the cytochrome c_6 / plastocyanin**
2 **switch in *Synechocystis* sp. PCC 6803**

3

4 Raquel García-Cañas^{1,2}, Joaquín Giner-Lamia^{1,3}, Francisco J. Florencio^{1,2} and
5 Luis López-Maury^{*1,2}.

6

7 1. Instituto de Bioquímica Vegetal y Fotosíntesis, Universidad de Sevilla- CSIC,
8 c/Américo Vespucio 49, 41092 Sevilla, Spain.

9

10 2. Departamento de Bioquímica Vegetal y Biología Molecular, Facultad de
11 Biología, Universidad de Sevilla, Avenida Reina Mercedes s/n, 41012 Sevilla,
12 Spain

13

14 3. Present address: Centro de Biotecnología y Genómica de Plantas UPM – INIA
15 Parque Científico y Tecnológico de la U.P.M. Campus de Montegancedo,
16 Autopista M-40, Km 38, Pozuelo de Alarcón, Madrid 28223, Spain

17

18 * Luis López-Maury

19 **Email:** llopez1@us.es

20 **Tel:** +34 954489501 ext. 909103

21 Raquel García-Cañas, ORCID 0000-0003-0062-6620; Joaquín Giner-Lamia,
22 ORCID 0000-0003-1553-8295; Francisco J. Florencio, ORCID 0000-0002-2068-
23 7861; Luis López-Maury, ORCID 0000-0002-3510-0621

24 **Classification**

25 Biological Sciences/Plant Biology

26 **Keywords**

27 Cyanobacteria, plastocyanin, cytochrome c_6

28 **Author Contributions**

29 Author contributions: R.M.G.C., J.G.L., F.J.F. and L.L.M. designed the research;
30 R.M.G.C., J.G.L. and L.L.M. performed the research; R.M.G.C., J.G.L., F.J.F.
31 and L.L.M. analyzed the data; L.L.M. wrote the manuscript with contributions
32 from all authors.

33

Abstract

After the Great Oxidation Event (GOE), iron availability was greatly decreased and photosynthetic organisms evolved several alternative proteins and mechanisms. One of these proteins, plastocyanin, is a type I blue-copper protein that can replace cytochrome c_6 as a soluble electron carrier between cytochrome b_6f and photosystem I. In most cyanobacteria, expression of these two alternative proteins is regulated by copper availability, but the regulatory system remains unknown. Herein, we provide evidence that the regulatory system is composed of a Blal/CopY family transcription factor (PetR) and a BlaR membrane protease (PetP). PetR represses *petE* (plastocyanin) expression and activates *petJ* (cytochrome c_6), while PetP controls PetR levels *in vivo*. Using whole-cell extracts, we demonstrated that PetR degradation requires both PetP and copper. Transcriptomic analysis revealed that the PetRP system regulates only four genes (*petE*, *petJ*, *slr0601*, and *slr0602*), highlighting its specificity. Furthermore, the presence of *petE* and *petRP* in early branching cyanobacteria indicates that acquisition of these genes could represent an early adaptation to decreased iron bioavailability following the GOE.

Significance Statement

After the appearance of oxygenic photosynthesis, Fe became oxidized and its solubility and availability were greatly decreased. This generated a problem for

55 most organisms since they are strongly dependent on Fe, especially
 56 photosynthetic organisms. In response, organisms evolved alternatives to Fe-
 57 containing proteins such as plastocyanin, a copper protein that substitutes for
 58 cytochrome c_6 in photosynthesis. Expression of these two proteins in
 59 cyanobacteria is regulated by Cu availability, but the regulatory system remains
 60 unknown. Herein, we describe the regulatory system for these alternative
 61 proteins in photosynthesis in cyanobacteria. The mechanism involves a
 62 transcription factor (PetR) and a membrane protease (PetP) that degrades PetR
 63 in the presence of Cu.

Main Text

Introduction

Cyanobacteria are prokaryotes that perform oxygenic photosynthesis, and their appearance on earth led to a dramatic increase in oxygen in the atmosphere. Although the timing of the Great Oxygenation Event (GOE) is well established, its extent is still a matter of dispute (1–4). The GOE caused a massive extinction because oxygen was toxic to most life forms (5, 6) and provoked Fe oxidation (Fe^{2+} to Fe^{3+}), which made Fe much less available since Fe^{3+} precipitates. This created an additional problem for all forms of life because Fe was already established as a major cofactor in biochemical reactions (7, 8). In fact, photosynthesis has high Fe requirements and photosynthetic organisms usually have a higher Fe quota than non-photosynthetic organisms because Fe is contained in proteins of both photosystems, in the cytochrome b_6f complex, in the soluble thylakoid lumen electron carrier cytochrome c_6 (Cyt c_6), and in ferredoxin outside the thylakoids (9). Nowadays, Fe is a limiting nutrient in most ecosystems, especially in aquatic environments (10). Consequently, photosynthetic organisms evolved several alternatives to iron-containing proteins. In most cyanobacteria and some algae, Cyt c_6 can be replaced by plastocyanin (PC), a type I blue-copper protein that acts as a thylakoid lumen soluble electron transporter. These two electron carriers are present in millimolar concentrations inside the thylakoid lumen (11), where Cyt c_6 is replaced by PC, and this substantially reduces the requirement for Fe. In addition, PC uses Cu

which is much more soluble (and therefore available) than Fe in an oxidizing atmosphere. Interestingly, plants have only retained PC for photosynthetic electron transfer (12).

Expression of these two alternative electron carriers is regulated by copper availability in both cyanobacteria and algae (13–15). The regulatory system has been well characterized in the green alga *Chlamydomonas reinhardtii*. CRR1, an SQUAMOSA promoter binding protein-like transcription factor (SBP), activates *CYC6* (encoding Cyt_{c6}) expression in response to copper limitation (16). CRR1 is able to bind both Cu⁺ and Cu²⁺, with the latter inhibiting the DNA-binding ability of the SBP domain (16). However, the regulatory system has not been identified in cyanobacteria, even though the Cyt_{c6}/PC switch was discovered more than 40 years ago (13, 14, 18–21), and it is frequently used to control the expression of genes in several model systems (22–24). The switch responds to the presence of copper at the transcriptional level; *petJ* (encoding Cyt_{c6}) is repressed while *petE* (encoding PC) is induced (18, 21, 25–27). Several regulatory systems that respond to copper have been described in cyanobacteria including CopRS (25, 27), InrS (27, 28), and BmtR (29), but none are involved in regulating these two genes.

Herein, we unveiled the regulatory mechanism for the Cyt_{c6}/PC switch in cyanobacteria. It involves two genes: one encoding a transcription factor of the Blal superfamily (PetR) that binds to both *petE* and *petJ* promoters, repressing *petE* and activating *petJ* in the absence of copper. The second gene encodes a

protease (PetP) that degrades PetR in response to copper. PetR degradation is activated by copper and requires PetP and copper *in vivo* and in an *in vitro* assay system, suggesting that PetP is responsible for PetR degradation. The system is widespread in all cyanobacteria, including early-branching cyanobacteria such as *Gloeobacter* and *Pseudoanabaena*, with the exception of most strains of the *Synechococcus/Prochlorococcus* groups.

Results

A Blal/CopY homolog regulates the *petJ/petE* switch in *Synechocystis*

Bioinformatics analysis of genes within the vicinity of *petE* and *petJ* in 169 cyanobacterial genomes revealed that a transcription factor of the Blal/CopY family (renamed PetR) was associated with *petJ* and/or *petE* genes in 13 genomes (SI Appendix, Figure S1). Although the overall number of genomes with this gene association was low, co-occurrence of *petR* with both *petE* and *petJ* was higher. Specifically, 75% of genomes harboring *petE* and *petJ* also contained *petR* (and a linked gene encoding a BlaR homolog; see below; SI Appendix, Figure S1), and these two genes were only present in genomes containing *petE* and *petJ* (SI Appendix, Figure S1; Supplementary dataset S1) with a few exceptions. This raised the possibility that *petR* could be related to the *petE/petJ* switch. In order to test this hypothesis, a mutant in the *slr0240* gene encoding PetR in *Synechocystis* sp. PCC 6803 (*Synechocystis* hereafter) was generated (PETR strain). The mutant strain fully segregated (SI Appendix, Figure

S2) and the phenotype was indistinguishable from that of the wild-type (WT) strain on BG11C plates (the standard defined medium for cyanobacteria; see below). Expression of *petJ* and *petE* genes was analyzed by RNA blot in both WT and PETR strains after copper addition to cells grown in BG11C lacking Cu (BG11C-Cu). In WT cells, *petE* was induced in response to copper and *petJ* was repressed (Figure 1A), as previously described (18, 25–27). By contrast, in the PETR strain *petE* was expressed even in the absence of copper and *petJ* was not expressed under any conditions (Figure 1A). Conversely, *copM* expression, which is induced in response to copper (25, 30), was similar in both strains. At the protein level, PC was induced in WT cells upon copper addition, while Cyt_c₆ abundance decreased after 4 and 24 h. CopM levels increased after copper treatment, with the unprocessed form of the protein accumulating first, followed by the processed at later timepoints (Figure 1B). By contrast, Cyt_c₆ was not detected under any conditions in the PETR strain, and PC was detected even in BG11C-Cu medium (Figure 1B). An increase in PC levels was still evident in the PETR strain after copper addition (Figure 1B), probably due to an increase in copper-bound PC, which is more stable than apoPC (25), since mRNA levels did not change after copper addition (Figure 1A). No changes in the CopM induction pattern were observed (Figure 1B). When an additional copy of *petR* was introduced in the PETR strain, either under its own promoter (PETR2 strain) or under the strong *cpcB* promoter (PETR3 strain), a WT phenotype was restored (SI Appendix, Figure S3). This indicates that *petR* was the only gene responsible for regulating these two genes. Similar results were obtained when protein levels

were analyzed in steady-state cells of the WT strain grown in media with differences in copper availability. PC was high in copper-containing medium (BG11C 330 nM Cu), lower in BG11C-Cu (~30 nM Cu), and much lower in BG11C-Cu+ bathocuproinedisulfonic acid (BCSA), a copper-specific chelator (no copper; Figure 1C). The PETR strain expressed PC in all conditions, but accumulated higher levels in copper-containing media. Meanwhile, Cyt_c₆ was not detected in the PETR strain in any media, and WT cells displayed higher levels in BG11C-Cu+BCSA than in BG11C-Cu (Cyt_c₆ was almost undetectable in copper-containing medium; Figure 1D).

Slr0241(PetP) is also involved in *petJ*/*petE* regulation

Downstream of *petR* there is another gene (*slr0241*, herein named *petP*) that encodes an M48/M56 superfamily of proteases (Pfam families PF01435 and PF05569). The *petP* start codon overlaps with the stop codon of *petR*, and both genes were expressed as a single transcriptional unit (SI Appendix, Figure S4). Analysis of the PetP sequence using TMHMM 2.0 software (31) predicted four transmembrane helices (residues 4–21, 28–50, 65–87, and 256–278), a 14 residue loop connecting TM2 and TM3, and a cytoplasmic domain (88–256) that contains the HEXXH conserved catalytic motif (SI Appendix, Figure S5). This domain organization is very similar to that of BlaR from *Staphylococcus aureus* (32), which is involved in regulating β -lactam resistance by degrading BlaI, although it lacks the C-terminal periplasmic domain that binds to β -lactams. Furthermore, the organization of *petRP* is conserved in all cyanobacterial

genomes (SI Appendix, Figure S1, and Supplementary dataset S1), suggesting that both genes might be functionally related. To test this hypothesis, we mutated the *petP* gene. The resulting mutant strain (PETP) was viable (SI Appendix, Figure S2) and exhibited a similar growth phenotype to the WT strain on BG11C plates (see below). When gene expression was tested in this mutant, *petE* was repressed both in the presence and absence of copper, while *petJ* was expressed constitutively (Figure 1A). The same results were observed at the protein level, with Cyt_c₆ detected both in the presence and absence of copper, while PC was undetectable under any conditions (Figure 1B). When steady-state proteins levels were analyzed in different copper-containing media, PC was not detected in the PETP strain in any media, while Cyt_c₆ was detected in all media. Despite this, Cyt_c₆ levels were lower in copper-containing medium than in media lacking copper (Figure 1D). These results indicate that *petP* is involved in the regulation of both genes, acting in an opposite way to *petR*. A double mutant lacking both *petR* and *petP* genes was also generated (PETRP; SI Appendix, Figure S2). The PETRP strain displayed the same phenotype as the PETR strain (SI Appendix, Figure S6), revealing that *petP* acts upstream of *petR*.

PetP regulates PetR levels in response to copper

Given that PetP belongs to the M48/M56 superfamily of proteases, this protein might regulate PetR through proteolysis. To investigate this hypothesis, we generated an antiserum against PetR that allowed us to monitor PetR protein

abundance in response to copper. In the WT strain, PetR was detected in cells cultivated in BG11C-Cu, and it was rapidly degraded after copper addition, since *petRP* mRNA levels did not change following copper addition (SI Appendix, Figure S4). Two bands were detected (~15 kDa and ~12 kDa, Figure 2A), suggesting an initial and specific cut, and subsequent degradation of the fragments. Indeed, in order to consistently detect PetR, whole cells were loaded onto gels (see Materials and Methods) because PetR was quickly degraded during protein extract preparation. The best studied homologs of PetRP are BlaIR proteins involved in antibiotic resistance from *S. aureus* (32). Processing of these proteins was initially identified in BlaI, it takes places after Phe100 in the amino acid sequence, and is completely conserved in all cyanobacterial PetR proteins (SI Appendix, Figure S7). As expected, the theoretical molecular mass of the fragment generated by this digestion was consistent with the expected size detected by immunoblotting (Figure 2A and SI Appendix, Figure S7; expected sizes of full-length and processed proteins are 15.80 and 12.38 kDa, respectively). In contrast to the WT strain, PetR levels did not change in response to copper in the PETP strain (Figure 2). Therefore, the initial cleavage is probably mediated by PetP, and the fragments are probably further degraded by other unspecific proteases. As expected, PetR was not detected in the PETR strain (Figure 2 and SI Appendix, Figure S3E). When PetR protein abundance was analyzed in steady-state cultures grown in different copper regimes, levels were clearly regulated in the WT strain, since they were higher in the presence of BCSA than in BG11-Cu, and not detected in copper-containing media (Figure 2B

and C). PetR levels were higher in the PETP strain than in the WT strain, even when WT cells were cultured in the presence of BCSA, although levels were decreased slightly at higher copper availability (Figure 2B and C). These results are consistent with the Cyt_c₆ and PC levels measured under the same conditions (Figure 1C and D), and demonstrate that PetR protein abundance is regulated by PetP *in vivo* according to Cu availability.

PetP degrades PetR directly in a copper-dependent manner

To further characterize this regulation, we performed an *in vitro* degradation assay in which recombinant GST-PetR was incubated with total cell extracts from WT, PETR, and PETP strains, and GST-PetR processing was followed by immunoblotting. GST-PetR was not degraded by whole-cells extracts prepared from WT, PETP, or PETR strains when copper was not added to the extracts (Figure 2D). In contrast, when copper was added to extracts, GST-PetR was degraded by WT and PETR extracts, but not by PETP extracts, indicating that a PetP-mediated PetR degradation was activated by copper (Figure 2D).

To confirm that PetP was directly involved in PetR degradation, and that this degradation was not mediated by an additional protein from *Synechocystis*, *petRP* genes under their own promoter were cloned in a plasmid and transformed into *E. coli*. Total extracts were prepared from cells carrying a plasmid containing *petRP* or an empty plasmid, and degradation of GST-PetR was monitored. The results were similar to those obtained using total extracts from *Synechocystis*: GST-PetR was degraded in total extracts prepared from

cells containing a plasmid that includes *petRP*, but not in extracts prepared from the strain carrying an empty plasmid (Figure 2E). PetR degradation also required copper supplementation of extracts, demonstrating that PetR degradation depends on this metal (Figure 2E). These results also showed that PetP can degrade PetR directly without the participation of any other protein from *Synechocystis*. Thus, PetP controls PetR levels directly via a copper-dependent mechanism.

PetR binds to both *petE* and *petJ* promoters

In order to establish whether PetR regulates *petJ* and *petE* directly, we performed Electrophoretic Mobility-Shift assays to analyze PetR binding to both promoter regions. As a first approach, we carried out band shift assays using a recombinant version of PetR (see Materials and Methods for details) and two probes corresponding to the *petJ* and *petE* promoters. PetR was able to bind to both promoters (Figure 3A and B), and the binding was not affected by Cu (Cu⁺ or Cu²⁺; SI Appendix, Figure S8). PetR exhibited a higher affinity toward *petE* than the *petJ* promoter, and was able to completely titrate the probe at lower concentrations (Figure 3A and B). These results indicate a dual role for PetR activating *petJ* expression and repressing *petE*.

In order to identify the PetR binding site, we carried out a bioinformatic analysis of the *petE* and *petJ* promoters in cyanobacterial genomes. A 300-bp fragment surrounding *petE* and *petJ* promoters was retrieved from genomes that contained *petE*, *petJ*, and *petRP* using MGcV database (<http://mgcv.cmbi.ru.nl/>;

SI Appendix, Supplementary dataset S2). Analysis of the sequences of *petE* and *petJ* promoters using MEME suite revealed that the motif shown in Figure 3C was significantly enriched in both. This motif contains the GACN₅GTC sequence, which was completely conserved, positioned 75 bp upstream from the starting ATG codon (and 37 bp from the tsp) in the *petJ* promoter, and at 118 bp from the ATG codon (and 23 bp from the tsp) in the *petE* promoter. In the *petJ* promoter, the putative binding site is centered around the -35 box in the promoter, in a location suitable as an activator site. In contrast, this sequence is located between the -10 and -35 boxes in the *petE* promoter, compatible with a repressor role of PetR. These locations are consistent with gene expression data obtained by RNA blotting (Figure 1) and transcriptomics analysis (see below) of PETR and PETP mutant strains, which suggests that PetR activates *petJ* and represses *petE* expression. Notably, when MEME analysis was repeated using only *petJ* promoters, an extended motif was identified (Figure 3D). This motif includes the core of the motif shown in Figure 3C, but also two direct repeats of the sequence TTTGAC-N₉-TTTGAC (Figure 3D). A different motif was also identified when the same analysis was performed using only *petE* promoter sequences (Figure 3E). This motif includes the core motif shown in Figure 3C, but in an extended form of the inverted repeat GACA-N₃-TGTC. However, the *Synechocystis* sequence in the *petE* promoter is GACG-N₃-TGTC (Figure 3F).

RNA-seq analysis of the PetR regulon

289 To fully characterize the PetR regulon, RNA-seq was carried out using RNA
 290 extracted from WT, PETR, and PETP strains grown in BG11C-Cu, and 2 h after
 291 addition of 0.5 μ M Cu addition (SI Appendix Supplementary dataset S3). Using a
 292 threshold of at least a 2-fold change and $P_{adj} < 0.01$, 14 genes were induced
 293 (*petE*, both plasmid and chromosomal copies of *copMRS*, *copBAC*, and *nrsBAC*)
 294 and three were repressed (*petJ*, *slr0601*, and *slr0602*) in the WT strain after
 295 copper addition (SI Appendix, Table S1; Supplementary dataset S4), in
 296 agreement with previous microarray analysis (27). A similar response was
 297 observed in the PETR strain with *copMRS* (both copies), *copBAC*, and *nrsBAC*
 298 operons, and three additional genes of unknown function were also differentially
 299 regulated (SI Appendix, Table S1). In the PETP strain, 16 genes were induced
 300 after copper treatment, including *copMRS* (both copies), *copBAC*, and *nrsBAC*
 301 operons (SI Appendix, Table S1). As expected from the results shown in Figure
 302 1, neither *petE* nor *petJ* were differentially expressed in PETR or PETP strains
 303 (SI Appendix, Figure S9). Furthermore, *slr0601* and *slr0602* were not
 304 differentially regulated in PETR or PETP (SI Appendix, Table S1). When gene
 305 expression was directly compared between WT and PETR strains, eight genes
 306 were differentially regulated in BG11C-Cu (seven expressed at higher levels in
 307 WT, and one downregulated), and six were differentially regulated after copper
 308 addition (SI Appendix, Supplementary dataset S4). As expected, this group of
 309 genes included *petRP* (*petR* was induced 5.93 and 4.73-fold, and *petP* was
 310 induced 37 and 38-fold for in BG11C-Cu and +Cu, respectively), *petE* (repressed
 311 41-fold in -Cu and 2-fold in +Cu), and *petJ* (induced 168-fold in -Cu and 8.7-fold

in +Cu), as well as *slr0601*, *slr0602*, *slr0242*, *sll1796*, *sll0494*, and *slr1896* (SI Appendix, Supplementary dataset S4). Similarly, when WT and PETP strains were compared, only three genes were differentially regulated in BG11C-Cu (all upregulated), and eight genes were differentially regulated in BG11C+Cu (also upregulated). These genes included *petP* (upregulated 5-fold in both conditions in WT), *petE* (also upregulated in WT, by 5.3-fold in -Cu and 116-fold in +Cu), *petJ* (repressed 25-fold in +Cu), and *slr0242*, *slr0601*, *slr0602*, *slr1896*, and *sll1926* (SI Appendix, Supplementary dataset S4). These results suggest that the PetRP regulon comprises *petJ*, *petE*, *slr0601*, and *slr0602*, as these genes were regulated by copper in WT cells, and affected in both PETR and PETP strains, although with opposite patterns in the two mutant strains.

The *slr0601* and *slr0602* genes are linked in the genome and probably form an operon (SI Appendix, Figure S10A). Expression of *slr0601* was analyzed by RNA blotting, and the pattern was similar to that of *petJ*: repressed after copper addition in WT, not expressed in PETR, and constitutively expressed in PETP strains (SI Appendix, Figure S10B). Furthermore, PetR could bind to the promoter sequence of *slr0601* (SI Appendix, Figure S10C), which contains a sequence matching the one identified in Figure 3 (SI Appendix, Figure S10D). Together, these results indicate that the *petRP* regulon is only composed of *petJ*, *petE*, and *slr0601-slr0602* in *Synechocystis*.

Physiological characterization of PETR and PETP mutant strains

334 All mutants generated in this study were fully segregated in BG11C, and were
335 able to grow in BG11C-Cu, but given their gene expression programs (Figures 1
336 and 2; SI Appendix, Figure S9; Table S1, supplementary dataset S4), growth
337 defects were expected in different copper regimes. We analyzed the growth of
338 the PETR and PETP strains on plates differing in copper availability: BG11C-
339 Cu+BCSA (no copper), BG11C-Cu (~30 nM Cu), and BG11C (330 nM Cu).
340 PETR was only affected in BG11C-Cu containing BCSA, similar to the PETJ
341 strain, which has an interrupted *petJ* gene (Figure 4A). By contrast, the PETP
342 strain was not affected under any conditions (Figure 4), because it expressed
343 *petJ* in all tested conditions (Figure 1). Finally, the PETE strain showed a
344 decreased growth rate (but survived) in BG11C, as previously reported (25, 26,
345 33) (Figure 4A and SI Appendix, Figure S11). A double mutant lacking both
346 genes (PETRP strain) was indistinguishable from the single mutant PETR
347 (Figure 4A and SI Appendix, Figure S6). Analysis of photosynthetic activity
348 revealed that PETJ and PETR strains reached lower oxygen evolution rates
349 ($16.8 \pm 0.3 \mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$ and $18.9 \pm 0.75 \mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$, respectively)
350 than WT ($23 \pm 3.82 \mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$), PETE ($24.5 \pm 2.46 \mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$),
351 and PETP ($23.2 \pm 5 \mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$) strains when cultured in BG11c-
352 Cu+BCSA, although they saturated at the same light intensity (Figure 4B). When
353 cultured in BG11C, the PETE strain exhibited a severe defect in oxygen evolution
354 (reaching only $5.3 \pm 2.7 \mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$ and saturating at a light intensity of
355 $180 \mu\text{mol m}^{-2}\text{s}^{-1}$) (Figure 4C). Surprisingly, the PETP strain exhibited a lower
356 maximum capacity, reaching only $13.24 \pm 2.63 \mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$, while WT

(24.11 $\pm$ 3.66 $\mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$), PETJ (22.25 $\pm$ 4.34 $\mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$), and PETR (21.87 $\pm$ 4.36 $\mu\text{mol min}^{-1}/\text{OD}_{750\text{nm}}$) strains all reached similar levels, equivalent to those observed in BG11C-Cu (Figure 4C). Despite this, all strains saturated at the same light intensity (500 $\mu\text{mol m}^{-2}\text{s}^{-1}$). The PETP strain expressed *petJ* constitutively (Figures 1 and 2), and was therefore not expected to show any photosynthetic defect.

Genetic interactions between *petRP*, *petJ*, and *petE*

Our results show that *petRP* is the main system responsible for the *petJ/petE* switch following copper addition. Either *petJ* or *petE* are required for photosystem I (PSI) reduction, and therefore for the growth of cyanobacteria (33), despite early studies showing growth of *Synechocystis* in the absence of both proteins (34). The results presented in Figures 1 and 4 suggest that the low levels of PC present in BG11C-Cu (even in the presence of BCSA) were enough to sustain growth of PETR and PETJ mutant strains, albeit at decreased rates (Figure 4A). Furthermore, although the PETE strain showed impaired growth in the presence of copper (Figure 4A), it was able to grow at decreased rates because all cells grew after prolonged incubation (SI Appendix, Figure S11), probably due to low levels of Cyt_c₆. Moreover, a double mutant lacking functional *petE* and *petR* genes (or both *petR* and *petP*, strain PETERP) could not be completely segregated (SI Appendix, Figure S12), probably due to low expression levels of *petJ* in the absence of PetR (Figures 1A and SI Appendix, Figure S9), reinforcing

the idea that at least one of the soluble transporters (even at low levels) is needed in *Synechocystis*. By contrast, a double mutant lacking *petE* and *petP* (PETEP) was able to grow at WT rates in the presence of copper (SI Appendix, Figure S11 and S12). Consistently, the double mutant PETJP did not segregate, while PETJR and PETJRP mutants fully segregated (SI Appendix, Figure S12). Together, these results suggest that a double mutant lacking both *petJ* and *petE* should not be viable. To test this hypothesis, a double mutant was generated by transforming the PETE mutant with pPETJCm. Merodiploid colonies were obtained, but they never segregated completely (SI Appendix, Figure S13), despite subculturing under several light regimes for more than 2 years. To confirm that PC and Cyt_c₆ were expressed in slow-growing mutants, we used concentrated extracts and loaded 150 µg of total protein per lane. Cyt_c₆ was detected at low levels in WT and PETE strains, but not in the PETR strain, in the presence of copper (SI Appendix, Figure S12). In WT, PETR, and PETJ strains, PC was detected (although at low levels; Figures 1, 2, and SI Appendix, Figure S12) in the absence of Cu, therefore allowing growth of these strains at decreased rates (Figure 4).

Discussion

Acquisition of PC as an alternative electron carrier was probably an early adaptation after oxygenic photosynthesis appeared, since local oxygenation (even before the GOE) presumably decreased Fe availability. This is reinforced

by the presence of both electron carriers in early-branching cyanobacteria such as *Gloeobacter* and *Pseudoanabaena* (SI Appendix, Figure S1, Supplementary dataset 1). Although known about for more than 40 years (13, 14), the regulatory system responsible for the *petJ/petE* switch in cyanobacteria remained undiscovered. Herein, we show that the *petRP* genes are responsible for this regulation (Figures 1 and 5). The system seems to have been acquired early during evolution, because like *petE*, it is present in early branching cyanobacteria, and it is widespread in most cyanobacterial clades (SI Appendix, Figure S1). The only exception is the *Synechococcus/Prochlorococcus* clade, in which most strains lack *petRP* genes, similar to previous observations on other regulatory systems (35). We showed that *petRP* act in a single pathway in which PetR positively regulates *petJ* and negatively regulates *petE*, while PetP regulates PetR protein levels in response to copper (Figure 5). PetR directly regulates *petE* and *petJ* gene expression by binding to the promoter regions of these two genes (Figure 3). However, the affinity for both promoters is differs markedly, with the *petJ* promoter requiring much higher PetR concentrations for binding than the *petE* promoter *in vitro*. In addition, binding sites are in different locations in the two promoters, replacing the -35 box in *petJ* and between -10 and 35 in *petE*, consistent with an activating role in *petJ* and a repressing role in *petE*, as confirmed by gene expression analysis (Figures 1, 2 and 5; SI Appendix, Figure S9, Table S1, supplementary dataset S4).

Furthermore, our bioinformatics analysis identified an inverted repeat as the core binding site for both promoters in cyanobacteria (Figure 3C), but when *petJ* and *petE* promoters were used, variations of the motif were identified (Figure 3D and E). Given that the protein abundance of PetR *in vivo* depends on copper concentration (Figure 2), and binding to the *petJ* (and *slr0601*) promoter requires higher PetR concentrations (Figure 3 and SI Appendix, Figure S10), *petJ* will only be fully activated when all copper is exhausted (and PetR accumulates at high levels), preventing induction of *petJ* if some copper remains available for PC synthesis. Although Blal/CopY homologs have been reported to act only as repressors, our results strongly suggest that PetR acts as an activator for *petJ* expression (Figures 1, 3, and SI Appendix, Figure S9B; Table S1). This indicates that Blal/CopY homologs may also act as activators, as suggested previously (36). Since the binding sites for PetR activation are different to the repressor sites (Figure 3), it is possible that they have remained elusive when studying PetR homologs (37). This dual role for DNA binding proteins that were initially considered to be repressors has been uncovered recently in proteins such as Fur, which was believed to be an activator in both its apo and holo forms (38–40).

Our RNA-seq analysis revealed that only four genes are under the control of the *petRP* system in *Synechocystis*: *petE*, *petJ*, *slr0601*, and *slr0602* (SI Appendix, Figures S9 and S10; Table S1). Inspection of the promoter sequences of these four genes reveals conservation of the core sequence, identified using

GACN₅GTC in the *petJ* and *petE* promoters, but also using only *petJ* promoter sequences (Figure 3 and SI Appendix, Figure S10). The *slr0601* and *slr0602* genes encode two small proteins restricted to cyanobacteria that share low sequence conservation. Slr0601 contains two transmembrane domains with both C-terminal and N-terminal regions predicted to be cytoplasmic, while Slr0602 contains a coiled-coil region but no other recognizable domains. The low levels of sequence conservation and the lack of known domains make it difficult to assign putative functions for these two proteins.

PetP belongs to the M48/M56 family of proteases, and we showed that this protein regulates PetR by controlling its protein abundance in response to copper *in vivo*. This regulation is probably direct as we showed that recombinant PetR is degraded *in vitro* by both *Synechocystis* and *E. coli* extracts, and that this degradation depends on the presence of PetP and copper (Figure 2). This regulatory system is homologous to the Blal/Blar regulatory system that regulates β -lactamase expression in *S. aureus*. We explored the mechanism by which copper is sensed by PetRP. PetR belongs to the Blal/CopY family of transcription factors, and CopY proteins have been demonstrated to bind copper directly (41). Therefore, it is possible that PetR may bind copper, and although this binding may not affect its DNA binding activity (SI Appendix, Figure S8), it may alter its conformation, making it more susceptible to degradation by PetP. Nevertheless, CopY copper binding sites are not conserved in PetR (28), making this hypothesis unlikely.

BlaR/MecR contains a C-terminal extra-cytoplasmic domain that is acylated by β -lactam antibiotics, and this modification activates BlaR, which subsequently degrades Blal. This domain is absent in PetP, and therefore copper sensing must be mediated by a different domain. The loop between the second and third transmembrane domains of PetP includes two methionine residues (M53 and M58 in *Synechocystis*) that are conserved in all PetP sequences (SI Appendix, Figure S5), which could be putative copper ligands. Interestingly, this domain in BlaR/MecR interacts with the periplasmic C-terminal domain after it binds to β -lactams (42–44). This interaction has been proposed to transduce the signal and activate the protease domain in BlaR. Therefore, these two methionine residues in PetP may bind copper and transduce the signal in a similar way.

Moreover, as PetP is an integral membrane protein, it would be interesting to identify which cyanobacterial membrane system this protein targets. Most copper in *Synechocystis* is presumably sequestered in the thylakoids, where it is bound to PC (45, 46), and CopS, the sensor protein in the copper resistance system, is located to both plasma and thylakoid membranes (25). If PetP is also targeted to both membrane systems, this could allow PetP to respond not only to an external surplus of copper, but also when the metal is released from internal PC. PetP localization to the thylakoids is reinforced by the fact that mutants lacking the P_1 -type-Cu-ATPases CtaA, PacS, or both (which are involved in Cu acquisition by PC) displayed decreased levels of *petE* mRNA after copper

addition (25, 47). This suggests that Cu needs to be transported to the thylakoid in order to be detected by the PetRP system.

Furthermore, our results showed that *Synechocystis* can synthesize PC even when copper levels are very low due to the presence of BCSA in BG11C-Cu. This indicates that some copper can be transported even in the presence of BCSA, suggesting that the copper import system (which remains unidentified) has a higher affinity for copper than the chelating agent. Accumulation of PC in the presence of BCSA was only observed in the PETR strain in which *petE* expression was constitutive (Figures 1, 5, and SI Appendix, Figure S9), because in the WT strain *petE* is repressed and PC cannot be synthesized, even if copper is available.

Finally, our results also suggest that there are other layers of regulation at the posttranscriptional level because Cyt_{c6} and PC protein abundance does not always correlate with mRNA levels. PC was more abundant in Cu-containing media in the PETR strain in which *petE* mRNA levels did not change in response to copper (Figure 1). Similarly, Cyt_{c6} levels were decreased in the PETP strain after copper addition, even when mRNA levels were not decreased. PC protein abundance may be related to the protein not being correctly translated and/or folded in the absence of available copper, or due to degradation of the apoprotein, but synthesis of Cyt_{c6} should not be altered by copper. It has been proposed that IsaR sRNA regulates *petJ* translation in response to Fe deprivation (24). We did not find *isaR* to be differentially regulated in our RNA-seq

experiment (SI Appendix, Figure S14), hence it is unlikely that this regulation is mediated by *petRP*. Furthermore, oxygen evolution was affected in the PETP strain (which expresses Cyt_c₆; Figure 4C) in the presence of copper, suggesting that additional changes in the photosynthetic transport chain might be present, as reported for *Chlamydomonas* (48). Finally, although single mutants of either *petJ* or *petE* were able to grow, even in a copper regime in which expression of the other gene was repressed, these strains exhibited growth and photosynthesis defects (Figure 4), in agreement with previous studies (33, 49–51). Double mutants of both *petJ* and *petE* were not viable (SI Appendix, Figure S13), and double mutants lacking one of the genes encoding the electron transport proteins and with decreased expression of the other due to the lack of *petP* (constitutive repression of *petE*; strain PETJP) or *petR* (unable to induce *petJ*; strain PETER) were not viable either (SI Appendix, Figure S12). Together, these results demonstrate that *petRP* are essential for expression of these genes, and that levels of PC or Cyt_c₆ proteins must be above a critical threshold for survival of the strains. These results also rule out the existence of another electron transfer mechanism between cytochrome *b*₆*f* and PSI.

Materials and Methods

Strains and culture conditions

All *Synechocystis* strains used in this work were grown photoautotrophically on BG11C (52) or BG11C-Cu (lacking CuSO₄) medium at 30°C under continuous

illumination using 4000–4500 K white-LED light ($50 \mu\text{mol photon m}^{-2} \text{ s}^{-1}$, measured at the flask surface) and bubbled with a stream of 1% (v/v) CO_2 in air. BG11C-Cu was supplemented with 300 μM BCSA as a chelating agent to trap any trace of copper, when required. For plate cultures, media were supplemented with 1% (w/v) agar. Kanamycin, nourseothricin, chloramphenicol, spectinomycin, and erythromycin were added to a final concentration of 50 $\mu\text{g mL}^{-1}$, 50 $\mu\text{g mL}^{-1}$, 20 $\mu\text{g mL}^{-1}$, 5 $\mu\text{g mL}^{-1}$, and 5 $\mu\text{g mL}^{-1}$, respectively. Experiments were performed using cultures from the logarithmic phase (3–4 $\mu\text{g chlorophyll mL}^{-1}$). *Synechocystis* strains are described in SI Appendix, Table S2.

E. coli DH5 α cells were grown in Luria Broth or M9 (supplemented with 1 mL LB per L of M9) media and supplemented with 100 $\mu\text{g mL}^{-1}$ ampicillin, 50 $\mu\text{g mL}^{-1}$ kanamycin, 50 $\mu\text{g mL}^{-1}$ nourseothricin, 20 $\mu\text{g mL}^{-1}$ chloramphenicol, and 100 $\mu\text{g mL}^{-1}$ spectinomycin when required.

Insertional mutagenesis of Synechocystis genes

For *slr0240* (*petR*) mutant construction, a 1524-bp fragment was amplified using oligonucleotides 223 and 224, cloned into pSPARKII (generating pPETR), and a Sp^R cassette was inserted in the *EcoRV* site at codon 31 in the *slr0240* open reading frame (ORF), generating plasmid pPETR_Sp. For the *slr0241* (*petP*) mutant, a 970-bp fragment was amplified using oligonucleotides 248 and 249 and cloned into pSPARKII, generating pPETP. A 582-bp *NheI* fragment was substituted by an *XbaI* erythromycin cassette, generating pPETP_Ery. For the *petRP* double mutant, a 1492-bp fragment was amplified by PCR using

oligonucleotides 256 and 257, and cloned into the pSPARKII vector, generating the pPETRP plasmid. A 650-bp *Bst*EI fragment from plasmid pPETRP was substituted by an *Hinc*II-digested SpR cassette from pRLSpec, generating pPETRP_Sp. To generate the *petJ* mutant, a 1023-bp fragment was amplified using oligos 235 and 236, and cloned into pSPARKII, generating pPETJ, and a *Xba*I CC1 Cm resistance cassette was inserted in the *Nhe*I site present in the *petJ* ORF, generating pPETJ_Cm. For PETR complementation, a 639-bp fragment (185 bp upstream of the *petR* STOP codon) was amplified using oligonucleotides 256 and 234, and cloned into pGLNN digested with *Eco*RI and *Not*I, generating pPETR_comp. For the PETR3 strain, a 463-bp fragment was amplified using oligonucleotides 258 and 234 (adding *Sa*I and the pET28 ribosome binding site upstream of *petR* ATG), digested with *Sa*I-*Not*I, and cloned into pGLN-PcpcB digested in the same way, generating pPETR_OE. These plasmids were used to transform WT *Synechocystis*, PETR, PETE, or PETJ strains, generating PETR, PETR2, PETR3, PETP, PETRP, PETER, PETEP, PETERP, PETJR, PETJP, PETJRP, and PETEJ strains. Segregation of the strains was verified by PCR with the appropriate primer pairs, as depicted in SI Appendix, Figures S2, S3, S11, and S13. All plasmids were sequenced to verify that no unwanted mutations were introduced. Sequences for all oligonucleotides are listed in SI Appendix, Table S3.

RNA isolation and RNA blot analysis

Total RNA was isolated from 30 mL samples of *Synechocystis* cultures in the mid-exponential growth phase (3 to 4 µg chlorophyll mL⁻¹). Extractions were performed by vortexing cells in the presence of phenol-chloroform and acid-washed baked glass beads (0.25–0.3 mm diameter) as previously described (53). A 5 µg sample of total RNA was loaded onto each lane of a 1.2% denaturing agarose formaldehyde gel (54), electrophoresed, and transferred to a nylon membrane (Hybond N-Plus; GE Biosciences). Prehybridization, hybridization, and washes were performed in accordance with the manufacturer's instructions. Probes for RNA blot hybridization were synthesized by PCR using oligonucleotide pairs 172-173, 174-175, 389-390, 275-276, 274-275, and 178-179 (SI Appendix, Table S3) for *petE*, *petJ*, *copM*, *slr0601*, and *mnpB*, respectively. DNA probes were ³²P-labeled with a random primer kit (Rediprime II catalog #RPN1633; GE Biosciences) using [α-³²P] dCTP (3,000 Ci/mmol). Hybridization signals were quantified with a Cyclone Phosphor System (Packard). Each experiment included at least three biological replicates.

Immunoblotting

For analysis, proteins were fractionated by SDS-PAGE and immunoblotted with antibodies against plastocyanin (1:3000), cytochrome *c*₆ (1:3000), CopM (1:7000), PetR (1:6000), or *Synechococcus* sp. PCC 6301 glutamine synthetase I (1:100,000). ECL Prime (catalog # RPN2232, GE Biosciences) was used to detect the different antigens with anti-rabbit secondary antibodies conjugated to horseradish peroxidase (1:25,000; catalog # A0545, Sigma). Either photographic

film or a ChemiDoc Imaging System (Bio-Rad) were used for signal detection and quantification.

Samples were prepared from whole cells or soluble extracts. For whole cells, cells were harvested from cultures with an optical density at 750 nm (OD_{750nm}) of 1, the supernatant was carefully removed, cells were resuspended in 100 μ L of 1 $\times$ Laemmli loading buffer, and boiled for 10 min. For soluble extracts, Cells equivalent to 20 OD_{750nm} were collected, resuspended in 300 μ L of buffer A (50 mM TRIS HCl pH 8, 50 mM NaCl, 1 mM phenylmethylsulfonyl fluoride) and broken using glass beads via two cycles, separated by 5 min on ice, of 1 min in a minibead beater. Cell extracts were recovered from the beads and samples were clarified by two sequential centrifugations: 5 min at 5000 g to eliminate cell debris, and 15 min at 15,000 g to remove membranes. Protein concentrations in cell-free extracts and purified protein preparations were quantified using Bradford reagent (catalog # 5000006, Bio-Rad), with ovalbumin as a standard, and specified amounts of protein were separated by SDS-PAGE.

PetR protein purification

The complete *petR* ORF was cloned from *Synechocystis* DNA after PCR amplification with oligonucleotides 233 and 234, and cloned into the *Bam*HI-*Not*I sites of pGEX6P, generating pGEX_PETR. The GST-PetR fusion protein was expressed in *E. coli* DH5 α cells. A 1 L culture was grown in Luria broth medium to an OD_{600nm} of 0.6, cooled to 4°C, 0.1 mM isopropyl-b-D-thiogalactopyranoside (IPTG) was added, and culturing was continued at 25°C overnight. Cells were

harvested by centrifugation, resuspended in 5 mL of PBS buffer (150 mM NaCl, 16 mM Na₂HPO₄, 4 mM NaH₂PO₄, 4 mM phenylmethylsulfonyl fluoride, 7 mM β-mercaptoethanol) supplemented with 0.1% Triton X-100, broken by sonication on ice, and insoluble debris was pelleted by centrifugation for 45 min at 25,000 *g*. Soluble extracts were mixed with 1 mL of glutathione agarose beads (catalogue number # 17075601, GE Healthcare) and incubated for 2 h at 4°C with gentle agitation. Beads were transferred to a column and washed extensively with PBS buffer (20–30 column volumes) until no more protein was eluted from the column. The column was equilibrated with 10 volumes of 50 mM TRIS HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, and 1 mM dithiothreitol, and beads were resuspended in two volumes of the same buffer. A 20 µL sample of Precision protease (2 U/µL; catalogue # 27084301, GE Healthcare) was added and beads were incubated overnight at 4°C. Beads were poured into an empty column and the flow through fraction was collected. This fraction was diluted three times in 50 mM TRS HCl pH 8, and applied to a HiTrap Heparin column (catalogue # 17040601, GE Healthcare) connected to an Amersham FPLC system. The column was washed with 10 volumes of 50 mM TRIS HCl, 100 mM NaCl, and a 20 mL gradient from 0.1–1 M NaCl was applied. Fractions containing PetR were pooled and concentrated. GST-PetR fusion protein was eluted from glutathione agarose beads with 3 mL of 50 mM TRIS HCl pH 8 containing 10 mM reduced glutathione after washing the column with PBS. A gel of the purified fractions is shown in SI Appendix, Figure S15.

Protease assays

Cells equivalent to 20 OD_{750nm} of *Synechocystis* strains were harvested from mid-exponential growth phase cultures, resuspended in 300 µL of buffer containing 50 mM TRIS-HCl pH 8, 50 mM NaCl, 10% glycerol, and 1 mM ZnSO₄, and broken with 2 volumes of glass beads using a minibead beater. Total cell extracts were collected by piercing the bottom of the tube with a needle. Extracts were centrifuged for 5 min at 5000 *g* to remove the remaining glass beads and unbroken cells, and the supernatant constituted the total extract. A 2-µg sample of GST-PetR was added to 150 µL of total extracts (containing 150 µg of total protein) and incubated at 30°C. At the indicated times, 20 µL samples were removed, mixed with 7 µL of 4× Laemmli buffer and boiled for 10 min. A 5 µL sample was used for immunoblotting using α-PetR antibodies.

For *E. coli* assays, DH5α cells transformed with pN_petRP_Nat or pN_Nat were grown to late exponential phase (OD_{595nm} 1–2), and cells at an OD_{595nm} value of 20 were collected and processed as described above for *Synechocystis*. The pN_petRP_Nat plasmid was constructed by excising a 1478-bp fragment from pPETRP using *EcoRI* and *XhoI*, containing petRP and the 185-bp region upstream from *petR* ATG, and inserting it into pN_Nat (30) cut in the same way.

Band shift and gel retardation assays

Probes were PCR-synthesized using oligonucleotides 53 and 92 for the *petE* promoter (217 bp), 299 and 262 for the *petJ* promoter (103 bp), and 295 and 296

for *slr0601* (177 bp), introducing a *SaI* restriction site in all cases. The resulting DNA was digested with *SaI* and end-labeled with [α -³²P]-dCTP (3000 Ci mmol⁻¹) using Sequenase v2.0 (Product #70775Y, Affymetrix). The binding reaction was carried out in a final volume of 20 μ L containing 4 ng of labeled DNA and 4 μ g of salmon sperm DNA in 20 mM TRIS-HCl (pH 8.0), 150 mM KCl, 10 mM DTT, 1 mM EDTA, 10% glycerol, and different amounts (from 0.001 μ g to 1 μ g) of purified PetR. The mixtures were incubated for 30 min at room temperature and loaded on a Tris/Borate/EDTA (TBE) buffer non-denaturing 6% polyacrylamide gel (acrylamide:bisacrylamide 30:0.6). Electrophoresis was carried out at 4°C and 200 V in 0.25 $\times$ TBE. Gels were transferred to a Whatman 3 MM paper, dried, and autoradiographed using a Cyclone Phosphor System (Packard). Each experiment was performed at least three times with two independent PetR preparations.

RNA-seq

The quantity and quality of total RNA were evaluated using RNA electropherograms acquired by an Agilent 2100 Bioanalyzer; Agilent Technologies, Santa Clara, CA, USA), and library construction of cDNA molecules was carried out using Illumina Kapa Stranded Total RNA with a RiboZero Library Preparation Kit. The resulting DNA fragments (DNA library) were sequenced on an Illumina HiSeq 4000 platform using 150-bp paired-end sequencing reads. Sequencing was carried out by STAB VIDA (Lda, Lisbon, Portugal). Reads were aligned against the NCBI genome sequence for

Synechocystis (NC_000911.1, NC_005229.1, NC_005230.1, NC_005231.1, and NC_005232.1) using bowtie2 version 2.2.4 (55). Raw read counts were calculated using the HTSeq-count function of HTSeq framework 0.11.0 (56). The Bioconductor DESeq2 R software package (57) was used to detect differentially expressed genes. An adjusted p -value of <0.01 was considered significant. For RNA-seq density profiles, data were normalized using the BamCoverage function of DeepTools2 0.8.0 (58), and visualized with Integrative Genomics Viewer (IGV) 2.3.6 (59). The RNA-seq dataset is available at GEO accession GSE155385.

Phylogenomic analysis

16S RNA from 169 cyanobacterial genomes (Supplementary Material S1; all completed genomes from NCBI and those included in (60)) were aligned with MUSCLE 3.8.31, and a maximum likelihood (ML) phylogeny was estimated with IQ-TREE 1.6.11 (61) using the best-fitting model (automatically selected by the ModelFinder function) and selecting the phylogeny with the highest likelihood from those found among independent searches, with 1000 bootstrap replicates. *petE*, *petJ*, *petR*, and *petP* genes were searched using TBLASTN. Only hits with $\geq 50\%$ query coverage and $>25\%$ identity were considered homologues, and checked manually. Phylogenetic trees were displayed and edited using the Interactive Tree Of Life webtool iTOL v4 (62).

Oxygen evolution

Oxygen evolution was measured by a Clark-type oxygen electrode (Hansatech Chlorolab 2) using mid-logarithmic ($OD_{750nm} = 0.8-1$) cultures adjusted to $OD_{750nm} = 0.5$ in BG11C or BG11C-Cu+BCSA media supplemented with 20 mM $NaHCO_3$ and white LED light.

Acknowledgments

We thank Jose Luis Crespo, Manuel J. Mallén, Miguel Roldán, and Sandra Díaz-Troya for critical reading of the manuscript. This research was funded by grant numbers BIO2016-75634-P and PID2019-104513GB-I00 from the Ministerio de Economía y Competividad (MINECO) and the Agencia Estatal de Investigación (AEI), respectively, and by Junta de Andalucía Group BIO-284, co-financed by European Regional Funds (FEDER), to Francisco J. Florencio. RMGC is the recipient of a predoctoral contract from Ministerio de Educación, Cultura y Deporte, Spain (contract #FPU2015/05025).

References

1. Schirrmeister BE, de Vos JM, Antonelli A, Bagheri HC (2013) Evolution of multicellularity coincided with increased diversification of cyanobacteria and the Great Oxidation Event. *Proc Natl Acad Sci U S A* 110(5):1791–6.
2. Koehler MC, Buick R, Kipp MA, Stüeken EE, Zaloumis J (2018) Transient surface ocean oxygenation recorded in the ~2.66-Ga Jeerinah Formation, Australia. *Proc Natl Acad Sci U S A* 115(30):7711–7716.

- 726 3. Fischer WW, Hemp J, Johnson JE (2016) Evolution of Oxygenic
727 Photosynthesis. *Annu Rev Earth Planet Sci* 44(1):647–683.
- 728 4. Alcott LJ, Mills BJW, Poulton SW (2019) Stepwise Earth oxygenation is an
729 inherent property of global biogeochemical cycling. *Science*
730 366(6471):1333–1337.
- 731 5. Hamilton TL (2019) The trouble with oxygen: The ecophysiology of extant
732 phototrophs and implications for the evolution of oxygenic photosynthesis.
733 *Free Radic Biol Med* 140:233–249.
- 734 6. Fischer WW, Hemp J, Valentine JS (2016) How did life survive Earth's
735 great oxygenation? *Curr Opin Chem Biol* 31:166–178.
- 736 7. Ilbert M, Bonnefoy V (2013) Insight into the evolution of the iron oxidation
737 pathways. *Biochim Biophys Acta - Bioenerg* 1827(2):161–175.
- 738 8. Crichton RR (2019) *Biological inorganic chemistry□: a new introduction to*
739 *molecular structure and function* (Academic Press, London).
- 740 9. Tortell PD, Maldonado MT, Granger J, Price NM (1999) Marine bacteria
741 and biogeochemical cycling of iron in the oceans. *FEMS Microbiol Ecol*
742 29(1):1–11.
- 743 10. Behrenfeld MJ, Kolber ZS (1999) Widespread iron limitation of
744 phytoplankton in the south pacific ocean. *Science* 283(5403):840–843.

- 745 11. Finazzi G, Sommer F, Hippler M (2005) Release of oxidized plastocyanin
746 from photosystem I limits electron transfer between photosystem I and
747 cytochrome b6f in vivo. *Proc Natl Acad Sci U S A* 102(19):7031–7036.
- 748 12. Weigel M, et al. (2003) Plastocyanin is indispensable for photosynthetic
749 electron flow in *Arabidopsis thaliana*. *J Biol Chem* 278(33):31286–9.
- 750 13. Sandmann G, Böger P (1980) Copper-induced exchange of plastocyanin
751 and cytochrome c-533 in cultures of *Anabaena variabilis* and *Plectonema*
752 *boryanum*. *Plant Sci Lett* 17(4):417–424.
- 753 14. Sandmann G (1986) Formation of plastocyanin and cytochrome c-553 in
754 different species of blue-green algae. *Arch Microbiol* 145(1):76–79.
- 755 15. WOOD PM (1978) Interchangeable Copper and Iron Proteins in Algal
756 Photosynthesis: Studies on Plastocyanin and Cytochrome c-552 in
757 *Chlamydomonas*. *Eur J Biochem* 87(1):9–19.
- 758 16. Kropat J, et al. (2005) A regulator of nutritional copper signaling in
759 *Chlamydomonas* is an SBP domain protein that recognizes the GTAC core
760 of copper response element. *Proc Natl Acad Sci U S A* 102(51):18730–
761 18735.
- 762 17. Sommer F, et al. (2010) The CRR1 nutritional copper sensor in
763 *chlamydomonas* contains two distinct metal-responsive domains. *Plant Cell*
764 22(12):4098–4113.

- 765 18. Zhang L, McSpadden B, Pakrasi HB, Whitmarsh J (1992) Copper-
766 mediated regulation of cytochrome c553 and plastocyanin in the
767 cyanobacterium *Synechocystis* 6803. *J Biol Chem* 267(27):19054–19059.
- 768 19. Ghassemian M, Wong B, Ferreira F, Markley JL, Straus NA (1994)
769 Cloning, sequencing and transcriptional studies of the genes for
770 cytochrome c-553 and plastocyanin from *Anabaena* sp. PCC 7120.
771 *Microbiology* 140(5):1151–9.
- 772 20. Van der Plas J, et al. (1989) The gene for the precursor of plastocyanin
773 from the cyanobacterium *Anabaena* sp. PCC 7937: isolation, sequence
774 and regulation. *Mol Microbiol* 3(3):275–84.
- 775 21. Briggs LM, Pecoraro VL, McIntosh L (1990) Copper-induced expression,
776 cloning, and regulatory studies of the plastocyanin gene from the
777 cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Mol Biol* 15(4):633–
778 42.
- 779 22. Buikema WJ, Haselkorn R (2001) Expression of the *Anabaena* hetR gene
780 from a copper-regulated promoter leads to heterocyst differentiation under
781 repressing conditions. *Proc Natl Acad Sci U S A* 98(5):2729–2734.
- 782 23. Tous C, Vega-Palas MA, Vioque A (2001) Conditional expression of
783 RNase P in the cyanobacterium *Synechocystis* sp. PCC6803 allows
784 detection of precursor RNAs. Insight in the in vivo maturation pathway of
785 transfer and other stable RNAs. *J Biol Chem* 276(31):29059–29066.

- 786 24. Georg J, et al. (2017) Acclimation of Oxygenic Photosynthesis to Iron
787 Starvation Is Controlled by the sRNA IsaR1. *Curr Biol* 27(10):1425-
788 1436.e7.
- 789 25. Giner-Lamia J, et al. (2012) The CopRS Two-Component System Is
790 Responsible for Resistance to Copper in the Cyanobacterium
791 *Synechocystis* sp. PCC 6803. *Plant Physiol* 159(4):1806–1818.
- 792 26. Giner-Lamia J, López-Maury L, Florencio FJ (2016) Ni interferes in the
793 Cu-regulated transcriptional switch petJ/petE in *Synechocystis* sp. PCC
794 6803. *FEBS Lett* 590(20):3639–3648.
- 795 27. Giner-Lamia J, López-Maury L, Florencio FJ (2014) Global transcriptional
796 profiles of the copper responses in the cyanobacterium *Synechocystis* sp.
797 PCC 6803. *PLoS One* 9(9):e108912.
- 798 28. Foster AW, Patterson CJ, Pernil R, Hess CR, Robinson NJ (2012)
799 Cytosolic Ni(II) sensor in cyanobacterium: Nickel detection follows nickel
800 affinity across four families of metal sensors. *J Biol Chem* 287(15):12142–
801 12151.
- 802 29. Liu T, et al. (2004) A novel cyanobacterial SmtB/ArsR family repressor
803 regulates the expression of a CPx-ATPase and a metallothionein in
804 response to both Cu(I)/Ag(I) and Zn(II)/Cd(II). *J Biol Chem* 279(17):17810–
805 8.

- 806 30. Giner-Lamia J, López-Maury L, Florencio FJ (2015) CopM is a novel
807 copper-binding protein involved in copper resistance in *Synechocystis* sp.
808 PCC 6803. *Microbiologyopen* 4(1):167–185.
- 809 31. Krogh A, Larsson B, Von Heijne G, Sonnhammer ELL (2001) Predicting
810 transmembrane protein topology with a hidden Markov model: Application
811 to complete genomes. *J Mol Biol* 305(3):567–580.
- 812 32. Peacock SJ, Paterson GK (2015) Mechanisms of Methicillin Resistance in
813 *Staphylococcus aureus*. *Annu Rev Biochem* 84(1):577–601.
- 814 33. Durán R V, Hervás M, De La Rosa MA, Navarro JA (2004) The efficient
815 functioning of photosynthesis and respiration in *Synechocystis* sp. PCC
816 6803 strictly requires the presence of either cytochrome c6 or plastocyanin.
817 *J Biol Chem* 279(8):7229–33.
- 818 34. Zhang L, Pakrasi HB, Whitmarsh J (1994) Photoautotrophic growth of the
819 cyanobacterium *Synechocystis* sp. PCC 6803 in the absence of
820 cytochrome c553 and plastocyanin. *J Biol Chem* 269(7):5036–5042.
- 821 35. Lambrecht SJ, Steglich C, Hess WR (2020) A minimum set of regulators to
822 thrive in the ocean. *FEMS Microbiol Rev* 44(2):232–252.
- 823 36. Pence MA, et al. (2015) Beta-lactamase repressor Blal modulates
824 *Staphylococcus aureus* cathelicidin antimicrobial peptide resistance and
825 virulence. *PLoS One* 10(8). doi:10.1371/journal.pone.0136605.

- 826 37. Sala C, et al. (2009) Genome-wide regulon and crystal structure of Blal
827 (Rv1846c) from *Mycobacterium tuberculosis*. *Mol Microbiol* 71(5):1102–16.
- 828 38. Seo SW, et al. (2014) Deciphering Fur transcriptional regulatory network
829 highlights its complex role beyond iron metabolism in *Escherichia coli*. *Nat*
830 *Commun* 5:4910.
- 831 39. Pinochet-Barros A, Helmann JD (2020) Bacillus subtilis Fur is a
832 transcriptional activator for the PerR-repressed *pfeT* gene encoding an iron
833 efflux pump. *J Bacteriol* 202(8):e00697-19.
- 834 40. Delany I, Rappuoli R, Scarlato V (2004) Fur functions as an activator and
835 as a repressor of putative virulence genes in *Neisseria meningitidis*. *Mol*
836 *Microbiol* 52(4):1081–1090.
- 837 41. Glauning H, et al. (2017) Metal-dependent allosteric activation and
838 inhibition on the same molecular scaffold: The copper sensor CopY from:
839 *Streptococcus pneumoniae*. *Chem Sci* 9(1):105–118.
- 840 42. Belluzo BS, et al. (2019) An experiment-informed signal transduction
841 model for the role of the *Staphylococcus aureus* MecR1 protein in β -lactam
842 resistance. *Sci Rep* 9(1):19558.
- 843 43. Llarrull LI, Toth M, Champion MM, Mobashery S (2011) Activation of BlaR1
844 protein of methicillin-resistant *Staphylococcus aureus*, its proteolytic
845 processing, and recovery from induction of resistance. *J Biol Chem*

- 846 286(44):38148–38158.
- 847 44. Staude MW, et al. (2015) Investigation of signal transduction routes within
848 the sensor/transducer protein BlaR1 of *Staphylococcus aureus*.
849 *Biochemistry* 54(8):1600–10.
- 850 45. Tottey S, et al. (2012) Cyanobacterial metallochaperone inhibits
851 deleterious side reactions of copper. *Proc Natl Acad Sci U S A* 109(1):95–
852 100.
- 853 46. Tottey S, et al. (2008) Protein-folding location can regulate manganese-
854 binding versus copper- or zinc-binding. *Nature* 455(7216):1138–42.
- 855 47. Tottey S, Rich PR, Rondet SA, Robinson NJ (2001) Two Menkes-type
856 atpases supply copper for photosynthesis in *Synechocystis* PCC 6803. *J*
857 *Biol Chem* 276(23):19999–20004.
- 858 48. Castruita M, et al. (2011) Systems biology approach in *Chlamydomonas*
859 reveals connections between copper nutrition and multiple metabolic steps.
860 *Plant Cell* 23(4):1273–1292.
- 861 49. Wang X-Q, Jiang H-B, Zhang R, Qiu B-S (2013) Inactivation of the *petE*
862 gene encoding plastocyanin causes different photosynthetic responses in
863 cyanobacterium *Synechocystis* PCC 6803 under light-dark photoperiod
864 and continuous light conditions. *FEMS Microbiol Lett* 341(2):106–114.
- 865 50. Ardelean I, Matthijs HC., Havaux M, Joset F, Jeanjean R (2002)

- 866 Unexpected changes in photosystem I function in a cytochrome c 6 -
867 deficient mutant of the cyanobacterium *Synechocystis* PCC 6803. *FEMS*
868 *Microbiol Lett* 213(1):113–119.
- 869 51. Ivanov AG, et al. (2012) Restricted capacity for PSI-dependent cyclic
870 electron flow in $\Delta petE$ mutant compromises the ability for acclimation to
871 iron stress in *Synechococcus* sp. PCC 7942 cells. *Biochimica et*
872 *Biophysica Acta - Bioenergetics*, pp 1277–1284.
- 873 52. Stanier RY, Deruelles J, Rippka R, Herdman M, Waterbury JB (1979)
874 Generic Assignments, Strain Histories and Properties of Pure Cultures of
875 Cyanobacteria. *Microbiology* 111(1):1–61.
- 876 53. García-Domínguez M, Florencio FJ (1997) Nitrogen availability and
877 electron transport control the expression of *glnB* gene (encoding PII
878 protein) in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Mol Biol*
879 35(6):723–734.
- 880 54. J.F. S, Russell D (2001) *Molecular Cloning: A Laboratory Manual* (3-
881 *Volume Set*).
- 882 55. Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie
883 2. *Nat Methods* 9(4):357–359.
- 884 56. Anders S, Pyl PT, Huber W (2015) Genome analysis HTSeq-a Python
885 framework to work with high-throughput sequencing data. 31(2):166–169.

886 57. Love MI, Huber W, Anders S (2014) Moderated estimation of fold change
887 and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15(12).
888 doi:10.1186/s13059-014-0550-8.

889 58. Ramírez F, et al. (2016) deepTools2: a next generation web server for
890 deep-sequencing data analysis. *Nucleic Acids Res* 44(W1):W160–W165.

891 59. Robinson JT, et al. (2011) Integrative genomics viewer. *Nat Biotechnol*
892 29(1):24–26.

893 60. Shih PM, et al. (2013) Improving the coverage of the cyanobacterial
894 phylum using diversity-driven genome sequencing. *Proc Natl Acad Sci U S*
895 *A* 110(3):1053–1058.

896 61. Nguyen L-T, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: A
897 Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood
898 Phylogenies. *Mol Biol Evol* 32(1):268–274.

899 62. Letunic I, Bork P (2019) Interactive Tree Of Life (iTOL) v4: recent updates
900 and new developments. *Web Serv issue Publ online* 47(W1):W256–W259.
901
902

Figure Legends

Figure 1. PetRP regulatory system controls *petJ/petE* switch in *Synechocystis* sp. PCC 6803.

A. RNA blot analysis of *petJ*, *petE* and *copM* in WT, PETR and PETP strains in response to 0.5 μ M copper addition. Total RNA was isolated from cells grown in BG11C-Cu medium at the indicated times after addition of 0.5 μ M of copper. The filters were hybridized with *petJ*, *petE* and *copM* probes and subsequently stripped and re-hybridized with a *rnpB* probe as a control.

B. Immunoblot analysis of Cyt_c₆, PC, CopM and GSI in WT, PETR and PETP strains in response to 0.5 μ M copper addition. Cells were grown in BG11C-Cu medium and cells were harvested at the indicated times addition of 0.5 μ M of copper. 5 μ g of total protein from soluble extracts was separated by 15 % SDS-PAGE and subjected to immunoblot to detect PC, Cyt_c₆, CopM or GSI.

C. Quantification of PC levels in WT, PETR and PETP strains grown in BG11C-Cu +BCSA (blue bars), BG11C-Cu (orange bars) or BG11C (grey bars). 10 μ g of total soluble proteins were loaded and compared to serial dilutions (100%, 50%, 25% and 12.5%) of the WT extract prepared from cells grown in BG11C. Data are the mean $\pm$ SE of three biologically independent experiments. Asterisk indicate significant difference to WT in the same condition (*t*-test; *p*<0.05).

D. Quantification of Cyt_c₆ levels in WT, PETR and PETP strains grown in BG11C-Cu +BCSA (blue bars), BG11C-Cu (orange bars) or BG11C (grey bars). 10 µg of total soluble proteins were loaded and compared to serial dilutions (100%, 50%, 25% and 12.5%) of the WT extract prepared from cells grown in BG11C-Cu+BCSA. Data are the mean ± SE of three biologically independent experiments. Asterisk indicate significant difference to WT in the same condition (*t*-test; *p*<0.05).

Figure 2. PetP is a copper activated protease that regulates PetR levels.

A. Immunoblot analysis of PetR in WT, PETR and PETP strains in response to 0.5 µM copper addition. Cells were grown in BG11C-Cu medium and harvested at the indicated times after addition of 0.5 µM of copper. Whole cells were loaded (0.2 OD_{750nm}), separated by 15 % SDS-PAGE and subjected to immunoblot to detect PetR or GSI as loading control.

B. Immunoblot analysis of Cyt_c₆, PC and PetR in WT, PETR and PETP strains grown in BG11C-Cu+BCSA (B), BG11C-Cu (-) or BG11C. 10 µg of total soluble proteins were separated by 15 % SDS-PAGE and subjected to immunoblot blot to detect PC, Cyt_c₆, or GSI as loading control. For PetR whole cell extracts were used and 20 µl (equivalent to 0.2 OD_{750nm}) were loaded per lane.

C. Quantification of PetR levels in WT, PETR and PETP strains grown in BG11C-Cu +BCSA (blue bars), BG11C-Cu (orange bars) or BG11C (grey bars). 20 µl of the whole cells extract were loaded and compared to serial

dilutions (100%, 50%, 25% and 12.5%) of the WT extract prepared from cells grown in BG11C-Cu+BCSA. Data are the mean $\pm$ SE of three biologically independent experiments. Asterisk indicate significant difference to WT in the same condition (t -test; $p > 0.05$).

D. GST-PetR was incubated for the indicated time with total extracts prepared from WT, PETR and PETP strains grown in BG11C-Cu. 0.5 μ M CuSO₄ was added to the extracts as indicated. Samples were taken at the indicated times, mixed with Laemmli buffer, boiled, separated in 12% SDS-PAGE gels and GST-PetR was detected using PetR antibodies.

E. GST-PetR was incubated for the indicated time with whole cell extracts prepared from *E. coli* carrying pN_petRP or pN_Nat (an empty plasmid) grown in M9 minimal media. 0.5 μ M CuSO₄ was added to the extracts as indicated. Samples were taken at the indicated times, mixed with Laemmli buffer, boiled, separated in 12% SDS-PAGE gels and GST-PetR was detected using PetR antibodies

Figure 3. PetR binds to *petJ* and *petE* promoters.

A. Electrophoretic mobility shift assay using recombinant PetR to a 103 bp *petJ* promoter probe. The indicated PetR concentration were used in each lane.

B. Electrophoretic mobility shift assay using recombinant PetR to a 217 bp *petE* promoter probe. The indicated PetR concentration were used in each lane.

- C. Sequence logo identified with MEME using upstream sequences (300 bp) from *petE* and *petJ* genes from genomes that also contained *petRP*.
- D. Sequence logo identified with MEME using upstream sequences (300 bp) from *petJ* genes from genomes that also contained *petRP*.
- E. Sequence logo identified with MEME using upstream sequences (300 bp) from *petE* from genomes that also contained *petRP*.
- F. Alignment of the *petJ* and *petE* promoter sequences from *Synechocystis*. Transcriptional start sites are in capital letters and underlined, -10 boxes are yellow underlined and conserved nucleotides from the motif found in C are in red.

Figure 4. Physiological characterization of *petR* and *petP* mutant strains.

- A. Growth of WT, PETE, PETJ, PETER, PETP and PETERP strains in different copper availability regimes. Tenfold serial dilutions of a 1 µg chlorophyll mL⁻¹ cells suspension were spotted onto BG11C-Cu+BCSA, BG11C-Cu o BG11C. Plates were photographed after 5 days of growth
- B. Oxygen evolution measured using a Clark electrode at increasing light intensities in exponential growing cultures (OD_{750nm}= 0.5-1) of WT (●), PETJ (●), PETE (●), PETER (●) and PETP (●) strains grown in BG11C-Cu+BCSA. Data are the mean ± SE of at least three biologically independent experiments.
- C. Oxygen evolution measured using a Clark electrode at increasing light intensities in exponential growing cultures (OD_{750nm}= 0.5-1) of WT (●),

994 PETJ (●), PETE (●), PETR (●) and PETP (●) strains grown in BG11C.

995 Data are the mean $\pm$ SE of at least three biologically independent
996 experiments.

997

998 **Figure 5. Model of the regulatory mechanism for *petE/petJ* switch mediated**
999 **by PetRP.**

1000

A

B

C

D

E

A

B

C