PH domain and leucine rich repeat phosphatase 1 (Phlpp1) suppresses parathyroid hormone receptor 1 (Pth1r) expression and signaling during bone growth

Samantha R. Weaver^{1*}, Earnest L. Taylor^{1*}, Elizabeth L. Zars¹, Katherine M. Arnold¹, Elizabeth W. Bradley², Jennifer J. Westendorf^{1,3}

1. Department of Orthopedic Surgery, Mayo Clinic, Rochester, MN, USA. 2. Department of Orthopedic Surgery and Stem Cell Institute, University of Minnesota, Minneapolis, MN. 3. Department of Biochemistry and Molecular Biology, Mayo Clinic, Rochester, MN, USA.

*indicates equal contribution of authors

Corresponding author:

Jennifer J. Westendorf, PhD

Mayo Clinic

200 First Street SW

Rochester, MN 55905

Phone: 507-538-5651

Email: westendorf.jennifer@mayo.edu

Running Title

Phlpp1 suppresses Pth1r in chondrocytes

Disclosures

The authors declare that they have no disclosures.

ABSTRACT

Endochondral ossification is tightly controlled by a coordinated network of signaling cascades including parathyroid hormone (PTH). PTH domain and leucine rich repeat phosphatase (Phlpp1) affects endochondral ossification by suppressing chondrocyte proliferation in the growth plate, longitudinal bone growth, and bone mineralization. As such, Phlpp1^{-/-} mice have shorter long bones, thicker growth plates, and proportionally larger growth plate proliferative zones. The goal of this study was to determine how Phlpp1 deficiency affects PTH signaling during bone growth. Transcriptomic analysis revealed greater Pth1r expression and H3K27ac enrichment at the Pth1r promoter in Phlpp1-deficient chondrocytes. PTH(1-34) enhanced and PTH(7-34) attenuated cell proliferation, cAMP signaling, CREB phosphorylation, and cell metabolic activity in Phlpp1-inhibited chondrocytes. To understand the role of Pth1r action in the endochondral phenotypes of Phlpp1-deficient mice, Phlpp1^{-/-} mice were injected with Pth1r ligand PTH(7-34) daily for the first four weeks of life. PTH(7-34) reversed the abnormal growth plate and long bone growth phenotypes of Phlpp1^{-/-} mice but did not rescue deficits in bone mineral density or trabecular number. These results demonstrate that elevated Pth1r expression and signaling contributes to increased proliferation in Phlpp1^{-/-} chondrocytes and shorter bones in Phlpp1-deficient mice. Our data reveal a novel molecular relationship between Phlpp1 and Pth1r in chondrocytes during growth plate development and longitudinal bone growth.

Keywords: Molecular pathways – development, Chondrocyte and cartilage biology, growth plate, PTH/Vit D/FGF23

INTRODUCTION

During development, long bones lengthen and harden through the process of endochondral ossification⁽¹⁾. In the epiphyseal growth plate, chondrocytes proliferate and undergo hypertrophy to drive appositional bone lengthening. Vascularization of the early cartilaginous bone brings osteoblast and osteoclast precursors which ossify, model, and remodel bone. Most hypertrophic chondrocytes will undergo apoptosis, but some survive and contribute to the osteogenic pool in bone marrow⁽²⁻⁴⁾. These dynamic and complex processes are orchestrated by numerous extracellular factors that induce transcriptional events and intracellular signaling.

PH domain and leucine rich repeat phosphatases (Phlpp1 and Phlpp2) control cell proliferation and survival through posttranslational modification of several intracellular substrates⁽⁵⁻⁷⁾. Originally identified as terminators of Akt signaling⁽⁸⁾, further work showed Phlpp1/2 regulation of protein kinase C (PKC)^(9,10), ribosomal protein S6 kinase (S6K)⁽¹¹⁾, and mitogen-activated protein kinase (Mapk/Erk)⁽⁵⁾. Additionally, Phlpp1/2 can translocate to the nucleus to modulate histone acetylation and phosphorylation⁽¹²⁾. Phlpp1/2 have been implicated as tumor suppressors^(9,13,14), but also regulate metabolic processes⁽¹⁵⁾, inflammation⁽¹⁶⁾, and tissue regeneration following injury⁽¹⁷⁻²⁴⁾. Phlpp1 is overexpressed in human osteoarthritic tissue and Phlpp1 inactivation improves murine post-traumatic osteoarthritis⁽¹⁷⁾. Phlpp1 deletion accelerates chondrocyte maturation in vitro, preserves articular cartilage in vivo, and increases mobility in mice with post-traumatic osteoarthritis^(17,18). In the developing appendicular skeleton, Phlpp1 controls chondrocyte proliferation and bone lengthening⁽⁵⁾. Phlpp1-depleted mice have short long bones and low bone mass⁽⁵⁾ and Phlpp1-deficient chondrocytes proliferate more and express higher levels of growth factors such as Fgf18 and growth factor receptors, including Pth1r.

Pth1r is a G-protein-copuled receptor for parathyroid hormone (PTH) and parathyroid hormone-related protein (PTHrP). Dysregulation of signaling from Pth1r is at the root of several genetic diseases and is the target of bone anabolic therapies⁽²⁵⁾. Pth1r activates Gα subunits to regulate cyclic AMP production⁽²⁵⁾, as well as Akt⁽²⁶⁾, PKC⁽²⁷⁾, and Erk1/2⁽²⁸⁾ activation. Pth1r is expressed in various musculoskeletal cells, including mesenchymal stem cells⁽²⁹⁾, osteocytes⁽³⁰⁾, osteoblasts^(31,32), and chondrocytes^(33,34). In the growth plate, Pth1r activation in proliferating cells inhibits premature hypertrophy and thus is crucial for

proper development prenatally and during early life^(1,35). PTH peptides (1-34 and 7-34) that bind Pth1r have been developed for clinical and research use. Both PTH(1-34) and PTH(7-34) cause Pth1r internalization with identical kinetics. However, only PTH(1-34) activates adenylyl cyclase and phospholipase C (PLC)⁽³⁶⁾. When injected intermittently, PTH(1-34) increases bone mass and bone formation, and reduces fracture risk⁽³⁷⁾. By contrast, continuous infusion of PTH(1-34) has catabolic effects on bone⁽³⁸⁾. PTH(7-34) does not appear to have any effects on bone when administered alone⁽³⁹⁾. We previously showed that Pth1r mRNA levels were elevated in Phlpp1 depleted chondrocytes⁽⁵⁾. Here, we demonstrate that Phlpp1 regulates Pth1r transcription and signaling and show how these pathways cooperate to regulate endochondral ossification in mice. PTH(7-34) administration from birth through four weeks of age reversed the abnormal bone growth phenotype characteristic of Phlpp1^{-/-} mice. In vitro, PTH(7-34) reversed the increases in cell proliferation, CREB phosphorylation, metabolic activity, and cAMP signaling evident in Phlpp1^{-/-} chondrocytes. Taken together, our results demonstrate that Phlpp1 suppresses Pth1r expression and signaling in chondrocytes to regulate endochondral ossification.

MATERIALS AND METHODS

In vitro Phlpp inhibitor experiments: For experiments involving small-molecule Phlpp inhibitors (Phlppi), cells were treated with 25μM NSC117079, 25μM NSC45586 (Glaxo Laboratories), or vehicle (0.0005% DMSO) for the indicated time.

In vitro PTH experiments: Primary WT or Phlpp1^{-/-} IMCs were plated and allowed to adhere overnight. Medium was replaced with DMEM containing either PTH(1-34), PTH(7-34) (Bachem), or vehicle (0.1% BSA in PBS) at the concentrations indicated.

ATDC5 cell culture: ATDC5 cells were seeded at a density of 5 x 10⁵ cells/well in a six-well plate and incubated overnight to allow for cell adhesion in DMEM supplemented with 10% FBS and 100 units/mL penicillin, 100 μg/mL streptomycin, and 0.25 μg/mL Amphotericin B (1% antibiotic/antimycotic; Gibco 15240112). Phlppi treatments were added as indicated. Each experiment included at least three technical replicates and was repeated at least three times. Results from a representative experiment are shown.

Immature chondrocyte (IMC) cell culture: Primary immature chondrocytes (IMCs) were collected from 5-day-old WT or Phlpp1^{-/-} mice as previously described^(5,40). To evaluate mRNA and proteins, IMCs were isolated from cartilage with an overnight digestion in 0.5 mg/mL collagenase in serum-free culture medium, washed in PBS, and lysed as described below. When cells required treatment in culture, chondrocytes were plated in monolayer at a seeding density of 5 x 10⁵ cells/well in a six-well plate and cultured in DMEM supplemented with 2% FBS and 1% antibiotic/antimycotic. For micromass experiments, 10 µL drops of IMC suspensions (2 x 10⁷ cells/mL) were plated in DMEM. Three micromasses were plated in each well of a 6-well plate. After 1 hour, micromasses were covered with 2 mL DMEM + 2% FBS + 1% antibiotic/antimycotic and allowed to grow for 3 days, after which time the culture medium was changed to 2% FBS + 1% antibiotic/antimycotic + 1x Insulin/Transferrin/Selenium (ITS) + 0.05 mg/mL ascorbic acid + 10 mM β-glycerophosphate + respective treatment^(5,41). Littermates were pooled according to genotype for each experiment. Each experiment was repeated at least three times and represents at least n=3 biological replicates. Data from a representative experiment are shown.

Transcriptional inhibition: Primary chondrocytes were plated and allowed to adhere overnight as described above. Actinomycin D (5µM, Gibco 11805017) was added for six hours and then replaced with medium containing Phlpp1 or vehicle for 24 hours. Cells were collected for RNA or protein extraction.

RNA isolation and real-time PCR: Total RNA was extracted from cell lines and primary chondrocytes using TRIzol (Invitrogen) and chloroform and 2µg was reverse transcribed using the iScript cDNA Synthesis Kit (Bio-Rad). Resulting cDNAs were used to assay gene expression via real-time PCR using the following gene-specific primers: Pth1r (5'-ACTTAGGCCGTTTCCTGTCC-3', 5'-GAGGAGCTGACTCAGGTTGG-3'), Ywhaz (5'-GCCCTAAATGGTCTGTCACC-3', 5'-GCTTTGGGTGTGACTTAGCC-3'). Fold changes in gene expression were calculated using the 2^{-ΔΔCt} method relative to control after normalization of gene-specific Ct values to Ywhaz Ct values⁽⁴²⁾.

Western blotting: Cell lysates were collected in a buffered SDS solution (0.1% glycerol, 0.01% SDS, 0.1M Tris, pH 6.8) on ice. Total protein concentrations were obtained using the Bio-Rad DC Assay (Bio-Rad). Proteins (20µg) were resolved by SDS-PAGE and transferred to a polyvinylidene difluoride membrane. Western blotting was performed with antibodies (1:1000 dilution) for Phlpp1 (Sigma 07-1341), Pth1r

(Sigma SAB4502493), H3pS10 (Abcam ab5176), H3K9ac (Abcam ab10812), H3K27ac (Abcam ab4729), H3pS28 (Cell Signaling Technology 9713S), H3K9K14ac (Cell Signaling Technology 9677S), total H3 (Abcam ab1791), pCREB-S133 (Cell Signaling Technology 9198S), total CREB (Cell Signaling Technology 4820), and Actin (Sigma A4700) and corresponding secondary antibodies conjugated to horseradish peroxidase (HRP) (Cell Signaling Technology). Antibody binding was detected with the Supersignal West Femto Chemiluminescent Substrate (Pierce Technology, Rockford, IL).

Live Cell Imaging: Primary chondrocytes (5×10^3 cells/well) were allowed to adhere in monolayer in a 48-well plate and cultured overnight. Medium was replaced with DMEM containing either 10nM PTH(1-34), 10nM PTH(7-34), or vehicle (0.1% BSA in PBS). Cell confluency was detected in real-time with the IncuCyte S3 Live Cell Analysis System (Roche Applied Sciences, Indianapolis, IN), with four captures per well every hour for 48 hours.

Cyclic AMP assay: WT and Phlpp1^{-/-} IMCs were allowed to attach to a 6-well plate overnight and then treated with 100nM PTH(1-34) or 100nM PTH(7-34) for 15 minutes. Cyclic AMP (cAMP) concentrations in IMC protein extracts (100 µg) were determined by ELISA (Cayman Chemical 581001).

Chondrocyte metabolic activity assay: Primary chondrocytes (1×10^3 cells/well) were cultured in monolayer in 96-well flat-well plates for 24 hours with 10nM PTH(1-34) or 10nM PTH(7-34). CellTiter 96® Aqueous One Solution Cell Proliferation Assays (Promega) were performed according to manufacturer specifications. Briefly, 20 µL of a reagent containing 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxy-phenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) was added to each well for 24 h at 37°C. Spectrophotometer readings were taken at OD490.

Micromass staining with Alcian Blue: Micromasses of WT or Phlpp1^{-/-} IMCs were plated as described above and incubated for three days in DMEM + 2% FBS. Medium was replaced with DMEM containing 1x ITS, 0.05 mg/mL ascorbic acid, and 10 mM β-glycerophosphate in the presence of 10nM PTH(1-34), 10nM PTH(7-34) for 9 days, with media changes every three days. Cells were washed with PBS, and 0.5% Alcian Blue (Sigma A5268) was applied for 2 hours.

Histology and Immunohistochemistry / In Situ Hybridization: Right tibiae were decalcified for 14 days in 15% EDTA, dehydrated, and embedded in paraffin for sectioning. Sections (5 µm thick) were stained with Safranin O (Sigma S2255) and Fast Green (Sigma F7252)⁽⁵⁾ and chosen for analysis based on anatomical landmarks in the bone. Cross-sectional areas of the proliferative and hypertrophic zones in the proximal tibia growth plate were found in MatLab (version R2019b)⁽⁴³⁾. A polygon was drawn around the respective zone and a binary image was formed where pixels inside the region of interest were set to one, and all other pixels were set to zero. The area of the binary image was calculated using the *bwarea* command and scaled to 3.45×10^{-6} mm² per pixel. Immunohistochemistry was performed with antibodies diluted in 1% bovine serum albumin in tris-buffered saline directed to Pth1r (1:50 dilution, Sigma SAB4502493) or with a nonspecific IgG (control). Chromagens were detected with a polyvalent secondary HRP kit (Abcam, ab93697) and 3,3'-diaminobenzidine (DAB) (Sigma-Aldrich, D3939). In situ hybridization was performed using the RNAScope[®] 2.5 HD Assay - Brown (ACD Biotechnne). Briefly, slides were deparaffinized and epitope retrieval was achieved using a Custom Pretreatment Reagent provided by the manufacturer. Pth1r (ACD Biotechnne 426191) and Col10a1 (ACD Biotechnne 426181) riboprobes were hybridized to the tissue for two hours and detected using RNAScope[®] 2.5 HD Assay detection reagents. DapB (ACD Biotechnne 310043) was the negative control. Imaging was completed using a Zeiss LSM 900 Confocal Microscope.

Chromatin Immunoprecipitation (ChIP) Sequencing and PCR: Primary chondrocytes were cultured in monolayer 10cm dishes for 24 hours. Cells were treated with 25µM NSC117079 or vehicle for 24 hours and then prepared for ChIP-Seq (2×10^7 cells / sample). ChIP-seq was performed as previously described⁽⁴¹⁾, utilizing the anti-H3K27ac antibody (CST 8173BC (D5E4)). Libraries were prepared from 10ng DNA using the ThruPLEX DNA-seq Kit V2 (Rubicon Genomics, Ann Arbor, MI) and sequenced to 51 base pairs from both ends on an Illumina HiSeq 4000 instrument at the Mayo Clinic Medical Genome Facility Sequencing Core. ChIP-qPCR was performed as previously described utilizing 2 µg of anti-H3K27ac (Abcam ab4729) or YY1 (Active Motif 61780)⁽⁴¹⁾. Primers used for ChIP-qPCR were: Pth1r Promoter (5'-CCGCAGACTGACACGGAGAC-3', 5'-CGACATTCATGGCAAGGCGG-3'), -1.47kb (5'-TGTGGAGTATCACACACTGCG-3', 5'-TTGGGTAAAGCGGTCCCATT-3'). The -1.47kb primer is labeled

to indicate the distance from the Pth1r promoter primer. The Pth1r promoter and upstream site were identified using Ensembl Genome Browser.

PTH injections to mice: All mice were maintained in an accredited facility with 12-h light/dark cycle and supplied with food (PicoLab[®] Rodent Diet 20 5058, LabDiet) *ad libitum*⁽⁴⁴⁾. All animal research was performed according to National Institute of Health and the Institute of Laboratory Animal Resources, National Research Council guidelines and the Mayo Clinic Institutional Animal Care and Use Committee approved all animal studies. Animals for these experiments were generated by crossing Phlpp1^{+/-} males and females. Beginning on the day after birth, entire litters of pups were injected daily with 100 µg/kg body weight/day PTH(7-34) or vehicle (0.1% BSA in PBS) through four weeks of age⁽⁴⁵⁾. Animals were genotyped at three weeks of age⁽⁵⁾. Sample size was determined based on a power calculation that provided an 80% chance of detecting a significant difference (P<0.05). Final male group sizes were: WT+Vehicle (n=5), WT+PTH(7-34) (n=5), HET+Vehicle (n=9), HET+PTH(7-34) (n=6), Phlpp1^{-/-}+Vehicle (n=4), and Phlpp1^{-/-}+PTH(7-34) (n=6). Final female group sizes were: WT+Vehicle (n=7), WT+PTH(7-34) (n=6), HET+Vehicle (n=8), HET+PTH(7-34) (n=7), Phlpp1^{-/-}+Vehicle (n=7), and Phlpp1^{-/-}+PTH(7-34) (n=8). At four weeks old, mice were euthanized and limbs were collected, fixed overnight in 10% neutral buffered formalin, and stored in 70% ethanol until further analysis.

X-ray Imaging: Radiographs of hind limbs were collected using a Faxitron X-ray imaging cabinet (Faxitron Bioptics, Tuscon, AZ). Limb length was measured on radiographs using ImageJ (1.52a) software.

MicroCT: Micro-CT imaging of the femur and tibiae were performed using a SkyScan 1276 scanner (Bruker, Kontich, Belgium). Bones were fixed in 10% NBF before storage in 70% ethanol. Scans were performed at 55kV, 200 µA, 10 µm pixel resolution, 0.4° rotation steps for 360°, 4 frames average imaging with a 0.25mm A1 filter. The acquired scans were reconstructed using the Skyscan NRecon software with beam hardening and post-alignment correction. Trabecular and cortical analyses of the femur were performed using Bruker CtAN software. The datasets were oriented in 3D to vertically align the longitudinal axis of each femur. As the bones were different lengths, a region of interest (ROI) for trabecular bone was defined as 5% the length of each bone, beginning 8% bone's-length distance away from the distal growth plate. A gray-value threshold of 70 was applied to trabecular segmentations.

Quantified outcomes were bone volume / total volume (BV/TV), trabecular thickness (TbTh), trabecular number (TbN), trabecular spacing (TbSp), and bone mineral density (BMD)⁽⁴⁶⁾. For cortical bone analyses, the ROI was defined as 5% of total femur length beginning at the femoral midpoint. Quantified outcomes for cortical bone were tissue mineral density (TMD), cortical thickness (CTh), total tissue cross-sectional area (TtAr), cortical bone area (CtAr), and cortical area fraction (CtAr/TtAr)⁽⁴⁶⁾. Semi-automatic segmentation of the proximal tibial growth plate was performed using 3D Slicer and a 3D region-growing method⁽⁴⁷⁾ as previously described⁽⁴⁸⁾. For one image slice, seeds were placed in the growth plate, bone, and empty space. This was repeated approximately 10 times in both the coronal and sagittal planes to define the segments in 3D. The “grow from seeds” feature was then applied to map the input image to user-specified segments. “Joint smoothing” was used on the resulting segmentation to preserve segment interfaces while removing noise. The growth plate segment was exported as a 3D patch object for processing in MatLab⁽⁴³⁾. In MatLab, thickness measurements were taken by applying a query grid with 50 micron spacing, and sampling the z values of the patch object at the query points, resulting in approximately 3500 measurements across the growth plate. From these values, thickness distribution was plotted as a color map onto the patch object.

Plasma and urine biochemical analyses: At four weeks of age, WT and Phlpp1^{-/-} mice were euthanized. Groups analyzed were: WT male (n=7), WT female (n=7), Phlpp1^{-/-} male (n=8), Phlpp1^{-/-} female (n=8). Whole blood was collected into EDTA-treated tubes (BD Biosciences 365974) and centrifuged for 10 minutes at 1,500 x g at 4°C to isolate plasma. Spot urine samples were collected immediately prior to euthanasia. Plasma intact PTH(1-84) was measured with an ELISA kit (Immutopics Inc., San Clemente, CA, USA). Urine creatinine and cAMP levels were measured via ELISA (Enzo Life Sciences, Farmingdale, NY, USA and Cayman Chemical, Ann Arbor, MI, USA). Urine was diluted 1:50 to fit within the standard curve of the creatinine kit and 1:300 for the cAMP kit.

BrdU injections to mice and immunohistochemical analysis: Beginning on postnatal day 1 (P1), entire litters of pups from Phlpp1^{+/-} breeding pairs were injected with either 100 µg/kg body weight/day PTH(7-34) or vehicle (0.1% BSA in PBS) daily through P5. Males and females were combined because no sex differences were observed in earlier studies. The final group sizes were: WT+Vehicle (n=6), WT+PTH(7-

34) ($n=5$), Phlpp1^{-/-}+Vehicle ($n=6$), Phlpp1^{-/-}+PTH(7-34) ($n=8$). On P5, pups were injected with 100 mL / 100 g body weight BrdU labeling reagent (Invitrogen 00-0103). Two hours later, pups were euthanized and hind limbs were collected, decalcified for 7 days in 15% EDTA, and processed for histology as described above. BrdU-positive cells were identified using the BrdU IHC kit (Millipore 2760). A representative area was chosen in the proximal tibia growth plate as described⁽⁴⁹⁾, and the number of BrdU-positive cells per total cell number was calculated.

Statistical analysis: Statistics were performed in Prism GraphPad (Version 8) using Student's t-test, one-way ANOVA, or two-way ANOVA as appropriate with the necessary post-hoc tests for multiple comparisons. Data are depicted as individual points with SD bars ($n=3$) or boxplots showing the median, interquartile distance, and min/max values ($n>3$).

RESULTS

Phlpp1 inhibition increases Pth1r expression in chondrocytes: We previously reported that Pth1r mRNA levels are elevated in Phlpp1^{-/-} chondrocytes⁽⁵⁾. Given the important role of both Phlpp1 and Pth1r in endochondral bone formation, we sought to validate Pth1r expression changes in Phlpp1-inactivated cells. Phlpp inhibitors NSC117079 and NSC45586 increased Pth1r mRNA and protein expression in ATDC5 cells (**Figs 1A,B; Supp Fig 1A**) and IMCs (**Figs 1C,D; Supp Fig 1B**) within 24 hours. Elevated Pth1r expression was also confirmed by RT-PCR, immunoblotting, and immunohistochemistry in Phlpp1^{-/-} compared to WT mice (**Figs 1E-H; Supp Fig 1C**). In situ staining for Pth1r was more intense in the proximal tibial growth plate of Phlpp1^{-/-} mice than WT mice at both the RNA and protein levels (**Fig 1G,H; Supp Fig 2A**). Thus, Phlpp1 inhibition rapidly elevates Pth1r expression in vitro and in vivo.

Phlpp1 regulates Pth1r through transcription: We next identified mechanisms by which Phlpp1 inactivation increases Pth1r levels. Phlpp1 is known to modulate histone 3 (H3) acetylation and phosphorylation^(12,50). Phlpp inhibitors NSC117079 and NSC45586 increased phosphorylation (p) of H3S10 (H3pS10) as well as acetylation (ac) of H3K9 and H3K27 in primary IMCs, but not H3pS28 or

H3K9K14ac (**Fig 2A; Supp Fig 1D-H**). Similar results were observed in *Phlpp1*^{-/-} IMCs (**Fig 2A; Supp Fig 1I-M**). ChIP-seq revealed several H3K27ac-enriched regions in control and NSC117079-treated cells. The abundance of H3K27ac was greater within the *Pth1r* promoter in IMCs treated with the Phlpp inhibitor NSC117079 (**Fig 2B**). When compared to publicly available datasets, H3K27ac peaks are similar to those found on embryonic limbs (**Supp Fig 3**)⁽⁵¹⁾. ChIP-PCR of chromatin from IMCs treated with NSC117079 and NSC45586 for 24 hours verified robust enrichment of H3K27ac in the *Pth1r* promoter of IMCs (**Fig 2C**). *Phlpp1*^{-/-} IMCs also had greater basal enrichment of H3K27ac in the *Pth1r* promoter compared to WT IMCs (**Fig 2D**). The RNA polymerase inhibitor, actinomycin D, prevented the increase in *Pth1r* mRNA and protein levels, indicating transcriptional regulation (**Fig 2F; Supp Fig 1N**). A search of two databases identified binding sites for over 70 transcription factors in the region of the *Pth1r* promoter that was hyperacetylated after Phlpp1 inhibition (**Supp Table 1**). The transcription factor Ying Yang 1 (YY1) was chosen for further evaluation because it was present in both databases, had multiple potential binding sites in the H3K27ac-enriched peak of the *Pth1r* promoter, and was detectable in primary chondrocytes via qPCR (data not shown). Primary chondrocytes treated with the Phlpp inhibitor NSC117079 for 30 minutes had increased binding of YY1 in the *Pth1r* promoter (**Fig 2F**). Together, these data demonstrate that Phlpp inactivation increases transcription of *Pth1r*.

PTH(7-34) reverses short limb length in Phlpp1^{-/-} mice: *Phlpp1* knockout mice have shorter long bones than WT littermates⁽⁵⁾ but similar plasma PTH and urine cAMP concentrations (**Supp Fig 4**). To determine the effect of *Pth1r* signaling on Phlpp1-mediated long bone growth, whole litters of pups from *Phlpp1*^{+/-} male and female pairs were injected daily with vehicle or PTH(7-34), which induces receptor internalization but not signaling, from the day of birth through four weeks of age. As expected, *Phlpp1*^{-/-} mice injected with vehicle had shorter tibiae (**Fig 3A,C,E,G; Supp Fig 5E**) and femurs (**Fig 3B,D,F,H; Supp Fig 5F**), with *Phlpp1*^{+/-} (HET) showing an intermediate phenotype between WT and *Phlpp1*^{-/-} mice (**Supp Fig 5A,B,E,F**). *Phlpp1*^{-/-} mice also had shorter tail-to-snout body length (**Supp Fig 5C,G; Supp Fig 6A,C**) and lower body weight (**Supp Figure 5D,H; Supp Fig 6B,D**) than WT mice. PTH(7-34) did not alter bone growth in WT mice but rescued the short femur and tibia lengths (**Fig 3; Supp Figure 5**) as well as in overall body length and body weight (**Supp Figure 5, 6**) in *Phlpp1*^{-/-} mice. As males and females showed nearly identical results, male mice were used for all subsequent analyses.

Pth1r and Phlpp1 coordinate to regulate growth plate size: Chondrocyte hypertrophy in the epiphyseal growth plate determines longitudinal bone growth. Four-week-old Phlpp1^{-/-} mice have an increased cell number in the proliferative zone of the growth plate, suggesting delayed entry into hypertrophy and therefore shorter bone length⁽⁵⁾. 3D rendering of the proximal tibial growth plate⁽⁴⁸⁾ confirmed that Phlpp1^{-/-} mice had thicker growth plates than WT littermates (**Fig 4A,B**). The difference in growth plate thickness was due to an increase in the size of the proliferative zone, which was attenuated by the administration of PTH(7-34) to Phlpp1^{-/-} mice (**Fig 4C,D; Supp Fig 2B**). Treatment with PTH(7-34) did not have any effect on the proliferative zone of WT mice. The hypertrophic zone area was not statistically different in any of the groups (**Fig 4C,D**) and there was no difference in the expression of Col10a1 as detected by in situ hybridization (**Fig 4E; Supp Fig 2C**). To more directly assess the effects of Phlpp1 deletion and PTH(7-34) administration on cell proliferation, WT and Phlpp1^{-/-} mice were injected with vehicle or PTH(7-34) at the same doses as the 4-week-old mice from postnatal day 1 (P1) daily through P5. BrdU-positive cells were labeled and quantified in the proximal tibial growth plate. As previously reported, Phlpp1^{-/-} P5 mice had a greater number of BrdU+ cells compared to WT mice. PTH(7-34) administration reversed the effect of Phlpp1 deletion, reducing the number of BrdU+ cells in the proximal tibial growth plate to levels similar to WT mice (**Fig 4F,G; Supp Fig 2D**).

Daily administration of PTH(7-34) does not affect bone mass: Given the effects of Phlpp1 and Pth1r modulation on bone length, it was pertinent to examine bone mass. Consistent with previous studies⁽⁵⁾, Phlpp1^{-/-} mice had lower bone mass than WT littermates. Specifically, Phlpp1^{-/-} mice had lower bone volume / tissue volume (**Fig 5A,B**), fewer trabeculae (**Fig 5C**), and lower trabecular bone mineral density (**Fig 5D**) compared to WT mice. In addition, Phlpp1^{-/-} mice had lower cortical tissue mineral density (**Fig 5E**) and thinner cortical bone (**Fig 5F**). The administration of PTH(7-34) did not affect trabecular bone volume / tissue volume (**Fig 5B**), trabecular number (**Fig 5C**), trabecular bone mineral density (**Fig 5D**), or cortical tissue mineral density (**Fig 5E**) in either WT or Phlpp1^{-/-} mice. However, daily PTH(7-34) administration did reverse the effect of Phlpp1 deletion on cortical thickness (**Fig 5F**). Total tissue cross-sectional area was reduced in Phlpp1^{-/-} mice compared to WT mice, which was reversed by PTH(7-34) administration (**Fig 5G**), as would be expected given the Phlpp1^{-/-} mice have smaller appendicular bones.

However, the cortical area fraction was unaffected by Phlpp1 deletion or PTH(7-34) modulation (**Fig 5H**). Trabecular spacing and trabecular thickness were not affected (data not shown).

Phlpp1 inhibition elevates Pth1r signaling cascades: Having established that Phlpp1 regulates Pth1r expression through transcription (**Fig 2**) and controls chondrocyte proliferation in vivo (**Fig 3**), we tested the effects of Phlpp inactivation on known Pth1r signaling targets in vitro. PTH(1-34) was included in these experiment as a positive control. Endogenous PTH and PTH(1-34) bind to Pth1r and stimulate intracellular cAMP, in turn activating PKA and inducing phosphorylation of cAMP-response element-binding protein (CREB) at S133^(52,53). IMCs isolated from Phlpp1^{-/-} mice had greater baseline levels of Pth1r and pS133-CREB (**Fig 6A; Supp Fig 10-Q**). PTH(1-34) enhanced and PTH(7-34) reduced Pth1r and pS133-CREB. Phlpp1^{-/-} IMCs had similar basal cAMP concentrations as WT IMCs, but were more responsive to PTH(1-34) than WT IMCs. PTH(7-34) had no effect (**Fig 6B**). Confluency of Phlpp1^{-/-} chondrocytes increased more rapidly over 48 hours than that of WT monolayer cultures. PTH(1-34) further enhanced confluency of both WT and Phlpp1^{-/-} chondrocyte monolayer cultures, while PTH(7-34) attenuated Phlpp1^{-/-} growth to WT rates (**Fig 6C**). While Phlpp1^{-/-} chondrocytes had only numerically higher MTS activity compared to WT, treatment with PTH(1-34) further enhanced the effects of Phlpp1^{-/-}, while PTH(7-34) reversed the effects of Phlpp1^{-/-} on MTS activity (**Fig 6D**). Similar to previous results⁽⁵⁾, Phlpp1^{-/-} micromasses have more intense staining of Alcian Blue compared to WT littermates. PTH(1-34) further intensified Alcian Blue staining in Phlpp1^{-/-} chondrocytes. By contrast, treating Phlpp1^{-/-} cells with PTH(7-34) diminished the intensity of Alcian Blue stain to WT levels (**Fig 6E**). Together, these data show that Phlpp1^{-/-} chondrocytes express more Pth1r and are more responsive to PTH ligands.

DISCUSSION

Our results show that greater Pth1r expression and signaling in Phlpp1^{-/-} chondrocytes contributes to growth delays in long bones. Administration of PTH(7-34) for the first four weeks of life reverses the short limb phenotype in the appendicular skeleton of Phlpp1^{-/-} mice. Specifically, PTH(7-34) attenuates the thicker growth plates and large proliferative zones in the epiphyseal growth plate of Phlpp1^{-/-} mice. Confirming previous results⁽⁵⁾, Phlpp1^{-/-} mice have lower bone mineral density than WT mice, and PTH(7-34) has no effect on bone mass or microarchitecture. Regulation of Pth1r by Phlpp1 occurs at the

transcriptional level, as H3K27 acetylation is increased at the Pth1r promoter in the Phlpp1-depleted chondrocytes and blocking transcription prevents the increase in Pth1r mRNA and protein levels associated with Phlpp1 inhibition. The transcription factor YY1 may contribute to the increased transcription of Pth1r in Phlpp1 inactivated cells. Additionally, Phlpp1 inhibition increases phosphorylation of Pth1r effector CREB and increases cell proliferation, each of which can be enhanced with Pth1r agonism and attenuated with Pth1r antagonism. Taken together, these results demonstrate that Phlpp1 inhibition increases Pth1r expression and signaling to stunt endochondral ossification.

Pth1r is critical for proper growth plate development. In human populations, both inactivating⁽⁵⁴⁾ and activating^(55,56) mutations of Pth1r are associated with shortened limbs and abnormal bone mineralization. Mice with a chondrocyte-specific deletion for Pth1r have reduced chondrocyte proliferation, accelerated hypertrophic differentiation, and premature growth plate closure, ultimately resulting in short stature^(35,57). Our results show that Pth1r is overexpressed in Phlpp1^{-/-} chondrocytes. In previous studies, Pth1r overexpression in chondrocytes delayed mineralization and decelerated the conversion of chondrocytes from proliferative to hypertrophic states, resulting in shorter limbs⁽⁵⁸⁾. Although less dramatic, Phlpp1^{-/-} mice demonstrate a similar phenotype in which bone mineral density is reduced, and the proliferative zone of the growth plate is larger than WT mice.

The altered growth plate phenotype in Phlpp1^{-/-} mice results in shorter femurs and tibiae and this is reversed by administration of PTH(7-34). Our results indicate that Phlpp1-deficiency promotes proliferation through Pth1r signaling and that PTH(7-34) suppression of Pth1r slows proliferation and promotes bone lengthening. Injecting mice with 100 µg/kg body weight/day PTH(7-34) did not fully rescue bone length in every parameter measured. For example, both male and female Phlpp1^{-/-} mice injected with PTH(7-34) had longer tibiae than vehicle-injected Phlpp1^{-/-} mice, but the length was not completely recovered to the control groups. There could be several explanations for these findings. PTH(7-34) was only injected postnatally. PTH signaling is active in utero⁽⁵⁹⁾ and it is possible that a basal level of elevated Pth1r signaling had already occurred in Phlpp1^{-/-} mice such that it was only partially attenuated by postnatal administration of PTH(7-34). Other doses and timing schemes of PTH(7-34) administration could be further pursued to determine if there is a more complete rescue. Although our results have

focused primarily on chondrocyte dynamics in the growth plate, various hormones, cytokines, and paracrine growth factors work in concert with PTH signaling to determine endochondral ossification and should be further examined in *Phlpp1*^{-/-} mice⁽⁶⁰⁾.

There were no apparent effects of PTH(7-34) on bone mineral density in either WT or *Phlpp1*^{-/-} mice, with the notable exception of reversing thin cortical bone in *Phlpp1*^{-/-} mice. PTH signaling plays a significant role in bone mineralization⁽⁶¹⁾ and perturbations in signaling to either enhance or decrease PTH signaling can have profound effects on bone⁽⁶²⁾. PTH(1-34) increases bone mass when administered intermittently during development and in more mature models^(37,45,63-65). Teriparatide (PTH(1-34)) and abaloparatide (PTHrP(1-34)) are widely utilized osteoanabolic treatments that rely on Pth1r signaling for bone accrual⁽⁶⁵⁾. Although *Phlpp1*^{-/-} mice have increased Pth1r expression and signaling, they have lower bone mineral density⁽⁵⁾. This is perhaps unsurprising, as constitutively overexpressing Pth1r delays bone mineralization during development⁽⁵⁸⁾. *Phlpp1* deletion specifically in osteoclasts increases bone mineralization⁽⁷⁾ while germline deletion of *Phlpp1* decreases bone mass⁽⁵⁾, demonstrating that *Phlpp1* effects are distinct in different musculoskeletal cell types. The fact that PTH(7-34) was unable to rescue low bone mineralization in *Phlpp1*^{-/-} mice suggests that *Phlpp1* differentially regulates Pth1r signaling in bone cells (osteoblasts, osteoclasts, and osteocytes) compared to chondrocytes. Future studies should probe the relationship between *Phlpp1* and Pth1r in individual musculoskeletal types to elucidate this relationship beyond chondrocytes.

In vitro, our results recapitulate the in vivo relationship between *Phlpp1* inhibition and enhanced Pth1r signaling in chondrocytes. While agonism of Pth1r with PTH(1-34) enhances *Phlpp1*^{-/-} responses, attenuation of PTH signaling with PTH(7-34) can reverse the effects of *Phlpp1* inhibition. *Phlpp1*-depleted cells show increased phosphorylation of CREB and treating *Phlpp1*^{-/-} cells with PTH(1-34) further enhanced pCREB, while PTH(7-34) reversed this indicator of Pth1r signaling. PTH(1-34) is more responsive in *Phlpp1*^{-/-} cells that have higher concentrations of the Pth1r receptor available for PTH(1-34) binding, as cell metabolic activity and cAMP concentrations were highest in *Phlpp1*^{-/-} cells treated with PTH(1-34). Similarly, cell proliferation, as well as cartilage matrix deposition, were increased in *Phlpp1*^{-/-} cells compared to wild type littermates, with PTH(7-34) attenuating elevated proliferation and

chondrogenesis associated with Phlpp1 inhibition. In vivo, BrdU incorporation in the growth plate of P5 mice confirmed the in vitro findings. Further enhancement of cell proliferation as a result of Phlpp1 inhibition has potential implications beyond chondrocytes. Phlpp1 deletion promotes cell proliferation in a model of intervertebral disc degeneration⁽¹⁹⁾ and regulation of cell apoptosis by Phlpp1 is implicated in ischemic brain and spinal cord injury^(20,22), colitis⁽²¹⁾, cardiac dysfunction⁽⁶⁶⁾, pancreatic beta-cell survival⁽⁶⁷⁾, and tumor activity⁽⁸⁾. As such, enhancement of signals associated with Phlpp1 inhibition could have a profound impact on various physiological and disease processes.

Phlpp1 modulates Pth1r signaling through chromatin remodeling of the ubiquitous Pth1r promoter resulting in increased transcription, as well as greater activation of Pth1r signaling⁽⁶⁸⁾. The mechanisms responsible for increased transcription of Pth1r via Phlpp inhibition remain to be determined, but appear to involve a modest increase in existing transcriptional events. A variety of transcription factors are known to bind to the Pth1r promoter and may be active in Phlpp-inhibited chondrocytes⁽⁶⁸⁻⁷²⁾. Numerous potential transcription factor binding sites within the 2,000 base pair H3K27ac enrichment peak in the Pth1r promoter were identified and are summarized in **Supplementary Table 1**⁽⁷³⁾. One promising transcription factor, YY1, showed increased binding at the Pth1r promoter in response to Phlpp inhibition. Ying Yang 1 (YY1) is a dual function transcription factor that regulates both transcriptional activation and repression and has been implicated in a host of cellular processes, including differentiation, DNA repair, cell division, and cell survival⁽⁷⁴⁾. Additionally, YY1 facilitates the interaction of active enhancers and promoter-proximal elements, serving as a general feature of mammalian gene control⁽⁷⁵⁾. In chondrocytes, Phlpp inhibitors increase binding of YY1 to the Pth1r promoter, suggesting that YY1 acts as a transcriptional activator in Phlpp-inactivated chondrocytes. However, because a large number of factors could associate with this region of the Pth1r promoter, it is likely that Phlpp1 repression of H3K27ac influences many additional transcription factors and co-regulators.

The relationship between Phlpp1 and Pth1r has applications beyond endochondral ossification during development. Pth1r^{-/-} mice have spontaneous cartilage degeneration and develop more severe post-traumatic osteoarthritis in the knee than wild type littermates, with greater chondrocyte hypertrophy⁽⁷⁶⁾. By contrast, Phlpp1^{-/-} mice and WT mice given an intra-articular injection of NSC117079 into the knee joint

are protected from the development of post-traumatic osteoarthritis, with improved allodynia and functional impairments. As such, Phlpp inhibitors have potential as novel disease modifying osteoarthritis drugs (DMOADs). Human recombinant PTH(1-34) (teriparatide) is currently in clinical trials as a DMOAD, following preclinical results of reduced cartilage degeneration and induction of matrix regeneration⁽⁷⁷⁾. As our findings indicate a synergistic relationship between Phlpp1 inhibition and enhanced Pth1r signaling, it is possible that combinations of Phlpp1 inhibitors and PTH(1-34) could be effective promoters of cartilage regeneration.

In summary, our results demonstrate a novel molecular relationship in which Phlpp1 inhibition enhances Pth1r expression and signaling in chondrocytes. Repression of Pth1r signaling reversed both the stunted growth of the appendicular skeleton characteristic of Phlpp1^{-/-} mice, and the increased cell proliferation of Phlpp1^{-/-} chondrocytes, both in vitro and in the proliferative zone of the epiphyseal growth plate. Our findings have implications on multiple processes, including endochondral ossification during development and musculoskeletal diseases such as osteoarthritis. Phlpp1 and Pth1r signaling is not restricted to the bone environment, as PTH controls mineral homeostasis⁽⁷⁸⁾ among other processes, and Phlpp1 is expressed throughout the mammalian body⁽⁶⁾. As such, the Phlpp1-Pth1r relationship could have a significant impact in the field of musculoskeletal health and beyond.

ACKNOWLEDGEMENTS

The authors thank Xiaodong Li and Carys Turner for technical assistance. The Mayo Clinic X-Ray Imaging Resources Core Facilities were essential for collection of the microCT data. We are grateful to the Mayo Clinic Medical Genome Facility Sequencing Core for performing and helping with the analysis of ChIP-Sequencing. This work was supported by research and training grants from the National Institutes of Health (AR068103, AR056950, DK07352, AR065397, AR072634).

Authors' roles: Study design: EWB, JJW. Study conducted: all authors. Data collection and analysis: all authors. Data interpretation: SRW, ELT, EWB, JJW. Writing – original draft: SRW. Revising and approving final version of manuscript: all authors. Supervision: EWB, JJW. Project leadership: SRW, ELT,

EWB, JJW. Funding acquisition: EWB, JJW. SRW and JJW take responsibility for the integrity of the data analysis.

FIGURE LEGENDS

Figure 1. *Phlpp inhibition increases Pth1r expression in chondrocytes.* (A-D) Pth1r mRNA expression and protein levels were measured in ATDC5 cells (A,B) and primary chondrocytes (C,D) cultured in the presence of Phlpp inhibitors for 24 hours. Pth1r (E) mRNA expression and (F) protein levels were measured in Phlpp1^{-/-} primary chondrocytes compared to WT. (G,H) The proximal tibial growth plates of 4-week-old male WT and Phlpp1^{-/-} mice are shown. (G) In situ hybridization was performed with a riboprobe targeted to Pth1r. Scale bar = 50 µm. (H) Immunohistochemistry was performed with an antibody targeted to Pth1r. Scale bar = 20 µm. Vertical black lines indicate either P = proliferative zone or H = hypertrophic zone. Statistically significant differences were determined with one-way ANOVA with Tukey's post-hoc test (A, C) or Student's t test (E).

Figure 2. *Phlpp1 regulates Pth1r expression through transcription.* (A) Primary WT chondrocytes were treated with 25 µM Phlpp inhibitors NSC117079 or NSC45586 for 24 hours. These lysates and those from primary chondrocytes from WT and Phlpp1^{-/-} mice were subjected to Western blotting for phosphorylation of S10 or S28 (pS10 or pS28), as well as K9, K14, and K27 acetylation (ac) of histone 3 (H3). Each blot was also probed with an antibody that recognizes all (total) H3 to control for loading. (B) H3K27ac ChIP-sequencing was performed on WT primary chondrocytes treated for 24 hours with vehicle (DMSO) or 25µM NSC117079. Two sites were identified for analysis by ChIP-qPCR, including in the Pth1r promoter and -1.47kb upstream. (C-D) ChIP-qPCR following pulldown with an H3K27ac antibody was performed on WT primary chondrocytes treated with vehicle or 25µM NSC117079 and NSC45586 after 24 hours (C) or on WT or Phlpp1^{-/-} primary chondrocytes (D). (E) Pth1r mRNA expression and protein levels were measured in primary chondrocytes after incubation with 5 µM transcriptional inhibitor actinomycin D for six hours and subsequent replacement of media containing Phlpp inhibitors for 24 hours. (F) WT primary chondrocytes were treated with vehicle or 25 µM NSC117079 and NSC45586 for 30 minutes. YY1-Pth1r promoter complexes were identified by ChIP-qPCR. Statistically significant differences were determined with two-way ANOVA with Tukey's post-hoc test.

Figure 3. Daily administration of PTH(7-34) reverses short limb length in *Phlpp1*^{-/-} mice. (A,C) Tibia and (B,D) femur lengths were evaluated in 4-week-old male WT or *Phlpp1*^{-/-} mice given daily injections of PTH(7-34) (100 mg/kg body weight/day) or vehicle (0.1% BSA in PBS). The same treatments were administered to female WT and *Phlpp1*^{-/-} mice and tibia (E,G) and femur (F,H) lengths were evaluated at 4 weeks of age. Scale bar = 5 mm. Statistically significant differences were determined with two-way ANOVA with Tukey's post-hoc test.

Figure 4. *Phlpp1* and *Pth1r* coordinate to regulate growth plate development. (A,B) 3D renderings of proximal tibial growth plates were generated from microCT scans of 4-week-old WT or *Phlpp1*^{-/-} male mice given daily injections of PTH(7-34) (100 mg/kg body weight/day) or vehicle. Scale bar on 2D capture = 1mm. Color scale bar is in mm. One representative 3D rendering is shown for each group. (C) Areas of proliferative and hypertrophic zones of the proximal tibia growth plate were quantified. (D) Safranin O / Fast Green staining was performed on the proximal tibial growth plate. Scale bar = 50 µm. Vertical black lines indicate either P = proliferative zone or H = hypertrophic zone. (E) In situ hybridization was performed for *Col10a1*. Scale bar = 50 µm. (F,G) WT or *Phlpp1*^{-/-} mice were injected with either vehicle or PTH(7-34) daily as described above from postnatal day 1 (P1) through P5. On P5, BrdU was administered 2 hours prior to euthanasia. (F) Percentage of cells that were BrdU-positive in the proximal tibial growth plate was quantified, as represented in (G). Scale bar in (G) = 50 µm. Statistically significant differences were determined with two-way ANOVA with Tukey's post-hoc test.

Figure 5. Daily administration of PTH(7-34) does not affect bone mass. (A) 2D reconstructions from microCT of the distal femur of 4-week-old WT or *Phlpp1*^{-/-} male mice given daily injections of PTH(7-34) (100 mg/kg body weight/day) or vehicle. Scale bar = 1 mm. Trabecular parameters analyzed via microCT were (B) bone volume / tissue volume, (C) trabecular number, and (D) bone mineral density. Cortical parameters included (E) tissue mineral density, (F) cortical thickness, (G) total tissue cross sectional area, and (H) cortical area fraction. Statistically significant differences were determined with two-way ANOVA with Tukey's post-hoc test or Student's t test as indicated.

Figure 6. *Phlpp1* inhibition elevates *Pth1r* signaling cascades. (A) Primary chondrocytes from *Phlpp1*^{-/-} and WT littermates were subjected to Western blotting for *Phlpp1*, *Pth1r*, pCREB (S133), total CREB, and

actin following 24 hours in culture, with the last 30 minutes in the presence of vehicle (0.1% BSA in PBS), PTH(1-34), or PTH(7-34) (100 nM). (B) cAMP levels in primary chondrocytes from Phlpp1^{-/-} and WT littermates were determined by ELISA following addition of PTH(1-34) or PTH(7-34) (100 nM) for 15 minutes. (C) Confluency of WT and Phlpp1^{-/-} primary chondrocytes was tracked every hour for 48 hours in the absence or presence of PTH(1-34) or PTH(7-34) (10 nM). Data in the three graphs are from the same experiment, but plotted separately for clarity. (D) Primary chondrocytes from Phlpp1^{-/-} and WT littermates were subjected to MTS assay following addition of PTH(1-34) or PTH(7-34) (10 nM) for 24 hours. (E) WT or Phlpp1^{-/-} micromasses were cultured for three days and PTH(1-34) or PTH(7-34) (10 nM) was applied for the next nine days at which time micromasses were stained with Alcian Blue. Statistically significant differences were determined with two-way ANOVA with Tukey's post-hoc test.

Supplementary Figure 1. Quantified Western Blots. (A-C) Pth1r protein levels were measured in (A) ATDC5s and (B) WT primary chondrocytes after incubation with vehicle or Phlpp inhibitors (25μM NSC117079 or NSC45586) for 24 hours, as well as in (C) WT or Phlpp1^{-/-} primary chondrocytes. Actin was the loading control. (D-M) (D-H) WT primary chondrocytes were treated with Phlpp inhibitors for 24 hours. (I-M) Primary chondrocytes from WT and Phlpp1^{-/-} were grown in culture for 24 hours. The protein levels measured were (D,I) H3pS10, (E,J) H3pS28, (F,K) H3K27ac, (G,L) H3K9ac, and (H,M) H3K9K14ac. Total histone 3 (H3) was the loading control. (N) Pth1r protein levels were measured in primary chondrocytes after incubation with 5 μM actinomycin D for six hours and subsequent replacement of media containing Phlpp inhibitors for 24 hours. (O-Q) Primary chondrocytes from Phlpp1^{-/-} and WT mice were subjected to Western blotting for Phlpp1, Pth1r, and pCREB (S133) with actin and total CREB as the loading controls following 24 hours in culture, with the last 30 minutes in the presence of vehicle, PTH(1-34) (100 nM), or PTH(7-34) (100 nM).

Supplementary Figure 2. Low magnification images of the growth plate. 5x magnification images of (A) the growth plates of 4-week-old WT or Phlpp1^{-/-} male mice after performing in situ hybridization (ISH) or immunohistochemistry (IHC) to detect Pth1r. Scale bar = 100 μm. (B,C) Four-week-old WT or Phlpp1^{-/-} mice were given daily injections of PTH(7-34) (100 mg/kg body weight/day) or vehicle. The proximal tibial growth plate was (B) stained with Safranin O / Fast Green and (C) probed for Col10a1 using ISH. (D) WT

or *Phlpp1*^{-/-} mice were injected with either vehicle or PTH(7-34) daily as described above from postnatal day 1 (P1) through P5. On P5, BrdU labeling reagent was administered 2 hours prior to euthanasia and BrdU-positive cells were identified.

Supplementary Figure 3. *Small-molecule Phlpp inhibitors increase H3K27ac in Pth1r.* H3K27ac ChIP-sequencing was performed on WT primary chondrocytes incubated for 24 hours with vehicle (DMSO) or 25µM NSC117079. Data were aligned with publicly available ChIP-Seq datasets from E10.5 and E15.5 limbs for Chr 9: 100,722,085 – 110,747,145. Limb data was retrieved from Ensembl for *Mus musculus*, version 100.38 (GRCm38.p6).

Supplementary Figure 4. *Plasma PTH and urine cAMP are the same in WT and Phlpp1*^{-/-} *mice.* (A) Plasma intact PTH(1-84) and (B) urine cAMP corrected for urine creatinine were measured in samples taken from four-week-old WT and *Phlpp1*^{-/-} mice. Statistically significant differences were determined via one-way ANOVA with Tukey's post-hoc test.

Supplementary Figure 5. *Daily administration of PTH(7-34) has an intermediate effect on rescuing limb length in Phlpp1 heterozygotes compared to WT or Phlpp1*^{-/-} *mice.* (A,E) Femur length, (B,F) tibia length, (C,G) body length, and (D,H) body weight were evaluated in 4-week-old male and female WT (*Phlpp1*^{+/+}), HET (*Phlpp1*^{+/-}), or KO (*Phlpp1*^{-/-}) mice given daily injections of PTH(7-34) (100 mg/kg body weight/day) or vehicle (0.1% BSA in PBS). Statistically significant differences were determined with two-way ANOVA with Tukey's post-hoc test.

Supplementary Figure 6. *Daily administration of PTH(7-34) reverses short body length and low body weight in Phlpp1*^{-/-} *mice.* (A,C) Tail-to-snout body length and (B,D) body weight were evaluated in 4-week-old male and female WT or *Phlpp1*^{-/-} mice given daily injections of PTH(7-34) (100 mg/kg body weight/day) or vehicle (0.1% BSA in PBS). Statistically significant differences were determined with two-way ANOVA with Tukey's post-hoc test.

REFERENCES

1. Kronenberg HM. Developmental regulation of the growth plate. *Nature*. May 15 2003;423(6937):332-6. Epub 2003/05/16.

2. Ono N, Ono W, Nagasawa T, Kronenberg HM. A subset of chondrogenic cells provides early mesenchymal progenitors in growing bones. *Nat Cell Biol.* Dec 2014;16(12):1157-67. Epub 2014/11/25.
3. Yang L, Tsang KY, Tang HC, Chan D, Cheah KSE. Hypertrophic chondrocytes can become osteoblasts and osteocytes in endochondral bone formation. *Proceedings of the National Academy of Sciences.* 2014;111(33):12097.
4. Zhou X, von der Mark K, Henry S, Norton W, Adams H, de Crombrughe B. Chondrocytes transdifferentiate into osteoblasts in endochondral bone during development, postnatal growth and fracture healing in mice. *PLoS Genet.* Dec 2014;10(12):e1004820. Epub 2014/12/05.
5. Bradley EW, Carpio LR, Newton AC, Westendorf JJ. Deletion of the PH-domain and Leucine-rich Repeat Protein Phosphatase 1 (Phlpp1) Increases Fibroblast Growth Factor (Fgf) 18 Expression and Promotes Chondrocyte Proliferation. *J Biol Chem.* Jun 26 2015;290(26):16272-80. Epub 2015/05/09.
6. Grzechnik AT, Newton AC. PHLPPing through history: a decade in the life of PHLPP phosphatases. *Biochem Soc Trans.* Dec 15 2016;44(6):1675-82. Epub 2016/12/04.
7. Mattson AM, Begun DL, Molstad DHH, Meyer MA, Oursler MJ, Westendorf JJ, et al. Deficiency in the phosphatase PHLPP1 suppresses osteoclast-mediated bone resorption and enhances bone formation in mice. *J Biol Chem.* Aug 2 2019;294(31):11772-84. Epub 2019/06/14.
8. Gao T, Furnari F, Newton AC. PHLPP: a phosphatase that directly dephosphorylates Akt, promotes apoptosis, and suppresses tumor growth. *Mol Cell.* Apr 1 2005;18(1):13-24. Epub 2005/04/06.
9. Liu J, Weiss HL, Rychahou P, Jackson LN, Evers BM, Gao T. Loss of PHLPP expression in colon cancer: role in proliferation and tumorigenesis. *Oncogene.* Feb 19 2009;28(7):994-1004. Epub 2008/12/17.
10. Gao T, Brognard J, Newton AC. The phosphatase PHLPP controls the cellular levels of protein kinase C. *J Biol Chem.* Mar 7 2008;283(10):6300-11. Epub 2007/12/29.
11. Liu J, Stevens PD, Li X, Schmidt MD, Gao T. PHLPP-mediated dephosphorylation of S6K1 inhibits protein translation and cell growth. *Mol Cell Biol.* Dec 2011;31(24):4917-27. Epub 2011/10/12.
12. Reyes G, Niederst M, Cohen-Katsenelson K, Stender JD, Kunkel MT, Chen M, et al. Pleckstrin homology domain leucine-rich repeat protein phosphatases set the amplitude of receptor tyrosine kinase output. *Proc Natl Acad Sci U S A.* Sep 23 2014;111(38):E3957-65. Epub 2014/09/10.
13. Nowak DG, Katsenelson KC, Watrud KE, Chen M, Mathew G, D'Andrea VD, et al. The PHLPP2 phosphatase is a druggable driver of prostate cancer progression. *J Cell Biol.* Jun 3 2019;218(6):1943-57. Epub 2019/05/17.
14. Xiong X, Wen YA, Mitov MI, M CO, Miyamoto S, Gao T. PHLPP regulates hexokinase 2-dependent glucose metabolism in colon cancer cells. *Cell Death Discov.* 2017;3:16103. Epub 2017/02/10.
15. Hribal ML, Mancuso E, Spiga R, Mannino GC, Fiorentino TV, Andreozzi F, et al. PHLPP phosphatases as a therapeutic target in insulin resistance-related diseases. *Expert Opin Ther Targets.* Jun 2016;20(6):663-75. Epub 2015/12/15.
16. Patterson SJ, Han JM, Garcia R, Assi K, Gao T, O'Neill A, et al. Cutting edge: PHLPP regulates the development, function, and molecular signaling pathways of regulatory T cells. *J Immunol.* May 15 2011;186(10):5533-7. Epub 2011/04/19.
17. Bradley EW, Carpio LR, McGee-Lawrence ME, Castillejo Becerra C, Amanatullah DF, Ta LE, et al. Phlpp1 facilitates post-traumatic osteoarthritis and is induced by inflammation and promoter demethylation in human osteoarthritis. *Osteoarthritis Cartilage.* 2016;24(6):1021-8. Epub 2015/12/31.
18. Hwang SM, Feigenson M, Begun DL, Shull LC, Culley KL, Otero M, et al. Phlpp inhibitors block pain and cartilage degradation associated with osteoarthritis. *J Orthop Res.* May 2018;36(5):1487-97. Epub 2017/10/27.
19. Zhang C, Smith MP, Zhou GK, Lai A, Hoy RC, Mroz V, et al. Phlpp1 is associated with human intervertebral disc degeneration and its deficiency promotes healing after needle puncture injury in mice. *Cell Death Dis.* Oct 3 2019;10(10):754. Epub 2019/10/05.

20. Shao Z, Lv G, Wen P, Cao Y, Yu D, Lu Y, et al. Silencing of PHLPP1 promotes neuronal apoptosis and inhibits functional recovery after spinal cord injury in mice. *Life Sci.* Sep 15 2018;209:291-9. Epub 2018/08/17.
21. Wen YA, Li X, Goretsky T, Weiss HL, Barrett TA, Gao T. Loss of PHLPP protects against colitis by inhibiting intestinal epithelial cell apoptosis. *Biochim Biophys Acta.* Oct 2015;1852(10 Pt A):2013-23. Epub 2015/07/19.
22. Chen B, Van Winkle JA, Lyden PD, Brown JH, Purcell NH. PHLPP1 gene deletion protects the brain from ischemic injury. *J Cereb Blood Flow Metab.* Feb 2013;33(2):196-204. Epub 2012/10/18.
23. Jackson TC, Verrier JD, Drabek T, Janesko-Feldman K, Gillespie DG, Uray T, et al. Pharmacological inhibition of pleckstrin homology domain leucine-rich repeat protein phosphatase is neuroprotective: differential effects on astrocytes. *J Pharmacol Exp Ther.* Nov 2013;347(2):516-28. Epub 2013/09/12.
24. Jackson TC, Dixon CE, Janesko-Feldman K, Vagni V, Kotermanski SE, Jackson EK, et al. Acute Physiology and Neurologic Outcomes after Brain Injury in SCOP/PHLPP1 KO Mice. *Sci Rep.* May 8 2018;8(1):7158. Epub 2018/05/10.
25. Cheloha RW, Gellman SH, Vilardaga JP, Gardella TJ. PTH receptor-1 signalling-mechanistic insights and therapeutic prospects. *Nat Rev Endocrinol.* Dec 2015;11(12):712-24. Epub 2015/08/26.
26. Yamamoto T, Kambe F, Cao X, Lu X, Ishiguro N, Seo H. Parathyroid hormone activates phosphoinositide 3-kinase-Akt-Bad cascade in osteoblast-like cells. *Bone.* Feb 2007;40(2):354-9. Epub 2006/10/19.
27. Miao D, Tong XK, Chan GK, Panda D, McPherson PS, Goltzman D. Parathyroid hormone-related peptide stimulates osteogenic cell proliferation through protein kinase C activation of the Ras/mitogen-activated protein kinase signaling pathway. *J Biol Chem.* Aug 24 2001;276(34):32204-13. Epub 2001/06/13.
28. Gesty-Palmer D, Chen M, Reiter E, Ahn S, Nelson CD, Wang S, et al. Distinct beta-arrestin- and G protein-dependent pathways for parathyroid hormone receptor-stimulated ERK1/2 activation. *J Biol Chem.* Apr 21 2006;281(16):10856-64. Epub 2006/02/24.
29. Fan Y, Hanai JI, Le PT, Bi R, Maridas D, DeMambro V, et al. Parathyroid Hormone Directs Bone Marrow Mesenchymal Cell Fate. *Cell Metab.* Mar 7 2017;25(3):661-72. Epub 2017/02/07.
30. Rhee Y, Allen MR, Condon K, Lezcano V, Ronda AC, Galli C, et al. PTH receptor signaling in osteocytes governs periosteal bone formation and intracortical remodeling. *J Bone Miner Res.* May 2011;26(5):1035-46. Epub 2010/12/09.
31. Qiu T, Xian L, Crane J, Wen C, Hilton M, Lu W, et al. PTH receptor signaling in osteoblasts regulates endochondral vascularization in maintenance of postnatal growth plate. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research.* 2015;30(2):309-17.
32. Calvi LM, Sims NA, Hunzelman JL, Knight MC, Giovannetti A, Saxton JM, et al. Activated parathyroid hormone/parathyroid hormone-related protein receptor in osteoblastic cells differentially affects cortical and trabecular bone. *J Clin Invest.* Feb 2001;107(3):277-86. Epub 2001/02/13.
33. Becher C, Szuwart T, Ronstedt P, Ostermeier S, Skwara A, Fuchs-Winkelmann S, et al. Decrease in the expression of the type 1 PTH/PTHrP receptor (PTH1R) on chondrocytes in animals with osteoarthritis. *J Orthop Surg Res.* 2010;5:28-.
34. Vortkamp A, Lee K, Lanske B, Segre GV, Kronenberg HM, Tabin CJ. Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein. *Science.* Aug 2 1996;273(5275):613-22. Epub 1996/08/02.
35. Hirai T, Chagin AS, Kobayashi T, Mackem S, Kronenberg HM. Parathyroid hormone/parathyroid hormone-related protein receptor signaling is required for maintenance of the growth plate in postnatal life. *Proc Natl Acad Sci U S A.* Jan 4 2011;108(1):191-6. Epub 2010/12/22.
36. Sneddon WB, Magyar CE, Willick GE, Syme CA, Galbiati F, Bisello A, et al. Ligand-selective dissociation of activation and internalization of the parathyroid hormone (PTH) receptor: conditional efficacy of PTH peptide fragments. *Endocrinology.* Jun 2004;145(6):2815-23. Epub 2004/03/16.

37. Silva BC, Costa AG, Cusano NE, Kousteni S, Bilezikian JP. Catabolic and anabolic actions of parathyroid hormone on the skeleton. *J Endocrinol Invest*. Nov 2011;34(10):801-10. Epub 2011/09/29.
38. Wein MN, Kronenberg HM. Regulation of Bone Remodeling by Parathyroid Hormone. *Cold Spring Harb Perspect Med*. Aug 1 2018;8(8). Epub 2018/01/24.
39. Gardinier JD, Mohamed F, Kohn DH. PTH Signaling During Exercise Contributes to Bone Adaptation. *J Bone Miner Res*. Jun 2015;30(6):1053-63. Epub 2014/12/23.
40. Gosset M, Berenbaum F, Thirion S, Jacques C. Primary culture and phenotyping of murine chondrocytes. *Nat Protoc*. 2008;3(8):1253-60. Epub 2008/08/21.
41. Carpio LR, Bradley EW, McGee-Lawrence ME, Weivoda MM, Poston DD, Dudakovic A, et al. Histone deacetylase 3 supports endochondral bone formation by controlling cytokine signaling and matrix remodeling. *Sci Signal*. Aug 9 2016;9(440):ra79. Epub 2016/08/11.
42. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*. Dec 2001;25(4):402-8. Epub 2002/02/16.
43. MatLab. 2019b ed. Natick, MA: Mathworks; 2019.
44. Masubuchi S, Gao T, Neill A, Eckel-Mahan K, Newton AC, Sassone-Corsi P. Protein phosphatase PHLPP1 controls the light-induced resetting of the circadian clock. *Proceedings of the National Academy of Sciences*. 2010;107(4):1642.
45. Xie Y, Su N, Jin M, Qi H, Yang J, Li C, et al. Intermittent PTH (1-34) injection rescues the retarded skeletal development and postnatal lethality of mice mimicking human achondroplasia and thanatophoric dysplasia. *Hum Mol Genet*. Sep 15 2012;21(18):3941-55. Epub 2012/05/29.
46. Bouxsein ML, Boyd SK, Christiansen BA, Guldberg RE, Jepsen KJ, Muller R. Guidelines for assessment of bone microstructure in rodents using micro-computed tomography. *J Bone Miner Res*. Jul 2010;25(7):1468-86. Epub 2010/06/10.
47. Fedorov A, Beichel R, Kalpathy-Cramer J, Finet J, Fillion-Robin JC, Pujol S, et al. 3D Slicer as an image computing platform for the Quantitative Imaging Network. *Magn Reson Imaging*. Nov 2012;30(9):1323-41. Epub 2012/07/10.
48. Shazeeb MS, Cox MK, Gupta A, Tang W, Singh K, Pryce CT, et al. Skeletal Characterization of the Fgfr3 Mouse Model of Achondroplasia Using Micro-CT and MRI Volumetric Imaging. *Scientific Reports*. 2018/01/11 2018;8(1):469.
49. Mead TJ, Lefebvre V. Proliferation assays (BrdU and EdU) on skeletal tissue sections. *Methods Mol Biol*. 2014;1130:233-43. Epub 2014/02/01.
50. Bradley EW, Carpio LR, Westendorf JJ. Histone deacetylase 3 suppression increases PH domain and leucine-rich repeat phosphatase (Phlpp)1 expression in chondrocytes to suppress Akt signaling and matrix secretion. *J Biol Chem*. Apr 5 2013;288(14):9572-82. Epub 2013/02/15.
51. Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, et al. Ensembl 2020. *Nucleic Acids Research*. 2019;48(D1):D682-D8.
52. Tyson DR, Swarthout JT, Partridge NC. Increased osteoblastic c-fos expression by parathyroid hormone requires protein kinase A phosphorylation of the cyclic adenosine 3',5'-monophosphate response element-binding protein at serine 133. *Endocrinology*. Mar 1999;140(3):1255-61. Epub 1999/03/06.
53. Qin L, Partridge NC. Stimulation of amphiregulin expression in osteoblastic cells by parathyroid hormone requires the protein kinase A and cAMP response element-binding protein signaling pathway. *J Cell Biochem*. Oct 15 2005;96(3):632-40. Epub 2005/08/10.
54. Jobert AS, Zhang P, Couvineau A, Bonaventure J, Roume J, Le Merrer M, et al. Absence of functional receptors for parathyroid hormone and parathyroid hormone-related peptide in Blomstrand chondrodysplasia. *J Clin Invest*. Jul 1 1998;102(1):34-40. Epub 1998/07/03.
55. Schipani E, Langman CB, Parfitt AM, Jensen GS, Kikuchi S, Kooh SW, et al. Constitutively activated receptors for parathyroid hormone and parathyroid hormone-related peptide in Jansen's metaphyseal chondrodysplasia. *N Engl J Med*. Sep 5 1996;335(10):708-14. Epub 1996/09/05.
56. Schipani E, Kruse K, Jüppner H. A constitutively active mutant PTH-PTHrP receptor in Jansen-type metaphyseal chondrodysplasia. *Science*. Apr 7 1995;268(5207):98-100. Epub 1995/04/07.
57. Chung UI, Lanske B, Lee K, Li E, Kronenberg H. The parathyroid hormone/parathyroid hormone-related peptide receptor coordinates endochondral bone development by directly controlling chondrocyte differentiation. *Proc Natl Acad Sci U S A*. Oct 27 1998;95(22):13030-5. Epub 1998/10/28.

58. Schipani E, Lanske B, Hunzelman J, Luz A, Kovacs CS, Lee K, et al. Targeted expression of constitutively active receptors for parathyroid hormone and parathyroid hormone-related peptide delays endochondral bone formation and rescues mice that lack parathyroid hormone-related peptide. *Proc Natl Acad Sci U S A*. Dec 9 1997;94(25):13689-94. Epub 1998/02/12.
59. Tobias JH, Cooper C. Perspective: PTH/PTHrP Activity and the Programming of Skeletal Development In Utero. *Journal of Bone and Mineral Research*. 2004;19(2):177-82.
60. Jee YH, Baron J. The Biology of Stature. *J Pediatr*. Jun 2016;173:32-8. Epub 2016/03/31.
61. Ryan BA, Kovacs CS. Calcitropic and phosphotropic hormones in fetal and neonatal bone development. *Semin Fetal Neonatal Med*. Feb 2020;25(1):101062. Epub 2019/12/02.
62. Qin L, Raggatt LJ, Partridge NC. Parathyroid hormone: a double-edged sword for bone metabolism. *Trends Endocrinol Metab*. Mar 2004;15(2):60-5. Epub 2004/03/24.
63. Alexander JM, Bab I, Fish S, Müller R, Uchiyama T, Gronowicz G, et al. Human parathyroid hormone 1-34 reverses bone loss in ovariectomized mice. *J Bone Miner Res*. Sep 2001;16(9):1665-73. Epub 2001/09/08.
64. Sheehan S, Muthusamy A, Paul E, Sikes RA, Gomes RR, Jr. Short-term intermittent PTH 1-34 administration enhances bone formation in SCID/Beige mice. *Endocr J*. 2010;57(5):373-82. Epub 2010/02/09.
65. Le Henaff C, Ricarte F, Finnie B, He Z, Johnson J, Warshaw J, et al. Abaloparatide at the Same Dose Has the Same Effects on Bone as PTH (1-34) in Mice. *Journal of Bone and Mineral Research*. 2020;35(4):714-24.
66. Zhang M, Wang X, Liu M, Liu D, Pan J, Tian J, et al. Inhibition of PHLPP1 ameliorates cardiac dysfunction via activation of the PI3K/Akt/mTOR signalling pathway in diabetic cardiomyopathy. *J Cell Mol Med*. Apr 2020;24(8):4612-23. Epub 2020/03/10.
67. Hribal ML, Mancuso E, Arcidiacono GP, Greco A, Musca D, Procopio T, et al. The Phosphatase PHLPP2 Plays a Key Role in the Regulation of Pancreatic Beta-Cell Survival. *Int J Endocrinol*. 2020;2020:1027386. Epub 2020/05/16.
68. Kawane T, Mimura J, Fujii-Kuriyama Y, Horiuchi N. Identification of the promoter region of the parathyroid hormone receptor gene responsible for transcriptional suppression by insulin-like growth factor-I. *Arch Biochem Biophys*. Jul 1 2005;439(1):61-9. Epub 2005/06/14.
69. Kawane T, Mimura J, Yanagawa T, Fujii-Kuriyama Y, Horiuchi N. Parathyroid hormone (PTH) down-regulates PTH/PTH-related protein receptor gene expression in UMR-106 osteoblast-like cells via a 3',5'-cyclic adenosine monophosphate-dependent, protein kinase A-independent pathway. *J Endocrinol*. Aug 2003;178(2):247-56. Epub 2003/08/09.
70. Minagawa M, Kwan MY, Bettoun JD, Mansour FW, Dassa J, Hendy GN, et al. Dissection of differentially regulated (G+C)-rich promoters of the human parathyroid hormone (PTH)/PTH-related peptide receptor gene. *Endocrinology*. Jul 2000;141(7):2410-21. Epub 2000/06/30.
71. Tohmonda T, Yoda M, Mizuochi H, Morioka H, Matsumoto M, Urano F, et al. The IRE1 α -XBP1 pathway positively regulates parathyroid hormone (PTH)/PTH-related peptide receptor expression and is involved in pth-induced osteoclastogenesis. *J Biol Chem*. Jan 18 2013;288(3):1691-5. Epub 2012/12/14.
72. Su S, Cao J, Meng X, Liu R, Ark AV, Woodford E, et al. Enzalutamide-induced PTH1R-mediated TGFBR2 decrease in osteoblasts contributes to resistance in prostate cancer bone metastases. *bioRxiv*. 2019:829044.
73. Kang R, Zhang Y, Huang Q, Meng J, Ding R, Chang Y, et al. EnhancerDB: a resource of transcriptional regulation in the context of enhancers. *Database*. 2019;2019.
74. Sarvagalla S, Kolapalli SP, Vallabhapurapu S. The Two Sides of YY1 in Cancer: A Friend and a Foe. *Front Oncol*. 2019;9:1230. Epub 2019/12/12.
75. Weintraub AS, Li CH, Zamudio AV, Sigova AA, Hannett NM, Day DS, et al. YY1 Is a Structural Regulator of Enhancer-Promoter Loops. *Cell*. Dec 14 2017;171(7):1573-88.e28. Epub 2017/12/12.
76. Kamal F, Schott EM, Carlson EL, El-Quadi M, Le Bleu HK, Hilton MJ, et al. Chondrocyte PTH1R anti-hypertrophic signaling is essential for articular cartilage maintenance and protection post trauma. *Osteoarthritis Cartilage*. 2017;25:S13-S4.
77. Sampson ER, Hilton MJ, Tian Y, Chen D, Schwarz EM, Mooney RA, et al. Teriparatide as a chondroregenerative therapy for injury-induced osteoarthritis. *Sci Transl Med*. Sep 21 2011;3(101):101ra93. Epub 2011/09/23.

773 78. Khundmiri SJ, Murray RD, Lederer E. PTH and Vitamin D. Compr Physiol. Mar 15 2016;6(2):561-
774 601. Epub 2016/04/12.

775

776 TABLES

777 **Supplementary Table 1.** *Candidate transcription factor binding sites within the Pth1r promoter.* Primary
778 chondrocytes were treated with NSC117079 for 24 hours and analyzed by ChIP-Seq. The DNA sequence
779 under the H3K27Ac peak in the Pth1r promoter (chromosome 9: 110742000-110744000) was used to
780 search EnhancerDB (<http://lcbj.swjtu.edu.cn/EnhancerDB/>) and identify candidate transcription factors
781 binding to this region.

TF Name	Start	End	Source	TF Name	Start	End	Source
ETS2	110743298	110743303	TRANSFAC	BHLHE40	110743562	110743572	JASPAR
	110743956	110743961	TRANSFAC	CEBPB	110743347	110743358	JASPAR
H1TF2	110743420	110743424	TRANSFAC	CLOCK	110743562	110743572	JASPAR
	110743687	110743691	TRANSFAC	DMRT3	110743247	110743258	JASPAR
MYB	110743245	110743248	TRANSFAC	EBF1	110743834	110743845	JASPAR
	110743255	110743260	TRANSFAC	EOMES	110743582	110743595	JASPAR
	110743373	110743378	TRANSFAC	FOXF2	110743351	110743365	JASPAR
	110743726	110743729	TRANSFAC	FOXG1	110743356	110743364	JASPAR
MYC	110743564	110743569	TRANSFAC	FOXP1	110743934	110743949	JASPAR
NFE	110743420	110743424	TRANSFAC		110743939	110743954	JASPAR
	110743620	110743624	TRANSFAC		110743944	110743959	JASPAR
	110743687	110743691	TRANSFAC		110743959	110743974	JASPAR
SRF	110743420	110743424	TRANSFAC		110743968	110743983	JASPAR
	110743687	110743691	TRANSFAC		110743974	110743989	JASPAR
TEAD2	110743724	110743728	TRANSFAC		110743976	110743991	JASPAR
USF1, USF2	110743564	110743569	TRANSFAC		110743977	110743992	JASPAR
YY1	110743204	110743208	TRANSFAC		110743982	110743997	JASPAR
	110743469	110743473	TRANSFAC		110743984	110744006	JASPAR
	110743485	110743489	TRANSFAC	HEY1	110743562	110743572	JASPAR
	110743558	110743562	TRANSFAC	HOXC10, HOXC11, HOXC12, HOXC13, HOXD11	110743440	110743451	JASPAR
	110743652	110743656	TRANSFAC	IRF1, IRF7,	110743382	110743394	JASPAR

	110743932	110743936	TRANSFAC	IRF8, IRF9	110743927	110743948	JASPAR
					110743933	110743954	JASPAR
					110743934	110743955	JASPAR
					110743938	110743959	JASPAR
					110743939	110743960	JASPAR
					110743943	110743964	JASPAR
					110743949	110743970	JASPAR
					110743954	110743975	JASPAR
					110743963	110743984	JASPAR
					110743967	110743988	JASPAR
					110743969	110743990	JASPAR
					110743973	110743994	JASPAR
					110743975	110743996	JASPAR
					110743976	110743997	JASPAR
					110743977	110743998	JASPAR
					110743981	11074402	JASPAR
					110743982	11074403	JASPAR
					110743983	110744004	JASPAR
				JUN	110743236	110743249	JASPAR
				MAFF, MAFK, MAFG	110743448	110743471	JASPAR
					110743623	110743638	JASPAR
					110743767	110743785	JASPAR
				MAX	110743562	110743572	JASPAR
				MEF2A, MEF2B, MEF2C, MEF2D	110743465	110743481	JASPAR
					110743728	110743740	JASPAR
				MFZ1	110743786	110743792	JASPAR
				MNT	110743562	110743572	JASPAR
				NFATC2, NFATC3	110743390	110743400	JASPAR
				NFE2L2	110743456	110743467	JASPAR
					110743902	110743913	JASPAR
				NRL	110743627	110743638	JASPAR
					110743769	110743780	JASPAR
				ONECUT3	110743509	110743523	JASPAR
				PAX5	110743338	110743357	JASPAR
				PHOX2A	110743397	110743408	JASPAR
				POU4F1, POU4F2, POU4F3	110743445	110743461	JASPAR

	POU2F1, POU3F1, POU3F2, POU3F4, POU5F1B	110743600	110743613	JASPAR
	PRDM1	110743945	110743960	JASPAR
		110743956	110743971	JASPAR
		110743965	110743980	JASPAR
		110743969	110743984	JASPAR
	PROP1	110743397	110743408	JASPAR
	RELA	110743605	110743615	JASPAR
	RREB1	110743563	110743583	JASPAR
	SPIC	110743966	110743980	JASPAR
	STAT1	110743952	110743966	JASPAR
		110743966	110743980	JASPAR
	STAT2	110743880	110743895	JASPAR
		110743935	110743950	JASPAR
		110743940	110743955	JASPAR
		110743945	110743960	JASPAR
		110743951	110743966	JASPAR
		110743956	110743971	JASPAR
		110743965	110743980	JASPAR
	STAT3	110743216	110743227	JASPAR
	TFAP2A, TFAP2B, TFAP2C	110743269	110743281	JASPAR
		110743711	110743726	JASPAR
		110743833	110743846	JASPAR
		110743845	110743860	JASPAR
	TBR1	110743583	110743593	JASPAR
	TBX20, TBX21	110743582	110743593	JASPAR
	YY1	110743436	110743442	JASPAR
	ZBTB7B, ZBTB7C	110743272	110743284	JASPAR
	ZNF263	110743938	110743959	JASPAR
		110743942	110743963	JASPAR
		110743946	110743967	JASPAR
		110743955	110743976	JASPAR
		110743958	110743979	JASPAR

Supplementary Table 2. Antibodies.

Product Name	Source and Catalog #	Host Species	Application	Dilution	Application specific details
Mouse Phlpp1 (PH)	Sigma 07-1341	Rabbit	WB	1:1000	5% BSA, overnight,

domain and Leucine rich repeat Protein Phosphatase 1)					4°C
Mouse Pth1r (Parathyroid hormone receptor)	Sigma SAB4502493	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
			IHC	1:50	Abcam ab93697 Mouse and Rabbit Specific HRP Plus (ABC) Detection IHC Kit
Mouse H3PS10 (histone 3, serine 10 phosphorylation)	Abcam ab5176	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
Mouse H3K9ac (histone 3, lysine 9 acetylation)	Abcam ab10812	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
Mouse H3K27ac (histone 3, lysine 27 acetylation)	Abcam ab4729	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
			ChIP	2 µg	Overnight, 4°C
Mouse H3K27ac (histone 3, lysine 27 acetylation)	Cell Signaling Technology 8173BC	Rabbit	ChIP-Sequencing	2 µg	Overnight, 4°C
Mouse H3PS28 (histone 3, serine 28 phosphorylation)	Cell Signaling Technology 9713S	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
Mouse H3K8K14ac (histone 3, lysine 8, lysine 14 acetylation)	Cell Signaling Technology 9677S	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
Mouse Total H3 (histone 3)	Abcam ab1791	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
Mouse YY1 (Ying Yang 1)	Active Motif 61780	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
Mouse pCREB-S133 (cAMP response element-binding protein, serine 133 phosphorylation)	Cell Signaling Technology 9198S	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
Mouse Total CREB (cAMP response element-binding protein)	Cell Signaling Technology 4820	Rabbit	WB	1:1000	5% BSA, overnight, 4°C
Mouse Actin	Sigma A4700	Mouse	WB	1:1000	5% BSA, overnight, 4°C
Anti-Rabbit IgG, HRP-linked	Cell Signaling Technology 7074S	Goat	WB	1:10,000	TBSTw, 1h, RT
Normal Rabbit IgG	Cell Signaling Technology 2729	Rabbit	IHC	1:50	Abcam ab93697 Mouse and Rabbit Specific HRP Plus (ABC) Detection IHC Kit

Supplementary Table 3. Biological Modulators.

Modulator	Source and Catalog #	Solvent / Vehicle	Concentration
NSC117079	Glix Laboratories	DMSO	25µM

	GLXC-03994		
NSC45586	Glxxx Laboratories GLXC-03991	DMSO	25µM
PTH(1-34)	Bachem 4011474	0.1% BSA in PBS	10nM or 100nM
PTH(7-34)	Bachem 4016931	0.1% BSA in PBS	In vivo: 100 µg/kg body weight/day In vitro: 10 nM or 100nM

786

A **B** **C** **D** **E**

Pth1r

Vehicle
NSC117079

Limb E10.5

Limb E15.5

Ensembl Mus musculus version 100.38 (GRCm38.p6) Chr 9: 110,722,085 - 110,747,145

A **B**

Male

Female

A **B** **C** **D**