Research article

Light-dependent induction of *Edn2* expression and attenuation of retinal pathology by endothelin receptor antagonists in *Prominin-1*-deficient mice

Yuka Kobayashi¹, Shizuka Watanabe², Manabu Shirai³, Chiemi Yamashiro¹, Tadahiko Ogata¹,
Fumiaki Higashijima¹, Takuya Yoshimoto¹, Takahide Hayano⁴, Yoshiyuki Asai⁴,
Noriaki Sasai^{2,5,*} and Kazuhiro Kimura^{1,5,*}

¹Department of Ophthalmology, Yamaguchi University Graduate School of Medicine, 1-1-1 Minami-kogushi, Ube 755-0046, Japan

²Developmental Biomedical Science, Division of Biological Sciences, Nara Institute of Science and Technology, 8916-5 Takayama-cho, Ikoma 630-0192, Japan

³Omics Research Center (ORC), National Cerebral and Cardiovascular Center, 6-1 Kishibe Shinmachi, Suita, Osaka 564-8565, Japan

⁴Department of Systems Bioinformatics, Yamaguchi University Graduate School of Medicine, 1-1-1 Minami-kogushi, Ube 755-0046, Japan

⁵ Co-senior authors

***Address for correspondence:**

Kazuhiro Kimura (e-mail: k.kimura@yamaguchi-u.ac.jp) or

Noriaki Sasai (e-mail: noriakisasai@bs.naist.jp)

Running Title

Blocking endothelin relieves retinopathy

Abstract

Retinitis pigmentosa (RP) and macular dystrophy (MD) are prevalent retinal degenerative diseases associated with gradual photoreceptor death. These diseases are often caused by genetic mutations that result in degeneration of the retina postnatally after it has fully developed. The *Prominin-1* gene (*Prom1*) is a causative gene for RP and MD, and *Prom1*- knockout (KO) mice recapitulate key features of these diseases including light-dependent retinal degeneration and stenosis of retinal blood vessels. The mechanisms underlying progression of such degeneration have remained unknown, however. We here analysed early events associated with retinal degeneration in *Prom1*-KO mice. We found that photoreceptor cell death and glial cell activation occur between 2 and 3 weeks after birth. High-throughput analysis revealed that expression of the endothelin-2 gene (*Edn2*) was markedly up-regulated in the *Prom1*-deficient retina during this period. Expression of *Edn2* was also induced by light stimulation in *Prom1*-KO mice that had been reared in the dark. Finally, treatment with endothelin receptor antagonists attenuated photoreceptor cell death, gliosis, and retinal vessel stenosis in *Prom1*-KO mice. Our findings suggest that inhibitors of endothelin signalling may delay the progression of RP and MD and therefore warrant further study as potential therapeutic agents for these diseases.

Keywords

prominin-1, photoreceptor, glial cell, retinal degeneration, endothelin-2, endothelin receptor antagonists

1. Introduction

Both retinitis pigmentosa (RP) and macular dystrophy (MD) are inherited retinal disorders associated with progressive photoreceptor cell death [1]. These diseases have a combined prevalence of 1 in 3000 to 4000 people worldwide. Initial symptoms include nyctalopia (night blindness) and visual field deficits, which are followed by loss of visual acuity and colour blindness and eventually by complete blindness. Although >60 genes encoding various types of protein - including membrane proteins, transcription factors, splicing regulators, and enzymes related to the visual cycle - have been implicated in RP and MD [1], these conditions remain incurable, with effective therapeutic strategies remaining to be established, and they have profound effects on the quality of life.

The *Prominin-1* gene (*Prom1*, also known as *AC133*, *CD133*, and *RP41*) encodes a pentaspan transmembrane glycoprotein that is expressed in photoreceptor cells of the retina as well as in kidney and testis [2]. Several mutations of *Prom1* have been identified in individuals with RP or MD [3-5], with all such mutations resulting in amino acid substitutions or carboxyl-terminal truncations of the encoded protein. The mechanisms underlying RP and MD associated with *Prom1* mutations have been investigated by studies of several lines of *Prom1*-knockout (KO) mice [5-7]. Although photoreceptor cells develop normally in these KO mice, they begin to degenerate after birth, resulting in a progressive loss of the outer nuclear layer (ONL) of the retina and recapitulation of the signs of RP and MD. The retinal vasculature also becomes attenuated with disease progression [7].

We previously showed that photoreceptor cells of the *Prom1*-KO mouse retina degenerate in response to light stimulation. Such mice reared in a completely dark setting thus manifested a marked delay in the loss of photoreceptor cells. We therefore suggested that the mutant retinal cells are hypersensitive to light stimulation and experience phototoxicity [6]. The visual cycle was also found to be impaired in the *Prom1*-KO cells, and treatment based on chemical compounds that modulate the visual cycle was found to mitigate the mutant phenotype [6].

The *Prom1* protein localises to the connecting cilium and outer segment of both rod and cone photoreceptors [3]. Ultrastructural analysis revealed the structure of the outer segment to be severely disorganised in photoreceptor cells of *Prom1*-KO mice, whereas other photoreceptor components - including the inner segment, nucleus, and axon - remained largely intact [6, 7]. Biochemical analysis has shown that two tyrosine residues in the carboxyl-terminal region of *Prom1* are phosphorylated by the tyrosine kinases Src and Fyn, although the physiological implications of such phosphorylation remain to be elucidated [8]. *Prom1* has also been shown to interact with the p85 regulatory subunit of phosphatidylinositol 3-kinase (PI3K) and to be essential for both the self-renewal and tumourigenic capacity of glioma stem cells [9]. In addition, *Prom1* has been detected in cilia, which are protrusive structures at the cell membrane and key signalling hubs [10], and to be essential for maximisation of Hedgehog signalling in neural stem cells [11]. We recently showed that *Prom1* activates the small GTPase Rho and regulates chloride conductance triggered by intracellular calcium uptake [12].

To characterise the mechanisms underlying the role of *Prom1* dysfunction in retinal degeneration and thereby to provide insight into potential treatments for *Prom1* mutation-associated RP and MD, we here investigated the initial manifestations of such degeneration. We analysed *Prom1* expression as well as the ONL transition in *Prom1*-KO mice. We then performed a high-throughput expression analysis to identify genes responsible for degeneration of the *Prom1*-deficient retina. Our results implicated an inflammatory pathway dependent on the endothelin 2 gene (*Edn2*), and we found that a chemical treatment targeted to endothelin signalling mitigated the deterioration of retinal structure and function in *Prom1*-KO mice.

2. Methods

2.1. Mice

Prom1-KO mice were established previously (CDB0623K, <http://www2.clst.riken.jp/arg/methods.html>), and they were reared on a hybrid genetic background of C57BL/6 and CBA/NSlc strains. The targeting vector for *Prom1* ablation contained the *lacZ* (β -galactosidase) gene, with the result that expression of this latter gene reflects that of *Prom1*. Both the *Prom1*-KO mice and their wild-type (WT) littermates were kept on a 12-hour-light, 12-hour-dark cycle, with the cage racks being covered with blackout curtains and all procedures including feeding and cage maintenance being performed in the absence of light (<0.5 lux) during the dark phase. For experiments involving light stimulation, mice were exposed for 3 h to a light panel (LED viewer 5000; Shinko, Tokyo, Japan) placed on top of the cage, which resulted in a light intensity of 3800 lux at the bottom of the cage. For chemical treatment, mice received intraperitoneal injections (2 mg/kg) of each of the endothelin receptor antagonists BQ-123 (ab141005, Abcam) and BQ-788 (ab144504, Abcam) on postnatal day (P) 14, P19, and P24. The mice were then subjected to analysis on P28.

2.2. RNA extraction and RT-qPCR analysis

The retina, retinal pigment epithelium (RPE), and testis were dissected from mice killed by cervical dislocation. Total RNA was extracted from the isolated tissue and was subjected to reverse transcription (RT) with the use of a NucleoSpin RNA extraction kit (U955C, Takara) and PrimeScript RT reagent kit (RR037, Takara), respectively. The resulting cDNA was subjected to quantitative polymerase chain reaction (qPCR) analysis with a CFX qPCR machine (Bio-Rad) and with primers listed in supplementary table S1. The amplification data were analysed with the comparative C_t method, and gene expression levels were normalised by that of the glyceraldehyde-3-phosphate dehydrogenase gene (*Gapdh*).

2.3. High-throughput expression analysis

Total RNA samples were prepared from three (P14) or four (P21) retinas of WT or *Prom1*-KO mice and were used to synthesise cDNA libraries with a TruSeq stranded-mRNA library preparation kit (Illumina, 20020594). The libraries were sequenced with the NextSeq 500 platform (Illumina). In total, approximately twenty million reads/sample were mapped with the CLC genomics workbench software (Qiagen) [13]. The sequencing data were deposited in the DNA Data Bank of Japan (DDBJ) public database, with the accession number of SSUB016168. Gene ontology (GO) term analysis was performed according to the Kyoto Encyclopaedia of Genes and Genomes database (KEGG, <https://www.genome.jp/kegg>).

2.4. Immunofluorescence analysis, β -galactosidase and isolectin staining, and TUNEL analysis

For immunofluorescence analysis, the enucleated retina was fixed for 2 h with a mixture of 1% paraformaldehyde and 0.2% glutaraldehyde in phosphate-buffered saline (PBS), incubated overnight in PBS containing 15% sucrose, embedded in O.C.T. compound (Sakura), and sectioned at a thickness of 12 μ m. The sections were exposed to mouse monoclonal antibodies to GFAP (G3893; Sigma) or rabbit polyclonal antibodies to Iba-1 (019-19741; Wako), and immune complexes were detected with Cy3-conjugated secondary antibodies (715-166-151 and 715-166-152 for mouse and rabbit, respectively; Jackson Immunoresearch). Nuclei were counterstained with 4',6-diamidino-2-phenylindole (DAPI) with the use of DAPI Fluoromount-G (0100-20; Southern Biotech). Sections were also stained for β -galactosidase (β -gal) activity with the use of a staining kit (11828673001, Roche). Apoptotic cells were detected by TUNEL analysis with digoxigenin-labelled dUTP (S7105, Merck Millipore), terminal deoxynucleotidyl transferase (3333566001, Merck), and rhodamine-conjugated antibodies to digoxigenin (11207750910, Roche). For preparation of flat-mount samples, the retina was fixed for 150 min with 4% paraformaldehyde and the RPE was peeled off. The samples were subjected to isolectin staining by consecutive exposure to 5% dried skim milk and Alexa Flour 488-conjugated GS-IB4 (I21411, Thermo Fisher Scientific) as described previously [14]. Images were acquired with an LSM 710 confocal microscope (Zeiss) for immunofluorescence, β -gal, and TUNEL staining, or with a BZ-X710 microscope (Keyence) for flat-mount preparations. Imaging data were processed and integrated with Photoshop (Adobe) and Illustrator (Adobe) software, respectively.

2.5. Statistical analysis

Quantitative data are presented as means $\pm$ s.e.m. Differences between two or among more than two groups were evaluated with the two-tailed Student's *t* test and by one-way analysis of variance (ANOVA) followed by Tukey's post hoc test, respectively. Statistical analysis was performed with Prism software (Graphpad), and a *p* value of <0.05 was considered statistically significant.

3. Results

3.1. *Prom1* is expressed in the retina from perinatal to adult stages

We previously showed that retinal cells in *Prom1*-KO mice appear to develop normally before the onset of degeneration [6]. We here first examined the spatiotemporal expression of *Prom1* in the mouse retina. Given that our *Prom1*-KO mice harbour the *lacZ* gene at the *Prom1* locus, we performed staining for β -gal activity in the heterozygous mutant mice at birth as well as at P2 (figure 1a-a''), P14 (figure 1b-b''), P21 (figure 1c-c''), and P42 (figure 1d-d''). At all the stages analysed, β -gal staining was localised predominantly to the outer layers in the retina, with more sporadic staining also apparent in the inner nuclear layer (INL). Given that retinal phenotypes of *Prom1*-KO mice are not obvious until 2 weeks after birth, these results suggested that *Prom1* expression precedes the onset of function of the encoded protein in postnatal retinal homeostasis.

3.2. The *Prom1*-KO mouse retina manifests both apoptosis and an inflammatory response at 3 weeks after birth

We previously showed that the retina of *Prom1*-KO mice appears normal at P14 and begins to degenerate soon after the animals first open their eyes at P14 [6]. We therefore investigated whether the *Prom1*-deficient retina might undergo apoptosis in response to light exposure. Whereas the TUNEL assay revealed few apoptotic cells in the retina of WT or *Prom1*-KO mice at P14 (figure 2a and b), a significant increase in the number of TUNEL-positive cells, located mainly in the ONL, was detected at P21 in the *Prom1*-KO retina (figure 2c-e). These results suggested that programmed cell death by apoptosis begins to occur in the ONL of the retina between 2 and 3 weeks after birth in *Prom1*-KO mice.

Glial fibrillary acidic protein (GFAP) is an intermediate filament protein that is expressed by Müller glia in response to retinal injury [15, 16]. Similarly, Iba-1 is a scaffold protein that is expressed in microglia and which is up-regulated during an inflammatory response [17, 18]. We therefore next examined whether the *Prom1*-KO retina might undergo light-induced inflammation by analysing the expression of these two proteins. Immunofluorescence analysis revealed that, whereas both GFAP and Iba-1 were essentially undetectable in the WT or *Prom1*-KO retina at P14 (figure 2f-i), a marked increase in the extent of staining for both proteins was observed in the *Prom1*-KO retina at P21 (figure 2j-m), suggesting that the increased cell death that occurs in the ONL of the mutant mice after birth is accompanied by the activation of glial cells.

3.3. Inflammation-related gene expression is up-regulated in the *Prom1*-KO mouse retina

We next sought to identify genes whose expression might be affected by *Prom1* deficiency by subjecting the retina of WT and *Prom1*-KO mice at P14 and P21 to high-throughput expression analysis based on RNA sequencing. Gene expression at P14 tended to vary within each genotype, and the only gene whose expression differed significantly between genotypes was *Prom1* itself (figure 3a, supplementary table S2), suggesting that *Prom1* does not significantly influence the gene expression

profile at P14. In contrast, the expression of various genes differed between the two genotypes at P21 (figure 3b, supplementary table S3). The expression of 1,081 and 766 genes was thus up- and down-regulated, respectively, in the *Prom1*-KO retina with a *p* value of <0.01. In particular, expression of *Edn2* was the most consistently and markedly up-regulated in the *Prom1*-KO retina. The expression of genes associated with the inflammatory response - such as *Ifi44l*, *Serpina3n*, *S100a6*, *Bcl3*, and *Gfap* - was also increased in the *Prom1*-KO retina at P21. Conversely, the expression of genes related to RP or of those essential for retinal development and functional homeostasis - including *Fscn2* (RP30) [19], *Prph2* (RP7) [20], *Nr2e3* (RP37) [21], *Kcnv2* [22], *Elovl2* [23], *Pde6b* (RD1) [24], and *Ttc21b* [25] - was down-regulated in the *Prom1*-KO retina at P21 (supplementary table S3). GO term analysis revealed that several signalling pathways, including apoptotic (TNF) and infectious-related signal (Epstein-Barr virus infection) signals, were affected by the loss of *Prom1* (figure 3c).

We also investigated whether the observed effects of *Prom1* deficiency on gene expression were specific to the retina. Given that *Prom1* is expressed in the retina, RPE, and testis [2], we performed RT-qPCR analysis of RNA prepared from these tissues of WT and *Prom1*-KO mice at P21. Consistent with the results of our RNA-sequencing analysis, the expression of *Edn2*, *Bcl3*, and *Gfap* was increased in the retina of *Prom1*-KO mice (figure 3d). However, the expression of these genes in the RPE and testis did not differ between the two genotypes, indicating that the effect of *Prom1* on their expression is specific to the retina. Together, these various data suggested that *Prom1* deficiency results in up-regulation of inflammation-related genes and down-regulation of genes essential for functional homeostasis of photoreceptor cells at 3 weeks after birth.

3.4. Inflammation-related gene expression is increased by light stimulation in the *Prom1*-KO mouse retina

To determine the mechanism underlying the up-regulation of specific gene expression apparent in the retina of *Prom1*-KO mice at P21, we examined whether light stimulation might play a role. We therefore compared such gene expression between P21 retinas obtained from *Prom1*-KO mice reared under a normal day-night cycle or in the dark. RT-qPCR analysis revealed that, whereas the expression of *Edn2*, *Bcl3*, and *Gfap* did not differ between *Prom1*-KO and WT mice reared in the dark condition, marked up-regulation of the expression of each of these genes was apparent specifically in *Prom1*-KO mice raised under the normal day-night condition (figure 4a). Consistent with these results, immunofluorescence analysis showed that the number of GFAP-positive cells in the retina was smaller for *Prom1*-KO mice reared in the dark compared with those reared under the normal condition (figure 4b and c). To examine further the effect of light on gene expression, we maintained *Prom1*-KO mice and their WT littermates under the dark condition for 3 weeks, exposed them to a bright light for 3 h, and then allowed them to recover for 3 days in the dark. The retina was then dissected and subjected to RT-qPCR and immunofluorescence analyses. Light stimulation resulted in a marked increase both in the expression of *Edn2* and *Bcl3* (Figure 4d) and in the number of GFAP-positive cells (figure 4e) in the retina of *Prom1*-

KO mice but not in that of WT mice. Collectively, these results thus suggested that the up-regulation of *Edn2*, *Bcl3*, and *Gfap* expression apparent in the retina of *Prom1*-KO mice is an immediate response to light stimulation, and that the inflammatory response mediated by these genes is one of the primary events leading to degeneration of the mutant retina.

3.5. Endothelin receptor antagonists attenuate *Gfap* expression and gliosis in the *Prom1*-KO mouse retina

Endothelin acts at specific receptors [26, 27] to increase both the number of GFAP-positive Müller cells [28] and retinal cell death [29]. Given the elevated expression of *Edn2* and *Gfap* apparent in the retina of *Prom1*-KO mice, we hypothesised that *Edn2* might induce aberrant proliferation of glial cells and GFAP expression in association with retinal degeneration in these animals. We therefore examined the possible effects of endothelin receptor antagonists in the mutant mice.

The drugs BQ-123 and BQ-788, which target endothelin receptors A and B, respectively [30], were both injected intraperitoneally into *Prom1*-KO mice at P14, P19, and P24, and the mice were analysed at P28. Whereas GFAP-positive cells were not observed in the retina of WT mice, they were detected in that of *Prom1*-KO mice treated with dimethyl sulphoxide (DMSO) vehicle (figure 5a and b). However, the number of GFAP-positive cells was markedly reduced in the mutant mice by treatment with BQ-123 and BQ-788 (figure 5c). Staining of retinal flat-mount preparations with fluorescently labelled isolectin to detect vascular endothelial cells also revealed fewer retinal vessels in *Prom1*-KO mice than in WT mice and that this difference was attenuated by treatment of the mutant animals with BQ-123 and BQ-788 (figure 5d–g).

RT-qPCR analysis showed that the expression of *Edn2*, *Bcl3*, and *Gfap* was increased in the retina of *Prom1*-KO mice at P28 compared with that in WT mice. Whereas the expression of *Edn2* and *Bcl3* in the mutant retina was not affected by treatment with BQ-123 and BQ-788, that of *Gfap* was significantly attenuated (figure 5h), suggesting that up-regulation of *Gfap* expression in the mutant retina is mediated by endothelin receptor signalling but that that of *Edn2* and *Bcl3* expression is not.

Finally, we examined the effect of BQ-123 and BQ-788 treatment on the number of apoptotic cells in the retina of *Prom1*-KO mice. The TUNEL assay revealed that the marked increase in the number of such cells apparent in the mutant retina at P28 was significantly attenuated by administration of the two drugs (figure 5i), suggesting that endothelin receptor signalling contributes to loss of retinal cell homeostasis.

4. Discussion

We have here described early manifestations of the retinal degeneration that occurs in *Prom1*-KO mice and identified related genes. We thus detected the aberrant presence of glial cells and the expression of genes associated with the inflammatory response in the mutant retina. Given that the expression of these genes was not activated in the retina of *Prom1*-KO mice maintained in the dark condition, this inflammatory response appears to be dependent on light stimulation. Finally, we found that the deterioration and gliosis characteristic of the mutant retina were ameliorated by the administration of endothelin receptor antagonists.

Although we found that *Prom1* is expressed in the retina from birth, the loss of *Prom1* did not substantially affect the expression level of any gene in the retina at P14, suggesting that *Prom1* may not play an essential role in the retina prior to light exposure. We previously showed by RT-qPCR analysis that the expression of both *Rdh12* and *Abca4*, two genes that contribute to the visual cycle, was reduced in the retina of *Prom1*-KO mice compared with that of WT mice at P14 [6], suggesting that impairment of the visual cycle might lead to retinal degeneration. Although this result is reproducible as assayed by RT-qPCR (supplementary figure S1), the difference in the expression level of each gene between the two genotypes was associated with a relatively high *p* value in the high-throughput expression analysis performed in the present study (figure 3, supplementary table 2), suggesting this decrease is not critical.

In contrast to the lack of an effect of *Prom1* deficiency on the gene expression profile of the retina at P14, we detected many genes, including those related to the inflammatory response, as well as signalling pathways whose activity was altered in the *Prom1*-KO retina at P21. The expression of genes related to phototransduction, for example, was significantly down-regulated in the *Prom1*-KO retina at P21 (figure 3c, supplementary table S3), indicating that *Prom1* may be essential for the transcription of such genes or may form a transcriptional network with them. Of note, we found that the expression of causal genes for RP was also down-regulated in the mutant retina at P21.

Of the genes whose expression was up-regulated in the *Prom1*-KO retina at P21, *Edn2* showed the largest fold change. *Edn2* encodes a secretory peptide that plays a role in a wide range of biological processes, including smooth muscle contraction and ovulation [31] as well as development of the enteric nervous system [32]. Its expression is also induced in association with the inflammatory response and promotes glial cell proliferation in the central nervous system [33]. Furthermore, consistent with the perturbation of the retinal vasculature in *Prom1*-KO mice apparent in both the present and a previous [7] study, *Edn2* has been found to inhibit retinal vascular development [34]. On the other hand, it was also shown to promote photoreceptor cell survival [35]. These various observations suggest that the role of *Edn2* in the photoreceptor degeneration associated with RP and MD is complex.

The expression of *Edn2* has also been shown to be up-regulated in other mouse models of RP [35], including retina-specific *Cdhr1*-KO mice [36], with *Prom1* and *Cdhr1* having been found to interact with each other [4]. In addition to *Edn2*, the other genes whose expression was affected in the *Prom1*-KO

mouse retina overlapped markedly with those affected in the conditional *Cdhr1*-KO mouse retina, suggesting that *Prom1* and *Cdhr1* may function in the same intracellular signalling pathways.

Although we found that the expression of *Edn2* and *Bcl3* in the *Prom1*-KO retina was induced by light stimulation, the mechanism underlying this effect remains unclear. Nevertheless, our study suggests the possibility that an imbalance in intracellular ions caused by the loss of *Prom1* (given that *Prom1* regulates chloride conductance activated by intracellular calcium uptake [12]) may impair the function of cytoplasmic organelles such as mitochondria and the endoplasmic reticulum, and thereby elicit a stress response. Studies to identify the transcriptional regulatory elements of *Edn2* and the corresponding transcription factors and upstream signalling pathways underlying its photoactivation are warranted.

Gliosis is a response to injury in the central nervous system and is associated with the appearance of GFAP-positive glial cells [26]. It is also a feature of certain neurodegenerative retinal diseases including RP [37], with gliosis in RP having been found to be related to several RP genes. Targeting of gliosis is therefore a potential clinical strategy to delay disease progression and ameliorate associated symptoms. We have now shown that administration of endothelin receptor antagonists attenuated both the appearance of GFAP-positive glial cells and vascular endothelial constriction in the retina of *Prom1*-KO mice. These findings indicate that blockade of endothelin signalling may be an effective clinical strategy for the treatment of gliosis. However, caution is warranted with such an approach for the treatment of RP, given the various functions of endothelins and the consequent potential for adverse systemic effects. Intravitreal injection of endothelin receptor antagonists may help to avoid such side effects. Gene therapy targeting endothelin receptor function is also a potential therapeutic approach for RP. Finally, replacement of dead tissue with functional cells through a regenerative medicine approach may be required for the successful treatment of RP and MD [38].

In conclusion, our results implicating up-regulation of *Edn2* expression in the retinal pathology of *Prom1*-KO mice suggest that localized pharmacological targeting of endothelin receptor signalling warrants further investigation as a clinical intervention for the prevention or treatment of retinal degenerative diseases such as RP and MD.

Ethics. All animal experiments were approved by the animal welfare and ethics committees of both Yamaguchi University (approval numbers J16021 and U16005 for K.K.) and Nara Institute of Science and Technology (approval numbers 1810 and 311 for N.S.) and were performed in accordance with the relevant guidelines and regulations.

Data availability. Data are available in the main text/figures and in the Supplementary Information.

Competing interests. The authors declare no competing interests.

Funding. This work was supported in part by grants-in-aid for scientific research from Japan Society for the Promotion of Science (17H03684 and 20H0326310 to N.S.; 20K09805 to K.K.), as well as by Novartis Pharma.

Acknowledgements. We thank Erika Yoshihara, Yukari Mizuno, and Ayaka Kataoka for technical assistance as well as other laboratory members for their support and discussion.

Author Contributions. KK and NS conceived the project; YK, NS, SW, MS, CY, TO, FH, TY performed experiments; YK, NS, MS, TH, YA analysed the data; All authors joined the discussion; NS, KK, YK wrote the manuscript.

References

- 1 Ferrari, S., Di Iorio, E., Barbaro, V., Ponzin, D., Sorrentino, F. S., Parmeggiani, F. 2011 Retinitis pigmentosa: genes and disease mechanisms. *Curr Genomics*. **12**, 238-249. (10.2174/138920211795860107)
- 2 Fargeas, C. A., Joester, A., Missol-Kolka, E., Hellwig, A., Huttner, W. B., Corbeil, D. 2004 Identification of novel Prominin-1/CD133 splice variants with alternative C-termini and their expression in epididymis and testis. *Journal of cell science*. **117**, 4301-4311. (10.1242/jcs.01315)
- 3 Maw, M. A., Corbeil, D., Koch, J., Hellwig, A., Wilson-Wheeler, J. C., Bridges, R. J., Kumaramanickavel, G., John, S., Nancarrow, D., Roper, K., *et al.* 2000 A frameshift mutation in prominin (mouse)-like 1 causes human retinal degeneration. *Human molecular genetics*. **9**, 27-34.
- 4 Yang, Z., Chen, Y., Lillo, C., Chien, J., Yu, Z., Michaelides, M., Klein, M., Howes, K. A., Li, Y., Kaminoh, Y., *et al.* 2008 Mutant prominin 1 found in patients with macular degeneration disrupts photoreceptor disk morphogenesis in mice. *The Journal of clinical investigation*. **118**, 2908-2916. (10.1172/JCI35891)
- 5 Michaelides, M., Gaillard, M. C., Escher, P., Tiab, L., Bedell, M., Borruat, F. X., Barthelmes, D., Carmona, R., Zhang, K., White, E., *et al.* 2010 The PROM1 mutation p.R373C causes an autosomal dominant bull's eye maculopathy associated with rod, rod-cone, and macular dystrophy. *Investigative ophthalmology & visual science*. **51**, 4771-4780. (10.1167/iovs.09-4561)
- 6 Dellett, M., Sasai, N., Nishide, K., Becker, S., Papadaki, V., Limb, G. A., Moore, A. T., Kondo, T., Ohnuma, S. 2015 Genetic background and light-dependent progression of photoreceptor cell degeneration in Prominin-1 knockout mice. *Investigative ophthalmology & visual science*. **56**, 164-176. (10.1167/iovs.14-15479)
- 7 Zacchigna, S., Oh, H., Wilsch-Brauninger, M., Missol-Kolka, E., Jaszai, J., Jansen, S., Tanimoto, N., Tonagel, F., Seeliger, M., Huttner, W. B., *et al.* 2009 Loss of the cholesterol-binding protein prominin-1/CD133 causes disk dysmorphogenesis and photoreceptor degeneration. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. **29**, 2297-2308. (10.1523/JNEUROSCI.2034-08.2009)
- 8 Boivin, D., Labbe, D., Fontaine, N., Lamy, S., Beaulieu, E., Gingras, D., Beliveau, R. 2009 The stem cell marker CD133 (prominin-1) is phosphorylated on cytoplasmic tyrosine-828 and tyrosine-852 by Src and Fyn tyrosine kinases. *Biochemistry*. **48**, 3998-4007. (10.1021/bi900159d)
- 9 Wei, Y., Jiang, Y., Zou, F., Liu, Y., Wang, S., Xu, N., Xu, W., Cui, C., Xing, Y., Liu, Y., *et al.* 2013 Activation of PI3K/Akt pathway by CD133-p85 interaction promotes tumorigenic capacity of glioma stem cells. *Proceedings of the National Academy of Sciences of the United States of America*. **110**, 6829-6834. (10.1073/pnas.1217002110)
- 10 Khatri, P., Obernier, K., Simeonova, I. K., Hellwig, A., Holzl-Wenig, G., Mandl, C., Scholl, C., Wolf, S., Winkler, J., Gaspar, J. A., *et al.* 2014 Proliferation and cilia dynamics in neural stem cells prospectively isolated from the SEZ. *Scientific reports*. **4**, 3803. (10.1038/srep03803)
- 11 Singer, D., Thamm, K., Zhuang, H., Karbanova, J., Gao, Y., Walker, J. V., Jin, H., Wu, X., Coveney, C. R., Marangoni, P., *et al.* 2019 Prominin-1 controls stem cell activation by orchestrating ciliary dynamics. *The EMBO journal*. **38**, (10.15252/embj.201899845)
- 12 Hori, A., Nishide, K., Yasukuni, Y., Haga, K., Kakuta, W., Ishikawa, Y., Hayes, M. J., Ohnuma, S. I., Kiyonari, H., Kimura, K., *et al.* 2019 Prominin-1 Modulates Rho/ROCK-Mediated Membrane Morphology and Calcium-Dependent Intracellular Chloride Flux. *Scientific reports*. **9**, 15911. (10.1038/s41598-019-52040-9)
- 13 Robinson, M. D., McCarthy, D. J., Smyth, G. K. 2010 edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. **26**, 139-140. (10.1093/bioinformatics/btp616)
- 14 Yamaguchi, M., Nakao, S., Arita, R., Kaizu, Y., Arima, M., Zhou, Y., Kita, T., Yoshida, S., Kimura, K., Isobe, T., *et al.* 2016 Vascular Normalization by ROCK Inhibitor: Therapeutic Potential of Ripasudil (K-115) Eye Drop in Retinal Angiogenesis and Hypoxia. *Investigative ophthalmology & visual science*. **57**, 2264-2276. (10.1167/iovs.15-17411)
- 15 Chang, M. L., Wu, C. H., Jiang-Shieh, Y. F., Shieh, J. Y., Wen, C. Y. 2007 Reactive changes of retinal astrocytes and Muller glial cells in kainate-induced neuroexcitotoxicity. *J Anat*. **210**, 54-65. (10.1111/j.1469-7580.2006.00671.x)

- 16 Lewis, G. P., Fisher, S. K. 2003 Up-regulation of glial fibrillary acidic protein in response to retinal injury: its potential role in glial remodeling and a comparison to vimentin expression. *Int Rev Cytol.* **230**, 263-290. (10.1016/s0074-7696(03)30005-1)
- 17 Rojas, B., Gallego, B. I., Ramirez, A. I., Salazar, J. J., de Hoz, R., Valiente-Soriano, F. J., Aviles-Trigueros, M., Villegas-Perez, M. P., Vidal-Sanz, M., Trivino, A., *et al.* 2014 Microglia in mouse retina contralateral to experimental glaucoma exhibit multiple signs of activation in all retinal layers. *J Neuroinflammation.* **11**, 133. (10.1186/1742-2094-11-133)
- 18 Omri, S., Behar-Cohen, F., de Kozak, Y., Sennlaub, F., Verissimo, L. M., Jonet, L., Savoldelli, M., Omri, B., Crisanti, P. 2011 Microglia/macrophages migrate through retinal epithelium barrier by a transcellular route in diabetic retinopathy: role of PKCzeta in the Goto Kakizaki rat model. *Am J Pathol.* **179**, 942-953. (10.1016/j.ajpath.2011.04.018)
- 19 Wada, Y., Abe, T., Takeshita, T., Sato, H., Yanashima, K., Tamai, M. 2001 Mutation of human retinal fascin gene (FSCN2) causes autosomal dominant retinitis pigmentosa. *Investigative ophthalmology & visual science.* **42**, 2395-2400.
- 20 Conley, S. M., Naash, M. I. 2014 Gene therapy for PRPH2-associated ocular disease: challenges and prospects. *Cold Spring Harbor perspectives in medicine.* **4**, a017376. (10.1101/cshperspect.a017376)
- 21 Cheng, H., Khanna, H., Oh, E. C., Hicks, D., Mitton, K. P., Swaroop, A. 2004 Photoreceptor-specific nuclear receptor NR2E3 functions as a transcriptional activator in rod photoreceptors. *Human molecular genetics.* **13**, 1563-1575. (10.1093/hmg/ddh173)
- 22 Holter, P., Kunst, S., Wolloscheck, T., Kelleher, D. K., Sticht, C., Wolfrum, U., Spessert, R. 2012 The retinal clock drives the expression of Kcnv2, a channel essential for visual function and cone survival. *Investigative ophthalmology & visual science.* **53**, 6947-6954. (10.1167/iovs.12-10234)
- 23 Chen, D., Chao, D. L., Rocha, L., Kolar, M., Nguyen Huu, V. A., Krawczyk, M., Dasyani, M., Wang, T., Jafari, M., Jabari, M., *et al.* 2020 The lipid elongation enzyme ELOVL2 is a molecular regulator of aging in the retina. *Aging Cell.* **19**, e13100. (10.1111/ace1.13100)
- 24 Yeo, J. H., Jung, B. K., Lee, H., Baek, I. J., Sung, Y. H., Shin, H. S., Kim, H. K., Seo, K. Y., Lee, J. Y. 2019 Development of a Pde6b Gene Knockout Rat Model for Studies of Degenerative Retinal Diseases. *Investigative ophthalmology & visual science.* **60**, 1519-1526. (10.1167/iovs.18-25556)
- 25 Liu, Q., Zhang, Q., Pierce, E. A. 2010 Photoreceptor sensory cilia and inherited retinal degeneration. *Advances in experimental medicine and biology.* **664**, 223-232. (10.1007/978-1-4419-1399-9_26)
- 26 Sarthy, V. P., Sawkar, H., Dudley, V. J. 2015 Endothelin2 Induces Expression of Genes Associated with Reactive Gliosis in Retinal Muller Cells. *Curr Eye Res.* **40**, 1181-1184. (10.3109/02713683.2014.982828)
- 27 Patel, C., Narayanan, S. P., Zhang, W., Xu, Z., Sukumari-Ramesh, S., Dhandapani, K. M., Caldwell, R. W., Caldwell, R. B. 2014 Activation of the endothelin system mediates pathological angiogenesis during ischemic retinopathy. *Am J Pathol.* **184**, 3040-3051. (10.1016/j.ajpath.2014.07.012)
- 28 Rattner, A., Toulabi, L., Williams, J., Yu, H., Nathans, J. 2008 The genomic response of the retinal pigment epithelium to light damage and retinal detachment. *The Journal of neuroscience : the official journal of the Society for Neuroscience.* **28**, 9880-9889. (10.1523/JNEUROSCI.2401-08.2008)
- 29 Kobayashi, T., Oku, H., Fukuhara, M., Kojima, S., Komori, A., Ichikawa, M., Katsumura, K., Kobayashi, M., Sugiyama, T., Ikeda, T. 2005 Endothelin-1 enhances glutamate-induced retinal cell death, possibly through ETA receptors. *Investigative ophthalmology & visual science.* **46**, 4684-4690. (10.1167/iovs.05-0785)
- 30 Fukuroda, T., Ozaki, S., Ihara, M., Ishikawa, K., Yano, M., Nishikibe, M. 1994 Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery. *Br J Pharmacol.* **113**, 336-338. (10.1111/j.1476-5381.1994.tb16901.x)
- 31 Cacioppo, J. A., Oh, S. W., Kim, H. Y., Cho, J., Lin, P. C., Yanagisawa, M., Ko, C. 2014 Loss of function of endothelin-2 leads to reduced ovulation and CL formation. *PloS one.* **9**, e96115. (10.1371/journal.pone.0096115)
- 32 Gershon, M. D. 1999 Endothelin and the development of the enteric nervous system. *Clin Exp Pharmacol Physiol.* **26**, 985-988. (10.1046/j.1440-1681.1999.03176.x)
- 33 Yuen, T. J., Johnson, K. R., Miron, V. E., Zhao, C., Quandt, J., Harrisingh, M. C., Swire, M., Williams, A., McFarland, H. F., Franklin, R. J., *et al.* 2013 Identification of endothelin 2 as an inflammatory factor that promotes central nervous system remyelination. *Brain.* **136**, 1035-1047. (10.1093/brain/awt024)

- 34 Rattner, A., Yu, H., Williams, J., Smallwood, P. M., Nathans, J. 2013 Endothelin-2 signaling in the neural retina promotes the endothelial tip cell state and inhibits angiogenesis. *Proceedings of the National Academy of Sciences of the United States of America*. **110**, E3830-3839. (10.1073/pnas.1315509110)
- 35 Bramall, A. N., Szego, M. J., Pacione, L. R., Chang, I., Diez, E., D'Orleans-Juste, P., Stewart, D. J., Hauswirth, W. W., Yanagisawa, M., McInnes, R. R. 2013 Endothelin-2-mediated protection of mutant photoreceptors in inherited photoreceptor degeneration. *PloS one*. **8**, e58023. (10.1371/journal.pone.0058023)
- 36 Rattner, A., Nathans, J. 2005 The genomic response to retinal disease and injury: evidence for endothelin signaling from photoreceptors to glia. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. **25**, 4540-4549. (10.1523/JNEUROSCI.0492-05.2005)
- 37 Roche, S. L., Ruiz-Lopez, A. M., Moloney, J. N., Byrne, A. M., Cotter, T. G. 2018 Microglial-induced Muller cell gliosis is attenuated by progesterone in a mouse model of retinitis pigmentosa. *Glia*. **66**, 295-310. (10.1002/glia.23243)
- 38 Stern, J. H., Tian, Y., Funderburgh, J., Pellegrini, G., Zhang, K., Goldberg, J. L., Ali, R. R., Young, M., Xie, Y., Temple, S. 2018 Regenerating Eye Tissues to Preserve and Restore Vision. *Cell Stem Cell*. **22**, 834-849. (10.1016/j.stem.2018.05.013)

Figure Legends

Figure 1. *Prom1* is expressed in the ONL of the retina from perinatal to adult stages. The retina of heterozygous *Prom1* mutant mice at P2 (*a-a''*), P14, (*b-b''*), P21 (*c-c''*), and P42 (*d-d''*) was subjected to staining of β -gal activity (*a,b,c,d*) as well as to staining of nuclei with DAPI (*a',b',c',d'*). Merged images are also shown (*a'',b'',c'',d''*). Data are representative of three retinas at each age. Scale bar in (*a*) is (50 μ m) and applies to all images. RPE, retinal pigment epithelium; NBL, neuroblast layer; GCL, ganglion cell layer; OS, outer segments; IS, inner segments; ONL, outer nuclear layer; OPL, outer plexiform layer; INL, inner nuclear layer; IPL, inner plexiform layer.

Figure 2. Programmed cell death and an inflammatory response in the postnatal *Prom1*-KO mouse retina. (*a-d''*) TUNEL staining of the WT (*a-a'',c-c''*) and *Prom1*-KO (*b-b'',d-d''*) mouse retina at P14 (*a-b''*) and P21 (*c-d''*). Nuclei were stained with DAPI (*a',b',c',d'*). Merged images of TUNEL (red) and DAPI (blue) staining are also shown (*a'',b'',c'',d''*). Arrowheads in (*d,d''*) indicate apoptotic cells. (*e*) Quantitation of the proportion of TUNEL-positive cells among all DAPI-stained cells for images similar to those in (*a*), (*b*), (*c*) and (*d*). Data are means $\pm$ s.e.m. for four retinas for each condition. ** $p < 0.01$; n.s., not significant (two-tailed Student's *t* test). (*f-m*) Immunofluorescence staining for GFAP (*f, h,j,l*) and Iba-1 (*g,i,k,m*) in the retina of WT (*f,g,j,k*) and *Prom1*-KO (*h,i,l,m*) mice at P14 (*f-i*) and P21 (*j-m*). Merged images with DAPI staining are also shown (*f',g',h',i',j',k',l',m'*). Arrowheads in (*m*) indicate Iba-1-positive cells. Data are representative of three (P14) or five (P21) retinas for each genotype. Scale bar in (*a*) is 50 μ m and applies to all images.

Figure 3. Effects of *Prom1* deficiency on gene expression in the retina. (*a,b*) Volcano plots for RNA-sequencing analysis of the retina of *Prom1*-KO mice relative to that of WT mice at P14 (*a*) and P21 (*b*). Genes with a p value of 1×10^{-10} are indicated with the blue arrowhead in (*b*). A cut-off p value of 1×10^{-2} is indicated by the green dashed line. Data are for three (P14) or four (P21) retinas of each genotype. (*c*) GO term analysis based on KEGG pathways for genes whose expression differed significantly between the retinas of *Prom1*-KO and WT mice in the RNA-sequencing analysis at P21. (*d*) RT-qPCR analysis of *Edn2*, *Bcl3*, and *Gfap* expression in the retina, RPE, and testis of WT and *Prom1*-KO mice at P21. Data are means $\pm$ s.e.m. for three retinas of each genotype. * $p < 0.05$, ** $p < 0.01$, n.s., not significant (two-tailed Student's *t* test).

Figure 4. Genes whose expression is increased by *Prom1* deficiency are up-regulated by light stimulation. (*a*) RT-qPCR analysis of *Edn2*, *Bcl3*, and *Gfap* expression in the P21 retina of WT or *Prom1*-KO mice that had been reared either under a normal day-night cycle or in the dark. Data are means $\pm$ s.e.m. for four retinas for each condition. * $p < 0.05$, ** $p < 0.01$, n.s., not significant, versus WT/normal (one-way ANOVA followed by Tukey's post hoc test). (*b* and *c*) Immunofluorescence analysis of GFAP expression in the retina of *Prom1*-KO mice raised as in (*a*). Merged images with DAPI staining are also shown. Scale

bar in (b) is 50 μ m and applies to all images. Data are representative of four (dark) or seven (normal day-night) retinas. (d) RT-qPCR analysis of *Edn2* and *Bcl3* expression in the retina of *Prom1*-KO and WT mice that had been reared in the dark condition for 3 weeks, exposed (or not) to a bright light for 3 h, and then allowed to recover in the dark for 3 days. Data are means $\pm$ s.e.m. for five retinas for each condition. $*p < 0.05$, n.s., not significant, versus WT/dark (one-way ANOVA followed by Tukey's post hoc test). (e) Immunofluorescence analysis of GFAP expression in the retina of *Prom1*-KO mice raised in the dark and stimulated with light as in (d). Merged images with DAPI staining are also shown. Data are representative of three retinas.

Figure 5. Endothelin receptor antagonists attenuate the increase in the number of GFAP-positive cells and vascular stenosis in the retina of *Prom1*-KO mice. (a–c) Immunofluorescence analysis of GFAP expression in the retina of WT (a) or *Prom1*-KO (b and c) mice treated with the combination of BQ-123 and BQ-788 (c) or with DMSO vehicle (a and b) at P14, P19, and P24 and analysed at P28. Merged images with DAPI staining are also shown. Scale bar in (a), 50 μ m. Data are representative of three retinas per condition. (d–f) Isolectin staining of the retina of mice as in (a) to (c). The boxed regions of the left panels are shown at higher magnification in the right panels. Scale bars, 100 μ m. (g) Area of blood vessels measured in images similar to those in (d) to (f). Data are means $\pm$ s.e.m. for X retinas per condition. $*p < 0.05$, $****p < 0.0001$ (one-way ANOVA followed by Tukey's post hoc test). (h) RT-qPCR analysis of *Edn2*, *Bcl3*, and *Gfap* expression in the retina of the treated mice. Data are means $\pm$ s.e.m. for three retinas per condition. $**p < 0.01$, n.s., not significant (one-way ANOVA followed by Tukey's post hoc test). (i–k) TUNEL staining for apoptotic cells in the retina of the treated mice. Merged images with DAPI staining are also shown. Scale bar in (i), 50 μ m. (l) Number of apoptotic cells determined from images similar to those in (i) to (k). Data are means $\pm$ s.e.m. for three retinas per condition. $**p < 0.01$, $***p < 0.001$ (one-way ANOVA followed by Tukey's post hoc test).

Supplementary Figure

Supplementary figure S1. Expression of *Rdh12* and *Abca4* is down-regulated in the retina of *Prom1*-KO mice at P14.

Supplementary Tables

Supplementary table S1. Primers used for this study.

Supplementary table S2. RNA-sequencing analysis of the retina of *Prom1*-KO and WT mice at P14.

Supplementary table S3. RNA-sequencing analysis of the retina of *Prom1*-KO and WT mice at P21.

Figure 1

Figure 2

Figure 3

Figure 4