Non-autophagy role of Atg5 and NBR1 in unconventional secretion of IL-12 prevents gut dysbiosis and inflammation

Seth D. Merkley^{1,#}, Samuel M. Goodfellow^{2,#}, Yan Guo¹, Zoe E.R. Wilton¹, Janie R. Byrum³, Kurt C. Schwalm⁴, Darrell L. Dinwiddie^{4,5}, Rama R. Gullapalli⁶, Vojo Deretic^{3,7}, Anthony Jimenez Hernandez¹, Steven B. Bradfute², Julie G. In^{1,8} and Eliseo F. Castillo^{*1,5,7}

¹ Division of Gastroenterology and Hepatology, Department of Internal Medicine, University of New Mexico Health Sciences, Albuquerque, NM.

² Center for Global Health, Department of Internal Medicine, University of New Mexico Health Sciences, Albuquerque, New Mexico.

³ Department of Molecular Genetics and Microbiology, University of New Mexico Health Sciences, Albuquerque, NM.

⁴ Department of Pediatrics, University of New Mexico Health Sciences, Albuquerque, NM.

⁵ Clinical and Translational Science Center, University of New Mexico Health Sciences, Albuquerque, New Mexico.

⁶ Department of Pathology, University of New Mexico Health Sciences, Albuquerque, NM.

⁷ Autophagy Inflammation and Metabolism Center of Biomedical Research Excellence, University of New Mexico Health Sciences, Albuquerque, NM.

⁸ Division of Gastroenterology and Hepatology, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD.

#Authors contributed equally to this study

*Correspondence should be addressed to: ecastillo@salud.unm.edu (E.F.C.)

Running title: Atg5 limits IL-12-mediated inflammation

Keywords: autophagy, macrophages, inflammation, cytokines, microbiota

ABSTRACT

Intestinal myeloid cells play a critical role in balancing intestinal homeostasis and inflammation. Here, we report that expression of the autophagy related 5 (*Atg5*) protein in myeloid cells prevents dysbiosis and excessive intestinal inflammation by limiting IL-12 production. Mice with a selective genetic deletion of *Atg5* in myeloid cells (*Atg5 Δ Mye*) showed signs of dysbiosis prior to colitis and exhibited severe intestinal inflammation upon colitis induction that was characterized by increased IFN γ production. This increase in IFN γ was due to excess IL-12 secretion from *Atg5*-deficient myeloid cells. *Atg5* functions to limit IL-12 secretion through modulation of late endosome (LE) acidity. Additionally, the autophagy cargo receptor NBR1, which accumulates in *Atg5*-deficient cells, played a role by delivering IL-12 to LE. Restoration of the intestinal microbiota and alleviation of intestinal inflammation was achieved by genetic deletion of IL-12 in *Atg5 Δ Mye* mice. In summary, *Atg5* expression in intestinal myeloid cells acts as an anti-inflammatory brake to regulate IL-12 thus preventing dysbiosis and uncontrolled IFN γ -driven intestinal inflammation.

INTRODUCTION

Immune system dysregulation, intestinal barrier defects and dysbiosis are believed to be driven in part by genetics according to recent genome-wide association studies (GWAS) (Schirmer, Garner et al., 2019). Many of the identified genes are involved in innate cell bacterial recognition and processing and appear to contribute to the pathogenesis observed in inflammatory bowel disease (IBD) such as *NOD2* and *ATG16L1* which are linked to autophagy (Schirmer et al., 2019). Autophagy is one such process associated with IBD susceptibility (Brest, Lapaquette et al., 2011, Hampe, Franke et al., 2007, Henckaerts, Cleynen et al., 2011, Lahiri, Hedl et al., 2015, Lassen & Xavier, 2018, Levine & Kroemer, 2019, McCarroll, Huett et al., 2008, Rioux, Xavier et al., 2007, Roberts, Gearry et al., 2007, Yamazaki, Onouchi et al., 2007). Autophagy is a conserved catabolic process that degrades protein aggregates, damaged organelles and numerous pathogens (Galluzzi, Baehrecke et al., 2017). Autophagy has proven critical for intestinal homeostasis. Defects in the autophagic pathway, specifically in intestinal epithelial cell lineages, results in increased intestinal permeability and Paneth and Goblet cell (GC) dysfunction (Cadwell, Liu et al., 2008, Cadwell, Patel et al., 2009, Matsuzawa-Ishimoto, Shono et al., 2017, Nighot, Hu et al., 2015, Patel, Miyoshi et al., 2013, Wong, Ganapathy et al., 2019). More recently, an IBD risk loci associated with autophagy was found to disrupt the microbiota

albeit it is unclear what cell-type mediates this effect (Lavoie, Conway et al., 2019).

Nevertheless, there is mounting evidence that autophagic genes are critical for intestinal homeostasis and defects in the autophagic process can lead to increased susceptibility to intestinal pathogens and overall enhanced IBD susceptibility (Bel, Pendse et al., 2017, Benjamin, Sumpter et al., 2013, Burger, Araujo et al., 2018, Lassen & Xavier, 2018).

Many of these IBD risk genes and pathways including autophagy are highly relevant to myeloid cell function in addition to that of epithelial cells (Chauhan, Mandell et al., 2015, Homer, Richmond et al., 2010, Singh, Davis et al., 2006, Zhang, Zheng et al., 2017). However, the specific role for autophagy and autophagy genes in myeloid cells in maintaining the balance between intestinal homeostasis and inflammation has yet to be fully explored. The prevailing hypothesis linking autophagy to IBD is through the IL-17 signaling pathway. IL-1, IL-17 and IL-23, all involved in IL-17-mediated inflammation, are upregulated in IBD (Angelidou, Chrysanthopoulou et al., 2018, Coccia, Harrison et al., 2012, Dragasevic, Stankovic et al., 2018, Jiang, Su et al., 2014, Kanai, Mikami et al., 2012, Liu, Yadav et al., 2011, Mao, Kitani et al., 2018, Menghini, Corridoni et al., 2019, Moschen, Tilg et al., 2019, Shouval, Biswas et al., 2016). Our previous work as well as that of others has demonstrated that autophagy regulates the production of the proinflammatory cytokines IL-1 α , IL-1 β and IL-23, primarily in infection models (Castillo, Dekonenko et al., 2012, Dupont, Jiang et al., 2011, Peral de Castro, Jones et al., 2012, Reed, Morris et al., 2015, Watson, Manzanillo et al., 2012, Zhang, Kenny et al., 2015). This observation is the canonical pathway frequently described linking autophagy dysregulation to excess inflammation via IL-1 and IL-17. IL-1 and IL-23 are key regulators of IL-17-mediated inflammation (Chung, Chang et al., 2009, McGeachy, Chen et al., 2009). However, the role of IL-17 in IBD pathogenesis has recently been questioned as therapeutic targeting of IL-17 exacerbates inflammation in both IBD patients and animal models of IBD (Fujino, Andoh et al., 2003, Hueber, Sands et al., 2012, Ogawa, Andoh et al., 2004, Targan, Feagan et al., 2016, Yang, Chang et al., 2008), suggesting that IL-17 may not be the predominant driver of autophagy-linked IBD pathogenesis. Thus, there is a clear gap in knowledge regarding which inflammatory pathway might underlie IBD pathology with respect to autophagy dysregulation.

This study assessed the role of the autophagy gene *Atg5* in myeloid cells in maintaining the balance between intestinal homeostasis and inflammation. *Atg5*'s most well-understood actions are via the Atg5-Atg12-Atg16L1 complex, which acts as the E3 enzyme conjugating PE to LC3, and along with the E1-like actions of Atg7 and the E2-like actions of Atg3, this pathway drives isolation membrane formation and eventual autophagosome maturation (Hanada, Noda

et al., 2007, Mizushima, Noda et al., 1998, Noda, Fujioka et al., 2013, Sakoh-Nakatogawa, Matoba et al., 2013). Atg5 is also embedded in autophagosomal membranes which allows it to interact with fusion proteins in lysosomal membranes such as Tectonic β -propeller repeat containing 1 (TECRP) that facilitate autophagosome-lysosome fusion (Chen, Fan et al., 2012, Ye, Zhou et al., 2018). Thus, Atg5 plays a major role in selective and bulk autophagy which are critical for cell-autonomous immunity. However, it is unclear how Atg5 expression, specifically in myeloid cells, functions outside of bacterial recognition and processing. Here, we show at steady state that mice with an *Atg5*-deficiency in myeloid cells (herein called *Atg5* Δ Mye mice) (Castillo et al., 2012, Zhao, Fux et al., 2008) show alterations in the gut microbiota as well as mucosal TH1 (IFN γ) skewing. *Atg5* Δ Mye mice were susceptible to chemically induced colitis that was characterized by an enhanced IFN γ response. Both TH1 skewing and microbiota changes were partly driven through IL-12 dysregulation in *Atg5*-deficient myeloid cells. Confirming our findings, *Atg5* Δ Mye mice crossed to *IL12p35*-deficient mice resulted in restoration of the gut microbiota and protection from dextran sodium sulfate (DSS)-induced colitis through the reduction of the IL-12 pathway. We further show IL-12 regulation was not mediated through autophagy but interactions with sequestosome-1-like receptor NBR1 and Atg5 action on late endosomes (LE). There has not been a single description of autophagic proteins regulating IL-12-driven immune responses. Thus, these data indicate a new autophagy-independent role for Atg5 and NBR1 in myeloid cells in influencing intestinal homeostasis through an IL-12 pathway.

RESULTS

Altered colonic microbiota in mice with an *Atg5*-deficiency in myeloid cells.

The gut microbiota is influenced by environmental factors, immune responses, and genetics which is highlighted in individuals with IBD. Numerous studies have reported a decrease in bacterial diversity in IBD patients with major alterations in the phyla Firmicutes and Bacteroidetes (Imhann, Vich Vila et al., 2018, Kang, Alvarado et al., 2020, Manichanh, Borruel et al., 2012, Manichanh, Rigottier-Gois et al., 2006, Ni, Wu et al., 2017, Willing, Dicksved et al., 2010). There is substantial evidence that intestinal myeloid cells are regulated by commensal microbiota (Bain, Bravo-Blas et al., 2014, Chang, Hao et al., 2014, Kang et al., 2020, Mortha, Chudnovskiy et al., 2014, Singh, Gurav et al., 2014); however, numerous studies have emerged demonstrating myeloid regulation of the microbiota (Bader, Enos et al., 2018, Niess, Brand et al., 2005, Rescigno, Urbano et al., 2001, Vallon-Eberhard, Landsman et al., 2006). Autophagy genes appear to be involved in the latter as it was shown the autophagy protein, Atg16L1,

expressed in myeloid cells from both humans and mice altered IgA-coated intestinal bacteria at steady-state and during inflammation (Zhang et al., 2017); however, it was not confirmed if this increase in IgA-coated bacteria affects the overall microbial community. More recently, another mouse model with a global knockin of the IBD risk allele *ATG16L1 T300A* had an altered gut microbiota prior to colitis induction (Lavoie et al., 2019).

Utilizing *Atg5* Δ Mye mice (myeloid specific loss of autophagy related 5, *Atg5*, gene), we investigated the importance of *Atg5* expression in myeloid cells in maintaining the intestinal microbiota (Castillo et al., 2012, Zhao et al., 2008). In comparison to littermate controls (*Atg5*-Wild-type, *Atg5*-Wt), *Atg5* Δ Mye mice showed significant differences in the intestinal microbial composition (Fig. 1 A). Similar to what is observed in IBD patients (Manichanh et al., 2012, Manichanh et al., 2006, Ni et al., 2017, Seksik, 2010, Willing et al., 2010), *Atg5* Δ Mye mice had a decrease in the phylum Firmicutes (Fig. 1 B) and an increase in the phylum Bacteroidetes (Fig. 1 C) compared to *Atg5*-Wt mice. No significant changes were observed for phyla Actinobacteria and Proteobacteria (Fig. 1, D and E). Additionally, the principal coordinate analysis (PCoA) plot and heatmap of bacterial communities revealed tight clustering of *Atg5*-Wt microbiota that was distinct from *Atg5* Δ Mye microbiota at the genus level (Fig. 1, F and G) and this was also observed for female mice (Supp. Fig. 1 A and B). Thus, *Atg5* expression in myeloid cells is critical for the maintenance of the intestinal microbiota.

Myeloid *Atg5* expression regulates IFN γ response in the intestinal microenvironment.

The proinflammatory cytokine IL-17 contributes to shaping and regulating the intestinal microbiota. We and others have shown T cells from mice wherein myeloid cells lack *Atg5* or other autophagy-related genes display skewing towards IL-17 polarization (Castillo et al., 2012, Peral de Castro et al., 2012, Reed et al., 2015, Watson et al., 2012). This T cell polarization is mediated through the dysregulation of cytokines or cellular components that promote IL-17 responses (Chung et al., 2009, McGeachy et al., 2009). Microbiota differences can also alter T cell polarization (Atarashi, Tanoue et al., 2011, Chen, Sun et al., 2019, Ivanov, Atarashi et al., 2009, Ivanov, Frutos Rde et al., 2008, Sun, Wu et al., 2018). To determine whether an enhanced level of IL-17-producing T cells populating the intestinal mucosa is influencing the microbiota of *Atg5* Δ Mye mice, we isolated and stimulated CD4⁺ T cells from the mesenteric lymph nodes (mLN) to determine TH polarization. Interestingly, we observed an increase in IFN γ -producing CD4⁺ T cells from *Atg5* Δ Mye mice (Supp. Fig. 1 C,D) with no significant

difference in the number of IL-17-producing CD4⁺ T cells (Supp. Fig. 1 E). Thus, at steady state, Atg5ΔMye mice show differences in CD4⁺ T cell polarization.

To examine the impact of TH1 skewing and microbiota alteration in Atg5ΔMye mice during an inflammatory state, we subjected both Atg5ΔMye and Atg5-Wt mice to DSS-induced acute colitis. Similar to what was reported in mice with an *Atg16l1*- or *Atg7*-deficiency specifically in myeloid cells (Lee, Kim et al., 2016a, Zhang et al., 2017), colitis induction in Atg5ΔMye mice resulted in an exacerbated inflammatory response compared to Atg5-Wt mice. Intestinal epithelial disruption via DSS caused significant weight loss (Fig. 2 A) as well as acute diarrheal response and mucosal bleeding (disease activity index, DAI, Fig. 2 B) in Atg5ΔMye mice. The length of the colon was measured to assess colonic mucosal injury and revealed colonic shortening and severe inflammation from the cecum to the distal colon of Atg5ΔMye mice (Fig. 2 C and Supp. Fig. 1, F and G). Additionally, we found that the spleen weight of colitic Atg5ΔMye mice was increased (Fig. 2 D). These results indicate Atg5 expression in myeloid cells, like Atg7 and Atg16L1, is critical to control acute intestinal inflammation.

The acute DSS model is an innate/wound repair model and although the acute DSS model is not solely dependent on B and T cell responses, both participate in the exaggerated presentation of the disease. In fact, T cells have been shown to be the major driver of colonic inflammation in the acute DSS model by day 4 with a peak at day 8 (Nunes, Kim et al., 2018). Interestingly, we do not see major changes in DAI in Atg5ΔMye mice until day 6 and 7 (Fig. 2B) when we would expect T cells to be responding. Furthermore, IFN γ has been shown to play an indispensable role in the initiation of acute DSS colitis and IFN γ -deficient mice are protected from acute DSS induced colitis (Ito, Shin-Ya et al., 2006). Thus, this model is sufficient to show IFN γ responses from T cells in Atg5ΔMye mice. Our observation that CD4⁺ T cells are skewed towards a type 1 immune response in Atg5ΔMye mice at steady state prompted an examination of cytokine expression and TH skewing during colitis. Colitic colons from Atg5ΔMye mice showed increased IFN γ gene and protein expression compared to controls (Fig. 2, E and G). Similar to steady state, there was also an increase in IFN γ -producing CD4⁺ T cells isolated from the mLN of colitic Atg5ΔMye mice (Fig. 2 I). While IL-17 was expressed in both Atg5ΔMye and Atg5-Wt mice, we found no significant difference in IL-17 expression in the colon (Fig. 2, F and H) or from mLN CD4⁺ T cells (Fig. 2 J). This discrepancy in IL-17 expression could be tissue-specific as models in which myeloid cells lack autophagic components and express high levels of IL-17 have been reported for respiratory infections (Castillo et al., 2012, Reed et al., 2015, Watson et al., 2012). However, others have reported TH1 skewing in ATG16L1T300A knockin mice as well as in conditional myeloid Atg7-deficient mice during intestinal inflammation (Lavoie

et al., 2019, Zhang et al., 2017). Thus, our data suggest *Atg5* expression in myeloid cells protects the intestinal microenvironment during inflammation through limiting IFN γ expression.

***Atg5* limits IL-12 production in myeloid cells independent of canonical autophagy.**

The current paradigm is that an autophagic defect (or loss of an autophagy-related gene) in myeloid cells leads to enhanced IL-1 α/β expression (Castillo et al., 2012, Dupont et al., 2011, Peral de Castro et al., 2012). The excess production of IL-1 skews lymphocytes to produce IL-17, subsequently promoting IL-17-mediated inflammation. However, we found colitic *Atg5* Δ Mye mice presented with increased IFN γ and no significant difference in IL-17 in the colon microenvironment. This suggests colonic *Atg5*-deficient myeloid cells are likely to produce other factors that promote TH1 skewing and IFN γ production. IL-12p70 (hereafter called IL-12) is a major cytokine involved in TH1 polarization and is a potent inducer of IFN γ (Eftychi, Schwarzer et al., 2019, Moschen et al., 2019, Perussia, Chan et al., 1992, Trinchieri, Wysocka et al., 1992). IL-12 is a heterodimeric protein comprised of IL-12p35 and IL-12p40 (IL-12p40 also dimerizes with IL-23p19 to form IL-23, a major cytokine maintaining TH17 cells) and is highly expressed by various myeloid cells including macrophages (Abdi & Singh, 2015, Carra, Gerosa et al., 2000, D'Andrea, Rengaraju et al., 1992, Moschen et al., 2019, Reitberger, Haimerl et al., 2017). We have previously shown *Atg5* Δ Mye mice also produce excess IL-12 during tuberculosis infection (Castillo et al., 2012). Utilizing bone marrow-derived macrophages (BMM) from *Atg5* Δ Mye and *Atg5*-Wt mice, we showed that *Atg5*-deficient macrophages secrete excess IL-12 upon LPS/IFN- γ stimulation compared to *Atg5*-Wt macrophages (Fig. 3A). For many cell types, low levels of IL-12p35 are constitutively expressed, while IL-12p40 expression occurs primarily in macrophages and dendritic cells, and both increase in response to microbial stimulation (Abdi & Singh, 2015, Carra et al., 2000, D'Andrea et al., 1992, Reitberger et al., 2017). We observed no difference in *Il12a* (IL-12p35) and *Il12b* (IL-12p40) gene expression (Supp. Fig. 2, A and B). Furthermore, there was no difference in cell surface expression of IFN γ -receptor, TLR4/MD2 or CD14 between *Atg5*-Wt and *Atg5*-deficient BMM (Supp. Fig. 2 C) that would enable an enhanced response. This data suggests *Atg5* has a post-transcriptional role in regulating IL-12 secretion.

As mentioned above, *Atg5* assists in autophagosome formation and autophagosome-lysosome fusion. Thus, the genetic deletion of *Atg5* would consequently affect several steps in autophagy. We next considered whether IL-12 was a direct target for autophagic removal. However, endogenous IL-12 and LC3 did not colocalize during stimulation and treatment with bafilomycin A1 (Baf. A1) (Supp. Fig. 2 D) in *Atg5*-Wt or *Atg5*-deficient BMM suggesting IL-12 is

unlikely to be a direct target for autophagic degradation. Baf. A1 is widely used to inhibit autophagic flux by targeting the V-ATPase ATP6V0C/V0 subunit c but can also deacidify endosome/lysosome vesicles through the same mechanism (Harada, Shakado et al., 1997, Yoshimori, Yamamoto et al., 1991). Atg5 also regulates acidification and deacidification of late endosomal compartments (Guo, Chitiprolu et al., 2017) and prior to stimulation *Atg5*-deficient BMM display increased lysotracker staining compared to *Atg5*-Wt BMM (Supp Fig. 2E) suggesting the loss of Atg5 affects the regulation of vesicles' pH levels (Guo et al., 2017, Peng, Zhang et al., 2014). Interestingly, Baf. A1 treatment decreased IL-12 secretion in *Atg5*-Wt BMM (Fig. 3 B). Furthermore, treatment of *Atg5*-deficient BMM with Baf. A1 also reduced IL-12 secretion (Fig. 3 C). These data suggest Atg5 regulates IL-12 secretion through vesicle acidification and Baf. A1 can compensate for the loss of Atg5 in regulating IL-12 secretion.

We next examined the intracellular localization of IL-12 after Baf. A1 treatment. As mentioned above, IL-12 consists of subunits IL-12p35 and IL-12p40. While the IL-12p40 subunit has conventional secretory sequences and can be secreted in its homodimeric form, the IL-12p35 peptide is leaderless, cannot be secreted as a monomer, and conventional secretion cannot be induced by the addition of a secretory sequence (Abdi & Singh, 2015, Carra et al., 2000, Reitberger et al., 2017). The purpose of IL-12p35's constitutive expression remains poorly understood (Abdi & Singh, 2015) but a recent report demonstrated IL-12p35 is trafficked to late endosomes (LE) before secretion (Chiaruttini, Piperno et al., 2016). As observed with total IL-12, Baf. A1 treatment resulted in IL-12p35 accumulation in LE as determined by IL-12p35 (red) and Lamp1 (green) colocalization (Fig. 3 D). White arrows in images indicate puncta co-localizing with insets displaying enlargement of indicated region. Interestingly, IL-12p35 (green) colocalized with Atg5 (red) (Fig. 3 E). IL-12 (red) colocalization with Lamp1 (green) was more evident in *Atg5*-deficient BMM (Fig. 3 F). Further analysis, revealed IL-12 (red) also colocalized with the LE marker Rab7 (green) (Supp. Fig. 2F) as previously reported (Chiaruttini et al., 2016). Taken together, our data suggest Lamp1⁺ LEs (Cheng, Xie et al., 2018, Dunster, Toh et al., 2002) contain IL-12 and the lack of Atg5 leads to increased levels of IL-12 in these vesicles.

The sequestosome-1-like receptor NBR1 is involved in IL-12 secretion.

While most cytokines are directed by their signal sequence through endoplasmic reticulum (ER)-golgi complex pathway for processing and trafficking, some inflammatory cytokines, including IL-1 β and IL-18, are known to be excreted via alternative strategies (Abdel Fattah, Bhattacharya et al., 2015, Claude-Taupin, Bissa et al., 2018, Dupont et al., 2011, Kimura, Jia et al., 2017, Murai, Okazaki et al., 2015). For IL-1 β , the selective autophagy cargo receptor TRIM16 directs IL-1 β to LC3-II⁺ sequestration membranes for secretion (Kimura et al., 2017).

This suggests other selective cargo receptors such as the sequestosome-1-like receptors (SLRs: p62/SQSTM1 and NBR1) could act as possible cargo receptors for alternative secretion (Claude-Taupin, Jia et al., 2017, Jiang, Dupont et al., 2013). We considered NBR1 might be responsible for delivering IL-12 to LE for secretion given a homolog of the mammalian NBR1, Nbr1 from *Schizosaccharomyces pombe* was shown to deliver proteins to LE (Mardakheh, Yekezare et al., 2009) and NBR1 accumulates upon autophagy inhibition (Kirkin, Lamark et al., 2009). Indeed, confocal imaging revealed NBR1 (red) co-localized with IL-12p35 (green) in WT BMM after LPS/IFN- γ and Baf. A1 treatment (Fig. 4 A, top row and B); white arrows indicate puncta co-localizing. However, this co-localization was reduced upon brefeldin A (BrefA) treatment, suggesting NBR1 (red) and IL-12 (green) interaction occurs after IL-12 leaves the ER (Fig. 4 A, bottom row and B). This reduction in colocalization was not due to enhanced secretion as IL-12 was undetectable by ELISA after LPS/IFN γ /brefA treatment (data not shown). To verify NBR1/IL-12 interaction in macrophages, cell lysates were immunoprecipitated with anti-IL12p35, anti-NBR1 or isotype control antibodies and subjected to western blot with anti-IL-12p35, anti-IL-12p40, anti-NBR1 and anti-p62 antibodies. Both IL-12p40 and NBR1 co-precipitated with anti-IL-12p35 antibodies (Fig. 4 C). Reciprocally, IL-12p35 and IL-12p40 co-precipitated with anti-NBR1 antibodies (Fig. 4 C); however, neither probe co-precipitated p62/SQSTM1 (Fig. 4C). This interaction was further verified *in silico* through ClusPro docking (Kozakov, Hall et al., 2017) as IL-12 (1F45) interacted with multiple peptide fragments (1WJ6, 2L8J, 2MGW, and 2MJ5) of NBR1 (Fig. 4 D and E). Lastly, a knock down of NBR1 via small interfering RNA (siRNA) in Atg5-Wt BMM reduced IL-12 secretion compared to scrambled siRNA control (Fig. 4 F, Supp. Fig. 2 G). A reduction in IL-12 secretion was also observed in Atg5-deficient BMM after NBR1 knockdown suggesting this interaction is independent of Atg5 (Fig. 4 G, Supp. Fig. 2 G). Taken together, our data suggest NBR1 is partially responsible for IL-12 secretion whereby it functions to direct IL-12p35 to LE for secretion.

Limiting IL-12 secretion restores the gut microbiota and protects against DSS-induced colitis in mice with an Atg5-deficiency in myeloid cells.

To determine if the change in the gut microbiota and susceptibility to DSS-induced colitis in Atg5 Δ Mye mice was due to the excess IL-12 secretion, we crossed Atg5 Δ Mye mice to IL-12p35 deficient mice (Mattner, Magram et al., 1996) to generate Atg5 Δ Mye-IL12KO mice. We first examined the colonic microbiota at steady state in all three groups of mice (i.e. Atg5-Wt, Atg5 Δ Mye and Atg5 Δ Mye-IL12KO). Atg5 Δ Mye-IL12KO mice showed a global change in the intestinal microbial composition in comparison to Atg5 Δ Mye mice, particularly an increase in the

phyla Firmicutes (Supp. Fig. 3 A). Furthermore, the PCoA plots and heatmap show clustering of the bacterial communities (genus level) of the Atg5ΔMye-IL12KO and Atg5-Wt colonic mice that was separated from the bacterial community in the Atg5ΔMye mice (Fig. 5, A and B).

We next examined Atg5ΔMye-IL12KO mice response to DSS-induced colitis. Atg5ΔMye-IL12KO mice were protected from DSS-induced colitis compared to Atg5ΔMye mice. Atg5ΔMye-IL12KO mice suffered minimal weight loss, showed no shortening of the colon after colitis induction, and had reduced spleen weights (Fig. 5, C-E). There was also a decrease in *Ifng* gene expression in the colon after colitis induction in Atg5ΔMye-IL12KO mice but no change in *Il17a* gene expression between both groups of mice. (Supp. Fig. 3, B and C). Additionally, we observed a decrease in the percent of IFN γ -producing CD4⁺ T cells from the mLN of Atg5ΔMye-IL12KO mice compared to Atg5ΔMye mice (Fig. 5 F). However, Atg5ΔMye-IL12KO mLN CD4⁺ T cells did produce more IL-17 (Fig. 5 G). This change in IL-17 expression is likely due to Atg5-deficient myeloid cells producing excess IL-1 (Castillo et al., 2012). Nevertheless, genetic deletion of IL-12p35 from mice in which myeloid cells also lack Atg5 restores the microbiota and protects against DSS-induced colitis.

DISCUSSION

This work identifies a role for Atg5 in myeloid cells in unconventional cytokine secretion that consequently affects intestinal homeostasis. Additionally, it adds to the plethora of functions described for Atg5 (Guo et al., 2017, Inomata, Into et al., 2013, Lee, Mattei et al., 2010, Ndoeye, Budina-Kolomets et al., 2017, Peng et al., 2014, Simon, Yousefi et al., 2014, Yousefi, Perozzo et al., 2006). Along with these studies, this work establishes a new non-autophagy, immunological role for Atg5 in promoting TH1 responses. In the context of IBD and autophagy-related proteins (Atg), GWAS has only identified *ATG16L1* variants (Hampe et al., 2007). Nevertheless, animal models with an Atg5-deficiency display a similar phenotype to Atg16L1-deficient mice (Cadwell et al., 2009). Interestingly, the levels of *ATG5* and the function of autophagy (and possibly other ATG5 functions) are decreased in IBD patient samples. This decrease in *ATG5* expression and autophagy activity in IBD patients is linked to an increased expression of the microRNA miR30C that acts to downregulate *ATG5* expression (Nguyen, Dalmaso et al., 2014, Ye et al., 2018). Therefore, inhibiting Atg5 expression and function, described here in only myeloid cells, ultimately alters the intestinal microenvironment.

As previously reported, Atg5-deficient myeloid cells produce excess proinflammatory cytokines including IL-1 α and IL-1 β (Castillo et al., 2012, Dupont et al., 2011). Thus, we cannot dismiss that either cytokine contributes to the excess IFN γ observed in Atg5cKO mice as IL-1

can synergize with IL-12 to enhance IFN γ production and TH1 skewing (Cooper, Fehniger et al., 2001, Tominaga, Yoshimoto et al., 2000). Although we and others did not find any changes in colonic IL-17-producing cells in mice with a genetic deletion of Atg5 or Atg7 in myeloid cells (Castillo et al., 2012, Lee et al., 2016a), both colonic TH17 and TH1 cells were found to be increased in *ATG16L1* T300A knockin mice (Lavoie et al., 2019). Given Atg5 Δ Mye-IL12KO mice were protected against colitis induction, these results together suggest IL-12 is a key cytokine driving intestinal pathology and TH1 skewing in our model.

This similarity in intestinal TH1 responses between these mouse models is likely linked to the functional role of Atg5, Atg7 and Atg16L1. All three proteins are involved in some aspect of the autophagy process as well as endosome and lysosome regulation (Fraser, Simpson et al., 2019, Galluzzi et al., 2017, Mardakheh et al., 2009). The role of the TH17 response in driving intestinal pathology is up for debate as there is strong evidence suggesting IL-17 may have a protective role in the intestine. Exacerbated intestinal inflammation was reported in *IL17a*-deficient mice or after *in vivo* blockade of IL-17 during DSS-induced colitis (Ogawa et al., 2004, Yang et al., 2008). Furthermore, in clinical trials for IBD patients, targeting IL-17 or IL-17R worsened symptoms leading to early clinical trial termination (Hueber et al., 2012, Targan et al., 2016). Nevertheless, this does not rule out the effects of increased levels of IL-17 during intestinal inflammation (Fujino et al., 2003, Jiang et al., 2014, Moschen et al., 2019). In fact, IL-17 plays a major role in regulating mucosal IgA production (Dann, Mantey et al., 2015, Kumar, Monin et al., 2016). Interestingly, increased IgA-coated bacteria were found in mice that lack Atg16L1 in myeloid cells as well as in the stool of Crohn's disease patients who were homozygous for the *ATG16L1* T300A risk allele (Zhang et al., 2017). *ATG16L1* T300A knockin mice also showed alteration in the microbiota. Similar to our findings, *ATG16L1* T300A knockin mice also had a decrease in the phyla Firmicutes and an increase in the phyla Bacteroidetes (Lavoie et al., 2019). It is unclear if these microbiota changes in *ATG16L1* T300A knockin mice are due specifically to myeloid cells, since all cells including the intestinal epithelium express the *ATG16L1* T300A gene in this mouse model. Additionally, deletions or polymorphisms of autophagy genes in myeloid cells alter bacterial clearance suggesting there are multiple mechanisms by which autophagy genes regulate host-microbiota interactions. Our data suggest myeloid cells likely have a strong contribution in maintaining the microbiota through Atg5's action on IL-12 secretion.

The autophagic process and the genes that regulate autophagy are crucial for intestinal homeostasis. Autophagy is required for the maintenance of tight junction integrity, gut-commensal homeostasis, and control of invasive bacteria at the intestinal epithelium (Bel et al.,

2017, Benjamin et al., 2013, Burger et al., 2018, Cadwell et al., 2008, Cadwell et al., 2009, Lee et al., 2016a, Matsuzawa-Ishimoto et al., 2017, Nighot et al., 2015, Patel et al., 2013, Wong et al., 2019, Wu, Wang et al., 2017). Multiple lines of evidence also suggest autophagy regulates inflammatory cytokines (Castillo et al., 2012, Dupont et al., 2011, Lee, Foote et al., 2016b, Merkley, Chock et al., 2018, Saitoh, Fujita et al., 2008). These attributes make modulation of autophagy or single autophagic proteins an excellent therapeutic target. Nevertheless, it is critical to understand the function of these individual proteins given they can modulate different TH responses. In fact, reagents that are known to target autophagy activation such as rapamycin, increase IL-12 secretion (Macedo, Turnquist et al., 2013, Ohtani, Nagai et al., 2008). Additionally, chloroquine, a known inhibitor of autophagosome-lysosome fusion, reduces IL-12 secretion (Said, Bock et al., 2014), likely through a similar mechanism as Baf. A1. These reagents have been used with some clinical efficacy in IBD patients and animal models of IBD (Goenka, Kochhar et al., 1996, Kanvinde, Chhonker et al., 2018, Nagar, Ranade et al., 2014, Park, Jang et al., 2019). It is unclear if these effects were due to IL-12 regulation but targeting IL-12, specifically, the IL-12p40 subunit that is shared with IL-23, has also proven beneficial for some IBD patients (Feagan, Sandborn et al., 2016, Moschen et al., 2019, Sands, Sandborn et al., 2019). Therefore, understanding how autophagy or single autophagic proteins regulate proinflammatory cytokines will allow precision in modulating the immune system in chronic inflammatory conditions like IBD. Lastly, targeting these autophagy proteins or their functions could be an alternative to mitigate problems associated with biologics including loss of function, immunogenicity and cost (D'Haens, 2007).

In conclusion, our data support a novel role for Atg5 expression in myeloid cells in regulating intestinal homeostasis. Atg5 expression in myeloid cells controls the IL-12-IFN γ pathway that influences the microbiota and limits this pathway during colitis (Supp. Fig. 3 D). Mechanistically, we show both Atg5 and the cargo receptor NBR1 regulate IL-12 secretion in myeloid cells. Specifically, we propose that NBR1 shuttles IL-12 to LE whereby Atg5 functions to control the pH of these IL-12 containing vesicles for secretion. A genetic deletion of Atg5 results in dysregulation of IL-12 secretion as well as the accumulation of SLR's such as NBR1 (Kirkin et al., 2009) which could potentially allow for increased accumulation of IL-12 in LE. Consequently, these attributes increase IL-12 secretion. A reduction in IL-12 secretion could be achieved by removing NBR1 or reducing LE/lysosomal pH with Baf. A1 even in the absence of Atg5 expression. Nevertheless, the dysregulation of IL-12 in Atg5 Δ Mye mice leads to alterations of the microbiota and severe colitis when the intestinal barrier is disrupted.

MATERIALS AND METHODS

Animals. The transgenic Atg5ΔMye mice (myeloid specific Atg5 deletion) and Atg5-Wt mice have previously been characterized (Zhao et al., 2008). B6.129S1-*Il12a*^{tm1jm}/J (IL-12p35-deficient mice) were purchased from JAX (002692) (Mattner et al., 1996) and crossed to Atg5ΔMye mice to generate Atg5ΔMye-IL12KO mice. All mice were genotyped for the presence of *Atg5*, *Il12a* or Cre expression by Transnetyx Inc. All experiments were approved by the Institutional Animal Care and Use Committee of the University of New Mexico Health Sciences Center, in accordance with the National Institutes of Health guidelines for use of live animals. The University of New Mexico Health Sciences Center is accredited by the American Association of Accreditation of Laboratory Animal Care.

Intestinal inflammation model. For Dextran Sodium Sulfate (DSS)-induced colitis, mice were provided 2.5% DSS (colitis grade, ~30,000 -50,000 MW; MP Biomedicals) in drinking water *ad libitum*. Reagent used for *in vivo* treatment as well as sample collection and stimulation are included in Table S1.

Microbiota analysis. Fecal samples from Atg5-Wt, Atg5ΔMye, and Atg5ΔMye-IL-12KO mice were collected fresh in sterile tubes and flash frozen. Microbial communities were determined by sequencing of the 16S rRNA as previously reported (Komesu, Richter et al., 2017) with minor modifications described below. Microbial DNA was isolated from feces using the ZymoBIOMICS DNA Miniprep Kit (Zymo Research) following manufacturer's recommendations. Variable regions V-3 through V-4 of the 16S rRNA gene were amplified by PCR using 100 ng input of DNA for each sample in duplicate using primers 319F- (5'-ACTCCTRCGGGAGGCAGCAG-3') and 806R- (5'-GACAGGACTACHVGGGTATCTAATCC-3') containing Nextera adapter overhangs. A second PCR was performed with Nextera® XT Index Kit v2 Set A (Illumina) to complete the adapter and add a unique sample-specific barcodes. After each PCR, a clean-up with AxyPrep Fragment Select-I magnetic beads (Axygen Biosciences) was completed, and all PCR reactions were run on an Applied Biosystems 2720 Thermal Cycler. Indexed samples were combined to yield duplicate 300 ng pools, followed by the creation and denaturation of a 4 nM library, and paired 250bp sequencing runs were completed on the Illumina MiSeq using v3 sequencing chemistry (Illumina). All reagents and kits are listed in Table S2.

Microscopy and image analysis. For confocal microscopy, macrophages were plated at 100k cells per well on 18 mm glass coverslips and stimulated with LPS (500 ng/mL) and IFN γ (10 ng/mL) for 8 hours and treated with BrefA (10 nM) or Baf. A1 (10 nM) for 2 hours. Cells were then fixed with 4% PFA followed by a wash with 1x PBS. Blocking buffer contained PBS with 50% FBS, 2% BSA and 0.1% saponin. After a 1-hour stain in primary antibody, cells were washed with PBS then followed by 1- hour stain in blocking buffer containing secondary antibody. Cells were mounted using ProLong Gold Antifade with DAPI and imaged using the Zeiss LSM 800 Airyscan Confocal microscope with a 63x oil objective lens. Images were processed using Zen Software and Adobe Photoshop (version CC 2019). All primary and secondary antibodies used for confocal staining are listed in Table S3.

Statistical Analysis. Statistical analysis was performed as described in figure legends and graphs generated display mean ($\pm$ SD) and were obtained using Prism software. Microbiome data was sequenced and processed by Illumina's service lab using their inhouse analysis pipeline. Cluster analysis was performed using heatmap3 (Zhao, Guo et al., 2014) package in R. T-test was used to measure specific microbiome species abundance between conditions. Adjust p-value > 0.05 was used as significant threshold. Principle coordinate analysis was conducted in R. Confocal Images Statistical Analysis and Additional Software - Pearson's Correlation Coefficient was acquired from BMM images using Huygens's Deconvolution Scientific Volume Image Software (UNM Fluorescence Microscopy and Cell Imaging shared resource). Quantification figures were also made using Prism, while confocal image figures were constructed using Adobe Illustrator (version CC 2019). All other data were analyzed using one -way ANOVA or two-tailed unpaired Student's t test (Prism).

Online supplemental material. Material and methods for cell and tissue preparation, RT-qPCR, flow cytometry, histology, immunoblotting are found in supplemental material.

Author contributions. SDM and SMG performed all analysis with the help from YG, ZERW, JRB, RRG, AJH, JGI and SBB. SDM, ZERW, YG, KCS and DLD contributed to microbiota sequencing and analysis. SMG, JRB and SBB participated in imaging and analysis. JGI, SBB and VD provided reagents and animals. SDM and SMG participated in writing the manuscript. EFC designed the study, analyzed data and wrote the paper. All authors approved the final version of the manuscript.

Declaration of Competing Interest

The authors declare no conflict of interest.

Acknowledgments

Supported in part by the National Center for Research Resources and the National Center for Advancing Translational Sciences of the National Institutes of Health (NIH) through grant no. UL1TR001449 (E.F.C.) and in part by NIH grant P20GM121176 (E.F.C.), and the Bioinformatics Shared Resource at University of New Mexico Comprehensive Cancer Center with grant P30CA118100 (Y.G.). S.M.G. was supported in part by the Infectious Disease and Inflammation Program pre-doctoral T32 training grant, NIH/NIAID grant T32AI007538.

References

- Abdel Fattah E, Bhattacharya A, Herron A, Safdar Z, Eissa NT (2015) Critical role for IL-18 in spontaneous lung inflammation caused by autophagy deficiency. *J Immunol* 194: 5407-16
- Abdi K, Singh NJ (2015) Making many from few: IL-12p40 as a model for the combinatorial assembly of heterodimeric cytokines. *Cytokine* 76: 53-7
- Angelidou I, Chrysanthopoulou A, Mitsios A, Arelaki S, Arampatzioglou A, Kambas K, Ritis D, Tsironidou V, Moschos I, Dalla V, Stakos D, Kouklakis G, Mitroulis I, Ritis K, Skendros P (2018) REDD1/Autophagy Pathway Is Associated with Neutrophil-Driven IL-1beta Inflammatory Response in Active Ulcerative Colitis. *J Immunol* 200: 3950-3961
- Atarashi K, Tanoue T, Shima T, Imaoka A, Kuwahara T, Momose Y, Cheng G, Yamasaki S, Saito T, Ohba Y, Taniguchi T, Takeda K, Hori S, Ivanov, II, Umesaki Y, Itoh K, Honda K (2011) Induction of colonic regulatory T cells by indigenous Clostridium species. *Science* 331: 337-41
- Bader JE, Enos RT, Velazquez KT, Carson MS, Nagarkatti M, Nagarkatti PS, Chatzistamou I, Davis JM, Carson JA, Robinson CM, Murphy EA (2018) Macrophage depletion using clodronate liposomes decreases tumorigenesis and alters gut microbiota in the AOM/DSS mouse model of colon cancer. *Am J Physiol Gastrointest Liver Physiol* 314: G22-G31
- Bain CC, Bravo-Blas A, Scott CL, Perdiguero EG, Geissmann F, Henri S, Malissen B, Osborne LC, Artis D, Mowat AM (2014) Constant replenishment from circulating monocytes maintains the macrophage pool in the intestine of adult mice. *Nat Immunol* 15: 929-937
- Bel S, Pendse M, Wang Y, Li Y, Ruhn KA, Hassell B, Leal T, Winter SE, Xavier RJ, Hooper LV (2017) Paneth cells secrete lysozyme via secretory autophagy during bacterial infection of the intestine. *Science* 357: 1047-1052
- Benjamin JL, Sumpter R, Jr., Levine B, Hooper LV (2013) Intestinal epithelial autophagy is essential for host defense against invasive bacteria. *Cell Host Microbe* 13: 723-34
- Brest P, Lapaquette P, Souidi M, Lebrigand K, Cesaro A, Vouret-Craviari V, Mari B, Barbry P, Mosnier JF, Hebuterne X, Harel-Bellan A, Mograbi B, Darfeuille-Michaud A, Hofman P (2011) A synonymous variant in IRGM alters a binding site for miR-196 and causes deregulation of IRGM-dependent xenophagy in Crohn's disease. *Nat Genet* 43: 242-5
- Burger E, Araujo A, Lopez-Yglesias A, Rajala MW, Geng L, Levine B, Hooper LV, Burstein E, Yarovinsky F (2018) Loss of Paneth Cell Autophagy Causes Acute Susceptibility to Toxoplasma gondii-Mediated Inflammation. *Cell Host Microbe* 23: 177-190 e4
- Cadwell K, Liu JY, Brown SL, Miyoshi H, Loh J, Lennerz JK, Kishi C, Kc W, Carrero JA, Hunt S, Stone CD, Brunt EM, Xavier RJ, Sleckman BP, Li E, Mizushima N, Stappenbeck TS, Virgin HWt (2008) A key role for autophagy and the autophagy gene Atg16L1 in mouse and human intestinal Paneth cells. *Nature* 456: 259-63
- Cadwell K, Patel KK, Komatsu M, Virgin HWt, Stappenbeck TS (2009) A common role for Atg16L1, Atg5 and Atg7 in small intestinal Paneth cells and Crohn disease. *Autophagy* 5: 250-2
- Carra G, Gerosa F, Trinchieri G (2000) Biosynthesis and posttranslational regulation of human IL-12. *J Immunol* 164: 4752-61
- Castillo EF, Dekonenko A, Arko-Mensah J, Mandell MA, Dupont N, Jiang S, Delgado-Vargas M, Timmins GS, Bhattacharya D, Yang H, Hutt J, Lyons CR, Dobos KM, Deretic V (2012) Autophagy protects against active tuberculosis by suppressing bacterial burden and inflammation. *Proc Natl Acad Sci U S A* 109: E3168-76
- Chang PV, Hao L, Offermanns S, Medzhitov R (2014) The microbial metabolite butyrate regulates intestinal macrophage function via histone deacetylase inhibition. *Proc Natl Acad Sci U S A* 111: 2247-52

- Chauhan S, Mandell MA, Deretic V (2015) IRGM governs the core autophagy machinery to conduct antimicrobial defense. *Mol Cell* 58: 507-21
- Chen D, Fan W, Lu Y, Ding X, Chen S, Zhong Q (2012) A mammalian autophagosome maturation mechanism mediated by TECPR1 and the Atg12-Atg5 conjugate. *Mol Cell* 45: 629-41
- Chen L, Sun M, Wu W, Yang W, Huang X, Xiao Y, Ma C, Xu L, Yao S, Liu Z, Cong Y (2019) Microbiota Metabolite Butyrate Differentially Regulates Th1 and Th17 Cells' Differentiation and Function in Induction of Colitis. *Inflamm Bowel Dis* 25: 1450-1461
- Cheng XT, Xie YX, Zhou B, Huang N, Farfel-Becker T, Sheng ZH (2018) Characterization of LAMP1-labeled nondegradative lysosomal and endocytic compartments in neurons. *J Cell Biol* 217: 3127-3139
- Chiaruttini G, Piperno GM, Jouve M, De Nardi F, Larghi P, Peden AA, Baj G, Muller S, Valitutti S, Galli T, Benvenuti F (2016) The SNARE VAMP7 Regulates Exocytic Trafficking of Interleukin-12 in Dendritic Cells. *Cell Rep* 14: 2624-36
- Chung Y, Chang SH, Martinez GJ, Yang XO, Nurieva R, Kang HS, Ma L, Watowich SS, Jetten AM, Tian Q, Dong C (2009) Critical regulation of early Th17 cell differentiation by interleukin-1 signaling. *Immunity* 30: 576-87
- Claude-Taupin A, Bissa B, Jia J, Gu Y, Deretic V (2018) Role of autophagy in IL-1 β export and release from cells. *Semin Cell Dev Biol* 83: 36-41
- Claude-Taupin A, Jia J, Mudd M, Deretic V (2017) Autophagy's secret life: secretion instead of degradation. *Essays Biochem* 61: 637-647
- Coccia M, Harrison OJ, Schiering C, Asquith MJ, Becher B, Powrie F, Maloy KJ (2012) IL-1 β mediates chronic intestinal inflammation by promoting the accumulation of IL-17A secreting innate lymphoid cells and CD4(+) Th17 cells. *J Exp Med* 209: 1595-609
- Cooper MA, Fehniger TA, Ponnappan A, Mehta V, Wewers MD, Caligiuri MA (2001) Interleukin-1 β costimulates interferon- γ production by human natural killer cells. *Eur J Immunol* 31: 792-801
- D'Andrea A, Rengaraju M, Valiante NM, Chehimi J, Kubin M, Aste M, Chan SH, Kobayashi M, Young D, Nickbarg E, et al. (1992) Production of natural killer cell stimulatory factor (interleukin 12) by peripheral blood mononuclear cells. *J Exp Med* 176: 1387-98
- D'Haens G (2007) Risks and benefits of biologic therapy for inflammatory bowel diseases. *Gut* 56: 725-32
- Dann SM, Manthey CF, Le C, Miyamoto Y, Gima L, Abraham A, Cao AT, Hanson EM, Kolls JK, Raz E, Cong Y, Eckmann L (2015) IL-17A promotes protective IgA responses and expression of other potential effectors against the lumen-dwelling enteric parasite *Giardia*. *Exp Parasitol* 156: 68-78
- Dragasevic S, Stankovic B, Sokic-Milutinovic A, Milosavljevic T, Milovanovic T, Lukic S, Drazilov SS, Klaassen K, Kotur N, Pavlovic S, Popovic D (2018) Importance of TLR9-IL23-IL17 axis in inflammatory bowel disease development: Gene expression profiling study. *Clin Immunol* 197: 86-95
- Dunster K, Toh BH, Sentry JW (2002) Early endosomes, late endosomes, and lysosomes display distinct partitioning strategies of inheritance with similarities to Golgi-derived membranes. *Eur J Cell Biol* 81: 117-24
- Dupont N, Jiang S, Pilli M, Ornatowski W, Bhattacharya D, Deretic V (2011) Autophagy-based unconventional secretory pathway for extracellular delivery of IL-1 β . *EMBO J* 30: 4701-11
- Eftychi C, Schwarzer R, Vlantis K, Wachsmuth L, Basic M, Wagle P, Neurath MF, Becker C, Bleich A, Pasparakis M (2019) Temporally Distinct Functions of the Cytokines IL-12 and IL-23 Drive Chronic Colon Inflammation in Response to Intestinal Barrier Impairment. *Immunity* 51: 367-380 e4
- Feagan BG, Sandborn WJ, Gasink C, Jacobstein D, Lang Y, Friedman JR, Blank MA, Johanns J, Gao LL, Miao Y, Adedokun OJ, Sands BE, Hanauer SB, Vermeire S, Targan S, Ghosh S, de Villiers WJ, Colombel JF, Tulassay Z, Seidler U et al. (2016) Ustekinumab as Induction and Maintenance Therapy for Crohn's Disease. *N Engl J Med* 375: 1946-1960
- Fraser J, Simpson J, Fontana R, Kishi-Itakura C, Ktistakis NT, Gammoh N (2019) Targeting of early endosomes by autophagy facilitates EGFR recycling and signalling. *EMBO Rep* 20: e47734

- Fujino S, Andoh A, Bamba S, Ogawa A, Hata K, Araki Y, Bamba T, Fujiyama Y (2003) Increased expression of interleukin 17 in inflammatory bowel disease. *Gut* 52: 65-70
- Galluzzi L, Baehrecke EH, Ballabio A, Boya P, Bravo-San Pedro JM, Cecconi F, Choi AM, Chu CT, Codogno P, Colombo MI, Cuervo AM, Debnath J, Deretic V, Dikic I, Eskelinen EL, Fimia GM, Fulda S, Gewirtz DA, Green DR, Hansen M et al. (2017) Molecular definitions of autophagy and related processes. *EMBO J* 36: 1811-1836
- Goenka MK, Kochhar R, Tandia B, Mehta SK (1996) Chloroquine for mild to moderately active ulcerative colitis: comparison with sulfasalazine. *Am J Gastroenterol* 91: 917-21
- Guo H, Chitiprolu M, Roncevic L, Javalet C, Hemming FJ, Trung MT, Meng L, Latreille E, Tanese de Souza C, McCulloch D, Baldwin RM, Auer R, Cote J, Russell RC, Sadoul R, Gibbins D (2017) Atg5 Disassociates the V1V0-ATPase to Promote Exosome Production and Tumor Metastasis Independent of Canonical Macroautophagy. *Dev Cell* 43: 716-730 e7
- Hampe J, Franke A, Rosenstiel P, Till A, Teuber M, Huse K, Albrecht M, Mayr G, De La Vega FM, Briggs J, Gunther S, Prescott NJ, Onnie CM, Hasler R, Sipos B, Folsch UR, Lengauer T, Platzer M, Mathew CG, Krawczak M et al. (2007) A genome-wide association scan of nonsynonymous SNPs identifies a susceptibility variant for Crohn disease in ATG16L1. *Nat Genet* 39: 207-11
- Hanada T, Noda NN, Satomi Y, Ichimura Y, Fujioka Y, Takao T, Inagaki F, Ohsumi Y (2007) The Atg12-Atg5 conjugate has a novel E3-like activity for protein lipidation in autophagy. *J Biol Chem* 282: 37298-302
- Harada M, Shakado S, Sakisaka S, Tamaki S, Ohishi M, Sasatomi K, Koga H, Sata M, Tanikawa K (1997) Bafilomycin A1, a specific inhibitor of V-type H⁺-ATPases, inhibits the acidification of endocytic structures and inhibits horseradish peroxidase uptake in isolated rat sinusoidal endothelial cells. *Liver* 17: 244-50
- Henckaerts L, Cleynen I, Brinar M, John JM, Van Steen K, Rutgeerts P, Vermeire S (2011) Genetic variation in the autophagy gene ULK1 and risk of Crohn's disease. *Inflamm Bowel Dis* 17: 1392-7
- Homer CR, Richmond AL, Rebert NA, Achkar JP, McDonald C (2010) ATG16L1 and NOD2 interact in an autophagy-dependent antibacterial pathway implicated in Crohn's disease pathogenesis. *Gastroenterology* 139: 1630-41, 1641 e1-2
- Hueber W, Sands BE, Lewitzky S, Vandemeulebroecke M, Reinisch W, Higgins PD, Wehkamp J, Feagan BG, Yao MD, Karczewski M, Karczewski J, Pezous N, Bek S, Bruin G, Mellgard B, Berger C, Londei M, Bertolino AP, Tougas G, Travis SP et al. (2012) Secukinumab, a human anti-IL-17A monoclonal antibody, for moderate to severe Crohn's disease: unexpected results of a randomised, double-blind placebo-controlled trial. *Gut* 61: 1693-700
- Imhann F, Vich Vila A, Bonder MJ, Fu J, Gevers D, Visschedijk MC, Spekhorst LM, Alberts R, Franke L, van Dullemen HM, Ter Steege RWF, Huttenhower C, Dijkstra G, Xavier RJ, Festen EAM, Wijmenga C, Zhernakova A, Weersma RK (2018) Interplay of host genetics and gut microbiota underlying the onset and clinical presentation of inflammatory bowel disease. *Gut* 67: 108-119
- Inomata M, Into T, Niida S, Murakami Y (2013) Atg5 regulates formation of MyD88 condensed structures and MyD88-dependent signal transduction. *Biochem Biophys Res Commun* 437: 509-14
- Ito R, Shin-Ya M, Kishida T, Urano A, Takada R, Sakagami J, Imanishi J, Kita M, Ueda Y, Iwakura Y, Kataoka K, Okanoue T, Mazda O (2006) Interferon-gamma is causatively involved in experimental inflammatory bowel disease in mice. *Clin Exp Immunol* 146: 330-8
- Ivanov, II, Atarashi K, Manel N, Brodie EL, Shima T, Karaoz U, Wei D, Goldfarb KC, Santee CA, Lynch SV, Tanoue T, Imaoka A, Itoh K, Takeda K, Umesaki Y, Honda K, Littman DR (2009) Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* 139: 485-98
- Ivanov, II, Frutos Rde L, Manel N, Yoshinaga K, Rifkin DB, Sartor RB, Finlay BB, Littman DR (2008) Specific microbiota direct the differentiation of IL-17-producing T-helper cells in the mucosa of the small intestine. *Cell Host Microbe* 4: 337-49

- Jiang S, Dupont N, Castillo EF, Deretic V (2013) Secretory versus degradative autophagy: unconventional secretion of inflammatory mediators. *J Innate Immun* 5: 471-9
- Jiang W, Su J, Zhang X, Cheng X, Zhou J, Shi R, Zhang H (2014) Elevated levels of Th17 cells and Th17-related cytokines are associated with disease activity in patients with inflammatory bowel disease. *Inflamm Res* 63: 943-50
- Kanai T, Mikami Y, Sujino T, Hisamatsu T, Hibi T (2012) RORgammat-dependent IL-17A-producing cells in the pathogenesis of intestinal inflammation. *Mucosal Immunol* 5: 240-7
- Kang B, Alvarado LJ, Kim T, Lehmann ML, Cho H, He J, Li P, Kim BH, Larochelle A, Kelsall BL (2020) Commensal microbiota drive the functional diversification of colon macrophages. *Mucosal Immunol* 13: 216-229
- Kanvinde S, Chhonker YS, Ahmad R, Yu F, Sleightholm R, Tang W, Jaramillo L, Chen Y, Sheinin Y, Li J, Murry DJ, Singh AB, Oupicky D (2018) Pharmacokinetics and efficacy of orally administered polymeric chloroquine as macromolecular drug in the treatment of inflammatory bowel disease. *Acta Biomater* 82: 158-170
- Kimura T, Jia J, Kumar S, Choi SW, Gu Y, Mudd M, Dupont N, Jiang S, Peters R, Farzam F, Jain A, Lidke KA, Adams CM, Johansen T, Deretic V (2017) Dedicated SNAREs and specialized TRIM cargo receptors mediate secretory autophagy. *EMBO J* 36: 42-60
- Kirkin V, Lamark T, Sou YS, Bjorkoy G, Nunn JL, Bruun JA, Shvets E, McEwan DG, Clausen TH, Wild P, Bilusic I, Theurillat JP, Overvatn A, Ishii T, Elazar Z, Komatsu M, Dikic I, Johansen T (2009) A role for NBR1 in autophagosomal degradation of ubiquitinated substrates. *Mol Cell* 33: 505-16
- Komesu YM, Richter HE, Dinwiddie DL, Siddiqui NY, Sung VW, Lukacz ES, Ridgeway B, Arya LA, Zyczynski HM, Rogers RG, Gantz M (2017) Methodology for a vaginal and urinary microbiome study in women with mixed urinary incontinence. *Int Urogynecol J* 28: 711-720
- Kozakov D, Hall DR, Xia B, Porter KA, Padhorny D, Yueh C, Beglov D, Vajda S (2017) The ClusPro web server for protein-protein docking. *Nat Protoc* 12: 255-278
- Kumar P, Monin L, Castillo P, Elsegeiny W, Horne W, Eddens T, Vikram A, Good M, Schoenborn AA, Bibby K, Montelaro RC, Metzger DW, Gulati AS, Kolls JK (2016) Intestinal Interleukin-17 Receptor Signaling Mediates Reciprocal Control of the Gut Microbiota and Autoimmune Inflammation. *Immunity* 44: 659-671
- Lahiri A, Hedl M, Abraham C (2015) MTMR3 risk allele enhances innate receptor-induced signaling and cytokines by decreasing autophagy and increasing caspase-1 activation. *Proc Natl Acad Sci U S A* 112: 10461-6
- Lassen KG, Xavier RJ (2018) Mechanisms and function of autophagy in intestinal disease. *Autophagy* 14: 216-220
- Lavoie S, Conway KL, Lassen KG, Jijon HB, Pan H, Chun E, Michaud M, Lang JK, Gallini Comeau CA, Dreyfuss JM, Glickman JN, Vlamakis H, Ananthakrishnan A, Kostic A, Garrett WS, Xavier RJ (2019) The Crohn's disease polymorphism, ATG16L1 T300A, alters the gut microbiota and enhances the local Th1/Th17 response. *Elife* 8
- Lee HK, Mattei LM, Steinberg BE, Alberts P, Lee YH, Chervonsky A, Mizushima N, Grinstein S, Iwasaki A (2010) In vivo requirement for Atg5 in antigen presentation by dendritic cells. *Immunity* 32: 227-39
- Lee HY, Kim J, Quan W, Lee JC, Kim MS, Kim SH, Bae JW, Hur KY, Lee MS (2016a) Autophagy deficiency in myeloid cells increases susceptibility to obesity-induced diabetes and experimental colitis. *Autophagy* 12: 1390-403
- Lee JP, Foote A, Fan H, Peral de Castro C, Lang T, Jones SA, Gavrilescu N, Mills KH, Leech M, Morand EF, Harris J (2016b) Loss of autophagy enhances MIF/macrophage migration inhibitory factor release by macrophages. *Autophagy* 12: 907-16
- Levine B, Kroemer G (2019) Biological Functions of Autophagy Genes: A Disease Perspective. *Cell* 176: 11-42

- Liu Z, Yadav PK, Xu X, Su J, Chen C, Tang M, Lin H, Yu J, Qian J, Yang PC, Wang X (2011) The increased expression of IL-23 in inflammatory bowel disease promotes intraepithelial and lamina propria lymphocyte inflammatory responses and cytotoxicity. *J Leukoc Biol* 89: 597-606
- Macedo C, Turnquist HR, Castillo-Rama M, Zahorchak AF, Shapiro R, Thomson AW, Metes D (2013) Rapamycin augments human DC IL-12p70 and IL-27 secretion to promote allogeneic Type 1 polarization modulated by NK cells. *Am J Transplant* 13: 2322-33
- Manichanh C, Borruel N, Casellas F, Guarner F (2012) The gut microbiota in IBD. *Nat Rev Gastroenterol Hepatol* 9: 599-608
- Manichanh C, Rigottier-Gois L, Bonnaud E, Gloux K, Pelletier E, Frangeul L, Nalin R, Jarrin C, Chardon P, Marteau P, Roca J, Dore J (2006) Reduced diversity of faecal microbiota in Crohn's disease revealed by a metagenomic approach. *Gut* 55: 205-11
- Mao L, Kitani A, Strober W, Fuss IJ (2018) The Role of NLRP3 and IL-1beta in the Pathogenesis of Inflammatory Bowel Disease. *Front Immunol* 9: 2566
- Mardakheh FK, Yekezare M, Machesky LM, Heath JK (2009) Spred2 interaction with the late endosomal protein NBR1 down-regulates fibroblast growth factor receptor signaling. *J Cell Biol* 187: 265-77
- Matsuzawa-Ishimoto Y, Shono Y, Gomez LE, Hubbard-Lucey VM, Cammer M, Neil J, Dewan MZ, Lieberman SR, Lazrak A, Marinis JM, Beal A, Harris PA, Bertin J, Liu C, Ding Y, van den Brink MRM, Cadwell K (2017) Autophagy protein ATG16L1 prevents necroptosis in the intestinal epithelium. *J Exp Med* 214: 3687-3705
- Mattner F, Magram J, Ferrante J, Launois P, Di Padova K, Behin R, Gately MK, Louis JA, Alber G (1996) Genetically resistant mice lacking interleukin-12 are susceptible to infection with *Leishmania major* and mount a polarized Th2 cell response. *Eur J Immunol* 26: 1553-9
- McCarroll SA, Huett A, Kuballa P, Chlewicki SD, Landry A, Goyette P, Zody MC, Hall JL, Brant SR, Cho JH, Duerr RH, Silverberg MS, Taylor KD, Rioux JD, Altshuler D, Daly MJ, Xavier RJ (2008) Deletion polymorphism upstream of IRGM associated with altered IRGM expression and Crohn's disease. *Nat Genet* 40: 1107-12
- McGeachy MJ, Chen Y, Tato CM, Laurence A, Joyce-Shaikh B, Blumenschein WM, McClanahan TK, O'Shea JJ, Cua DJ (2009) The interleukin 23 receptor is essential for the terminal differentiation of interleukin 17-producing effector T helper cells in vivo. *Nat Immunol* 10: 314-24
- Menghini P, Corridoni D, Butto LF, Osme A, Shivaswamy S, Lam M, Bamias G, Pizarro TT, Rodriguez-Palacios A, Dinarello CA, Cominelli F (2019) Neutralization of IL-1alpha ameliorates Crohn's disease-like ileitis by functional alterations of the gut microbiome. *Proc Natl Acad Sci U S A*
- Merkley SD, Chock CJ, Yang XO, Harris J, Castillo EF (2018) Modulating T Cell Responses via Autophagy: The Intrinsic Influence Controlling the Function of Both Antigen-Presenting Cells and T Cells. *Front Immunol* 9: 2914
- Mizushima N, Noda T, Yoshimori T, Tanaka Y, Ishii T, George MD, Klionsky DJ, Ohsumi M, Ohsumi Y (1998) A protein conjugation system essential for autophagy. *Nature* 395: 395-8
- Mortha A, Chudnovskiy A, Hashimoto D, Bogunovic M, Spencer SP, Belkaid Y, Merad M (2014) Microbiota-dependent crosstalk between macrophages and ILC3 promotes intestinal homeostasis. *Science* 343: 1249288
- Moschen AR, Tilg H, Raine T (2019) IL-12, IL-23 and IL-17 in IBD: immunobiology and therapeutic targeting. *Nat Rev Gastroenterol Hepatol* 16: 185-196
- Murai H, Okazaki S, Hayashi H, Kawakita A, Hosoki K, Yasutomi M, Sur S, Ohshima Y (2015) *Alternaria* extract activates autophagy that induces IL-18 release from airway epithelial cells. *Biochem Biophys Res Commun* 464: 969-974
- Nagar J, Ranade S, Kamath V, Singh S, Karunanithi P, Subramani S, Venkatesh K, Srivastava R, Dudhgaonkar S, Vikramadithyan RK (2014) Therapeutic potential of chloroquine in a murine model of inflammatory bowel disease. *Int Immunopharmacol* 21: 328-35

- Ndoye A, Budina-Kolomets A, Kugel CH, 3rd, Webster MR, Kaur A, Behera R, Rebecca VW, Li L, Brafford PA, Liu Q, Gopal YNV, Davies MA, Mills GB, Xu X, Wu H, Herlyn M, Nicastri MC, Winkler JD, Soengas MS, Amaravadi RK et al. (2017) ATG5 Mediates a Positive Feedback Loop between Wnt Signaling and Autophagy in Melanoma. *Cancer Res* 77: 5873-5885
- Nguyen HT, Dalmaso G, Muller S, Carriere J, Seibold F, Darfeuille-Michaud A (2014) Crohn's disease-associated adherent invasive Escherichia coli modulate levels of microRNAs in intestinal epithelial cells to reduce autophagy. *Gastroenterology* 146: 508-19
- Ni J, Wu GD, Albenberg L, Tomov VT (2017) Gut microbiota and IBD: causation or correlation? *Nat Rev Gastroenterol Hepatol* 14: 573-584
- Niess JH, Brand S, Gu X, Landsman L, Jung S, McCormick BA, Vyas JM, Boes M, Ploegh HL, Fox JG, Littman DR, Reinecker HC (2005) CX3CR1-mediated dendritic cell access to the intestinal lumen and bacterial clearance. *Science* 307: 254-8
- Nighot PK, Hu CA, Ma TY (2015) Autophagy enhances intestinal epithelial tight junction barrier function by targeting claudin-2 protein degradation. *J Biol Chem* 290: 7234-46
- Noda NN, Fujioka Y, Hanada T, Ohsumi Y, Inagaki F (2013) Structure of the Atg12-Atg5 conjugate reveals a platform for stimulating Atg8-PE conjugation. *EMBO Rep* 14: 206-11
- Nunes NS, Kim S, Sundby M, Chandran P, Burks SR, Paz AH, Frank JA (2018) Temporal clinical, proteomic, histological and cellular immune responses of dextran sulfate sodium-induced acute colitis. *World J Gastroenterol* 24: 4341-4355
- Ogawa A, Andoh A, Araki Y, Bamba T, Fujiyama Y (2004) Neutralization of interleukin-17 aggravates dextran sulfate sodium-induced colitis in mice. *Clin Immunol* 110: 55-62
- Ohtani M, Nagai S, Kondo S, Mizuno S, Nakamura K, Tanabe M, Takeuchi T, Matsuda S, Koyasu S (2008) Mammalian target of rapamycin and glycogen synthase kinase 3 differentially regulate lipopolysaccharide-induced interleukin-12 production in dendritic cells. *Blood* 112: 635-43
- Park TY, Jang Y, Kim W, Shin J, Toh HT, Kim CH, Yoon HS, Leblanc P, Kim KS (2019) Chloroquine modulates inflammatory autoimmune responses through Nurr1 in autoimmune diseases. *Sci Rep* 9: 15559
- Patel KK, Miyoshi H, Beatty WL, Head RD, Malvin NP, Cadwell K, Guan JL, Saitoh T, Akira S, Seglen PO, Dinanier MC, Virgin HW, Stappenbeck TS (2013) Autophagy proteins control goblet cell function by potentiating reactive oxygen species production. *EMBO J* 32: 3130-44
- Peng J, Zhang R, Cui Y, Liu H, Zhao X, Huang L, Hu M, Yuan X, Ma B, Ma X, Takashi U, Masaaki K, Liang X, Yu L (2014) Atg5 regulates late endosome and lysosome biogenesis. *Sci China Life Sci* 57: 59-68
- Peral de Castro C, Jones SA, Ni Cheallaigh C, Hearnden CA, Williams L, Winter J, Lavelle EC, Mills KH, Harris J (2012) Autophagy regulates IL-23 secretion and innate T cell responses through effects on IL-1 secretion. *J Immunol* 189: 4144-53
- Perussia B, Chan SH, D'Andrea A, Tsuji K, Santoli D, Pospisil M, Young D, Wolf SF, Trinchieri G (1992) Natural killer (NK) cell stimulatory factor or IL-12 has differential effects on the proliferation of TCR-alpha beta+, TCR-gamma delta+ T lymphocytes, and NK cells. *J Immunol* 149: 3495-502
- Reed M, Morris SH, Owczarczyk AB, Lukacs NW (2015) Deficiency of autophagy protein Map1-LC3b mediates IL-17-dependent lung pathology during respiratory viral infection via ER stress-associated IL-1. *Mucosal Immunol* 8: 1118-30
- Reitberger S, Haimerl P, Aschenbrenner I, Esser-von Bieren J, Feige MJ (2017) Assembly-induced folding regulates interleukin 12 biogenesis and secretion. *J Biol Chem* 292: 8073-8081
- Rescigno M, Urbano M, Valzasina B, Francolini M, Rotta G, Bonasio R, Granucci F, Kraehenbuhl JP, Ricciardi-Castagnoli P (2001) Dendritic cells express tight junction proteins and penetrate gut epithelial monolayers to sample bacteria. *Nat Immunol* 2: 361-7
- Rioux JD, Xavier RJ, Taylor KD, Silverberg MS, Goyette P, Huett A, Green T, Kuballa P, Barmada MM, Datta LW, Shugart YY, Griffiths AM, Targan SR, Ippoliti AF, Bernard EJ, Mei L, Nicolae DL, Regueiro M,

- Schumm LP, Steinhart AH et al. (2007) Genome-wide association study identifies new susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis. *Nat Genet* 39: 596-604
- Roberts RL, Gearry RB, Hollis-Moffatt JE, Miller AL, Reid J, Abkevich V, Timms KM, Gutin A, Lanchbury JS, Merriman TR, Barclay ML, Kennedy MA (2007) IL23R R381Q and ATG16L1 T300A are strongly associated with Crohn's disease in a study of New Zealand Caucasians with inflammatory bowel disease. *Am J Gastroenterol* 102: 2754-61
- Said A, Bock S, Lajqi T, Muller G, Weindl G (2014) Chloroquine promotes IL-17 production by CD4+ T cells via p38-dependent IL-23 release by monocyte-derived Langerhans-like cells. *J Immunol* 193: 6135-43
- Saitoh T, Fujita N, Jang MH, Uematsu S, Yang BG, Satoh T, Omori H, Noda T, Yamamoto N, Komatsu M, Tanaka K, Kawai T, Tsujimura T, Takeuchi O, Yoshimori T, Akira S (2008) Loss of the autophagy protein Atg16L1 enhances endotoxin-induced IL-1 β production. *Nature* 456: 264-8
- Sakoh-Nakatogawa M, Matoba K, Asai E, Kirisako H, Ishii J, Noda NN, Inagaki F, Nakatogawa H, Ohsumi Y (2013) Atg12-Atg5 conjugate enhances E2 activity of Atg3 by rearranging its catalytic site. *Nat Struct Mol Biol* 20: 433-9
- Sands BE, Sandborn WJ, Panaccione R, O'Brien CD, Zhang H, Johanns J, Adedokun OJ, Li K, Peyrin-Biroulet L, Van Assche G, Danese S, Targan S, Abreu MT, Hisamatsu T, Szapary P, Marano C, Group US (2019) Ustekinumab as Induction and Maintenance Therapy for Ulcerative Colitis. *N Engl J Med* 381: 1201-1214
- Schirmer M, Garner A, Vlamakis H, Xavier RJ (2019) Microbial genes and pathways in inflammatory bowel disease. *Nat Rev Microbiol* 17: 497-511
- Seksik P (2010) [Gut microbiota and IBD]. *Gastroenterol Clin Biol* 34 Suppl 1: S44-51
- Shouval DS, Biswas A, Kang YH, Griffith AE, Konnikova L, Mascanfroni ID, Redhu NS, Frei SM, Field M, Doty AL, Goldsmith JD, Bhan AK, Loizides A, Weiss B, Yerushalmi B, Yanagi T, Lui X, Quintana FJ, Muise AM, Klein C et al. (2016) Interleukin 1 β Mediates Intestinal Inflammation in Mice and Patients With Interleukin 10 Receptor Deficiency. *Gastroenterology* 151: 1100-1104
- Simon HU, Yousefi S, Schmid I, Friis R (2014) ATG5 can regulate p53 expression and activation. *Cell Death Dis* 5: e1339
- Singh N, Gurav A, Sivaprakasam S, Brady E, Padia R, Shi H, Thangaraju M, Prasad PD, Manicassamy S, Munn DH, Lee JR, Offermanns S, Ganapathy V (2014) Activation of Gpr109a, receptor for niacin and the commensal metabolite butyrate, suppresses colonic inflammation and carcinogenesis. *Immunity* 40: 128-39
- Singh SB, Davis AS, Taylor GA, Deretic V (2006) Human IRGM induces autophagy to eliminate intracellular mycobacteria. *Science* 313: 1438-41
- Sun M, Wu W, Chen L, Yang W, Huang X, Ma C, Chen F, Xiao Y, Zhao Y, Ma C, Yao S, Carpio VH, Dann SM, Zhao Q, Liu Z, Cong Y (2018) Microbiota-derived short-chain fatty acids promote Th1 cell IL-10 production to maintain intestinal homeostasis. *Nat Commun* 9: 3555
- Targan SR, Feagan B, Vermeire S, Panaccione R, Melmed GY, Landers C, Li D, Russell C, Newmark R, Zhang N, Chon Y, Hsu YH, Lin SL, Klekotka P (2016) A Randomized, Double-Blind, Placebo-Controlled Phase 2 Study of Brodalumab in Patients With Moderate-to-Severe Crohn's Disease. *Am J Gastroenterol* 111: 1599-1607
- Tominaga K, Yoshimoto T, Torigoe K, Kurimoto M, Matsui K, Hada T, Okamura H, Nakanishi K (2000) IL-12 synergizes with IL-18 or IL-1 β for IFN- γ production from human T cells. *Int Immunol* 12: 151-60
- Trinchieri G, Wysocka M, D'Andrea A, Rengaraju M, Aste-Amezaga M, Kubin M, Valiante NM, Chehimi J (1992) Natural killer cell stimulatory factor (NKSF) or interleukin-12 is a key regulator of immune response and inflammation. *Prog Growth Factor Res* 4: 355-68
- Vallon-Eberhard A, Landsman L, Yegorov N, Verrier B, Jung S (2006) Transepithelial pathogen uptake into the small intestinal lamina propria. *J Immunol* 176: 2465-9

- Watson RO, Manzanillo PS, Cox JS (2012) Extracellular M. tuberculosis DNA targets bacteria for autophagy by activating the host DNA-sensing pathway. *Cell* 150: 803-15
- Willing BP, Dicksved J, Halfvarson J, Andersson AF, Lucio M, Zheng Z, Jarnerot G, Tysk C, Jansson JK, Engstrand L (2010) A pyrosequencing study in twins shows that gastrointestinal microbial profiles vary with inflammatory bowel disease phenotypes. *Gastroenterology* 139: 1844-1854 e1
- Wong M, Ganapathy AS, Suchanec E, Laidler L, Ma T, Nighot P (2019) Intestinal epithelial tight junction barrier regulation by autophagy-related protein ATG6/beclin 1. *Am J Physiol Cell Physiol* 316: C753-C765
- Wu Y, Wang Y, Zou H, Wang B, Sun Q, Fu A, Wang Y, Wang Y, Xu X, Li W (2017) Probiotic *Bacillus amyloliquefaciens* SC06 Induces Autophagy to Protect against Pathogens in Macrophages. *Front Microbiol* 8: 469
- Yamazaki K, Onouchi Y, Takazoe M, Kubo M, Nakamura Y, Hata A (2007) Association analysis of genetic variants in IL23R, ATG16L1 and 5p13.1 loci with Crohn's disease in Japanese patients. *J Hum Genet* 52: 575-583
- Yang XO, Chang SH, Park H, Nurieva R, Shah B, Acero L, Wang YH, Schluns KS, Broaddus RR, Zhu Z, Dong C (2008) Regulation of inflammatory responses by IL-17F. *J Exp Med* 205: 1063-75
- Ye X, Zhou XJ, Zhang H (2018) Exploring the Role of Autophagy-Related Gene 5 (ATG5) Yields Important Insights Into Autophagy in Autoimmune/Autoinflammatory Diseases. *Front Immunol* 9: 2334
- Yoshimori T, Yamamoto A, Moriyama Y, Futai M, Tashiro Y (1991) Bafilomycin A1, a specific inhibitor of vacuolar-type H(+)-ATPase, inhibits acidification and protein degradation in lysosomes of cultured cells. *J Biol Chem* 266: 17707-12
- Yousefi S, Perozzo R, Schmid I, Ziemiecki A, Schaffner T, Scapozza L, Brunner T, Simon HU (2006) Calpain-mediated cleavage of Atg5 switches autophagy to apoptosis. *Nat Cell Biol* 8: 1124-32
- Zhang H, Zheng L, McGovern DP, Hamill AM, Ichikawa R, Kanazawa Y, Luu J, Kumagai K, Cilluffo M, Fukata M, Targan SR, Underhill DM, Zhang X, Shih DQ (2017) Myeloid ATG16L1 Facilitates Host-Bacteria Interactions in Maintaining Intestinal Homeostasis. *J Immunol* 198: 2133-2146
- Zhang M, Kenny SJ, Ge L, Xu K, Schekman R (2015) Translocation of interleukin-1beta into a vesicle intermediate in autophagy-mediated secretion. *Elife* 4
- Zhao S, Guo Y, Sheng Q, Shyr Y (2014) Advanced heat map and clustering analysis using heatmap3. *Biomed Res Int* 2014: 986048
- Zhao Z, Fux B, Goodwin M, Dunay IR, Strong D, Miller BC, Cadwell K, Delgado MA, Ponpuak M, Green KG, Schmidt RE, Mizushima N, Deretic V, Sibley LD, Virgin HW (2008) Autophagosome-independent essential function for the autophagy protein Atg5 in cellular immunity to intracellular pathogens. *Cell Host Microbe* 4: 458-69

Non-autophagy role of Atg5 and NBR1 in unconventional secretion of IL-12 prevents gut dysbiosis and inflammation

Merkley et al.,

Contents: Figures 1-5 and figure legends.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure legends

Figure 1. Alterations in the composition of the colonic microbial community in mice with a selective *Atg5*-deficiency in myeloid cells. Bacterial composition as assessed by 16S rRNA sequencing of DNA extracted from freshly collected stool of *Atg5*-Wt and *Atg5* Δ Mye mice (n=5 per group). **(A)** Pie chart showing the average proportion of Firmicutes, Bacteroidetes, Actinobacteria, Proteobacteria and all other phyla in 8-week old mice. **(B-E)** Quantification of the phyla Firmicutes, Bacteroidetes, Actinobacteria, and Proteobacteria. **(F)** Heatmap of the relative abundance of colonic microbes (Genus level). **(G)** Principal coordinates analysis (PCoA) plot of microbiota composition (Genus level). Data is shown as mean ($\pm$ 95%CI) and a T-test was used to measure specific microbiome species abundance between groups. Adjusted p-value > 0.05 was used as significant threshold.

Figure 2. Myeloid *Atg5* expression prevents excessive IFN γ -mediated intestinal inflammation. DSS-induced colitis and T cell response in *Atg5*-Wt and *Atg5* Δ Mye mice. **(A)** Percent weight loss between *Atg5*-Wt and *Atg5* Δ Mye mice. **(B)** Disease activity index (DAI) as determined by weight loss, behavior, acute diarrheal response and mucosal bleeding. **(C)** Colon length after colitis. **(D)** Spleen weight after colitis. **(E and F)** Colonic *Ifng* and *Il17a* gene expression after colitis induction. **(G and H)** Colonic IFN γ and IL-17 protein expression as determined by ELISA. **(I and J)** Percent of IFN γ ⁺ and IL-17⁺ CD4⁺ T cells isolated from the mLN of colitic mice. Representative of two independent experiments, Graphs indicate mean ($\pm$ SD). * P<0.05, **P<0.01. Two-tailed unpaired Student's t tests or by 2-way ANOVA with Tukey's post hoc test.

Figure 3. *Atg5* regulates IL-12 secretion in macrophages. Analysis of IL-12 secretion in macrophages. **(A)** Detection of IL-12 by ELISA from *Atg5*-Wt or *Atg5* Δ Mye BMM after LPS and IFN γ stimulation. **(B and C)** Detection of IL-12 by ELISA from *Atg5*-Wt **(B)** or *Atg5* Δ Mye **(C)** BMM after LPS and IFN γ stimulation in the presence or absence of Baf. A1. Representative of two independent of experiments. **(D - F)** 1-3 μ m Z-stack images were performed on BMM stimulated with LPS, IFN γ and Baf. A1 using immunofluorescence at 1.6 Zoom by a 63X oil immersion objective. **(D)** *Atg5*-Wt BMM were stained for LAMP1 (green) and IL-12p35 (red). **(E)** *Atg5*-Wt BMM were stained for ATG5 (red) and IL-12p35 (green). **(F)** *Atg5* Δ Mye BMM were stained for LAMP1 (green) and IL-12p35 (red). Representative from 65 images from 15 slides from two independent of experiments. Arrows in images indicate puncta co-localizing with insets displaying enlargement of indicated region. Scale bars: 10 μ m. Graphs indicate mean ($\pm$ SD). * P<0.05, **P<0.01, ***P<0.001. Two-tailed unpaired Student's t tests.

Figure 4. NBR1 and IL-12 interaction. Analysis of NBR1 and IL-12 interaction in macrophages. **(A)** Atg5-Wt BMM were stained for NBR1 (red) and IL-12p35 (green) after LPS and IFN γ stimulation and the addition of Bafilomycin A1 (Top Row) or Brefeldin A (Bottom Row). **(B)** Quantification was performed using Pearson's correlation coefficient (co-localization) using image analysis. Representative of two independent of experiments and of 10 images from 5 slides. Arrows in images indicate puncta co-localizing with insets displaying enlargement of indicated region. Scale bars: 10 μ m. **(C)** Anti-IL-12p35 antibody co-immunoprecipitated NBR1 and IL-12p40, and anti-NBR1 antibody co-immunoprecipitated IL-12p35 and IL-12p40. **(D and E)** Protein docking between IL12A (PDB ID: 1F45) and NBR1. **(D)** the docking models and **(E)** ClusPro docking results for four candidate interaction models between 1F45 and four peptide fragments of NBR1 (PDB IDs: 1WJ6, 2L8J, 2MGW, and 2MJ5). **(F and G)** Effects of NBR1 knockdown on IL-12 secretion in Atg5-Wt or Atg5 Δ Mye BMM. Graphs indicate mean ($\pm$ SD). * $P < 0.05$, ** $P < 0.01$. Two-tailed unpaired Student's t tests.

Figure 5. Limiting IL-12 in mice with a selective Atg5-deficiency in myeloid cells restores the intestinal microbiota and alleviates inflammation Bacterial composition as assessed by 16S rRNA sequencing from freshly collected stool of Atg5 Δ Mye and Atg5 Δ Mye-IL12KO mice prior to colitis induction (n=5 per group). **(A)** Principal coordinates analysis (PCoA) plot of microbiota composition (Genus level). **(B)** Heatmap of the relative abundance of colonic microbes (Genus level). A T-test was used to measure specific microbiome species abundance between groups. Adjusted p-value > 0.05 was used as significant threshold. **(C)** Percent weight loss between Atg5 Δ Mye and Atg5 Δ Mye-IL12KO mice. **(D)** Colon length after colitis. **(E)** Spleen weight after colitis. **(F and G)** Percent of IFN γ ⁺ and IL-17⁺ CD4⁺ T cells isolated from the mLN of colitic mice. Graphs indicate mean ($\pm$ SD). * $P < 0.05$, ** $P < 0.01$. Two-tailed unpaired Student's t tests or by 2-way ANOVA with Tukey's post hoc test.

Supplementary Information

Non-autophagy role of Atg5 and NBR1 in unconventional secretion of IL-12 prevents gut dysbiosis and inflammation

Seth D. Merkley et al.,

Contents: Supplementary materials and methods, supplementary figures, supplementary figures legends, and supplementary tables

Cells and tissue preparation. Mesenteric lymph nodes (mLN) were collected and manually homogenized into single cell suspensions in RPMI prior to plating and 4-hour stimulation with PMA (50 ng/mL), Ionomycin (500 ng/mL), Brefeldin A/GolgiPlug (1:1000, BD Biosciences), and Monesin/GolgiStop, (1:1000, BD Biosciences) in preparation for intracellular cytokine staining and analysis by flow cytometry. Colon samples were cryopreserved in RNA-Later (Invitrogen), or cultured in RPMI for 4 hrs for cytokine analysis by ELISA. The histopathological evaluation was performed on the mouse colon sections using Hematoxylin and Eosin (H and E) stained slides after formalin fixation. Standard five-micron thick H and E sections obtained from paraffin embedded blocks fixed in 10% formalin were used for assessment. The large intestine of the mice was embedded in paraffin using the 'swiss-roll' technique. The generation of BMM were prepared as briefly described, marrow was collected from the femur and tibia, washed and differentiated in DMEM (Gibco), FBS (VWR), and murine L929-fibroblast supernatant containing CSF-1 for 10 days, after which cellular morphology was evaluated to confirm differentiation. BMM's rested for 16 hours in media without CSF-1 prior to stimulation as specified for each experimental procedure below where relevant. All reagents used in cell isolation, culture, and stimulation are listed in Table S1. Flow cytometric protocol is described below, and all antibodies used for flow cytometric analysis are included in Table S7.

RNA Isolation, Quantification, and RT-qPCR

RNA isolation was performed on cells or tissues cryopreserved in RNA-Later (Invitrogen) using the RNA Purelink Minikit (Invitrogen) according to the manufacturer's protocols. RNA was quantified on Nanodrop2000 and all samples yielded a 260/280 of 2 ± 0.15 . cDNA synthesis was performed using Oligo(dT) Primer and SSIV Reverse Transcriptase in the presence of

Cloned Ribonuclease Inhibitor (all ThermoFisher). Reverse transcription reaction and RT-qPCR runs utilized Taqman MasterMix (ThermoFisher) on a viiA7 Thermal Cycler (ThermoFisher) using QuantStudio 7. RT-qPCR primers and reagents used are listed in Table S3.

ClusPro web server

The ClusPro web server for protein-protein docking was utilized to assess the interaction between IL-12p (PDB ID: 1F45) and NBR1 (PDB IDs: 1WJ6, 2L8J, 2MGW, and 2MJ5).

<https://cluspro.org/jobdetail.php?job=458409>

<https://cluspro.org/jobdetail.php?job=458405>

<https://cluspro.org/jobdetail.php?job=458404>

<https://cluspro.org/jobdetail.php?job=458403>

siRNA knockdown, Western blot, co-immunoprecipitation analysis and protein detection assays

Small interfering RNA knockdowns were achieved using Amaxa Mouse Macrophage Nucleofector™ Kit (Lonza). Nucleofection was performed on Nucleofector 2b Device (Lonza) per manufacturer's protocols, using 20 µM reconstitutions of Non-Targeting (Dharmacon) and 5 µM reconstitutions of NBR1 (Dharmacon) siRNA. Following Nucleofection, cells were plated and rested for 16 hours prior to stimulation. All macrophages were stimulated with LPS (500 ng/mL), IFN γ (10 ng/mL) for 6 hours, with one of two duplicates also receiving BafA1 for 5.5 hours. siRNA reagents are listed in Table S5. All ELISA were performed using Quantikine ELISA Kits (R&D) according to the manufacturer's specifications. For all analyses, undiluted cell media was collected immediately prior to cell harvesting and placed on ice prior to freezing. ELISA quantification was performed on Synergy Neo2 (BioTek) and plate washes using Quantikine wash buffer were performed on Multiwash Advantage (Tricontinent). All reagents used were taken from the Quantikine kits listed in Table S6. For flow cytometry, cells were washed post-stimulation with FACS Buffer [90% by volume 1x PBS, (Life Technologies), 10%FBS (VWR), and 0.05% 0.5 M EDTA, (Invitrogen)], and incubated for 20 min with antibody stain, followed by further FACS wash, and fixation in 1% PFA. To examine intracellular proteins, cells were permeabilize and incubated with intracellular cytokines and washed and resuspended FACS buffer. Sample analysis was performed on LSR Fortessa (BD Biosciences). Antibodies used for flow cytometric analysis are listed in Table S7. For immunoblots, after stimulation, cells were washed in 1x PBS and lysed in 20 µL of Pierce RIPA Lysis Buffer (ThermoFisher). All samples were boiled for 5 min at 65C in 1:1 Pierce RIPA Buffer: 2x Laemmli Buffer prior to gel electrophoresis at 90V (10% Mini-PROTEAN® TGX™ Precast Protein Gels, Mini-PROTEAN® Tetra Vertical Electrophoresis Cell for Mini Precast Gels, 2-gel, BioRad). Immunoblot transfer used the Trans-Blot® Turbo™ Transfer System (Bio-Rad), and all blots were probed in 5% dry

milk. All blots were imaged on Odyssey Clx (Licor Biosciences). For co-immunoprecipitated of IL-12p35 to NBR1 and IL-12p40 or NBR1 to IL-12p35 and IL-12p40, lysates were incubated with protein A/G magnetic beads with 2µg of anti-IL-12p35 or anti-NBR1 antibody or isotype control for 4 hours at 4°C. The bead-antibody complexes were washed with cold PBS followed by incubation with whole cell lysates overnight at 4°C. Following washes with lysis buffer and cold PBS, the bead-immune complexes were then resuspended in Laemmli's sample buffer and boiled for 5 minutes. Samples were then prepared for immunoblot analysis. All antibodies used for Western Blot are listed in Table S8.

Supplementary Figure 1. Steady State CD4⁺ T cell polarization and colitic response in Atg5cKO mice. (A) Principal coordinates analysis (PCoA) plot and (B) heatmap of microbiota composition in female mice (Genus level). (C) Flow cytometric gating of mLN CD4⁺ T cells for intracellular cytokine staining. Representative dot plots of intracellular IFNγ expression in mLN CD4⁺ T cells isolated from Atg5-Wt and Atg5ΔMyc mice. (D and E) Graph of IFNγ⁺ and IL-17⁺ CD4⁺ T cells isolated from the mLN of Atg5-Wt and Atg5ΔMyc mice. (F) Representative gross anatomy of the cecum and colon of colitic Atg5-Wt and Atg5ΔMyc mice. (G) H&E staining of colons. Representative of two-three independent experiments, Graphs indicate mean (±SD). Student's t test, P<0.05.

Supplementary Figure 2. Regulation of IL-12 secretion in macrophages. (A) *Il12a* and (B) *Il12b* gene expression in unstimulated macrophages. (C) Representative histograms of cell surface expression on IFN γ R (left), TLR4/MD2 (center), and CD14 (right) between unstimulated Atg5-Wt (gray) and Atg5ΔMyc (red) macrophages. (D) Atg5-Wt and Atg5ΔMyc macrophages were stained for LC3 (autophagosome marker) and IL-12p35 (subunit of cytokine IL12) after LPS, IFN γ and BafA1. Representative from 20 images from 5 slides. (E) Mean fluorescence intensity (MFI) of Lysotracker green in unstimulated Atg5-Wt and Atg5ΔMyc macrophages. (F) Atg5ΔMyc macrophages were stained for Rab7 (late endosome marker) and IL-12p35 (subunit of cytokine IL12) after LPS, IFN γ and BafA1. Immunoblot of IL-12p35 and actin expression in Atg5-Wt and Atg5ΔMyc macrophages after stimulation with LPS and IFN γ or LPS, IFN γ and BafA1. Representative from 20 images from 3 slides. Arrows in images indicate puncta co-localizing with insets displaying enlargement of indicated region. (G) Immunoblot of NBR1 expression for scrambled (Scr) or knockdown (KD) siRNA targeted of NBR1 in Atg5-Wt and Atg5ΔMyc macrophages after stimulation with LPS and IFN γ . Representative of two-three independent experiments, Graphs indicate mean ($\pm$ SD).

Supplementary Figure 3. Comparison of Atg5ΔMye and Atg5ΔMye-IL12KO mice.

Metagenomic sequencing and 16S profiling of DNA extracted from freshly collected stool of Atg5ΔMye and Atg5ΔMye-IL12KO mice (n=5 per group). **(A)** Pie chart showing the average proportion of Firmicutes, Bacteroidetes, Actinobacteria, Proteobacteria and all other phyla in 8-week old mice. **(B and C)** Colonic *Ifng* and *Il17a* gene expression after colitis induction. **(D)** Graphical abstract (BioRender). Representative of two independent experiments, Graphs indicate mean (±SD). * P<0.05, Two-tailed unpaired Student's t tests.