1 **Characterization and description of *Faecalibacterium***
2 ***butyricigenerans* sp. nov. and *F. longum* sp. nov., isolated**
3 **from human faeces**

4
5 Yuanqiang Zou^{1, 2, 3, 4§*}, Xiaoqian Lin^{1, 5§}, Wenbin Xue¹, Ying Dai¹, Karsten Kristiansen^{1, 2, 4}, Liang
6 Xiao^{1, 3, 4*}

7
8 ¹ BGI-Shenzhen, Shenzhen 518083, China

9 ² Laboratory of Genomics and Molecular Biomedicine, Department of Biology, University of
10 Copenhagen, Universitetsparken 13, 2100 Copenhagen, Denmark

11 ³ Shenzhen Engineering Laboratory of Detection and Intervention of human intestinal microbiome,
12 BGI-Shenzhen, Shenzhen, China

13 ⁴ Qingdao-Europe Advanced Institute for Life Sciences, BGI-Shenzhen, Qingdao 266555, China

14 ⁵ School of Bioscience and Biotechnology, South China University of Technology, Guangzhou
15 510006, China

16

17 * **Corresponding authors:**

18 Liang Xiao.

19 Address: BGI-Shenzhen, Beishan Industrial Zone, Shenzhen 518083, China.

20 E-mail: xiaoliang@genomics.cn

21 Yuanqiang Zou.

22 Address: BGI-Shenzhen, Beishan Industrial Zone, Shenzhen 518083, China.

23 E-mail: zouyuanqiang@genomics.cn

24

25 [†] Yuanqiang Zou and Xiaoqian Lin contributed equally to this work.

26 Running title: *Faecalibacterium butyricigenerans* sp. nov. and *Faecalibacterium longum* sp. nov.

27 Contents Category: New Taxa –*Firmicutes* and related organisms

28

29 **Keywords:** *Faecalibacterium*, *Faecalibacterium butyricigenerans* sp. nov., human faeces,
30 taxonomy, genome sequencing, phylogenetic analysis, average nucleotide identity

31

32 Abstract

33 Exploiting a pure culture strategy to investigate the composition of human gut microbiota,
34 two novel anaerobes, designated strains AF52-21^T and CM04-06^T, were isolated from faeces
35 of two healthy Chinese donors and characterized using a polyphasic approach. The two
36 strains were Gram-stain-negative, non-motile, and rod-shaped. Both strains grew optimally
37 at 37°C and pH 7.0. Phylogenetic analysis based on 16S rRNA gene sequences revealed that
38 the two strain clustered with species of the genus *Faecalibacterium* and were most closely
39 related to *Faecalibacterium prausnitzii* ATCC 27768^T with sequence similarity of 97.18% and
40 96.87%, respectively. The two isolates shared a 16S rRNA gene sequence identity of 98.69%.
41 Draft genome sequencing was performed for strains AF52-21^T and CM04-06^T, generating
42 genome sizes of 2.85 Mbp and 3.01 Mbp. The calculated average nucleotide identity values
43 between the genomes of the strains AF52-21^T and CM04-06^T compared to *Faecalibacterium*

44 *prausnitzii* ATCC 27768^T were 83.20% and 82.54%, respectively, and 90.09% when
 45 comparing AF52-21^T and CM04-06^T. Both values were below the previously proposed
 46 species threshold (95%), supporting their recognition as novel species in the genus
 47 *Faecalibacterium*. The genomic DNA G+C contents of strain AF52-21^T and CM04-06^T
 48 calculated from genome sequences were 57.77 mol% and 57.51 mol%, respectively. Based on
 49 the phenotypic, chemotaxonomic and phylogenetic characteristics, we conclude that both
 50 strains represent two new *Faecalibacterium* species, for which the names *Faecalibacterium*
 51 *butyricigenerans* sp. nov. (type strain AF52-21^T = CGMCC 1.5206^T = DSM 103434^T) and
 52 *Faecalibacterium longum* sp. nov. (type strain CM04-06^T = CGMCC 1.5208^T = DSM 103432^T)
 53 are proposed.

54

55 Introduction

56 The human gastrointestinal¹ tract harbours complex microbial communities², dominated by
 57 bacteria from the phyla *Bacteroidetes* and *Firmicutes*³⁻⁶. The composition and diversity of the gut
 58 microbiota are affected by numerous factors, including host genetics⁷, long-term diet^{8,9}, drugs^{1,10,11}
 59 and several other environmental factors¹². Evidence suggests that the composition of the
 60 microbiota is associated with the development of obesity^{4,13-15}, diabetes^{16,17}, inflammatory bowel
 61 disease^{18,19}, colorectal cancer^{20,21}, and non-alcoholic fatty liver disease^{22,23}. Therefore, the
 62 composition and function of the microbial species living in our gut are crucial importance for
 63 maintenance of health. Short-chain fatty acids (SCFAs), produced by fermentation of dietary fibre
 64 by several abundant genera of the intestinal microbiota, including *Roseburia*, *Eubacterium* and
 65 *Faecalibacterium*²⁴, have been reported to elicit beneficial effects on energy metabolism and for

66 prevention of colonization of pathogens²⁵. The genus *Faecalibacterium* as an abundant butyric
67 acid-producing bacterium colonizing the human gut displays anti-inflammatory effects and may be
68 used as a potential probiotics for treatment of gut inflammation^{26,27}.
69 The genus *Faecalibacterium*, belonging to the family *Ruminococcaceae* within the order
70 *Clostridiales*, comprises only one validated species, *Faecalibacterium prausnitzii*²⁸, and two
71 non-validly published species, *Faecalibacterium moorei*²⁹ and *Faecalibacterium hominis*³⁰, all
72 originally isolated from human faeces. *F. prausnitzii* is a gram-negative non-spore-forming and
73 strictly anaerobic rod-shaped bacterium. The genomic G+C content of genus *Faecalibacterium*
74 ranges from 47% to 57%³¹. The fermentation products from glucose are butyrate, D-lactate and
75 formate. In the present study, we describe two novel species of the genus *Faecalibacterium* by
76 using polyphasic taxonomy along with whole genome sequence analysis.

77

78 **Results and discussion**

79 **Phenotypic and Chemotaxonomic Characterization**

80 Both strains (AF52-21^T and CM04-06^T) were obligate anaerobic, Gram-stain-negative,
81 non-spore-forming, non-motile and rod-shaped bacteria (**Fig. 1**). After incubation on MPYG agar
82 at 37°C for 2 days, the colonies appeared 1.0-2.0 mm in diameter, round, creamy white to
83 yellowish, convex and opaque with entire margins for AF52-21^T and 2.0 mm in diameter, round,
84 yellowish, slightly convex and opaque with entire margins for CM04-06^T. The growth temperature
85 was 20-42°C (optimum 37°C) for AF52-21^T and 30-45°C (optimum 37°C) for CM04-06^T. Growth
86 was observed at pH 6.0-7.5 (optimum 7.0-7.5) for AF52-21^T and pH 5.0-8.0 (optimum 7.0-7.5) for
87 CM04-06^T. Strains AF52-21^T and CM04-06^T grew with 0-1% and 0-3% NaCl, respectively. Both

88 strains were catalase-negative. The major metabolic end products for strains AF52-21^T and
89 CM04-06^T were acetic acid, formic acid, butyric acid and lactic acid. Differential physiological
90 and biochemical characteristics of strains AF52-21^T and CM04-06^T with the closest related species
91 of genus *Faecalibacterium* are listed in the species description and in **Table 1**.

92

93 **Table 1. Differential phenotypic characteristics of strains AF52-21^T, CM04-06^T, and the**
94 **related species *F. prausnitzii* ATCC 27768^T.**

95 Strains: 1, *F. butyricigenerans* AF52-21^T; 2, *F. longum* CM04-06^T; 3, *F. prausnitzii* ATCC
96 27768^T. +, positive; w, weakly positive; –, negative.

Phenotypic features	1	2	3
Growth:			
Temperature range (optimum) (°C)	20-42 (37)	30-45 (37)	20-42 (37)
pH range	6.0-7.5	5.0-8.0	6.0-7.5
Salt tolerance (%)	1	3	3
Hydrolysis of:			
Aesculin	+	–	+
Gelatin	–	+	–
Acid from (API 20A and API 50CHL):			
Cellobiose	+	–	w
D-Fructose	w	–	+
D-Fucose	w	–	w
D-Galactose	w	–	–
D-Glucose	w	–	+
D-Lactose	+	–	–
D-Maltose	+	+	w
D-Fannitol	+	–	–
D-Fannose	w	–	–
D-Mannose	+	+	–
D-Raffinose	–	w	–
D-Trehalose	+	w	w
Gluconate	–	–	+
Glycogen	+	–	–

Inositol	w	—	—
Inulin	+	—	+
Methyl- β -D-Xylopyranoside	w	—	—
Enzyme activity (API ZYM):			
<i>N</i> -acetyl- β -Glucosaminidase	—	w	—
Naphthol-AS-BI-Phosphohydrolase	+	—	+
α -Glucosidase	—	—	+
β -Galactosidase	—	—	w
β -Glucosidase	+	—	—
β -Glucuronidase	+	w	+
DNA G+C (mol %)	57.77	57.51	52–57

97 All data are from this study.

98

99 The result of cellular fatty acid profiles of strain AF52-21^T, CM04-06^T and related species are
100 shown in **Table 2**. The major components of fatty acids (constituting >5% of the total) present in
101 strain AF52-21^T were C_{14:0} (5.9%), C_{16:0} (16.3%), C_{18:1} ω 7c (8.1), C_{18:1} ω 9c (39.0%) and iso-C_{19:0}
102 (12.9%). The profiles including C_{16:0} (25.5%), C_{18:1} ω 7c (7.5%), C_{18:1} ω 9c (32.5%), iso-C_{19:0} (5.9%)
103 and iso-C_{17:1} I/anteiso B (9.7%) were detected as the predominant fatty acids for strain CM04-06^T.
104 The highest levels of fatty acids, including C_{16:0} and C_{18:1} ω 9c, were similar, but not identical
105 comparing strain AF52-21^T, CM04-06^T and ATCC 27768^T. Furthermore, strains AF52-21^T,
106 CM04-06^T and ATCC 27768^T could be differentiated by less abundant fatty acids, such as C_{18:1}
107 2OH, anteiso-C_{15:0}, anteiso-C_{17:0}, C_{13:0} 3OH/Iso-C_{15:1} I, C_{16:1} ω 7c/C_{16:1} ω 6c and anteiso-C_{18:0}/C_{18:2}
108 ω 6, 9c (**Table 2**).

109

110 **Table 2. Fatty acid profile of strains AF52-21^T, CM04-06^T and the closest related species *F.***
111 ***prausnitzii* ATCC 27768^T.**

112 Numbers represent percentages of the total fatty acids. —, not detected (<1%).

Fatty acids composition	<i>F. butyricigenans</i> AF52-21 ^T	<i>F. longum</i> CM04-06 ^T	<i>F. prausnitzii</i> ATCC 27768 ^T
C _{12:0}	1.5	1.8	1.9
C _{13:1}	—	—	1.25
C _{14:0}	5.9	4.6	11.8
C _{16:0}	16.3	25.5	21.1
C _{17:1} ω8c	1.3	—	1.1
C _{18:1} ω7c	8.1	7.5	5.7
C _{18:1} ω9c	39.0	32.5	31.4
C _{18:0}	4.5	3.5	4.1
C _{18:1} 2OH	2.9	—	2.0
Iso-C _{19:1} I	1.2	1.1	2.1
Iso-C _{19:0}	12.9	5.9	—
Anteiso-C _{15:0}	—	2.6	—
Anteiso-C _{17:0}	—	2.1	—
C _{13:0} 3OH/ Iso-C _{15:1} I	—	—	2.1
C _{16:1} ω7c/ C _{16:1} ω6c	1.5	1.9	4.0
Iso-C _{17:1} I/anteiso B	4.7	9.7	7.6
Antei-C _{18:0} /C _{18:2} ω6, 9c	—	1.9	1.3

113

114 Phylogenetic Analysis

115 The almost complete 16S rRNA gene sequences of strains AF52-21^T and CM04-06^T, comprising

116 1,382bp and 1,374bp, respectively, were obtained. BLAST analysis of the 16S rRNA gene

117 sequences against the EzBioCloud server showed that the two strains grouped in the genus

118 *Faecalibacterium* within the family *Ruminococcaceae* and were most closely related to *F.*

119 *prausnitzii* ATCC 27768^T, which is the sole valid species of the genus *Faecalibacterium*, with

120 similarity values of 97.18% and 96.87%, respectively. *Faecalibacterium hominis* 4P-15^T, an

121 unrecognized species of the genus *Faecalibacterium*, was also used as a related taxa for 16S rRNA

122 gene analysis. Strains AF52-21^T and CM04-06^T share 16S rRNA gene sequence similarity of

123 98.65% and 97.68% with *F. hominis* 4P-15^T. The 16S rRNA gene sequence similarity between

124 strains AF52-21^T and CM04-06^T was 98.69% (**Table 3**), both these values were lower than the
125 recommended thresholds (98.7%) for classification of human-associated bacterial isolates at the
126 species level ³². Phylogenetic analysis based on the neighbour-joining, maximum-likelihood and
127 minimum-evolution (**Fig. 2, Supplementary Fig. S1 and Fig. S2**, respectively) confirmed the
128 result of affiliation of the novel isolates with the species of the genus *Faecalibacterium* and
129 revealed that the two isolates formed a distinct lineage with *F. prausnitzii* ATCC 27768^T.

130

131 **Table 3. Levels of 16S rRNA gene sequence similarity and ANI values (in percentages) based**
132 **on BLAST for strains AF52-21^T, CM04-06^T and the phylogenetically related species *F.***
133 ***prausnitzii* ATCC 27768^T and the unrecognized species *Faecalibacterium hominis* 4P-15^T.**

134 Taxa: 1, *F. butyricigenans* AF52-21^T; 2, *F. longum* CM04-06^T; 3, *F. prausnitzii* ATCC
135 27768^T; 4, *Faecalibacterium hominis* 4P-15^T.

Strain	Accession no.	1	2	3	4
16S rRNA gene sequence similarity (%)					
AF52-21 ^T	KX146426	100	98.69	97.18	98.65
CM04-06 ^T	KX150462	98.69	100	96.87	97.68
ATCC 27768 ^T	AJ413954	97.18	96.87	100	98.35
4P-15 ^T	NMDCN000012L	98.65	97.68	98.35	100
ANI values (%)					
AF52-21 ^T	CNA0017730	100	90.01	83.16	85.72
CM04-06 ^T	CNA0017731	90.19	100	82.53	85.40
ATCC 27768 ^T	CNA0017732	83.32	82.58	100	85.7985
4P-15 ^T	NMDC60014083	85.72	85.40	85.7985	100

136

137 **Genome Analysis and function annotation**

138 The assembled draft genomes of strains AF52-21^T and CM04-06^T comprised total lengths of
139 2,851,918bp and 3,011,178bp with 73 and 47 scaffolds, respectively (**Table 4**). The G+C contents
140 calculated from genome sequences were 57.77% and 57.51%, which were slightly higher than the
141 range reported previously for the genus *Faecalibacterium* (47-57 mol%)²⁸. CheckM analysis of
142 the genomes showed high completeness (>90%) and low contamination (<5%) (**Table 4**),
143 indicating these are high-quality genomes sequences. The genome comparison between strains
144 AF52-21^T, CM04-06^T, ATCC 27768^T and 4P-15^T showed ANI values ranged from 82.53% to
145 90.19% (**Table 3**), which were significantly below the proposed cutoff value of 95–96% for
146 delineating bacterial species, indicating that strains AF52-21^T and CM04-06^T represented novel
147 species in the genus *Faecalibacterium*. Circular maps of the two strains AF52-21^T and CM04-06^T
148 are shown in **Fig. 3** and **Fig. 4**.

149
150 **Table 4. Genome properties of *F. butyricigenerans* AF52-21^T and *F. longum* CM04-06^T.**

Feature	AF52-21 ^T	CM04-06 ^T
Accession No.	CNA0017730	CNA0017731
Approximate Genome Size (bp)	2,851,918	3,011,178
G+C content (mol%)	57.77	57.51
DNA scaffolds	73	47
N50 Length	191,233	119,299
Completeness	100	99.32
Contamination	0	0
Genes total number	2,291	2,506
Gene average length (bp)	939	920
rRNAs (5S, 16S, 23S)	4	5

tRNAs	60	61
sRNA	0	0
Genes assigned to COGs	2029	2164

151

152 For genome annotation, the distributions of the genes into clusters of orthologous groups (COGs)
153 functional categories are depicted in **Fig. 5** and **Table S1**. Both strain strains AF52-21^T and
154 CM04-06^T share identical COGs functional categories. The abundant categories comprise amino
155 acid transport and metabolism (E), carbohydrate transport and metabolism (G), cell
156 wall/membrane/envelope biogenesis (M), energy production and conversion (C), general function
157 prediction only (R), replication, recombination and repair (L), signal transduction mechanisms (T),
158 transcription (K) and translation, ribosomal structure and biogenesis (J). Annotated genes
159 associated with synthesis of diaminopimelic acid, teichoic and lipoteichoic acids and
160 lipopolysaccharides, and metabolism of polar lipids and polyamines by RAST annotation,
161 comparing strains AF52-21^T, CM04-06^T with ATCC 27768^T, are shown in **Table 5** and **Table S2**.

162

163 **Table 5. Number of genes associated with biosynthetic pathway from whole genome**
164 **sequences of strain *F. butyricigenerans* AF52-21^T and *F. longum* CM04-06^T and *F. prausnitzii***
165 **ATCC 27768^T identified by RAST.**

166 Taxa: 1, AF52-21^T; 2, CM04-06^T; 3, *F. prausnitzii* ATCC 27768^T. Data are for type strains.

167 Numbers of genes identified for lipopolysaccharides and mycolic acids were zero for all taxa
168 studied.

Genes responsible for biosynthesis	1	2	3
---	----------	----------	----------

Diaminopimelic acid	11	12	12
Polar lipids	18	19	22
Polyamines	12	13	11
Quinones	14	16	15
Teichoic and lipoteichoic acids	3	2	3

169

170 The annotation showed that AF52-21^T, CM04-06^T, and ATCC 27768^T contained a complete
171 acetyl-CoA to butyrate synthesis pathway, but possessed butyryl-CoA:acetate CoA-transferase
172 activity only in the final step (**Fig. 6**), as discussed previously^{33,34}. Prophages were identified
173 using the PHAST software, and the results are shown in **Fig. 7**. Two incomplete phage sequences
174 were detected in the AF52-21^T genome, one of which encodes the Phd_YefM protein, an antitoxin
175 component. Three incomplete phage sequences and two intact prophages were detected in the
176 CM04-06^T genome, encoding the Phd_YefM protein, relaxase/mobilisation nuclease domain,
177 bacterial mobilisation protein (MobC) /ribbon-helix-helix protein, helix-turn-helix, and predicted
178 transcriptional regulators. Moreover, the antibiotic resistance analysis indicated that AF52-21^T
179 contained macrolide antibiotic, lincosamide antibiotic, and streptogramin antibiotic genes, while
180 CM04-06^T and ATCC 27768^T contained aminoglycoside antibiotic genes (**Fig. 8**).

181

182 Discussion

183 The 16S rRNA genes phylogenetic, physiological results and genome description showed that the
184 two new isolates AF52-21^T and CM04-06^T represent two novel species. The ANI values between
185 AF52-21^T, CM04-06^T and the closest related species ATCC 27768^T were 82.54% and 90.09%,
186 respectively, which is in support of a new species delineation. The result of biochemical and

187 genomic functional analyses showed that both AF52-21^T and CM04-06^T are butyric
188 acid-producing bacteria.

189 Most strains in the genus *Faecalibacterium* exhibit a common ability to produce butyric acid,
190 peptides and other anti-inflammatory substances, which have immunomodulatory effects^{26,27,35}.

191 Some studies have confirmed that the decreased abundance of this genus is related to the
192 occurrence and development of inflammatory bowel diseases³⁶⁻³⁸. Accordingly, *Faecalibacterium*
193 is receiving much attention as one of the candidate next-generation probiotics (NGPs) , which can
194 be used for disease treatment^{39,40}.

195 Previous studies based on comparative genomics from isolates suggested the wide diversity of this
196 genus, with the presence of at least two phylotypes in *F. prausnitzii*²⁹. A recent study analysing the
197 *Faecalibacterium*-like MAGs, proposed that *Faecalibacterium* from the human gut can be divided
198 into 12 clades⁴⁰. These studies have expanded the diversity of *Faecalibacterium* and proposed that
199 different phylotypes have different functions, which results in different contributions to health or
200 diseases.

201 Moreover, as a candidate taxa for the NGPs, the *Faecalibacterium* isolates can be used for in-vitro
202 functional verification and animal model experiments to further explore its probiotic functions,
203 and ultimately expected to be used in clinical disease intervention.

204

205 **Description of *Faecalibacterium butyricigenerans* sp. nov.**

206 *Faecalibacterium butyricigenerans* (bu.ty.ri.ci.ge'ne.rans. N.L. n. *acidum butyricum* butyric
207 acid; L. part. adj. *generans*, producing; N.L. adj. *butyricigenerans*, butyric acid-producing;
208 referring to its production of butyric acid)

Cells of strain AF52-21^T are Gram-stain-negative, non-motile, non-spore-forming and rod-shaped. Strictly anaerobic and catalase negative. Colonies on PYG agar are round, creamy white to yellowish, convex and opaque with entire margins and colony size is approximately 1.0-2.0 mm in diameter after incubation at 37°C for 2 days. Cells are able to grow at 20-42°C with optimum temperature at 37°C. The pH range for growth is 6.0-7.5 (optimum at 7.0-7.5). Growth occurs at NaCl concentrations 0-1%. The strain is positive for the assimilation of cellobiose, D-lactose, D-maltose, D-mannitol, D-mannose, D-trehalose, glycogen, inulin, D-fructose, D-fucose, D-galactose, D-glucose, inositol and methyl- β -D-xylopyranoside, but negative for amygdalin, arbutin, D-adonitol, D-arabinose, D-arabitol, D-lyxose, D-melezitose, D-melibiose, D-raffinose, D-ribose, D-sorbitol, D-tagatose, D-turanose, dulcitol, D-xylose, erythritol, gentiobiose, gluconate, glycerol, L-arabinose, L-arabitol, L-fucose, L-rhamnose, L-sorbose, L-xylose, methyl-D-glucopyranoside, methyl- α -D-mannopyranoside, *N*-acetyl-glucosamine, salicin, starch, sucrose, xylitol, 2-ketogluconate and 5-ketogluconate. In enzymatic activity tests, strain AF52-21^T is positive for naphthol-AS-BI-phosphohydrolase, β -glucuronidase and β -glucosidase, and negative for alkaline phosphatase, esterase (C4), esterase lipase (C8), lipase (C14), leucine arylamidase, valine arylamidase, cystine arylamidase, trypsin, α -chymotrypsin, acid phosphatase, naphthol-AS-BI-phosphohydrolase, α -galactosidase, β -galactosidase, α -glucosidase, α -mannosidase and β -fucosidase. Indole is not produced. Positive for hydrolysis of esculin and negative for gelatin. Formic acid, acetic acid, butyric acid and lactic acid are the fermentation products. The major fatty acids are C_{14:0}, C_{16:0}, C_{18:1} ω 7c, C_{18:1} ω 9c and iso-C_{19:0}.

In the result of RAST annotation, 11 genes/proteins are associated with biosynthesis of DAP, including 4-hydroxy-tetrahydronicotinate reductase (EC 1.17.1.8),

231 4-hydroxy-tetrahydrodipicolinate synthase (EC 4.3.3.7), aspartate-semialdehyde dehydrogenase
232 (EC 1.2.1.11), aspartokinase (EC 2.7.2.4) (2 copies), diaminopimelate decarboxylase (EC
233 4.1.1.20), diaminopimelate epimerase (EC 5.1.1.7), L, L-diaminopimelate aminotransferase (EC
234 2.6.1.83), *N*-acetyl-L, L-diaminopimelate deacetylase (EC 3.5.1.47),
235 UDP-*N*-acetylmuramoylalanyl-D-glutamate-2, 6-diaminopimelate ligase (EC 6.3.2.13) and
236 UDP-*N*-acetylmuramoylalanyl-D-glutamyl-2, 6-diaminopimelate-D-alanyl-D-alanine ligase (EC
237 6.3.2.10). 18 genes/proteins are associated with biosynthesis of polar lipids, including
238 1-acyl-sn-glycerol-3-phosphate acyltransferase (EC 2.3.1.51) (2 copies), ABC-type
239 multidrug/protein/lipid transport system, ATPase component, acyl carrier protein (3 copies),
240 acyl-phosphate:glycerol-3-phosphate O-acyltransferase PlsY, cardiolipin synthetase (EC 2.7.8.-) (3
241 copies), CDP-diacylglycerol--glycerol-3-phosphate 3-phosphatidyltransferase (EC 2.7.8.5),
242 dihydroxyacetone kinase family protein, glycerol kinase (EC 2.7.1.30), glycerol-3-phosphate
243 dehydrogenase [NAD(P)⁺] (EC 1.1.1.94), phosphate:acyl-ACP acyltransferase PlsX,
244 phosphatidate cytidyltransferase (EC 2.7.7.41) and phosphatidylglycerophosphatase B (EC
245 3.1.3.27) (2 copies). 12 genes/proteins are associated with biosynthesis of polyamines, including
246 5'-methylthioadenosine nucleosidase (EC 3.2.2.16), S-adenosylhomocysteine nucleosidase (EC
247 3.2.2.9), ABC transporter, periplasmic spermidine putrescine-binding protein PotD (TC
248 3.A.1.11.1), agmatinase (EC 3.5.3.11), arginine decarboxylase (EC 4.1.1.19), arginine/ornithine
249 antiporter ArcD (2 copies), carboxynorspermidine decarboxylase, putative (EC 4.1.1.-),
250 carboxynorspermidine dehydrogenase, putative (EC 1.1.1.-), putrescine transport ATP-binding
251 protein PotA (TC 3.A.1.11.1), spermidine putrescine ABC transporter permease component PotB
252 (TC 3.A.1.11.1), spermidine putrescine ABC transporter permease component potC

(TC..3.A.1.11.1) and spermidine synthase (EC 2.5.1.16). 3 genes/proteins are associated with biosynthesis of teichoic and lipoteichoic acids, including cell wall teichoic acid glycosylation protein gtcA, teichoic acid export ATP-binding protein TagH (EC 3.6.3.40) and membrane protein involved in the export of O-antigen, teichoic acid lipoteichoic acids. 14 genes/proteins are associated with biosynthesis of quinones, including 2-heptaprenyl-1,4-naphthoquinone methyltransferase (EC 2.1.1.163), electron transport complex protein RnfA (2 copies), electron transport complex protein RnfB, electron transport complex protein RnfC, electron transport complex protein RnfD (2 copies), electron transport complex protein RnfE (2 copies), electron transport complex protein RnfG, F420H2:quinone oxidoreductase, heptaprenyl diphosphate synthase component I (EC 2.5.1.30), microsomal dipeptidase (EC 3.4.13.19) and undecaprenyl diphosphate synthase (EC 2.5.1.31). There are no genes responsible for biosynthesis of lipopolysaccharides or mycolic acids. Additional annotations showed that the AF52-21^T genome contains a complete butyrate synthesis pathway, two prophage remnants, and three antibiotic genes.

The type strain, AF52-21^T (=CGMCC 1.5206^T = DSM 103434^T), was isolated from human gut. The G+C content of the genomic DNA is 57.77 mol% as calculated from whole genome sequencing.

270

271 **Description of *Faecalibacterium longum* sp. nov.**

272 *Faecalibacterium longum* (lon'gum. L. neut. adj. *longum* long, the shape of the cells)

273 Cells are Gram-stain-negative, non-motile, non-spore forming, long rod in shape. Strictly
274 anaerobic. Catalase and urease are negative. Colonies are round, yellowish, slightly convex,

275 opaque with entire margins with 2.0 mm in diameter on PYG agar for incubation at 37°C for 48 h
276 under anaerobic condition. The strain showed growth at 30-45°C (optimum temperature is 37°C).
277 Growth is observed at pH 5.0-8.0 (optimum pH is 7.0-7.5). NaCl is tolerated with concentrations
278 up to 3%. Acid is produced from D-maltose, D-mannose, D-raffinose, D-trehalose and salicin, but
279 not from amygdalin, arbutin, cellobiose, D-adonitol, D-arabinose, D-arabitol, D-cellobiose,
280 D-fructose, D-fucose, D-galactose, D-glucose, D-lactose, D-lyxose, D-maltose, D-mannitol,
281 D-mannose, D-melezitose, D-melibiose, D-raffinose, D-ribose, D-sorbitol, D-sucrose, D-tagatose,
282 D-turanose, dulcitol, D-xylose, erythritol, gentiobiose, gluconate, glycerol, glycogen, inositol,
283 inulin, L-arabinose, L-arabitol, L-fucose, L-rhamnose, L-sorbose, L-xylose,
284 methyl-D-glucopyranoside, methyl- α -D-mannopyranoside, methyl- β -D-xylopyranoside,
285 *N*-acetyl-glucosamine, salicin, starch, sucrose, xylitol, 2-ketogluconate and 5-ketogluconate. In the
286 API ZYM strip, strain showed weakly positive enzyme activities for β -glucuronidase and
287 *N*-acetyl- β -glucosaminidase, but negative for alkaline phosphatase, esterase (C4), esterase lipase
288 (C8), lipase (C14), leucine arylamidase, valine arylamidase, cystine arylamidase, trypsin,
289 α -chymotrypsin, acid phosphatase, naphthol-AS-BI-phosphohydrolase, α -galactosidase,
290 β -galactosidase, α -glucosidase, β -glucosidase, α -mannosidase and β -fucosidase. Indole is not
291 produced. Gelatin is hydrolysed, but aesculin is not. Major end products are acetic acid, formic
292 acid, butyric acid and lactic acid. The major fatty acids (constituting >5% of the total) are C_{16:0},
293 C_{18:1} ω 7c, C_{18:1} ω 9c, iso-C_{19:0} and iso-C_{17:1} I/anteiso B.

294 In the result of RAST annotation, 12 genes/proteins are associated with biosynthesis of DAP,
295 including 4-hydroxy-tetrahydronicotinate reductase (EC 1.17.1.8),
296 4-hydroxy-tetrahydronicotinate synthase (EC 4.3.3.7) (2 copies), aspartate-semialdehyde

297 dehydrogenase (EC 1.2.1.11), aspartokinase (EC 2.7.2.4) (2 copies), diaminopimelate
298 decarboxylase (EC 4.1.1.20), diaminopimelate epimerase (EC 5.1.1.7), L, L-diaminopimelate
299 aminotransferase (EC 2.6.1.83), *N*-acetyl-L, L-diaminopimelate deacetylase (EC 3.5.1.47),
300 UDP-*N*-acetylmuramoylalanyl-D-glutamate-2, 6-diaminopimelate ligase (EC 6.3.2.13) and
301 UDP-*N*-acetylmuramoylalanyl-D-glutamyl-2, 6-diaminopimelate-D-alanyl-D-alanine ligase (EC
302 6.3.2.10). 19 genes/proteins are associated with biosynthesis of polar lipids, including
303 1-acyl-sn-glycerol-3-phosphate acyltransferase (EC 2.3.1.51) (2 copies), ABC-type
304 multidrug/protein/lipid transport system, ATPase component, acyl carrier protein (3 copies),
305 acyl-phosphate:glycerol-3-phosphate O-acyltransferase PlsY, cardiolipin synthetase (EC 2.7.8.-) (2
306 copies), CDP-diacylglycerol--glycerol-3-phosphate 3-phosphatidyltransferase (EC 2.7.8.5),
307 dihydroxyacetone kinase family protein, glycerate kinase (EC 2.7.1.31), glycerol kinase (EC
308 2.7.1.30), glycerol-3-phosphate dehydrogenase [NAD(P)⁺] (EC 1.1.1.94), octaprenyl diphosphate
309 synthase (EC 2.5.1.90) / geranylgeranyltransferase (EC 2.5.1.1) / (2E,6E)-farnesyl diphosphate
310 synthase (EC 2.5.1.10) / geranylgeranyl pyrophosphate synthetase (EC 2.5.1.29),
311 phosphate:acyl-ACP acyltransferase PlsX, phosphatidate cytidyltransferase (EC 2.7.7.41) and
312 phosphatidylglycerophosphatase B (EC 3.1.3.27) (2 copies). 13 genes/proteins are associated with
313 biosynthesis of polyamines, including 5'-methylthioadenosine nucleosidase (EC 3.2.2.16) @
314 S-adenosylhomocysteine nucleosidase (EC 3.2.2.9), ABC transporter, periplasmic spermidine
315 putrescine-binding protein PotD (TC 3.A.1.11.1), agmatinase (EC 3.5.3.11), arginine
316 decarboxylase (EC 4.1.1.19), arginine/ornithine antiporter ArcD (3 copies), carboxynorspermidine
317 decarboxylase, putative (EC 4.1.1.-), carboxynorspermidine dehydrogenase, putative (EC 1.1.1.-),
318 putrescine transport ATP-binding protein PotA (TC 3.A.1.11.1), spermidine putrescine ABC

319 transporter permease component PotB (TC 3.A.1.11.1), spermidine putrescine ABC transporter
320 permease component potC (TC..3.A.1.11.1) and spermidine synthase (EC 2.5.1.16). 2
321 genes/proteins are associated with biosynthesis of teichoic and lipoteichoic acids, including cell
322 wall teichoic acid glycosylation protein gtcA and teichoic acid export ATP-binding protein TagH
323 (EC 3.6.3.40). 16 genes/proteins are associated with biosynthesis of quinones, including
324 2-heptaprenyl-1,4-naphthoquinone methyltransferase (EC 2.1.1.163), electron transport complex
325 protein RnfA (2 copies), electron transport complex protein RnfB, electron transport complex
326 protein RnfC, electron transport complex protein RnfD (2 copies), electron transport complex
327 protein RnfE (2 copies), electron transport complex protein RnfG, heptaprenyl diphosphate
328 synthase component I (EC 2.5.1.30), microsomal dipeptidase (EC 3.4.13.19), octaprenyl
329 diphosphate synthase (EC 2.5.1.90) / dimethylallyltransferase (EC 2.5.1.1) / (2E,6E)-farnesyl
330 diphosphate synthase (EC 2.5.1.10) / geranylgeranyl pyrophosphate synthetase (EC 2.5.1.29)
331 ubiquinone/menaquinone biosynthesis methyltransferase UbiE (EC 2.1.1.-) @
332 2-heptaprenyl-1,4-naphthoquinone methyltransferase MenG (EC 2.1.1.163) (2 copies) and
333 undecaprenyl diphosphate synthase (EC 2.5.1.31). There are no genes responsible for biosynthesis
334 of lipopolysaccharides or mycolic acids. Additional annotations showed that the CM04-06^T
335 genome contains a complete butyrate synthesis pathway, three phage remnants, two intact
336 prophages, and aminoglycoside antibiotic genes.

337 The type strain, CM04-06^T (=CGMCC 1.5208^T = DSM 103432^T), was isolated from human
338 gut. The G+C content of the genomic DNA is 57.51 mol% as calculated from whole genome
339 sequencing.

340

341 **Materials and Methods**

342 **Origin of bacterial strains**

343 Faeces samples were collected from two healthy donors living in Shenzhen, Guangdong province,
344 China (GPS positioning of the samples collection site is 37°35'37"N/114°15'32"E) and preserved
345 refrigerated and anaerobically until processed. The collection of the samples was approved by the
346 Institutional Review Board on Bioethics and Biosafety of BGI under number BGI-IRB17005-T1.
347 All protocols were in compliance with the Declaration of Helsinki and explicit informed consent
348 was obtained from the participants. 1 g of faecal sample was diluted with 0.1 M PBS (pH 7,
349 supplemented with 0.5% cysteine) and spread onto modified peptone-yeast extract-glucose
350 (MPYG, supplemented with 5g/L sodium acetate in DSMZ 104 medium) agar plates in an
351 anaerobic box (Bactron Anaerobic Chamber, Bactron[®]-2, shellab, USA). The plates were
352 incubated at 37°C under anaerobic conditions (90% N₂, 5% CO₂ and 5% H₂, v/v) for 3 to 5 days.
353 Single colonies were randomly picked and purified by repetitive subculturing on the new plates
354 containing the same medium and incubated under the same conditions as described above. Among
355 the pure cultures, two isolates, designated as AF52-21^T and CM04-06^T, respectively, were
356 obtained and subsequently maintained in 20% (v/v) glycerol and frozen at -80°C.

357 **Phenotypic characterization**

358 The morphological characteristics of strains AF52-21^T and CM04-06^T were performed on cultures
359 grown on MPYG medium at 37°C. Bacterial cell shape was examined by phase contrast
360 microscopy (Olympus BX51, Japan) during the exponential phase of growth. Cell motility was
361 examined using semi-solid MPYG medium containing 0.5% agar⁴¹. The Gram reaction was
362 carried out using a Gram-staining kit (Solarbio, China). Spore formation and presence of flagella

were determined by staining using spore stain kit and flagella stain kit supplied by Solarbio (China) following the manufacturer's instructions. Colony morphology was observed following growth of the cultures on PYG agar for 2 days at 37°C. Optimal temperature for growth was determined using growth in MPYG medium at 4, 10, 20, 25, 30, 35, 37, 45 and 50°C for 7 days. The pH range for growth was also measured in MPYG medium covering the range of pH 3.0–10.0 (at interval of 0.5 pH units) at 37°C for 7 days. Growth at various NaCl concentrations (0–6%, in increments of 1.0%) was performed for determining tolerance to NaCl. Bile tolerance was measured at different bile salt concentrations (0–5%, at an interval of 1.0%) in the MPYG medium. Catalase activity was assessed by gas formation after dropping the fresh cells in 3% H₂O₂ solution. Biochemical properties, including utilization of substrates, acid production from carbohydrates, enzyme activities, hydrolytic activities, were determined using the API 20A, API 50CHL and API ZYM systems ((bioMérieux Inc., Marcy-l'Étoile, France) according to the manufacturer's instructions with modification by adding sodium acetate at concentration of 0.5% in all tests. The reference type strain was tested under the same condition with strains AF52-21^T and CM04-06^T.

Chemotaxonomic characteristics

Chemotaxonomic features were investigated by analyses of cellular fatty acids. Biomasses of strains AF52-21^T and CM04-06^T were harvested from cells growing in MPYG at 37°C for 2 days. Whole cell fatty acid methyl esters (FAMES) were extracted, separated and identified according to the MIDI Microbial Identifications System and performed by CGMGG (China General Microbiological Culture Collection Center, Beijing, China) identification service.

Fermentation products analysis

For analysis the metabolic end products from glucose fermentation, including SCFAs and organic

385 acids, cells were cultured in MPYG broth at 37°C for 2 days. Supernatant harvested from the
386 cultures centrifuged at 10000g for 10min was used for determining SCFAs and organic acids.
387 SCFAs detection was performed using a gas chromatograph (GC-7890B, Agilent) equipped with a
388 flame ionization detector (FID) and capillary column packed with Agilent
389 19091N-133HP-INNOWax porapak HP-INNOWax (30m × 0.25mm × 0.25um). Organic acids
390 were analysed by equipping capillary column packed with Agilent 122-5532G DB-5ms (40m ×
391 0.25mm × 0.25um).

392 **PCR of bacterial 16S rRNA genes and phylogenetic analysis**

393 Total genomic DNA of strains AF52-21^T and CM04-06^T were extracted using the standard
394 phenol:chloroform method as described by Cheng and Jiang⁴². The complete 16S rRNA genes
395 were amplified and sequenced according to the method previously described⁴³. Primers used for
396 amplification of 16S rRNA genes were 27f (5'-AGAGTTTGATCATGGCTCAG-3') and 1492r
397 (5'-TAGGGTTACCTTGTTACGACTT-3'). The obtained 16S rRNA gene sequences of strains
398 AF52-21^T and CM04-06^T were compared with the sequences of type strains retrieved from the
399 EzBioCloud database (<https://www.ezbiocloud.net/>)⁴⁴ using the BLAST program to determine the
400 nearest phylogenetic neighbours and 16S rRNA gene sequence similarity values. Phylogenetic
401 trees were reconstructed by using the neighbour-joining method⁴⁵, maximum-likelihood method⁴⁶
402 and minimum-evolution method⁴⁷ with the MEGA X program package⁴⁸, after Clustal W multiple
403 alignment of the sequences. Robustness of the phylogenetic trees was evaluated by using the
404 bootstrap resampling method (1000 resamplings) of Felsenstein⁴⁹.

405 **Genome sequencing , assembly, and annotation of isolates**

406 For genome sequences of strains AF52-21^T and CM04-06^T, genome DNA was prepared following

the method described above. The draft genome was sequenced on an Ion Proton Technology (Life Technologies) platform at BGI-Shenzhen (Shenzhen, China) after constructing a paired-end DNA library with insert size of 500 bp. The resulting reads were assembled using the SOAPdenovo 2 package⁵⁰. CheckM (v1.1.2) was used to estimate genome completeness and contamination⁵¹. Genome assemblies were visualized using CGView Server⁵² (http://stothard.afns.ualberta.ca/cgview_server/index.html). Annotation of the assembled genome was performed using the Rapid Annotation Using Subsystem Technology (RAST) server⁵³ and COG database⁵⁴. The G+C content in genomic DNA was calculated from the whole genome sequence. The genes in known pathways from acetyl-CoA to butyrate were annotated by BLAST (evalue=1e-5, identity≥60%, coverage≥90%)³³. A search for prophages was performed by PHAST (<http://phast.wishartlab.com/>)⁵⁵. Antibiotic resistance was analysed using the CARD database⁵⁶.

Average nucleotide identities

Genome relatedness was investigated by calculating average nucleotide identity (ANI)⁵⁷, with a value of 95-96% proposed for delineating bacterial species, corresponding to the traditional 70% DNA-DNA reassociation standard^{58,59}. The ANI values between strains AF52-21^T, CM04-06^T, and closely related species were determined using the FastANI⁶⁰.

References

- 1 Russell, S. L. *et al.* Early life antibiotic-driven changes in microbiota enhance susceptibility to allergic asthma. *EMBO Rep* **13**, 440-447, doi:10.1038/embor.2012.32 (2012).
- 2 Costello, E. K. *et al.* Bacterial community variation in human body habitats across space and time. *Science* **326**, 1694-1697, doi:10.1126/science.1177486 (2009).
- 3 Robles Alonso, V. & Guarner, F. Linking the gut microbiota to human health. *Br J Nutr* **109** Suppl 2, S21-26, doi:10.1017/S0007114512005235 (2013).
- 4 Tagliabue, A. & Elli, M. The role of gut microbiota in human obesity: recent findings and future perspectives. *Nutr Metab Cardiovasc Dis* **23**, 160-168,

doi:10.1016/j.numecd.2012.09.002 (2013).

5 Ringel-Kulka, T. *et al.* Intestinal microbiota in healthy U.S. young children and adults--a high throughput microarray analysis. *PLoS One* **8**, e64315, doi:10.1371/journal.pone.0064315 (2013).

6 Voreades, N., Kozil, A. & Weir, T. L. Diet and the development of the human intestinal microbiome. *Front Microbiol* **5**, 494, doi:10.3389/fmicb.2014.00494 (2014).

7 Benson, A. K. *et al.* Individuality in gut microbiota composition is a complex polygenic trait shaped by multiple environmental and host genetic factors. *Proc Natl Acad Sci U S A* **107**, 18933-18938, doi:10.1073/pnas.1007028107 (2010).

8 Claesson, M. J. *et al.* Gut microbiota composition correlates with diet and health in the elderly. *Nature* **488**, 178-184, doi:10.1038/nature11319 (2012).

9 De Filippo, C. *et al.* Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci U S A* **107**, 14691-14696, doi:10.1073/pnas.1005963107 (2010).

10 Antunes, L. C. & Finlay, B. B. A comparative analysis of the effect of antibiotic treatment and enteric infection on intestinal homeostasis. *Gut Microbes* **2**, 105-108 (2011).

11 Wilson, I. D. & Nicholson, J. K. The role of gut microbiota in drug response. *Curr Pharm Des* **15**, 1519-1523 (2009).

12 Adlerberth, I. Factors influencing the establishment of the intestinal microbiota in infancy. *Nestle Nutr Workshop Ser Pediatr Program* **62**, 13-29; discussion 29-33, doi:10.1159/000146245 (2008).

13 DiBaise, J. K. *et al.* Gut microbiota and its possible relationship with obesity. *Mayo Clin Proc* **83**, 460-469, doi:10.4065/83.4.460 (2008).

14 Ley, R. E., Turnbaugh, P. J., Klein, S. & Gordon, J. I. Microbial ecology: human gut microbes associated with obesity. *Nature* **444**, 1022-1023, doi:10.1038/4441022a (2006).

15 Parekh, P. J., Arusi, E., Vinik, A. I. & Johnson, D. A. The role and influence of gut microbiota in pathogenesis and management of obesity and metabolic syndrome. *Front Endocrinol (Lausanne)* **5**, 47, doi:10.3389/fendo.2014.00047 (2014).

16 Larsen, N. *et al.* Gut microbiota in human adults with type 2 diabetes differs from non-diabetic adults. *PLoS One* **5**, e9085, doi:10.1371/journal.pone.0009085 (2010).

17 Qin, J. *et al.* A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature* **490**, 55-60, doi:10.1038/nature11450 (2012).

18 Conte, M. P. *et al.* Gut-associated bacterial microbiota in paediatric patients with inflammatory bowel disease. *Gut* **55**, 1760-1767, doi:10.1136/gut.2005.078824 (2006).

19 Joossens, M. *et al.* Dysbiosis of the faecal microbiota in patients with Crohn's disease and their unaffected relatives. *Gut* **60**, 631-637, doi:10.1136/gut.2010.223263 (2011).

20 Feng, Q. *et al.* Gut microbiome development along the colorectal adenoma-carcinoma sequence. *Nat Commun* **6**, 6528, doi:10.1038/ncomms7528 (2015).

21 Louis, P., Hold, G. L. & Flint, H. J. The gut microbiota, bacterial metabolites and colorectal cancer. *Nat Rev Microbiol* **12**, 661-672, doi:10.1038/nrmicro3344 (2014).

22 Aron-Wisniewsky, J., Gaborit, B., Dutour, A. & Clement, K. Gut microbiota and non-alcoholic fatty liver disease: new insights. *Clin Microbiol Infect* **19**, 338-348, doi:10.1111/1469-0691.12140 (2013).

23 Gkolfakis, P., Dimitriadis, G. & Triantafyllou, K. Gut microbiota and non-alcoholic fatty liver

477 disease. *Hepatobiliary Pancreat Dis Int* **14**, 572-581 (2015).

478 24 Ríos-Covián, D. *et al.* Intestinal Short Chain Fatty Acids and their Link with Diet and Human
479 Health. *Frontiers in Microbiology* **7**, doi:10.3389/fmicb.2016.00185 (2016).

480 25 Puertollano, E., Kolida, S. & Yaqoob, P. Biological significance of short-chain fatty acid
481 metabolism by the intestinal microbiome. *Curr Opin Clin Nutr Metab Care* **17**, 139-144,
482 doi:10.1097/MCO.000000000000025 (2014).

483 26 Quevrain, E. *et al.* Identification of an anti-inflammatory protein from *Faecalibacterium*
484 *prausnitzii*, a commensal bacterium deficient in Crohn's disease. *Gut* **65**, 415-425,
485 doi:10.1136/gutjnl-2014-307649 (2016).

486 27 Sokol, H. *et al.* *Faecalibacterium prausnitzii* is an anti-inflammatory commensal bacterium
487 identified by gut microbiota analysis of Crohn disease patients. *Proc Natl Acad Sci U S A* **105**,
488 16731-16736, doi:10.1073/pnas.0804812105 (2008).

489 28 Duncan, S. H., Hold, G. L., Harmsen, H. J., Stewart, C. S. & Flint, H. J. Growth requirements
490 and fermentation products of *Fusobacterium prausnitzii*, and a proposal to reclassify it as
491 *Faecalibacterium prausnitzii* gen. nov., comb. nov. *Int J Syst Evol Microbiol* **52**, 2141-2146,
492 doi:10.1099/00207713-52-6-2141 (2002).

493 29 Fitzgerald, C. B. *et al.* Comparative analysis of *Faecalibacterium prausnitzii* genomes shows a
494 high level of genome plasticity and warrants separation into new species-level taxa. *BMC*
495 *Genomics* **19**, 931, doi:10.1186/s12864-018-5313-6 (2018).

496 30 Chang Liu, M.-X. D., Rexiding Abuduaini, Hai-Ying Yu, Dan-Hua Li, Yu-Jing Wang, Nan
497 Zhou, Min-Zhi Jiang, Peng-Xia Niu, Shan-Shan Han, Hong-He Chen, Wen-Yu Shi, Linhuan
498 Wu, Yu-Hua Xin, Juncai Ma, Yuguang Zhou, Cheng-Ying Jiang, Hong-Wei Liu, Shuang-Jiang
499 Liu. Enlightening the Taxonomy Darkness of Human Gut Microbiomes With Cultured
500 Biobank. *Research Square*, doi:10.21203/rs.3.rs-74101/v1 (2020).

501 31 Miquel, S. *et al.* *Faecalibacterium prausnitzii* and human intestinal health. *Curr Opin*
502 *Microbiol* **16**, 255-261, doi:10.1016/j.mib.2013.06.003 (2013).

503 32 Rossi-Tamisier, M., Fournier, P.-E., Benamar, S. & Raoult, D. Cautionary tale of using 16S
504 rRNA gene sequence similarity values in identification of human-associated bacterial species.
505 *International Journal of Systematic and Evolutionary Microbiology* **65**, 1929-1934,
506 doi:10.1099/ijs.0.000161 (2015).

507 33 Vital, M., Howe, A. C. & Tiedje, J. M. Revealing the bacterial butyrate synthesis pathways by
508 analyzing (meta)genomic data. *mBio* **5**, e00889, doi:10.1128/mBio.00889-14 (2014).

509 34 Zou, Y. *et al.* 1,520 reference genomes from cultivated human gut bacteria enable functional
510 microbiome analyses. *Nat Biotechnol* **37**, 179-185, doi:10.1038/s41587-018-0008-8 (2019).

511 35 Zhou, L. *et al.* *Faecalibacterium prausnitzii* Produces Butyrate to Maintain Th17/Treg Balance
512 and to Ameliorate Colorectal Colitis by Inhibiting Histone Deacetylase 1. *Inflamm Bowel Dis*
513 **24**, 1926-1940, doi:10.1093/ibd/izy182 (2018).

514 36 Fujimoto, T. *et al.* Decreased abundance of *Faecalibacterium prausnitzii* in the gut microbiota
515 of Crohn's disease. *J Gastroenterol Hepatol* **28**, 613-619, doi:10.1111/jgh.12073 (2013).

516 37 Machiels, K. *et al.* A decrease of the butyrate-producing species *Roseburia hominis* and
517 *Faecalibacterium prausnitzii* defines dysbiosis in patients with ulcerative colitis. *Gut* **63**,
518 1275-1283, doi:10.1136/gutjnl-2013-304833 (2014).

519 38 Lopez-Siles, M. *et al.* Alterations in the Abundance and Co-occurrence of *Akkermansia*
520 *muciniphila* and *Faecalibacterium prausnitzii* in the Colonic Mucosa of Inflammatory Bowel

521 Disease Subjects. *Front Cell Infect Microbiol* **8**, 281, doi:10.3389/fcimb.2018.00281 (2018).

522 39 Chang, C. J. *et al.* Next generation probiotics in disease amelioration. *J Food Drug Anal* **27**,
523 615-622, doi:10.1016/j.jfda.2018.12.011 (2019).

524 40 De Filippis, F., Pasolli, E. & Ercolini, D. Newly Explored Faecalibacterium Diversity Is
525 Connected to Age, Lifestyle, Geography, and Disease. *Curr Biol*,
526 doi:10.1016/j.cub.2020.09.063 (2020).

527 41 Tittsler, R. P. & Sandholzer, L. A. The Use of Semi-solid Agar for the Detection of Bacterial
528 Motility. *J Bacteriol* **31**, 575-580 (1936).

529 42 Cheng, H. R. & Jiang, N. Extremely rapid extraction of DNA from bacteria and yeasts.
530 *Biotechnol Lett* **28**, 55-59, doi:10.1007/s10529-005-4688-z (2006).

531 43 Zou, Y. *et al.* Lactobacillus shenzhenensis sp. nov., isolated from a fermented dairy beverage.
532 *Int J Syst Evol Microbiol* **63**, 1817-1823, doi:10.1099/ijs.0.041111-0 (2013).

533 44 Yoon, S. H. *et al.* Introducing EzBioCloud: a taxonomically united database of 16S rRNA
534 gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* **67**, 1613-1617,
535 doi:10.1099/ijsem.0.001755 (2017).

536 45 Saitou, N. & Nei, M. The neighbor-joining method: a new method for reconstructing
537 phylogenetic trees. *Mol Biol Evol* **4**, 406-425 (1987).

538 46 Felsenstein, J. Evolutionary trees from DNA sequences: a maximum likelihood approach. *J*
539 *Mol Evol* **17**, 368-376 (1981).

540 47 Rzhetsky, A. & Nei, M. Theoretical foundation of the minimum-evolution method of
541 phylogenetic inference. *Mol Biol Evol* **10**, 1073-1095 (1993).

542 48 Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary
543 Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**, 1547-1549,
544 doi:10.1093/molbev/msy096 (2018).

545 49 Felsenstein, J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*
546 **39**, 783-791, doi:10.2307/2408678 (1985).

547 50 Luo, R. *et al.* SOAPdenovo2: an empirically improved memory-efficient short-read de novo
548 assembler. *Gigascience* **1**, 18, doi:10.1186/2047-217X-1-18 (2012).

549 51 Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
550 assessing the quality of microbial genomes recovered from isolates, single cells, and
551 metagenomes. *Genome Res* **25**, 1043-1055, doi:10.1101/gr.186072.114 (2015).

552 52 Grant, J. R. & Stothard, P. The CGView Server: a comparative genomics tool for circular
553 genomes. *Nucleic Acids Res* **36**, W181-184, doi:10.1093/nar/gkn179 (2008).

554 53 Aziz, R. K. *et al.* The RAST Server: rapid annotations using subsystems technology. *BMC*
555 *Genomics* **9**, 75, doi:10.1186/1471-2164-9-75 (2008).

556 54 Galperin, M. Y., Makarova, K. S., Wolf, Y. I. & Koonin, E. V. Expanded microbial genome
557 coverage and improved protein family annotation in the COG database. *Nucleic Acids Res* **43**,
558 D261-269, doi:10.1093/nar/gku1223 (2015).

559 55 Zhou, Y., Liang, Y., Lynch, K. H., Dennis, J. J. & Wishart, D. S. PHAST: a fast phage search
560 tool. *Nucleic Acids Res* **39**, W347-352, doi:10.1093/nar/gkr485 (2011).

561 56 Jia, B. *et al.* CARD 2017: expansion and model-centric curation of the comprehensive
562 antibiotic resistance database. *Nucleic Acids Res* **45**, D566-D573, doi:10.1093/nar/gkw1004
563 (2017).

564 57 Richter, M. & Rossello-Mora, R. Shifting the genomic gold standard for the prokaryotic

- species definition. *Proc Natl Acad Sci U S A* **106**, 19126-19131, doi:10.1073/pnas.0906412106 (2009).
- 58 Goris, J. *et al.* DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* **57**, 81-91, doi:10.1099/ijs.0.64483-0 (2007).
- 59 Kim, M., Oh, H. S., Park, S. C. & Chun, J. Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. *Int J Syst Evol Microbiol* **64**, 346-351, doi:10.1099/ijs.0.059774-0 (2014).
- 60 Jain, C., Rodriguez, R. L., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* **9**, 5114, doi:10.1038/s41467-018-07641-9 (2018).
- 61 Guo, X. *et al.* CNSA: a data repository for archiving omics data. *Database (Oxford)* **2020**, doi:10.1093/database/baaa055 (2020).
- 62 Chen, F. Z. *et al.* CNGBdb: China National GeneBank DataBase. *Yi Chuan* **42**, 799-809, doi:10.16288/j.ycz.20-080 (2020).

Acknowledgment

This work was supported by grants from National Key Research and Development Program of China (No. 2018YFC1313801) and Natural Science Foundation of Guangdong Province, China (No. 2019B020230001). We also thank the colleagues at BGI-Shenzhen for sample collection, and discussions, and China National Genebank (CNGB) Shenzhen for DNA extraction, library construction, and sequencing.

Author contributions

Conceived and designed the experiments: Y.Z. and L.X. Performed the experiments: Y.Z., W.X., and Y.D. Analyzed the data: Y.Z., L.X., and X.L. Contributed reagents/materials/analysis tools: Y.Z., W.X. and Y.D. Wrote the paper: Y.Z. and X.L. Revised the paper: K.K.

Data Availability Statement

The GenBank/EMBL/DDBJ accession numbers for the 16S rRNA gene sequences determined in this study are: AF52-21^T (KX146426) and CM04-06^T (KX150462). The data of draft genome

594 sequences have been deposited into CNGB Sequence Archive (CNSA) ⁶¹ of China National
595 GeneBank DataBase (CNGBdb) ⁶² with accession number CNA0017730 and CNA0017731 for
596 strains AF52-21^T and CM04-06^T, respectively.

597

Figure Legends

Figure 1. Micrographs of strains AF52-21^T, CM04-06^T after Gram staining.

A, AF52-21^T; B, CM04-06^T.

Figure 2. Neighbour-joining phylogenetic tree based on 16S rRNA gene sequences showing the phylogenetic relationships of strains AF52-21^T, CM04-06^T and the representatives of several other related taxa within the family *Ruminococcaceae*.

Clostridium butyricum DSM 10702^T (AQQF01000149) was used as an out-group. Bootstrap values based on 1000 replications higher than 70% are shown at the branching points. Bar, substitutions per nucleotide position.

Figure 3. Circular map of AF52-21^T.

Innermost circle, GC skew; circle 2, G+C content; circle 3, contigs; circles 4, predicted prophage remnants; circle 5, tmRNA, tRNA and rRNA genes; circles 6, CDS; circles 7-9, homologous genomic segments from CM04-06^T, *F. prausnitzii* ATCC 27768^T and *F. hominis* 4P-15^T.

Figure 4. Circular map of CM04-06^T.

Innermost circle, GC skew; circle 2, G+C content; circle 3, contigs; circles 4, predicted prophage remnants; circle 5, tmRNA, tRNA and rRNA genes; circles 6, CDS; circles 7-9, homologous genomic segments from AF52-21^T, *F. prausnitzii* ATCC 27768^T and *F. hominis* 4P-15^T.

Figure 5. Comparison of COG functional categories of strains AF52-21^T, CM04-06^T and the

620 closest related species *F. prausnitzii* ATCC 27768^T.

621

622 **Figure 6. The synthesis pathways from Acetyl-CoA to Butyrate. Strains AF52-21^T,**
623 **CM04-06^T and ATCC 27768^T were annotated as blue, red, and yellow.**

624 Thl, thiolase; Hdb, β-hydroxybutyryl-CoA dehydrogenase; Cro, crotonase; Bcd, butyryl-CoA
625 dehydrogenase; But, butyryl-CoA:acetate CoA transferase; Ptb, phosphate butyryltransferase; Buk,
626 butyrate kinase.

627

628 **Figure 7. Distribution of prophage in strains AF52-21T and CM04-06T.**

629

630 **Figure 8. Comparison of resistance genes in strains AF52-21^T, CM04-06^T, and *F.***
631 ***prausnitzii* ATCC 27768^T.**

632

633

634

635

636

637

638

639

640

641

642 **Figure 1**

643

644

645 **Figure 2**

646

647

648 **Figure 3**

Length:2,851,918bp

649

650

651 **Figure 4**

653 **Figure 5**

654

655

656 **Figure 6**

657

658

659 **Figure 7**

660

661

662 **Figure 8**

663

664

665 **Supplementary Material**

666 **Supplementary Table S1. Number of genes associated with general COG functional**
 667 **categories in the genome of *F. butyricigenerans* AF52-21^T and *F. longum* CM04-06^T.**

668 **Supplementary Table S2. The specific genes/protein related to biosynthesis of DAP, polar**
 669 **lipids, polyamines and lipoteichoic and teichoic acids and their positions in the genome in**
 670 **comparison of strains AF52-21^T, CM04-06^T and related organism, ATCC 27768^T identified**
 671 **by Rapid Annotation Subsystem Technology (RAST).**

672
 673 **Supplementary Figure S1. Maximum-likelihood phylogenetic tree based on 16S rRNA gene**
 674 **sequences showing the phylogenetic relationships of strains AF52-21^T, CM04-06^T and the**
 675 **representatives of related taxa. *Clostridium butyricum* DSM 10702^T (AQQF01000149) was used**
 676 **as an out-group. Bootstrap values based on 1000 replications higher than 70% are shown at the**
 677 **branching points. Bar, substitutions per nucleotide position.**

678
 679 **Supplementary Figure S2. Minimum-evolution phylogenetic tree based on 16S rRNA gene**
 680 **sequences showing the phylogenetic relationships of strains AF52-21^T, CM04-06^T and the**
 681 **representatives of related taxa. *Clostridium butyricum* DSM 10702^T (AQQF01000149) was used**
 682 **as an out-group. Bootstrap values based on 1000 replications higher than 70% are shown at the**
 683 **branching points. Bar, substitutions per nucleotide position.**

684
 685 **Supplementary Figure S3. Certification. Deposit certification of AF52-21^T in CGMCC.**

686 **Supplementary Figure S4. Certification. Deposit certification of AF52-21^T in DSMZ.**

687 **Supplementary Figure S5. Certification. Deposit certification of CM04-06^T in CGMCC.**

688 **Supplementary Figure S6. Certification. Deposit certification of CM04-06^T in DSMZ.**