1 Zebrafish cooperate while inspecting predators: experimental evidence

2 for conditional approach

3 Ana Flávia Nogueira Pimentel¹, Monica Gomes Lima-Maximino², Marta Candeias
4 Soares³, Caio Maximino^{1*}

5

6 1 – Laboratório de Neurociências e Comportamento “Frederico Guilherme Graeff”,
7 Faculdade de Psicologia, Instituto de Estudos em Saúde e Biológicas, Universidade
8 Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil

9 2 – Laboratório de Neurofarmacologia e Biofísica, Centro de Ciências Biológicas e da
10 Saúde, Universidade do Estado do Pará – Campus VIII, Marabá/PA, Brazil

11 3 – CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos,
12 Universidade do Porto, Vairão, Portugal

13

14 * Corresponding author

15 Caio Maximino

16 Laboratório de Neurociências e Comportamento – Instituto de Estudos em Saúde e
17 Biológicas, Universidade Federal do Sul e Sudeste do Pará, Unidade III

18 Av. dos Ipês, S/N, s/ CEP, Bairro Cidade Jardim, Marabá/PA, Brazil

19 cmaximino@unifesspa.edu.br

Different fish species employ a conditional approach strategy during predator inspection; the risk of approaching a predator is distributed across all inspectors, but is not shared with the animals which keep its distance. Zebrafish, a highly social fish, is increasingly being used in behavioural neuroscience, but it is not known whether it displays conditional approach. In the predator inspection task, animals are observed in a tank with a refuge in one extremity, and an animated image of a predator in the other extremity, with a mirror positioned in parallel to the tank, simulating a perfectly reciprocating conspecific. In Experiment 1, animals spent more time **in an inspection zone** when the image was turned on, but also displayed more erratic swimming, suggesting cooperation under fear. In Experiment 2, animals spent more time inspecting predators when the mirror was parallel to the tank (“cooperating mirror”) than when the mirror was in an angle (“defecting mirror”), suggesting retaliatory behaviour; in both conditions, animals displayed more freezing and erratic swimming. **In Experiment 3, no changes in behaviour were observed, regardless of mirror position, when the image was turned off, suggesting that the choice of specific zones in Experiment 2 was not due to shoaling tendencies.** These results suggest that predator inspection is associated with conditional approach, while at the same time inducing fear-like behaviour in the animal.

Keywords: Predator inspection; Altruism; Zebrafish; Social behaviour

1. Introduction

While being widespread among animals, **altruism** is an adaptive/evolutionary conundrum, as it may involve reducing one’s own fitness in order to benefit another individual (Kropotkin, 1902). Different explanations have been given to this phenomenon, including kin selection (W. D. Hamilton, 1963, 1964) and reciprocal altruism (Trivers, 1971). Reciprocal altruism predicts that **this type of cooperative behaviour** only occurs in populations when it increases the likelihood of reciprocation at the receiving end – that is, the individual decreases its own fitness in the expectation that the other organism will act in a similar manner in a later time (Trivers, 1971). Game theoretical approaches suggest a “tit-for-tat” strategy, in which an individual cooperates in the first encounter and continues to cooperate as long as the other agent does as well (Trivers, 1971).

Experimental evidence of “tit-for-tat” has been hard to produce because the payoff (i.e., possible positive and negative outcomes) of cooperating vs. defecting is known, but the benefits (and risks) cooperating are usually not known. Nonetheless, “reciprocal altruism-like” behaviour has been observed in many species (Bshary & Oliveira, 2015). An interesting example is conditional approach, a strategy used by social fish when approaching a potential predator, leaving the shoal to obtain information on putative threat and/or try to intimidate the dangerous predator (Dugatkin, 2013). For two interacting fish approaching a predator, defecting (staying behind) is beneficial because the defecting individual decreases its risk but increases its payoff (e.g., information gathered) by watching the fate of the other fish. Conditional approach predicts that inspecting fish should immediately retaliate defecting conspecifics by also defecting (Bshary & Oliveira, 2015; Dugatkin, 1988; Pimentel et al., 2019). The strategy appears to function as an incentive for predator inspection, given that the risk of approaching it is distributed between all inspectors but not shared with those that remain at a distance. Since this behaviour implies costs to the individual(s) interacting with the predator, but contributes to the shoal’s fitness, it has been interpreted as cooperative-like and altruistic (Dugatkin, 1988, 1997; Dugatkin & Alfieri, 1991b, 1992; Pimentel et al., 2019). When choosing whether or not to inspect a predator, individuals leaving the shoal are exposed to increased predation risk (Dugatkin, 1992; Milinski et al., 1997), and therefore are expected to display threat-sensitive behaviours such as freezing and erratic swimming. In guppies, predator inspection is accompanied by increased freezing duration, suggesting that this is indeed the case (Pimentel et al., 2019).

In conditional approach, prey approaching predators inspect the predator at the first “move”, immediately head back away from the predator if the conspecific falls behind or disappears, and approach the predator again if the conspecific’s move of “swimming away and staying behind” is followed by “swim parallel” (Dugatkin, 1988). However, differently from tit-for-tat (Axelrod & Hamilton, 1981), conditional approach does not require that the actual payoff matrix is known (Dugatkin, 1988). Conditional approach has been shown in sticklebacks (*Gasterosteus aculeatus* Linnaeus 1758) (Huntingford et al., 1994; Milinski, 1987) and guppies (*Poecilia reticulata* Peter 1859) (Dugatkin, 1988, 1991; Dugatkin & Alfieri, 1991b, 1992; Pimentel et al., 2019), but not for zebrafish, an important model organism in the neurosciences.

Zebrafish (*Danio rerio* Hamilton 1822) is a model organism that is widely used in developmental biology (Parichy, 2015) and which has been introduced in the field of behavioural neuroscience and behavioural ecology (Bonan & Norton, 2015; Stewart et al., 2015). This fish is also a good model organism for biological psychiatry (Stewart et al., 2015), due to the availability of behavioural bioassays and the advantages associated with being a model organism, including low cost of acquisition and maintenance, easy handling and housing, short lifespan, and easy reproduction in the laboratory. As a highly social animal that lives in groups with well-structured social relations – including marginalization, dominance hierarchies, and territoriality (Spence et al., 2008) –, zebrafish are also known to display predator inspection (Dugatkin et al., 2010; Pannia et al., 2014); however, it is not known whether zebrafish employ conditional approach when inspecting predators.

In addition to helping to better understand complex social behaviour in this species, here we aim to introduce new behavioural techniques that can also advance research in the field of neuropsychopharmacology, including the development of drugs to treat disorders of social behaviour (Soares et al., 2018). While the focus of this field on zebrafish social behaviour has relied mainly on shoaling, more complex behaviours, such as cooperative-like behaviour, could represent an important addition to the field. Our main aim was to investigate preliminary evidence of conditional approach in zebrafish in a predator inspection model, based in Milinski (1987) and Dugatkin (1988). In Experiment 1, a mirror was placed in parallel to a tank, and the animated image of a predator (Gangetic leaffish *Nandus nandus* Hamilton 1822): when the animation was turned on, animals spent more time in the inspection area than when it was turned off, suggesting that the presence of both a predator and a cooperating conspecific (mirror image) are necessary for inspection behaviour. In Experiment 2, the mirror was either placed in parallel (“cooperating mirror”) or in an angle of 45° with the side of the tank which was opposite to the predator (“defecting mirror”): animals then spent less time in the inspection zone during the defecting mirror condition. In Experiment 3, animals were exposed to conditions of Experiment 2, but the video was not turned on. Overall, animals displayed freezing and erratic swimming when the animation was visible, suggesting fear. This manuscript is the first complete report of all the studies performed to test the hypothesis that zebrafish display conditional approach during predator inspection. We report all data exclusions (if any), all manipulations, and all measures in the study.

118

119 2. Methods

120 2.1. Animals and housing

121 31 animals were used in Experiments 1 and 2 described below; data from one
 122 animal was excluded from each experiment due to poor health after the end of the
 123 experiments, which could impact the results. In Experiment 3, 32 animals were
 124 used. Thus, a total of 94 animals were used across all three experiments.
 125 Animals were bought from a commercial vendor, and arrived in the laboratory with an
 126 approximate age of 3 months (standard length = 13.2 ± 1.4 mm), and were quarantined
 127 for two weeks; the experiment began when animals had an approximate age of 4 months
 128 (standard length = 23.0 ± 3.2 mm). Animals were kept in mixed-sex tanks during
 129 acclimation, with an approximate sex ratio of 1:1 (confirmed by body morphology).
 130 Adult zebrafish from the wildtype strain (longfin phenotype) were used in the
 131 experiments. Outbred populations were used due to their increased genetic variability,
 132 decreasing the effects of random genetic drift which could lead to the development of
 133 uniquely heritable traits (Parra et al., 2009; Speedie & Gerlai, 2008). Thus, the animals
 134 used in the experiments are expected to better represent the natural populations in the
 135 wild. The breeder was licensed for aquaculture under Ibama's (Instituto Brasileiro do
 136 Meio Ambiente e dos Recursos Naturais Renováveis) Resolution 95/1993. Animals
 137 were group-housed in 40 L tanks, with a maximum density of 25 fish per tank, for at
 138 least 2 weeks before experiments begun. Tanks were filled with non-chlorinated water
 139 at room temperature (28 °C) and a pH of 7.0-8.0 (coefficient of variation = 5.01%).
 140 Lighting was provided by fluorescent lamps in a cycle of 14-10 hours (LD), according
 141 to standards of care for zebrafish (Lawrence, 2007). Water quality parameters were as
 142 follows: pH 7.0-8.0; hardness 100-150 mg/L CaCO₃; dissolved oxygen 7.5-8.0 mg/L;
 143 ammonia and nitrite < 0.001 ppm. All manipulations minimized their potential suffering
 144 of animals, and followed Brazilian legislation (Diretriz Brasileira Para o Cuidado e a
 145 Utilização de Animais Para Fins Científicos e Didáticos - DBCA. Anexo I. Peixes
 146 Mantidos Em Instalações de Instituições de Ensino Ou Pesquisa Científica, 2017).
 147 Animals were used for only one experiment and in a single behavioural test, to reduce
 148 interference from apparatus exposure. Experiments were approved by UEPA's IACUC
 149 under protocol 06/18.

150

151 2.2. Sample size determination

152 Sample sizes were determined based on results regarding conditional approach
153 in guppies(Pimentel et al., 2019). Time on the inspection zone was chosen as the
154 primary endpoint, and calculations for sample sizes are valid only for that endpoint.
155 Calculations were based on Rosner's (2016) method for comparing two means(Rosner,
156 2016), and assumed $\alpha = 0.05$ and power 80% on a two-tailed analysis. Based on these
157 calculations, 15 animals were used in each group in both experiments. Animals were
158 randomly drawn from the tank immediately before testing, and the order with which
159 they were allocated to different conditions was randomized *via* generation of random
160 numbers using the randomization tool in <http://www.randomization.com/>. Blinding was
161 not possible, due to the positioning of the camera.

162

163 2.3. Experimental apparatus

164 The experimental apparatus was based on that described by Pimentel et al.
165 (2019) for guppies (Pimentel et al., 2019) (Fig. 1A-B), consisting in a glass tank with
166 40.6 cm x 18 cm x 25 cm (l X w X h). The bottom of the tank was divided in 10
167 equally-sized quadrants; the quadrant that was farther from the side in which the
168 stimulus was presented was numbered "section 1" (refuge zone), and an artificial plant
169 was placed there to serve as refuge. Parallel to the quadrant that was farther from
170 section 1, a computer screen (Samsung T20c310lb, 20", LED screen, nominal
171 brightness 20 cd/m³) allowed the presentation of animated images. These animations
172 were produced on LibreOffice Impress (Pimentel et al., 2019), and consisted of a
173 moving image of a leaffish (*Nandus nandus*), a sympatric predator for zebrafish
174 (Engeszer et al., 2007; Parichy, 2015) that has been shown to induce antipredator
175 behavior (Bass & Gerlai, 2008; Gerlai et al., 2009). Parallel to one of the longer walls of
176 the tank (Fig. 1B), a mirror, with the length of 40.6 cm, was positioned. The mirror,
177 therefore, occupied the whole length of the experimental apparatus and, as a
178 consequence, created an image of a second zebrafish swimming in parallel to the focal
179 animal. In Experiments 2 and 3, the mirror was either parallel to the tank (Cooperating
180 mirror) or forming an angle of 45° with the tank (Defecting mirror).

181

182 2.4. General design

183 Animals were tested individually and only once. Before beginning a trial, animals were
 184 individually transferred to the experimental tank, placed on quadrant 5 (in the center of
 185 the tank), and behaviour was filmed for 5 min while swimming freely. During this
 186 acclimation period, the screen was turned off. Animals were randomly allocated to
 187 groups, using the random number generator at <https://randomization.com>, to reduce
 188 allocation bias. Moreover, the order with which individuals from each group were tested
 189 was also randomized. Each trial took 15 min, and consisted in an acclimation stage
 190 (min. 0-5), immediately after the animal was introduced in the tank and during which
 191 the video was turned off, followed by the stimulation stage (min. 6-15), during which
 192 the video was turned on. Trials were recorded in digital format by a camera (Sony DCR-
 193 DVD610) mounted above the tank. Given the position of the camera in relation to the
 194 mirror, it was not possible to blind video transcription to treatment. Digital videos were
 195 later viewed, and behaviour was manually recorded using XPlo-Rat
 196 (<http://scotty.ffclrp.usp.br>). The following variables were recorded:

- 197 • **Time in the inspection zone:** Time spent in Sections 6 to 10 (nearest to the
 198 computer screen), expressed as percent change (time after stimulus onset – time
 199 before stimulus onset) and normalized by time block duration (i.e., time after
 200 stimulus onset was scaled to 600 s, and time before stimulus onset was scaled to
 201 300 s);
- 202 • **Time in the avoidance zone:** Time spent in Sections 2 to 5, expressed as
 203 percent change and normalized by time block duration;
- 204 • **Time in the refuge:** Time spent in Section 1, expressed as percent change and
 205 normalized by time block duration;
- 206 • **Freezing:** Time spent immobile, except for eye and opercular movements;
- 207 • **Erratic swimming:** Number of events of zig-zagging, fast swimming episodes.

208

209 2.5. Experiment 1

Experiment 1 was a conceptual replication of experiments in guppies (Pimentel et al., 2019), and attempted to understand whether approach and defect behaviours were potential responses to the specific predator inspection context, or whether the animal was simply responding to the conspecific (i.e., following its own mirror image). In the “Predator” condition, the animation was shown after 5 minutes of acclimation, as described in “General design”, above; in the “No-predator” condition, no animation was shown.

2.6. Experiment 2

Experiment 2 was a conceptual replication of studies on cooperation vs. defection made on guppies (Dugatkin, 1988) and sticklebacks (Milinski, 1987). The rationale for the experiment is that, for two interacting fish approaching a predator, defecting (staying behind) is beneficial because the defecting individual decreases its risk but increases its payoff (e.g., information gathered) by watching the fate of the other fish. Conditional approach predicts that inspecting fish should immediately retaliate defecting conspecifics by also defecting (Bshary & Oliveira, 2015; Dugatkin, 1988; Pimentel et al., 2019). We attempt to simulate this situation by using two mirror conditions, a “cooperating” mirror (parallel to the tank) and a “defecting” mirror (in an angle of 45° to the tank); in this last situation, as the fish approached the predator, its mirror image would appear to turn away.

2.7. Experiment 3

Experiment 3 was an attempt to untangle shoaling tendencies from results from Experiment 2. The setup and design was exactly as in Experiment 2, except that no predator animation was shown to either mirror group.

2.8. Statistical analysis

For all experiments, data for time on the inspection, avoidance, and refuge zones were assessed using independent samples *t*-tests. For data on freezing and erratic swimming, one-way repeated measures analyses of variance (ANOVAs) were made,

with subject as random variable. Bayes Factors (BF) were also calculated, using the ‘BayesFactor’ package in R (Morey & Rouder, 2018) (v. 0.9.12; <https://richarddmorey.github.io/BayesFactor/>). The prior is described by a Cauchy distribution centred around zero and with a width parameter of 0.707. This corresponds to a probability of 80% that the effect size lies between -1.25 and 1.25 for changes in time in inspection zone (prior = 0.672); between -0.49 and 0.49 for changes in time in the avoidance zone (prior = 0.385); between -0.82 and 0.82 for changes in time in the refuge zone (prior = 0.547); and between -1.36 and 1.36 for freezing (prior = 0.695). These effect sizes were based on previous work on guppies (Pimentel et al., 2019). Bayes factors were reported alongside frequentist statistics under the assumption that this can help clarify results that may be difficult to interpret using frequentist statistics alone (Malone & Coyne, 2020). *A posteriori* effect sizes were calculated as estimates of Cohen’s *d* using a bootstrap sampling distribution, with 5000 bootstrap samples, and reported with bias-corrected and accelerated confidence intervals; these calculations were made using the ‘dabestr’ package in R (Ho et al., 2018) (v. 0.2.1; <https://github.com/ACCLAB/dabestr>).

2.8. Open science practices

Experiments were not formally pre-registered. Datapackages and analysis scripts can be found on GitHub (<https://github.com/lanec-unifesspa/zebrafishTFT/>). Preprint versions of this paper have been submitted to bioRxiv (doi:10.1101/814434).

3. Results

In Experiment 1, a significant difference was found for time spent in inspection zone (Fig. 2A), with animals in the Predator condition spending more time in that area than animals in the No-Predator condition ($t(17.410) = -3.065$, $p = 0.00687$, $d = 1.13$ [95%CI 0.233, 1.99], BF = 10.623). No differences were found for time in the avoidance zone (Fig. 2B; $t(21.219) = -1.097$, $p = 0.28746$, $d = 0.402$ [95%CI -0.324, 1.17], BF = 0.546). Likewise, no differences were found for time in the refuge zone (Fig. 2C; $t(26.968) = -1.291$, $p = 0.2076$, $d = 0.467$ [95%CI -0.396, 1.13], BF = 0.644). Freezing was increased after animation in animals in the Predator condition compared to

animals in the No-Predator condition (main effects of group: $F[1, 29] = 4.606$, $p = 0.0404$, $BF = 1.313$; main effects of time block: $F[1, 29] = 4.983$, $p = 0.0445$, $BF = 2.239$; interaction: $F[1, 29] = 2.689$, $p = 0.119$, $BF = 3.33$; d for Non-predator condition = 0.446 [95%CI $-0.238, 1.02$]; d for Predator condition = 0.637 [95%CI $-0.207, 1.07$]); while the RM-ANOVA was not significant for the interaction effect, the higher Bayes Factor suggests that the interaction model is preferred against all models.

In Experiment 2, a significant difference was found for time spent in inspection zone (Fig. 3A), with animals in the Cooperating Mirror condition spending more time in this zone than animals in the Defecting Mirror condition ($t(29) = -2.572$, $p = 0.0155$, $d = 0.922$ [95%CI $0.187, 1.61$]). Again, no differences were found for time in the avoidance zone (Fig. 3B; $t(29) = -0.925$, $p = 0.3667$, $d = -0.34$ [95%CI $-1.17, 0.559$]). Moreover, no significant differences were found for time in the refuge zone (Fig. 3C; $t(29) = -0.866$, $p = 0.4$, $d = -0.311$ [95%CI $-0.965, 0.572$]). An effect of time block ($F[1, 29] = 30.126$, $p = 6.35 \times 10^{-6}$; BF against model with subject only = 257.202) was found for freezing, but no effect of group ($F[1, 29] = 0.855$, $p = 0.363$; BF against model with subject only = 0.32) or interaction ($F[1, 29] = 1.894$, $p = 0.179$; BF against model with subject only = 614.199) was found (paired d for Defecting Mirror = 0.546 [95%CI $0.139, 0.891$]; paired d for Cooperating Mirror = 0.789 [95%CI $0.383, 1.22$]). Bayes Factors suggest the superiority of a model in which time block is the main variable driving the variation in erratic swimming, An effect of time block ($F[1, 30] = 10.72$, $p = 0.00268$; BF against model with subject only = 39.967) was found for erratic swimming, but no effect of group ($F[1, 30] = 0.113$, $p = 0.739$; BF against model with subject only = 0.31) or interaction ($F[1, 30] = 1.72$, $p = 0.2$; BF against model with subject only = 9.259) was found (paired d for Defecting Mirror = 1.09 [95%CI $0.358, 1.67$]; paired d for Cooperating Mirror = 1.41 [95%CI $0.851, 1.85$]). Bayes Factors suggest the superiority of a model in which time block is the main variable driving the variation in erratic swimming and freezing, reinforcing the hypothesis that both cooperating and defecting mirror configurations induce fear-like behaviour.

In Experiment 3, no significant differences were found between groups for time spent in inspection zone (Fig. 4A), with animals in the Cooperating Mirror condition spending a similar amount of time in this zone as animals in the Defecting Mirror condition when the animation was not presented (Fig. 4A; $t_{(28)} = -0.499$, $p = 0.621$, $d = -$

0.182 [95%CI -0.898, 0.540], BF = 0.392). Again, no significant differences were found between groups for time in avoidance (Fig. 4B; $t_{(28)} = 0.789$, $p = 0.437$, $d = 0.288$ [95%CI -0.440, 1.007], BF = 0.534) and refuge (Fig. 4C; $t_{(28)} = -1.064$, $p = 0.296$, $d = -0.389$ [95%CI -1.112, 0.348], BF = 0.604). No main effects of time block ($F_{[1, 30]} = 0.142$, $p = 0.709$, BF against model with subject only = 0.271) or mirror position ($F_{[1, 30]} = 0.01$, $p = 0.921$, BF against model with subject only = 0.256) were found for freezing (Fig. 4D); no interaction effect was found as well ($F_{[1, 31]} = 0.191$, $p = 0.665$, BF against model with subject only = 0.07). Finally, no main effects of time block ($F_{[1, 30]} = 2.04$, $p = 0.164$, BF against model with subject only = 0.64) nor mirror position ($F_{[1, 30]} = 3.019$, $p = 0.093$, BF against model with subject only = 0.74) were found for erratic swimming (Fig. 4E); no interaction effect was found as well ($F_{[1, 31]} = 1.966$, $p = 0.171$, BF against model with subject only = 0.47). Thus, the data provide evidence against the hypothesis that, when the predator image is turned off, mirror position affects position in the tank (e.g., time in inspection or refuge zones) or fear-like responses.

4. Discussion

In Experiment 1, we found that the animal spends more time in the inspection zone than in other zones when the predator stimulus is present, suggesting that the presence of a predator in the concomitant presence of a conspecific induces predator inspection, while simultaneously observing that the individual has a fear response in the form of increased erratic swimming and freezing behaviour. In experiment 2 we observed that the animal spends more time in inspection in the cooperating mirror condition than in the defecting mirror condition, and in both conditions we observed that it had fear-like behaviour. In experiment 3, we observed no changes in behaviour in any mirror position when the video was turned off, thus discarding shoaling tendencies as a driver of social behaviour under these conditions. The results of these experiments suggest that zebrafish exhibits conditional approach behaviour in situations in which also elicit fear.

In the present experiments, animals behaved towards conspecifics by approaching the mirror, and towards the predator by keeping a distance and freezing and/or displaying erratic swimming; however, when both stimuli are present, the animal

emits predator inspection, entering the inspection zone and orienting towards the predator image. While it has been previously shown that visual contact with conspecifics decrease fear responses in zebrafish (Faustino et al., 2017), the presence of the conspecific image did not inhibit fear in the present experiment, as the animal still displays freezing and erratic swimming. We reported similar effects on guppies (Pimentel et al., 2019). Moreover, guppies selectively bred for high leading in a similar paradigm show no alterations in “boldness” in a simulated aerial predator task (Dimitriadou et al., 2019), suggesting that conditional approach is not related to less fear. However, field experiments show that predation risk is a factor in the evolution of predator inspection across natural guppy populations (Dugatkin & Alfieri, 1992; Edenbrow et al., 2017), suggesting that conditional approach is sensitive to threat levels.

In the cooperating mirror condition in Experiment 2, zebrafish spent more time inspecting the predator than fish in the defecting mirror condition, suggesting that zebrafish employs a “conditional approach” strategy. Similarly to “tit-for-tat”, in conditional approach the individual approaches the predator in the first movement, and subsequently approaches only if the second fish (the mirror image) swims along, retreating if the conspecific also retreated (Dugatkin, 1988, 1991, 1997). Results from our Experiment 2 suggest that the animal retaliates after defection (i.e., reduces inspection in the defecting mirror condition).

Zebrafish were thus shown to be able to change their strategy from avoiding to approaching predators when conspecifics are present. If in visual contact with a conspecific (their own reflection), they proceed to spend more time inspecting predator (cooperation mirror condition), a situation that significantly changed in the defecting mirror condition, as their company was absent. This looks like a typical “conditional approach” situation: conditionally, the individual swims towards the predator in the first movement, but subsequent close inspection only occurs if the second fish (the mirror image) swims along, retreating immediately if the accompanying conspecific also abandons (Dugatkin, 1988). This is an indication of the social conditions necessary the evolution of predator approach behaviour as a cooperative strategy in zebrafish, as in guppies and sticklebacks (Brosnan et al., 2003; Croft et al., 2006; Dugatkin & Alfieri, 1991a; Edenbrow et al., 2017; Huntingford et al., 1994): strong social bonds between conspecifics that enable their first risky move (cooperative behaviour) which can confer benefits to those inspecting and to the shoal (Bshary & Oliveira, 2015; Croft et al.,

2006; Dugatkin, 2013). However, if facing defecting shoal-mates, individuals may retaliate, ceasing inspection but are able to “forgive” if baseline conditions are restored. Our findings allow us to include zebrafish in the restrict group of teleost fish species capable of complex social decision making, responding to the quality of social bonds by being able to change strategy as conditions change.

Currently, the benefits for the inspecting zebrafish are unknown. Reciprocal altruism suggests that the benefit of any cooperative action involve increased probability of reciprocation in the future (Trivers, 1971) – that is, an individual would be incentivized to inspect a predator if a shoal-mate is more likely to inspect in the future (Edenbrow et al., 2017). **Reciprocal altruism can also evolve by generalized reciprocity, in which cooperation evolves in small groups when individuals base their decision on the outcome of previous interactions with anonymous partners (i.e., without the need for individual recognition and specific social memory) (Pfeiffer et al., 2005). Generalized reciprocity is thought to be favoured by contexts in which cooperation is more advantageous in a cooperative than in a noncooperative environment, leading to stable levels of cooperation when the decision to stay or leave a group co-evolves with the decision to cooperate (I. M. Hamilton & Taborsky, 2005).** Indeed, both Experiments 1 and 2 suggest that, as zebrafish are more likely to inspect when a “shoal-mate” (mirror image) is present, and when the shoal-mate reciprocates inspection (i.e., in the “cooperating”, but not in the “defecting” mirror). This does not discard, of course, other benefits of inspecting, although currently these have not been investigated in zebrafish. **Indeed, our results are also consistent with generalized reciprocity, in which simple decision rules such as “defect when encountering noncooperation” is sufficient to lead to stable cooperation levels (Aktipis, 2004).** In guppies, cooperative animals (i.e., animals with high levels of predator inspection) are preferred by conspecifics (Dugatkin & Alfieri, 1991a; Godin & Dugatkin, 1996), and assortative interactions and social selection are expected to play a role (Edenbrow et al., 2017). It has been previously suggested that conditional approach is better understood in a “social competence” approach (Bshary & Oliveira, 2015; Taborsky & Oliveira, 2012), and therefore does not require some form of kin or group selection to be at work (Pimentel et al., 2019). In this context, situational cues (such as presence or absence of females, or varying risk levels), together with other factors, allow individuals to assess the situation and make behavioural decisions.

Zebrafish has been previously shown to display predator inspection, at low levels, without the presence of conspecifics (Dugatkin et al., 2010; Pannia et al., 2014). Our results are the first demonstration that predator inspection is related to cooperative-like behaviour in zebrafish, even when the animal displays fear-like behaviour towards the predator. These results suggest that the social behaviour of zebrafish is more complex than previously thought, and can be further exploited in the study of the neural bases of altruism in an interesting model organism.

Acknowledgements

AFP was the recipient of a CNPq studentship.

References

- Aktipis, C. A. (2004). Know when to walk away: Contingent movement and the evolution of cooperation. *Journal of Theoretical Biology*, 231, 249–260. <https://doi.org/10.1016/j.jtbi.2004.06.020>
- Axelrod, R., & Hamilton, W. (1981). The evolution of cooperation. *Science*, 211, 1390–1396. <https://doi.org/10.1126/science.7466396>
- Bass, S. L. S., & Gerlai, R. (2008). Zebrafish (*Danio rerio*) responds differentially to stimulus fish: The effects of sympatric and allopatric predators and harmless fish. *Behavioural Brain Research*, 186, 107–117. <https://doi.org/10.1016/j.bbr.2007.07.037>
- Bonan, C. D., & Norton, W. H. J. (2015). The utility of zebrafish as a model for behavioural genetics. *Current Opinion in Behavioral Sciences*, 2(Table 1), 34–38. <https://doi.org/10.1016/j.cobeha.2014.07.003>
- Brosnan, S. F., Earley, R. L., & Dugatkin, L. A. (2003). Observational learning and predator inspection in guppies (<i>Poecilia reticulata</i>). *Ethology*, 109, 823–833.
- Bshary, R., & Oliveira, R. F. (2015). Cooperation in animals: Toward a game theory within the framework of social competence. *Current Opinion in Behavioral Sciences*, 3, 31–37. <https://doi.org/10.1016/j.cobeha.2015.01.008>
- Diretriz brasileira para o cuidado e a utilização de animais para fins científicos e didáticos - DBCA. Anexo I. Peixes mantidos em instalações de instituições de ensino ou pesquisa científica, Pub. L. No. Resolução Normativa n. 30/2016 (2017).

- 434 Croft, D. P., James, R., Thomas, P. O. R., Hathaway, C., Mawdsley, D., Laland, K. N.,
435 & Krause, J. (2006). Social structure and co-operative interactions in a wild
436 population of guppies (*Poecilia reticulata*). *Behavioral Ecology and Sociobiology*,
437 59, 644–650. <https://doi.org/10.1007/s00265-005-0091-y>
- 438 Dimitriadou, S., Croft, D. P., & Darden, S. K. (2019). Divergence in social traits in
439 Trinidadian guppies selectively bred for high and low leadership in a cooperative
440 context. *Scientific Reports*, 9, 17194. <https://doi.org/10.1038/s41598-019-53748-4>
- 441 Dugatkin, L. A. (1988). Do guppies play TIT FOR TAT during predator inspection
442 visits? *Behavioral Ecology and Sociobiology*, 23, 395–399.
443 <https://doi.org/10.1007/BF00303714>
- 444 Dugatkin, L. A. (1991). Dynamics of the TIT FOR TAT strategy during predator
445 inspection in the guppy (*Poecilia reticulata*). *Behavioral Ecology and*
446 *Sociobiology*, 29, 127–132. <https://doi.org/10.1007/BF00166487>
- 447 Dugatkin, L. A. (1992). Tendency to inspect predators predict mortality risk in the
448 guppy, *Poecilia reticulata*. *Behavioral Ecology*, 3, 124–128.
- 449 Dugatkin, L. A. (1997). The evolution of cooperation. *BioScience*, 47, 355–362.
- 450 Dugatkin, L. A. (2013). The evolution of risk-taking. *Cerebrum*, 1, 1–9.
- 451 Dugatkin, L. A., & Alfieri, M. (1991a). Guppies and the TIT FOR TAT strategy:
452 Preference based on past interaction. *Behavioral Ecology and Sociobiology*, 28,
453 243–246. <https://doi.org/10.1007/BF00175096>
- 454 Dugatkin, L. A., & Alfieri, M. (1991b). Tit-For-Tat in guppies (*Poecilia reticulata*): The
455 relative nature of cooperation and defection during predator inspection.
456 *Evolutionary Ecology*, 5, 300–309. <https://doi.org/10.1007/BF02214234>
- 457 Dugatkin, L. A., & Alfieri, M. S. (1992). Interpopulational differences in the use of the
458 Tit- For-Tat strategy during predator inspection in the guppy, *Poecilia reticulata*.
459 *Evolutionary Ecology*, 6, 519–526. <https://doi.org/10.1007/BF02270695>
- 460 Dugatkin, L. A., McCall, M. A., Gregg, R. G., Cavanaugh, A., Christensen, C., &
461 Unseld, M. (2010). Zebrafish (*Danio rerio*) exhibit individual differences in risk-
462 taking behavior during predator inspection. *Ethology, Ecology & Evolution*, 17,
463 77–81. <https://doi.org/10.1080/08927014.2005.9522617>
- 464 Edenbrow, M., Bleakley, B. H., Tyler, C. R., Ramnarine, I. W., & Croft, D. P. (2017).
465 The evolution of cooperation: Interacting phenotypes among social partners.
466 *American Naturalist*, 189, 630–643. <https://doi.org/10.1086/691386>
- 467 Engeszer, R. E., Patterson, L. B., Rao, A. A., & Parichy, D. M. (2007). Zebrafish in the
468 wild: a review of natural history and new notes from the field. *Zebrafish*, 4(1), 21–
469 40. <https://doi.org/10.1089/zeb.2006.9997>

- 470 Faustino, A. I., Tacão-Monteiro, A., Oliveira, R. F., Parichy, D. M., Landgraf, R., &
471 Herculano, A. M. (2017). Mechanisms of social buffering of fear in zebrafish.
472 *Scientific Reports*, 7, 44329. <https://doi.org/10.1038/srep44329>
- 473 Gerlai, R., Fernandes, Y., & Pereira, T. (2009). Zebrafish (*Danio rerio*) responds to the
474 animated image of a predator: Towards the development of an automated aversive
475 task. *Behavioural Brain Research*, 201, 318–324.
476 <https://doi.org/10.1016/j.bbr.2009.03.003>
- 477 Godin, J. G., & Dugatkin, L. A. (1996). Female mating preference for bold males in the
478 guppy, *Poecilia reticulata*. *Proceedings of the National Academy of Sciences of the*
479 *United States of America*, 93, 10262–10267.
- 480 Hamilton, I. M., & Taborsky, M. (2005). Contingent movement and cooperation evolve
481 under generalized reciprocity. *Proceedings of the Royal Society B: Biological*
482 *Sciences*, 272, 2259–2267. <https://doi.org/10.1098/rspb.2005.3248>
- 483 Hamilton, W. D. (1963). The evolution of altruistic behavior. *American Naturalist*, 97,
484 354–356.
- 485 Hamilton, W. D. (1964). The genetical evolution of social behaviour. I. *Journal of*
486 *Theoretical Biology*, 7, 1–16. [https://doi.org/10.1016/0022-5193\(64\)90038-4](https://doi.org/10.1016/0022-5193(64)90038-4)
- 487 Ho, J., Tumkaya, T., Aryal, S., Choi, H., & Claridge-Chang, A. (2018). Moving beyond
488 P values: Everyday data analysis with estimation plots. *BioRxiv Pre-Print*.
489 <https://doi.org/10.1101/377978>
- 490 Huntingford, F. A., Lazarus, J., Barrie, B. D., & Webb, S. (1994). A dynamic analysis
491 of cooperative predator inspection in sticklebacks. *Animal Behaviour*, 47, 413–423.
- 492 Kropotkin, P. (1902). *Mutual aid*. William Heinemann.
- 493 Malone, H. E., & Coyne, I. (2020). Complementing the P-value from null-hypothesis
494 significance testing with a Bayes factor from null-hypothesis Bayesian testing.
495 *Nurse Researcher*, 28. <https://doi.org/10.7748/nr.2020.e1756>
- 496 Milinski, M. (1987). TIT FOR TAT in sticklebacks and the evolution of cooperation.
497 *Nature*, 325, 433–435.
- 498 Milinski, M., Lüthi, J. H., Eggler, R., & Parker, G. A. (1997). Cooperation under
499 predation risk: experiments on costs and benefits. *Proceedings of the Royal Society*
500 *of London. Series B: Biological Sciences*, 264, 831–837.
501 <https://doi.org/10.1098/rspb.1997.0116>
- 502 Morey, R. D., & Rouder, J. N. (2018). *BayesFactor: Computation of Bayes Factors for*
503 *Common Designs* (0.9.12-4.2). <https://cran.r-project.org/package=BayesFactor>
- 504 Pannia, E., Tran, S., Rampersad, M., & Gerlai, R. (2014). Acute ethanol exposure
505 induces behavioural differences in two zebrafish (*Danio rerio*) strains: A time

506 course analysis. *Behavioural Brain Research*, 259, 174–185.
507 <https://doi.org/10.1016/j.bbr.2013.11.006>

508 Parichy, D. M. (2015). Advancing biology through a deeper understanding of zebrafish
509 ecology and evolution. *ELife*, 4, 1–11. <https://doi.org/10.7554/eLife.05635>

510 Parra, K. V, Adrian Jr, J. C., & Gerlai, R. (2009). The synthetic substance hypoxanthine
511 3-N-oxide elicits alarm reactions in zebrafish (*Danio rerio*). *Behavioural Brain*
512 *Research*, 205, 336–341. <https://doi.org/10.1016/j.bbr.2009.06.037>

513 Pfeiffer, T., Rutte, C., Killingback, T., Taborsky, M., & Bonhoeffer, S. (2005).
514 Evolution of cooperation by generalized reciprocity. *Proceedings of the Royal*
515 *Society B: Biological Sciences*, 272, 1115–1120.
516 <https://doi.org/10.1098/rspb.2004.2988>

517 Pimentel, A., Carvalho, T. dos S., Lima, F., Lima-Maximino, M., Soares, M. C., &
518 Maximino, C. (2019). Conditional approach as cooperation in predator inspection:
519 a role for serotonin? *Behavioural Brain Research*, 365, 164–169.
520 <https://doi.org/10.1016/j.bbr.2019.03.005>

521 Rosner, B. (2016). *Fundamentals of biostatistics*. Cengage Learning.

522 Soares, M. C., Cardoso, S. C., Carvalho, T. dos S., & Maximino, C. (2018). Using
523 model fish to study the biological mechanisms of cooperative behaviour: A future
524 for translational research concerning social anxiety disorders? *Progress in Neuro-*
525 *Psychopharmacology and Biological Psychiatry*, 82, 205–215.
526 <https://doi.org/10.1016/j.pnpbp.2017.11.014>

527 Speedie, N., & Gerlai, R. (2008). Alarm substance induced behavioral responses in
528 zebrafish (*Danio rerio*). *Behavioural Brain Research*, 188, 168–177.
529 <https://doi.org/10.1016/j.bbr.2007.10.031>

530 Spence, R., Gerlach, G., Lawrence, C., & Smith, C. (2008). The behaviour and ecology
531 of the zebrafish, *Danio rerio*. *Biological Reviews*, 83, 13–34.
532 <https://doi.org/10.1111/j.1469-185X.2007.00030.x>

533 Stewart, A. M., Ullmann, J. F. P., Norton, W. H. J., Parker, M. O., Brennan, C. H.,
534 Gerlai, R., & Kalueff, A. V. (2015). Molecular psychiatry of zebrafish. *Molecular*
535 *Psychiatry*, 20, 2–17. <https://doi.org/10.1038/mp.2014.128>

536 Taborsky, B., & Oliveira, R. F. (2012). Social competence: An evolutionary approach.
537 *Trends in Ecology and Evolution*, 27, 679–688.
538 <https://doi.org/10.1016/j.tree.2012.09.003>

539 Trivers, R. L. (1971). The evolution of reciprocal altruism. *Quarterly Review in*
540 *Biology*, 46, 35–57. <https://doi.org/10.1086/406755>

541

542

543

544 **Figure captions**

545 Figure 1 – Experimental apparatus (A) Viewed from the side; (B) Viewed from above.

546 In (B), the positions of the mirror in Experiment 2 are also indicated as CM
547 (“Cooperating mirror”) and DM (“Defecting mirror”).

548 Figure 2 – Concomitant presence of a predator animation and a cooperating mirror
549 elicits predator approach. (A) Percent of change in the time spent in the inspection
550 zone. (B) Percent of change in the time spent in the avoidance zone. (C) Percent of
551 change in the time spent in the refuge zone. (D) Freezing duration. (D’) Cumming
552 estimation plots for freezing. (E) Frequency of erratic swimming events. (E’)
553 Cumming estimation plots for erratic swimming. In (A-C), percent changes are
554 normalized to time block duration, individual points are shown in the left-most
555 panel, and Cohen's d between No predator and Predator groups are shown in a
556 Gardner-Altman estimation plot. Both groups are plotted on the left axes; the mean
557 difference is plotted on a floating axes on the right as a bootstrap sampling
558 distribution. The mean difference is depicted as a dot; the 95% confidence interval
559 is indicated by the ends of the vertical error bar. In (D-E), the raw data is plotted on
560 the upper axes; each paired set of observations is connected by a line. In (D’-E’),
561 the paired mean differences for comparisons between groups are shown in a
562 Cumming estimation plot, with each paired mean difference plotted as a bootstrap
563 sampling distribution of Cohen's d . Mean differences are depicted as dots; the 95%
564 confidence interval is indicated by the ends of the vertical error bar.

565 Figure 3 – Zebrafish retaliate defections during predator inspection. (A) Percent of
566 change in the time spent in the inspection zone. (B) Percent of change in the time
567 spent in the avoidance zone. (C) Percent of change in the time spent in the refuge
568 zone. (D) Freezing duration. (D’) Cumming estimation plots for freezing. (E)
569 Frequency of erratic swimming events. (E’) Cumming estimation plots for erratic
570 swimming. In (A-C), percent changes are normalized to time block duration,
571 individual points are shown in the left-most panel, and Cohen's d between
572 Defecting mirror (DM) and Cooperating mirror (CM) groups are shown in a
573 Gardner-Altman estimation plot. Both groups are plotted on the left axes; the mean
574 difference is plotted on a floating axes on the right as a bootstrap sampling
575 distribution. The mean difference is depicted as a dot; the 95% confidence interval
576 is indicated by the ends of the vertical error bar. In (D-E), the raw data is plotted on

the upper axes; each paired set of observations is connected by a line. In (D'-E'), the paired mean differences for comparisons between groups are shown in a Cumming estimation plot, with each paired mean difference plotted as a bootstrap sampling distribution of Cohen's *d*. Mean differences are depicted as dots; the 95% confidence interval is indicated by the ends of the vertical error bar.

Figure 4 – Both the mirror image and predator animation are necessary to elicit inspection behaviour and fear-like responses. (A) Percent of change in the time spent in the inspection zone. (B) Percent of change in the time spent in the avoidance zone. (C) Percent of change in the time spent in the refuge zone. (D) Freezing duration. (D') Cumming estimation plots for freezing. (E) Frequency of erratic swimming events. (E') Cumming estimation plots for erratic swimming. In (A-C), percent changes are normalized to time block duration, individual points are shown in the left-most panel, and Cohen's *d* between Defecting mirror (DM) and Cooperating mirror (CM) groups are shown in a Gardner-Altman estimation plot. Both groups are plotted on the left axes; the mean difference is plotted on a floating axes on the right as a bootstrap sampling distribution. The mean difference is depicted as a dot; the 95% confidence interval is indicated by the ends of the vertical error bar. In (D-E), the raw data is plotted on the upper axes; each paired set of observations is connected by a line. In (D'-E'), the paired mean differences for comparisons between groups are shown in a Cumming estimation plot, with each paired mean difference plotted as a bootstrap sampling distribution of Cohen's *d*. Mean differences are depicted as dots; the 95% confidence interval is indicated by the ends of the vertical error bar.

A **B** **C**

A

Change in predator inspection (%)

B

Cohen's d

D

Freezing (s)

D'Paired
cohen's d **E**

Erratic swimming (N)

E'Paired
cohen's d

A **B** **C** **D** **D'** **E** **E'**