Existence and Features of the Myodural Bridge in Gentoo Penguins: a morphological study

Cheng Chen¹, Sheng-bo Yu¹, Yan-yan Chi¹, Guang-yuan Tan², Bao-cheng Yan², Nan Zheng^{1*}, Hong-Jin Sui^{1,3*}

¹ Department of Anatomy, College of Basic Medicine, Dalian Medical University, Dalian, China

² Haichang Ocean Park Holdings.,Ltd, Biological Conservation Center, Shanghai, China

³ Dalian Hoffen Preservation Institution, Dalian, China

*Corresponding to:

Prof. Dr. Hong-Jin Sui, Department of Anatomy, College of Basic Medicine, Dalian Medical University, Dalian, China. E-mail address: suihj@hotmail.com

Prof. Dr. Nan Zheng, Department of Anatomy, College of Basic Medicine, Dalian Medical University, Dalian, China. E-mail address: zhengnan831016@163.com

20

21

22

23 **Abstract**

24

25 Recent studies have evidenced that the anatomical structure now known as the myodural bridge
 26 (MDB) connects the suboccipital musculature to the cervical spinal dura mater (SDM). In humans,
 27 the MDB passes through both the posterior atlanto-occipital and the posterior atlanto-axial
 28 interspaces. The present authors suggest that the MDB has important physiological functions in
 29 humans. The existence of the MDB in various mammals, including flying birds (*Rock pigeons and*
 30 *Gallus domesticus*) has been previously validated. Gentoo penguins are marine birds, able to make
 31 450 dives per day, reaching depths of up to 660 feet. Gentoo penguins are also the world's fastest
 32 diving birds. The present study was therefore carried out to investigate the existence and
 33 characteristics of the MDB in Gentoo penguin (*Pygoscelis papua*), a non-flying, marine bird that
 34 can dive. While foraging, this penguin is able to reach speeds of up to 22 miles per hour. For this
 35 study, six Gentoo penguin specimens were dissected to observe the existence and composition of
 36 their MDB. Histological staining was also performed to analyze the anatomic relationships and
 37 characteristic of the MDB in the Gentoo penguin. In this study, it was found that the suboccipital
 38 musculature in the Gentoo penguin consists of the rectus capitis dorsalis minor (RCDmi) muscle
 39 and rectus capitis dorsalis major (RCDma) muscle. Dense connective tissue fibers were observed
 40 connecting these two suboccipital muscles to the spinal dura mater (SDM). This dense connective

tissue bridge consists of primarily type I collagen fibers. Thus, this penguin's MDB appears to be analogous to the MDB previously observed in humans. The present study evidences that the MDB not only exists in penguins but it also has unique features that distinguishes it from that of flying birds. Thus, this study advances the understanding of the morphological characteristics of the MDB in flightless, marine birds.

Key words: Myodural bridge, Gentoo penguins, Morphology, Suboccipital muscles, Histologic staining

Introduction

The tissue bridge connecting a suboccipital muscle (RCPmi) to the cervical spinal dura mater (SDM) in humans was identified in the atlanto-occipital interspace by Hack et al., naming it the “myodural bridge” [1]. The myodural bridge (MDB) is now described as a fibrous, dense structure connecting the suboccipital musculature to the SDM, including fibers originating from the rectus capitis posterior minor (RCPmi) muscle, the rectus capitis posterior major (RCPma) muscle, the obliques capitis inferior (OCI) muscle, and the nuchae ligament (NL) [2-7]. With these in-depth studies, many researchers now speculate that the MDB may play a significant role in physiological functions. It has been proposed that the MDB's fibers connecting the suboccipital muscles to the cervical SDM [1] might prevent dural infolding, thus maintaining the normal flow of the cerebrospinal fluid within the cisterna magna or cerebellomedullaris cistern [4, 7-12]. Sui et al.

(2013) proposed that the MDB may play an important roles in modulating the circulation of cerebrospinal fluid. According to recent reports, MDB dysfunction may occur with pathological conditions of the RCPmi muscle, resulting in the generation of cervicogenic headache, and other craniofacial disorders [13-17].

In recent years, researchers have found that the MDB is a universal structure in mammals [18], including marine mammals (*Nephocaena phocaenoides* and *sperm whales*) [19, 20]. The MDB also exists in reptiles (*Siamese crocodile* and *Trachemys scripta elegans*) [21, 22], as well as in birds (*Rock pigeons* and *Gallus domesticus*) [23, 24]. Therefore the universal existence of the MDB among reptiles, birds, and mammals suggests that the MDB is an evolutionarily conserved anatomic structure having important biological functions. Moreover, we suggest that there may be structural differences in among various animals to adapt to different environmental conditions.

Penguins are unique birds that can both walk on land and dive and swim in deep water. Penguins are flightless birds, unlike the Rock pigeon and the *Gallus domesticus*. This marine bird's ability to swim presents it with different living environments and habits, then those of non-marine birds. Therefore, Gentoo penguins will be studied in this paper. This study will provide a comparative anatomical foundation for the future study of the MDB in different animals.

Materials and Methods

For this scientific research study, six adult Gentoo penguin specimens from Haichang Ocean Park Holdings., Ltd., that died of natural causes, were obtained with approval from both the Chinese

83 Authorities for Animal Protection and also approved by from the Ethics Committee of Dalian
84 Medical University.

85 In addition, all experiments were conducted in accordance with the guidelines and regulations of
86 Dalian Medical University. The collected Gentoo penguin carcasses were immobilized in a 10%
87 formalin solution for subsequent experiments.

88

89 **Anatomical dissection in the suboccipital region**

90

91 Three of the six penguin specimens were used for gross dissection. The epidermis was cut
92 longitudinally along the dorsal midline of the specimens, and the suboccipital musculature was
93 fully exposed. The deep suboccipital musculature, including the RCDmi, (analogous to the RCPmi
94 muscle found in humans) and the RCDma, (analogous to the RCPma muscle also found in
95 humans) was exposed. The deep suboccipital muscles (RCDmi and RDCma) were then both cut
96 from the bony occipital crest to observe their connections with the dorsal atlanto-occipital
97 membrane (DAOM, analogous to the PAOM observed in humans) and the dorsal atlanto-axial
98 membrane. Next the DAOM was also cut along its cranial attachment to observe the connection
99 between the DAOM and the SDM. Lateral incision of the posterior arch of the atlas was also made
100 to observe the connection between the dorsal atlanto-axial membrane and the SDM. The
101 photographic materials were taken with a Canon 7D camera (Canon Inc., Tokyo, Japan).

102

Histological slices and staining

Three specimens were used for histological studies. Tissue samples of the occiput and the cervical region were immobilized in a 10% formalin solution for ten days. Subsequently, the specimens were transferred to Jiang Weizhong's decalcification solution for decalcification. The Jiang Weizhong's decalcification solution was changed at three-day intervals until the bone was easily punctured by a needle (21 days). The decalcified tissue samples were then washed overnight in running water, and then dehydrated with increasing grades of alcohol, transparent in xylene, and then infiltrated with melted paraffin, and embedded. A rotary microtome (Leica Micro HM450; Lei Microsystems GmbH, Wetzlar, Germany) was used to cut 8- μ m-thick slices. These tissue slices were then divided into three groups: Group 1 slices were used for Hematoxylin and Eosin (HE) staining, which can show the general structural features of the MDB; Group 2 slices were used for Masson trichrome staining (Masson), to detect any collagen fibers in the MDB; Group 3 slices were used for Picrosirius Red (PRS) staining, which can also detect collagen fibers in the MDB, and most importantly, collagen types can be distinguished when viewed with a polarized light microscope. Microscopic examination and photography were carried out with the Nikon ECLIPSE80i research light microscope, and multiple images of each section were stitched together using the Microsoft Image Synthesis Editor of Nikon ECLIPSE80i Image Processing and Analysis System. The results of Picric acid-Sirius red staining were observed with a light microscope as well as a polarized light microscope.

Results

Anatomical dissection of the suboccipital region

The RCDmi and the RCDma muscles, located in the deep suboccipital region, were observed to be tightly connected to each other (Fig 1a). The cranial end of RCDmi muscle attached to the medial part of the occipital crest, and the caudal end attached to the dorsal side of the posterior tubercle of atlas. Moreover multiple dense fibrous tissues were observed to originate from the ventral part of the RCDmi connect to the DAOM (Fig 1b). The DAOM and the SDM were intimately connected to each other by numerous trabecular fibrous bundles (Fig 1c). The cranial end of the RCDma muscle attached to the lateral aspect of the occipital crest, and the caudal end attached to the spinous process of the axis. Some of the dense fibrous bundles originating from the ventral part of the RCDma muscle connected to the dorsal atlanto-axial membrane (Fig 2a). The dorsal atlanto-axial membrane adheres tightly to the SDM via several dense cord-like fibrous tissue (Fig 2b). This cord-like trabecular fibrous tissue was found in all the three specimens.

Fig 1. Gentoo penguin, dorsal views of an anatomical dissection of the deep suboccipital

space. a: Superficial view of the deep suboccipital region. The RCDmi and the RCDma are connected tightly. b: The connection between the RCDmi and the DAOM. The ventral surface of the RCDmi is connected by multiple dense fibrous tissues (arrow) to the DAOM (hollow star). c: The connection between the DAOM and the SDM. The DAOM was separated from its cranial

attachment at the foramen magnum, and the ventral surface of the DAOM is connected by trabecular fibrous bundles (arrow) to the SDM (hollow triangle). Abbreviation: OCCI=occipital bone; RCDmi=rectus capitis dorsalis minor muscle; RCDma=rectus capitis dorsalis major muscle; DAOM=dorsal atlanto-occipital membrane; SDM=spinal dura mater; C1=posterior arch of atlas.

Fig 2. Gentoo penguin, dorsal views of an anatomical dissection of the Atlanto-axial

interspace. a: The connection between the RCDma and the dorsal atlanto-axial membrane. The ventral surface of RCDma is connected by dense fibrous bundles (arrow) to the dorsal atlanto-axial membrane. b: The connection between the dorsal atlanto-axial membrane and the SDM. The dorsal atlanto-axial membrane was separated from its posterior arch of atlas, and the ventral surface of the the dorsal atlanto-axial membrane is connected by dense cord-like fibrous tissues (arrow) to the SDM (hollow crescent). Abbreviation: RCDmi=rectus capitis dorsalis minor muscle; RCDma=rectus capitis dorsalis major muscle; SDM=spinal dura mater; C1=posterior arch of atlas.

Histology studies

In the HE-stained sections, multiple dense fibers originating from the ventral anterior aspect of RCDmi pass through the atlanto-occipital interspace (Fig 3a). These fibers extend from the ventral side of the RCDmi muscle and terminate in the DAOM. The DAOM emitted dense fibrous bundles and connected with the SDM passing through several venous sinuses (Fig 3b,c). However, different origins of these connective fibers in the atlanto-axial interspace were found. Some of the

fibrous bundles originating from the ventral part of the RCDmi connect with the dorsal atlanto-axial membrane directly (Fig 3d). Fibers originating from the ventral aspect of RCDma as well as fibers originating from the OCP fuse together then pass through the dorsal atlanto-axial membrane and terminate on the SDM (Fig 3d). These connective tissue fibers observed between the muscles and the SDM are analogous to the MDB fibers previously observed in humans. In the Masson-stained sections, the muscular fibers were stained red, and the MDB fibers were stained blue (Fig 4). The results of the Masson staining demonstrate that the MDB fibers present in Gentoo penguins are collagenous fibers. In the PSR-stained sections, the fibers of the MDB were stained red under when viewed the ordinary optical microscope (Fig 5a), which implies that the penguin MDB is composed primarily of collagen fibers. Viewed with the polarizing microscope, the fibrous tissues of MDB were stained either red or yellow (Fig 5b,c,d). This observation implies that the MDB is composed of primarily type I collagen fibers.

Fig 3. Gentoo penguin, prepared sagittal section of the suboccipital region with stained with HE. Area within square in a is shown enlarged in b, c, and d. The fibers (arrow) originate from the ventral anterior of RCDmi, extend directly into the atlanto-occipital interspace, then fuse with the DAOM, and the DAOM extends into a fiber bundle through the venous sinus to connect with the SDM (hollow triangle). In the atlanto-axial interspace, the dense fibers bundles (arrow) originating from the ventral of RCDmi run ventrally, and integrate into the dorsal atlanto-axial membrane (hollow crescent) together with the dense fibers(filled arrowhead) originating from the RCDma and OCP, and send out dense fibers bundles to closely connect with the SDM. Abbreviation: OCCI=occipital bone; RCDmi=rectus capitis dorsalis minor muscle; OCP=oblique capitis

posterior; DAOM=dorsal atlanto-occipital membrane; SDM=spinal dura mater; C1= atlas;

C2=axial;V=venous sinus

Fig 4. Gentoo penguin, prepared sagittal section of the suboccipital region with stained with

Masson. Area within square in a is shown enlarged in b, c, and d. The fibers (arrow) originate from the ventral anterior of RCDmi, extend directly into the atlanto-occipital interspace, then fuse with the DAOM, and the DAOM extends into a fiber bundle through the venous sinus to connect with the SDM (hollow triangle). In the atlanto-axial interspace, the dense fibers bundles (arrow) originating from the ventral of RCDmi run ventrally, and integrate into the dorsal atlanto-axial membrane (hollow crescent) together with the dense fibers(filled arrowhead) originating from the RCDma and OCP, and send out dense fibers bundles to closely connect with the SDM.

Abbreviation: OCCI=occipital bone; RCDmi=rectus capitis dorsalis minor muscle; OCP=oblique capitis posterior; DAOM=dorsal atlanto-occipital membrane; SDM=spinal dura mater; C1= atlas; C2=axial; V=venous sinus.

Figure 5. Gentoo penguin, prepared sagittal section of the suboccipital region with stained

with PSR-stained. Area within square in a is shown enlarged in b, c and d. The fibers (arrow) originate from the ventral anterior of RCDmi, extend directly into the atlanto-occipital interspace, then fuse with the DAOM, and the DAOM extends into a fiber bundle through the venous sinus to connect with the SDM (hollow triangle). In the atlanto-axial interspace, the dense fibers bundles (arrow) originating from the ventral of RCDmi run ventrally, and integrate into the dorsal atlanto-axial membrane (hollow crescent) together with the dense fibers(filled arrowhead) originating from the RCDma and OCP, and send out dense fibers bundles to closely connect with the SDM. Abbreviation: OCCI=occipital bone; RCDmi=rectus capitis dorsalis minor muscle;

OCP=oblique capitis posterior; DAOM=dorsal atlanto-occipital membrane; SDM=spinal dura mater; C1= atlas; C2=axial; V=venous sinus.

Discussion

In 1995 Hack et al. initially proposed the concept of the MDB [1]. Numerous researchers have since studied the MDB in detail, confirming the MDB's existence in both humans [1-7] and animals [18-24]. These researchers have further hypothesized about the putative physiological functions of the MDB [8-17]. The MDB is described as a dense connective tissue bridge connecting the suboccipital musculature to the cervical SDM, while passing through the posterior atlanto-occipital and the atlanto-axial interspaces. Therefore, what is unique about the penguin as a natatores.

According to the observation of the gross anatomy of the penguin, evidenced in the present study, the deep suboccipital musculature of the Gentoo penguin consists of the RCDmi and the RCDma muscles. The cranial ends of both the RCDmi and the RCDma muscles respectively attach to the medial and lateral aspects of the occipital crest. They were intimately related to each other and not easily separated. Interestingly, it was previously found that the RCDmi muscle of *Nephocaena phocaenoides* was located on the deep surface of the RCDma, and the RCDmi was observed originating from occiput [19]. Furthermore, the RCDmi of the *sperm whale* was previously found to originate from the dorsal aspect of the occipital squama [20]. This spatial distribution of the

RCDmi in marine mammals (*Nephocaena phocaenoides* and *sperm whale*) is similar to that of the Gentoo penguin. However, the cranial attachments of both the RCDmi and RCDma muscle of the Gentoo penguin are distinct from that observed in other birds (*Rock pigeons* and *Gallus domesticus*). The cranial aspect of the RCDmi attached below the transverse nuchal crest, and the both the RCDmi and RCDma muscle were easily detached [23, 24]. This difference may be related to the survival and feeding behavior of the penguin. These behaviors can be inferred from skull morphology [28].

In the present study, we have confirmed the existence of the MDB in the Gentoo penguin through multiple methods. In the atlanto-occipital interspace of the Gentoo penguins, the dense MDB fibers originated primarily from the ventral aspect of the RCDmi. These fibers extend from the ventral side and extend superiorly to fuse with the DAOM. Then these fibers connect with the SDM passing through the venous sinus. In the atlanto-axial interspace of the Gentoo penguin, the dense MDB fibers originate from the ventral of the RCDmi, the RCDma, and the OCP, all converging to form the MDB, and then passing through the dorsal atlanto-axial membrane, which is tightly connected to the SDM. The arrangement of the fibers of the MDB, observed in the Gentoo penguin is similar to that of other birds [23, 24] and mammals [18]. Although penguins live in different conditions from that of the *Rock pigeons* and the *Gallus domesticus*, they all possess an MDB. Once again, the MDB is seen as a highly conserved structure. During the course of biological evolution, structures that are not functionally important tend to degrade, and this evolutionary conservation of the MDB demonstrates the significance of MDB. In addition, the MDB observed in the Gentoo penguin presents as a dense fibrous structure that is composed of

type I collagen fibers.

While, being a marine bird, penguin not only has the same type of MDB as that observed in flying birds, but also has the same spatial distribution of the suboccipital muscles like marine mammals. What then are the similarities and differences between the MDB of penguins and marine mammals? Research has confirmed that no DAOM found in the posterior interspace between the occipital bone and atlas of *Nephocaena phocaenoides* [19]. The tendinous fibers of the RCDmi observed in the *Nephocaena phocaenoides* projected through the atlanto-occipital interspace attaching directly to the SDM. This direct connection between the RCDmi and the SDM of the *Nephocaena phocaenoides* might create a more pronounced effect of the RCDmi muscle on the SDM [19]. Furthermore, the present authors have confirmed that *sperm whales* have two different origins of their MDB: one origin is from the ODB (occipital dural bridge) which originates from the periosteal surface of the occiput and fuses with the dura mater [20], and the other bridge (MDB) originates from the RCDmi, which transmits the tension from the RCDmi to the dura mater. The MDB stabilizes the dura mater, the spinal cord, and the ODB [20]. The MDB, observed in the atlanto-occipital interspace of penguins, is primarily composed of dense connective tissue fibers originating from the RCDmi muscle. Therefore, there is a type of the MDB, originating from the RCDmi, in penguins, *sperm whales*, and *Nephocaena phocaenoides* which all pass through the atlanto-occipital interspace. Moreover, they have similar distribution of suboccipital muscles and the presence of numerous suboccipital venous sinuses. As these different animals live in similar conditions, some of their anatomic structures tend to be similar. This is referred to as the “similar effect”. Therefore, as the penguin is capable of navigating in deep

water, this ability may be associated with cerebrospinal fluid circulation. This also demonstrates that the MDB is a significant and highly conserved anatomic structure.

Interestingly, the *sperm whale* has an atlas and six fused vertebra (C2-C7) [27], while their MDB exists only in the atlanto-occipital interspace, with extensive venous plexus among the MDB fibers within their atlanto-occipital interspace [20]. As *sperm whales* swim to great depths, the MDB of these animals may contribute to transferring the tensile forces generated by the suboccipital muscles to the cervical dura mater and thereby continuously alter the volume of the CSF contained within the subarachnoid space, acting as a unique mechanism to facilitate circulation of the CSF [20]. The penguin's MDB exists in both the atlanto-occipital and the atlanto-axial interspaces, and there are numerous venous sinuses among the MDB fibers. As the penguin dives to great ocean, the contraction of MDB fibers via the suboccipital musculature may accelerate venous blood flow back to the heart and also promote cerebrospinal fluid circulation. According to previous research, the penguin can dive to approximately 100 meters within 3 minutes [29, 30]. Therefore, the penguin's MDB might be a key structure which helps the penguin to maintain normal cerebrospinal fluid circulation during deep dives.

In this study, the present authors validated that there are dense connective tissues (MDB) in both the atlanto-occipital and atlanto-axial interspaces of Gentoo penguin, that are composed of primarily type I collagen fibers. Therefore, this finding implies that the MDB is not only a highly conserved anatomic structure, from an evolutionary point of view, but also has its variant structure

for acclimatization. These findings provide supporting evidence for research into the physiological function of the MDB.

Acknowledgements

We are grateful to Haichang Ocean Park Holdings., Ltd for provision of cadaveric specimens. Thanks to Gary D. Hack of University of Maryland School of dentistry, USA, for helping with language editing.

References

1. Hack, G. D., Koritzer, R. T., Robinson, W. L., Hallgren, R. C., & Greenman, P. E. 1995. Anatomic Relation between the Rectus Capitis Posterior Minor Muscle and the Dura Mater.– Spine. 20(23), 2484–2486. doi: 10.1097/00007632-199512000-00003.
2. Mitchell BS, Humphreys BK, Osullivan E. 1998. Attachments of the ligamentum nuchae to cervical posterior spinal dura and the lateral part of the occipital bone.–J Manipulative Physiol Ther. 21(3):145–148.
3. Nash L, Nicholson H, Lee AS, Johnson GM, Zhang M. 2005. Configuration of the Connective Tissue in the Posterior Atlanto-Occipital Interspace: A Sheet Plastination and Confocal Microscopy Study. – Spine. 30(12):1359–1366. doi: 10.1097/01.brs.0000166159.31329.92.

- 316 4. Scali, F., Marsili, E. S., & Pontell, M. E. 2011. Anatomical Connection Between the Rectus
317 Capitis Posterior Major and the Dura Mater.—Spine. 36(25):E1612–E1614.
318 doi:10.1097/brs.0b013e31821129df.
- 319 5. Pontell ME, Scali F, Marshall E, Enix D. 2013. The obliquus capitis inferior myodural
320 bridge.—Clin Anat. 26(4):450–454. doi: 10.1002/ca.22134.
- 321 6. Palomeque-del-Cerro, L., Arráez-Aybar, L. A., Rodríguez-Blanco, C., Guzmán-García, R.,
322 Menendez-Aparicio, M., & Oliva-Pascual-Vaca, Á. 2017. A Systematic Review of the
323 Soft-Tissue Connections Between Neck Muscles and Dura Mater. — Spine. 42(1):49–54.
324 doi:10.1097/brs.0000000000001655.
- 325 7. Zheng N, Yuan XY, Li YF, Chi YY, Gao HB, Zhao X, Yu SB, Sui HJ, John Sharkey. 2014.
326 Definition of the to be named ligament and vertebroductal ligament and their possible effects
327 on the circulation of CSF. — PloS one. 9(8): e103451. doi:10.1371/journal.pone.0103451.
- 328 8. Rutten HP, Szpak K, van Mameren H, Ten Holter J, de Jong JC. 1997. Anatomic relation
329 between the rectus capitis posterior minor muscle and the dura mater. — Spine. 22(8):924–926.
330 doi: 10.1097/00007632-199704150-00018.
- 331 9. Sui HJ, Yu SB, Yuan XY, Gao HB, Xu Q, Li XF. 2013. Anatomical study on the connections
332 between the suboccipital structures and the spinal dura mater. — Chinese journal of clinical
333 anatomy. 31:489–490.
- 334 10. Pontell ME, Scali F, Enix DE, Battaglia PJ, Marshall E. 2013. Histological examination of
335 the human obliquus capitis inferior myodural bridge. — Ann Anat. 195(6), 522–526.
336 doi:10.1016/j.aanat.2013.04.013.

- 337 11. Scali F, Pontell ME, Enix DE, Marshall E. 2013. Histological analysis of the rectus capitis
338 posterior major's myodural bridge. — Spine J. 13(5):558–563.
339 doi:10.1016/j.spinee.2013.01.015.
- 340 12. Scali F, Pontell ME, Welk AB, Malmstrom TK, Marshall E, Kettner NW. 2013. Magnetic
341 resonance imaging investigation of the atlanto-axial interspace. — Clin Anat. 2013
342 May;26(4):444-449. doi: 10.1002/ca.22094.
- 343 13. Hack GD, Hallgren RC. 2014. Chronic headache relief after section of suboccipital muscle
344 dural connections:a case report. — Headache. 44(1):84–89. doi:
345 10.1111/j.1526-4610.2004.04015.x.
- 346 14. Fernández-de-las-Peñas, C., Bueno, A., Ferrando, J., Elliott, J., Cuadrado, M., Pareja, J.
347 2007. Magnetic Resonance Imaging Study of The Morphometry of Cervical Extensor
348 Muscles in Chronic Tension-Type Headache. — Cephalalgia. 27(4), 355–362.
349 doi:10.1111/j.1468-2982.2007.01293.x.
- 350 15. Bogduk N. 2001. Cervicogenic headache: anatomic basis and pathophysiologic mechanisms.
351 — Curr Pain Headache Rep. 5(4):382-386. doi: 10.1007/s11916-001-0029-7.
- 352 16. Fernández-de-las-Peñas, C., Cuadrado, M. L., Arendt-Nielsen, L., Ge, H.-Y., Pareja, J. A.
353 2008. A proposed etiology of cervicogenic headache: the neurophysiologic basis and
354 anatomic relationship between the dura mater and the rectus posterior capitis minor muscle.
355 — Am J Phys Med Rehabil. 87(3):197-203. doi: 10.1097/PHM.0b013e3181619766.
- 356 17. Yuan XY, Yu SB, Liu C, Xu Q, Zheng N, Zhang JF, Chi YY, Wang XG, Lin XT, Sui HJ.
357 2016. Correlation between chronic headaches and the rectus capitis posterior minor muscle:

- 358 A comparative analysis of cross-sectional trail. — Cephalalgia. 37(11):1051-1056. doi:
359 10.1177/0333102416664775.
- 360 18. Zheng N, Yuan XY, Chi YY, Liu P, Wang B, Sui JY, Han SH, Yu SB, Sui HJ. 2017. The
361 universal existence of myodural bridge in mammals: an indication of a necessary function. —
362 Sci Rep. 7(1):8248. doi: 10.1038/s41598-017-06863-z.
- 363 19. Liu P, Li C, Zheng N, Xu Q, Yu SB, Sui HJ. 2017. The myodural bridge existing in the
364 Nephocaena phocaenoides. — PLoS One. 12(3):e0173630. doi: 10.1371/journal.pone.0173630.
- 365 20. Liu P, Li C, Zheng N, Yuan Xy, Zhou Y, Chun P, Chi YY, Gilmore Campbell, Yu SB, Sui
366 HJ. 2018. The myodural bridges' existence in the sperm whale. — PLoS One. 13(7):e0200260.
367 doi: 10.1371/journal.pone.0200260.
- 368 21. Zhang JH, Tang W, Zhang ZX, Luan BY, Yu SB, Sui HJ. 2016. Connection of the Posterior
369 Occipital Muscle and Dura Mater of the Siamese Crocodile. — Anat Rec (Hoboken).
370 299(10):1402-1408. doi: 10.1002/ar.23445.
- 371 22. Zhao Huangfu, Xiao Zhang, Jia-Ying Sui, Qi-Qi Zhao, Xiao-Ying Yuan, Chan Li, Ya-Ru
372 Dou, Wei Tang, Mei-Ling Du, Nan Zheng, Yan-Yan Chi, Sheng-Bo Yu, Hong-Jin Sui. 2019.
373 Existence of Myodural Bridge in the Trachemys scripta elegans: Indication of its Important
374 Physiological Function. — Int J Morphol. 37(4):1353-1360.
- 375 23. Chukwuemeka Samuel Okoye, Zheng N, Yu SB, Sui HJ. 2018. The myodural bridge in the
376 common rock pigeon (Columbia livia) : Morphology and possible physiological
377 implications. — J Morphol. 279(10):1524-1531. doi: 10.1002/jmor.20890.

- 378 24. Dou YR, Zheng N, Gong J, Tang W, Chukwuemeka Samuel Okoye, Zhang Y, Chen YX ,
379 Zhang Y, Pi SY, Qu LC, Yu SB, Sui HJ. 2019. Existence and features of the myodural bridge
380 in Gallus domesticus: indication of its important physiological function. — Anat Sci Int.
381 94(2):184-191. doi: 10.1007/s12565-018-00470-2.
- 382 25. Zumpano MP, Hartwell S, Jagos CS. 2006. Soft tissue connection between rectus capitus
383 posterior minor and the posterior atlanto-occipital membrane: a cadaveric study. — Clin Anat.
384 19(6):522-7. doi: 10.1002/ca.20220.
- 385 26. Zheng Nan, Chi Yan-Yan, Yang Xiao-Han, Wang Nan-Xing, Li Yi-Lin, Ge Yang-Yang, Zhang
386 Lan-Xin, Liu Tai-Yuan, Yuan Xiao-Ying, Yu Sheng-Bo, Sui Hong-Jin. 2018. Orientation and
387 property of fibers of the myodural bridge in humans. — Spine J. 18(6):1081-1087. doi:
388 10.1016/j.spinee.2018.02.006.
- 389 27. Shadwick RE, Gosline JM. 1995. Arterial Windkessels in marine mammals. — Symp Soc Exp
390 Biol. 49:243-52.
- 391 28. Acosta Hospitaleche, C., Tambussi, C. 2006. Skull morphometry of Pygoscelis
392 (Sphenisciformes): inter and intraspecific variations. — Polar Biol. 29(9):728–734.
393 doi:10.1007/s00300-006-0109-6.
- 394 29. Nigel J. Adams, Christopher R. Brown. 1983. Diving Depths of the Gentoo Penguin
395 (Pygoscelis papua). — Condor. 85(4):503-504.
- 396 30. Williams, T. D., Kato, A., Croxall, J. P., Briggs, D. R., & Barton, T. R. 1992. Diving Pattern
397 and Performance in Nonbreeding Gentoo Penguins (Pygoscelis papua) during Winter. — Auk.
398 109(2), 223–234. doi:10.2307/4088190.

399

400 **Supporting information**

401 S1 File. Staining methods.

402

403

404

405

406

407

408

409

410

411

Figure

Figure

Figure

Figure

Figure