Risk-related decision making evokes distinct brain activation patterns in reward evaluation regions in substance use naïve versus non-naïve adolescents

Goldie A. McQuaid¹, Valerie L. Darcey^{1,2}, Amanda E. Patterson¹, Emma J. Rose³, John W. VanMeter¹, Diana H. Fishbein^{3,4}

¹Center for Functional and Molecular Imaging, Georgetown University Medical Center, Washington, DC, USA

²The Interdisciplinary Program in Neuroscience, Georgetown University, Washington DC, USA

³Bennett Pierce Prevention Research Center and The Department of Human Development and Family Studies, College of Health and Human Development, The Pennsylvania State University, University Park, PA, USA

⁴Frank Porter Graham Child Development Institute, University of North Carolina, Chapel Hill, NC, USA

Corresponding author: Goldie A. McQuaid, Center for Functional and Molecular Imaging, Georgetown University Medical Center, Washington, DC, USA, gam58@georgetown.edu

ABSTRACT

Identifying brain and behavioral precursors to substance use (SU) may guide interventions that delay initiation in youth at risk for SU disorders (SUD). Heightened reward-sensitivity and risk-taking may confer risk for SUD. In a longitudinal, prospective study, we characterized behavioral and neural profiles associated with reward-sensitivity and risk-taking in substance-naïve adolescents, examining whether they differed as a function of SU initiation at 18- and 36-months follow-up.

Adolescents ($N=70$; 11.1-14.0 years) completed a reward-related decision-making task (Wheel of Fortune (WOF)) while undergoing functional MRI. Measures of reward sensitivity (Behavioral Inhibition System-Behavioral Approach System; BIS-BAS), impulsive decision-making (delay discounting task), and SUD risk (Drug Use Screening Inventory, Revised (DUSI-R)) were collected at baseline. Baseline metrics were compared for youth who did (SI; $n=27$) and did not (SN; $n=43$) initiate SU at follow-up.

While groups displayed similar discounting and risk taking behavior, SI youth showed more variable patterns of activation in left insular cortex during high-risk selections, and left anterior cingulate cortex in response to rewarded outcomes. SI participants scored higher on the DUSI-R and BAS subscales. Results suggest differences in brain regions critical in the development and experience of SUDs may precede SU and serve as a biomarker for SUD risk.

Keywords: Adolescence; Anterior Cingulate Cortex; Decision-making; Insula; Reward; Substance use

INTRODUCTION

Adolescence is commonly characterized as a period of increased risk-taking coupled with heightened reward sensitivity^{1,2}. Risk-taking during this evolutionarily conserved developmental period may have positive outcomes³. For instance, increased exploratory behaviors occurring during adolescence allow for adaptive risk-taking⁴⁻⁷, which facilitates the achievement of key developmental milestones in preparation for the transition to adulthood⁸. However, brain changes that condition adaptive risk-taking also render adolescents vulnerable to risk-taking that leads to negative outcomes, including substance use (SU)⁹.

Early SU initiation is associated with a constellation of other negative risk-taking behaviors and related adverse outcomes¹⁰, including delinquency or criminal activity¹¹, risky sexual behavior^{12,13}, physical assault¹⁴, accidental injury¹⁵ and death¹⁶. Given the potential for deleterious outcomes, SU among adolescents has been identified as a global health concern¹⁷. Critically, while earlier SU initiation is associated with greater risk for development of lifetime SU disorders (SUDs)¹⁸⁻²⁷, any delay in SU initiation decreases risk for development of SUDs^{19,28}. For instance, each year of delayed alcohol initiation is associated with a 5-9% decrease in risk for alcohol use disorder²⁰. Identifying factors which facilitate prediction of early initiation, therefore, may be beneficial in targeting prevention efforts to delay SU onset.

Developmental neuroscience models offer potential explanations for increased risk-taking during adolescence that leads to SU initiation. The dual systems^{29,30}, triadic^{31,32}, and imbalance models^{33,34}, generally postulate that subcortical brain regions associated with reward processing (e.g., ventral striatum, amygdalae) develop earlier than neocortical regions (e.g., prefrontal cortex (PFC)) associated with cognitive control. Development of the PFC and its functional networks, which continues throughout adolescence and into early adulthood^{35,36}, is associated

with improvements in top-down control of behavior^{37–39}. Such a protracted course of development may render the PFC vulnerable to the impacts of abused substances during adolescence⁴⁰, with early SU potentially altering neurodevelopmental trajectories^{41–44} and, ultimately, adversely affecting adult neurobiology and behavior^{9,45}.

Given the proximal and distal negative outcomes associated with early SU, it is critical for the design of effective interventions to identify risk markers that *precede* initiation. Adolescents who initiate SU early often demonstrate preexisting heightened impulsivity,^{46,47} sensation seeking⁴⁸ and reward sensitivity^{49–51}. Such traits are associated with poor self-regulation of emotion⁵², behavioral dyscontrol⁵¹, and a relative imperviousness to punishment⁵³, along with increased susceptibility to deviant peer influences⁵⁴. However, these associations do not explain individual differences in outcomes nor do they identify mechanisms that may underlie those differences that would otherwise provide malleable targets for intervention. Brain metrics may, in contrast, predict risk for psychopathology with greater specificity and sensitivity than behavioral measures alone⁵⁵. Thus, it is important to understand underlying neurobiology associated with increased risk⁵⁶ and identify neuroendophenotypes concerning risk for and protection from SUDs^{57,58}.

To address whether such biomarkers are observable prior to SU initiation, we followed SU-naïve early adolescents ($N=70$; aged 11.1–14.0 years) over 36 months, and compared participants who did and did not report initiation of alcohol and/or drugs at follow-up. Our central aim was to characterize behavioral and neural profiles associated with reward-sensitivity/risk-taking in relation to SU initiation. Adolescent participants completed measures probing reward sensitivity and risk aversion (Behavioral Inhibition System/Behavioral Activation System (BIS/BAS) Scales), and tasks to assess risk-taking and impulsivity in the context of rewards (Wheel of Fortune and delay discounting tasks, respectively). We predicted that at baseline, those who

would go on to initiate SU at 18- or 36-months follow-up would demonstrate greater hedonic and behavioral responsivity to rewards, overvalue immediate rewards, and make riskier choices in a reward-related decision-making task compared to adolescents who remained SU-naïve throughout the study. Further, we predicted that, prior to SU onset, reward-based decision making would be associated with differences between subsequent SU initiators and non-initiators in brain regions implicated in decision-making under uncertainty (i.e., medial prefrontal cortex⁵⁹⁻⁶²), and in the modulation of reward processing/sensitivity (i.e., ventral striatum and amygdalae^{59,63}).

METHODS

Participants were recruited as part of the Adolescent Development Study (ADS), a prospective longitudinal investigation of the neurodevelopmental precursors to and consequences of early SU initiation and escalation. Detailed information on ADS study methods and aims is presented elsewhere⁶⁴. Briefly, a total of 135 typically developing, SU-naïve early adolescents were recruited from the Metropolitan Washington D.C. region and followed longitudinally. Demographic, cognitive, behavioral, and imaging assessments were conducted at an initial (“baseline”) visit and during two follow-up visits, approximately 18- and 36-months later. The mean time elapsed between the baseline visit and the planned 18-months visit (Wave 2) was 18.4 (SD=3.6) months; and the mean time between baseline and 36-months follow-up (Wave 3) was 36.7 (SD=4.4) months. Imaging and behavioral data reported here were collected during the initial SU-naïve baseline assessment. Exclusionary criteria for the study included adolescent self-report of alcohol (>1 full drink of alcohol at any time) or, with the exception of nicotine, any SU prior to the initial visit; *in utero* exposure to alcohol or illicit drugs (parent-reported); a diagnosed

neurodevelopmental disorder (e.g., Autism Spectrum Disorder); left-handedness; a sibling of a current participant; history of head injury resulting in loss of consciousness >5 minutes; or MRI contraindication. The Georgetown University IRB approved all procedures, and written consent and assent were obtained from the parent and adolescent, respectively.

Participants

Of the 135 participants enrolled in the study, 70 adolescents aged 11.1–14.0 years ($M=12.7$ years, $SD=0.66$; female=40 (57%)) were included in the analyses reported here. One enrolled participant was excluded due to neurodevelopmental disorder. Participants were excluded from analyses due to missing or incomplete imaging data ($n=15$) and/or excessive head motion during imaging ($n=24$). Additionally, since a primary aim was to examine neural activation during risk-taking, participants who did not make any ‘high-reward/risk’ selections during the Wheel of Fortune task (WOF; see below for paradigm description) were excluded from analyses ($n=4$). Groups were defined based on SU status at follow-up, as detailed below. Participants for whom SU status could not be determined due to attrition or survey discrepancies ($n=21$) were also excluded from analyses reported here. (Table S1 provides a detailed summary of exclusions/inclusions.)

Family/caregiver measures

Socioeconomic (SES) index

An SES Index was calculated by averaging the mean of two standard scores (mean household income bracket before taxes and mean cumulative years of parental education), and re-standardizing these to obtain a z-score distribution with a 0-centered mean and a standard

deviation of 1 for the sample analyzed ($N=70$) (method adapted from⁶⁵).

Adolescent measures

SU initiation status

At baseline and at the Wave 2 and Wave 3 follow-up visits, adolescents completed two self-report surveys to determine SU status: the Tobacco Alcohol and Drug (TAD) survey and the Drug Use Screening Inventory Revised (DUSI-R)^{66,67}. The study-specific TAD included the alcohol and drug portion of the Semi-Structured Interview for the Genetics of Alcoholism⁶⁸ and asked about the use of substances, including tobacco, alcohol, and illicit drugs (i.e., marijuana, cocaine, methamphetamine, ecstasy, opiates, salvia, synthetic marijuana, inhalants, and illegally used prescription drugs), along with an open-ended “any other substances” question.

Adolescents also completed the DUSI-R, a survey with demonstrated psychometric validity^{69–71} and reliability⁷² for assessing SU and factors associated with risk for SUD later in adolescence. The DUSI-R includes 20 questions concerning use of specific substances (e.g., alcohol, marijuana, prescription painkillers, smoking tobacco, chewing tobacco) or substance classes (e.g., over the counter medications, tranquilizer pills, stimulants).

For the purposes of the analyses reported here, affirmative SU responses on both the TAD *and* the DUSI-R were used in determining SU status. Participants who reported SU on both the TAD and DUSI-R at either Wave 2 or Wave 3 follow-up were categorized as SU initiators (SI). Those who reported no SU on both the TAD and DUSI-R at both follow-up assessments were categorized as SU non-initiators (SN). As detailed above, participants for whom SU status could not be determined were excluded from analyses reported here.

DUSI-R absolute problem density (APD) score

In addition to questions concerning SU, the DUSI-R probes experiences and behaviors known to precede and co-occur with SU. The survey includes eight domains comprised of 159 *yes-no* items that are relevant for early adolescents: SU, behavior, health, social competence, psychiatric symptoms, school performance, family and peer relationships, and recreation⁷³. An absolute problem density (APD) score, which reflects overall risk for SU, is calculated by dividing the total number of “yes” questions by the total number of DUSI-R items. Here, group comparisons were conducted for the DUSI-R APD score only.

Delay Discounting (DD) task

Adolescents completed the DD⁷⁴ task outside of the scanner. The task was implemented in E-Prime 2.0. Participants were instructed to choose between receipt of an immediate reward of variable value ($\leq \$10$, in increments of \$0.50), versus receipt of \$10 after a specified temporal delay (e.g., *Would you rather have \$2 now, or \$10 in 30 days*). Discounting was assessed at six delays: 1, 2, 10, 30, 180, and 365 days. Participants were instructed to make their selections with care, as they would receive a reward ($\leq \$10$) based on a random selection of one of their choices⁷⁵.

Values for which the participant demonstrated no preference for immediate versus delayed receipt (i.e., the ‘indifference point’) were normalized to the static delayed reward value (\$10)⁷⁶ and plotted against each delay. To adjust for unequal weighting of indifference points at longer delays (a limitation of conventional methods of calculating area under the discounting curve; AUC), while preserving the notion of subjective experience of time via delay scaling (an appeal of conventional AUC metrics), data were log10-transformed (AUClog⁷⁷). Values ranged from 0

to 1, with smaller AUClogd values representing steeper discounting and thus preference for immediate (smaller, sooner) reward.

Behavioral Inhibitory System/Behavioral Activation System (BIS/BAS) Scale

Adolescent participants completed the BIS/BAS⁷⁸. The BIS/BAS is a 20-item self-report measure answered on a 4-point Likert scale. The BIS is a single, 7-question scale that probes behavioral and emotional responsivity to punishment. Conversely, the BAS is comprised of 3 subscales: Reward Responsiveness, Drive and Fun Seeking. A higher BIS score reflects aversion to and avoidance of potential punishment, while higher BAS subscale scores reflect positive emotionality (Reward Responsiveness) and behavioral approach (Drive and Fun Seeking) in the context of potential rewards.

IQ and pubertal development measures

Full-scale IQ (FSIQ) was estimated using the Kaufman Brief Intelligence Test (KBIT), Second Edition⁷⁹. Adolescents completed the Pubertal Development Scale (PDS)^{80,81} as a proxy assessment of physical development via Tanner stage⁸⁰.

Wheel of Fortune (WOF) Task

The WOF task was completed during functional neuroimaging. This well-validated paradigm has been used to probe the neural bases of reward responsivity and risky-decision making under conditions of probabilistic reward versus penalty in both adults^{60,61,82,83} and adolescents^{60,83,84}. A modified version of this task was used in this study to probe reinforcing outcomes (i.e., winning or losing) (Figure 1; see Supplement, section S1.1, for further description of the WOF task).

Participants were guided through an in-scanner practice (during the structural MRI scan)

prior to the actual task to ensure their understanding of how to perform the task. Prior to each run, participants were encouraged to maximize their hypothetical gains and/or exceed their previous total winnings. The task was implemented in E-Prime 2.0, and stimuli were presented via back-projection onto a screen viewed in a mirror mounted to the head coil. A slow event-related design with temporal jitter provided by a variable inter-trial fixation of 2500–10,000 ms based on a Poisson distribution was utilized.

Contrasts of interest for the selection and feedback phases were High-reward/risk>Low-reward/risk and Win>Lose, respectively. Behavioral data analyses considered the percentage of high-reward/risk selections and the average response times (RT) for high-reward/risk and low-reward/risk selections as well as the average RT across all selections.

Figure 1. *Wheel of fortune (WOF) task. (A) Task stimuli illustrating two trial types (90/10 and 70/30), and highlighting high-reward/risk selections for each of these trial types. (B) Example of a single trial and its timing. Pressing the left button, corresponding to the smaller, magenta portion of the wheel, represents selection of the high-reward/risk option (10% chance of receiving \$9), over the low-reward/risk option depicted in blue (90% chance of receiving \$1). Analyses reported examined Selection and Feedback phases of the task.*

MRI Protocol

Data acquisition

During the baseline visit, structural and functional images were acquired on a Siemens TIM Trio 3T scanner using a 12-channel head coil. During three runs of the WOF task, functional images were collected using a T2*-weighted gradient-echo planar imaging (EPI) sequence (interleaved slice acquisition, 47 axial slices per volume, TR=2500ms, TE=30ms, TA=2.48ms, slice thickness=3mm, voxel size= $3.0 \times 3.0 \times 3.0\text{mm}^3$, FoV= $192 \times 192\text{mm}^2$, flip angle= 90°).

High-resolution structural images were obtained using a T1-weighted magnetization prepared rapid acquisition gradient echo (MPRAGE) sequence (176 sagittal slices:

TR/TE/TI=920/2.52/900 ms, flip angle= 9° , slice thickness=1.0 mm, FOV= $250 \times 250\text{mm}^2$, matrix of 256×256 for an effective spatial resolution of $0.97 \times 0.97 \times 1.0\text{mm}^3$).

fMRI Data Preprocessing

Image preprocessing and statistical analyses were carried out using SPM8 (<http://www.fil.ion.ucl.ac.uk/spm>). Preprocessing included correction for interleaved slice timing, realignment of all images to the mean fMRI image to correct for head motion artifacts between images, and coregistration of realigned images to the anatomical MPRAGE. The MPRAGE was segmented and transformed into Montreal Neurological Institute (MNI) standard stereotactic space using non-linear warping. Lastly, these transformation parameters were

applied to normalize the functional images into MNI space, and the data were spatially smoothed using a Gaussian kernel of 6 mm³ FWHM. A scrubbing algorithm utilizing framewise displacement was implemented to assess participant movement during the fMRI scans⁸⁵. Participants included in analyses demonstrated less than 1mm displacement in fewer than 20% of their total volumes across all three runs of the task.

Statistical Analyses

Imaging data

First-level statistical analyses of imaging data included regressors encoding for trials during which the subject chose either the 10% or 30% probability (High-reward/risk) or the 70% or 90% probability (Low-reward/risk). Regressors of interest also included feedback trials on which subjects won (Win) or lost (Lose). Six translations and rotations modeling participant motion calculated during realignment were included as nuisance regressors.

Contrasts of interest examined whole brain activation for high-reward/risk compared to low-reward/risk trials (High-reward/risk>Low-reward/risk), and winning versus losing outcomes (Win > Lose). Regressors were convolved with the canonical hemodynamic response function. A temporal high-pass filter of 128s was applied to the data to eliminate low-frequency noise (e.g., MRI signal drift). First-level contrasts of interest were used in a second-level analyses for comparisons between SI and SN groups. The initial cluster defining threshold was $p<.001$, with a cluster extent of 10 voxels (voxel size=2.0mm isotropic). Corrections for multiple comparisons were made using a cluster-level FWE threshold of $p<.05$. Macro-anatomical labels reported are based on peak coordinates and were assigned by the Harvard-Oxford Cortical/Subcortical Structural atlases^{86–89}, supplemented with labels from *Atlas of the Human Brain, 4th edition*⁹⁰.

Demographics and behavioral data

Statistical analyses were performed using R v.3.5.1. Dependent variables were free from outliers and normality was examined. In the SN group, SES was negatively skewed (Shapiro-Wilk $W=.88$, $p<.001$). BAS fun-seeking scores were non-normally distributed in both groups (SI: $W=.876$, $p<.05$; SN: $W=.934$, $p<.05$); BIS was non-normal for SI ($W=.897$, $p<.05$) and BAS reward responsivity for SN ($W=.923$, $p<.05$). Further, the percent of high-reward/risk selections in the WOF task was positively skewed in both groups (SI: $W=.757$, $p<.001$; SN: $W=.873$, $p<.001$).

Standard transformations for the above dependent variables did not correct distributions; thus, the between-group comparisons were performed using non-parametric Mann-Whitney U test. Alpha was set at $p=0.05$, and Bonferroni correction for multiple comparisons was applied where noted (e.g., BAS adjusted statistical threshold of $.05/3=.0167$).

With the exception of the group comparisons for DUSI-R APD, all statistical tests were two-tailed. We used one-tailed tests in comparing groups on this measure given a priori evidence of directionality (i.e., DUSI-R APD severity, reflected by higher scores, positively predicts SU⁶⁷).

RESULTS

Demographics

SI and SN groups were similar for age, sex, PDS, SES, and race/ethnicity (Table 1). Although mean IQ was lower in SI than SN youth, the difference did not reach statistical significance ($t(68)=1.92$, $p=.059$). Based on reported associations between cognitive ability and measures of decision-making as well as risk-taking^{91–95}, we explored whether FSIQ was associated with risk-taking behavior on the WOF task. Given that the percentage of high risk decisions across all runs

of the task was positively correlated with IQ ($r_s=.28$, $p=.021$) (see Figure S1A, Table S2), an effect driven by the percent of high risk decisions in Run 1 (Figure S1B), we treated IQ as a covariate of no interest in imaging analyses presented here. Imaging results for analyses without IQ as a covariate are presented in the supplementary materials (section S2.1, Table S5. Figure S2).

Table 1. Participant characteristics at initial assessment

	All participants <i>N</i> =70	SI <i>n</i> =27	SN <i>n</i> =43	Test statistic	<i>p</i>
Age at scan					
Mean(SD)	12.7(.66)	12.9(.61)	12.7(.69)	$t(68)=-1.57$	.12
Range	11.1-14.0	11.5-14.0	11.1-14.0		
Sex				$\chi^2=.045(1)$	.83
Females, <i>n</i> (%)	40(57%)	15(56%)	25(58%)		
Males, <i>n</i> (%)	30(43%)	12(44%)	18(42%)		
PDS					
Mean(SD)	2.3(.67)	2.4(.64)	2.2(.68)	$t(68)=-1.42$	.16
Females	2.58(.67)	2.72(.65)	2.49(.68)	$t(38)=-1.07$	.29
Males	1.94 (.47)	2.1(.46)	1.83(.47)	$t(28)=-1.55$	.13
Race, <i>n</i>(%)				$\chi^2(3)=3.75$	.29
African American	19(27%)	6(22.2%)	13(20.2%)		
Caucasian	42(60%)	15(55.6%)	27(62.8%)		
Hispanic/Latino	4(6%)	3(11.1%)	1(2.3%)		
Other	5(7%)	3(11.1%)	2(4.7%)		
FSIQ				$t(68)=1.92$	.059
Mean(SD)	112.1(15.3)	107.7(14.4)	114.8(15.4)		
SES Index <i>z</i>-score					
Median(range)	.22(-2.7-1.45)	.22(-2.0-1.45)	.26 (-2.68-1.36)	$U=599$	.82
Parental education, years, Mean(SD)	16.6(2.8)	16.5 (2.8)	16.6(2.8)		
Household income, Median	100,000-149,000	100,000-149,000	100,000-149,000		

PDS=pubertal development scale, FSIQ=full-scale IQ, SES=socioeconomic status.

Among SI adolescents ($n=27$), 12 (44%) and 15 (56%) reported initiation at Wave 2 and Wave 3 follow-up, respectively. While different in age of initiation (Wave 2: $M=14.79$, $SD=.41$; Wave 3: $M=15.78$, $SD=.70$; $t(25)=-4.3$, $p<.001$), initiators at both timepoints were similar in demographic characteristics (sex ($\chi^2(1)=1.08$, $p=.299$); race/ethnicity ($\chi^2(1)=4.72$, $p=.19$); age at initial assessment ($t(25)=.50$, $p=.62$); IQ ($t(25)=-1.69$, $p=.10$); pubertal development ($t(25)=1.66$, $p=.11$); SES ($t(25)=-1.26$, $p=.22$)). (See Supplement section, S2.2–S2.4 for SI group SU details).

Behavioral Results

DUSI-R APD, DD, and BIS/BAS

A one-tailed independent samples t -test showed adolescents in the SI group had significantly higher scores on the DUSI-R APD compared to the SN group ($t(67)=-1.89$, $p=.03$) (Table 2), suggestive of increased problematic behavior in domains predictive of a future SUD. The SI and SN groups did not differ in discounting behavior (Table 2) indicating similar preferences for immediate rewards. Compared to the SN group, SI adolescents had significantly higher scores on the BAS Drive ($t(68)=-2.6$, $p=.012$) and Fun Seeking ($U=362.5$, $p=.008$) scales, but did not differ for BAS Reward Responsiveness (Table 2).

Table 2. *DUSI-R APD, DD, and BIS/BAS results*

	All participants $N=70$	SI $n=27$	SN $n=43$	Test statistic	p
DUSI-R APD		$n=26$			
Mean (SD) $N=69$	16.04(10.28)	19(11.5)	14.25(9.1)	$t(67)=-1.9$	.031*
DD, AUClogd		$n=25$	$n=36$		
Mean(SD) $N=61$	.55(.19)	.54(.19)	.55(.19)	$t(59)=.22$	.83
BAS Drive	9.7	10.7	9.1	$t(68)=-2.6$	.012†

Mean(SD)	(2.7)	(2.4)	(2.7)		
BAS Fun-seeking	11	13	11	$U=362.5$	.008†
Median(Range)	(4–15)	(6–15)	(4–15)		
BAS Reward Responsivity				$U=516.5$	.44
Median(Range)	18(14–20)	18(14–20)	18(14–20)		
BIS				$U=552.5$	.74
Median(Range)	20(12–27)	20(12–24)	18(15–27)		

*Note. Group comparisons for the DUSI-R APD used a one-tailed independent samples t-test. Comparisons for AUClogd and the BAS Drive scale used two-tailed independent samples t-tests; and those for the BAS Fun Seeking and Reward Responsivity scales and the BIS were conducted using two-tailed Mann-Whitney U-tests. *= $p<.05$; †= $p<.0167$ Bonferroni-corrected threshold for BAS subscales. DUSI-R APD=Drug Use Screening Inventory, Revised, Absolute Problem Density; DD, AUC=delay discounting, area under the curve; BAS=Behavioral Activation System; BIS=Behavioral Inhibition System.*

WOF task behavior

The groups made similar proportions of high-reward/risk selections ($Z=.537$, $p=.70$) (Table 3). A two-way repeated measures ANOVA was used to examine the effect of group (SI vs. SN) and selection type (high-reward/risk vs. low-reward/risk) on RT. A main effect of selection type was found, with both SI and SN groups taking significantly more time to make riskier selections compared to safer ones ($F(1,68)=65.7$, $p<.0001$). There was no significant main effect of group ($F(1,68)=.0006$, $p=.98$), nor was there a significant group $\times$ selection type interaction ($F(1,68)=.007$, $p=.93$).

Table 3. WOF task behavior: descriptive statistics

	All participants $N=70$	SI $n=27$	SN $n=43$
High-reward/risk selections, %			
Median(range)	10(1-59)	10(1-59)	11(1-55)
High-reward/risk selections			
RT(ms)			
Mean(SD)	1240(410)	1240(390)	1240(420)
Low-reward/risk selections			
RT(ms)			
Mean(SD)	990(300)	990(270)	990(310)

All selections			
RT(ms)			
Mean(SD)	1120(330)	1110(320)	1120 (350)

FMRI results

Compared to SN youth, SI adolescents demonstrated less activation in the left insula when selecting high-reward/risk versus low-reward/risk options. Additionally, when presented with winning versus losing feedback, SI adolescents showed less activation in the left cingulate gyrus, relative to their SN peers (Table 4, Figure 2). There were no regions for which SI youth showed greater activation compared to SN youth for either contrast (see Supplement, Section S21 and Table S5 for presentation of within-group fMRI results). Results surviving corrections for multiple comparisons are presented in Table 4 and Figure 2.

Table 4. *Summary of SN > SI cluster-level corrected results for two contrasts of interest (IQ as covariate of no interest). Initial cluster defining threshold= $p < 0.001$, $k = 10$ voxels. Reported results survive FWE cluster-correction ($p < .05$).*

Region	Cluster size	MNI coordinates			Z	t	Corrected p-value (FWE)
		x	y	z			
High-reward/risk > Low-reward/risk							
Left insular cortex	492	-34	8	6	3.91	4.16	.001
Win > Lose							
Left anterior cingulate gyrus	207	-4	20	38	4.22	4.54	.031

Figure 2. Between-group results for which SN participants demonstrate increased activation relative to SI adolescents. Interaction charts depict mean parameter estimates (error bars represent standard errors) for A) High-reward/risk>Low-reward/risk, left insula; and B) Win>Lose, left anterior cingulate cortex (ACC). FSIQ as covariate of no interest. Initial cluster defining threshold= $p < 0.001$, $k = 10$ voxels. Results survive FWE cluster-correction at $p < .05$.

Exploratory analyses: Post-hoc tests of parameter estimates

Visual inspection of the parameter estimates in Figure 2 suggest that the significant between-group results for the contrasts of interest may be driven by differences in how the groups are

processing each trial type during selection and feedback. To probe these potential differences, exploratory tests were conducted. All tests were two-tailed. Follow-up independent-samples *t*-tests showed groups significantly differed in activity during low risk ($t(68)=2.97, p=.004$) but not high-risk ($t(68)=-1.25, p=.22$) trials (Figure 2A). Paired-samples *t*-tests examining within-group processing of high- compared to low-risk trials showed significant differences for SN ($t(42)=-3.89, p=.0004$) but not SI youth ($t(26)=1.95, p=.062$). Independent-samples *t*-tests showed groups significantly differed for losing ($t(68)=-2.17, p=.034$) but not winning ($t(68)=1.84, p=.07$) feedback (Figure 2B). Paired-samples *t*-tests revealed SI youth ($t(26)=-4.35, p=.0002$) significantly differed for activity when processing winning and losing trial types, while SN do not ($t(42)=1.59, p=.12$). Implications of these exploratory analyses as they relate to the between-group findings for High-risk>Low-risk and Win>Lose contrasts are treated in the Discussion.

DISCUSSION

This study aimed to elucidate neural and behavioral factors that may identify adolescents who are at risk for SU initiation. We examined neural activity during a reward-based decision-making task in SU-naïve early adolescents who either endorsed or denied initiating alcohol and/or drugs during follow-up assessments. Despite similar performance on the risk-taking task, adolescents who went on to report SU initiation showed distinct patterns of activity in left-lateralized insula during risky decisions, and in the left anterior cingulate cortex (ACC) when presented with rewarded outcomes, compared to their non-initiating peers. Furthermore, adolescents who endorsed SU initiation at follow-up reported early increased problematic behavior in domains predictive of a future SUD (DUSI-R APD) as well as greater affective and behavioral responsivity to cues of impending reward (BAS Fun Seeking and Drive) compared to their SN peers. Overall, these findings are consistent with the premise that neural differences in PFC

regions may occur *prior* to SU initiation and confer vulnerability to SUDs^{96,97}. The findings reported here furthermore lend support to models suggesting divergent neurodevelopmental trajectories may be present prior to SU; as such, due to their malleability, these structures may be targets for early intervention⁹⁸. Critically, the results here underscore the potential utility of neuroimaging in identifying potential precursors of risk for early SU initiation and SUDs.

WOF task: Behavior and brain

Task-based risk-taking was similar between groups. Compared to low-reward/risk options, selection of high-reward/risk options was accompanied by longer deliberation, an effect consistent with previous studies^{61,82,99,100}. Despite equivalent performance, SI and SN groups differed in neural activity underlying *making* risky selections and *processing* rewarded outcomes.

The contrast high-reward/risk > low-reward/risk was greater in left insular cortex for SN compared to SI adolescents (Figure 2A). Post-hoc exploratory analyses indicated that during risk-taking SI and SN groups differed in patterns of activation depending on whether they chose a high- or low-risk option, suggesting that the marked difference in responsivity to low-reward/risk trials drives the significant between-group result for the High-reward/risk>Low-reward/risk contrast. Within-group exploratory analyses revealed only SN adolescents showed significantly different activation for high-reward/risk compared to low-reward/risk trials, while activation for the two trial types did not significantly differ for the SI group.

During decision-making, the insular cortex plays a role in refocusing attention based on salience, evaluating risk, inhibiting action, and processing outcomes^{101–104}. Attenuated insula activity is associated with increased real-world risk-taking among adolescents^{105,106}, and aberrant insula engagement in processing salient stimuli is observed in individuals with addiction^{107–109}.

Reduced activation of the anterior insula has been found to play an important role in adolescent risky decision making in comparison to adults and is linked to more emotionally driven decisions¹¹⁰. Considered within the context of such studies, our results may be indicative of relative immaturity in the SI group in a region that plays an important role in evaluating degree of risk¹¹¹ and as such may have been a contributing factor in the early initiation of substances of abuse in these adolescents.

The contrast Win>Lose was greater for SN compared to SI adolescents in the left ACC. Inspection of parameter plots suggested that while processing outcomes related to gain or loss, SN and SI adolescents demonstrated differing patterns of responses in this region (Figure 2B). Post-hoc exploratory analyses revealed groups significantly differed for activation during losing, but not winning, feedback. Further, only SI youth showed significantly different activation for Lose compared to Win trials. Increased ACC activity has previously been associated with processing gains (relative to no gains) in a gambling task in adolescents¹⁰⁰. Individuals with established SUDs show impairments in decision-making¹¹², altered ACC structure¹¹³ and differences in brain activity during risk-taking¹¹⁴. Specifically, individuals with SUDs display not only greater substance-related cue-induced ACC activity during active use¹¹⁵, but also blunted ACC activity during decision-making while abstinent¹¹⁶, an effect which predicts craving, length of time to relapse and relapse severity¹¹⁷. Importantly, some of these differences may be evident prior to development of AUD/SUD including alterations in ACC neuroanatomy¹¹⁸ and suggest increased vulnerability to SUDs (e.g., youth with positive family history of alcoholism demonstrate hyper-activation during risk taking compared to youth with negative family history)¹¹⁹.

Both the insula and ACC are implicated in reward-related decision-making^{60,82,105,111,120,121} and the ACC exhibits age-related decreases in activity during risky-decision-making¹²², while adolescents, compared to adults, show increased activation in both ACC and insula during reward processing¹²³. Moreover, as hubs in the functional salience network, the anterior insula and ACC^{124,125} are believed to integrate automatic, bottom-up detection of relevant internal and external stimuli with cognitive, top-down processing¹²⁶. The salience network is implicated not only in altered cue-reactivity among individuals with SUD^{107,108}, but may play a contributory etiological role in early SU and transition to SUD¹²⁷. While others have established that adults with SUD demonstrate aberrant patterns of insular and cingulate activity during risky decision-making¹²⁸ and that reduced insular activity during risk-related processing is predictive of relapse¹²⁹, our results suggest that variability in insular and ACC activity is present in individuals at risk for SUD even *before* substance initiation.

The exploratory results intriguingly suggest SI youth may be more sensitive to the distinction between wins and losses, though this remains to be empirically tested in planned comparisons correcting for multiple comparisons. It is also possible that steep hypoactivation of the ACC in SI adolescents in the context of rewarding outcomes indicates an increased threshold for rewarding stimuli (consistent with elevated BAS fun-seeking scores SI youth). These group differences may reflect differences in outcome monitoring and processing¹³⁰ and awareness of outcomes¹³¹, which serve in part to guide behavior^{132,133}. A notable consistency between present findings and previous studies is that youth with differential risk for SU demonstrate similar task performance but differences in patterns of brain activation across a variety of tasks^{134,135}. Thus, early disruptions in PFC function, including ACC, may contribute to a constellation of

impairments including aberrant response inhibition and salience assignment⁹⁷ and ultimately to real-world risky decision making.

Measures of risk and reward-sensitivity: DUSI-R, BIS/BAS, and DD

Compared to their SN peers, SI adolescents showed elevated DUSI-R absolute problem density (APD) scores. Because the APD score is a multi-dimensional construct, quantifying adolescent difficulties across health, psychosocial and psychiatric domains associated with SUD⁶⁶, elevated APD scores in SI adolescents may reflect increased relative risk for SU. It should be noted, however, that 69% of SI youth would *not be considered high risk* according to the previously established cut-off score of 24⁶⁷; further, similar proportions of adolescents in each group scored ≥ 24 (SI: 30% (8/27); SN: 19% (8/43); $\chi^2=.70(1)$, $p=.40$). In contrast, the DUSI APD validation study in youth aged 12-14 years predicted SUD (according to DSM-III-R criteria) by age 19 with 73% accuracy⁶⁷. Although early initiation is itself a key risk factor for SUDs, as noted earlier, indices of brain function may be more sensitive and specific indicators of risk than outward behavior and may be apparent *prior to SU onset*. If so, such measures may identify potent targets for effective interventions to prevent SUDs.

Prior to SU, SI adolescents (relative to their SN peers) showed elevated BAS Fun Seeking and Drive scores. Elevated scores on these two BAS scales have been associated with low levels of harm avoidance⁷⁸ and problematic alcohol use, including among adolescents^{136–139}. Further, these subscales were positively correlated with adolescent risky choices during a win-only (but not a lose-only) version of a WOF task⁹⁹. Thus, elevated scores on these BAS subscales prior to SU may reflect a propensity towards affective and behavioral responsivity to rewarding stimuli in SI

youth, which may bias individuals toward greater real-world situational risk-taking and decreased harm avoidance¹⁰¹.

SI and SN youth did not differ on performance on the DD task, suggesting similar preference for immediate rewards under current task parameters. Unlike the BIS/BAS and DUSI-R which assess real-world preference and situationally-based behavior, the laboratory DD task (like WOF) may lack the sensitivity to detect group differences prior to initiation¹⁴⁰. The absence of differences between SN and SI participants on the DD and WOF task performance, in the context of functional differences during WOF performance, lends further support to the notion that brain indices may be more sensitive to risk prior to the onset of SU, compared to behavioral measures.

Strengths and Limitations

An important strength of the current study is the stringent inclusion criteria implemented to ensure SU-naïve status of youth at initial assessment, and the requirement of convergent responses on two SU measures (DUSI-R and TAD) to classify youth at follow-up. In contrast, previous studies examining “SU-naïve” youth include those who report “little to no” alcohol use¹⁴¹, or who do not report “significant”^{59,63,142} or “heavy” alcohol or drug use¹⁴³. Others rely on urine drug screening at scanning time⁸⁴, which, for many drugs, capture only recent use¹⁴⁴ and are not reflective of patterns of use over time.

Another strength of the current study is the narrow age range at baseline (11-13-year-olds). While previous studies enrolled participants with a more distributed age range^{60,84,142}, we restricted eligibility at enrollment to a much smaller range in an effort to capture information regarding early initiation and minimize potential age-related confounds in neurodevelopment. Finally, the current sample of adolescents were well-characterized using a battery capturing a

variety of factors presumed to confer risk for or resilience to early SU, including preference for immediate gratification (DD), affective and behavioral responsivity to rewards and punishment (BIS/BAS), and multidimensional risk for SU problems (DUSI-R).

On the other hand, by analyzing the selection phase of the WOF in a version of the task that consistently coupled high reward with low probability and low reward with high probability we were unable to dissociate between patterns of activation associated with reward versus risk. Although this limitation is not unique to the current study¹⁴⁵, it is unclear here whether between-group differences in insular cortex were driven by reward sensitivity or risky decision-making. Inclusion of choices with equal probability of high/low reward (i.e., 50/50 wheels), as in the ‘classic’ WOF task, would have permitted testing the relative contributions of reward magnitude independent of perceived risk⁶¹. It is important to note, however, that estimation of reward value and tendency towards risk outside of the laboratory may not be not entirely separable either; decisions with greater reward potential, whether adaptive (e.g., approaching a classmate to initiate a conversation) or maladaptive (e.g., underage alcohol consumption) are inherently accompanied by risk (e.g., social consequences such as peer rejection; or adverse physiological impacts of alcohol consumption and parental or school punishment for drinking).

The current study identified youth who initiated at different ages (initiation at approximately 18- vs. 36-months follow-up), which may also limit the interpretation of our outcomes. Although the two SI subgroups (i.e., earlier vs. later initiators) were similar in demographic, physical and cognitive characteristics, as well as task-based behavior and BIS/BAS scores, those who reported earlier initiation (approximately 18 month follow-up) scored higher on DUSI-APD, indicative of greater risk in domains that precede or co-occur with problematic SU. Due to concerns regarding statistical power, we were unable to compare SI

subgroups on brain activation during the WOF task. It is recommended that future studies recruit greater numbers of participants who are likely to be assigned to one of these two SU subgroups, in order to increase the statistical power to prospectively examine group differences in neural activation among individuals who initiate at different stages of adolescence.

Relatedly, the current study is unable to determine pathways to SU escalation, and SUD. SU initiation itself is not necessarily indicative of continued or escalated use or the eventual entrenchment of pathways that might be specific to SUD risk. The elucidation of factors that give rise to such pathways, including early brain biomarkers, may provide a much richer understanding of how brain functioning in SU-naïve adolescents portends subsequent life course outcomes.

CONCLUSIONS

During adolescence, while exploratory behavior like risk-taking is normative and even adaptive under certain circumstances, the brain changes that drive risk-taking during this developmental period may also facilitate vulnerabilities for maladaptive risk-taking including early SU initiation and escalation. Earlier SU is associated with adverse outcomes and dramatically elevated risk for SUDs, which present a lifelong burden to the individual. Early SU may also alter the neurodevelopment of regions critical to impulse control, further compounding predispositions of youth to risk-taking^{33,44,146}. Thus, identification of neuroendophenotypes for early SU initiation is crucial to inform targeted interventions among at-risk youth *prior to* SU entrenchment. At a minimum, delaying SU in these individuals would allow time for cognitive control functions supported by the developing PFC to come on board and may thus confer protection against SUDs in adulthood¹²⁷. The current study corroborates and extends the

literature by demonstrating that variability in activity in ACC and insula—key regions known to support reward- and risk-related decision-making—may distinguish SU-naïve youth who initiate SU early from those who remain abstinent.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was supported by the National Institutes of Health/National Institute on Alcohol Abuse and Alcoholism [1R01AA019983-01; 3R01AA019983-02S1; 5F31AA023462-02], NIH/NCATS [1KL2RR031974-01, and NIH/NICHHD [2P30HD040677-11].

REFERENCES

1. Arnett, J. (1992) Reckless behavior in adolescence: A developmental perspective. *Developmental Review*, **12**, 339–73. [https://doi.org/10.1016/0273-2297\(92\)90013-R](https://doi.org/10.1016/0273-2297(92)90013-R)
2. Steinberg, L. (2004) Risk taking in adolescence: What changes, and why? *Annals of the New York Academy of Sciences*, **1021**, 51–8. <https://doi.org/10.1196/annals.1308.005>
3. Duell, N. and Steinberg, L. (2019) Positive risk taking in adolescence. *Child Development Perspectives*, **13**, 48–52. <https://doi.org/10.1111/cdep.12310>
4. Telzer, E.H. (2016) Dopaminergic reward sensitivity can promote adolescent health: A new perspective on the mechanism of ventral striatum activation. *Developmental Cognitive Neuroscience*, **17**, 57–67. <https://doi.org/10.1016/j.dcn.2015.10.010>
5. van Duijvenvoorde, A.C.K., Peters, S., Braams, B.R. and Crone, E.A. (2016) What

- motivates adolescents? Neural responses to rewards and their influence on adolescents' risk taking, learning, and cognitive control. *Neuroscience & Biobehavioral Reviews*, **70**, 135–47. <https://doi.org/10.1016/j.neubiorev.2016.06.037>
6. Ellis, B.J., Del Giudice, M., Dishion, T.J., Figueredo, A.J., Gray, P., Griskevicius, V. et al. (2012) The evolutionary basis of risky adolescent behavior: Implications for science, policy, and practice. *Developmental Psychology*, **48**, 598–623. <https://doi.org/10.1037/a0026220>
 7. Gullo, M.J. and Dawe, S. (2008) Impulsivity and adolescent substance use: Rashly dismissed as “all-bad”? *Neuroscience and Biobehavioral Reviews*, **32**, 1507–18. <https://doi.org/10.1016/j.neubiorev.2008.06.003>
 8. Roisman, G.I., Masten, A.S., Coatsworth, J.D. and Tellegen, A. (2004) Salient and emerging developmental tasks in the transition to adulthood. *Child Development*, **75**, 123–33. <https://doi.org/10.1111/j.1467-8624.2004.00658.x>
 9. Crews, F.T., Vetreno, R.P., Broadwater, M.A. and Robinson, D.L. (2016) Adolescent Alcohol Exposure Persistently Impacts Adult Neurobiology and Behavior. *Pharmacological Reviews*, **68**, 1074–109. <https://doi.org/10.1124/pr.115.012138>
 10. Ewing Feldstein, S.W. and Miller, W.R. (2006) Substance use and risk-taking among adolescents. *Journal of Mental Health*, **15**, 633–43. <https://doi.org/10.1080/09638230600998896>
 11. French, M.T. and Maclean, J.C. (2006) Underage alcohol use, delinquency, and criminal activity. *Health Economics*, **15**, 1261–81. <https://doi.org/10.1002/hec.1126>
 12. Crockett, L.J., Raffaelli, M. and Shen, Y.L. (2006) Linking self-regulation and risk proneness to risky sexual behavior: Pathways through peer pressure and early substance

- use. *Journal of Research on Adolescence*, **16**, 503–25. <https://doi.org/10.1111/j.1532-7795.2006.00505.x>
13. Tapert, S.F., Arons, G.A., Sedlar, G.R. and Brown, S.A. (2001) Adolescent substance use and sexual risk-taking behavior. *Journal of Adolescent Health*, **28**, 181–9. [https://doi.org/10.1016/S1054-139X\(00\)00169-5](https://doi.org/10.1016/S1054-139X(00)00169-5)
14. Cunningham, R.M., Ranney, M., Newton, M., Woodhull, W., Zimmerman, M. and Walton, M.A. (2014) Characteristics of youth seeking emergency care for assault injuries. *Pediatrics*, **133**. <https://doi.org/10.1542/peds.2013-1864>
15. Bonomo, Y., Coffey, C., Wolfe, R., Lynskey, M., Bowes, G. and Patton, G. (2001) Adverse outcomes of alcohol use in adolescents. *Addiction*, **96**, 1485–96. <https://doi.org/10.1046/j.1360-0443.2001.9610148512.x>
16. Toumbourou, J.W., Stockwell, T., Neighbors, C., Marlatt, G.A., Sturge, J. and Rehm, J. (2007) Interventions to reduce harm associated with adolescent substance use. *Lancet*, **369**, 1391–401. <https://doi.org/10.1016/S0140>
17. Hall, W.D., Patton, G., Stockings, E., Weier, M., Lynskey, M., Morley, K.I. et al. (2016) Why young people’s substance use matters for global health. *The Lancet Psychiatry*, **3**, 265–79. [https://doi.org/10.1016/S2215-0366\(16\)00013-4](https://doi.org/10.1016/S2215-0366(16)00013-4)
18. Dawson, D.A., Goldstein, R.B., Patricia Chou, S., June Ruan, W. and Grant, B.F. (2008) Age at first drink and the first incidence of adult-onset DSM-IV alcohol use disorders. *Alcoholism: Clinical and Experimental Research*, **32**, 2149–60. <https://doi.org/10.1111/j.1530-0277.2008.00806.x>
19. Grant, B.F. and Dawson, D.A. (1998) Age of onset of drug use and its association with DSM-IV drug abuse and dependence: Results from the national longitudinal alcohol

- epidemiologic survey. *Journal of Substance Abuse*, **10**, 163–73.
[https://doi.org/10.1016/S0899-3289\(99\)80131-X](https://doi.org/10.1016/S0899-3289(99)80131-X)
20. Grant, B.F., Stinson, F.S. and Harford, T.C. (2001) Age at onset of alcohol use and DSM-IV alcohol abuse and dependence: A 12-year follow-up. *Journal of Substance Abuse*, **13**, 493–504. [https://doi.org/10.1016/S0899-3289\(01\)00096-7](https://doi.org/10.1016/S0899-3289(01)00096-7)
 21. King, K.M. and Chassin, L. (2007) A prospective study of the effects of age of initiation of alcohol and drug use on young adult substance dependence. *Journal of Studies on Alcohol and Drugs*, **68**, 256–65. <https://doi.org/10.15288/jsad.2007.68.256>
 22. Stone, A.L., Becker, L.G., Huber, A.M. and Catalano, R.F. (2012) Review of risk and protective factors of substance use and problem use in emerging adulthood. *Addictive Behaviors*, **37**, 747–75. <https://doi.org/10.1016/j.addbeh.2012.02.014>
 23. Anthony, J.C. and Petronis, K.R. (1995) Early-onset drug use and risk of later drug problems. *Drug and Alcohol Dependence*, **40**, 9–15. [https://doi.org/10.1016/0376-8716\(95\)01194-3](https://doi.org/10.1016/0376-8716(95)01194-3)
 24. Brook, D.W., Brook, J.S., Zhang, C., Cohen, P. and Whiteman, M. (2002) Drug Use and the Risk of Major Depressive Disorder, Alcohol Dependence, and Substance Use Disorders. *Archives of General Psychiatry*, **59**, 1039.
<https://doi.org/10.1001/archpsyc.59.11.1039>
 25. Donoghue, K., Rose, H., Boniface, S., Deluca, P., Coulton, S., Alam, M.F. et al. (2017) Alcohol Consumption, Early-Onset Drinking, and Health-Related Consequences in Adolescents Presenting at Emergency Departments in England. *Journal of Adolescent Health*, **60**, 438–46. <https://doi.org/10.1016/j.jadohealth.2016.11.017>
 26. DuRant, R.H., Smith, J.A., Kreiter, S.R. and Krowchuk, D.P. (1999) The relationship

- between early age of onset of initial substance use and engaging in multiple health risk behaviors among young adolescents. *Archives of Pediatrics & Adolescent Medicine*, **153**, 286–91. <https://doi.org/10.1001/archpedi.153.3.286>
27. Substance Abuse and Mental Health Services Administration. (2014) The TEDS Report: Age of Substance Use Initiation among Treatment Admissions Aged 18 to 30. Rockville, MD.
https://www.samhsa.gov/data/sites/default/files/WebFiles_TEDS_SR142_AgeatInit_07-10-14/TEDS-SR142-AgeatInit-2014.pdf
 28. Wells, J.E., McGee, M.A., Baxter, J., Agnew, F. and Kokaua, J. (2009) Onset and lifetime use of drugs in New Zealand: Results from Te Rau Hinengaro: The New Zealand Mental Health Survey 2003-2004. *Drug and Alcohol Review*, **28**, 166–74.
<https://doi.org/10.1111/j.1465-3362.2008.00043.x>
 29. Steinberg, L. (2008) A social neuroscience perspective on adolescent risk-taking. *Developmental Review*, **28**, 78–106. <https://doi.org/10.1016/j.dr.2007.08.002>
 30. Shulman, E.P., Smith, A.R., Silva, K., Icenogle, G., Duell, N., Chein, J. et al. (2016) The dual systems model: Review, reappraisal, and reaffirmation. *Developmental Cognitive Neuroscience*, **17**, 103–17. <https://doi.org/10.1016/j.dcn.2015.12.010>
 31. Ernst, M. (2014) The triadic model perspective for the study of adolescent motivated behavior. *Brain and Cognition*, **89**, 104–11. <https://doi.org/10.1016/j.bandc.2014.01.006>
 32. Ernst, M., Pine, D.S. and Hardin, M. (2006) Triadic model of the neurobiology of motivated behavior in adolescence. *Psychological Medicine*, **36**, 299–312.
<https://doi.org/10.1017/S0033291705005891>
 33. Casey, B.J. and Jones, R.M. (2010) Neurobiology of the adolescent brain and behavior:

- Implications for substance use disorders. *Journal of the American Academy of Child and Adolescent Psychiatry*, **49**, 1189–201. <https://doi.org/10.1016/j.jaac.2010.08.017>
34. Somerville, L.H. and Casey, B.J. (2010) Developmental neurobiology of cognitive control and motivational systems. *Current Opinion in Neurobiology*, **20**, 236–41.
 35. Sowell, E.R., Thompson, P.M., Tessner, K.D. and Toga, A.W. (2001) Mapping continued brain growth and gray matter density reduction in dorsal frontal cortex: Inverse relationships during postadolescent brain maturation. *Journal of Neuroscience*, **21**, 8819–29. <https://doi.org/10.1523/jneurosci.21-22-08819.2001>
 36. Sowell, E.R., Thompson, P.M., Holmes, C.J., Jernigan, T.L. and Toga, A.W. (1999) In vivo evidence for post-adolescent brain maturation in frontal and striatal regions. *Nature Neuroscience*, **2**, 859–61. <https://doi.org/10.1038/13154>
 37. Satterthwaite, T.D., Wolf, D.H., Erus, G., Ruparel, K., Elliott, M.A., Gennatas, E.D. et al. (2013) Functional maturation of the executive system during adolescence. *Journal of Neuroscience*, **33**, 16249–61. <https://doi.org/10.1523/JNEUROSCI.2345-13.2013>
 38. Luna, B., Marek, S., Larsen, B., Tervo-Clemmens, B. and Chahal, R. (2015) An integrative model of the maturation of cognitive control. *Annual Review of Neuroscience*, **38**, 151–70. <https://doi.org/10.1146/annurev-neuro-071714-034054>
 39. Ordaz, S.J., Foran, W., Velanova, K. and Luna, B. (2013) Longitudinal growth curves of brain function underlying inhibitory control through adolescence. *Journal of Neuroscience*, **33**, 18109–24. <https://doi.org/10.1523/JNEUROSCI.1741-13.2013>
 40. Jadhav, K.S. and Boutrel, B. (2019) Prefrontal cortex development and emergence of self-regulatory competence: The two cardinal features of adolescence disrupted in context of alcohol abuse. *European Journal of Neuroscience*, **50**, 2274–81.

<https://doi.org/10.1111/ejn.14316>

41. Pfefferbaum, A., Kwon, D., Brumback, T., Thompson, W.K., Cummins, K., Tapert, S.F. et al. (2018) Altered Brain Developmental Trajectories in Adolescents After Initiating Drinking. *American Journal of Psychiatry*, **175**, 370–80.
<https://doi.org/10.1176/appi.ajp.2017.17040469>
42. Cservenka, A. and Brumback, T. (2017) The burden of binge and heavy drinking on the brain: Effects on adolescent and young adult neural structure and function. *Frontiers in Psychology*, **8**. <https://doi.org/10.3389/fpsyg.2017.01111>
43. Spear, L.P. (2018) Effects of adolescent alcohol consumption on the brain and behaviour. *Nature Reviews Neuroscience*, **19**, 197–214. <https://doi.org/10.1038/nrn.2018.10>
44. Squeglia, L.M., Jacobus, J. and Tapert, S.F. (2009) The influence of substance use on adolescent brain development. *Clinical EEG and Neuroscience*, **40**, 31–8.
<https://doi.org/10.1177/155005940904000110>
45. Risher, M.L., Fleming, R.L., Risher, W.C., Miller, K.M., Klein, R.C., Wills, T. et al. (2015) Adolescent intermittent alcohol exposure: Persistence of structural and functional hippocampal abnormalities into adulthood. *Alcoholism: Clinical and Experimental Research*, **39**, 989–97. <https://doi.org/10.1111/acer.12725>
46. McGue, M., Iacono, W.G., Legrand, L.N., Malone, S. and Elkins, I. (2001) Origins and consequences of age at first drink. I. Associations with substance-use disorders, disinhibitory behavior and psychopathology, and P3 amplitude. *Alcoholism: Clinical and Experimental Research*, **25**, 1156–65. <https://doi.org/10.1111/j.1530-0277.2001.tb02330.x>
47. Nees, F., Tzschoppe, J., Patrick, C.J., Vollstädt-Klein, S., Steiner, S., Poustka, L. et al. (2012) Determinants of early alcohol use in healthy adolescents: The differential

- contribution of neuroimaging and psychological factors. *Neuropsychopharmacology*, **37**, 986–95. <https://doi.org/10.1038/npp.2011.282>
48. Crawford, A.M., Pentz, M.A., Chou, C.P., Li, C. and Dwyer, J.H. (2003) Parallel developmental trajectories of sensation seeking and regular substance use in adolescents. *Psychology of Addictive Behaviors*, **17**, 179–92. <https://doi.org/10.1037/0893-164X.17.3.179>
49. Genovese, J.E.C. and Wallace, D. (2007) Reward sensitivity and substance abuse in middle school and high school students. *Journal of Genetic Psychology*, **168**, 465–9. <https://doi.org/10.3200/GNTP.168.4.465-469>
50. Urošević, S., Collins, P., Muetzel, R., Schissel, A., Lim, K.O. and Luciana, M. (2015) Effects of reward sensitivity and regional brain volumes on substance use initiation in adolescence. *Social Cognitive and Affective Neuroscience*, **10**, 106–13. <https://doi.org/10.1093/scan/nsu022>
51. Kim-Spoon, J., Deater-Deckard, K., Holmes, C., Lee, J., Chiu, P. and King-Casas, B. (2016) Behavioral and neural inhibitory control moderates the effects of reward sensitivity on adolescent substance use. *Neuropsychologia*, **91**, 318–26. <https://doi.org/10.1016/j.neuropsychologia.2016.08.028>
52. Smith, G.T. and Cyders, M.A. (2016) Integrating affect and impulsivity: The role of positive and negative urgency in substance use risk. *Drug and Alcohol Dependence*, **163**, S3–12. <https://doi.org/10.1016/j.drugalcdep.2015.08.038>
53. Morgan, J.E., Bowen, K.L., Moore, S.C. and van Goozen, S.H.M. (2014) The relationship between reward and punishment sensitivity and antisocial behavior in male adolescents. *Personality and Individual Differences*, **63**, 122–7.

- <https://doi.org/10.1016/j.paid.2014.01.054>
54. Hawkins, J.D., Catalano, R.F. and Miller, J.Y. (1992) Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: Implications for substance abuse prevention. *Psychological Bulletin*, **112**, 64–105.
<https://doi.org/10.1037/0033-2909.112.1.64>
 55. Gabrieli, J.D.E., Ghosh, S.S. and Whitfield-Gabrieli, S. (2015) Prediction as a humanitarian and pragmatic contribution from human cognitive neuroscience. *Neuron*, Cell Press. **85**, 11–26. <https://doi.org/10.1016/j.neuron.2014.10.047>
 56. Gray, K.M. and Squeglia, L.M. (2018) Research Review: What have we learned about adolescent substance use? *Journal of Child Psychology and Psychiatry*, **59**, 618–27.
<https://doi.org/10.1111/jcpp.12783>
 57. Ersche, K.D., Jones, P.S., Williams, G.B., Turton, A.J., Robbins, T.W. and Bullmore, E.T. (2012) Abnormal brain structure implicated in stimulant drug addiction. *Science*, **335**, 601–4. <https://doi.org/10.1126/science.1214463>
 58. Squeglia, L.M. and Cservenka, A. (2017) Adolescence and drug use vulnerability: findings from neuroimaging. *Current Opinion in Behavioral Sciences*, **13**, 164–70.
<https://doi.org/10.1016/j.cobeha.2016.12.005>
 59. Morales, A.M., Jones, S.A., Ehlers, A., Lavine, J.B. and Nagel, B.J. (2018) Ventral striatal response during decision making involving risk and reward is associated with future binge drinking in adolescents. *Neuropsychopharmacology*, **43**, 1884–90.
<https://doi.org/10.1038/s41386-018-0087-8>
 60. Eshel, N., Nelson, E.E., Blair, J., Pine, D.S. and Ernst, M. (2007) Neural substrates of choice selection in adults and adolescents: Development of the ventrolateral prefrontal and

- anterior cingulate cortices. *Neuropsychologia*, **45**, 1270–9.
<https://doi.org/10.1016/j.neuropsychologia.2006.10.004>
61. Ernst, M., Nelson, E.E., McClure, E.B., Monk, C.S., Munson, S., Eshel, N. et al. (2004) Choice selection and reward anticipation: An fMRI study. *Neuropsychologia*, **42**, 1585–97. <https://doi.org/10.1016/j.neuropsychologia.2004.05.011>
 62. Ernst, M. and Paulus, M.P. (2005) Neurobiology of Decision Making: A Selective Review from a Neurocognitive and Clinical Perspective. *Biological Psychiatry*, **58**, 597–604.
<https://doi.org/10.1016/j.biopsych.2005.06.004>
 63. Morales, A.M., Jones, S.A., Harman, G., Patching-Bunch, J. and Nagel, B.J. (2019) Associations between nucleus accumbens structural connectivity, brain function, and initiation of binge drinking. *Addiction Biology*, e12767. <https://doi.org/10.1111/adb.12767>
 64. Fishbein, D.H., Rose, E.J., Darcey, V.L., Belcher, A.M. and VanMeter, J.W. (2016) Neurodevelopmental precursors and consequences of substance use during adolescence: Promises and pitfalls of longitudinal neuroimaging strategies. *Frontiers in Human Neuroscience*, **10**, 296. <https://doi.org/10.3389/fnhum.2016.00296>
 65. Manuck, S.B., Phillips, J.E., Gianaros, P.J., Flory, J.D. and Muldoon, M.F. (2010) Subjective socioeconomic status and presence of the metabolic syndrome in midlife community volunteers. *Psychosomatic Medicine*, **72**, 35–45.
<https://doi.org/10.1097/PSY.0b013e3181c484dc>
 66. Tarter, R.E. (1990) Evaluation and treatment of adolescent substance abuse: A decision tree method. *American Journal of Drug and Alcohol Abuse*, **16**, 1–46.
<https://doi.org/10.3109/00952999009001570>
 67. Tarter, R.E. and Kirisci, L. (2001) Validity of the Drug Use Screening Inventory for

- predicting DSM-III-R substance use disorder. *Journal of Child and Adolescent Substance Abuse*, **10**, 45–53. https://doi.org/10.1300/J029v10n04_05
68. Bucholz, K.K., Cadoret, R., Cloninger, C.R., Dinwiddie, S.H., Hesselbrock, V.M., Nurnberger, J.I. et al. (1994) A new, semi-structured psychiatric interview for use in genetic linkage studies: A report on the reliability of the SSAGA. *Journal of Studies on Alcohol*, **55**, 149–58. <https://doi.org/10.15288/jsa.1994.55.149>
69. Tarter, R.E., Laird, S.B., Bukstein, O. and Kaminer, Y. (1992) Validation of the adolescent Drug Use Screening Inventory: Preliminary findings. *Psychology of Addictive Behaviors*, **6**, 233–6. <https://doi.org/10.1037/h0080632>
70. Kirisci, L., Mezzich, A. and Tarter, R.E. (1995) Norms and sensitivity of the adolescent version of the drug use screening inventory. *Addictive Behaviors*, **20**, 149–57. [https://doi.org/10.1016/0306-4603\(94\)00058-1](https://doi.org/10.1016/0306-4603(94)00058-1)
71. Siewert, E.A., Stallings, M.C. and Hewitt, J.K. (2004) Factor structure and concurrent validity of the Drug Use Screening Inventory in a community adolescent sample. *Addictive Behaviors*, **29**, 627–38. <https://doi.org/10.1016/J.ADDBEH.2003.08.027>
72. Tarter, R.E., Mezzich, A.C., Kirisci, L. and Kaczyński, N. (1994) Reliability of the Drug Use Screening Inventory among adolescent alcoholics. *Journal of Child & Adolescent Substance Abuse*, **3**, 25–36. https://doi.org/10.1300/J029v03n01_03
73. Tarter, R.E. and Hegedus, A.M. (1991) The Drug Use Screening Inventory: Its applications in the evaluation and treatment of alcohol and other drug abuse. *Alcohol Health & Research World*, **15**, 65–75.
74. Richards, J.B., Zhang, L., Mitchell, S.H. and de Wit, H. (1999) Delay or probability discounting in a model of impulsive behavior: Effect of alcohol. *Journal of the*

- Experimental Analysis of Behavior*, **71**, 121–43. <https://doi.org/10.1901/jeab.1999.71-121>
75. Olson, E.A., Hooper, C.J., Collins, P. and Luciana, M. (2007) Adolescents' performance on delay and probability discounting tasks: Contributions of age, intelligence, executive functioning, and self-reported externalizing behavior. *Personality and Individual Differences*, **43**, 1886–97. <https://doi.org/10.1016/J.PAID.2007.06.016>
 76. Myerson, J., Green, L. and Warusawitharana, M. (2001) Area under the curve as a measure of discounting. *Journal of the Experimental Analysis of Behavior*, **76**, 235–43. <https://doi.org/10.1901/jeab.2001.76-235>
 77. Borges, A.M., Kuang, J., Milhorn, H. and Yi, R. (2016) An alternative approach to calculating Area-Under-the-Curve (AUC) in delay discounting research. *Journal of the Experimental Analysis of Behavior*, **106**, 145–55. <https://doi.org/10.1002/jeab.219>
 78. Carver, C.S. and White, T.L. (1994) Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS Scales. *Journal of Personality and Social Psychology*, **67**, 319–33. <https://doi.org/10.1037/0022-3514.67.2.319>
 79. Kaufman, A.S. and Kaufman, N.L. (2004) Kaufman Brief Intelligence Test. 2nd ed. AGS Publishing, Circle Pines, MN.
 80. Carskadon, M.A. and Acebo, C. (1993) A self-administered rating scale for pubertal development. *Journal of Adolescent Health*, **14**, 190–5. [https://doi.org/10.1016/1054-139X\(93\)90004-9](https://doi.org/10.1016/1054-139X(93)90004-9)
 81. Petersen, A.C., Crockett, L., Richards, M. and Boxer, A. (1988) A self-report measure of pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*, **17**, 117–33. <https://doi.org/10.1007/BF01537962>

82. Smith, B.W., Mitchell, D.G.V., Hardin, M.G., Jazbec, S., Fridberg, D., Blair, J. et al.
(2009) Neural substrates of reward magnitude, probability, and risk during a wheel of fortune decision-making task. *NeuroImage*, **44**, 600–9.
<https://doi.org/10.1016/j.neuroimage.2008.08.016>
83. Ernst, M., Nelson, E.E., Jazbec, S., McClure, E.B., Monk, C.S., Leibenluft, E. et al.
(2005) Amygdala and nucleus accumbens in responses to receipt and omission of gains in adults and adolescents. *NeuroImage*, **25**, 1279–91.
<https://doi.org/10.1016/j.neuroimage.2004.12.038>
84. Shad, M.U., Bidesi, A.S., Chen, L.-A., Thomas, B.P., Ernst, M. and Rao, U. (2011)
Neurobiology of decision-making in adolescents. *Behavioural Brain Research*, **217**, 67–76. <https://doi.org/10.1016/J.BBR.2010.09.033>
85. Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L. and Petersen, S.E. (2012)
Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage*, **59**, 2142–54.
<https://doi.org/10.1016/j.neuroimage.2011.10.018>
86. Makris, N., Goldstein, J.M., Kennedy, D., Hodge, S.M., Caviness, V.S., Faraone, S. V. et al. (2006) Decreased volume of left and total anterior insular lobule in schizophrenia. *Schizophrenia Research*, **83**, 155–71. <https://doi.org/10.1016/j.schres.2005.11.020>
87. Frazier, J.A., Chiu, S., Breeze, J.L., Makris, N., Lange, N., Kennedy, D.N. et al. (2005)
Structural brain magnetic resonance imaging of limbic and thalamic volumes in pediatric bipolar disorder. *American Journal of Psychiatry*, **162**, 1256–65.
<https://doi.org/10.1176/appi.ajp.162.7.1256>
88. Desikan, R.S., Ségonne, F., Fischl, B., Quinn, B.T., Dickerson, B.C., Blacker, D. et al.

- (2006) An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*, **31**, 968–80.
<https://doi.org/10.1016/j.neuroimage.2006.01.021>
89. Goldstein, J.M., Seidman, L.J., Makris, N., Ahern, T., O'Brien, L.M., Caviness, V.S. et al. (2007) Hypothalamic abnormalities in schizophrenia: sex effects and genetic vulnerability. *Biological Psychiatry*, **61**, 935–45. <https://doi.org/10.1016/j.biopsych.2006.06.027>
 90. Mai, J.K., Majtanik, M. and Paxinos, G. (2016) Atlas of the human brain. 4th ed. Academic Press, Amsterdam.
 91. Benjamin, D.J., Brown, S.A. and Shapiro, J.M. (2013) WHO is “behavioral”? Cognitive ability and anomalous preferences. *Journal of the European Economic Association*, **11**, 1231–55. <https://doi.org/10.1111/jeea.12055>
 92. Burks, S. V., Carpenter, J.P., Goette, L. and Rustichini, A. (2009) Cognitive skills affect economic preferences, strategic behavior, and job attachment. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 7745–50.
<https://doi.org/10.1073/pnas.0812360106>
 93. Dohmen, T., Falk, A., Huffman, D. and Sunde, U. (2018) On the relationship between cognitive ability and risk preference. *Journal of Economic Perspectives*, **32**, 115–34.
<https://doi.org/10.1257/jep.32.2.115>
 94. Cokely, E.T. and Kelley, C.M. (2009) Cognitive abilities and superior decision making under risk: A protocol analysis and process model evaluation. *Judgment and Decision Making*, **4**, 20–33.
 95. Donati, M.A., Panno, A., Chiesi, F. and Primi, C. (2014) A mediation model to explain decision making under conditions of risk among adolescents: The role of fluid intelligence

- and probabilistic reasoning. *Journal of Clinical and Experimental Neuropsychology*, **36**, 588–95. <https://doi.org/10.1080/13803395.2014.918091>
96. Zilverstand, A. and Goldstein, R.Z. (2020) Dual models of drug addiction: the impaired response inhibition and salience attribution model. In: Verdejo-Garcia ABT-C and A, editor. *Cognition and Addiction: A Researcher's Guide from Mechanisms towards Interventions*, Academic Press. p. 17–23. <https://doi.org/https://doi.org/10.1016/B978-0-12-815298-0.00003-4>
 97. Goldstein, R.Z. and Volkow, N.D. (2011) Dysfunction of the prefrontal cortex in addiction: Neuroimaging findings and clinical implications. *Nature Reviews Neuroscience*, **12**, 652–69. <https://doi.org/10.1038/nrn3119>
 98. Rose, E.J., Picci, G. and Fishbein, D.H. (2019) Neurocognitive precursors of substance misuse corresponding to risk, resistance, and resilience pathways: Implications for prevention science. *Frontiers in Psychiatry*, **10**, 399. <https://doi.org/10.3389/fpsy.2019.00399>
 99. Rao, U., Sidhartha, T., Harker, K.R., Bidesi, A.S., Chen, L.A. and Ernst, M. (2011) Relationship between adolescent risk preferences on a laboratory task and behavioral measures of risk-taking. *Journal of Adolescent Health*, **48**, 151–8. <https://doi.org/10.1016/j.jadohealth.2010.06.008>
 100. Blankenstein, N.E., Schreuders, E., Peper, J.S., Crone, E.A. and van Duijvenvoorde, A.C.K. (2018) Individual differences in risk-taking tendencies modulate the neural processing of risky and ambiguous decision-making in adolescence. *NeuroImage*, **172**, 663–73. <https://doi.org/10.1016/j.neuroimage.2018.01.085>
 101. Drouman, V., Bechara, A. and Read, S.J. (2015) Roles of the different sub-regions of the

- insular cortex in various phases of the decision-making process. *Frontiers in Behavioral Neuroscience*, **9**. <https://doi.org/10.3389/fnbeh.2015.00309>
102. Mohr, P.N.C., Biele, G. and Heekeren, H.R. (2010) Neural processing of risk. *Journal of Neuroscience*, **30**, 6613–9. <https://doi.org/10.1523/JNEUROSCI.0003-10.2010>
 103. Kuhn, C.M. and Knutson, B. (2005) The neural basis of financial risk taking. *Neuron*, **47**, 763–70. <https://doi.org/10.1016/j.neuron.2005.08.008>
 104. Clark, L., Bechara, A., Damasio, H., Aitken, M.R.F., Sahakian, B.J. and Robbins, T.W. (2008) Differential effects of insular and ventromedial prefrontal cortex lesions on risky decision-making. *Brain*, **131**, 1311–22. <https://doi.org/10.1093/brain/awn066>
 105. Li, M., Lauharatanahirun, N., Steinberg, L., King-Casas, B., Kim-Spoon, J. and Deater-Deckard, K. (2019) Longitudinal link between trait motivation and risk-taking behaviors via neural risk processing. *Developmental Cognitive Neuroscience*, **40**. <https://doi.org/10.1016/j.dcn.2019.100725>
 106. Kim-Spoon, J., Deater-Deckard, K., Lauharatanahirun, N., Farley, J.P., Chiu, P.H., Bickel, W.K. et al. (2017) Neural interaction between risk sensitivity and cognitive control predicting health risk behaviors among late adolescents. *Journal of Research on Adolescence*, **27**, 674–82. <https://doi.org/10.1111/jora.12295>
 107. Engelmann, J.M., Versace, F., Robinson, J.D., Minnix, J.A., Lam, C.Y., Cui, Y. et al. (2012) Neural substrates of smoking cue reactivity: A meta-analysis of fMRI studies. *NeuroImage*, **60**, 252–62. <https://doi.org/10.1016/j.neuroimage.2011.12.024>
 108. Kühn, S. and Gallinat, J. (2011) Common biology of craving across legal and illegal drugs - a quantitative meta-analysis of cue-reactivity brain response. *European Journal of Neuroscience*, **33**, 1318–26. <https://doi.org/10.1111/j.1460-9568.2010.07590.x>

109. Liang, X., He, Y., Salmeron, B.J., Gu, H., Stein, E.A. and Yang, Y. (2015) Interactions between the salience and default-mode networks are disrupted in cocaine addiction. *Journal of Neuroscience*, **35**, 8081–90. <https://doi.org/10.1523/JNEUROSCI.3188-14.2015>
110. Smith, A.R., Steinberg, L. and Chein, J. (2014) The role of the anterior insula in adolescent decision making. *Developmental Neuroscience*, **36**, 196–209. <https://doi.org/10.1159/000358918>
111. Korucuoglu, O., Harms, M.P., Kennedy, J.T., Golosheykin, S., Astafiev, S. V, Barch, D.M. et al. (2020) Adolescent decision-making under risk: Neural correlates and sex differences. *Cerebral Cortex*, **30**, 2690–706. <https://doi.org/10.1093/cercor/bhz269>
112. Le Berre, A.P., Rauchs, G., La Joie, R., Mézenge, F., Boudehent, C., Vabret, F. et al. (2014) Impaired decision-making and brain shrinkage in alcoholism. *European Psychiatry*, **29**, 125–33. <https://doi.org/10.1016/j.eurpsy.2012.10.002>
113. Klaming, R., Harlé, K.M., Infante, M.A., Bomyea, J., Kim, C. and Spadoni, A.D. (2019) Shared gray matter reductions across alcohol use disorder and posttraumatic stress disorder in the anterior cingulate cortex: A dual meta-analysis. *Neurobiology of Stress*, **10**, 100132.
114. Gowin, J.L., Mackey, S. and Paulus, M.P. (2013) Altered risk-related processing in substance users: Imbalance of pain and gain. *Drug and Alcohol Dependence*, **132**, 13–21. <https://doi.org/10.1016/j.drugalcdep.2013.03.019>
115. Schacht, J.P., Anton, R.F. and Myrick, H. (2013) Functional neuroimaging studies of alcohol cue reactivity: A quantitative meta-analysis and systematic review. *Addiction Biology*, **18**, 121–33. <https://doi.org/10.1111/j.1369-1600.2012.00464.x>

116. Fishbein, D.H., Eldreth, D.L., Hyde, C., Matochik, J.A., London, E.D., Contoreggi, C. et al. (2005) Risky decision making and the anterior cingulate cortex in abstinent drug abusers and nonusers. *Cognitive Brain Research*, **23**, 119–36.
<https://doi.org/10.1016/j.cogbrainres.2004.12.010>
117. Seo, D., Lacadie, C.M., Tuit, K., Hong, K.I., Todd Constable, R. and Sinha, R. (2013) Disrupted ventromedial prefrontal function, alcohol craving, and subsequent relapse risk. *JAMA Psychiatry*, **70**, 727–39. <https://doi.org/10.1001/jamapsychiatry.2013.762>
118. Cheetham, A., Allen, N.B., Whittle, S., Simmons, J., Yücel, M. and Lubman, D.I. (2014) Volumetric differences in the anterior cingulate cortex prospectively predict alcohol-related problems in adolescence. *Psychopharmacology*, **231**, 1731–42.
<https://doi.org/10.1007/s00213-014-3483-8>
119. Acheson, A., Robinson, J.L., Glahn, D.C., Lovallo, W.R. and Fox, P.T. (2009) Differential activation of the anterior cingulate cortex and caudate nucleus during a gambling simulation in persons with a family history of alcoholism: Studies from the Oklahoma Family Health Patterns Project. *Drug and Alcohol Dependence*, **100**, 17–23.
<https://doi.org/10.1016/j.drugalcdep.2008.08.019>
120. Bush, G., Vogt, B.A., Holmes, J., Dale, A.M., Greve, D., Jenike, M.A. et al. (2002) Dorsal anterior cingulate cortex: A role in reward-based decision making. *Proceedings of the National Academy of Sciences of the United States of America*, **99**, 523–8.
<https://doi.org/10.1073/pnas.012470999>
121. Van Duijvenvoorde, A.C.K., Huizenga, H.M., Van Duijvenvoorde, A.C.K., Van Duijvenvoorde, A.C.K., Huizenga, H.M., Somerville, L.H. et al. (2015) Neural correlates of expected risks and returns in risky choice across development. *Journal of*

- Neuroscience*, **35**, 1549–60. <https://doi.org/10.1523/JNEUROSCI.1924-14.2015>
122. Van Leijenhorst, L., Moor, B.G., Op de Macks, Z.A., Rombouts, S.A.R.B., Westenberg, P.M. and Crone, E.A. (2010) Adolescent risky decision-making: Neurocognitive development of reward and control regions. *NeuroImage*, **51**, 345–55.
<https://doi.org/10.1016/j.neuroimage.2010.02.038>
 123. Silverman, M.H., Jedd, K. and Luciana, M. (2015) Neural networks involved in adolescent reward processing: An activation likelihood estimation meta-analysis of functional neuroimaging studies. *NeuroImage*, **122**, 427–39.
<https://doi.org/10.1016/j.neuroimage.2015.07.083>
 124. Seeley, W.W., Menon, V., Schatzberg, A.F., Keller, J., Glover, G.H., Kenna, H. et al. (2007) Dissociable intrinsic connectivity networks for salience processing and executive control. *Journal of Neuroscience*, **27**, 2349–56.
<https://doi.org/10.1523/JNEUROSCI.5587-06.2007>
 125. Zielinski, B.A., Gennatas, E.D., Zhou, J. and Seeley, W.W. (2010) Network-level structural covariance in the developing brain. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 18191–6.
<https://doi.org/10.1073/pnas.1003109107>
 126. Menon, V. and Uddin, L.Q. (2010) Saliency, switching, attention and control: A network model of insula function. *Brain Structure & Function*, **214**, 655–67.
<https://doi.org/10.1007/s00429-010-0262-0>
 127. Jordan, C.J. and Andersen, S.L. (2017) Sensitive periods of substance abuse: Early risk for the transition to dependence. *Developmental Cognitive Neuroscience*, **25**, 29–44.
<https://doi.org/10.1016/j.dcn.2016.10.004>

128. Gowin, J.L., Stewart, J.L., May, A.C., Ball, T.M., Wittmann, M., Tapert, S.F. et al. (2014) Altered cingulate and insular cortex activation during risk-taking in methamphetamine dependence: Losses lose impact. *Addiction*, **109**, 237–47.
<https://doi.org/10.1111/add.12354>
129. Gowin, J.L., Harlé, K.M., Stewart, J.L., Wittmann, M., Tapert, S.F. and Paulus, M.P. (2014) Attenuated insular processing during risk predicts relapse in early abstinent methamphetamine-dependent individuals. *Neuropsychopharmacology*, **39**, 1379–87.
<https://doi.org/10.1038/npp.2013.333>
130. Stewart, J.L., Butt, M., May, A.C., Tapert, S.F. and Paulus, M.P. (2017) Insular and cingulate attenuation during decision making is associated with future transition to stimulant use disorder. *Addiction*, **112**, 1567–77. <https://doi.org/10.1111/add.13839>
131. Hester, R., Nestor, L. and Garavan, H. (2009) Impaired error awareness and anterior cingulate cortex hypoactivity in chronic cannabis users. *Neuropsychopharmacology*, **34**, 2450–8. <https://doi.org/10.1038/npp.2009.67>
132. Umemoto, A., HajiHosseini, A., Yates, M.E. and Holroyd, C.B. (2017) Reward-based contextual learning supported by anterior cingulate cortex. *Cognitive, Affective, & Behavioral Neuroscience*, **17**, 642–51. <https://doi.org/10.3758/s13415-017-0502-3>
133. Holroyd, C.B. and Yeung, N. (2012) Motivation of extended behaviors by anterior cingulate cortex. *Trends in Cognitive Sciences*, **16**, 122–8.
<https://doi.org/10.1016/j.tics.2011.12.008>
134. Ewing Feldstein, S.W., Sakhardande, A. and Blakemore, S.J. (2014) The effect of alcohol consumption on the adolescent brain: A systematic review of MRI and fMRI studies of alcohol-using youth. *NeuroImage: Clinical*, **5**, 420–37.

<https://doi.org/10.1016/j.nicl.2014.06.011>

135. Silveri, M.M., Dager, A.D., Cohen-Gilbert, J.E. and Sneider, J.T. (2016) Neurobiological signatures associated with alcohol and drug use in the human adolescent brain. *Neuroscience and Biobehavioral Reviews*, **70**, 244–59.
<https://doi.org/10.1016/j.neubiorev.2016.06.042>
136. Keough, M.T. and O'Connor, R.M. (2014) Clarifying the measurement and the role of the Behavioral Inhibition System in alcohol misuse. *Alcoholism: Clinical and Experimental Research*, **38**, 1470–9. <https://doi.org/10.1111/acer.12387>
137. O'Connor, R.M., Stewart, S.H. and Watt, M.C. (2009) Distinguishing BAS risk for university students' drinking, smoking, and gambling behaviors. *Personality and Individual Differences*, **46**, 514–9. <https://doi.org/10.1016/j.paid.2008.12.002>
138. Willem, L., Bijttebier, P., Claes, L. and Uytterhaegen, A. (2012) Temperament and problematic alcohol use in adolescence: An examination of drinking motives as mediators. *Journal of Psychopathology and Behavioral Assessment*, **34**, 282–92.
<https://doi.org/10.1007/s10862-012-9279-4>
139. Torsheim, T., Sørli, M.-A., Olseth, A. and Bjørnebekk, G. (2015) Environmental and temperamental correlates of alcohol user patterns in grade 7 students. *Nordic Studies on Alcohol and Drugs*, **32**, 605–22. <https://doi.org/10.1515/nsad-2015-0057>
140. Demidenko, M.I., Huntley, E.D., Martz, M.E. and Keating, D.P. (2019) Adolescent health risk behaviors: Convergent, discriminant and predictive validity of self-report and cognitive measures. *Journal of Youth and Adolescence*, **48**, 1765–83.
<https://doi.org/10.1007/s10964-019-01057-4>
141. Cservenka, A. and Nagel, B.J. (2012) Risky decision-making: An fMRI study of youth at

- high risk for alcoholism. *Alcoholism: Clinical and Experimental Research*, **36**, 604–15.
<https://doi.org/10.1111/j.1530-0277.2011.01650.x>
142. Alarcón, G., Cservenka, A. and Nagel, B.J. (2017) Adolescent neural response to reward is related to participant sex and task motivation. *Brain and Cognition*, **111**, 51–62.
<https://doi.org/10.1016/j.bandc.2016.10.003>
143. Cservenka, A., Herting, M.M., Seghete, K.L.M., Hudson, K.A. and Nagel, B.J. (2013) High and low sensation seeking adolescents show distinct patterns of brain activity during reward processing. *NeuroImage*, **66**, 184–93.
<https://doi.org/10.1016/j.neuroimage.2012.11.003>
144. Hadland, S.E. and Levy, S. (2016) Objective testing: Urine and other drug tests. *Child and Adolescent Psychiatric Clinics of North America*, **25**, 549–65.
<https://doi.org/10.1016/j.chc.2016.02.005>
145. Crone, E.A., van Duijvenvoorde, A.C.K. and Peper, J.S. (2016) Annual Research Review: Neural contributions to risk-taking in adolescence - developmental changes and individual differences. *Journal of Child Psychology and Psychiatry*, **57**, 353–68.
<https://doi.org/10.1111/jcpp.12502>
146. He, J., Hodge, C., Crews, F., He, J. and Hodge, C. (2007) Adolescent cortical development: A critical period of vulnerability for addiction. *Pharmacology Biochemistry and Behavior*, **86**, 189–99.