Title: Lipooligosaccharide, Vag8, and pertussis toxin of *Bordetella pertussis* cooperatively cause coughing in mice

Authors: Yukihiro Hiramatsu¹, Koichiro Suzuki^{1,2}, Takashi Nishida¹, Naoki Onoda¹, Takashi Satoh^{3#}, Shizuo Akira³, Masahito Ikawa⁴, Hiroko Ikeda⁵, Junzo Kamei^{5,6}, Sandra Derouiche^{7,8}, Makoto Tominaga^{7,8}, Yasuhiko Horiguchi^{1*}

Affiliations:

¹Department of Molecular Bacteriology, Research Institute for Microbial Diseases, Osaka University, 3-1 Yamada-oka, Suita, Osaka 565-0871, Japan.

²The Research Foundation for Microbial Diseases of Osaka University (BIKEN), 3-1 Yamada-oka, Suita, Osaka 565-0871, Japan.

³Laboratory of Host Defense, World Premier Institute Immunology Frontier Research Center, Osaka University, 3-1 Yamada-oka, Suita, Osaka 565-0871, Japan.

⁴Department of Experimental Genome Research, Research Institute for Microbial Diseases, Osaka University, 3-1 Yamada-oka, Suita, Osaka 565-0871, Japan.

⁵Department of Pathophysiology and Therapeutics, Hoshi University School of Pharmacy and Pharmaceutical Sciences, 2-4-41 Ebara, Shinagawa-ku, Tokyo 142-8501, Japan.

⁶Global Research Center for Innovative Life Science, Hoshi University School of Pharmacy and Pharmaceutical Sciences, 2-4-41 Ebara, Shinagawa-ku, Tokyo 142-8501, Japan.

⁷Division of Cell Signaling, National Institute for Physiological Sciences, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan.

⁸Thermal Biology Group, Exploratory Research Center on Life and Living Systems (ExCELLS), 5-1 Higashiyama, Myodaiji, Okazaki 444-8787, Japan.

#Current address: The Department of Immune Regulation, Graduate School and Faculty of Medicine, Tokyo Medical and Dental University, 1-5-45, Yushima, Bunkyo-ku, Tokyo 113-8510, Japan.

*Correspondence to: horiguti@biken.osaka-u.ac.jp (Y.H.)

Summary: Whooping cough, a contagious respiratory disease caused by *Bordetella pertussis*, is characterized by paroxysmal coughing; however, the mechanism has not been studied because of the lack of versatile animal models that reproduce the cough. Here, we present a mouse model that reproduces coughing after intranasal inoculation with the bacteria or its components and demonstrate that lipooligosaccharide (LOS), pertussis toxin (PTx), and Vag8 of the bacteria cooperatively function to cause coughing. LOS-induced bradykinin sensitized a transient receptor potential ion channel, TRPV1, which acts as a sensor to evoke the cough reflex. Vag8 further increased bradykinin levels by inhibiting the C1 esterase inhibitor, the major downregulator of the contact system, which generates bradykinin. PTx inhibits intrinsic negative regulation systems for TRPV1 through inactivation of G_i GTPases. Our findings provide a basis for answering long-standing questions on the pathophysiology of the pertussis cough.

Keywords: *Bordetella pertussis*, Cough, Mouse model, Lipooligosaccharide, Vag8, Pertussis toxin, TRPV1, Bradykinin

Introduction: Whooping cough, also referred to as pertussis, is a highly contagious respiratory disease caused by the gram-negative bacterium *Bordetella pertussis* (Kilgore et al., 2016). Generally, the disease can be prevented through vaccination; however, the number of pertussis cases is significantly increasing worldwide, hypothetically because of the rapid waning of immunity induced by recent acellular vaccines and adaptation of the bacteria to escape vaccine-induced immunity (Tan et al., 2015; Twillert et al., 2015). Patients with the disease exhibit various clinical manifestations, including bronchopneumonia, pulmonary hypertension, hypoglycemia, leukocytosis, and paroxysmal coughing. Among these, paroxysmal coughing, which persists for several weeks and imposes a significant burden on infants, is the hallmark of pertussis. However, etiological agents and the mechanism of pertussis cough remain unknown.

B. pertussis is highly adapted to humans; therefore, convenient animal models that reproduce coughing induced by the bacterial infection are difficult to establish. Baboons were recently reported to replicate various pertussis symptoms, including paroxysmal coughing (Warfel et al., 2012a; 2012b; Warfel and Merkel, 2014); however, these large primates are difficult to use in a sufficient number for analytical experiments because of ethical and cost issues. Rats were reported to cough after intrabronchial administration of agarose-embedded or naked *B. pertussis* (Hall et al., 1999; 1998; 1997; 1994; Hornibrook and Ashburn, 1939; Parton et al., 1994; Wardlaw et al., 1993; Woods et al., 1989); however, there have been no reports on further studies with rats that analyze the mechanism underlying the coughing. Meanwhile, we re-established the coughing model of rats infected with *B. bronchiseptica*, which produces many common virulence factors shared with *B. pertussis*, and observed that BspR/BtrA, an anti- σ factor, regulates the ability of *B. bronchiseptica* to cause coughing in rats (Nakamura et al., 2019). Additionally, our rat model exhibited coughing in response to *B. pertussis* infection; however, the cough frequencies were lower than that caused by *B. bronchiseptica*, and the cough production was not reproduced well. Therefore, we attempted to develop an alternative animal model for pertussis cough and focused on mice because of their multiple genetically modified mutants. Mice have not been used for analyses of *B. pertussis*-induced coughing, because they have long been believed to be unable to cough (Elahi et al., 2007; Mackenzie et al., 2004; Scanlon et al., 2019). However, recently, many reports have demonstrated that vagal sensory neurons, which are involved in the cough reflex, innervate the mouse airway (Dinh et al., 2005; Thai Dinh et al., 2005; J. W. Zhang et al., 2006). Mouse coughing in response to certain tussive stimuli has been detected and recorded by independent research groups (Chen et al., 2013; Kamei et al., 1993; 2006; Lin et al., 2019; C. Zhang et al., 2017). In the present study, our mouse-coughing model revealed that lipooligosaccharide (LOS), Vag8, and pertussis toxin (PTx) cooperatively function to produce coughing through the pathway from bradykinin (Bdk) generation to transient receptor potential vanilloid 1 (TRPV1) sensitization.

Results:

C57BL/6 mice respond to *B. pertussis* infection by coughing

We intranasally inoculated mice with *B. pertussis* in a manner similar to that in our previous study on rats (Nakamura et al., 2019) and observed that C57BL/6J mice coughed approximately one week after inoculation (Movies S1 and S2, and Fig. 1A–1C). The incidence of coughing varied depending on the bacterial strains and mouse strains. *B. pertussis* 18323, a laboratory strain, markedly caused coughing while another classical strain, Tohama, did not. Clinical isolates caused varying degrees of coughing. This effect did not correlate with their ability

to colonize the trachea and lungs. Unlike C57BL/6J mice, BALB/c mice did not exhibit coughing upon infection with *B. pertussis* 18323 (Fig. S1A–S1B). The extent of cough production did not differ between the sexes of the mice (data not shown). Thus, we further explored the mechanism of pertussis cough using *B. pertussis* 18323 and male C57BL/6J mice.

Pertussis toxin (PTx) contributes to cough production

Previous studies using rat or baboon models of *B. pertussis* infection suggested the involvement of PTx in cough production based on the observation that infection with a PTx-deficient strain did not cause coughing and that immunization with acellular pertussis vaccines containing pertussis toxoid protected the animals from coughing after the bacterial infection (Hall et al., 1998; Parton et al., 1994; Warfel et al., 2013). Therefore, we examined this idea by immunizing mice with the DPT-IPV (diphtheria-pertussis-tetanus-inactivated polio) vaccine. Consistent with previous studies, the immunized mice were colonized by the bacteria but exhibited no or less coughing (Fig. 1D, 1E, S1C, and S1D). A PTx-deficient strain (Δptx) did not cause coughing in mice (Fig. 1F and 1G). A mutant strain of *B. pertussis* that lacked *bspR*, a transcriptional regulator associated with the cough-inducing ability of *B. bronchiseptica* in rats (Nakamura et al., 2019), caused coughing. This result indicates that the regulatory systems of bacterial gene expression that cause coughing differ between *B. pertussis* and *B. bronchiseptica*. We previously observed that not only living *B. bronchiseptica* but also the bacterial lysates caused coughing in rats (Nakamura et al., 2019). Similarly, in the present study, the lysates of wild-type *B. pertussis* caused coughing similar to the infection with living bacteria (Fig. 2A and 2B). In contrast, the lysates from neither the Δptx mutant nor the mutant producing enzymatically inactive PTx (PTx_{ED}) caused coughing (Fig. 2A, 2B and S2A). The Δptx lysate complemented with purified PTx caused coughing to the same extent as the wild-type lysate (Fig. 2B and 2D). These results indicate that the enzyme (ADP-ribosylating) activity of PTx is required for cough production. However, contrary to our expectation, intranasal inoculation of purified PTx hardly caused coughing, implying that bacterial factors other than PTx are required (Fig. 2A and 2B). The involvement of FhaB, which is also included in the DPT-IPV vaccine, was concluded to be negligible because the lysate of the FhaB-deficient mutant caused coughing as well (Fig. 2C).

Vag8 and lipooligosaccharide (LOS) along with PTx contribute to cough production

To identify additional bacterial factors contributing to cough production, we examined various combinations of bacterial lysates and purified bacterial components. *B. pertussis* exhibits two distinct phenotypic phases, Bvg⁺ and Bvg⁻, in response to environmental alterations (Cotter and Jones, 2003). In the Bvg⁺ phase, the bacteria produce a set of virulence factors including PTx, adenylate cyclase toxin, dermonecrotic toxin, and the machinery and effectors of the type III secretion system. In contrast, in the Bvg⁻ phase, the bacteria shut down the expression of the virulence factors and, instead, express several factors specific to this phenotype. Therefore, in general, the Bvg⁺ phase is considered to represent the virulent phenotype of the organism. Consistently, the lysate from the Bvg⁺ phase-locked mutant (Bvg⁺ lysate) caused coughing, while that from the Bvg⁻ phase-locked mutant (Bvg⁻ lysate) did not (Fig. 2C). The addition of PTx into the Bvg⁻ lysate only partially restored the cough production, compared to the wild-type lysate and the Δptx lysate complemented with PTx (Fig. 2D). The wild-type lysate and Δptx lysate were obtained from the bacteria in the Bvg⁺ phase. Therefore, these results suggest that at least two distinct factors in addition to PTx are involved in cough production. One factor is present in the Bvg⁺ lysate, and the other is a Bvg-independent molecule present both in the Bvg⁺ and Bvg⁻ lysates.

Heat treatment at 56°C for 1 h abrogated the cough induced by the wild-type lysate (Fig. 2D). The addition of PTx to the heat-treated lysate moderately restored the cough production, compared to the wild-type lysate and the Δptx lysate complemented with PTx. Considering that purified PTx alone did not cause coughing, we hypothesize that one factor is heat-labile and the other factor is heat-stable.

We next examined various mutants of *B. pertussis* kept in our laboratory and observed that a mutant strain ($\Delta vag8$) that is deficient in the autotransporter protein Vag8 exhibited only a moderate ability to cause coughing, whereas another autotransporter mutant, $\Delta brkA$, was fully active (Fig. S2B and S2C). Similarly, the $\Delta vag8$ lysate caused moderate coughing compared to the wild-type lysate (Fig. 2E). The addition of a recombinant Vag8 protein (Vag8p) compensated for the activity of the $\Delta vag8$ lysate. Vag8p alone or in combination with PTx did not cause coughing (Fig. 2E). The Bvg⁻ lysate complemented with PTx and Vag8p caused coughing to the same extent as the wild-type lysate (Fig. 2G). These results revealed that Vag8 is the second factor contributing to cough production. Because Vag8 is heat-labile and specific to the Bvg⁺-phase bacteria, we narrowed the third factor to one that is heat-stable and present independently of the Bvg phases. We considered LOS, which is a heat-stable and biologically active outer membrane component, as a probable candidate. The wild-type lysate in which LOS was eliminated with polymyxin B treatment exhibited a reduced ability to cause coughing (Fig. 2F). The addition of a purified LOS preparation restored the coughing. When the lysate from *E. coli* DH5 α was inoculated, instead of that from *B. pertussis* in combination with PTx and Vag8, the mice exhibited coughing (Fig. 2G). Finally, the combination of PTx, Vag8, and LOS caused coughing to the same extent as the wild-type lysate (Fig. 2G and 2H). Thus, we concluded that LOS, PTx, and Vag8 cooperatively cause coughing in mice; however, the combination of any two of these three factors caused no or only a few coughs (Fig. 2H). Notably, commercially available synthetic lipid A of *E. coli* could serve as a substitute for LOS for producing coughing in combination with PTx and Vag8 (Fig. 2H). These results indicate that the lipid A moiety of LOS, which is virtually identical among gram-negative bacteria, is essential to cause coughing. LOS, PTx, and Vag8 from the *B. pertussis* Tohama strain caused coughing in mice, while the living bacteria or the lysate of this strain did not (Fig. 1A, 1B, and 2H). PTx from the Tohama strain compensated for the ability of the Δptx lysate of the 18323 strain to cause coughing (Fig. S2D). In contrast, PTx from both Tohama and 18323 did not compensate the ability of the Δptx lysate of Tohama strain. These results suggest that the Tohama strain may produce an inhibitory factor or multiple inhibitory factors against cough production.

Cough-evoking pathways stimulated by LOS, Vag8, and PTx

The cough reflex involves vagal afferent fibers innervating the airways from the larynx to the proximal bronchi; however, the mechanism through which the afferent nerves are activated to evoke coughing is not fully understood. Nevertheless, it is accepted that transient receptor potential (TRP) ion channels on sensory nerve terminals participate in the cough reflex (Bonvini and Belvisi, 2017; Grace et al., 2011). TRP channels are modulated by cell signals from G protein-coupled receptors (GPCRs) that are activated by ligands such as Bdk, prostanoids, and tachykinins including neurokinins and substance P (Bonvini and Belvisi, 2017; Carr et al., 2003; Geppetti et al., 2009; Grace et al., 2012; 2011; Kumar et al., 2017; Maher et al., 2012; 2011; Mizumura et al., 2009; Sugiura et al., 2002). To explore the stimulatory mechanism through which the *B. pertussis* components cause coughing, we utilized various antagonists for GPCRs (Bdk B1 and B2 receptors, Neurokinin 1–3 receptors, and prostaglandin E2 receptor EP3) and TRP channels (TRPV1, TRPV4, and TRPA1) in our mouse model of coughing. Mice pre-administered antagonists against the B2

receptor (B2R) and TRPV1 exhibited reduced levels of coughing in response to bacterial lysate; however, other reagents were apparently ineffective (Fig. 3A). Similar results were obtained in mice inoculated with LOS, PTx, and Vag8 (Fig. 3B, S3A, and S3B). When both antagonists against B2R and TRPV1 were simultaneously administered, the inhibitory effects were not additively augmented, suggesting that the pathway upstream of B2R to TRPV1 may neither diverge nor converge (Fig. S3C). TRPV1-deficient mice, but not TRPA1-deficient mice, were less responsive to the combination of LOS, PTx, and Vag8 (Fig. 3C); this observation is consistent with the results of the experiments using the antagonists. Thus, we considered LOS, PTx, and Vag8 to cooperatively stimulate the afferent nerves through a pathway from Bdk–B2R to TRPV1 (Fig. 6E). Bdk was reported to enhance the cough reflex in guinea pigs (Al-Shamlan and El-Hashim, 2019; Featherstone et al., 1996; Fox et al., 1996). Bdk concentrations in the bronchoalveolar lavage fluid (BALF) of mice were increased 4 days after intranasal inoculation with LOS–PTx–Vag8 (Fig. 3D). We further analyzed the roles of each bacterial component in the stimulating pathway from Bdk to TRPV1.

Role of LOS

Because the lipid A moiety of LOS was determined to be essential for cough production, we examined the role of the specific receptor for lipid A, toll-like receptor 4 (TLR4), by using TLR4-deficient mice. *B. pertussis* colonized the respiratory organs of TLR4-deficient mice, similar to that in isogenic wild-type mice (Fig. S4A); however, it caused low cough production in the deficient mice (Fig. 3E). Similar results were obtained with intranasal inoculation of the mice with the bacterial lysates or the combination of LOS–PTx–Vag8 (Fig. 3E). The increase in Bdk levels in the BALF of TLR4-deficient mice inoculated with LOS–PTx–Vag8 was considerably lower than that of wild-type mice (Fig. 3F). These results suggest that LOS stimulates Bdk generation via interaction with TLR4. In addition to Bdk, proinflammatory cytokines such as IL-1 β , IL-6, and TNF α tended to increase in wild-type mice but not in TLR4-deficient mice inoculated with LOS–PTx–Vag8 (Fig. S4B). Furthermore, increases in these cytokines were not observed without inoculation of LOS (Fig. S4C), indicating that LOS leads to slight inflammation via TLR4 under these experimental conditions. Bdk is a potent inflammatory mediator that is released from high-molecular-weight kininogen (HK) through the proteolytic activity of plasma kallikrein (PK). HK-deficient (*Kngr1*^{-/-}) mice, which do not produce Bdk, were less responsive to infection with *B. pertussis* or inoculation of the bacterial lysate than isogenic wild-type mice (Fig. 4A and 4B), demonstrating that Bdk participates in a cascade leading to *B. pertussis*-induced cough.

Role of Vag8

Bdk, HK, and PK comprise the kallikrein–kinin system in the plasma contact system, which is initiated and accelerated by two proteases, factor XII (FXII) and plasma prekallikrein (PPK) (Maas et al., 2011; Weidmann et al., 2017; Wu, 2018) (Fig. 6E). Certain stimuli convert zymogen FXII into an active enzyme, FXIIa. FXIIa cleaves PPK to generate PK, which releases Bdk by cleaving HK. In turn, PK cleaves FXII to release FXIIa, providing a positive feedback cycle. FXIIa and PK are inhibited by C1 esterase inhibitor (C1-Inh), the major regulator of the contact system, which is present in the human plasma (Sanrattana et al., 2019). This inhibitory action of C1-Inh was reportedly inhibited by Vag8 (Hovingh et al., 2017; Marr et al., 2011). If this is the case, Vag8 is likely to exacerbate the cough response of mice by upregulating the Bdk level through the inhibition of C1-Inh activity. Indeed, the Bdk levels in the BALF of mice inoculated with LOS and PTx were increased upon the administration of Vag8p (Fig. 4C). These results are consistent with the results described above showing that Vag8p increased the number of coughs

induced by the $\Delta vag8$ lysate or the combination of LOS and PTx (Fig. 2E, 2H, and 4D). This exacerbating effect of Vag8p on cough production was not observed in *Kngr1^{-/-}* mice (Fig. 4D). In addition, the truncated fragment of Vag8p ranging from amino acid positions 102 to 479 ($\Delta C479$), which is unable to bind and inactivate C1-Inh, did not increase the number of coughs in mice inoculated with LOS and PTx; in contrast, other types of truncated Vag8, which retain the ability to inhibit C1-Inh increased the number of coughs similar to the whole passenger-domain fragment of Vag8 (Vag8p) (Onoda et al., 2020) (Fig. 4E–4G). These results indicate that Vag8 exacerbates the cough response of mice by upregulating the Bdk level through the inhibition of C1-Inh activity.

Role of PTx in Bdk-induced sensitization of TRPV1

The above results indicate that LOS and Vag8 function to raise the level of Bdk. Bdk sensitizes TRPV1, which is directly activated by capsaicin (Cap) to evoke coughing in animals (Bonvini and Belvisi, 2017; Caterina et al., 1997; Choi and Hwang, 2018; Fox et al., 1996; Grace et al., 2012; Shin et al., 2002; Trevisani et al., 2004). Previous reports indicated that PTx enhanced the Bdk action by affecting the GTPases of the G_i family, which are the target molecules of the toxin (Diehl et al., 2014; Moss et al., 1988). We therefore examined the role of PTx in the effect of Bdk on TRPV1 using a whole-cell patch clamp technique in HEK293T cells expressing B2R and TRPV1. The cells without ectopic expression did not respond to Bdk and Cap (data not shown). Cap induced a TRPV1-dependent current that was desensitized in the second Cap application (Fig. 5A and 5B) (Bhave et al., 2002; Mohapatra, 2003; Mohapatra and Nau, 2005; Sanz-Salvador et al., 2012). When the cells were treated with Bdk before the second application, the Cap-induced currents were not reduced but were enhanced in the second application, as reported previously (Fox et al., 1996). We observed that this enhancement was further augmented in cells pretreated with PTx but not in those pretreated with PTx_{ED}. Without Bdk, PTx did not influence the desensitization induced by the second Cap application. Similar results were obtained in independent experiments that evaluated the intracellular Ca^{2+} level of dorsal root ganglion (DRG) neurons (Fig. 5D).

The sensitization of TRPV1 by Bdk is considered to be mediated by the protein kinase C (PKC)-dependent pathway, which is activated via B2R-coupled $G_{q/11}$ (Bhave et al., 2003; Mizumura et al., 2009; Numazaki et al., 2002). In contrast, the desensitization (tachyphylaxis) of TRPV1, which partly results from calcineurin-induced dephosphorylation of the channel, is reversed through phosphorylation by the cAMP-dependent protein kinase (PKA) (Bhave et al., 2002; Mohapatra, 2003; Mohapatra and Nau, 2005) (Fig. 6E). We confirmed that the Bdk-induced sensitization of TRPV1 was inhibited by calphostin C (CalpC), a PKC inhibitor, but not by H-89, a PKA inhibitor (Fig. 5C and S5B). Exacerbation of the Bdk-induced sensitization of TRPV1 by PTx was prevented by both inhibitors (Fig. 5C and S5C). An additional inhibitory effect was observed when CalpC and H-89 were simultaneously applied (Fig. 5C). Considering that the PKC inhibitor reduced the Bdk-induced sensitization of TRPV1 and that PTx inactivates G_i GTPases, leading to PKA activation through an increase in the intracellular cAMP level (Katada, 2012), we concluded that H-89 is likely to inhibit PTx-related events. To verify that PTx exacerbated the Bdk-induced sensitization of TRPV1 through the phosphorylation of TRPV1 by PKA, we examined HEK293T cells expressing B2R and a PKA-insensitive mutant of TRPV1 whose Ser¹¹⁷ and Thr³⁷¹, which are phosphorylated by PKA (Bhave et al., 2002; Mohapatra, 2003; Mohapatra and Nau, 2005), were substituted by Ala (Fig. 5B and S5A). In these cells, the effects of PTx were abrogated. These results indicate that PTx exacerbates the Bdk-induced sensitization of TRPV1 through an increase in intracellular cAMP levels, followed by PKA-mediated phosphorylation of TRPV1. B2R has been reported to couple with $G_{q/11}$, G_i , and G_s GTPases (Ewald et al., 1989; Fumi et al., 1991; Gutowski et al., 1991; Lamorte et al., 1993; Liebmann et al., 1996; Mizumura et al., 2009). Our

results indicate that Bdk appears to simultaneously stimulate $G_{q/11}$ - and G_i -dependent pathways through B2R as if it simultaneously applies the accelerator and brakes on TRPV1. We conclude that PTx reverses only the inhibitory effect of Bdk on TRPV1 by uncoupling G_i GTPases from B2R, thereby exacerbating the Bdk-induced sensitization mediated by $G_{q/11}$ GTPases (Fig. 6E). The involvement of G_s GTPases, which stimulate adenylate cyclase and subsequently PKA, in these events is probably negligible, since H-89 did not influence the Bdk-induced sensitization in the absence of PTx.

Another possible role of PTx in the antinociceptive system

Coughs, as well as pain sensation, are initiated by nociceptive stimuli that act on nociceptors including $A\delta$ and C fibers of the sensory nerves, where TRP channels are located. In pain sensation, nociceptive signals are modulated by antinociceptive systems, including the descending antinociceptive system conducted by serotonergic and noradrenergic neurons (Pertovaara, 2013; Tao et al., 2019). In addition, because of the similarities between the neural processing systems for pain- and cough-related inputs, several researchers have recently provided evidence for the existence of possible neural pathways that modulate the processing of the cough reflex, as well as pain sensation (Cinelli et al., 2013; Kubin et al., 2006; McGovern et al., 2017; Mutolo et al., 2014). The modulating systems for respiratory reflexes are suggested to involve the α_2 -adrenergic receptor, which is coupled with G_i GTPases, the target molecule for PTx (Cinelli et al., 2013; Kubin et al., 2006). If PTx inhibits such modulating systems to suppress the cough reflex, coughing will be further exacerbated. We examined this possibility using a recently reported experimental system in which noradrenaline (NA), a ligand for the α_2 adrenergic receptor, reduced the TRPV1 activity of DRG neurons isolated from rats (Chakraborty et al., 2017; Matsushita et al., 2018) (Fig. 6A, 6B, and S6). Consistent with previous studies (Chakraborty et al., 2017; Matsushita et al., 2018), in HEK293T cells expressing TRPV1, B2R and α_{2A} adrenergic receptor (ADR_{A2A}), NA inhibited the Cap-stimulated TRPV1 activity; this inhibitory effect was cancelled by pretreatment of the cells with PTx, but not with PTx_{ED} (Fig. S6). In addition, NA also inhibited the Bdk-induced sensitization of TRPV1 action and that this inhibitory effect of NA was reversed by pretreatment with PTx (Fig. 6A and 6B). Consistent results were obtained using the calcium imaging technique with DRG neurons (Fig. 6C and 6D).

Discussion

Here, we present a series of evidence showing that LOS, Vag8, and PTx of *B. pertussis* cooperatively function to produce coughing in mice (Fig. 6E). (1) LOS stimulates Bdk generation by the kallikrein–kinin system through interaction with TLR4. (2) Vag8 accelerates Bdk generation by inhibiting C1-Inh, which is the major negative regulator of the contact system/kallikrein–kinin system. (3) Bdk sensitizes TRPV1 action, which is regulated by the phosphorylation states mediated by both PKC and PKA; phosphatases (PPase) such as calcineurin desensitize TRPV1 through dephosphorylation. Sensitization of TRPV1 by Bdk depends on B2R– G_q -mediated PKC activation. Additionally, Bdk stimulates the inhibitory pathway of TRPV1 through B2R– G_i -mediated inhibition of adenylate cyclase (AC) and subsequently PKA. (4) PTx reverses the B2R– G_i -mediated inhibition and exacerbates B2R– G_q -mediated sensitization of TRPV1. Consequently, TRPV1 remains in the sensitized state and readily increase nervous excitation to evoke coughing. The sensitized state of TRPV1 continues until G_i , which is ADP-ribosylated by PTx, is replaced by intact ones. (5) In addition, it is possible that PTx inhibits predicted negative regulation systems for the cough reflex that are

relayed by G_i GTPase-coupled receptors, such as NA-stimulated α_2 adrenergic receptors. The negative regulation systems for the cough reflex are yet to be clearly defined; however, step 5 deserves consideration because it may explain the long-lasting coughing in pertussis.

The causative agent of pertussis cough has long been subject to debate. One theory postulated an unidentified “cough toxin,” which is shared by the classical *Bordetella*, *B. pertussis*, *B. parapertussis*, and *B. bronchiseptica*. This idea originated from the fact that infections with the classical *Bordetella* commonly exhibit characteristic paroxysmal coughing in host animals, including humans. According to this idea, the “cough toxin” is not PTx, which is not produced by *B. parapertussis* or *B. bronchiseptica* (Cherry and Paddock, 2014). In contrast, another theory proposed that PTx significantly contributes to pertussis cough, which is supported by previous observations that rats and baboons that are experimentally infected with PTx-deficient *B. pertussis* do not exhibit coughing (Parton et al., 1994) and that immunization with pertussis toxoid protected animals from coughing but not from bacterial colonization (Hall et al., 1998; Warfel et al., 2013); these results are reaffirmed by the present study. Our results demonstrate that PTx is necessary but not sufficient to cause coughing. LOS (lipid A) as the ligand for TLR4 is common among gram-negative bacteria. Vag8 of *B. pertussis* is 97.3% identical to Vag8 of *B. parapertussis* and *B. bronchiseptica*, which themselves are completely identical. Therefore, lipid A and Vag8 are possibly the “cough toxins” shared by the classical *Bordetella*. *B. parapertussis* and *B. bronchiseptica*, but not *B. pertussis*, may produce another virulence factor that corresponds to PTx, which modulates the activity of the ion channels that evoke action potentials for the cough reflex. Identification of such a factor may provide an insight regarding the mechanism of cough production by the classical *Bordetella* other than *B. pertussis*.

Cough production was markedly reduced but not completely eliminated in *Tlr4*^{-/-}, *Kngr1*^{-/-}, or *Trpv1*^{-/-} mice, indicating that the TLR4–Bdk–TRPV1 pathway is not the only pathway that evokes pertussis cough. LOS (lipid A) is essential for cough production. We demonstrated that lipid A-bound TLR4 slightly induced inflammatory responses; however, the mechanism through which it triggers Bdk generation remains unknown. *Tlr4*^{-/-} mice still exhibited slight coughing in response to LOS–PTx–Vag8, probably because LOS marginally induces Bdk or other cough mediators in a TLR4-independent manner. The former possibility is supported by previous studies demonstrating that lipopolysaccharide and lipid A induced Bdk generation in a mixture of FXII, PPK, and HK (Kalter et al., 1983; Roese et al., 1988). The latter possibility is supported by the present study, which demonstrated that *Kngr1*^{-/-} mice still exhibited slight coughing in response to inoculation with the bacteria or bacterial components. In addition to TRPV1, other ion channels, including TRPV4, TRPA1, TRPM8, and the purinergic P2X3 receptor are reportedly involved in signals that evoke the cough reflex (Bonvini and Belvisi, 2017). Additionally, arachidonic acid metabolites, such as prostaglandins and hydroxyeicosatetraenoic acids, and tachykinins including neurokinins and substance P are known to mediate the activation of TRP ion channels (Bonvini and Belvisi, 2017; Carr et al., 2003; Choi and Hwang, 2018; Geppetti et al., 2009; Grace et al., 2012; 2011; Ishiura et al., 2009; Kumar et al., 2017; Maher et al., 2012; 2011; Mizumura et al., 2009; Moriyama et al., 2005; Shin et al., 2002). In the present study, we negated the involvement of TRPV4, TRPA1, neurokinin receptors, and the prostaglandin E2 receptor EP3 through experiments using *Trpa1*^{-/-} mice and antagonists against these ion channels and receptors. However, the cough reflex observed in various diseases is evoked via a network or crosstalk of events involving mediators and ion channels that are not fully understood. Furthermore, the physiology of the cough reflex is not necessarily identical among animal species (Canning, 2008; Mackenzie et al., 2004). Thus, further work is required for understanding the complete mechanism of pertussis cough and applying our results to human

cases.

References:

- Abbasi, F., Miyata, H., Shimada, K., Morohoshi, A., Nozawa, K., Matsumura, T., Xu, Z., Pratiwi, P., Ikawa, M., 2018. RSPH6A is required for sperm flagellum formation and male fertility in mice. *J Cell Sci* 131, jcs221648. doi:10.1242/jcs.221648
- Al-Shamlan, F., El-Hashim, A.Z., 2019. Bradykinin sensitizes the cough reflex via a B2 receptor dependent activation of TRPV1 and TRPA1 channels through metabolites of cyclooxygenase and 12-lipoxygenase. *Respir Res* 20, 46–15. doi:10.1186/s12931-019-1060-8
- Bautista, D.M., Jordt, S.-E., Nikai, T., Tsuruda, P.R., Read, A.J., Poblete, J., Yamoah, E.N., Basbaum, A.I., Julius, D., 2006. TRPA1 Mediates the Inflammatory Actions of Environmental Irritants and Proalgesic Agents. *Cell* 124, 1269–1282. doi:10.1016/j.cell.2006.02.023
- Bhave, G., Hu, H.-J., Glauner, K.S., Zhu, W., Wang, H., Brasier, D.J., Oxford, G.S., Gereau, R.W., 2003. Protein kinase C phosphorylation sensitizes but does not activate the capsaicin receptor transient receptor potential vanilloid 1 (TRPV1). *Proc Natl Acad Sci USA* 100, 12480–12485. doi:10.1073/pnas.2032100100
- Bhave, G., Zhu, W., Wang, H., Brasier, D.J., Oxford, G.S., Gereau, R.W., 2002. cAMP-dependent protein kinase regulates desensitization of the capsaicin receptor (VR1) by direct phosphorylation. *Neuron* 35, 721–731.
- Bonvini, S.J., Belvisi, M.G., 2017. Cough and airway disease: The role of ion channels. *Pulm Pharmacol Ther* 47, 21–28. doi:10.1016/j.pupt.2017.06.009
- Buasri, W., Impoolsup, A., Boonchird, C., Luengchaichawange, A., Prompiboon, P., Petre, J., Panbangred, W., 2012. Construction of Bordetella pertussis strains with enhanced production of genetically-inactivated Pertussis Toxin and Pertactin by unmarked allelic exchange. *BMC Microbiol* 12, 61–16. doi:10.1186/1471-2180-12-61
- Canning, B.J., 2008. The Cough Reflex in Animals: Relevance to Human Cough Research. *Lung* 186, 23–28. doi:10.1007/s00408-007-9054-6
- Carr, M.J., Kollarik, M., Meeker, S.N., Udem, B.J., 2003. A role for TRPV1 in bradykinin-induced excitation of vagal airway afferent nerve terminals. *J Pharmacol Exp Ther* 304, 1275–1279. doi:10.1124/jpet.102.043422
- Caterina, M.J., Leffler, A., Malmberg, A.B., Martin, W.J., Trafton, J., Petersen-Zeit, K.R., Koltzenburg, M., Basbaum, A.I., Julius, D., 2000. Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 288, 306–313. doi:10.1126/science.288.5464.306
- Caterina, M.J., Schumacher, M.A., Tominaga, M., Rosen, T.A., Levine, J.D., Julius, D., 1997. The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 389, 816–824. doi:10.1038/39807
- Chakraborty, S., Elvezio, V., Kaczocha, M., Rebecchi, M., Puopolo, M., 2017. Presynaptic inhibition of transient receptor potential vanilloid type 1 (TRPV1) receptors by noradrenaline in nociceptive neurons. *J Physiol (Lond)* 595, 2639–2660. doi:10.1113/JP273455
- Chen, L., Lai, K., Lomask, J.M., Jiang, B., Zhong, N., 2013. Detection of Mouse Cough Based on Sound Monitoring and Respiratory Airflow Waveforms. *PLoS ONE* 8, e59263–10. doi:10.1371/journal.pone.0059263
- Cherry, J.D., Paddock, C.D., 2014. Pathogenesis and histopathology of pertussis: implications for immunization. *Expert Review of Vaccines* 13, 1115–1123.

doi:10.1586/14760584.2014.935766

Choi, S.-I., Hwang, S.W., 2018. Depolarizing Effectors of Bradykinin Signaling in Nociceptor
Excitation in Pain Perception. *Biomolecules & Therapeutics* 26, 255–267.

doi:10.4062/biomolther.2017.127

Cinelli, E., Bongiani, F., Pantaleo, T., Mutolo, D., 2013. Suppression of the cough reflex by
 α 2-adrenergic receptor agonists in the rabbit. *Physiol Rep* 1, 215–18. doi:10.1002/phy2.122

Cotter, P.A., Jones, A.M., 2003. Phosphorelay control of virulence gene expression in *Bordetella*.
Trends Microbiol 11, 367–373.

Diehl, S.A., McElvany, B., Noubade, R., Seeberger, N., Harding, B., Spach, K., Teuscher, C.,
2014. G proteins Gai1/3 are critical targets for *Bordetella pertussis* toxin-induced vasoactive
amine sensitization. *Infect Immun* 82, 773–782. doi:10.1128/IAI.00971-13

Dinh, Q.T., Mingomataj, E., Quarcoo, D., Groneberg, D.A., Witt, C., Klapp, B.F., Braun, A.,
Fischer, A., 2005. Allergic airway inflammation induces tachykinin peptides expression in
vagal sensory neurons innervating mouse airways. *Clin Exp Allergy* 35, 820–825.

doi:10.1111/j.1365-2222.2005.02264.x

Elahi, S., Holmstrom, J., Gerds, V., 2007. The benefits of using diverse animal models for
studying pertussis. *Trends Microbiol* 15, 462–468. doi:10.1016/j.tim.2007.09.003

Ewald, D.A., Pang, I.H., Sternweis, P.C., Miller, R.J., 1989. Differential G protein-mediated
coupling of neurotransmitter receptors to Ca^{2+} channels in rat dorsal root ganglion neurons
in vitro. *Neuron* 2, 1185–1193. doi:10.1016/0896-6273(89)90185-2

Featherstone, R.L., Parry, J.E., Evans, D.M., Jones, D.M., Olsson, H., Szelke, M., Church, M.K.,
1996. Mechanism of irritant-induced cough: studies with a kinin antagonist and a kallikrein
inhibitor. *Lung* 174, 269–275.

Figurski, D.H., Helinski, D.R., 1979. Replication of an origin-containing derivative of plasmid
RK2 dependent on a plasmid function provided in trans. *Proc Natl Acad Sci USA* 76,
1648–1652.

Fox, A.J., Laloo, U.G., Belvisi, M.G., Bernareggi, M., Chung, K.F., Barnes, P.J., 1996.

Bradykinin-evoked sensitization of airway sensory nerves: a mechanism for ACE-inhibitor
cough. *Nat Med* 2, 814–817.

Fumi, Y., Masato, H., Toshitaka, K., 1991. Evidence for coupling of bradykinin receptors to a
guanine-nucleotide binding protein to stimulate arachidonate liberation in the osteoblast-like
cell line, MC3T3-E1. *Biochim Biophys Acta* 1094, 139–146.

doi:10.1016/0167-4889(91)90001-E

Geppetti, P., Patacchini, R., Nassini, R., Materazzi, S., 2009. Cough: The Emerging Role of the
TRPA1 Channel. *Lung* 188, 63–68. doi:10.1007/s00408-009-9201-3

Grace, M., Birrell, M.A., Dubuis, E., Maher, S.A., Belvisi, M.G., 2012. Transient receptor
potential channels mediate the tussive response to prostaglandin E2 and bradykinin. *Thorax*
67, 891–900. doi:10.1136/thoraxjnl-2011-201443

Grace, M.S., Dubuis, E., Birrell, M.A., Belvisi, M.G., 2011. TRP Channel Antagonists as
Potential Antitussives. *Lung* 190, 11–15. doi:10.1007/s00408-011-9322-3

Gutowski, S., Smrcka, A., Nowak, L., Wu, D.G., Simon, M., Sternweis, P.C., 1991. Antibodies
to the alpha q subfamily of guanine nucleotide-binding regulatory protein alpha subunits
attenuate activation of phosphatidylinositol 4,5-bisphosphate hydrolysis by hormones. *J Biol*
Chem 266, 20519–20524.

Hall, E., Parton, R., Wardlaw, A.C., 1999. Time-course of infection and responses in a coughing
rat model of pertussis. *J Med Microbiol* 48, 95–98. doi:10.1099/00222615-48-1-95

Hall, E., Parton, R., Wardlaw, A.C., 1998. Responses to acellular pertussis vaccines and

- component antigens in a coughing-rat model of pertussis. *Vaccine* 16, 1595–1603.
- Hall, E., Parton, R., Wardlaw, A.C., 1997. Differences in coughing and other responses to intrabronchial infection with *Bordetella pertussis* among strains of rats. *Infect Immun* 65, 4711–4717.
- Hall, E., Parton, R., Wardlaw, A.C., 1994. Cough production, leucocytosis and serology of rats infected intrabronchially with *Bordetella pertussis*. *J Med Microbiol* 40, 205–213. doi:10.1099/00222615-40-3-205
- Hiramatsu, Y., Suzuki, K., Motooka, D., Nakamura, S., Horiguchi, Y., 2020. Expression of small RNAs of *Bordetella pertussis* colonizing murine tracheas. *Microbiology and Immunology* 64, 469–475. doi:10.1111/1348-0421.12791
- Hornibrook, J., Ashburn, L., 1939. A study of experimental pertussis in the young rat. *Public Health Rep* 54, 439–444.
- Hoshino, K., Takeuchi, O., Kawai, T., Sanjo, H., Ogawa, T., Takeda, Y., Takeda, K., Akira, S., 1999. Cutting edge: Toll-like receptor 4 (TLR4)-deficient mice are hyporesponsive to lipopolysaccharide evidence for TLR4 as the Lps gene product. *J Immunol* 162, 3749–3752.
- Hovingh, E.S., van den Broek, B., Kuipers, B., Pinelli, E., Rooijackers, S.H.M., Jongerius, I., 2017. Acquisition of C1 inhibitor by *Bordetella pertussis* virulence associated gene 8 results in C2 and C4 consumption away from the bacterial surface. *PLoS Pathog* 13, e1006531. doi:10.1371/journal.ppat.1006531
- Ishiura, Y., Fujimura, M., Yamamoto, H., Ishiguro, T., Ohkura, N., Myou, S., 2009. COX-2 inhibition attenuates cough reflex sensitivity to inhaled capsaicin in patients with asthma. *J Investig Allergol Clin Immunol* 19, 370–374.
- Kalter, E.S., van Dijk, W.C., Timmerman, A., Verhoef, J., Bouma, B.N., 1983. Activation of purified human plasma prekallikrein triggered by cell wall fractions of *Escherichia coli* and *Staphylococcus aureus*. *J. Infect. Dis.* 148, 682–691. doi:10.1093/infdis/148.4.682
- Kamei, J., Iwamoto, Y., Kawashima, N., Suzuki, T., Nagase, H., Misawa, M., Kasuya, Y., 1993. Possible involvement of μ_2 -mediated mechanisms in μ -mediated antitussive activity in the mouse. *Neuroscience Letters* 149, 169–172. doi:10.1016/0304-3940(93)90763-B
- Kamei, J., Yoshikawa, Y., Saitoh, A., 2006. Effect of N-arachidonoyl-(2-methyl-4-hydroxyphenyl) amine (VDM11), an anandamide transporter inhibitor, on capsaicin-induced cough in mice. *Cough* 2, 2. doi:10.1186/1745-9974-2-2
- Katada, T., 2012. The inhibitory G protein Gi identified as pertussis toxin-catalyzed ADP-ribosylation. *Biological and Pharmaceutical Bulletin* 35, 2103–2111. doi:10.1248/bpb.b212024
- Kilgore, P.E., Salim, A.M., Zervos, M.J., Schmitt, H.-J., 2016. Pertussis: Microbiology, Disease, Treatment, and Prevention. *Clinical Microbiology Reviews* 29, 449–486. doi:10.1128/CMR.00083-15
- Kittaka, H., Uchida, K., Fukuta, N., Tominaga, M., 2017. Lysophosphatidic acid-induced itch is mediated by signalling of LPA 5receptor, phospholipase D and TRPA1/TRPV1. *J Physiol (Lond)* 595, 2681–2698. doi:10.1113/JP273961
- Kubin, L., Alheid, G.F., Zuperku, E.J., McCrimmon, D.R., 2006. Central pathways of pulmonary and lower airway vagal afferents. *Journal of Applied Physiology* 101, 618–627. doi:10.1152/japplphysiol.00252.2006
- Kumar, R., Hazan, A., Geron, M., Steinberg, R., Livni, L., Matzner, H., Priel, A., 2017. Activation of transient receptor potential vanilloid 1 by lipoxygenase metabolites depends on PKC phosphorylation. *The FASEB Journal* 31, 1238–1247. doi:10.1096/fj.201601132R
- Lamorte, V.J., Harootunian, A.T., Spiegel, A.M., Tsien, R.Y., Feramisco, J.R., 1993. Mediation

of growth factor induced DNA synthesis and calcium mobilization by G_q and G_{i2} . J Cell Biol 121, 91–99. doi:10.1083/jcb.121.1.91

Li, T., Saito, C.T., Hikitsuchi, T., Inoguchi, Y., Mitsuishi, H., Saito, S., Tominaga, M., 2019. Diverse sensitivities of TRPA1 from different mosquito species to thermal and chemical stimuli. Sci Rep 9, 20200–13. doi:10.1038/s41598-019-56639-w

Liebmman, C., Graness, A., Ludwig, B., Adomeit, A., Boehmer, A., Boehmer, F.D., Nürnberg, B., Wetzker, R., 1996. Dual bradykinin B_2 receptor signalling in A431 human epidermoid carcinoma cells: Activation of protein kinase C is counteracted by a G_s -mediated stimulation of the cyclic AMP pathway. Biochem J 313, 109–118. doi:10.1042/bj3130109

Lin, A.-H., Athukorala, A., Gleich, G.J., Lee, L.-Y., 2019. Cough responses to inhaled irritants are enhanced by eosinophil major basic protein in awake mice. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology 317, R93–R97. doi:10.1152/ajpregu.00081.2019

Maas, C., Oschatz, C., Renné, T., 2011. The plasma contact system 2.0. Semin. Thromb. Hemost. 37, 375–381. doi:10.1055/s-0031-1276586

Mackenzie, A.J., Spina, D., Page, C.P., 2004. Models used in the development of antitussive drugs. Drug Discovery Today: Disease Models 1, 297–302. doi:10.1016/j.ddmod.2004.10.009

Maher, S.A., Birrell, M.A., Belvisi, M.G., 2012. Prostaglandin E2 Mediates Cough via the EP3 Receptor. Am. J. Respir. Crit. Care Med. 180, 923–928. doi:10.1164/rccm.200903-0388OC

Maher, S.A., Dubuis, E.D., Belvisi, M.G., 2011. G-protein coupled receptors regulating cough. Curr Opin Pharmacol 11, 248–253. doi:10.1016/j.coph.2011.06.005

Marr, N., Shah, N.R., Lee, R., Kim, E.J., Fernandez, R.C., 2011. *Bordetella pertussis* autotransporter Vag8 binds human C1 esterase inhibitor and confers serum resistance. PLoS ONE 6, e20585. doi:10.1371/journal.pone.0020585

Maruyama, K., Takayama, Y., Kondo, T., Ishibashi, K.-I., Sahoo, B.R., Kanemaru, H., Kumagai, Y., Martino, M.M., Tanaka, H., Ohno, N., Iwakura, Y., Takemura, N., Tominaga, M., Akira, S., 2017. Nociceptors Boost the Resolution of Fungal Osteoinflammation via the TRP Channel-CGRP-Jdp2 Axis. Cell Reports 19, 2730–2742. doi:10.1016/j.celrep.2017.06.002

Matsushita, Y., Manabe, M., Kitamura, N., Shibuya, I., 2018. Adrenergic receptors inhibit TRPV1 activity in the dorsal root ganglion neurons of rats. PLoS ONE 13, e0191032–15. doi:10.1371/journal.pone.0191032

McGovern, A.E., Ajayi, I.E., Farrell, M.J., Mazzone, S.B., 2017. A neuroanatomical framework for the central modulation of respiratory sensory processing and cough by the periaqueductal grey. J. Thorac. Dis. 9, 4098–4107. doi:10.21037/jtd.2017.08.119

Minnick, M.F., 1994. Identification of outer membrane proteins of Bartonella bacilliformis. Infect Immun 62, 2644–2648.

Mizumura, K., Sugiura, T., Katanosaka, K., Banik, R.K., Kozaki, Y., 2009. Excitation and sensitization of nociceptors by bradykinin: what do we know? Exp Brain Res 196, 53–65. doi:10.1007/s00221-009-1814-5

Mohapatra, D.P., 2003. Desensitization of Capsaicin-activated Currents in the Vanilloid Receptor TRPV1 Is Decreased by the Cyclic AMP-dependent Protein Kinase Pathway. J Biol Chem 278, 50080–50090. doi:10.1074/jbc.M306619200

Mohapatra, D.P., Nau, C., 2005. Regulation of Ca^{2+} -dependent desensitization in the vanilloid receptor TRPV1 by calcineurin and cAMP-dependent protein kinase. J Biol Chem 280, 13424–13432. doi:10.1074/jbc.M410917200

- Moriyama, T., Higashi, T., Togashi, K., Iida, T., Segi, E., Sugimoto, Y., Tominaga, T., Narumiya, S., Tominaga, M., 2005. Sensitization of TRPV1 by EP 1 and IP reveals peripheral
555 nociceptive mechanism of prostaglandins. *Molecular Pain* 2005 1:1 1, 3.
doi:10.1186/1744-8069-1-3
- Moss, J., Hom, B.E., Hewlett, E.L., Tsai, S.C., Adamik, R., Halpern, J.L., Price, S.R., Manganiello, V.C., 1988. Mechanism of enhanced sensitivity to bradykinin in pertussis toxin-treated fibroblasts: toxin increases bradykinin-stimulated prostaglandin formation. *Mol*
560 *Pharmacol* 34, 279–285.
- Mutolo, D., Cinelli, E., Bongianini, F., Pantaleo, T., 2014. Inhibitory control of the cough reflex by galanin receptors in the caudal nucleus tractus solitarius of the rabbit. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 307, R1358–R1367.
doi:10.1152/ajpregu.00237.2014
- 565 Nakamura, K., Shinoda, N., Hiramatsu, Y., Ohnishi, S., Kamitani, S., Ogura, Y., Hayashi, T., Horiguchi, Y., 2019. BspR/BtrA, an Anti- σ factor, regulates the ability of *Bordetella bronchiseptica* to cause cough in rats. *mSphere* 4. doi:10.1128/MSPHERE.00093-19
- Nishikawa, S., Shinzawa, N., Nakamura, K., Ishigaki, K., Abe, H., Horiguchi, Y., 2016. The bvg-repressed gene brtA, encoding biofilm-associated surface adhesin, is expressed during
570 host infection by *Bordetella bronchiseptica*. *Microbiology and Immunology* 60, 93–105.
doi:10.1111/1348-0421.12356
- Numazaki, M., Tominaga, T., Toyooka, H., Tominaga, M., 2002. Direct phosphorylation of capsaicin receptor VR1 by protein kinase C ϵ and identification of two target serine residues. *J Biol Chem* 277, 13375–13378. doi:10.1074/jbc.C200104200
- 575 Onoda, N., Hiramatsu, Y., Teruya, S., Suzuki, K., Horiguchi, Y., 2020. Identification of the minimum region of *Bordetella pertussis* Vag8 required for interaction with C1 inhibitor. *Microbiology and Immunology* 1348–0421.12799. doi:10.1111/1348-0421.12799
- Park, J., Zhang, Y., Buboltz, A.M., Zhang, X., Schuster, S.C., Ahuja, U., Liu, M., Miller, J.F., Sebahia, M., Bentley, S.D., Parkhill, J., Harvill, E.T., 2012. Comparative genomics of the
580 classical *Bordetella* subspecies: the evolution and exchange of virulence-associated diversity amongst closely related pathogens. *BMC Genomics* 13, 545. doi:10.1186/1471-2164-13-545
- Parkhill, J., Sebahia, M., Preston, A., Murphy, L.D., Thomson, N., Harris, D.E., Holden, M.T.G., Churcher, C.M., Bentley, S.D., Mungall, K.L., Cerdeño-Tárraga, A.M., Temple, L., James, K., Harris, B., Quail, M.A., Achtman, M., Atkin, R., Baker, S., Basham, D., Bason, N.,
585 Cherevach, I., Chillingworth, T., Collins, M., Cronin, A., Davis, P., Doggett, J., Feltwell, T., Goble, A., Hamlin, N., Hauser, H., Holroyd, S., Jagels, K., Leather, S., Moule, S., Norberczak, H., O'Neil, S., Ormond, D., Price, C., Rabinowitsch, E., Rutter, S., Sanders, M., Saunders, D., Seeger, K., Sharp, S., Simmonds, M., Skelton, J., Squares, R., Squares, S., Stevens, K., Unwin, L., Whitehead, S., Barrell, B.G., Maskell, D.J., 2003. Comparative
590 analysis of the genome sequences of *Bordetella pertussis*, *Bordetella parapertussis* and *Bordetella bronchiseptica*. *Nat Genet* 35, 32–40. doi:10.1038/ng1227
- Parton, R., Hall, E., Wardlaw, A.C., 1994. Responses to *Bordetella pertussis* mutant strains and to vaccination in the coughing rat model of pertussis. *J Med Microbiol* 40, 307–312.
- Pertovaara, A., 2013. The noradrenergic pain regulation system_ A potential target for pain
595 therapy. *European Journal of Pharmacology* 716, 2–7. doi:10.1016/j.ejphar.2013.01.067
- Roeise, O., Bouma, B.N., Stadaas, J.O., Aasen, A.O., 1988. Dose dependence of endotoxin-induced activation of the plasma contact system: An in vitro study. *Circ. Shock* 26, 419–430.
- Sanrattana, W., Maas, C., de Maat, S., 2019. SERPINs—From Trap to Treatment. *Front. Med.* 6,

- D624–8. doi:10.3389/fmed.2019.00025
- Sanz-Salvador, L., Andres-Borderia, A., Ferrer-Montiel, A., Planells-Cases, R., 2012. Agonist- and Ca²⁺-dependent desensitization of TRPV1 targets the receptor to lysosomes for degradation. *J Biol Chem*. doi:10.1074/jbc.M111.289751
- Scanlon, K., Skerry, C., Carbonetti, N., 2019. Association of Pertussis Toxin with Severe Pertussis Disease. *Toxins* 11, 373–14. doi:10.3390/toxins11070373
- Sekiya, K., Ohishi, M., Ogino, T., Tamano, K., Sasakawa, C., Abe, A., 2001. Supermolecular structure of the enteropathogenic *Escherichia coli* type III secretion system and its direct interaction with the EspA-sheath-like structure. *Proc Natl Acad Sci USA* 98, 11638–11643. doi:10.1073/pnas.191378598
- Shin, J., Cho, H., Hwang, S.W., Jung, J., Shin, C.Y., Lee, S.-Y., Kim, S.H., Lee, M.G., Choi, Y.H., Kim, J., Haber, N.A., Reichling, D.B., Khasar, S., Levine, J.D., Oh, U., 2002. Bradykinin-12-lipoxygenase-VR1 signaling pathway for inflammatory hyperalgesia. *Proc Natl Acad Sci USA* 99, 10150–10155. doi:10.1073/pnas.152002699
- Skelton, S.K., Wong, K.H., 1990. Simple, efficient purification of filamentous hemagglutinin and pertussis toxin from *Bordetella pertussis* by hydrophobic and affinity interaction. *J Clin Microbiol* 28, 1062–1065.
- Stainer, D.W., Scholte, M.J., 1970. A Simple Chemically Defined Medium for the Production of Phase I *Bordetella pertussis*. *Microbiology* 63, 211–220. doi:10.1099/00221287-63-2-211
- Sugiura, T., Tominaga, M., Katsuya, H., Mizumura, K., 2002. Bradykinin lowers the threshold temperature for heat activation of vanilloid receptor 1. *J. Neurophysiol.* 88, 544–548.
- Tan, T., Dalby, T., Forsyth, K., Halperin, S.A., Heininger, U., Hozbor, D., Plotkin, S., Ulloa-Gutierrez, R., Wirsing von König, C.H., 2015. Pertussis across the globe: Recent epidemiologic trends from 2000 to 2013. *Pediatr. Infect. Dis. J.* 34, e222–32. doi:10.1097/INF.0000000000000795
- Tao, Z.-Y., Wang, P.-X., Wei, S.-Q., Traub, R.J., Li, J.-F., Cao, D.-Y., 2019. The Role of Descending Pain Modulation in Chronic Primary Pain: Potential Application of Drugs Targeting Serotonergic System. *Neural Plasticity* 2019, 1–16. doi:10.1155/2019/1389296
- Thai Dinh, Q., Groneberg, D.A., Peiser, C., Joachim, R.A., Frossard, N., Arck, P.C., Klapp, B.F., Fischer, A., 2005. Expression of substance P and nitric oxide synthase in vagal sensory neurons innervating the mouse airways. *Regul. Pept.* 126, 189–194. doi:10.1016/j.regpep.2004.09.006
- Trevisani, M., Milan, A., Gatti, R., Zanasi, A., Harrison, S., Fontana, G., Morice, A.H., Geppetti, P., 2004. Antitussive activity of iodo-resiniferatoxin in guinea pigs. *Thorax* 59, 769–772. doi:10.1136/thx.2003.012930
- Twillert, I.V., Han, W.G.H., van Els, C.A.C.M., 2015. Waning and aging of cellular immunity to *B. pertussis*. *Pathogens and Disease* ftv071–10. doi:10.1093/femspd/ftv071
- Wang, S., Dai, Y., Fukuoka, T., Yamanaka, H., Kobayashi, K., Obata, K., Cui, X., Tominaga, M., Noguchi, K., 2008. Phospholipase C and protein kinase A mediate bradykinin sensitization of TRPA1: a molecular mechanism of inflammatory pain. *Brain* 131, 1241–1251. doi:10.1093/brain/awn060
- Wardlaw, A.C., Hall, E., Parton, R., 1993. Coughing Rat Model of Pertussis. *Biologicals* 21, 27–29. doi:10.1006/biol.1993.1024
- Warfel, J.M., Beren, J., Kelly, V.K., Lee, G., Merkel, T.J., 2012a. Nonhuman primate model of pertussis. *Infect Immun* 80, 1530–1536. doi:10.1128/IAI.06310-11
- Warfel, J.M., Beren, J., Merkel, T.J., 2012b. Airborne Transmission of *Bordetella pertussis*. *J Infect Dis* 206, 902–906. doi:10.1093/infdis/jis443

Warfel, J.M., Merkel, T.J., 2014. The baboon model of pertussis: effective use and lessons for pertussis vaccines. *Expert Review of Vaccines* 13, 1241–1252. doi:10.1586/14760584.2014.946016

Warfel, J.M., Zimmerman, L.I., Merkel, T.J., 2013. Acellular pertussis vaccines protect against disease but fail to prevent infection and transmission in a nonhuman primate model. *Proc Natl Acad Sci USA* 111, 787–792. doi:10.1073/pnas.1314688110

Weidmann, H., Heikau, L., Long, A.T., Naudin, C., Schlüter, H., Renné, T., 2017. The plasma contact system, a protease cascade at the nexus of inflammation, coagulation and immunity. *BBA - Molecular Cell Research* 1864, 2118–2127. doi:10.1016/j.bbamcr.2017.07.009

Woods, D.E., Franklin, R., Cryz, S.J., Ganss, M., Peppler, M., Ewanowich, C., 1989. Development of a rat model for respiratory infection with *Bordetella pertussis*. *Infect Immun* 57, 1018–1024.

Wu, Y., 2018. The plasma contact system as a modulator of innate immunity. *Current Opinion in Hematology* 25, 389–394. doi:10.1097/MOH.0000000000000448

Zhang, C., Lin, R.-L., Hong, J., Khosravi, M., Lee, L.-Y., 2017. Cough and expiration reflexes elicited by inhaled irritant gases are intensified in ovalbumin-sensitized mice. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 312, R718–R726. doi:10.1152/ajpregu.00444.2016

Zhang, J.W., Walker, J.F., Guardiola, J., Yu, J., 2006. Pulmonary sensory and reflex responses in the mouse. *Journal of Applied Physiology* 101, 986–992. doi:10.1152/japplphysiol.00161.2006

Acknowledgments: We thank K. Kamachi for the clinical strains of *B. pertussis*, K. Minamisawa for pRK2013, and A. Abe for pABB-CR2-Gm. We additionally acknowledge E. Mekada for critically reviewing the manuscript, and E. Hosoyamada, N. Sugi, and N. Ishihara for technical and secretarial assistance. **Funding:** This study was supported by JSPS KAKANHI grants JP26293096, JP17H04075, JP20H03485, JP15H06276, and JP16H06276 (AdAMS); Hyogo Science and Technology Association research grant 2035; and Ohyama Health Foundation research grant.

Author contributions: Yu.H., K.S., and N.O. performed the main experiments. T.N., H.I., and J.K. analyzed mouse coughing using a plethysmograph. T.S., S.A., M.I., and M.T. were involved in the generation of genetically modified mice and in designing the mouse experiments. D.S. performed the electrophysiological study. Yu.H., M.T., and Ya.H. outlined the study and wrote the manuscript with contributions from the other authors. **Declaration of interests:** The authors declare no competing interests.

Fig. 1. Coughing of mice inoculated with *B. pertussis*.

(A–C) Cough production in male C57BL/6J mice inoculated with *B. pertussis* 18323, Tohama, BP140, BP141, BP142, or BP144 (n = 3). (D and E) Cough production in mice immunized with DPT-IPV. Male C57BL/6J mice were immunized with 1/4, 1/10, or 1/25 the human dose of DPT-IPV or 4 mg of aluminum hydroxide (Alum). (F and G) Cough production in male C57BL/6J mice inoculated with wild-type (18323), Δ ptx, or Δ bspR strains of *B. pertussis*. Mice were intranasally inoculated with 5×10^6 CFU of *B. pertussis* strains in 50 μ l of SS medium (Mock). The number of coughs was counted for 5 min/mouse/day for 9 days from days 6 to 14 post-inoculation (A), and the total number is expressed (B, D, and F). The number of bacteria recovered from the

tracheas and lungs was counted on day 14 (C, E, and G). Each plot represents the mean $\pm$ SEM (A).
 695 Each horizontal bar represents the mean $\pm$ SEM (B, D, and F) or geometric mean $\pm$ SD (C, E, and G). The number of mice in each test group is presented in parentheses (D–G).

Fig. 2. Coughing of mice inoculated with *B. pertussis* components.

(A–C) Cough production in mice inoculated with cell lysates from *B. pertussis* wild-type and mutant strains. Mice were intranasally inoculated with cell lysates (50 μ g) of *B. pertussis* 18323

wild-type, Δptx strains, or PTx (200 ng) with or without Δptx cell lysate on days 0 to 4 (arrows in A). The number of coughs was counted for 11 days from days 4 to 14 (A) and expressed as the sum from days 6 to 14 (B). Mice were similarly inoculated with cell lysates of *B. pertussis* wild-type, Bvg⁺-phase locked, Bvg⁻-phase locked, Δptx , or $\Delta fhaB$ strains. The total number of coughs from days 6 to 14 is shown (C). (D–H) Cough production in mice inoculated with various preparations of bacterial components. The number of coughs was counted, and the sum of coughs from days 6 to 14 is shown. The inoculated preparations were as follows: cell lysates (50 μ g) from *B. pertussis* 18323 wild-type (WT) (D–H), Δptx (D), Bvg⁻-phase locked (D and G), $\Delta vag8$ (E), and *E. coli* DH5 α (G); cell lysates of *B. pertussis* 18323 wild-type that were incubated at 56°C for 1 h (D, Heat +) or pretreated with Detoxi-GelTM (polymyxin B, Polmx +) (F); PTx (D, E, G, and H, 200 ng), Vag8 (E, G, and H, 500 ng), and LOS (F–H, 4×10^4 EU) of the 18323 strain; and synthetic lipid A of *E. coli* (H, 4×10^4 EU). PTx, Vag8, and LOS of the Tohama strain are indicated as “Thm”. PBS was used for mock-inoculation. Each plot represents the mean $\pm$ SEM (A). Each horizontal bar represents the mean $\pm$ SEM (B–H). The number of mice in each test group is as follows: n = 5 (A and B), n = 3 or 4 (C), n = 4 (D), n = 5 (E and G), n = 10 (F), n = 4 or 5 (H).

Fig. 3. Involvement of B2R, TRPV1, and TLR4 in *B. pertussis*-induced coughing.

(A and B) Effects of antagonists on cough reflex-related pathways in *B. pertussis*-induced coughing. Mice were inoculated with each antagonist prior to inoculation with *B. pertussis* 18323 cell lysate (A, 50 μ g) or combinations (B) of PTx (200 ng), LOS (4×10^4 EU), and Vag8 (500 ng). The number of coughs was counted as described in Materials and Methods. The sum of coughs from days 6 to 14 is shown. V1, TRPV1; V4, TRPV4; A1, TRPA1 (A). (C and E) Cough production of mice deficient in TRP ion channels (C) or TLR4 (E). Wild-type (WT), *Trpv1*^{-/-}, *Trpa1*^{-/-}, *Trpv1*^{-/-} *Trpa1*^{-/-}, or *Tlr4*^{-/-} mice were inoculated with *B. pertussis* 18323 (F, left panel), the cell lysate of the bacteria (F, center panel), or the combination (C and F, right panel) of PTx, LOS, and Vag8. The sum of coughs from days 6 to 14 is shown. (D and F) Bdk concentrations in the BALF of mice inoculated with the combination of PTx, LOS, and Vag8. Wild-type (WT) and *Tlr4*^{-/-} mice were similarly inoculated with the sample, and the concentrations of Bdk in BALF on days 1, 4, and 10 (D) or 4 (F) were determined using ELISA. Each horizontal bar represents the mean $\pm$ SEM. The number of mice in each test group is as follows: n = 5 (A, B, D, and F), n = 7 to

10 (C), n = 4 or 5 (E). Two-way ANOVA with (F) or without (D) Sidak's multiple-comparison test or unpaired *t*-test (E) was performed for the statistical analyses.

735

Fig. 4. Role of Vag8 in cough production.

(A, B, and D) Cough production of *Kng1*^{-/-} mice. Wild-type (WT) and *Kng1*^{-/-} mice were inoculated with *B. pertussis* 18323 (A); the cell lysate of the bacteria (B); or combinations (D) of

740 PTx, LOS, and Vag8. The sum of coughs from days 6 to 14 (A left panel, B, and D) and the number of the bacteria recovered from the trachea and lungs on day 14 (A right panel) are shown. (C) Bdk concentrations in the BALF of mice inoculated with PTx, LOS, and/or Vag8. (E) Schematic representations of Vag8 and recombinant proteins. The passenger domain of Vag8 (Vag8_p) and truncated derivatives are listed with their names and amino-acid positions. (F) The binding of the Vag8 recombinant proteins to C1-Inh. Relative binding levels of the Vag8 recombinant proteins are expressed as OD₄₅₀ values normalized to those for Vag8_p in the ELISA-based binding assay. Bars represent the means and SEM (n = 3). (G) Cough production of mice inoculated with Vag8_p (500 ng, ca. 8 pmol) or the truncated derivatives (Δ C596, Δ C548, or Δ C479, 24 pmol) along with PTx and LOS. The sum of coughs from days 6 to 14 are shown. Each horizontal bar represents the mean $\pm$ SEM (A left panel, B–D, and G) or geometric mean $\pm$ SD (A right panel). Unpaired *t*-test was performed for the statistical analyses (A and B).

745

750

Fig. 5. Role of PTx in Bdk-induced sensitization of TRPV1.

(A) Current responses of HEK293T cells expressing B2R and wild-type TRPV1 or mutant (TRPV1_{mut}) to repetitive applications of Cap (10 nM) and an intervening application of Bdk (100 nM). Representative traces of Cap-evoked currents are shown. Scales at the left bottom of each trace indicate 1 nA and 10 sec on the ordinate and abscissa, respectively. Horizontal bars below the reagent names indicate the incubation period of each reagent. (B and C) Ratio of the second peak to the initial peak of the current densities induced by Cap. The cells were preincubated with or without 10 ng/ml PTx or the enzymatically inactive derivative of PTx (PTx_{ED}) for 24–30 h prior to the recording (A–C). The cells were stimulated with Bdk and subsequently Cap in the presence of 1 μ M CalpC and/or H-89 (C). Values represent the means $\pm$ SEM (n = 3). See also Fig. S5. (D) Intracellular calcium levels of DRG cells changed in response to repetitive applications of Cap. Isolated DRG cells pretreated with or without 10 ng/ml PTx or PTx_{ED} for 24–30 h were subjected to calcium imaging with transient applications of Cap (1 μ M for 1 min) and Bdk (100 nM for 3 min). Each line represents a single cell isolated from DRG. Nine to ten independent cells were tested for each experiment. The results for Cap-sensitive neurons are strongly colored (upper and lower left panels). (Lower right panel) Fluorescence-intensity ratios of the second peak to the initial peak increased in response to Cap. Values represent the means $\pm$ SEM (n = 3).

Fig. 6. Reversal of the inhibitory effect of NA on Cap-evoked and Bdk-sensitized TRPV1 by PTx.

(A and B) Current responses of HEK293T cells expressing B2R, TRPV1, and ADRA2A to repetitive applications of Cap (10 nM) and intervening applications of Bdk (100 nM) in the presence or absence of NA (100 nM). The cells were treated with 10 ng/ml PTx or PTx_{ED} or untreated prior to the whole-cell patch-clamp recording. Representative traces of Cap-evoked currents (A) and the ratio of the second peak to the initial peak of current densities induced by Cap (B) are shown. Scales at the left bottom of each trace indicate 1 nA and 10 sec on the ordinate and abscissa, respectively. Horizontal bars below the reagent names indicate the incubation period of each reagent (A). Plotted data in (B) represent means $\pm$ SEM (n = 3). (C) Intracellular calcium levels of DRG cells changed in response to repetitive applications of Cap (1 μ M, 1 min) and an intervening application of Bdk (100 nM, 3 min) in the presence of NA (100 nM, 5 min). Temporal changes in the calcium levels are shown. Each line represents a single cell isolated from DRG, and the results from ten independent cells are depicted in each panel. The cells that responded to Cap are represented in highlight colors. The cells that responded to both Cap and NA are represented by thick lines with highlighted colors. The highlighted thin lines indicate cells responsive to Cap but irresponsive to NA. (D) The number of DRG cells that responded to Cap and NA. Thirty-one DRG cells were respectively examined in PTx-untreated and PTx-treated test groups. Note that no cells responded to NA in the PTx treatment group, whereas 5 out of 31 cells responded to NA in the untreated group. (E) The action of LOS, Vag8, and PTx to produce coughing. See Discussion in the main text.